

Pacific Journal of Mathematics

**GEOMETRIC PROPERTIES OF LEVEL CURVES
OF HARMONIC FUNCTIONS AND MINIMAL GRAPHS
IN 2-DIMENSIONAL SPACE FORMS**

JINJU XU AND WEI ZHANG

GEOMETRIC PROPERTIES OF LEVEL CURVES OF HARMONIC FUNCTIONS AND MINIMAL GRAPHS IN 2-DIMENSIONAL SPACE FORMS

JINJU XU AND WEI ZHANG

We study the geometric properties of level curves of harmonic functions and minimal graphs in 2-dimensional space forms using the maximum principle. More precisely, we find two auxiliary functions which consist of tangential derivatives of the curvature of level curves and the norms of the gradient of the solution functions. Then we prove that they satisfy certain elliptic partial differential equations.

1. Introduction

The geometric properties of the level surfaces of solutions of elliptic partial differential equations have been studied for a long time. For instance, a book by Ahlfors [1973] contains the well-known result that level curves of the Green function of a 2-dimensional convex domain are convex curves. Gergen [1931] proved the level surfaces of the Green function of a 3-dimensional star-shaped domain are also star-shaped. Shiffman [1956] studied the convexity of the level curves of immersed minimal surfaces in \mathbb{R}^3 . He proved that if two convex curves in parallel planes in \mathbb{R}^3 bound a minimal surface S then the intersections of all other parallel planes with S are also convex curves. In particular, he obtained that if the boundaries are two circles then intermediate level curves are also circles. Gabriel [1957] proved that the level surfaces of the Green function of a 3-dimensional convex domain are strictly convex. Later, Lewis [1977] extended Gabriel's results to p -harmonic functions in high dimensions. For more related extensions and a survey on this subject, see [Bianchini et al. 2009; Caffarelli and Spruck 1982; Kawohl 1985].

There is also a lot of literature on the quantitative curvature estimates of level surfaces of solutions of elliptic partial differential equations. For 2-dimensional harmonic functions, Talenti [1983] got the following result. Let $\Omega \subset \mathbb{R}^2$ be a domain

Xu was supported by the National Natural Science Foundation of China under Grant 11601311. Zhang was supported by the National Natural Science Foundation of China under Grants 11401278 and 11401280.

MSC2010: primary 35B45, 35B50; secondary 35J05, 35J93.

Keywords: level curves, harmonic functions, minimal graphs, space forms.

and u be a harmonic function with no critical points in Ω . Then the function $\kappa/|\nabla u|$ is harmonic in Ω . Here

$$\kappa = \frac{2u_1u_2u_{12} - u_1^2u_{22} - u_2^2u_{11}}{|\nabla u|^3}$$

is the curvature of the level curves of u . Throughout the paper we use subscripts to represent the derivatives with respect to any orthonormal frames. Similar results can also be seen in [Ortel and Schneider 1983; Longinetti 1983]. Recently, Ma, Ou and Zhang [Ma et al. 2010] generalized the above results to n -dimensional harmonic functions ($2 \leq n < \infty$) and obtained the sharp Gaussian curvature estimates of the level surfaces. See also [Chang et al. 2010; Ma and Zhang 2013; 2014; Wang and Zhang 2012; Zhang and Zhang 2013].

More recently, Kong and Xu [2015] found that if u is a harmonic function of two variables with no critical points, then the function $(\kappa_1u_2 - \kappa_2u_1)/|\nabla u|^3$ is also harmonic. Using this fact, they proved that all the level curves of solutions of the Laplace equation with homogeneous Dirichlet boundary conditions on an annulus are circles. This result can be viewed as a generalization of Shiffman's result on minimal surfaces. In this paper, we extend Kong and Xu's and Shiffman's results to harmonic functions and minimal graphs in 2-dimensional space forms. More precisely, we obtain the following results.

Theorem 1.1. *Suppose that $M^2(c)$ is a 2-dimensional Riemannian manifold with constant sectional curvature c . Let $\Omega \subset M^2(c)$ be a domain and u be a harmonic function with no critical points in Ω . Let κ be the curvature of the level curves of u . Then the function $\varphi = (\kappa_1u_2 - \kappa_2u_1)/|\nabla u|^3$ is also harmonic in Ω .*

For minimal graphs, we have the following similar result.

Theorem 1.2. *Suppose that $M^2(c)$ is a 2-dimensional Riemannian manifold with constant sectional curvature c . Let $\Omega \subset M^2(c)$ be a domain and u satisfy the minimal surface equation*

$$\sum_{ij} a_{ij}u_{ij} = 0 \quad \text{in } \Omega,$$

where $a_{ij} = (1 + |\nabla u|^2)\delta_{ij} - u_iu_j$. Furthermore, assume that there are no critical points of u in Ω . Let κ be the curvature of the level curves of u . Set

$$\psi = \frac{(1 + |\nabla u|^2)^{3/2}}{|\nabla u|^3} \cdot (\kappa_1u_2 - \kappa_2u_1).$$

Then the function ψ satisfies the differential equation

$$\sum_{ij} a_{ij}\psi_{ij} + \sum_i b_i\psi_i = 0 \quad \text{in } \Omega.$$

Here the b_i are bounded functions.

Based on the above theorems, we have the following characterization of geodesic circles.

Remark 1.3. Since $(\kappa_1 u_2 - \kappa_2 u_1)/|\nabla u|$ is the tangential derivative of the curvature of the level curves, the auxiliary functions φ and ψ are independent of the choice of orthonormal frames. Similar to the case of Euclidean space, by the maximum principle, we know that all the level curves of solutions of the Laplace equation or the minimal surface equation with homogeneous Dirichlet boundary conditions on an annulus are geodesic circles.

Now we give the derivative commutation formulas in Riemannian geometry. Let u be a smooth function and R_{ijkl} be coefficients of the Riemannian curvature tensor under orthonormal frames. Here for 2-dimensional space forms $M^2(c)$, we adopt $R_{1212} = c$. Then we have

$$(1-1) \quad u_{ij} - u_{ji} = 0,$$

$$(1-2) \quad u_{ijk} - u_{ikj} = \sum_m u_m R_{mijk},$$

$$(1-3) \quad u_{ijkl} - u_{ijlk} = \sum_m u_{mj} R_{mikl} + \sum_m u_{im} R_{mjkl}.$$

For more details, one can consult any book on Riemannian geometry, such as [Chern et al. 1999].

In this paper, all the summation indices i, j, k, l and m run from 1 to 2. In Section 2, we prove Theorem 1.1. In Section 3, we prove Theorem 1.2.

2. Level curves of harmonic functions

In this section, we focus on the calculation of harmonic functions in 2-dimensional space forms.

Let $\Omega \subset M^2(c)$ be a domain and u be a harmonic function defined in Ω with no critical points. Set

$$\varphi = f(|\nabla u|)(\kappa_1 u_2 - \kappa_2 u_1),$$

where κ is the curvature of the level curves and f is a smooth function of one variable defined on the interval $(0, +\infty)$ which will be determined later. For a suitable choice of f , we will prove that φ is also a harmonic function in Ω , i.e., the function φ satisfies

$$(2-1) \quad \Delta \varphi = 0 \quad \text{in } \Omega.$$

In order to prove (2-1) at an arbitrary point $x_0 \in \Omega$, we may choose the orthonormal frames such that

$$u_1(x_0) = 0, \quad u_2(x_0) = |\nabla u|(x_0) > 0.$$

From now on, all the calculations will be done at the fixed point x_0 unless otherwise specified.

By taking the first derivative of φ , we have

$$(2-2) \quad \varphi_i = f'(|\nabla u|)_i \cdot (\kappa_1 u_2 - \kappa_2 u_1) + f \cdot (\kappa_{1i} u_2 + \kappa_1 u_{2i} - \kappa_{2i} u_1 - \kappa_2 u_{1i}).$$

Differentiating (2-2) once more, we have

$$\begin{aligned} \varphi_{ii} &= f''(|\nabla u|)_i^2 \cdot \kappa_1 u_2 + f'(|\nabla u|)_{ii} \cdot \kappa_1 u_2 + 2f'(|\nabla u|)_i \cdot (\kappa_{1i} u_2 + \kappa_1 u_{2i} - \kappa_{2i} u_1 - \kappa_2 u_{1i}) \\ &\quad + f \cdot (\kappa_{1ii} u_2 + 2\kappa_{1i} u_{2i} + \kappa_1 u_{2ii} - 2\kappa_{2i} u_{1i} - \kappa_2 u_{1ii}); \end{aligned}$$

hence

$$\begin{aligned} (2-3) \quad \Delta \varphi &= u_2 f \sum_i k_{1ii} + [2u_2 f'(|\nabla u|)_1 + 2fu_{12}] \cdot \kappa_{11} \\ &\quad + [2u_2 f'(|\nabla u|)_2 + 2fu_{22} - 2fu_{11}] \cdot \kappa_{12} + [-2fu_{12}] \cdot \kappa_{22} \\ &\quad + \left[u_2 f'' \sum_i (|\nabla u|)_i^2 + u_2 f' \sum_i (|\nabla u|)_{ii} \right. \\ &\quad \left. + 2f' \sum_i (|\nabla u|)_i u_{2i} + f \sum_i u_{2ii} \right] \cdot \kappa_{11} \\ &\quad + \left[-2f' \sum_i (|\nabla u|)_i u_{1i} - f \sum_i u_{1ii} \right] \cdot \kappa_2. \end{aligned}$$

Direct calculation yields

$$\begin{aligned} (2-4) \quad (|\nabla u|)_i &= \frac{1}{|\nabla u|} \sum_j u_j u_{ji}, \\ (|\nabla u|)_{ii} &= \frac{1}{|\nabla u|} \sum_j u_{ji}^2 + \frac{1}{|\nabla u|} \sum_j u_j u_{jii} - \frac{1}{|\nabla u|^3} \sum_{jk} u_j u_{ji} u_k u_{ki}. \end{aligned}$$

Then at the point x_0 ,

$$(2-5) \quad (|\nabla u|)_i = u_{2i}, \quad (|\nabla u|)_{ii} = \frac{u_{1i}^2}{u_2} + u_{2ii}.$$

By the commutation formulas (1-1)–(1-2), we have

$$(2-6) \quad \sum_i u_{1ii} = \sum_i u_{i1i} = \sum_i \left[u_{ii1} + \sum_m u_m R_{mi1i} \right] = 0,$$

$$(2-7) \quad \sum_i u_{2ii} = \sum_i u_{i2i} = \sum_i \left[u_{ii2} + \sum_m u_m R_{mi2i} \right] = u_2 \cdot c.$$

Putting (2-5)–(2-7) into (2-3), we obtain

$$(2-8) \quad \Delta\varphi = u_2 f \sum_i \kappa_{1ii} + (2u_2 f' + 2f) u_{12} \cdot \kappa_{11} + (-2u_2 f' - 4f) u_{11} \cdot \kappa_{12} \\ + (-2f) u_{12} \cdot \kappa_{22} + (u_2 f'' + 3f') (u_{11}^2 + u_{12}^2) \cdot \kappa_1 + (u_2^2 f' + u_2 f) \cdot \kappa_1 \cdot c.$$

To compute the first term in (2-8), we should get the formula for $\Delta\kappa$ at a general point in advance. Recalling the curvature formula for the level curves, we have

$$(2-9) \quad |\nabla u|^3 \cdot \kappa = 2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11}.$$

By applying the Laplace operator on both sides of (2-9) and then using (2-4), we obtain

$$(2-10) \quad \Delta\kappa = \frac{1}{|\nabla u|^3} \sum_i (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11})_{ii} - \frac{6}{|\nabla u|} \sum_i (|\nabla u|)_i \cdot \kappa_i \\ - \frac{6}{|\nabla u|^2} \sum_i (|\nabla u|)_i^2 \cdot \kappa - \frac{3}{|\nabla u|} \sum_i (|\nabla u|)_{ii} \cdot \kappa \\ = \frac{1}{|\nabla u|^3} \sum_i (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11})_{ii} - \frac{6}{|\nabla u|^2} (u_1 u_{11} + u_2 u_{12}) \cdot \kappa_1 \\ - \frac{6}{|\nabla u|^2} (u_1 u_{12} - u_2 u_{11}) \cdot \kappa_2 - \frac{1}{|\nabla u|^2} \left[9(u_{11}^2 + u_{12}^2) + 3 \sum_{ij} u_j u_{jii} \right] \cdot \kappa.$$

Now the commutation formulas (1-1)–(1-2) yield

$$(2-11) \quad \sum_{ij} u_j u_{jii} = \sum_{ij} u_j \left[u_{iij} + \sum_m u_m R_{miji} \right] = \sum_{ijm} u_j u_m R_{miji} = |\nabla u|^2 \cdot c.$$

By inserting (2-11) into (2-10), we have

$$(2-12) \quad \Delta\kappa = \frac{1}{|\nabla u|^3} \sum_i (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11})_{ii} - \frac{6}{|\nabla u|^2} (u_1 u_{11} + u_2 u_{12}) \cdot \kappa_1 \\ - \frac{6}{|\nabla u|^2} (u_1 u_{12} - u_2 u_{11}) \cdot \kappa_2 - \frac{9}{|\nabla u|^2} (u_{11}^2 + u_{12}^2) \cdot \kappa - 3\kappa \cdot c.$$

Straightforward computation gives

$$(2-13) \quad \sum_i (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11})_{ii} \\ = \sum_i [2u_{1ii} u_2 u_{12} + 2u_1 u_{2ii} u_{12} + 2u_1 u_2 u_{12ii} - 2u_1 u_{1ii} u_{22} \\ - u_1^2 u_{22ii} - 2u_2 u_{2ii} u_{11} - u_2^2 u_{11ii} + 4u_{1i} u_{2i} u_{12} + 4u_{1i} u_{2i} u_{12i} \\ + 4u_1 u_{2i} u_{12i} - 2u_{1i}^2 u_{22} - 4u_1 u_{1i} u_{22i} - 2u_{2i}^2 u_{11} - 4u_2 u_{2i} u_{11i}]$$

$$\begin{aligned}
&= 2(u_2 u_{12} + u_1 u_{11}) \sum_i u_{1ii} + 2(u_1 u_{12} - u_2 u_{11}) \sum_i u_{2ii} \\
&\quad + 2u_1 u_2 \sum_i u_{12ii} - u_1^2 \sum_i u_{22ii} - u_2^2 \sum_i u_{11ii} \\
&\quad + 4 \sum_i (u_2 u_{1i} + u_1 u_{2i}) u_{12i} + 4 \sum_i (u_1 u_{1i} - u_2 u_{2i}) u_{11i} \\
&\triangleq I_1 + I_2 + I_3,
\end{aligned}$$

where

$$\begin{aligned}
I_1 &= 2(u_2 u_{12} + u_1 u_{11}) \sum_i u_{1ii} + 2(u_1 u_{12} - u_2 u_{11}) \sum_i u_{2ii}, \\
I_2 &= 2u_1 u_2 \sum_i u_{12ii} - u_1^2 \sum_i u_{22ii} - u_2^2 \sum_i u_{11ii}, \\
I_3 &= 4 \sum_i (u_2 u_{1i} + u_1 u_{2i}) u_{12i} + 4 \sum_i (u_1 u_{1i} - u_2 u_{2i}) u_{11i}.
\end{aligned}$$

We deal with the terms I_1 , I_2 and I_3 consecutively. By (2-6)–(2-7), we have

$$\begin{aligned}
(2-14) \quad I_1 &= 2(u_2 u_{12} + u_1 u_{11}) \sum_{im} u_m R_{mi1i} + 2(u_1 u_{12} - u_2 u_{11}) \sum_{im} u_m R_{mi2i} \\
&= 2(u_2 u_{12} + u_1 u_{11}) \cdot u_1 \cdot c + 2(u_1 u_{12} - u_2 u_{11}) \cdot u_2 \cdot c \\
&= 2|\nabla u|^3 \cdot \kappa \cdot c.
\end{aligned}$$

By the commutation formulas (1-1)–(1-3), we have

$$u_{jkii} = u_{iijk} + \sum_m u_{mk} R_{miji} + \sum_m u_{mj} R_{miki} + 2 \sum_m u_{mi} R_{mjki}.$$

It follows that

$$\begin{aligned}
(2-15) \quad I_2 &= 2u_1 u_2 \cdot \left[\sum_{im} u_{m2} R_{mi1i} + \sum_{im} u_{m1} R_{mi2i} + 2 \sum_{im} u_{mi} R_{m12i} \right] \\
&\quad - u_1^2 \cdot \left[2 \sum_{im} u_{m2} R_{mi2i} + 2 \sum_{im} u_{mi} R_{m22i} \right] \\
&\quad - u_2^2 \cdot \left[2 \sum_{im} u_{m1} R_{mi1i} + 2 \sum_{im} u_{mi} R_{m11i} \right] \\
&= 2u_1 u_2 \cdot 4u_{12} \cdot c - u_1^2 \cdot (-4u_{11}) \cdot c - u_2^2 \cdot 4u_{11} \cdot c \\
&= (8u_1 u_2 u_{12} + 4u_1^2 u_{11} - 4u_2^2 u_{11}) \cdot c.
\end{aligned}$$

By the commutation formulas (1-1)–(1-2),

$$(2-16) \quad u_{121} = u_{112} + \sum_m u_m R_{m121} = u_{112} + u_2 \cdot c,$$

$$(2-17) \quad u_{122} = u_{221} + \sum_m u_m R_{m212} = -u_{111} + u_1 \cdot c.$$

Then we have

$$(2-18) \quad I_3 = 8(u_1 u_{11} - u_2 u_{12})u_{111} + 8(u_1 u_{12} + u_2 u_{11})u_{112} + (8u_1 u_2 u_{12} + 4u_2^2 u_{11} - 4u_1^2 u_{11}) \cdot c.$$

Combining (2-13)–(2-15) and (2-18), we get

$$(2-19) \quad \begin{aligned} & \sum_i (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11})_{ii} \\ &= 8(u_1 u_{11} - u_2 u_{12})u_{111} + 8(u_1 u_{12} + u_2 u_{11})u_{112} \\ & \quad + 2|\nabla u|^3 \cdot \kappa \cdot c + 16u_1 u_2 u_{12} \cdot c. \end{aligned}$$

Now let us explore the relations between u_{111} , u_{112} and κ_1 , κ_2 . Taking the first derivative on both sides of (2-9) and using (2-4), (2-16) and (2-17), we obtain

$$\begin{aligned} & (u_1^2 - u_2^2) \cdot u_{111} + 2u_1 u_2 \cdot u_{112} \\ &= |\nabla u|^3 \cdot \kappa_1 + 3|\nabla u|(u_1 u_{11} + u_2 u_{12}) \cdot \kappa - 2u_1(u_{11}^2 + u_{12}^2) - 2u_1 u_2^2 \cdot c, \\ & -2u_1 u_2 \cdot u_{111} + (u_1^2 - u_2^2) \cdot u_{112} \\ &= |\nabla u|^3 \cdot \kappa_2 + 3|\nabla u|(u_1 u_{12} - u_2 u_{11}) \cdot \kappa - 2u_2(u_{11}^2 + u_{12}^2) - 2u_1^2 u_2 \cdot c. \end{aligned}$$

Thus we have

$$(2-20) \quad u_{111} = \frac{u_1^2 - u_2^2}{|\nabla u|} \cdot \kappa_1 - \frac{2u_1 u_2}{|\nabla u|} \cdot \kappa_2 + \frac{3}{|\nabla u|}(u_1 u_{11} - u_2 u_{12}) \cdot \kappa - \frac{2u_1(u_1^2 - 3u_2^2)}{|\nabla u|^4}(u_{11}^2 + u_{12}^2) + \frac{2u_1 u_2^2}{|\nabla u|^2} \cdot c,$$

and

$$(2-21) \quad u_{112} = \frac{2u_1 u_2}{|\nabla u|} \cdot \kappa_1 + \frac{u_1^2 - u_2^2}{|\nabla u|} \cdot \kappa_2 + \frac{3}{|\nabla u|}(u_2 u_{11} + u_1 u_{12}) \cdot \kappa - \frac{2u_2(3u_1^2 - u_2^2)}{|\nabla u|^4}(u_{11}^2 + u_{12}^2) - \frac{2u_1^2 u_2}{|\nabla u|^2} \cdot c.$$

Hence the formula (2-19) reduces to

$$(2-22) \quad \begin{aligned} \sum_i (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11})_{ii} \\ = 8|\nabla u|(u_1 u_{11} + u_2 u_{12}) \cdot \kappa_1 + 8|\nabla u|(u_1 u_{12} - u_2 u_{11}) \cdot \kappa_2 \\ + 8|\nabla u|(u_{11}^2 + u_{12}^2) \cdot \kappa + 2|\nabla u|^3 \cdot \kappa \cdot c. \end{aligned}$$

By (2-12) and (2-22), we have

$$(2-23) \quad \Delta\kappa = \frac{2}{|\nabla u|^2}(u_1 u_{11} + u_2 u_{12}) \cdot \kappa_1 + \frac{2}{|\nabla u|^2}(u_1 u_{12} - u_2 u_{11}) \cdot \kappa_2 \\ - \frac{1}{|\nabla u|^2}(u_{11}^2 + u_{12}^2) \cdot \kappa - \kappa \cdot c.$$

Then at the point x_0 , we take the first derivative of (2-23). With (2-5) and (2-16) in hand, we obtain

$$(2-24) \quad (\Delta\kappa)_1 = \frac{2}{u_2} u_{12} \cdot \kappa_{11} - \frac{2}{u_2} u_{11} \cdot \kappa_{12} + \left[\frac{2}{u_2} u_{112} + \frac{1}{u_2^2} u_{11}^2 - \frac{3}{u_2^2} u_{12}^2 \right] \cdot \kappa_1 \\ + \left[-\frac{2}{u_2} u_{111} + \frac{4}{u_2^2} u_{11} u_{12} \right] \cdot \kappa_2 \\ + \left[-\frac{2}{u_2^2} (u_{11} u_{111} + u_{12} u_{112}) + \frac{2}{u_2^3} (u_{11}^2 + u_{12}^2) u_{12} \right] \cdot \kappa \\ + \left[\kappa_1 - \frac{2}{u_2} u_{12} \cdot \kappa \right] \cdot c.$$

Now, the equations (2-20) and (2-21) are simplified as

$$(2-25) \quad u_{111} = -u_2 \kappa_1 + \frac{3}{u_2} u_{11} u_{12},$$

$$(2-26) \quad u_{112} = -u_2 \kappa_2 - \frac{1}{u_2} u_{11}^2 + \frac{2}{u_2} u_{12}^2.$$

Putting (2-25)–(2-26) into (2-24), one achieves

$$\begin{aligned} (\Delta\kappa)_1 = \frac{2}{u_2} u_{12} \cdot \kappa_{11} - \frac{2}{u_2} u_{11} \cdot \kappa_{12} + \left[-\frac{3}{u_2^2} u_{11}^2 + \frac{1}{u_2^2} u_{12}^2 \right] \cdot \kappa_1 + \left[-\frac{4}{u_2^2} u_{11} u_{12} \right] \cdot \kappa_2 \\ + \left[-\frac{2}{u_2^3} (u_{11}^2 + u_{12}^2) u_{12} \right] \cdot \kappa + \left[\kappa_1 - \frac{2}{u_2} u_{12} \cdot \kappa \right] \cdot c. \end{aligned}$$

Therefore, by the commutation formulas (1-1)–(1-2), we get

$$\begin{aligned}
 (2-27) \quad & \sum_i \kappa_{1ii} \\
 &= \sum_i \left[\kappa_{ii1} + \sum_m \kappa_m R_{mi1i} \right] \\
 &= (\Delta\kappa)_1 + \kappa_1 \cdot c \\
 &= \frac{2}{u_2} u_{12} \cdot \kappa_{11} - \frac{2}{u_2} u_{11} \cdot \kappa_{12} + \left[-\frac{3}{u_2^2} u_{11}^2 + \frac{1}{u_2^2} u_{12}^2 \right] \cdot \kappa_1 \\
 &\quad + \left[-\frac{4}{u_2^2} u_{11} u_{12} \right] \cdot \kappa_2 + \left[-\frac{2}{u_2^3} (u_{11}^2 + u_{12}^2) u_{12} \right] \cdot \kappa + \left[2\kappa_1 - \frac{2}{u_2} u_{12} \cdot \kappa \right] \cdot c.
 \end{aligned}$$

Thanks to (2-27), the formula (2-8) reduces to

$$\begin{aligned}
 \Delta\varphi = & [(2u_2 f' + 4f)u_{12}] \cdot \kappa_{11} + [(-2u_2 f' - 6f)u_{11}] \cdot \kappa_{12} + [-2fu_{12}] \cdot \kappa_{22} \\
 & + \left[\left(u_2 f'' + 3f' - \frac{3f}{u_2} \right) u_{11}^2 + \left(u_2 f'' + 3f' + \frac{f}{u_2} \right) u_{12}^2 \right] \cdot \kappa_1 \\
 & + \left[-\frac{4f}{u_2} u_{11} u_{12} \right] \cdot \kappa_2 + \left[-\frac{2f}{u_2^2} (u_{11}^2 + u_{12}^2) u_{12} \right] \cdot \kappa \\
 & + [(u_2^2 f' + 3u_2 f) \cdot \kappa_1 - 2fu_{12} \cdot \kappa] \cdot c.
 \end{aligned}$$

At the point x_0 , by (2-23), we have

$$\kappa_{22} = -\kappa_{11} + \frac{2}{u_2} u_{12} \cdot \kappa_1 - \frac{2}{u_2} u_{11} \cdot \kappa_2 - \frac{1}{u_2^2} (u_{11}^2 + u_{12}^2) \cdot \kappa - \kappa \cdot c.$$

Thus

$$\begin{aligned}
 (2-28) \quad \Delta\varphi = & (2u_2 f' + 6f)u_{12} \cdot \kappa_{11} + (-2u_2 f' - 6f)u_{11} \cdot \kappa_{12} \\
 & + \left(u_2 f'' + 3f' - \frac{3f}{u_2} \right) (u_{11}^2 + u_{12}^2) \cdot \kappa_1 + (u_2^2 f' + 3u_2 f) \cdot \kappa_1 \cdot c.
 \end{aligned}$$

By (2-2), we have

$$(2-29) \quad \kappa_{11} = \frac{1}{u_2 f} [\varphi_1 - (u_2 f' + f)u_{12} \cdot \kappa_1 + fu_{11} \cdot \kappa_2],$$

$$(2-30) \quad \kappa_{12} = \frac{1}{u_2 f} [\varphi_2 + (u_2 f' + f)u_{11} \cdot \kappa_1 + fu_{12} \cdot \kappa_2].$$

Putting (2-29)–(2-30) into (2-28), we finally get

$$(2-31) \quad \Delta\varphi = \left(\frac{2f'}{f} + \frac{6}{u_2} \right) \cdot (u_{12}\varphi_1 - u_{11}\varphi_2) \\ + \left(u_2 f'' - \frac{2u_2 f'^2}{f} - 5f' - \frac{9f}{u_2} \right) (u_{11}^2 + u_{12}^2) \cdot \kappa_1 + (u_2^2 f' + 3u_2 f) \cdot \kappa_1 \cdot c.$$

If we let

$$f(t) = t^{-3},$$

then all of the terms on the right-hand side of (2-31) vanish. This completes the proof of [Theorem 1.1](#). \square

3. Level curves of minimal graphs

Along the same lines as in [Section 2](#), in this section we deal with the minimal graphs in 2-dimensional space forms.

Let $\Omega \subset M^2(c)$ be a domain and u be a solution with no critical points of the minimal surface equation

$$(3-1) \quad \sum_{ij} a_{ij} u_{ij} = 0 \quad \text{in } \Omega,$$

where

$$a_{ij} = (1 + |\nabla u|^2) \delta_{ij} - u_i u_j.$$

Set

$$\psi = g(|\nabla u|)(\kappa_1 u_2 - \kappa_2 u_1),$$

where κ is the curvature of the level curves and g is a smooth function of one variable defined on the interval $(0, +\infty)$ to be determined later. For a suitable choice of g , we will prove that the function ψ satisfies

$$(3-2) \quad \sum_{ij} a_{ij} \psi_{ij} + \sum_i b_i \psi_i = 0 \quad \text{in } \Omega.$$

Here the b_i are bounded functions.

In order to prove (3-2) at an arbitrary point $x_0 \in \Omega$, we may choose the orthonormal frames such that

$$u_1(x_0) = 0,$$

$$u_2(x_0) = |\nabla u|(x_0) > 0.$$

From now on, all the calculations will be done at the fixed point x_0 unless otherwise specified.

By taking the derivative of ψ , we have

$$(3-3) \quad \psi_i = g'(|\nabla u|)_i \cdot (\kappa_1 u_2 - \kappa_2 u_1) + g \cdot (\kappa_{1i} u_2 + \kappa_{1u2i} - \kappa_{2i} u_1 - \kappa_{2u1i}).$$

Differentiating (3-3) once more, we have

$$\begin{aligned} \psi_{ij} &= g''(|\nabla u|)_j (|\nabla u|)_i \cdot \kappa_1 u_2 + g'(|\nabla u|)_{ij} \cdot \kappa_1 u_2 + g'(|\nabla u|)_i \cdot (\kappa_{1j} u_2 + \kappa_{1u2j} - \kappa_{2u1j}) \\ &\quad + g'(|\nabla u|)_j \cdot (\kappa_{1i} u_2 + \kappa_{1u2i} - \kappa_{2u1i}) \\ &\quad + g \cdot (\kappa_{1ij} u_2 + \kappa_{1i} u_{2j} + \kappa_{1j} u_{2i} + \kappa_{1u2ij} - \kappa_{2i} u_{1j} - \kappa_{2j} u_{1i} - \kappa_{2u1ij}); \end{aligned}$$

hence

$$\begin{aligned} (3-4) \quad \sum_{ij} a_{ij} \psi_{ij} &= u_2 g \sum_{ij} a_{ij} k_{1ij} + \left[2u_2 g' \sum_j a_{1j} (|\nabla u|)_j + 2g \sum_j a_{1j} u_{2j} \right] \cdot \kappa_{11} \\ &\quad + \left[2u_2 g' \sum_j a_{2j} (|\nabla u|)_j + 2g \sum_{2j} a_{2j} u_{2j} - 2g \sum_j a_{1j} u_{1j} \right] \cdot \kappa_{12} \\ &\quad + \left[-2g \sum_j a_{2j} u_{1j} \right] \cdot \kappa_{22} \\ &\quad + \left[u_2 g'' \sum_{ij} a_{ij} (|\nabla u|)_i (|\nabla u|)_j + u_2 g' \sum_{ij} a_{ij} (|\nabla u|)_{ij} \right. \\ &\quad \quad \left. + 2g' \sum_{ij} a_{ij} (|\nabla u|)_j u_{2i} + g \sum_{ij} a_{ij} u_{2ij} \right] \cdot \kappa_1 \\ &\quad + \left[-2g' \sum_{ij} a_{ij} (|\nabla u|)_j u_{1i} - g \sum_{ij} a_{ij} u_{1ij} \right] \cdot \kappa_2. \end{aligned}$$

Direct calculation yields

$$\begin{aligned} (3-5) \quad (|\nabla u|)_i &= \frac{1}{|\nabla u|} \sum_k u_k u_{ki}, \\ (|\nabla u|)_{ij} &= \frac{1}{|\nabla u|} \sum_k u_{kj} u_{ki} + \frac{1}{|\nabla u|} \sum_k u_k u_{kij} - \frac{1}{|\nabla u|^3} \sum_{kl} u_l u_{lj} u_k u_{ki}. \end{aligned}$$

Then at the point x_0 ,

$$\begin{aligned} (3-6) \quad (|\nabla u|)_i &= u_{2i}, \\ (|\nabla u|)_{ij} &= \frac{u_{1j} u_{1i}}{u_2} + u_{2ij}. \end{aligned}$$

By the commutation formulas (1-1)–(1-2), we obtain

$$\begin{aligned}
 (3-7) \quad \sum_{ij} a_{ij} u_{1ij} &= \sum_{ij} a_{ij} u_{i1j} \\
 &= \sum_{ij} a_{ij} \left[u_{ij1} + \sum_m u_m R_{mi1j} \right] \\
 &= - \sum_{ij} a_{ij,1} u_{ij} + \sum_{ijm} a_{ij} u_m R_{mi1j} \\
 &= -2u_1(u_{11}u_{22} - u_{12}^2) + u_1(1 + |\nabla u|^2) \cdot c
 \end{aligned}$$

and

$$\begin{aligned}
 (3-8) \quad \sum_{ij} a_{ij} u_{2ij} &= \sum_{ij} a_{ij} u_{i2j} \\
 &= \sum_{ij} a_{ij} \left[u_{ij2} + \sum_m u_m R_{mi2j} \right] \\
 &= - \sum_{ij} a_{ij,2} u_{ij} + \sum_{ijm} a_{ij} u_m R_{mi2j} \\
 &= -2u_2(u_{11}u_{22} - u_{12}^2) + u_2(1 + |\nabla u|^2) \cdot c.
 \end{aligned}$$

Hence at the point x_0 , we have

$$(3-9) \quad \sum_{ij} a_{ij} u_{1ij} = 0,$$

$$(3-10) \quad \sum_{ij} a_{ij} u_{2ij} = -2u_2(u_{11}u_{22} - u_{12}^2) + u_2(1 + u_2^2) \cdot c.$$

On the other hand,

$$(3-11) \quad a_{11} = 1 + u_2^2, \quad a_{12} = 0, \quad a_{21} = 0, \quad a_{22} = 1,$$

$$(3-12) \quad u_{22} = -(1 + u_2^2)u_{11}.$$

Inserting (3-6), (3-9)–(3-12) into (3-4), we obtain

$$\begin{aligned}
 (3-13) \quad \sum_{ij} a_{ij} \psi_{ij} &= u_2 g \sum_{ij} a_{ij} k_{1ij} + [2u_2(1 + u_2^2)g' + 2(1 + u_2^2)g]u_{12} \cdot \kappa_{11} \\
 &\quad + [-2u_2(1 + u_2^2)g' - 4(1 + u_2^2)g]u_{11} \cdot \kappa_{12} + [-2g]u_{12} \cdot \kappa_{22} \\
 &\quad + [u_2(1 + u_2^2)g'' + (3 + 4u_2^2)g' + 2u_2 g] \cdot [(1 + u_2^2)u_{11}^2 + u_{12}^2] \cdot \kappa_1 \\
 &\quad + [u_2^2(1 + u_2^2)g' + u_2(1 + u_2^2)g] \cdot \kappa_1 \cdot c.
 \end{aligned}$$

To compute the first term in (3-13), we should get the formula for $\sum_{ij} a_{ij} \kappa_{ij}$ at a general point in advance. Recalling the curvature formula for the level curves, we have

$$(3-14) \quad |\nabla u|^3 \cdot \kappa = 2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11}.$$

On both sides of (3-14), we take the second derivative with respect to ij , multiply by a_{ij} , and then sum with respect to ij . We obtain

$$(3-15) \quad \begin{aligned} \sum_{ij} a_{ij} \kappa_{ij} &= \frac{1}{|\nabla u|^3} \sum_{ij} a_{ij} (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11})_{ij} \\ &\quad - \frac{6}{|\nabla u|} \sum_{ij} a_{ij} (|\nabla u|)_j \cdot \kappa_i - \frac{6}{|\nabla u|^2} \sum_{ij} a_{ij} (|\nabla u|)_i (|\nabla u|)_j \cdot \kappa \\ &\quad - \frac{3}{|\nabla u|} \sum_{ij} a_{ij} (|\nabla u|)_{ij} \cdot \kappa. \end{aligned}$$

Recalling the minimal surface equation (3-1), we have

$$(3-16) \quad a_{11} = 1 + u_2^2, \quad a_{12} = -u_1 u_2, \quad a_{21} = -u_1 u_2, \quad a_{22} = 1 + u_1^2,$$

and

$$(3-17) \quad u_{22} = \frac{2u_1 u_2 u_{12} - (1 + u_2^2) u_{11}}{1 + u_1^2}.$$

Inserting (3-5) and (3-16)–(3-17) into (3-15), we get

$$(3-18) \quad \begin{aligned} \sum_{ij} a_{ij} \kappa_{ij} &= \frac{1}{|\nabla u|^3} \sum_{ij} a_{ij} (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11})_{ij} \\ &\quad - \frac{6(1 + |\nabla u|^2)}{|\nabla u|^2(1 + u_1^2)} [(1 + u_2^2)u_1 u_{11} + (1 - u_1^2)u_2 u_{12}] \cdot \kappa_1 \\ &\quad - \frac{6(1 + |\nabla u|^2)}{|\nabla u|^2} (u_1 u_{12} - u_2 u_{11}) \cdot \kappa_2 \\ &\quad - \frac{1}{|\nabla u|^2} \left\{ \frac{9 + 6|\nabla u|^2}{1 + u_1^2} [(1 + u_2^2)u_{11}^2 - 2u_1 u_2 u_{11} u_{12} + (1 + u_1^2)u_{12}^2] \right. \\ &\quad \left. + 3 \sum_{ijk} a_{ij} u_k u_{kij} \right\} \cdot \kappa. \end{aligned}$$

Now the commutation formulas (1-1)–(1-2) and relations (3-16)–(3-17) yield

$$\begin{aligned}
 (3-19) \quad \sum_{ijk} a_{ij} u_k u_{kij} &= \sum_{ijk} a_{ij} u_k u_{ijk} + \sum_{ijkm} a_{ij} u_k u_m R_{mikj} \\
 &= - \sum_{ijk} a_{ij,k} u_k u_{ij} + \sum_{ijkm} a_{ij} u_k u_m R_{mikj} \\
 &= \frac{2|\nabla u|^2}{1+u_1^2} [(1+u_2^2)u_{11}^2 - 2u_1 u_2 u_{11} u_{12} + (1+u_1^2)u_{12}^2] \\
 &\quad + |\nabla u|^2 (1 + |\nabla u|^2) \cdot c.
 \end{aligned}$$

Putting (3-19) into (3-18), we have

$$\begin{aligned}
 (3-20) \quad \sum_{ij} a_{ij} \kappa_{ij} &= \frac{1}{|\nabla u|^3} \sum_{ij} a_{ij} (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11})_{ij} \\
 &\quad - \frac{6(1+|\nabla u|^2)}{|\nabla u|^2(1+u_1^2)} [(1+u_2^2)u_{11}u_{11} + (1-u_1^2)u_2u_{12}] \cdot \kappa_1 \\
 &\quad - \frac{6(1+|\nabla u|^2)}{|\nabla u|^2} (u_1 u_{12} - u_2 u_{11}) \cdot \kappa_2 \\
 &\quad - \frac{9+12|\nabla u|^2}{|\nabla u|^2(1+u_1^2)} [(1+u_2^2)u_{11}^2 - 2u_1 u_2 u_{11} u_{12} + (1+u_1^2)u_{12}^2] \cdot \kappa \\
 &\quad - 3(1+|\nabla u|^2) \cdot \kappa \cdot c.
 \end{aligned}$$

Straightforward computation gives

$$\begin{aligned}
 (3-21) \quad \sum_{ij} a_{ij} (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11})_{ij} &= 2u_1 u_2 \sum_{ij} a_{ij} u_{12ij} - u_1^2 \sum_{ij} a_{ij} u_{22ij} - u_2^2 \sum_{ij} a_{ij} u_{11ij} \\
 &\quad + 2(u_2 u_{12} - u_1 u_{22}) \sum_{ij} a_{ij} u_{1ij} + 2(u_1 u_{12} - u_2 u_{11}) \sum_{ij} a_{ij} u_{2ij} \\
 &\quad - 4u_2 \sum_{ij} a_{ij} u_{2i} u_{11j} - 4u_1 \sum_{ij} a_{ij} u_{1i} u_{22j} + 4u_2 \sum_{ij} a_{ij} u_{1i} u_{12j} \\
 &\quad + 4u_1 \sum_{ij} a_{ij} u_{2i} u_{12j} + 4u_{12} \sum_{ij} a_{ij} u_{1i} u_{2j} \\
 &\quad - 2u_{22} \sum_{ij} a_{ij} u_{1i} u_{1j} - 2u_{11} \sum_{ij} a_{ij} u_{2i} u_{2j} \\
 &\triangleq J_1 + J_2 + J_3 + J_4,
 \end{aligned}$$

where

$$\begin{aligned} J_1 &= 2u_1u_2 \sum_{ij} a_{ij}u_{12ij} - u_1^2 \sum_{ij} a_{ij}u_{22ij} - u_2^2 \sum_{ij} a_{ij}u_{11ij}, \\ J_2 &= 2(u_2u_{12} - u_1u_{22}) \sum_{ij} a_{ij}u_{1ij} + 2(u_1u_{12} - u_2u_{11}) \sum_{ij} a_{ij}u_{2ij}, \\ J_3 &= -4u_2 \sum_{ij} a_{ij}u_{2i}u_{11j} - 4u_1 \sum_{ij} a_{ij}u_{1i}u_{22j} \\ &\quad + 4u_2 \sum_{ij} a_{ij}u_{1i}u_{12j} + 4u_1 \sum_{ij} a_{ij}u_{2i}u_{12j}, \\ J_4 &= 4u_{12} \sum_{ij} a_{ij}u_{1i}u_{2j} - 2u_{22} \sum_{ij} a_{ij}u_{1i}u_{1j} - 2u_{11} \sum_{ij} a_{ij}u_{2i}u_{2j}. \end{aligned}$$

We deal with the terms J_1 , J_2 , J_3 and J_4 consecutively. If we differentiate the minimal surface equation (3-1) twice, then we have

$$(3-22) \quad \sum_{ij} a_{ij,k}u_{ij} + \sum_{ij} a_{ij}u_{ijk} = 0,$$

$$(3-23) \quad \sum_{ij} a_{ij,kl}u_{ij} + \sum_{ij} a_{ij,k}u_{ijl} + \sum_{ij} a_{ij,l}u_{ijk} + \sum_{ij} a_{ij}u_{ijkl} = 0.$$

By the commutation formulas (1-1)–(1-3), we have

$$u_{klij} = u_{ijkl} + \sum_m u_{mi} R_{mklj} + \sum_m u_{mj} R_{mkli} + \sum_m u_{mk} R_{milj} + \sum_m u_{ml} R_{mikj}.$$

It follows that

$$\begin{aligned} (3-24) \quad \sum_{ij} a_{ij}u_{klij} &= - \left[\sum_{ij} a_{ij,kl}u_{ij} + \sum_{ij} a_{ij,k}u_{ijl} + \sum_{ij} a_{ij,l}u_{ijk} \right] \\ &\quad + \sum_{ijm} a_{ij}u_{mi} R_{mklj} + \sum_{ijm} a_{ij}u_{mj} R_{mkli} \\ &\quad + \sum_{ijm} a_{ij}u_{mk} R_{milj} + \sum_{ijm} a_{ij}u_{ml} R_{mikj}. \end{aligned}$$

Note that $a_{ij} = (1 + |\nabla u|^2)\delta_{ij} - u_i u_j$. It is easy to get

$$a_{ij,k} = 2 \sum_m u_m u_{mk} \delta_{ij} - u_{ik} u_j - u_i u_{jk},$$

$$a_{ij,l} = 2 \sum_m u_m u_{ml} \delta_{ij} - u_{il} u_j - u_i u_{jl},$$

$$a_{ij,kl} = 2 \sum_m u_m u_{mk} \delta_{ij} + 2 \sum_m u_m u_{mkl} \delta_{ij} - u_{ikl} u_j - u_{ik} u_{jl} - u_{il} u_{jk} - u_i u_{jkl}.$$

Thus we have

$$(3-25) \quad \sum_{ij} a_{ij,kl} u_{ij} = 2(u_1 u_{22} - u_2 u_{12}) u_{1kl} + 2(u_2 u_{11} - u_1 u_{12}) u_{2kl} \\ + 2u_{1l} u_{1k} u_{22} + 2u_{2l} u_{2k} u_{11} - 2u_{1k} u_{2l} u_{12} - 2u_{2k} u_{1l} u_{12},$$

$$(3-26) \quad \sum_{ij} a_{ij,k} u_{ijl} = 2u_2 u_{2k} u_{11l} + 2u_1 u_{1k} u_{22l} - 2(u_1 u_{2k} + u_2 u_{1k}) u_{12l},$$

$$(3-27) \quad \sum_{ij} a_{ij,l} u_{ijk} = 2u_2 u_{2l} u_{11k} + 2u_1 u_{1l} u_{22k} - 2(u_1 u_{2l} + u_2 u_{1l}) u_{12k}.$$

Putting (3-25)–(3-27) into (3-24), we obtain

$$(3-28) \quad \sum_{ij} a_{ij} u_{klij} = -[2(u_1 u_{22} - u_2 u_{12}) u_{1kl} + 2(u_2 u_{11} - u_1 u_{12}) u_{2kl} + 2u_{1l} u_{1k} u_{22} \\ + 2u_{2l} u_{2k} u_{11} - 2u_{1k} u_{2l} u_{12} - 2u_{2k} u_{1l} u_{12} + 2u_2 u_{2k} u_{11l} \\ + 2u_1 u_{1k} u_{22l} - 2(u_1 u_{2k} + u_2 u_{1k}) u_{12l} + 2u_2 u_{2l} u_{11k} \\ + 2u_1 u_{1l} u_{22k} - 2(u_1 u_{2l} + u_2 u_{1l}) u_{12k}] \\ + \sum_{ijm} a_{ij} u_{mi} R_{mklj} + \sum_{ijm} a_{ij} u_{mj} R_{mklj} \\ + \sum_{ijm} a_{ij} u_{mk} R_{milj} + \sum_{ijm} a_{ij} u_{ml} R_{mikj}.$$

By the commutation formulas (1-1)–(1-2),

$$(3-29) \quad u_{121} = u_{112} + \sum_m u_m R_{m121} = u_{112} + u_2 \cdot c,$$

$$(3-30) \quad u_{122} = u_{221} + \sum_m u_m R_{m212} = u_{221} + u_1 \cdot c.$$

With (3-16) and (3-29)–(3-30) in hand, formula (3-28) is equivalent to

$$\sum_{ij} a_{ij} u_{12ij} = -2u_2 u_{22} u_{111} + 2u_2 u_{12} u_{112} + 2u_1 u_{12} u_{221} \\ - 2u_1 u_{11} u_{222} - 2u_{12} (u_{11} u_{22} - u_{12}^2) \\ + [u_1 u_2 u_{11} + (4 + 5u_1^2 + 5u_2^2) u_{12} + u_1 u_2 u_{22}] \cdot c,$$

$$\sum_{ij} a_{ij} u_{22ij} = -4u_2 u_{22} u_{112} + 2(3u_2 u_{12} + u_1 u_{22}) u_{221} \\ - 2(u_2 u_{11} + u_1 u_{12}) u_{222} - 2u_{22} (u_{11} u_{22} - u_{12}^2) \\ + [-2(1 + u_2^2) u_{11} + 10u_1 u_2 u_{12} + 2(1 + u_1^2 + u_2^2) u_{22}] \cdot c,$$

$$\sum_{ij} a_{ij} u_{11ij} = -2(u_2 u_{12} + u_1 u_{22}) u_{111} + 2(u_2 u_{11} + 3u_1 u_{12}) u_{112} \\ - 4u_1 u_{11} u_{221} - 2u_{11} (u_{11} u_{22} - u_{12}^2) \\ + [2(1 + u_1^2 + u_2^2) u_{11} + 10u_1 u_2 u_{12} - 2(1 + u_1^2) u_{22}] \cdot c.$$

Therefore,

(3-31)

$$\begin{aligned} J_1 = & [2u_2^3u_{12} - 2u_1u_2^2u_{22}] \cdot u_{111} + [-2u_2^3u_{11} - 2u_1u_2^2u_{12} + 4u_1^2u_2u_{22}] \cdot u_{112} \\ & + [4u_1u_2^2u_{11} - 2u_1^2u_2u_{12} - 2u_1^3u_{22}] \cdot u_{221} + [-2u_1^2u_2u_{11} + 2u_1^3u_{12}] \cdot u_{222} \\ & - 2(u_{11}u_{22} - u_{12}^2) \cdot (2u_1u_2u_{12} - u_1^2u_{22} - u_2^2u_{11}) \\ & + [2(u_1^2 - u_2^2 + u_1^2u_2^2 - u_2^4)u_{11} + 8u_1u_2u_{12} + 2(-u_1^2 + u_2^2 + u_1^2u_2^2 - u_1^4)u_{22}] \cdot c. \end{aligned}$$

Let us handle the term J_2 . By (1-1)–(1-2), (3-16) and (3-22), we have

$$\begin{aligned} \sum_{ij} a_{ij}u_{1ij} &= \sum_{ij} a_{ij}u_{ij1} + \sum_{ijm} a_{ij}u_m R_{mi1j} \\ &= -\sum_{ij} a_{ij,1}u_{ij} + \sum_{ijm} a_{ij}u_m R_{mi1j} \\ &= -2u_1(u_{11}u_{22} - u_{12}^2) + u_1(1 + u_1^2 + u_2^2) \cdot c \end{aligned}$$

and

$$\begin{aligned} \sum_{ij} a_{ij}u_{2ij} &= \sum_{ij} a_{ij}u_{ij2} + \sum_{ijm} a_{ij}u_m R_{mi2j} \\ &= -\sum_{ij} a_{ij,2}u_{ij} + \sum_{ijm} a_{ij}u_m R_{mi2j} \\ &= -2u_2(u_{11}u_{22} - u_{12}^2) + u_2(1 + u_1^2 + u_2^2) \cdot c. \end{aligned}$$

Thus

$$(3-32) \quad \begin{aligned} J_2 = & -4(u_{11}u_{22} - u_{12}^2) \cdot (2u_1u_2u_{12} - u_1^2u_{22} - u_2^2u_{11}) \\ & + 2(1 + u_1^2 + u_2^2) \cdot (2u_1u_2u_{12} - u_1^2u_{22} - u_2^2u_{11}) \cdot c. \end{aligned}$$

For the term J_3 , by (3-16) and (3-29)–(3-30), we have

$$\begin{aligned} (3-33) \quad J_3 = & [(-4u_2 - 4u_2^3)u_{12} + 4u_1u_2^2u_{22}] \cdot u_{111} \\ & + [(4u_2 + 4u_2^3)u_{11} + (4u_1 + 4u_1u_2^2)u_{12} + (-4u_2 - 8u_1^2u_2)u_{22}] \cdot u_{112} \\ & + [(-4u_1 - 8u_1u_2^2)u_{11} + (4u_2 + 4u_1^2u_2)u_{12} + (4u_1 + 4u_1^3)u_{22}] \cdot u_{221} \\ & + [4u_1^2u_2u_{11} + (-4u_1 - 4u_1^3)u_{12}] \cdot u_{222} \\ & + [(4u_2^2 - 4u_1^2u_2^2 + 4u_2^4)u_{11} + 8u_1u_2u_{12} + (4u_1^2 - 4u_1^2u_2^2 + 4u_1^4)u_{22}] \cdot c. \end{aligned}$$

Moreover, straightforward computation yields

$$(3-34) \quad J_4 = -2(u_{11}u_{22} - u_{12}^2)[(1 + u_2^2)u_{11} - 2u_1u_2u_{12} + (1 + u_1^2)u_{22}] = 0.$$

Combining (3-21) and (3-31)–(3-34), we obtain

$$\begin{aligned}
 (3-35) \quad & \sum_{ij} a_{ij} (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11})_{ij} \\
 &= [(-4u_2 - 2u_2^3)u_{12} + 2u_1 u_2^2 u_{22}] \cdot u_{111} \\
 &\quad + [(4u_2 + 2u_2^3)u_{11} + (4u_1 + 2u_1 u_2^2)u_{12} + (-4u_2 - 4u_1^2 u_2)u_{22}] \cdot u_{112} \\
 &\quad + [(-4u_1 - 4u_1 u_2^2)u_{11} + (4u_2 + 2u_1^2 u_2)u_{12} + (4u_1 + 2u_1^3)u_{22}] \cdot u_{221} \\
 &\quad + [2u_1^2 u_2 u_{11} + (-4u_1 - 2u_1^3)u_{12}] \cdot u_{222} \\
 &\quad - 6(u_{11} u_{22} - u_{12}^2) \cdot (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11}) \\
 &\quad + [(2u_1^2 - 4u_1^2 u_2^2)u_{11} + (20u_1 u_2 + 4u_1 u_2^3 + 4u_1^3 u_2)u_{12} \\
 &\quad \quad \quad + (2u_2^2 - 4u_1^2 u_2^2)u_{22}] \cdot c.
 \end{aligned}$$

Now let us explore the relations between $u_{111}, u_{112}, u_{221}, u_{222}$ and κ_1, κ_2 . If we take the first derivative on both sides of (3-14) and (3-1), respectively, then using (3-5), (3-29)–(3-30) and (3-16), we obtain

$$\begin{aligned}
 & -u_2^2 \cdot u_{111} + 2u_1 u_2 u_{12} \cdot u_{112} - u_1^2 \cdot u_{221} - |\nabla u|^3 \cdot \kappa_1 \\
 & \quad - 3|\nabla u|(u_1 u_{11} + u_2 u_{12}) \cdot \kappa - 2u_1(u_{11} u_{22} - u_{12}^2) + 2u_1 u_2^2 \cdot c = 0, \\
 & -u_2^2 \cdot u_{112} + 2u_1 u_2 u_{12} \cdot u_{221} - u_1^2 \cdot u_{222} - |\nabla u|^3 \cdot \kappa_2 \\
 & \quad - 3|\nabla u|(u_1 u_{12} + u_2 u_{22}) \cdot \kappa - 2u_2(u_{11} u_{22} - u_{12}^2) + 2u_1^2 u_2 \cdot c = 0,
 \end{aligned}$$

and

$$\begin{aligned}
 & (1 + u_2^2) \cdot u_{111} - 2u_1 u_2 \cdot u_{112} + (1 + u_1^2) \cdot u_{221} + 2u_1(u_{11} u_{22} - u_{12}^2) - 2u_1 u_2^2 \cdot c = 0, \\
 & (1 + u_2^2) \cdot u_{112} - 2u_1 u_2 \cdot u_{221} + (1 + u_1^2) \cdot u_{222} + 2u_2(u_{11} u_{22} - u_{12}^2) - 2u_1^2 u_2 \cdot c = 0.
 \end{aligned}$$

Thus we have

$$\begin{aligned}
 (3-36) \quad u_{111} = & (-u_2^2 + u_1^2 + 3u_1^2 u_2^2 + u_1^4) |\nabla u|^{-1} \cdot \kappa_1 + (-2u_1 u_2 - 2u_1^3 u_2) |\nabla u|^{-1} \cdot \kappa_2 \\
 & + 3|\nabla u|^{-3} [(-u_1 u_2^2 + u_1^3 + 3u_1^3 u_2^2 + u_1^5) u_{11} \\
 & \quad + (-u_2^3 - u_1^2 u_2 + 3u_1^2 u_2^3 - u_1^4 u_2) u_{12} \\
 & \quad \quad \quad + (-2u_1 u_2^2 - 2u_1^3 u_2^2) u_{22}] \cdot \kappa \\
 & + (-6u_1 u_2^2 + 2u_1^3) |\nabla u|^{-4} \cdot (u_{11} u_{22} - u_{12}^2) + 2u_1 u_2^2 |\nabla u|^{-2} \cdot c,
 \end{aligned}$$

$$\begin{aligned}
 (3-37) \quad u_{112} = & (2u_1 u_2 + 2u_1 u_2^3) |\nabla u|^{-1} \cdot \kappa_1 + (-u_2^2 + u_1^2 - u_1^2 u_2^2 + u_1^4) |\nabla u|^{-1} \cdot \kappa_2 \\
 & + 3|\nabla u|^{-3} [(2u_1^2 u_2 + 2u_1^2 u_2^3) u_{11} \\
 & \quad + (u_1 u_2^2 + u_1^3 + 2u_1 u_2^4 - u_1^3 u_2^2 + u_1^5) u_{12} \\
 & \quad \quad \quad + (-u_2^3 + u_1^2 u_2 - u_1^2 u_2^3 + u_1^4 u_2) u_{22}] \cdot \kappa \\
 & + (-2u_2^3 + 6u_1^2 u_2) |\nabla u|^{-4} \cdot (u_{11} u_{22} - u_{12}^2) - 2u_1^2 u_2 |\nabla u|^{-2} \cdot c,
 \end{aligned}$$

and

$$(3-38) \quad u_{221} = (u_2^2 + u_2^4 - u_1^2 - u_1^2 u_2^2) |\nabla u|^{-1} \cdot \kappa_1 + (2u_1 u_2 + 2u_1^3 u_2) |\nabla u|^{-1} \cdot \kappa_2 \\ + 3|\nabla u|^{-3} [(u_1 u_2^2 + u_1 u_2^4 - u_1^3 - u_1^3 u_2^2) u_{11} \\ + (u_2^3 + u_2^5 + u_1^2 u_2 - u_1^2 u_2^3 + 2u_1^4 u_2) u_{12} \\ + (2u_1 u_2^2 + 2u_1^3 u_2^2) u_{22}] \cdot \kappa \\ + (6u_1 u_2^2 - 2u_1^3) |\nabla u|^{-4} \cdot (u_{11} u_{22} - u_{12}^2) - 2u_1 u_2^2 |\nabla u|^{-2} \cdot c,$$

$$(3-39) \quad u_{222} = (-2u_1 u_2 - 2u_1 u_2^3) |\nabla u|^{-1} \cdot \kappa_1 + (u_2^2 + u_2^4 - u_1^2 + 3u_1^2 u_2^2) |\nabla u|^{-1} \cdot \kappa_2 \\ + 3|\nabla u|^{-3} [(-2u_1^2 u_2 - 2u_1^2 u_2^3) u_{11} \\ + (-u_1 u_2^2 - u_1 u_2^4 - u_1^3 + 3u_1^3 u_2^2) u_{12} \\ + (u_2^3 + u_2^5 - u_1^2 u_2 + 3u_1^2 u_2^3) u_{22}] \cdot \kappa \\ + (2u_2^3 - 6u_1^2 u_2) |\nabla u|^{-4} \cdot (u_{11} u_{22} - u_{12}^2) + 2u_1^2 u_2 |\nabla u|^{-2} \cdot c.$$

Inserting (3-36)–(3-39) and (3-17) into (3-35), after some tedious calculation, we get

$$(3-40) \quad \sum_{ij} a_{ij} (2u_1 u_2 u_{12} - u_1^2 u_{22} - u_2^2 u_{11})_{ij} \\ = \frac{2|\nabla u|(4+3|\nabla u|^2)}{1+u_1^2} [(1+u_2^2)u_1 u_{11} + (1-u_1^2)u_2 u_{12}] \cdot \kappa_1 \\ + 2|\nabla u|(4+3|\nabla u|^2)(u_1 u_{12} - u_2 u_{11}) \cdot \kappa_2 \\ + \frac{4|\nabla u|(2+3|\nabla u|^2)}{1+u_1^2} [(1+u_2^2)u_{11}^2 - 2u_1 u_2 u_{11} u_{12} + (1+u_1^2)u_{12}^2] \cdot \kappa \\ + 2|\nabla u|^3(1+|\nabla u|^2) \cdot \kappa \cdot c.$$

By (3-20) and (3-40), we have

$$(3-41) \quad \sum_{ij} a_{ij} \kappa_{ij} = \frac{2}{|\nabla u|^2(1+u_1^2)} [(1+u_2^2)u_1 u_{11} + (1-u_1^2)u_2 u_{12}] \cdot \kappa_1 \\ + \frac{2}{|\nabla u|^2} (u_1 u_{12} - u_2 u_{11}) \cdot \kappa_2 \\ - \frac{1}{|\nabla u|^2(1+u_1^2)} [(1+u_2^2)u_{11}^2 - 2u_1 u_2 u_{11} u_{12} + (1+u_1^2)u_{12}^2] \cdot \kappa \\ - (1+|\nabla u|^2)\kappa \cdot c.$$

Then at the point x_0 , we take the first derivative of (3-41). Note that

$$\kappa(x_0) = -\frac{u_{11}}{u_2}.$$

With (3-6), (3-16) and (3-29) in hand, we obtain

$$(3-42) \quad \sum_{ij} a_{ij} \kappa_{ij1} = \frac{2(1-u_2^2)}{u_2} u_{12} \cdot \kappa_{11} - \frac{2(1-u_2^2)}{u_2} u_{11} \cdot \kappa_{12} \\ + \left[\frac{2}{u_2} u_{112} + \frac{1+u_2^2}{u_2^2} u_{11}^2 - \frac{3}{u_2^2} u_{12}^2 \right] \cdot \kappa_1 \\ + \left[-\frac{2}{u_2} u_{111} + \frac{4}{u_2^2} u_{11} u_{12} \right] \cdot \kappa_2 + \frac{2(1+u_2^2)}{u_2^3} u_{11}^2 u_{111} \\ + \frac{2}{u_2^3} u_{11} u_{12} u_{112} - \frac{2(1+u_2^2)}{u_2^4} u_{11}^3 u_{12} - \frac{2}{u_2^4} u_{11} u_{12}^3 \\ + \left[(1-u_2^2) \cdot \kappa_1 + \frac{2(1+u_2^2)}{u_2^2} u_{11} u_{12} \right] \cdot c.$$

Now, the equations (3-36) and (3-37) are simplified as

$$(3-43) \quad u_{111} = -u_2 \kappa_1 + \frac{3}{u_2} u_{11} u_{12},$$

$$(3-44) \quad u_{112} = -u_2 \kappa_2 - \frac{1+u_2^2}{u_2} u_{11}^2 + \frac{2}{u_2} u_{12}^2.$$

Putting (3-43)–(3-44) into (3-42), one obtains

$$\sum_{ij} a_{ij} \kappa_{ij1} = \frac{2(1-u_2^2)}{u_2} u_{12} \cdot \kappa_{11} - \frac{2(1-u_2^2)}{u_2} u_{11} \cdot \kappa_{12} + \left[-\frac{3(1+u_2^2)}{u_2^2} u_{11}^2 + \frac{1}{u_2^2} u_{12}^2 \right] \cdot \kappa_1 \\ - \frac{4}{u_2^2} u_{11} u_{12} \cdot \kappa_2 + \frac{2(1+u_2^2)}{u_2^4} u_{11}^3 u_{12} + \frac{2}{u_2^4} u_{11} u_{12}^3 \\ + \left[(1-u_2^2) \cdot \kappa_1 + \frac{2(1+u_2^2)}{u_2^2} u_{11} u_{12} \right] \cdot c.$$

Therefore, by commutation formulas (1-1)–(1-2), we get

$$(3-45) \quad \sum_{ij} a_{ij} \kappa_{1ij} = \sum_{ij} a_{ij} \left[\kappa_{ij1} + \sum_m \kappa_m R_{mi1j} \right] = \sum_{ij} a_{ij} \kappa_{ij1} + \kappa_1 \cdot c \\ = \frac{2(1-u_2^2)}{u_2} u_{12} \cdot \kappa_{11} - \frac{2(1-u_2^2)}{u_2} u_{11} \cdot \kappa_{12} \\ + \left[-\frac{3(1+u_2^2)}{u_2^2} u_{11}^2 + \frac{1}{u_2^2} u_{12}^2 \right] \cdot \kappa_1 - \frac{4}{u_2^2} u_{11} u_{12} \cdot \kappa_2 + \frac{2(1+u_2^2)}{u_2^4} u_{11}^3 u_{12} \\ + \frac{2}{u_2^4} u_{11} u_{12}^3 + \left[(2-u_2^2) \cdot \kappa_1 + \frac{2(1+u_2^2)}{u_2^2} u_{11} u_{12} \right] \cdot c.$$

Thanks to (3-45), the formula (3-13) reduces to

$$\begin{aligned}
 (3-46) \quad & \sum_{ij} a_{ij} \psi_{ij} \\
 &= [2u_2(1+u_2^2)g' + 4g] u_{12} \cdot \kappa_{11} \\
 &\quad - [2u_2(1+u_2^2)g' + (6+2u_2^2)g] u_{11} \cdot \kappa_{12} - 2gu_{12} \cdot \kappa_{22} \\
 &\quad + \left\{ \left[u_2(1+u_2^2)^2 g'' + (3+4u_2^2)(1+u_2^2)g' - \frac{(3-2u_2^2)(1+u_2^2)}{u_2}g \right] u_{11}^2 \right. \\
 &\quad \quad \left. + \left[u_2(1+u_2^2)g'' + (3+4u_2^2)g' + \frac{1+2u_2^2}{u_2}g \right] u_{12}^2 \right\} \cdot \kappa_1 \\
 &\quad - \frac{4g}{u_2} u_{11} u_{12} \cdot \kappa_2 + \frac{2g}{u_2^3} [(1+u_2^2)u_{11}^2 + u_{12}^2] u_{11} u_{12} \\
 &\quad + \left\{ [u_2^2(1+u_2^2)g' + 3u_2g] \cdot \kappa_1 + \frac{2(1+u_2^2)g}{u_2} u_{11} u_{12} \right\} \cdot c.
 \end{aligned}$$

By (3-3) and (3-41), we have

$$(3-47) \quad \kappa_{11} = \frac{1}{u_2 g} [\psi_1 - (u_2 g' + g) u_{12} \cdot \kappa_1 + g u_{11} \cdot \kappa_2],$$

$$(3-48) \quad \kappa_{12} = \frac{1}{u_2 g} [\psi_2 + (u_2 g' + g)(1+u_2^2) u_{11} \cdot \kappa_1 + g u_{12} \cdot \kappa_2],$$

and

$$\begin{aligned}
 (3-49) \quad \kappa_{22} = \frac{1}{u_2 g} \left\{ -(1+u_2^2) \psi_1 + [u_2(1+u_2^2)g' + (3+u_2^2)g] u_{12} \cdot \kappa_1 \right. \\
 \quad \quad \quad \left. - (3+u_2^2)g u_{11} \cdot \kappa_2 + \frac{g}{u_2^2} u_{11} [(1+u_2^2)u_{11}^2 + u_{12}^2] \right. \\
 \quad \quad \quad \left. + (1+u_2^2)g u_{11} \cdot c \right\}.
 \end{aligned}$$

Putting (3-47)–(3-49) into (3-46), we finally get

$$\begin{aligned}
 (3-50) \quad & \sum_{ij} a_{ij} \psi_{ij} \\
 &= \left[\frac{2(1+u_2^2)g'}{g} + \frac{6+2u_2^2}{u_2} \right] \cdot (u_{12} \psi_1 - u_{11} \psi_2) \\
 &\quad + \left[u_2(1+u_2^2)g'' - \frac{2u_2(1+u_2^2)g'^2}{g} - 5g' - \frac{9g}{u_2} \right] \cdot [(1+u_2^2)u_{11}^2 + u_{12}^2] \cdot \kappa_1 \\
 &\quad + [u_2^2(1+u_2^2)g' + 3u_2g] \cdot \kappa_1 \cdot c.
 \end{aligned}$$

If we let

$$g(t) = \frac{(1+t^2)^{3/2}}{t^3},$$

then the last two terms on the right-hand side of (3-50) vanish. Namely,

$$\sum_{ij} a_{ij} \psi_{ij} = 2u_2 \cdot (u_{12}\psi_1 - u_{11}\psi_2).$$

This completes the proof of [Theorem 1.2](#). □

Acknowledgement

The authors would like to thank Professor Xi-nan Ma for suggesting this problem to them and for some helpful discussions.

References

- [Ahlfors 1973] L. V. Ahlfors, *Conformal invariants: topics in geometric function theory*, McGraw-Hill, New York, 1973. [MR](#) [Zbl](#)
- [Bianchini et al. 2009] C. Bianchini, M. Longinetti, and P. Salani, “Quasiconcave solutions to elliptic problems in convex rings”, *Indiana Univ. Math. J.* **58**:4 (2009), 1565–1589. [MR](#) [Zbl](#)
- [Caffarelli and Spruck 1982] L. A. Caffarelli and J. Spruck, “Convexity properties of solutions to some classical variational problems”, *Comm. Partial Differential Equations* **7**:11 (1982), 1337–1379. [MR](#) [Zbl](#)
- [Chang et al. 2010] S.-Y. A. Chang, X.-N. Ma, and P. Yang, “Principal curvature estimates for the convex level sets of semilinear elliptic equations”, *Discrete Contin. Dyn. Syst.* **28**:3 (2010), 1151–1164. [MR](#) [Zbl](#)
- [Chern et al. 1999] S. S. Chern, W. H. Chen, and K. S. Lam, *Lectures on differential geometry*, Series on University Mathematics **1**, World Scientific, River Edge, NJ, 1999. [MR](#) [Zbl](#)
- [Gabriel 1957] R. M. Gabriel, “A result concerning convex level surfaces of 3-dimensional harmonic functions”, *J. London Math. Soc.* **32** (1957), 286–294. [MR](#) [Zbl](#)
- [Gergen 1931] J. J. Gergen, “Note on the green function of a star-shaped three dimensional region”, *Amer. J. Math.* **53**:4 (1931), 746–752. [MR](#) [Zbl](#)
- [Kawohl 1985] B. Kawohl, *Rearrangements and convexity of level sets in PDE*, Lecture Notes in Mathematics **1150**, Springer, 1985. [MR](#) [Zbl](#)
- [Kong and Xu 2015] S. Kong and J. Xu, “A remark on the level sets of the graph of harmonic functions bounded by two circles in parallel planes”, *J. Partial Differ. Equ.* **28**:3 (2015), 197–207. [MR](#) [Zbl](#)
- [Lewis 1977] J. L. Lewis, “Capacitary functions in convex rings”, *Arch. Rational Mech. Anal.* **66**:3 (1977), 201–224. [MR](#) [Zbl](#)
- [Longinetti 1983] M. Longinetti, “Convexity of the level lines of harmonic functions”, *Boll. Un. Mat. Ital. A* (6) **2**:1 (1983), 71–75. In Italian. [MR](#) [Zbl](#)
- [Ma and Zhang 2013] X.-N. Ma and W. Zhang, “The concavity of the Gaussian curvature of the convex level sets of p -harmonic functions with respect to the height”, *Commun. Math. Stat.* **1**:4 (2013), 465–489. [MR](#) [Zbl](#)

- [Ma and Zhang 2014] X.-N. Ma and Y. Zhang, “The convexity and the Gaussian curvature estimates for the level sets of harmonic functions on convex rings in space forms”, *J. Geom. Anal.* **24**:1 (2014), 337–374. [MR](#) [Zbl](#)
- [Ma et al. 2010] X.-N. Ma, Q. Ou, and W. Zhang, “Gaussian curvature estimates for the convex level sets of p -harmonic functions”, *Comm. Pure Appl. Math.* **63**:7 (2010), 935–971. [MR](#) [Zbl](#)
- [Ortel and Schneider 1983] M. Ortel and W. Schneider, “Curvature of level curves of harmonic functions”, *Canad. Math. Bull.* **26**:4 (1983), 399–405. [MR](#) [Zbl](#)
- [Shiffman 1956] M. Shiffman, “On surfaces of stationary area bounded by two circles, or convex curves, in parallel planes”, *Ann. of Math.* (2) **63** (1956), 77–90. [MR](#) [Zbl](#)
- [Talenti 1983] G. Talenti, “On functions, whose lines of steepest descent bend proportionally to level lines”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **10**:4 (1983), 587–605. [MR](#) [Zbl](#)
- [Wang and Zhang 2012] P. Wang and W. Zhang, “Gaussian curvature estimates for the convex level sets of solutions for some nonlinear elliptic partial differential equations”, *J. Partial Differ. Equ.* **25**:3 (2012), 239–275. [MR](#) [Zbl](#)
- [Zhang and Zhang 2013] T. Zhang and W. Zhang, “On convexity of level sets of p -harmonic functions”, *J. Differential Equations* **255**:7 (2013), 2065–2081. [MR](#) [Zbl](#)

Received May 25, 2016. Revised September 24, 2016.

JINJU XU
DEPARTMENT OF MATHEMATICS
SHANGHAI NORMAL UNIVERSITY
SHANGHAI 200234
CHINA
jjxujane@shu.edu.cn

WEI ZHANG
SCHOOL OF MATHEMATICS & STATISTICS
LANZHOU UNIVERSITY
GANSU 730000
CHINA
zhangw@lzu.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)

Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 288 No. 1 May 2017

<i>C¹</i> -umbilics with arbitrarily high indices	1
NAOYA ANDO, TOSHIFUMI FUJIYAMA and MASAAKI UMEHARA	
Well-posedness of second-order degenerate differential equations with finite delay in vector-valued function spaces	27
SHANQUAN BU and GANG CAI	
On cusp solutions to a prescribed mean curvature equation	47
ALEXANDRA K. ECHART and KIRK E. LANCASTER	
Radial limits of capillary surfaces at corners	55
MOZHGAN (NORA) ENTEKHABI and KIRK E. LANCASTER	
A new bicommutant theorem	69
ILIJAS FARAH	
Noncompact manifolds that are inward tame	87
CRAIG R. GUILBAULT and FREDERICK C. TINSLEY	
<i>p</i> -adic variation of unit root <i>L</i> -functions	129
C. DOUGLAS HAESSIG and STEVEN SPERBER	
Bavard's duality theorem on conjugation-invariant norms	157
MORIMICHI KAWASAKI	
Parabolic minimal surfaces in $\mathbb{M}^2 \times \mathbb{R}$	171
VANDERSON LIMA	
Regularity conditions for suitable weak solutions of the Navier–Stokes system from its rotation form	189
CHANGXING MIAO and YANQING WANG	
Geometric properties of level curves of harmonic functions and minimal graphs in 2-dimensional space forms	217
JINJU XU and WEI ZHANG	
Eigenvalue resolution of self-adjoint matrices	241
XUWEN ZHU	