

*Pacific
Journal of
Mathematics*

**ASYMPTOTIC ORDER-OF-VANISHING FUNCTIONS
ON THE PSEUDOEFFECTIVE CONE**

SHIN-YAO JOW

Volume 288 No. 2

June 2017

ASYMPTOTIC ORDER-OF-VANISHING FUNCTIONS ON THE PSEUDOEFFECTIVE CONE

SHIN-YAO JOW

Let v be a discrete valuation on the function field of a normal projective variety X . Ein, Lazarsfeld, Mustařă, Nakamaye, and Popa showed that v induces a nonnegative real-valued continuous function on the big cone of X , which they called the asymptotic order of vanishing along v . The case where v is given by the order of vanishing along a prime divisor was studied earlier by Nakayama, who extended the domain of the function to the pseudoeffective cone and investigated the continuity of the extended function.

Here we generalize Nakayama's results to any discrete valuation v , using an approach inspired by Lazarsfeld and Mustařă's construction of the global Okounkov body, which has a quite different flavor from the arguments employed by Nakayama.

A corollary is that the asymptotic order-of-vanishing function can be extended continuously to the pseudoeffective cone $\text{PE}(X)$ of X if $\text{PE}(X)$ is polyhedral (note that we do *not* require $\text{PE}(X)$ to be *rational polyhedral*).

Let X be a normal projective variety over an algebraically closed field k , and let $K(X)$ be the function field of X . Let v be a discrete valuation of $K(X)$ over k , and let Z be the center of v on X . Ein, Lazarsfeld, Mustařă, Nakamaye, and Popa gave the following definitions:

Definition 1 [Ein et al. 2006]. Let D be an effective big Cartier divisor on X . We establish the following notation:

- (i) $v(D) = v(f)$, where f is a local equation of D at the generic point of Z .
- (ii) $v(|D|) = \min\{v(D') : D' \in |D|\} = v(D')$ for general $D' \in |D|$.
- (iii) $v(\|D\|) = \lim_{m \rightarrow \infty} v(|mD|)/m$. This is called the asymptotic order of vanishing of D along v .

MSC2010: 14C20.

Keywords: asymptotic order of vanishing, pseudoeffective cone, global Okounkov body, Nakayama's σ -decomposition.

By taking m to be sufficiently divisible, the definition of $v(\|D\|)$ can also be extended to big \mathbb{Q} -divisors D . It is proved in [Ein et al. 2006, Theorem A] that $v(\|D\|)$ depends only on the numerical equivalence class of D , so it induces a function on the set of numerical equivalence classes of big \mathbb{Q} -divisors. Moreover, this function extends uniquely to a continuous function on the cone $\text{Big}(X)$ of numerical equivalence classes of big \mathbb{R} -divisors. In view of this result, it is natural to ask whether this function can be extended continuously to the pseudoeffective cone $\text{PE}(X)$, the closure of the big cone $\text{Big}(X)$ in the Néron–Severi space $N^1(X)_{\mathbb{R}}$.

The case where v is a divisorial valuation was investigated earlier by Nakayama [2004] during his study of Zariski decomposition in higher dimensions. More precisely, let Γ be a prime divisor on a smooth projective variety X , and let v be the discrete valuation of $K(X)$ given by the order of vanishing at the generic point of Γ . Nakayama used the notation $\sigma_{\Gamma}(D)$ to denote the asymptotic order of vanishing $v(\|D\|)$ of a big divisor class $D \in \text{Big}(X)$. If $D \in \text{PE}(X)$ is a pseudoeffective class, he defined $\sigma_{\Gamma}(D)$ by picking an arbitrary ample class $A \in N^1(X)_{\mathbb{R}}$ and setting $\sigma_{\Gamma}(D)$ to be the limit

$$\sigma_{\Gamma}(D) = \lim_{\epsilon \rightarrow 0^+} \sigma_{\Gamma}(D + \epsilon A),$$

after establishing that this limit does not depend on the choice of A , in [Nakayama 2004, III.1.5]. In III.1.7 of the same work, Nakayama showed that the function $\sigma_{\Gamma} : \text{PE}(X) \rightarrow \mathbb{R}_{\geq 0}$ is lower semicontinuous, and he gave an example where it is not continuous in IV.2.8. It is interesting to note that in his example $\text{PE}(X)$ is not polyhedral. The goal of this short note is to generalize Nakayama’s results to any discrete valuation v of $K(X)/k$, using an approach inspired by Lazarsfeld and Mustață’s construction [2009] of the global Okounkov body, which has a quite different flavor from the arguments employed by Nakayama. In addition, we will see that the function $v(\|\cdot\|) : \text{Big}(X) \rightarrow \mathbb{R}_{\geq 0}$ can be extended continuously to $\text{PE}(X)$ if $\text{PE}(X)$ is polyhedral.

Theorem 2. *Let X be a normal projective variety over an algebraically closed field k , and let v be a discrete valuation of $K(X)$ over k . If $D \in \text{PE}(X)$ is a pseudoeffective class, then for any ample class $A \in N^1(X)_{\mathbb{R}}$, $\lim_{\epsilon \rightarrow 0^+} v(\|D + \epsilon A\|)$ does not depend on the choice of A . Moreover, if we denote this limit by $\sigma_v(D)$, then the function*

$$\sigma_v : \text{PE}(X) \rightarrow \mathbb{R}_{\geq 0} \cup \{+\infty\}$$

is lower semicontinuous, and is continuous at every point where $\text{PE}(X)$ is locally polyhedral.

A subset S of \mathbb{R}^n is said to be *locally polyhedral* at a point $x \in S$ if there exist a polytope $P \subset \mathbb{R}^n$ and an open subset U of \mathbb{R}^n containing x such that $U \cap S = U \cap P$. It follows from Theorem 2 that the function $v(\|\cdot\|) : \text{Big}(X) \rightarrow \mathbb{R}_{\geq 0}$ can be extended

continuously to $\text{PE}(X)$ if $\text{PE}(X)$ is polyhedral, which is the case, for example, when the Picard number of X is 2. Note that we do *not* require $\text{PE}(X)$ to be *rational* polyhedral (cf. [Ein et al. 2006, Theorem D]).

Remark 3. If v is divisorial, the limit $\lim_{\epsilon \rightarrow 0^+} v(\|D + \epsilon A\|)$ in Theorem 2 is finite [Nakayama 2004, III.1.5]. We do not know if this is true for all discrete valuations v , which is why we include $+\infty$ in the target of σ_v . In case the value of σ_v is $+\infty$ at a point of $\text{PE}(X)$, the (semi)continuity of σ_v should be interpreted with respect to the usual order topology on $\mathbb{R}_{\geq 0} \cup \{+\infty\}$.

Let us introduce some notions from convex analysis which will be useful in the proof of Theorem 2. Let $f : S \rightarrow \mathbb{R} \cup \{+\infty\}$ be a function on a convex subset S of \mathbb{R}^n . We say that f is *convex* if

$$f(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda f(x_1) + (1 - \lambda)f(x_2)$$

for all $x_1, x_2 \in S$ and all $0 \leq \lambda \leq 1$. The *epigraph* of f is the set

$$\{(x, y) \in S \times \mathbb{R} : y \geq f(x)\}.$$

A convex function f is said to be *closed* if its epigraph is a closed subset of \mathbb{R}^{n+1} . It is not difficult to show that if f is a closed convex function, then f is lower semicontinuous.

Proof of Theorem 2. As mentioned earlier, our approach is inspired by the construction of the global Okounkov body due to [Lazarsfeld and Mustařă 2009]. The strategy is to construct the epigraph of the asymptotic order-of-vanishing function as the closed convex cone spanned by a certain lattice semigroup. To see how this works for one big divisor D , let \mathbb{N} denote the set of nonnegative integers, and let

$$S(D) = \{(m, y) \in \mathbb{N}^2 : y \geq v(|mD|)\},$$

which is a subsemigroup of \mathbb{N}^2 . Let $C(D) = \overline{\text{cone}}(S(D))$ be the closed convex cone spanned by $S(D)$ in \mathbb{R}^2 . Then $C(D)$ is the epigraph of the function $x \mapsto v(\|x D\|)$. In order to get the epigraph of the function $v(\|\cdot\|) : \text{Big}(X) \rightarrow \mathbb{R}_{\geq 0}$, pick a \mathbb{Z} -basis D_1, \dots, D_n for $N^1(X)$ such that, after identifying $N^1(X)_{\mathbb{R}}$ with \mathbb{R}^n by this basis, we have $\text{PE}(X) \subseteq \mathbb{R}_{\geq 0}^n$. Let

$$S(X) = \{(m_1, \dots, m_n, y) \in \mathbb{N}^n \times \mathbb{N} : y \geq v(|m_1 D_1 + \dots + m_n D_n|)\},$$

and let

$$C(X) = \overline{\text{cone}}(S(X)) \subseteq \mathbb{R}_{\geq 0}^n \times \mathbb{R}_{\geq 0}$$

be the closed convex cone spanned by $S(X)$. Let

$$f : \text{PE}(X) \rightarrow \mathbb{R}_{\geq 0} \cup \{+\infty\}$$

be the function whose epigraph is $C(X)$. Then f is a closed convex function since $C(X)$ is a closed convex cone. Moreover, on the big cone $\text{Big}(X)$, f coincides with the function $v(\|\cdot\|)$ by [Lazarsfeld and Mustață 2009, Proposition 4.9].

To see what $f(D)$ is if D is on the boundary of $\text{PE}(X)$, we invoke a theorem of Gale, Klee, and Rockafellar, which states that a closed convex function is continuous at every point where its domain is locally polyhedral ([Gale et al. 1968, Theorem 2]; see also the introduction of [Ernst 2013]). It follows that for any ample $A \in N^1(X)_{\mathbb{R}}$, the restriction of f to the half-line $D + \mathbb{R}_{\geq 0}A$ is continuous. Hence

$$f(D) = \lim_{\epsilon \rightarrow 0^+} f(D + \epsilon A) = \lim_{\epsilon \rightarrow 0^+} v(\|D + \epsilon A\|).$$

This shows that the limit on the right does not depend on the choice of A , and that in fact $f = \sigma_v$. Since σ_v is a closed convex function, it is lower semicontinuous, and is continuous at every point where $\text{PE}(X)$ is locally polyhedral by the theorem of Gale, Klee and Rockafellar. \square

Acknowledgments

The author gratefully acknowledges the support of MOST (Ministry of Science and Technology, Taiwan), NCTS (National Center for Theoretical Sciences), and the Kenda Foundation.

References

- [Ein et al. 2006] L. Ein, R. Lazarsfeld, M. Mustață, M. Nakamaye, and M. Popa, “Asymptotic invariants of base loci”, *Ann. Inst. Fourier (Grenoble)* **56**:6 (2006), 1701–1734. MR Zbl
- [Ernst 2013] E. Ernst, “A converse of the Gale–Klee–Rockafellar theorem: continuity of convex functions at the boundary of their domains”, *Proc. Amer. Math. Soc.* **141**:10 (2013), 3665–3672. MR Zbl
- [Gale et al. 1968] D. Gale, V. Klee, and R. T. Rockafellar, “Convex functions on convex polytopes”, *Proc. Amer. Math. Soc.* **19** (1968), 867–873. MR Zbl
- [Lazarsfeld and Mustață 2009] R. Lazarsfeld and M. Mustață, “Convex bodies associated to linear series”, *Ann. Sci. Éc. Norm. Supér. (4)* **42**:5 (2009), 783–835. MR Zbl
- [Nakayama 2004] N. Nakayama, *Zariski-decomposition and abundance*, MSJ Memoirs **14**, Mathematical Society of Japan, Tokyo, 2004. MR Zbl

Received May 13, 2016.

SHIN-YAO JOW
 DEPARTMENT OF MATHEMATICS
 NATIONAL TSING HUA UNIVERSITY
 NO. 101, SEC. 2, GUANGFU RD
 HSINCHU 300
 TAIWAN
 syjow@math.nthu.edu.tw

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 288 No. 2 June 2017

Order on the homology groups of Smale spaces	257
MASSOUD AMINI, IAN F. PUTNAM and SARAH SAEIDI GHOLIKANDI	
Characterizations of immersed gradient almost Ricci solitons	289
CÍCERO P. AQUINO, HENRIQUE F. DE LIMA and JOSÉ N. V. GOMES	
Weighted Sobolev regularity of the Bergman projection on the Hartogs triangle	307
LIWEI CHEN	
Knots of tunnel number one and meridional tori	319
MARIO EUDAVE-MUÑOZ and GRISEL SANTIAGO-GONZÁLEZ	
On bisectional nonpositively curved compact Kähler–Einstein surfaces	343
DANIEL GUAN	
Effective lower bounds for $L(1, \chi)$ via Eisenstein series	355
PETER HUMPHRIES	
Asymptotic order-of-vanishing functions on the pseudoeffective cone	377
SHIN-YAO JOW	
Augmentations and rulings of Legendrian links in $\#^k(S^1 \times S^2)$	381
CAITLIN LEVERSON	
The Faber–Krahn inequality for the first eigenvalue of the fractional Dirichlet p -Laplacian for triangles and quadrilaterals	425
FRANCO OLIVARES CONTADOR	
Topological invariance of quantum quaternion spheres	435
BIPUL SAURABH	
Gap theorems for complete λ -hypersurfaces	453
HUIJUAN WANG, HONGWEI XU and ENTAO ZHAO	
Bach-flat h -almost gradient Ricci solitons	475
GABJIN YUN, JINSEOK CO and SEUNGSU HWANG	
A sharp height estimate for the spacelike constant mean curvature graph in the Lorentz–Minkowski space	489
JINGYONG ZHU	



0030-8730(201706)288:2;1-R