

*Pacific
Journal of
Mathematics*

**CRITICALITY OF THE AXIALLY SYMMETRIC
NAVIER–STOKES EQUATIONS**

ZHEN LEI AND QI S. ZHANG

CRITICALITY OF THE AXIALLY SYMMETRIC NAVIER–STOKES EQUATIONS

ZHEN LEI AND QI S. ZHANG

Smooth solutions to the axisymmetric Navier–Stokes equations obey the following maximum principle:

$$\sup_{t \geq 0} \|rv^\theta(t, \cdot)\|_{L^\infty} \leq \|rv^\theta(0, \cdot)\|_{L^\infty}.$$

We prove that all solutions with initial data in $H^{1/2}$ are smooth globally in time if rv^θ satisfies a kind of form boundedness condition (FBC) which is invariant under the natural scaling of the Navier–Stokes equations. In particular, if rv^θ satisfies

$$\sup_{t \geq 0} |rv^\theta(t, r, z)| \leq C_* |\ln r|^{-2}, \quad \text{where } r \leq \delta_0 \in (0, \frac{1}{2}), \quad C_* < \infty,$$

then our FBC is satisfied. Here δ_0 and C_* are independent of neither the profile nor the norm of the initial data. So the gap from regularity is logarithmic in nature. We also prove the global regularity of solutions if $\|rv^\theta(0, \cdot)\|_{L^\infty}$ or $\sup_{t \geq 0} \|rv^\theta(t, \cdot)\|_{L^\infty(r \leq r_0)}$ is small but the smallness depends on a certain dimensionless quantity of the initial data.

1. Introduction

The global regularity problem of three-dimensional incompressible Navier–Stokes equations is commonly considered as supercritical because the a priori estimates based on energy equality become worse when looking into finer and finer scales; see, for instance, [Tao 2007]. Such a “supercriticality” barrier is one of the main reasons why this is such a hard problem.

Recently, the axisymmetric Navier–Stokes equations have attracted tremendous interest from experts. See, for instance, [Burke Loftus and Zhang 2010; Chae and Lee 2002; Chen et al. 2008; 2009; 2015; Hou and Li 2008; Hou et al. 2008; Jiu and Xin 2003; Koch et al. 2009; Lei et al. 2013; Lei and Zhang 2011b; 2011a; Leonardi et al. 1999; Neustupa and Pokorný 2000; 2001; Pan 2016; Seregin and Šverák 2009; Tian and Xin 1998; Zhang and Zhang 2014]. These results heavily depend on the maximum principle of the dimensionless quantity $\Gamma = rv^\theta$, which

MSC2010: 35K55.

Keywords: axially symmetric Navier–Stokes equations, regularity condition, criticality.

makes the axisymmetric Navier–Stokes equations partially critical (only the swirl component v^θ of the velocity field satisfies a dimensionless a priori estimate). Although the axially symmetric Navier–Stokes equation is a special case of the full three-dimensional one, our level of understanding had been roughly the same, with the previously mentioned difficulty unresolved because the effective a priori bound available is still the energy estimate, which has positive dimension $\frac{1}{2}$.

The aim of this article is to show that the axisymmetric Navier–Stokes equation is, in fact, fully critical. More precisely, we prove all solutions with initial data in $H^{1/2}$ are smooth globally in time if rv^θ satisfies a kind of FBC which is invariant under the natural scaling of the Navier–Stokes equations. In particular, if rv^θ satisfies $\sup_{t \geq 0} |rv^\theta(t, r, z)| \leq C_* |\ln r|^{-2}$, where $r \leq \delta_0 \in (0, \frac{1}{2})$, $C_* < \infty$, then our FBC is satisfied. Here δ_0 and C_* are independent of neither the profile nor the norm of the initial data. The proof is based on the observation that the vorticity equations can be transformed into a system such that the vortex-stretching terms are critical. This means that the potentials in front of unknown functions scale as $1/|x|^2$. For example, in (1-8) below, the function J is regarded as unknown and the potential in front of it is $-2v^\theta/r$ which scales as $1/|x|^2$.

We also prove the global regularity of solutions if $\sup_{t \geq 0} \|rv^\theta(t, \cdot)\|_{L^\infty(r \leq r_0)}$ or $\|rv^\theta(0, \cdot)\|_{L^\infty}$ is small but the smallness depends on a certain dimensionless quantity of the initial data. Our work is inspired by the recent interesting result of Chen, Fang and Zhang in [Chen et al. 2015] where, among other things, global regularity is obtained if $rv^\theta(t, \cdot, z)$ is Hölder continuous in the r variable.

To state our result more precisely, let us recall that in cylindrical coordinates r, θ, z with $(x_1, x_2, x_3) = (r \cos \theta, r \sin \theta, z)$, axially symmetric solutions of the Navier–Stokes equations are of the following form:

$$\begin{cases} v(t, x) = v^r(t, r, z)e_r + v^\theta(t, r, z)e_\theta + v^z(t, r, z)e_z, \\ p(t, x) = p(t, r, z). \end{cases}$$

The components v^r, v^θ, v^z are all independent of the angle of rotation θ . Here e_r, e_θ, e_z are the basis vectors for \mathbb{R}^3 given by

$$e_r = \left(\frac{x_1}{r}, \frac{x_2}{r}, 0 \right)^\top, \quad e_\theta = \left(\frac{-x_2}{r}, \frac{x_1}{r}, 0 \right)^\top, \quad e_z = (0, 0, 1)^\top.$$

In terms of (v^r, v^θ, v^z, p) , the axisymmetric Navier–Stokes equations are

$$(1-1) \quad \begin{cases} \partial_t v^r + (v^r e_r + v^z e_z) \cdot \nabla v^r - \frac{(v^\theta)^2}{r} + \partial_r p = \left(\Delta - \frac{1}{r^2} \right) v^r, \\ \partial_t v^\theta + (v^r e_r + v^z e_z) \cdot \nabla v^\theta + \frac{v^r v^\theta}{r} = \left(\Delta - \frac{1}{r^2} \right) v^\theta, \\ \partial_t v^z + (v^r e_r + v^z e_z) \cdot \nabla v^z + \partial_z p = \Delta v^z, \\ \partial_r v^r + \frac{v^r}{r} + \partial_z v^z = 0. \end{cases}$$

It is well known that finite energy smooth solutions of the Navier–Stokes equations satisfy the following energy identity

$$(1-2) \quad \|v(t, \cdot)\|_{L^2}^2 + 2 \int_0^t \|\nabla v(s, \cdot)\|_{L^2}^2 ds = \|v_0\|_{L^2}^2, \quad \forall t \geq 0.$$

Set

$$\Gamma = rv^\theta.$$

One can easily check that

$$(1-3) \quad \partial_t \Gamma + (v^r e_r + v^z e_z) \cdot \nabla \Gamma = \left(\Delta - \frac{2}{r} \partial_r \right) \Gamma.$$

A significant consequence of (1-3) is that smooth solutions of the axisymmetric Navier–Stokes equations satisfy the following maximum principle; see, for instance, [Chae and Lee 2002; Hou and Li 2008; Chen et al. 2008; Neustupa and Pokorný 2000; 2001]:

$$(1-4) \quad \sup_t \|\Gamma(t, \cdot)\|_{L^\infty} \leq \|\Gamma_0\|_{L^\infty}.$$

We emphasize that $\|\Gamma(t, \cdot)\|_{L^\infty}$ is a dimensionless quantity with respect to the natural scaling of the Navier–Stokes equations. From this point of view, the axisymmetric Navier–Stokes equations can be seen as partially critical, while the general Navier–Stokes equations are known to be supercritical; see [Tao 2007].

Now let us introduce the function class where v^θ lives. It is defined in an integral way which is usually called the form boundedness condition (FBC), which is similar to a condition that a certain Hardy-type inequality holds.

Definition 1.1. We say that the angular velocity $v^\theta(t, r, z)$ is in a (δ_*, C_*) -critical class if

$$(1-5) \quad \int \frac{|v^\theta|}{r} |f|^2 dx \leq C_* \int |\partial_r f|^2 dx + C_0 \int_{r \geq r_0} |f|^2 dx,$$

$$(1-6) \quad \int |v^\theta|^2 |f|^2 dx \leq \delta_* \int |\partial_r f|^2 dx + C_0 \int_{r \geq r_0} |f|^2 dx,$$

hold for some $r_0 > 0$, some $C_0 > 0$ and for all $t \geq 0$ and all axisymmetric scalar and vector functions $f \in H^1$.

Clearly, under the natural scaling of the Navier–Stokes equations, namely,

$$v^\lambda(t, x) = \lambda v(\lambda^2 t, \lambda x), \quad p^\lambda(t, x) = \lambda^2 p(\lambda^2 t, \lambda x),$$

the above definition of FBC is invariant: $(v^\lambda)^\theta$ satisfies (1-5)–(1-6) if v^θ also does.

We now state the first result of this article.

Theorem 1.2. *For any $C_* > 1$, there exists a constant $\delta_* > 0$ depending on C_* such that the following conclusion holds for all local strong solutions to the axially symmetric Navier–Stokes equations with initial data $\|v_0\|_{H^{1/2}} < \infty$ and $\|\Gamma_0\|_{L^\infty} < \infty$. If the angular velocity field v^θ is in the (δ_*, C_*) -critical class, i.e., v^θ satisfies the critical form boundedness condition in (1-5)–(1-6), then v is regular globally in time.*

An important corollary of [Theorem 1.2](#) is this:

Corollary 1.3. *Let $\delta_0 \in (0, \frac{1}{2})$ and $C_1 > 1$. Let v be the local strong solution of the axially symmetric Navier–Stokes equations with initial data $v_0 \in H^{1/2}$ and $\|\Gamma_0\|_{L^\infty} < \infty$. If*

$$(1-7) \quad \sup_{0 \leq t < T} |\Gamma(t, r, z)| \leq C_1 |\ln r|^{-2}, \quad r \leq \delta_0,$$

then v is regular globally in time.

We emphasize that C_* in [Theorem 1.2](#) and C_1 in [Corollary 1.3](#) are independent of neither the profile nor the norm of the given initial data. The proof of this corollary will be given at the end of [Section 2](#). The point is that if (1-7) is satisfied, then the FBC of (1-5)–(1-6) is true. Then one can apply [Theorem 1.2](#) to get the desired conclusion.

Our work is inspired by a recent very interesting work by Chen, Fang and Zhang [\[2015\]](#) where, among other things, the authors proved that v is regular if Γ is Hölder continuous. Let

$$\Omega = \frac{\omega^\theta}{r}, \quad J = -\frac{\partial_z v^\theta}{r}.$$

We emphasize that J was introduced in [\[Chen et al. 2015\]](#), while Ω appeared much earlier and can be at least tracked back to the book of Majda and Bertozzi [\[2002\]](#). Both of the two new variables are of great importance in our work. Following [\[Majda and Bertozzi 2002; Hou and Li 2008; Chen et al. 2015\]](#), we also study the equations for J and Ω :

$$(1-8) \quad \begin{cases} \partial_t J + (b \cdot \nabla) J = \left(\Delta + \frac{2}{r} \partial_r \right) J + (\omega^r \partial_r + \omega^z \partial_z) \frac{v^r}{r}, \\ \partial_t \Omega + (b \cdot \nabla) \Omega = \left(\Delta + \frac{2}{r} \partial_r \right) \Omega - 2 \frac{v^\theta}{r} J. \end{cases}$$

Here ω^θ is the angular component of the vorticity $\omega = \nabla \times v$, which reads

$$\omega(t, x) = \omega^r e_r + \omega^\theta e_\theta + \omega^z e_z,$$

with

$$\omega^r = -\partial_z v^\theta, \quad \omega^\theta = \partial_z v^r - \partial_r v^z, \quad \omega^z = \partial_r v_\theta + \frac{v^\theta}{r}.$$

Our new observation is that the axisymmetric Navier–Stokes equations exhibit certain critical nature when being formulated in terms of a new set of unknowns,

J and Ω . Our second observation is that, with the FBC assumptions (1-5)–(1-6), the stretching term $(\omega^r \partial_r + \omega^z \partial_z)v^r/r$ in the equation for J could be arbitrarily small, by using the relation of v^r , v^z and Ω in Lemma 2.1 (which was originally proved by Hou, Lei and Li [Hou et al. 2008] in the periodic case and later on extended to the general case by Lei [2015]. Alternatively, one may also use the magic formula given by Miao and Zheng [2013] to prove it). Then we can derive a closed a priori estimate for J and Ω using the first two observations and the structure of the stretching term in the equation for Ω .

Our second goal is to prove that the smallness of $\sup_{t \geq 0} \|\Gamma(t, \cdot)\|_{L^\infty(r \leq r_0)}$ or $\|\Gamma_0\|_{L^\infty}$ implies the global regularity of the solutions. Recently, Chen, Fang and Zhang [2015] proved that, among many other interesting results, if $\Gamma(t, \cdot, z)$ is Hölder continuous in the r variable, then the solution of the axisymmetric Navier–Stokes equations is smooth. Both of the results depend on given initial data. More precisely, the smallness of $\sup_{t \geq 0} \|\Gamma(t, \cdot)\|_{L^\infty(r \leq r_0)}$ or $\|\Gamma_0\|_{L^\infty}$ in our Theorem 1.4 depends on other dimensionless norms of the initial data. From this point of view, our result improves the one in [Chen et al. 2015].

We define

$$V = \frac{v^\theta}{\sqrt{r}}.$$

Here is the second main result:

Theorem 1.4. *Let $r_0 > 0$. Suppose that $v_0 \in H^{1/2}$ such that $\Omega_0 \in L^2$, $V_0^2 \in L^2$ and $\Gamma_0 \in L^2 \cap L^\infty$. Denote*

$$(\|\Omega_0\|_{L^2} + \|V_0^2\|_{L^2})\|\Gamma_0\|_{L^2} = M_0$$

and

$$(\|V_0^2\|_{L^2} + \|\Omega_0\|_{L^2} + r_0^{-2}\|v_0\|_{L^2}\|\Gamma_0\|_{L^\infty}^{3/2})\|\Gamma_0\|_{L^2} = M_1.$$

There exists an absolute positive (small) constant $\delta > 0$ such that if either

$$\|\Gamma_0\|_{L^\infty} \leq \delta M_0^{-1},$$

or

$$\sup_{t \geq 0} \|\Gamma(t, \cdot)\|_{L^\infty(r \leq r_0)} \leq \delta M_1^{-1},$$

then the axially symmetric Navier–Stokes equations are globally well-posed.

The proof of Theorem 1.4 is based on a new formulation of the axisymmetric Navier–Stokes equations (1-1) in terms of V and $\Omega = \omega^\theta/r$, and also on the estimate of v^r/r in terms of Ω and its derivative (see Lemma 2.1).

Now let us recall some highlights on the study of the axisymmetric Navier–Stokes equations. It has been known since the late 1960s (see [Ladyzhenskaya 1968; Ukhovskii and Iudovich 1968]) that if the swirl $v_\theta = 0$, then finite energy solutions to (1-1) are smooth for all time. See also [Leonardi et al. 1999], by

Leonardi, Málek, Nečas and Pokorný. In the presence of swirl, it is not known in general if finite energy solutions blow up or not in finite time. Hou and Li [2008] constructed a family of large solutions based on some deep insights on a one-dimensional model. See also some extended results in [Hou et al. 2008] by Hou, Lei and Li. We also mention various a priori estimates of smooth solutions by Chae and Lee [2002] and Burke Loftus and Zhang [2010]. To the best of our knowledge, the best a priori bound of the velocity field is given in [Lei et al. 2013]:

$$|v(t, x)| \leq C_* r^{-2} |\ln r|^{1/2}.$$

In [Chen et al. 2008], Chen, Strain, Tsai and Yau obtained a lower bound for the possible blow up rate of singularities: if

$$|v(t, x)| \leq \frac{C_*}{r},$$

then v is regular. This seems to be the first time that people have been able to exclude possible singularities in the presence of assumptions on $|x|^{-1}$ -type nonsmallness quantities. Soon afterward, Chen, Strain, Yau and Tsai [Chen et al. 2009] and Koch, Nadirashvili, Seregin and Šverák [Koch et al. 2009] extended the result of [Chen et al. 2008] and in particular, excluded the possibility of type I singularities of v . See also a local version by Seregin and Šverák [2009] and various extensions by Pan [2016]. We also mention that Lei and Zhang [2011b] excluded the possibility of singularities under

$$v^r e_r + v^z e_z \in L^\infty([0, T], \text{BMO}^{-1})$$

based on an observation in [Lei and Zhang 2011a]. This solves the regularity problem of $L^\infty([0, T], \text{BMO}^{-1})$ solutions of Navier–Stokes equations in the axisymmetric case. Moreover, it extends the result of [Chen et al. 2008] and [Koch et al. 2009] since the assumptions on the axial component of velocity $|v^z| \leq C_* r^{-1}$ itself imply $v^r e_r + v^z e_z \in L^\infty([0, T], \text{BMO}^{-1})$ (see [Lei et al. 2013] for details).

Let us also mention that Neustupa and Pokorný [2000] proved that the regularity of one component (either v^r or v^θ) implies regularity of the other components of the solution. The work of Jiu and Xin [2003] also proves regularity under an assumption of sufficiently small zero-dimension scaled norms. See more refined results in [Neustupa and Pokorný 2001] and the work of Ping Zhang and Ting Zhang [Zhang and Zhang 2014]. Chae and Lee [2002] also proved regularity results assuming finiteness of another certain zero-dimensional integral. Tian and Xin [1998] constructed a family of singular axially symmetric solutions with singular initial data.

The remainder of the paper is simply organized as follows. In Section 2 we recall two basic lemmas and prove Corollary 1.3 by assuming the validity of Theorem 1.2. In Section 3 we prove Theorem 1.2. Section 4 is devoted to the proof of Theorem 1.4.

Remark. This paper was posted on the arXiv in May 2015. In August 2015, Dongyi Wei [2016] improved one of our regularity conditions by a factor of $\sqrt{|\ln r|}$.

2. Notations and lemmas

For abbreviation, we denote

$$b(t, x) = v^r e_r + v^z e_z.$$

The last equation in (1-1) shows that b is divergence-free. The Laplacian operator Δ and the gradient operator ∇ in the cylindrical coordinate are

$$\Delta = \partial_r^2 + \frac{1}{r} \partial_r + \frac{1}{r^2} \partial_\theta^2 + \partial_z^2, \quad \nabla = e_r \partial_r + \frac{e_\theta}{r} \partial_\theta + e_z \partial_z.$$

For the scalar axisymmetric function $f(t, r, z)$, we often use the commutation property:

$$\nabla \partial_r f(r, z) = \partial_r \nabla f(r, z).$$

Throughout the proof, we will denote

$$\|f\|_{L^2}^2 = \int |f|^2 r \, dr \, dz, \quad dx = r \, dr \, dz.$$

The estimate in Lemma 2.1 will be used often. It was originally proved in [Hou et al. 2008] in the periodic case and then extended to the general case in [Lei 2015] (see (4.5)–(4.6) there, noting the relation $v^r = -\partial_z \psi^\theta$). Alternatively, one may also use the magic formula given by Miao and Zheng [2013] to prove it.

Lemma 2.1. *Let v^r be the radial component of the velocity field and $\Omega = \omega^\theta/r$. Then there exists an absolute positive constant $K_0 > 0$ such that*

$$\left\| \nabla \frac{v^r}{r} \right\|_{L^2} \leq K_0 \|\Omega\|_{L^2}, \quad \left\| \nabla^2 \frac{v^r}{r} \right\|_{L^2} \leq K_0 \|\partial_z \Omega\|_{L^2}.$$

Lemma 2.2 gives the uniform decay estimate for the angular component of vorticity in the r direction for large r . We point out that a weaker estimate for ω^θ has appeared in [Chae and Lee 2002]. Even though we don't need to use the estimate for ω^r and ω^z in this paper, we will include them below for possible future use.

Lemma 2.2. *Suppose that $v_0 \in L^2$ is an axially symmetric divergence-free vector and $(r\omega_0^r, r^2\omega_0^\theta, r\omega_0^z) \in L^2$. Then the smooth solution of the Navier–Stokes equation with initial data v_0 satisfies the following a priori estimates:*

$$\begin{aligned} (2-1) \quad & \sup_{0 \leq t < T} (\|r\omega^r(t, \cdot)\|_{L^2}^2, \|r\omega^z(t, \cdot)\|_{L^2}^2) \\ & + \int_0^T (\|\nabla[r\omega^r(t, \cdot)]\|_{L^2}^2, \|\nabla[r\omega^z(t, \cdot)]\|_{L^2}^2) \, dt \\ & \leq \|r\omega_0^r\|_{L^2}^2 + \|r\omega_0^z\|_{L^2}^2 + 4(\|\Gamma_0\|_{L^\infty}^2 + 1) \|v_0\|_{L^2}^2, \end{aligned}$$

and

$$(2-2) \quad \sup_{0 \leq t < T} \|r^2 \omega^\theta(t, \cdot)\|_{L^2}^2 + \int_0^T \|\nabla(r^2 \omega^\theta)\|_{L^2}^2 dt \\ \leq C_0 \left(\|r^2 \omega_0^\theta\|_{L^2}^2 + (\|v_0\|_{L^2}^4 + \|\Gamma_0\|_{L^3}^2) \|v_0\|_{L^2}^2 \right) \exp \left\{ \frac{T}{\|\Gamma_0\|_{L^3}^2 + \|v_0\|_{L^2}^4} \right\},$$

where C_0 is a generic positive constant.

Proof. First of all, let us recall that

$$(2-3) \quad \begin{cases} \partial_t \omega^r + b \cdot \nabla \omega^r - \partial_r v^r \omega^r = \left(\Delta - \frac{1}{r^2} \right) \omega^r + \partial_z v^r \omega^z, \\ \partial_t \omega^\theta + b \cdot \nabla \omega^\theta - \frac{v^r}{r} \omega^\theta = \left(\Delta - \frac{1}{r^2} \right) \omega^\theta + \partial_z \frac{(v^\theta)^2}{r}, \\ \partial_t \omega^z + b \cdot \nabla \omega^z - \partial_z v^z \omega^z = \Delta \omega^z + \partial_r v^z \omega^r. \end{cases}$$

Let us first prove (2-1). Taking the L^2 inner product of the first equation of (2-3) with $r^2 \omega^r$, and of the third equation with $r^2 \omega^z$, we have

$$\frac{1}{2} \frac{d}{dt} \int (\omega^r)^2 r^2 dx - \int r^2 \omega^r \left(\Delta - \frac{1}{r^2} \right) \omega^r dx \\ = - \int r^2 \omega^r b \cdot \nabla \omega^r dx + \int r^2 \partial_r v^r (\omega^r)^2 dx + \int r^2 \omega^r \partial_z v^r \omega^z dx$$

and

$$\frac{1}{2} \frac{d}{dt} \int (\omega^z)^2 r^2 dx - \int r^2 \omega^z \Delta \omega^z dx \\ = - \int r^2 \omega^z b \cdot \nabla \omega^z dx + \int r^2 \partial_z v^z (\omega^z)^2 dx + \int r^2 \omega^z \partial_r v^z \omega^r dx.$$

Using integration by parts, we have

$$- \int r^2 \omega^r \left(\Delta - \frac{1}{r^2} \right) \omega^r dx = \int |\nabla(r \omega^r)|^2 dx$$

and

$$- \int r^2 \omega^z \Delta \omega^z dx = \int |\nabla(r \omega^z)|^2 dx - 2 \int |\omega^z|^2 dx.$$

Using integration by parts and the incompressibility constraint, one has

$$- \int r^2 \omega^r b \cdot \nabla \omega^r dx - \int r^2 \omega^z b \cdot \nabla \omega^z dx + \int r^2 \partial_r v^r (\omega^r)^2 dx + \int r^2 \partial_z v^z (\omega^z)^2 dx \\ = \int (r v^r + r^2 \partial_r v^r) (\omega^r)^2 + (r v^r + r^2 \partial_z v^z) (\omega^z)^2 dx \\ = - \int [\partial_z v^z (r \omega^r)^2 + \partial_r v^r (r \omega^z)^2] dx.$$

Consequently, we have

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int [(\omega^r)^2 + (\omega^z)^2] r^2 dx + \int (|\nabla(r\omega^r)|^2 + |\nabla(r\omega^z)|^2) dx \\ &= \int |\omega^z|^2 dx - \int [\partial_z v^z (r\omega^r)^2 + \partial_r v^r (r\omega^z)^2] dx + \int (r^2 \omega^r \partial_z v^r \omega^z + r^2 \omega^z \partial_r v^z \omega^r) dx \\ &\leq 2 \int |\omega^z|^2 dx + \|\nabla b\|_{L^2} (\|r\omega^r\|_{L^4}^2 + \|r\omega^z\|_{L^4}^2). \end{aligned}$$

Note that by Gagliardo–Nirenberg’s inequality and the maximum principle $\|\Gamma\|_{L^\infty} \leq \|\Gamma_0\|_{L^\infty}$, one has

$$\begin{aligned} \|r\omega^r\|_{L^4}^2 + \|r\omega^z\|_{L^4}^2 &= \|\nabla\Gamma\|_{L^4}^2 \\ &= \left(\int -\Gamma \nabla \cdot (\nabla\Gamma |\nabla\Gamma|^2) dx \right)^{1/2} \leq 3\|\Gamma\|_{L^\infty} \|\Delta\Gamma\|_{L^2} \\ &\leq 3\|\Gamma_0\|_{L^\infty} (\|\partial_r(r\omega^z)\|_{L^2} + \|\partial_z(r\omega^r)\|_{L^2} + \|\omega^z\|_{L^2}). \end{aligned}$$

Hence, by Hölder’s inequality, we have

$$\frac{d}{dt} \int [(\omega^r)^2 + (\omega^z)^2] r^2 dx + \int (|\nabla(r\omega^r)|^2 + |\nabla(r\omega^z)|^2) dx \leq 4(\|\Gamma_0\|_{L^\infty}^2 + 1) \|\nabla b\|_{L^2}^2.$$

Integrating the above differential inequality with respect to time and recalling the basic energy estimate, one gets (2-1).

Next, let us prove (2-2). Let us first write the second equation of (2-3) as:

$$\partial_t(r^2\omega^\theta) + b \cdot \nabla(r^2\omega^\theta) - 3rv^r\omega^\theta = \frac{\partial_z \Gamma^2}{r} + \Delta(r^2\omega^\theta) - \frac{4}{r} \partial_r(r^2\omega^\theta) + 3\omega^\theta.$$

The standard energy estimate gives that

$$\frac{1}{2} \frac{d}{dt} \|r^2\omega^\theta\|_{L^2}^2 + \|\nabla(r^2\omega^\theta)\|_{L^2}^2 = 3 \int rv^r\omega^\theta r^2\omega^\theta dx + \int \partial_z \Gamma^2 r\omega^\theta dx + 3 \int \omega^\theta r^2\omega^\theta dx.$$

It is easy to estimate that

$$\begin{aligned} \int \partial_z \Gamma^2 r\omega^\theta dx &\leq 2\|\Gamma\|_{L^3} \|\nabla v^\theta\|_{L^2} \|r^2\omega^\theta\|_{L^6} \\ &\leq 4\|\Gamma_0\|_{L^3}^2 \|\nabla v^\theta\|_{L^2}^2 + \frac{1}{4} \|\nabla(r^2\omega^\theta)\|_{L^2}^2. \end{aligned}$$

Next, one also has

$$\begin{aligned} \int \omega^\theta r^2\omega^\theta dx &\leq \|\omega^\theta\|_{L^2(r \leq R(t))}^2 R^2(t) + \|r^2\omega^\theta\|_{L^2(r > R(t))}^2 R^{-2}(t) \\ &\leq \|\omega^\theta\|_{L^2}^2 R^2(t) + \|r^2\omega^\theta\|_{L^2}^2 R^{-2}(t). \end{aligned}$$

Finally, we estimate that

$$\begin{aligned} \int r v^r \omega^\theta r^2 \omega^\theta dx &\leq \|v^r\|_{L^2} \|(r^2 \omega^\theta)^{3/2}\|_{L^4} \|(\omega^\theta)^{1/2}\|_{L^4} \\ &\leq \|v_0\|_{L^2}^4 \|\omega^\theta\|_{L^2}^2 + \frac{1}{4} \|\nabla(r^2 \omega^\theta)\|_{L^2}^2. \end{aligned}$$

By taking $R(t) = \|\Gamma_0\|_{L^3} + \|u_0\|_{L^2}^2$, we arrive at

$$\begin{aligned} \frac{d}{dt} \|r^2 \omega^\theta\|_{L^2}^2 + \|\nabla(r^2 \omega^\theta)\|_{L^2}^2 \\ \lesssim (\|\Gamma_0\|_{L^3}^2 + \|v_0\|_{L^2}^4) \|\nabla v\|_{L^2}^2 + (\|\Gamma_0\|_{L^3}^2 + \|u_0\|_{L^2}^4)^{-1} \|r^2 \omega^\theta\|_{L^2}^2. \end{aligned}$$

Clearly, (2-2) follows by the basic energy estimate and applying Gronwall's inequality to the above differential inequality. \square

Finally, let us prove Corollary 1.3 by using Theorem 1.2.

Proof. It suffices to check the validity of FBC in (1-5)–(1-6) under the assumptions in Corollary 1.3. Let $\delta_0 \in (0, \frac{1}{2})$ and $C_1 > 1$ be arbitrarily large. Noting $\|\Gamma_0\|_{L^\infty} < \infty$ and using the maximum principle, we have

$$\|\Gamma(t, \cdot)\|_{L^\infty} \leq \|\Gamma_0\|_{L^\infty}.$$

Take a smooth cut-off function of r such that

$$\phi \equiv 1 \text{ if } 0 \leq r \leq 1, \quad \phi \equiv 0 \text{ if } r \geq 2.$$

For all $\delta < \delta_0/2$, using (1-7), one has

$$\int \frac{|v^\theta|}{r} \left| \phi\left(\frac{r}{\delta}\right) f \right|^2 r dr dz \leq \int \frac{C_1}{r^2 |\ln r|^2} \left| \phi\left(\frac{r}{\delta}\right) f \right|^2 r dr dz.$$

Using integration by parts, one has

$$\begin{aligned} \int \frac{1}{r^2 |\ln r|^2} \left| \phi\left(\frac{r}{\delta}\right) f \right|^2 r dr dz \\ = \int \left| \phi\left(\frac{r}{\delta}\right) f \right|^2 d|\ln r|^{-1} dz = \int |\ln r|^{-1} \phi\left(\frac{r}{\delta}\right) f \partial_r \left[\phi\left(\frac{r}{\delta}\right) f \right] dr dz \\ \leq \frac{1}{2} \int \frac{1}{r^2 |\ln r|^2} \left| \phi\left(\frac{r}{\delta}\right) f \right|^2 r dr dz + \frac{1}{2} \int \left| \partial_r \left[\phi\left(\frac{r}{\delta}\right) f \right] \right|^2 r dr dz. \end{aligned}$$

Hence, we have

$$\int \frac{1}{r^2 |\ln r|^2} \left| \phi\left(\frac{r}{\delta}\right) f \right|^2 r dr dz \leq \int \left| \partial_r \left[\phi\left(\frac{r}{\delta}\right) f \right] \right|^2 r dr dz,$$

which further gives that

$$\int \frac{|v^\theta|}{r} \left| \phi\left(\frac{r}{\delta}\right) f \right|^2 r dr dz \leq C_1 \int \left| \partial_r \left[\phi\left(\frac{r}{\delta}\right) f \right] \right|^2 r dr dz.$$

On the other hand, it is easy to see that

$$\int \frac{|v^\theta|}{r} \left| \left[1 - \phi\left(\frac{r}{\delta}\right) \right] f \right|^2 r dr dz \leq \|\Gamma_0\|_{L^\infty} \delta^{-2} \int \left| \left[1 - \phi\left(\frac{r}{\delta}\right) \right] f \right|^2 r dr dz.$$

Consequently, we have

$$(2-4) \quad \int \frac{|v^\theta|}{r} |f|^2 r dr dz \leq 2C_1 \int \left| \partial_r \left[\phi\left(\frac{r}{\delta}\right) f \right] \right|^2 r dr dz + 2\|\Gamma_0\|_{L^\infty} \delta^{-2} \int \left| \left[1 - \phi\left(\frac{r}{\delta}\right) \right] f \right|^2 r dr dz \leq 4C_1 \int |\partial_r f|^2 r dr dz + C\delta^{-2} \int_{r \geq \delta} |f|^2 r dr dz.$$

Here and in the next inequality we use C to denote some generic positive constant whose value may change from line to line and which may depend on $\|\Gamma_0\|_{L^\infty}$ and C_1 .

Next, using (2-4), we have

$$(2-5) \quad \int |v^\theta|^2 |f|^2 r dr dz \leq 2 \int |rv^\theta| \frac{|v^\theta|}{r} \left| \phi\left(\frac{r}{\delta}\right) f \right|^2 r dr dz + 2 \int |rv^\theta|^2 r^{-2} \left| \left[1 - \phi\left(\frac{r}{\delta}\right) \right] f \right|^2 r dr dz \leq 2C_1 |\ln \delta|^{-2} \int \frac{|v^\theta|}{r} \left| \phi\left(\frac{r}{\delta}\right) f \right|^2 r dr dz + 2\|\Gamma_0\|_{L^\infty}^2 \delta^{-2} \int_{r \geq \delta} |f|^2 r dr dz \leq C\delta^{-2} \int_{r \geq \delta} |f|^2 r dr dz + 8C_1^2 |\ln \delta|^{-2} \int |\partial_r f|^2 r dr dz.$$

Hence, one may choose δ small enough so that $16C_1^2 |\ln \delta|^{-2} \leq \delta_*$ and choose $C_* = 4C_1$. Then it is clear from (2-4) and (2-5) that the assumptions in equations (1-5)–(1-6) are satisfied. Using Theorem 1.2, one concludes that v is smooth for all $t > 0$. \square

3. Criticality of axisymmetric Navier–Stokes equations

Proof of Theorem 1.2. First of all, for initial data $v_0 \in H^{1/2}$, by the classical results of Leray [1934] and Fujita and Kato [1964], there exists a unique local strong solution v to the Navier–Stokes equations (1-1). Moreover, $v(t, \cdot) \in H^s$ for any $s \geq 0$, at least on a short time interval $[\epsilon, 2\epsilon]$. In particular, $\nabla \omega(t, \cdot) \in L^2$, at least on a short time interval $[\epsilon, 2\epsilon]$. A consequence is that $\nabla \omega^r$, $\nabla \omega^\theta$, $\nabla \omega^z$, ω^r/r , ω^θ/r are all L^2 -functions. In particular, recalling that

$$J = \frac{\omega^r}{r} \quad \text{and} \quad \Omega = \frac{\omega^\theta}{r},$$

one has $J(t, \cdot) \in L^2$ and $\Omega(t, \cdot) \in L^2$ for $t \in [\epsilon, 2\epsilon]$. Inductively, one also has $J(t, \cdot) \in H^2$ and $\Omega(t, \cdot) \in H^2$. Without loss of generality, we may assume that

$$J_0 \in H^2 \quad \text{and} \quad \Omega_0 \in H^2.$$

Otherwise we may start from $t = \epsilon$. As long as the solution is still smooth, one has

$$\|J(t, \cdot)\|_{L^2} + \|\Omega(t, \cdot)\|_{L^2} < \infty, \quad \|\nabla J(t, \cdot)\|_{L^2}^2 + \|\nabla \Omega(t, \cdot)\|_{L^2}^2 < \infty$$

and

$$\int_{-\infty}^{\infty} (|J(t, 0, z)|^2 + |\Omega(t, 0, z)|^2) dz \lesssim \|J(t, \cdot)\|_{H^2}^2 + \|\Omega(t, \cdot)\|_{H^2}^2 < \infty.$$

So all calculations below are legal as long as the solution is still smooth. Our task is to derive a certain sufficiently strong a priori estimate.

By applying the standard energy estimate to the first equation in (1-8), we have

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|J\|_{L^2}^2 &= - \int J(b \cdot \nabla) J r dr dz + \int J \left(\Delta + \frac{2}{r} \partial_r \right) J \\ &\quad + \int J (\omega^r \partial_r + \omega^z \partial_z) \frac{v^r}{r} r dr dz. \end{aligned}$$

Using the incompressibility constraint, one has

$$- \int J(b \cdot \nabla) J r dr dz = \frac{1}{2} \int J^2 \nabla \cdot b r dr dz = 0.$$

On the other hand, by direct calculations, one has

$$\int J \left(\Delta + \frac{2}{r} \partial_r \right) J = - \|\nabla J\|_{L^2}^2 - \int |J(t, 0, z)|^2 dz.$$

Consequently, we have

$$(3-1) \quad \frac{1}{2} \frac{d}{dt} \|J\|_{L^2}^2 + \|\nabla J\|_{L^2}^2 + \int_{-\infty}^{\infty} |J(t, 0, z)|^2 dz = \int J (\omega^r \partial_r + \omega^z \partial_z) \frac{v^r}{r} r dr dz.$$

Similarly, by applying the energy estimate to the second equation in (1-8), one obtains that

$$(3-2) \quad \frac{1}{2} \frac{d}{dt} \|\Omega\|_{L^2}^2 + \|\nabla \Omega\|_{L^2}^2 + \int_{-\infty}^{\infty} |\Omega(t, 0, z)|^2 dz = -2 \int \frac{v^\theta}{r} J \Omega r dr dz.$$

In the remaining part of the proof of [Theorem 1.2](#), we will use C to denote a generic positive constant whose value may change from line to line and which may depend on $\|\Gamma_0\|_{L^\infty}$, C_0 , C_* and r_0 . Using $\|\Gamma\|_{L^\infty} \leq \|\Gamma_0\|_{L^\infty}$ and the form boundedness condition in (1-5), one has

$$\begin{aligned}
(3-3) \quad & \left| \int \frac{v^\theta}{r} J \Omega r \, dr \, dz \right| \\
& \leq \frac{1}{4C_*} \int \left| \frac{v^\theta}{r} \right| \Omega^2 r \, dr \, dz + C_* \int \left| \frac{v^\theta}{r} \right| J^2 r \, dr \, dz \\
& \leq \frac{1}{4} \int |\partial_r \Omega|^2 r \, dr \, dz + C_*^2 \int |\partial_r J|^2 r \, dr \, dz + C \int_{r \geq r_0} (|J|^2 + |\Omega|^2) r \, dr \, dz.
\end{aligned}$$

Inserting (3-3) into (3-2), one has

$$(3-4) \quad \frac{d}{dt} \|\Omega\|_{L^2}^2 + \|\nabla \Omega\|_{L^2}^2 + 2 \int_{-\infty}^{\infty} |\Omega(t, 0, z)|^2 \, dz \leq 2C_*^2 \|\nabla J\|_{L^2}^2 + C \|\omega\|_{L^2}^2.$$

Next, we estimate that

$$\begin{aligned}
\left| \int J(\omega^r \partial_r + \omega^z \partial_z) \frac{v^r}{r} \, dx \right| &= \left| \int [\nabla \times (v^\theta e_\theta)] \cdot \left(J \nabla \frac{v^r}{r} \right) \, dx \right| \\
&\leq \|\nabla J\|_{L^2} \left\| v^\theta \nabla \frac{v^r}{r} \right\|_{L^2}.
\end{aligned}$$

Again, using the form boundedness condition in (1-6), one has

$$\left\| v^\theta \nabla \frac{v^r}{r} \right\|_{L^2}^2 \leq \delta_* \left\| \partial_r \nabla \frac{v^r}{r} \right\|_{L^2}^2 + C_0 \int_{r \geq r_0} \left| \nabla \frac{v^r}{r} \right|^2 \, dr \, dr \, dz.$$

Using Lemma 2.1 and the identity

$$\nabla \frac{v^r}{r} = \frac{\nabla v^r}{r} - e_r \frac{v^r}{r^2} = \frac{\nabla v^r}{r} + e_r \frac{\partial_r v^r + \partial_z v^z}{r},$$

we have

$$\begin{aligned}
(3-5) \quad & \left| \int J(\omega^r \partial_r + \omega^z \partial_z) \frac{v^r}{r} \, dr \, dz \right| \\
& \leq \frac{1}{2} \|\nabla J\|_{L^2}^2 + \frac{\delta_*}{2} \left\| \partial_r \nabla \frac{v^r}{r} \right\|_{L^2}^2 + \frac{C_0}{2} \int_{r \geq r_0} \left| \nabla \frac{v^r}{r} \right|^2 \, dr \, dr \, dz \\
& \leq \frac{1}{2} \|\nabla J\|_{L^2}^2 + \frac{K_0 \delta_*}{2} \|\partial_z \Omega\|_{L^2}^2 + C \int |\nabla v|^2 \, dr \, dr \, dz.
\end{aligned}$$

Inserting (3-5) into (3-1), we have

$$\begin{aligned}
(3-6) \quad & \frac{d}{dt} \|J\|_{L^2}^2 + \|\nabla J\|_{L^2}^2 + 2 \int_{-\infty}^{\infty} |J(t, 0, z)|^2 \, dz \\
& \leq K_0 \delta_* \|\partial_z \Omega\|_{L^2}^2 + C \int |\nabla v|^2 \, dr \, dr \, dz.
\end{aligned}$$

Multiplying (3-6) by $3C_*^2$ and then adding it to (3-4), we have

$$\begin{aligned} \frac{d}{dt} (3C_*^2 \|J\|_{L^2}^2 + \|\Omega\|_{L^2}^2) + (C_*^2 \|\nabla J\|_{L^2}^2 + \|\nabla \Omega\|_{L^2}^2) \\ + \int_{-\infty}^{\infty} (6C_*^2 |J(t, 0, z)|^2 + 2|\Omega(t, 0, z)|^2) dz \\ \leq 3C_*^2 K_0 \delta_* \|\nabla \Omega\|_{L^2}^2 + C \int |\nabla v|^2 r dr dz. \end{aligned}$$

Integrating the above inequality with respect to time, we have

$$\begin{aligned} 3C_*^2 \|J(t, \cdot)\|_{L^2}^2 + \|\Omega(t, \cdot)\|_{L^2}^2 + \int_0^t (C_*^2 \|\nabla J\|_{L^2}^2 + \|\nabla \Omega\|_{L^2}^2) ds \\ + \int_0^t \int_{-\infty}^{\infty} (6C_*^2 |J(t, 0, z)|^2 + 2|\Omega(t, 0, z)|^2) dz ds \\ \leq C C_*^2 (\|J_0\|_{L^2}^2 + \|\Omega_0\|_{L^2}^2) + 3C_*^2 K_0 \delta_* \int_0^t \|\nabla \Omega\|_{L^2}^2 ds + C \|v_0\|_{L^2}. \end{aligned}$$

Here we used the basic energy identity (1-2). Recall that K_0 is an absolute positive constant determined in Lemma 2.1. Hence, we may take δ_* so that $3C_*^2 K_0 \delta_* < \frac{1}{2}$. Consequently, we have

$$(3-7) \quad \sup_{0 \leq t < T} (\|J(t)\|_{L^2}^2 + \|\Omega(t)\|_{L^2}^2) + \int_0^T (\|\nabla J\|_{L^2}^2 + \|\nabla \Omega\|_{L^2}^2) dt \\ + \int_0^T \int_{-\infty}^{\infty} (|J(t, 0, z)|^2 + |\Omega(t, 0, z)|^2) dz dt < \infty$$

for all $T < \infty$.

Clearly, the a priori estimate (3-7) and Sobolev imbedding theorem imply that

$$(3-8) \quad \sup_{0 \leq t < T} \|b(t, \cdot)\|_{L^p(r \leq 1)} < \infty, \quad 3 \leq p \leq 6, \quad T < \infty.$$

Remark 3.1. There are several ways to prove the regularity of v from here. For instance, the easiest way is just to use the result in [Neustupa and Pokorný 2000]. If one would like to assume further decay properties on the initial data so that the conditions in Lemma 2.2 are satisfied, then one can easily derive an $L^\infty([0, T], L^p)$ estimate for b when $r \geq 1$ and $3 \leq p \leq 6$, by using the basic energy estimate and Sobolev imbedding theorem. This, combined with the local L^p estimate of v in (3-8), immediately gives that b is in $L^\infty([0, T], L^3)$ for any $T < \infty$. Using the Sobolev imbedding theorem once more, one has $b \in L^\infty([0, T], \text{BMO}^{-1})$. Then the result in [Lei and Zhang 2011b] implies that v is regular up to time T .

Let us give an alternative and self-contained proof. We first derive an L^4 a priori estimate for v^θ without using the result of [Lei and Zhang 2011b] and Lemma 2.2. Using the equation of v^θ in (1-1) and the standard energy estimate, one has

$$\begin{aligned} \frac{d}{dt} \|v^\theta\|_{L^4}^4 + \|\nabla(v^\theta)^2\|_{L^2}^2 + \|r^{-1}(v^\theta)^2\|_{L^2}^2 &\leq C \left| \int \frac{v^r (v^\theta)^4}{r} r dr dz \right| \\ &\leq C \|r^{-1}v^r\|_{L^\infty} \|v^\theta\|_{L^4}^4. \end{aligned}$$

Hence, by using [Lemma 2.1](#) and the three-dimensional interpolation inequality $\|f\|_{L^\infty}^2 \lesssim \|\nabla f\|_{L^2} \|\nabla^2 f\|_{L^2}$, one has

$$\left\| \frac{v^r}{r} \right\|_{L^\infty} \lesssim \left\| \nabla \partial_z \frac{\psi^\theta}{r} \right\|_{L^2}^{1/2} \left\| \nabla^2 \partial_z \frac{\psi^\theta}{r} \right\|_{L^2}^{1/2} = \left\| \nabla \frac{v^r}{r} \right\|_{L^2}^{1/2} \left\| \nabla^2 \frac{v^r}{r} \right\|_{L^2}^{1/2} \lesssim \|\Omega\|_{L^2}^{1/2} \|\partial_z \Omega\|_{L^2}^{1/2},$$

and using (3-7), one concludes from Gronwall's inequality that

$$\|v^\theta\|_{L^4} < \infty, \quad \text{and} \quad \int_0^T \|r^{-1}(v^\theta)^2\|_{L^2}^2 dt < \infty, \quad 0 \leq t \leq T.$$

Then we use the second equation of (2-3) to derive that

$$\begin{aligned} \frac{d}{dt} \|\omega^\theta\|_{L^2}^2 + 2\|\nabla \omega^\theta\|_{L^2}^2 + 2\|r^{-1}\omega^\theta\|_{L^2}^2 \\ = - \int \frac{v^r}{r} (\omega^\theta)^2 r dr dz + \int \omega^\theta \partial_z \frac{(v^\theta)^2}{r} r dr dz \\ \leq C \|r^{-1}v^r\|_{L^\infty} \|\omega^\theta\|_{L^2}^2 + \|\partial_z \omega^\theta\|_{L^2}^2 + \frac{1}{4} \|r^{-1}(v^\theta)^2\|_{L^2}^2. \end{aligned}$$

Hence, Gronwall's inequality similarly gives that

$$\omega^\theta \in L^2, \quad T < \infty.$$

By the basic energy identity (1-2) and Sobolev imbedding, one has

$$b \in L^p, \quad 2 \leq p \leq 6.$$

Hence, $v \in L_T^\infty(L_x^4)$. So the Serrin-type criterion implies v is regular up to time T . \square

4. Small $\|\Gamma_0\|_{L^\infty}$ or $\|\Gamma(t, \cdot)\|_{L^\infty(r \leq r_0)}$ global regularity

This section is devoted to proving [Theorem 1.4](#).

Proof of Theorem 1.4. Recall that

$$V = \frac{v^\theta}{\sqrt{r}}, \quad \Omega = \frac{\omega^\theta}{r}.$$

Let us first formulate the axisymmetric Navier-Stokes equations (1-1) in terms of V and Ω as follows:

$$(4-1) \quad \begin{cases} \partial_t V + b \cdot \nabla V + \frac{3v^r}{2r} V = \left(\Delta + \frac{1}{r} \partial_r - \frac{3}{4r^2} \right) V, \\ \partial_t \Omega + b \cdot \nabla \Omega = \left(\Delta + \frac{2}{r} \partial_r \right) \Omega + \frac{2\partial_z V^2}{r}. \end{cases}$$

By the energy estimate, one has

$$(4-2) \quad \frac{1}{2} \frac{d}{dt} \|\Omega\|_{L^2}^2 + \|\nabla \Omega\|_{L^2}^2 \leq \frac{1}{2} \|\partial_z \Omega\|_{L^2}^2 + \frac{1}{2} \|r^{-1} |V|^2\|_{L^2}^2.$$

Let us first prove the global regularity under

$$\|\Gamma_0\|_{L^\infty} \leq \delta M_0^{-1}.$$

Using [Lemma 2.1](#), one has

$$\left\| \frac{v^r}{r} \right\|_{L^\infty} \lesssim \|\Omega\|_{L^2}^{1/2} \|\partial_z \Omega\|_{L^2}^{1/2}.$$

Noting that

$$\|V\|_{L^4}^4 \lesssim \|r^{-1} |V|^2\|_{L^2}^{3/2} \|\Gamma\|_{L^4},$$

one can apply the L^4 energy estimate for V to get

$$(4-3) \quad \begin{aligned} \frac{d}{dt} \| |V|^2 \|_{L^2}^2 + \|\nabla |V|^2\|_{L^2}^2 + \|r^{-1} |V|^2\|_{L^2}^2 \\ \lesssim \left\| \frac{v^r}{r} \right\|_{L^\infty} \|V\|_{L^4}^4 \lesssim \|\Omega\|_{L^2}^{1/2} \|\partial_z \Omega\|_{L^2}^{1/2} \|r^{-1} |V|^2\|_{L^2}^{3/2} \|\Gamma\|_{L^4} \\ \lesssim \|\Omega\|_{L^2}^{1/2} \|\Gamma\|_{L^2}^{1/2} \|\Gamma\|_{L^\infty}^{1/2} (\|\partial_z \Omega\|_{L^2}^2 + \|r^{-1} |V|^2\|_{L^2}^2). \end{aligned}$$

Combining (4-2) and (4-3), we arrive at

$$(4-4) \quad \begin{aligned} \frac{d}{dt} (\| |V|^2 \|_{L^2}^2 + \|\Omega\|_{L^2}^2) + (\|\nabla |V|^2\|_{L^2}^2 + \|\nabla \Omega\|_{L^2}^2) + \|r^{-1} |V|^2\|_{L^2}^2 \\ \lesssim \|\Omega\|_{L^2}^{1/2} \|\Gamma\|_{L^2}^{1/2} \|\Gamma\|_{L^\infty}^{1/2} (\|\partial_z \Omega\|_{L^2}^2 + \|r^{-1} |V|^2\|_{L^2}^2). \end{aligned}$$

Recall that we have the following a priori estimate:

$$\|\Gamma\|_{L^2} \leq \|\Gamma_0\|_{L^2}, \quad \|\Gamma\|_{L^\infty} \leq \|\Gamma_0\|_{L^\infty}.$$

Hence, under the condition of the theorem, there exists $T > 0$ such that

$$\| |V|^2 \|_{L^2}^2 + \|\Omega\|_{L^2}^2 < 2\| |V_0|^2 \|_{L^2}^2 + 2\|\Omega_0\|_{L^2}^2, \quad \forall 0 \leq t < T.$$

If δ is a suitably small positive constant and

$$\|\Gamma_0\|_{L^\infty} \leq \delta M_0^{-1}$$

is satisfied, then we have

$$\|\Omega\|_{L^2}^{1/2} \|\Gamma\|_{L^2}^{1/2} \|\Gamma\|_{L^\infty}^{1/2} \lesssim M_0^{1/2} (\delta M_0^{-1})^{1/2} \lesssim \delta^{1/2}, \quad \forall 0 \leq t < T.$$

Hence, by (4-4), we derive that

$$\frac{d}{dt} (\| |V|^2 \|_{L^2}^2 + \|\Omega\|_{L^2}^2) \leq 0, \quad \forall 0 \leq t < T,$$

which implies that

$$\| |V|^2 \|_{L^2}^2 + \|\Omega\|_{L^2}^2 \leq \| |V_0|^2 \|_{L^2}^2 + \|\Omega_0\|_{L^2}^2, \quad 0 \leq t \leq T.$$

The above argument implies, by the standard continuation method that

$$\| |V|^2 \|_{L^2}^2 + \|\Omega\|_{L^2}^2 \leq \| |V_0|^2 \|_{L^2}^2 + \|\Omega_0\|_{L^2}^2, \quad \forall t \geq 0.$$

Hence the proof for first part of the theorem is finished.

On the other hand, if

$$\|\Gamma(t, \cdot)\|_{L^\infty(r \leq r_0)} \leq \delta M_1^{-1}$$

is satisfied, then one may treat (4-3) as follows:

$$\begin{aligned} \frac{d}{dt} \| |V|^2 \|_{L^2}^2 + \|\nabla |V|^2 \|_{L^2}^2 + \|r^{-1}|V|^2 \|_{L^2}^2 \\ \lesssim \left\| \frac{v^r}{r} \right\|_{L^\infty} \|V\|_{L^4(r \leq r_0)}^4 + \int_{r \geq r_0} \left| \frac{v^r}{r} \frac{(v^\theta)^4}{r^2} \right| r \, dr \, dz \\ \lesssim \|\Omega\|_{L^2}^{1/2} \|\Gamma\|_{L^2}^{1/2} \|\Gamma\|_{L^\infty(r \leq r_0)}^{1/2} (\|\partial_z \Omega\|_{L^2}^2 + \|r^{-1}|V|^2 \|_{L^2}^2) \\ + r_0^{-4} \left\| \frac{v^r}{r} \right\|_{L^2} \left\| \frac{v^\theta}{r} \right\|_{L^2} \|\Gamma\|_{L^\infty(r \geq r_0)}^3, \end{aligned}$$

which, combined with (4-2), gives that

$$\begin{aligned} \| |V(t, \cdot)|^2 \|_{L^2}^2 + \|\Omega(t, \cdot)\|_{L^2}^2 + \int_0^t (\|\nabla |V|^2 \|_{L^2}^2 + \|\nabla \Omega\|_{L^2}^2) \, ds \\ \lesssim \| |V_0|^2 \|_{L^2}^2 + \|\Omega_0\|_{L^2}^2 + r_0^{-4} \|v_0\|_{L^2}^2 \|\Gamma_0\|_{L^\infty}^3 \\ + (\|\Gamma\|_{L^\infty(r \leq r_0)} \|\Gamma_0\|_{L^2} \sup_{0 \leq s \leq t} \|\Omega(s, \cdot)\|_{L^2})^{1/2} (\|\partial_z \Omega\|_{L^2}^2 + \|r^{-1}|V|^2 \|_{L^2}^2). \end{aligned}$$

Here $r_0 > 0$ is arbitrary. Then similar continuation arguments as in the proof used in the first part imply that if δ is a suitable small absolute positive constant, then the solution v is regular. Here M_1 is given in the statement of [Theorem 1.4](#). This shows that the smallness of Γ locally in r implies the regularity of the solutions. \square

Remark. Since $\| |V_0|^2 \|_{L^2}$ and $\|\Omega_0\|_{L^2}$ have dimension $-\frac{3}{2}$, and $\|\Gamma_0\|_{L^2}$ has dimension $\frac{3}{2}$, the constant M_0 in [Theorem 1.4](#) has dimension 0. Similarly, one can check that M_1 is also dimensionless if one assigns r_0 dimension 1.

Acknowledgements

The authors would like to thank Prof. Daoyuan Fang, Dr. Bin Han, Mr. Xinghong Pan and Prof. Ting Zhang for discussions. Z. Lei was in part supported by NSFC (Grants No.11171072, 11421061 and 11222107), the Shanghai Shu Guang project, the Shanghai Talent Development Fund and SGST 09DZ2272900. Qi S. Zhang is grateful for the support of the Siyuan Foundation through Nanjing University, and the Simons foundation.

References

- [Burke Loftus and Zhang 2010] J. Burke Loftus and Q. S. Zhang, “A priori bounds for the vorticity of axially symmetric solutions to the Navier–Stokes equations”, *Adv. Differential Equations* **15**:5/6 (2010), 531–560. [MR](#) [Zbl](#)
- [Chae and Lee 2002] D. Chae and J. Lee, “On the regularity of the axisymmetric solutions of the Navier–Stokes equations”, *Math. Z.* **239**:4 (2002), 645–671. [MR](#) [Zbl](#)
- [Chen et al. 2008] C.-C. Chen, R. M. Strain, H.-T. Yau, and T.-P. Tsai, “Lower bound on the blow-up rate of the axisymmetric Navier–Stokes equations”, *Int. Math. Res. Not.* **2008**:9 (2008), art. id. rnn016, 31 pp. [MR](#) [Zbl](#)
- [Chen et al. 2009] C.-C. Chen, R. M. Strain, T.-P. Tsai, and H.-T. Yau, “Lower bounds on the blow-up rate of the axisymmetric Navier–Stokes equations, II”, *Comm. Partial Differential Equations* **34**:1-3 (2009), 203–232. [MR](#) [Zbl](#)
- [Chen et al. 2015] H. Chen, D. Fang, and T. Zhang, “Regularity of 3D axisymmetric Navier–Stokes equations”, preprint, 2015. [arXiv](#)
- [Fujita and Kato 1964] H. Fujita and T. Kato, “On the Navier–Stokes initial value problem, I”, *Arch. Rational Mech. Anal.* **16** (1964), 269–315. [MR](#) [Zbl](#)
- [Hou and Li 2008] T. Y. Hou and C. Li, “Dynamic stability of the three-dimensional axisymmetric Navier–Stokes equations with swirl”, *Comm. Pure Appl. Math.* **61**:5 (2008), 661–697. [MR](#) [Zbl](#)
- [Hou et al. 2008] T. Y. Hou, Z. Lei, and C. Li, “Global regularity of the 3D axisymmetric Navier–Stokes equations with anisotropic data”, *Comm. Partial Differential Equations* **33**:7-9 (2008), 1622–1637. [MR](#) [Zbl](#)
- [Jiu and Xin 2003] Q. Jiu and Z. Xin, “Some regularity criteria on suitable weak solutions of the 3D incompressible axisymmetric Navier–Stokes equations”, pp. 119–139 in *Lectures on partial differential equations*, edited by S.-Y. A. Chang et al., New Stud. Adv. Math. **2**, International Press, Somerville, MA, 2003. [MR](#) [Zbl](#)
- [Koch et al. 2009] G. Koch, N. Nadirashvili, G. A. Seregin, and V. Šverák, “Liouville theorems for the Navier–Stokes equations and applications”, *Acta Math.* **203**:1 (2009), 83–105. [MR](#) [Zbl](#)
- [Ladyzhenskaya 1968] O. A. Ladyzhenskaya, “Unique global solvability of the three-dimensional Cauchy problem for the Navier–Stokes equations in the presence of axial symmetry”, *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **7** (1968), 155–177. In Russian; translated in *Sem. Math. V. A. Steklov Math. Inst. Leningrad* **7** (1970), 70–79. [MR](#) [Zbl](#)
- [Lei 2015] Z. Lei, “On axially symmetric incompressible magnetohydrodynamics in three dimensions”, *J. Differential Equations* **259**:7 (2015), 3202–3215. [MR](#) [Zbl](#)
- [Lei and Zhang 2011a] Z. Lei and Q. S. Zhang, “Structure of solutions of 3D axisymmetric Navier–Stokes equations near maximal points”, *Pacific J. Math.* **254**:2 (2011), 335–344. [MR](#) [Zbl](#)
- [Lei and Zhang 2011b] Z. Lei and Q. S. Zhang, “A Liouville theorem for the axially-symmetric Navier–Stokes equations”, *J. Funct. Anal.* **261**:8 (2011), 2323–2345. [MR](#) [Zbl](#)
- [Lei et al. 2013] Z. Lei, E. A. Navas, and Q. S. Zhang, “A priori bound on the velocity in axially symmetric Navier–Stokes equations”, preprint, 2013. [arXiv](#)
- [Leonardi et al. 1999] S. Leonardi, J. Málek, J. Nečas, and M. Pokorný, “On axially symmetric flows in \mathbb{R}^3 ”, *Z. Anal. Anwendungen* **18**:3 (1999), 639–649. [MR](#) [Zbl](#)
- [Leray 1934] J. Leray, “Sur le mouvement d’un liquide visqueux emplissant l’espace”, *Acta Math.* **63**:1 (1934), 193–248. [MR](#) [Zbl](#)
- [Majda and Bertozzi 2002] A. J. Majda and A. L. Bertozzi, *Vorticity and incompressible flow*, Cambridge Texts in Applied Mathematics **27**, Cambridge University Press, 2002. [MR](#) [Zbl](#)

- [Miao and Zheng 2013] C. Miao and X. Zheng, “On the global well-posedness for the Boussinesq system with horizontal dissipation”, *Comm. Math. Phys.* **321**:1 (2013), 33–67. [MR](#) [Zbl](#)
- [Neustupa and Pokorný 2000] J. Neustupa and M. Pokorný, “An interior regularity criterion for an axially symmetric suitable weak solution to the Navier–Stokes equations”, *J. Math. Fluid Mech.* **2**:4 (2000), 381–399. [MR](#) [Zbl](#)
- [Neustupa and Pokorný 2001] J. Neustupa and M. Pokorný, “Axisymmetric flow of Navier–Stokes fluid in the whole space with non-zero angular velocity component”, *Math. Bohem.* **126**:2 (2001), 469–481. [MR](#) [Zbl](#)
- [Pan 2016] X. Pan, “Regularity of solutions to axisymmetric Navier–Stokes equations with a slightly supercritical condition”, *J. Differential Equations* **260**:12 (2016), 8485–8529. [MR](#) [Zbl](#)
- [Seregin and Šverák 2009] G. Seregin and V. Šverák, “On type I singularities of the local axisymmetric solutions of the Navier–Stokes equations”, *Comm. Partial Differential Equations* **34**:1-3 (2009), 171–201. [MR](#) [Zbl](#)
- [Tao 2007] T. Tao, “Why global regularity for Navier–Stokes is hard”, blog entry, 2007, available at <http://tinyurl.com/tao-navier-stokes>.
- [Tian and Xin 1998] G. Tian and Z. Xin, “One-point singular solutions to the Navier–Stokes equations”, *Topol. Methods Nonlinear Anal.* **11**:1 (1998), 135–145. [MR](#) [Zbl](#)
- [Ukhovskii and Iudovich 1968] M. R. Ukhovskii and V. I. Iudovich, “Axially symmetric flows of ideal and viscous fluids filling the whole space”, *J. Appl. Math. Mech.* **32** (1968), 52–61. [MR](#) [Zbl](#)
- [Wei 2016] D. Wei, “Regularity criterion to the axially symmetric Navier–Stokes equations”, *J. Math. Anal. Appl.* **435**:1 (2016), 402–413. [MR](#) [arXiv](#)
- [Zhang and Zhang 2014] P. Zhang and T. Zhang, “Global axisymmetric solutions to three-dimensional Navier–Stokes system”, *Int. Math. Res. Not.* **2014**:3 (2014), 610–642. [MR](#)

Received July 19, 2016. Revised September 26, 2016.

ZHEN LEI
SCHOOL OF MATHEMATICAL SCIENCES
FUDAN UNIVERSITY
SHANGHAI, 200433
CHINA
zlei@fudan.edu.cn

QI S. ZHANG
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF CALIFORNIA, RIVERSIDE
RIVERSIDE, CA 92521
UNITED STATES
qizhang@math.ucr.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 289 No. 1 July 2017

Restricted Poisson algebras	1
YAN-HONG BAO, YU YE and JAMES J. ZHANG	
Remarks on GJMS operator of order six	35
XUEZHANG CHEN and FEI HOU	
On the asymptotic behavior of Bergman kernels for positive line bundles	71
TIEN-CUONG DINH, XIAONAN MA and VIÊT-ANH NGUYÊN	
Molino theory for matchbox manifolds	91
JESSICA DYER, STEVEN HURDER and OLGA LUKINA	
Cyclic pursuit on compact manifolds	153
DMITRI GEKHTMAN	
Criticality of the axially symmetric Navier–Stokes equations	169
ZHEN LEI and QI S. ZHANG	
Convexity of the entropy of positive solutions to the heat equation on quaternionic contact and CR manifolds	189
DIMITER VASSILEV	
On handlebody structures of rational balls	203
LUKE WILLIAMS	
On certain Fourier coefficients of Eisenstein series on G_2	235
WEI XIONG	