

*Pacific
Journal of
Mathematics*

**LOCAL CONSTANCY OF DIMENSION OF SLOPE SUBSPACES
OF AUTOMORPHIC FORMS**

JOACHIM MAHNKOPF

Volume 289 No. 2

August 2017

LOCAL CONSTANCY OF DIMENSION OF SLOPE SUBSPACES OF AUTOMORPHIC FORMS

JOACHIM MAHNKOPF

We prove an analogue of a Gouvêa–Mazur conjecture on local constancy of dimension of slope subspaces of modular forms on the upper half plane for automorphic forms on reductive algebraic groups \tilde{G}/\mathbb{Q} having discrete series. The proof uses a comparison of Bewersdorff’s elementary trace formula for pairs of congruent weights and does not make use of methods from p -adic Banach space theory, overconvergent forms or rigid analytic geometry.

We also compare two Goresky–MacPherson trace formulas computing Lefschetz numbers on weighted cohomology for pairs of congruent weights; this has an application to a more explicit version of the Gouvêa–Mazur conjecture for symplectic groups of rank 2.

Introduction

0.1. We fix a prime $p \in \mathbb{N}$ and an integer N not divisible by p . Generalizing Hida’s theory [1993; 1988] of ordinary modular forms, Gouvêa and Mazur [1992] conjectured that the dimension $d(\beta, k)$ of the slope β subspace of the space $S_k(\Gamma_0(pN))$ of cuspidal modular forms of level pN and weight k is locally constant in the p -adic topology as a function of k . More precisely, they conjectured that there is a linear polynomial $m(x)$ such that the conditions $k, k' \geq 2\beta + 2$ and $k \equiv k' \pmod{(p-1)p^{m-1}}$ with $m \geq m(\beta)$ imply

$$(1) \quad d(\beta, k) = d(\beta, k').$$

Using work of Coleman [1997] which is based on rigid analytic geometry and p -adic spectral theory, as well as Katz’s theory of p -adic modular forms and results of Gouvêa and Mazur, Wan [1998] proved that there is a quadratic polynomial $m(x)$ such that equation (1) holds. On the other hand, Buzzard and Calegari [2004] showed that in general there is no linear polynomial $m(x)$ such that (1) holds, hence, Wan’s result is best possible.

MSC2010: 11F75.

Keywords: Gouvêa–Mazur conjecture, cohomology of arithmetic groups.

0.2. In this article we prove a higher-rank analogue of the Gouvêa–Mazur conjecture. To describe this in more detail, we denote by \mathbb{A} the adèles of \mathbb{Q} and fix a prime $p \in \mathbb{N}$. We let $\tilde{\mathbf{G}}/\mathbb{Q}$ be a connected reductive algebraic group which contains a maximal torus $\tilde{\mathbf{T}}/\mathbb{Q}$ which splits over \mathbb{Q}_p . We select a basis Δ of the root system Φ of $\tilde{\mathbf{G}}/F$ where F/\mathbb{Q} is a (minimal) splitting field for $\tilde{\mathbf{T}}/\mathbb{Q}$. We let $\tilde{\mathbf{K}} \leq \tilde{\mathbf{G}}(\mathbb{A}_f)$ be a compact open subgroup with p -component $\tilde{\mathbf{K}}_p$ equal to the Iwahori subgroup $\tilde{\mathcal{I}}$ of $\tilde{\mathbf{G}}(\mathbb{Q}_p)$. We denote by \mathbb{T} the Hecke operator attached to the double coset $\tilde{\mathbf{K}}h^{-1}r\tilde{\mathbf{K}}$ where $h \in \tilde{\mathbf{T}}(\mathbb{Q})^{++} \leq \tilde{\mathbf{T}}(\mathbb{Q}_p)^{++}$ is a strictly dominant element and $r \in \tilde{\mathbf{G}}(\mathbb{A}_f)^{(p)}$, i.e., r has trivial p -component. For any dominant weight $\tilde{\lambda} \in X(\tilde{\mathbf{T}})$ we understand by $L_{\tilde{\lambda}}$ the irreducible representation of $\tilde{\mathbf{G}}/F$ of highest weight $\tilde{\lambda}$. The normalization $\mathbb{T}_{\tilde{\lambda}}$ of \mathbb{T} acts on full cohomology $H^i(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{Q}_p))$ as well as on cuspidal cohomology $H_{\text{cusp}}^i(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{C}))$ where $S_{\tilde{\mathbf{K}}} = \tilde{\mathbf{G}}(\mathbb{Q}) \backslash \tilde{\mathbf{G}}(\mathbb{A}) / \tilde{\mathbf{K}}\tilde{\mathbf{K}}_{\infty}$ is the locally symmetric space. We write $H^i(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{\beta}$ (resp. $H^i(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{\leq \beta}$) for the subspace of slope β (resp. slope $\leq \beta$) w.r.t. $\mathbb{T}_{\tilde{\lambda}}$ and we use analogous notation for cuspidal cohomology. Our main results then are as follows.

Theorem A (see 3.10 Corollary, 4.11.4 Theorem). *Let $s = |\Phi^+|$ be the number of positive roots of $\tilde{\mathbf{G}}/\mathbb{Q}_p$, $\sigma = \max_{\alpha \in \Phi^+} \text{ht}(\alpha)$ the maximal height of a positive root and \mathbf{g}_i the number of i -cells in a finite cell complex \mathcal{Z} which is homotopy equivalent to the Borel Serre compactification $\bar{S}_{\tilde{\mathbf{K}}}$ of $S_{\tilde{\mathbf{K}}}$. Then for all $\beta \in \mathbb{Q}_{\geq 0}$, $\tilde{\lambda} \in X(\tilde{\mathbf{T}})^{\text{dom}}$ and $i \in \mathbb{N}_0$ we obtain*

$$\dim H^i(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{\leq \beta} \leq \mathfrak{m}\beta^s + \mathfrak{n};$$

here, $\mathfrak{m} = 12\mathbf{g}_i\sigma^{s+1}/s$ and $\mathfrak{n} \in \mathbb{N}$ is an integer which also only depends on $\tilde{\mathbf{K}}$ (and, hence, on $\tilde{\mathbf{G}}$ and p) and on i .

Theorem B (see 5.2 Theorem). *We assume that $\tilde{\mathbf{G}}$ has discrete series and we denote by $d = d_{\tilde{\mathbf{G}}}$ the middle degree. There are polynomials $\mathbf{m}_1(x), \mathbf{m}_2(x) \in \mathbb{Q}[x]$ both of degree $s + 1$ and leading term $12\mathbf{g}_d\sigma^{s+1}/s$ which only depend on $\tilde{\mathbf{K}}$ (hence, on $\tilde{\mathbf{G}}$ and p) and $h \in \tilde{\mathbf{T}}(\mathbb{Q})^{++}$ with the following property. Let $\beta \in \mathbb{Q}_{\geq 0}$. Suppose the dominant weights $\tilde{\lambda}, \tilde{\lambda}' \in X(\tilde{\mathbf{T}})$ satisfy*

- $\langle \tilde{\lambda}, \alpha^{\vee} \rangle > 2\mathbf{m}_1(\beta)$ and $\langle \tilde{\lambda}', \alpha^{\vee} \rangle > 2\mathbf{m}_1(\beta)$ for all $\alpha \in \Delta_{\tilde{\mathbf{G}}}$;
- $\tilde{\lambda} \equiv \tilde{\lambda}' \pmod{(p-1)p^{m-1}X(\tilde{\mathbf{T}})}$ with $m \geq \mathbf{m}_2(\beta)$ ($m \in \mathbb{N}$).

Then

$$\dim H_{\text{cusp}}^d(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{C}))^{\gamma} = \dim H_{\text{cusp}}^d(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}'}(\mathbb{C}))^{\gamma} \quad \text{for all } 0 \leq \gamma \leq \beta.$$

Remark. In the \mathbf{GL}_2 -case $\mathbf{m}_2(x)$ is a quadratic polynomial; i.e., we obtain the same growth as that of $\mathbf{m}(x)$ in [Wan 1998], except that the weights have to satisfy a stronger lower bound (quadratic in β instead of linear as in that paper).

0.3. To prove Theorems A and B we will mostly work in a non-adelic setting; i.e., $\Gamma \leq \tilde{G}(\mathbb{Q})$ denotes an arithmetic subgroup contained in $\tilde{\mathcal{I}}$. Theorem A then is an extension of the main result of [Mahnkopf 2014] (see Section 3.1) and the proof is based on an extension of the notion of truncation of an irreducible representation of \tilde{G}/\mathbb{Q}_p introduced in [Mahnkopf 2013; 2014].

Using the boundedness result of Theorem A the proof of Theorem B reduces to proving certain congruences between traces of powers of the Hecke operator $\mathbb{T}_{\tilde{\lambda}}$ on $H_{\text{cusp}}^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))$ and on $H_{\text{cusp}}^i(S_{\tilde{K}}, L_{\tilde{\lambda}'})$ for p -adically close weights $\tilde{\lambda}, \tilde{\lambda}'$. We first verify these congruences on full cohomology and our principal tool for this is a comparison of a simple and elementary trace formula of Bewersdorff [1985] for cohomology with coefficients in $L_{\tilde{\lambda}}$ and in $L_{\tilde{\lambda}'}$. The equality of mod p^n reductions of geometric sides essentially follows from p -adic properties of the diagonalization of elements in $\tilde{\mathcal{I}}h^e\tilde{\mathcal{I}} \subseteq \tilde{G}(\mathbb{Q}_p)$, $e \in \mathbb{N}$, which are proved using basic algebra (see 4.3 Lemma and 4.4 Proposition); we note that $\tilde{\mathcal{I}}h^{-e}\tilde{\mathcal{I}}$ is the p -component of \mathbb{T}^e . To obtain congruences on cuspidal cohomology we directly prove congruences on the Eisenstein part of full cohomology and subtract from congruences on full cohomology.

Since the Bewersdorff trace formula is elementary we obtain an elementary proof of the congruences on full cohomology and the proofs of Theorems A and B do not make use of methods from p -adic Banach space theory, overconvergent cohomology or rigid analytic geometry (but use the spectral decomposition of full cohomology for regular weight).

0.4. Weighted cohomology. Goresky and MacPherson [Goresky and MacPherson 2003] proved a trace formula for Lefschetz numbers of Hecke operators on weighted cohomology. Unlike Bewersdorff’s formula it contains contributions not only from \tilde{G} but from all \mathbb{Q} -parabolic subgroups of \tilde{G} . Nevertheless, the same diagonalization of elements in $\tilde{\mathcal{I}}h^e\tilde{\mathcal{I}} \subseteq \tilde{G}(\mathbb{Q}_p)$ as in 0.3 allows to compare two Goresky–MacPherson trace formulas for pairs of congruent weights. This then yields certain congruences on weighted cohomology groups and has an application to a version of the Gouvêa–Mazur conjecture for symplectic groups of rank 2 which is more explicit since we avoid use of the spectral decomposition of full cohomology (see Section 5.8). We note that this depends on properties of the root system C_2 but using instead the Goresky–Kottwitz–MacPherson trace formula [Goresky et al. 1997] together with the calculations of Spallone [2009] it might be possible to extend this to arbitrary reductive groups \tilde{G}/\mathbb{Q} .

0.5. Buzzard [Buzzard 2001] gave an elementary proof of boundedness of dimension of slope subspaces in the case \mathbf{GL}_2/\mathbb{Q} also based on an analysis of representations of $\mathbf{GL}_2(\mathbb{Z}_p)$. In the case of quaternion algebras over \mathbb{Q} he also proved in [Buzzard 1998] local constancy of dimension of slope subspaces and his

results were generalized to \mathbf{GL}_2 over totally real fields by Pande [2009]. Following the method of Ash and Stevens [2008], who introduced overconvergent cohomology, Urban [2011] obtained p -adic families of systems of Hecke eigenvalues; he uses this to also derive a p -adic trace formula on overconvergent cohomology. Andreatta, Iovita and Pilloni also proved existence of p -adic families of eigenforms using rigid analytic geometry; see [Andreatta et al. 2015].

More closely related to our approach is work of Koike [1975; 1976] (and some unpublished work of Clozel); like Buzzard, Koike does not make use of methods from rigid analytic geometry or p -adic Banach space theory. In the case of cuspidal modular forms, i.e., in the case \mathbf{GL}_2/\mathbb{Q} he uses a Selberg trace formula which yields an explicit expression for the trace of Hecke operators to deduce congruences between traces of Hecke operators. Since the Selberg trace formula becomes much more involved this seems difficult to generalize to higher rank. We therefore do not attempt to determine an explicit expression for the trace of Hecke operators but only equate mod p^n reductions of traces for p -adically close weights $\tilde{\lambda}, \tilde{\lambda}'$. This can be done even in higher rank by comparing the simple (non-explicit) trace formula of Bewersdorff for weights $\tilde{\lambda}$ and $\tilde{\lambda}'$.

1. Chevalley groups

We recall some basic facts from the theory of Chevalley groups and their representations and we give proofs for some (technical) results for which we do not know a reference.

1.1. Complex semisimple Lie algebras. Let \mathfrak{g} be a complex semisimple Lie algebra. We denote by \mathfrak{h} a Cartan subalgebra of \mathfrak{g} and by $\Phi = \Phi(\mathfrak{g}, \mathfrak{h})$ the set of roots of \mathfrak{g} w.r.t. to \mathfrak{h} . We choose a basis Δ of Φ and we denote by Φ^+ the set of positive roots. For each root $\alpha \in \Phi$ we write $\mathfrak{g}(\alpha)$ for the corresponding root subspace of \mathfrak{g} and we select elements $h_\alpha \in \mathfrak{h}$, $\alpha \in \Delta$, and $x_\alpha \in \mathfrak{g}(\alpha)$, $\alpha \in \Phi$, such that $\{h_\alpha, \alpha \in \Delta, x_\beta, \beta \in \Phi\}$ is a *Chevalley basis* of \mathfrak{g} . In particular, h_α is the coroot corresponding to $\alpha \in \Delta$. The Chevalley basis yields \mathbb{Z} -forms $\mathfrak{g}(\mathbb{Z}) = \bigoplus_{\beta \in \Phi} \mathbb{Z}x_\beta \oplus \bigoplus_{\alpha \in \Delta} \mathbb{Z}h_\alpha$ (resp. $\mathfrak{h}(\mathbb{Z}) = \bigoplus_{\alpha \in \Delta} \mathbb{Z}h_\alpha$) of \mathfrak{g} (resp. of \mathfrak{h}). We denote by $\mathcal{U}_{\mathbb{Z}}$ the \mathbb{Z} -form of the universal enveloping algebra \mathcal{U} of \mathfrak{g} which as a ring is generated by the elements $x_\alpha^n/n!$, $\alpha \in \Phi$, $n \in \mathbb{N}_0$ (see [Humphreys 1972, Theorem 26.4, p. 156]). We set $\mathfrak{g}(R) = \mathfrak{g}(\mathbb{Z}) \otimes R$, $\mathfrak{h}(R) = \mathfrak{h}(\mathbb{Z}) \otimes R$ and $\mathcal{U}_R = \mathcal{U}_{\mathbb{Z}} \otimes R$, R a \mathbb{Z} -algebra. We set $s = |\Phi^+|$ and we fix an ordering $\Phi^+ = \{\alpha_1, \dots, \alpha_s\}$ of the set of positive roots and we set

$$X_{\pm}^{\mathbf{n}} = \frac{x_{\pm\alpha_1}^{n_1}}{n_1!} \dots \frac{x_{\pm\alpha_s}^{n_s}}{n_s!} \in \mathcal{U}_{\mathbb{Z}},$$

where $\mathbf{n} = (n_1, \dots, n_s) \in \mathbb{N}_0^s$. The \mathbb{Z} -span of the elements $X_{\pm}^{\mathbf{n}}$, where $\mathbf{n} \in \mathbb{N}_0^s$, is a

\mathbb{Z} -form $\mathcal{U}_{\mathbb{Z}}^-$ of the universal enveloping algebra \mathcal{U}^- of $\mathfrak{n}^- = \bigoplus_{\alpha < 0} \mathfrak{g}(\alpha)$. Finally, $\mu \leq \lambda$, for $\lambda, \mu \in \mathfrak{h}^*$, means that $\lambda - \mu$ is a linear combination of positive roots with nonnegative coefficients. For any $\lambda \in \mathfrak{h}^*$ we define a relative height function $\text{ht}_{\lambda} : \{\mu \in \mathfrak{h}^* : \mu \leq \lambda\} \rightarrow \mathbb{N}_0$ by $\text{ht}_{\lambda}(\mu) = \text{ht}(\lambda - \mu)$ (see [Mahnkopf 2013, 1.3]); here, $\text{ht} = \text{ht}_{\Delta}$ is the height function corresponding to Δ , i.e., $\text{ht}(\lambda - \mu) = \sum_{\alpha \in \Delta} n_{\alpha}$ if $\lambda - \mu = \sum_{\alpha \in \Delta} n_{\alpha} \alpha$. By $\omega_{\alpha} \in \mathfrak{h}^*$, $\alpha \in \Delta$, we understand the fundamental (dominant) weights, i.e., $\omega_{\beta}(h_{\alpha}) = \delta_{\alpha, \beta}$. The fundamental weights span the weight lattice Γ_{sc} of \mathfrak{g} which contains the root lattice Γ_{ad} .

For any integral and dominant weight $\lambda \in \mathfrak{h}^*$ we denote by $(\rho_{\lambda}, L_{\lambda})$ the complex irreducible \mathfrak{g} -module of highest weight λ . We denote by Γ_{λ} the subgroup of the weight lattice Γ_{sc} of \mathfrak{g} which is generated by the (finite) set of weights P_{λ} of L_{λ} . The representation ρ_{λ} is defined over \mathbb{Z} , i.e., $L_{\lambda} = L_{\lambda}(\mathbb{Z}) \otimes \mathbb{C}$ where $L_{\lambda}(\mathbb{Z})$ is $\mathcal{U}_{\mathbb{Z}}$ -invariant. We select a highest weight vector $v_{\lambda} \in L_{\lambda}$; the lattice then is defined as $L_{\lambda}(\mathbb{Z}) = \mathcal{U}_{\mathbb{Z}} v_{\lambda}$ (see [Humphreys 1972, proof of Theorem 27.1, p. 158]). Moreover, we set $L_{\lambda}(\mathbb{Z}, \mu) = L_{\lambda}(\mathbb{Z}) \cap L_{\lambda}(\mu)$ where $L_{\lambda}(\mu) \subseteq L_{\lambda}$ is the weight μ subspace and obtain $L_{\lambda}(\mathbb{Z}) = \bigoplus_{\mu \leq \lambda} L_{\lambda}(\mathbb{Z}, \mu)$ by [Humphreys 1972, Theorem 27.1, p. 158]. More generally, for any \mathbb{Z} -algebra R we put $L_{\lambda}(R) = R \otimes L_{\lambda}(\mathbb{Z})$ and $L_{\lambda}(R, \mu) = R \otimes L_{\lambda}(\mathbb{Z}, \mu)$. The space $L_{\lambda}(R)$ is a \mathcal{U}_R -module and $L_{\lambda}(R, \mu)$ is a $\mathfrak{h}(R)$ -module and the weight decomposition of $L_{\lambda}(R)$ w.r.t. $\mathfrak{h}(R)$ reads

$$L_{\lambda}(R) = \bigoplus_{\mu \leq \lambda} L_{\lambda}(R, \mu).$$

More generally, let (π, L_{π}) be a faithful complex finite dimensional representation of \mathfrak{g} . Since $\pi = \bigoplus_i \rho_{\lambda_i}$ is semisimple (by the theorem just cited) there is a $\mathcal{U}_{\mathbb{Z}}$ -invariant lattice $L_{\pi}(\mathbb{Z})$ in L_{π} i.e., $L_{\pi} = L_{\pi}(\mathbb{Z}) \otimes \mathbb{C}$. Furthermore, for any weight $\mu \in \mathfrak{h}^*$ we set $L_{\pi}(\mathbb{Z}, \mu) = L_{\pi}(\mathbb{Z}) \cap L_{\pi}(\mu)$ and $L_{\pi}(R, \mu) = R \otimes L_{\pi}(\mathbb{Z}, \mu)$ (R a \mathbb{Z} -algebra) and obtain

$$L_{\pi}(R) = \bigoplus_{\mu \in P_{\pi}} L_{\pi}(R, \mu),$$

where $P_{\pi} \subseteq \mathfrak{h}^*$ is the set of weights of π . We note that $P_{\pi} = \bigcup_i P_{\lambda_i}$ and we set $\Gamma_{\pi} = \langle P_{\pi} \rangle$.

1.2. Chevalley groups / \mathbb{Z}_p . From now on we fix an algebraic closure $\bar{\mathbb{Q}}_p$ of \mathbb{Q}_p and we denote by $\mathcal{O}_{\bar{\mathbb{Q}}_p}$ the integer ring in $\bar{\mathbb{Q}}_p$. We recall some basic facts from the theory of Chevalley groups. Let (π, L_{π}) be a finite dimensional complex representation of \mathfrak{g} and let R be a \mathbb{Z}_p -algebra. For any $t \in R$ and any root $\alpha \in \Phi$ we define the element $x_{\alpha}(t) = x_{\alpha}^{\pi}(t) = \exp(\pi(t x_{\alpha})) \in \text{Aut}(L_{\pi}(R))$. The subgroup

$$G_{\pi, R} = \langle x_{\alpha}(t_{\alpha}), \alpha \in \Phi, t_{\alpha} \in R \rangle \leq \text{Aut}(L_{\pi}(R))$$

is called the *Chevalley group* attached to π and R .

The group $G_{\pi, \bar{\mathbb{Q}}_p}$ is a semisimple connected algebraic group, i.e., it is the set of $\bar{\mathbb{Q}}_p$ -points of an algebraic group (group scheme) \mathbf{G}_π which is defined over \mathbb{Z}_p . To make this more precise, we denote by \mathbf{GL}_n the general linear group with canonical \mathbb{Z}_p -structure $\mathbb{Z}_p[\mathbf{GL}_n] = \mathbb{Z}_p[x_{ij}, \det^{-1}]$, i.e., for any \mathbb{Z}_p -algebra R we obtain

$$\mathbf{GL}_n(R) = \text{Mor}_{\mathbb{Z}_p\text{-alg}}(\mathbb{Z}_p[\mathbf{GL}_n], R) = \{(x_{ij}) \in R^{n^2} : \det(x_{ij}) \in R^*\}.$$

We select a basis \mathcal{B} of the free \mathbb{Z}_p -module $L_\pi(\mathbb{Z}_p)$ and obtain for any \mathbb{Z}_p -algebra R an identification

$$\text{Aut}(L_\pi(R)) \stackrel{\mathcal{B}}{\cong} \mathbf{GL}_n(R) \quad (n = \dim L_\pi).$$

In particular, $G_{\pi, \bar{\mathbb{Q}}_p}$ is a subset of $\text{Aut}(L_\pi(\bar{\mathbb{Q}}_p)) = \mathbf{GL}_n(\bar{\mathbb{Q}}_p)$ and it is the set of $\bar{\mathbb{Q}}_p$ -points $\mathbf{G}_\pi(\bar{\mathbb{Q}}_p)$ of a closed algebraic subgroup $\mathbf{G}_\pi = \mathbf{G}_\pi/\mathbb{Q}_p$ of $\mathbf{GL}_n/\mathbb{Q}_p$ which is defined over \mathbb{Q}_p (see [Borel 1970, 3.3(1), p. 14 and 3.4, p. 18]); in particular, $\mathbb{Q}_p[\mathbf{G}_\pi] = \mathbb{Q}_p[\mathbf{GL}_n]/J'$ for some ideal $J' \leq \mathbb{Q}_p[\mathbf{GL}_n]$.

We set $J = J' \cap \mathbb{Z}_p[\mathbf{GL}_n]$ and $\mathbb{Z}_p[\mathbf{G}_\pi] := \mathbb{Z}_p[\mathbf{GL}_n]/J$ then is a \mathbb{Z}_p -form on $\mathbb{Q}_p[\mathbf{G}_\pi]$ which yields a \mathbb{Z}_p -structure on \mathbf{G}_π , i.e., which yields a \mathbb{Z}_p -group scheme $\mathbf{G}_\pi/\mathbb{Z}_p$ whose extension to \mathbb{Q}_p is \mathbf{G}_π (ibid., 3.4, p. 18). Thus, for any \mathbb{Z}_p -algebra R contained in $\bar{\mathbb{Q}}_p$ the group of R -points $\mathbf{G}_\pi(R)$ is defined and

$$\mathbf{G}_\pi(R) = \mathbf{G}_\pi(\bar{\mathbb{Q}}_p) \cap \mathbf{GL}_n(R) = \mathbf{G}_{\pi, \bar{\mathbb{Q}}_p} \cap \text{Aut}(L_\pi(R)).$$

In particular, since $x_\alpha(t_\alpha) \in \text{Aut}(L_\pi(R))$, $t_\alpha \in R$, we obtain $x_\alpha(t_\alpha) \in \mathbf{G}_\pi(R)$ if $t_\alpha \in R$ which yields

$$(2) \quad G_{\pi, R} \subseteq \mathbf{G}_\pi(R).$$

For each $\alpha \in \Phi$ there is a unique morphism $\mu_\alpha = \mu_\alpha^\pi : \mathbf{SL}_2(\bar{\mathbb{Q}}_p) \rightarrow \mathbf{G}_\pi(\bar{\mathbb{Q}}_p)$ such that $\begin{pmatrix} 1 & t \\ & 1 \end{pmatrix} \mapsto x_\alpha^\pi(t)$ and $\begin{pmatrix} 1 & \\ & t^{-1} \end{pmatrix} \mapsto x_{-\alpha}^\pi(t)$ ($t \in \bar{\mathbb{Q}}_p$) map (see [Borel 1970, 3.2(1), p. 13]). The μ_α is defined over \mathbb{Z}_p (ibid., 3.3(2), p. 15 and 4.3, p. 22). We denote by $h_\alpha(t) = h_\alpha^\pi(t)$ the image of $\begin{pmatrix} t & \\ & t^{-1} \end{pmatrix}$ under μ_α (ibid., 3.2(1), p. 13). The algebraic group $\mathbf{G}_\pi/\mathbb{Q}_p$ contains a \mathbb{Q}_p -split maximal torus $\mathbf{T}/\mathbb{Q}_p = \mathbf{T}^\pi/\mathbb{Q}_p$ such that the group of $\bar{\mathbb{Q}}_p$ -rational points of \mathbf{T} is given as

$$\mathbf{T}(\bar{\mathbb{Q}}_p) = \langle h_\alpha(t_\alpha), \alpha \in \Delta, t_\alpha \in \bar{\mathbb{Q}}_p^* \rangle$$

(ibid., 3.2(1), p. 13 and 3.3(3), p. 15). To any $\lambda \in \Gamma_\pi$ we attach a rational character $\lambda^\circ \in X(\mathbf{T})$ by setting

$$\lambda^\circ \left(\prod_{\alpha \in \Delta} h_\alpha(t_\alpha) \right) = \prod_{\alpha \in \Delta} t_\alpha^{\lambda(h_\alpha)}$$

for all $t_\alpha \in \bar{\mathbb{Q}}_p^*$ (ibid., 3.3, p. 15). This defines an isomorphism $\Gamma_\pi \rightarrow X(\mathbf{T})$ (ibid., 3.3(3), p. 15). We note that $d\lambda^\circ = \lambda$ (ibid., 3.3 equation (2), p. 16). The characters α° , $\alpha \in \Phi$, are the roots of $\mathbf{G}_\pi/\mathbb{Q}_p$ with respect to \mathbf{T} (i.e., the weights of the adjoint

action of T on $\text{Lie}(\mathbf{G}_\pi/\mathbb{Q}_p)$). To simplify notation we denote the exponential $\alpha^\circ : T(\bar{\mathbb{Q}}_p) \rightarrow \bar{\mathbb{Q}}_p^*$ of a root α also by α .

There are closed subgroups $N = N^\pi$ and $N^- = N^{-,\pi}$ of $\mathbf{G}_\pi/\mathbb{Q}_p$ such that

$$(3) \quad \begin{aligned} N(\bar{\mathbb{Q}}_p) &= \langle x_\alpha(t), t_\alpha \in \bar{\mathbb{Q}}_p, \alpha \in \Phi^+ \rangle, \\ N^-(\bar{\mathbb{Q}}_p) &= \langle x_\alpha(t), t_\alpha \in \bar{\mathbb{Q}}_p, \alpha \in \Phi^- \rangle. \end{aligned}$$

The subgroups N and N^- are defined over \mathbb{Q}_p and they are maximal unipotent (see [Borel 1970, 3.3(3), p. 15]). In the same way as above the \mathbb{Z}_p -structure on \mathbf{G}_π induces \mathbb{Z}_p -structures on closed subgroups H of $\mathbf{G}_\pi/\mathbb{Q}_p$ such that

$$H(\mathbb{Z}_p) = H(\bar{\mathbb{Q}}_p) \cap \mathbf{G}_\pi(\mathbb{Z}_p) = H(\bar{\mathbb{Q}}_p) \cap \text{Aut}(L_\pi(\mathbb{Z}_p)).$$

For example, this applies to the groups N, N^-, T , which thus have \mathbb{Z}_p -structures.

We set $\mathbf{B} = \mathbf{TN}$, which is a subgroup of \mathbf{G}_π defined over \mathbb{Q}_p . Thus, $\mathbf{B}(\bar{\mathbb{Q}}_p)$ is the subgroup of $\mathbf{G}_\pi(\bar{\mathbb{Q}}_p)$ which is generated by the root subgroups $\mathbf{G}_{\pi,\alpha}(\bar{\mathbb{Q}}_p)$ of $\mathbf{G}_\pi(\bar{\mathbb{Q}}_p)$ with $\alpha \in \Phi^+$ together with $T(\bar{\mathbb{Q}}_p)$ and its existence as a closed subgroup defined over \mathbb{Q}_p also follows from [Popov and Vinberg 1994, 5.3.4 Proposition, p. 70]. In particular, \mathbf{B} is a minimal parabolic subgroup. We also define the subgroup $\mathbf{B}^- = \mathbf{TN}^-$ of $\mathbf{G}_\pi/\mathbb{Q}_p$. Since $h_\alpha(t) \in \mathbf{G}_\pi(\mathbb{Z}_p)$, $t \in \mathbb{Z}_p^*$, because μ_α is defined over \mathbb{Z}_p we obtain

$$h_\alpha(t) \in T(\bar{\mathbb{Q}}_p) \cap \mathbf{G}_\pi(\mathbb{Z}_p) = T(\mathbb{Z}_p) \quad (t \in \mathbb{Z}_p^*).$$

Analogously, we obtain for any $t \in \mathbb{Z}_p$ and any positive root α

$$x_\alpha(t) \in N(\bar{\mathbb{Q}}_p) \cap \mathbf{G}_\pi(\mathbb{Z}_p) = N(\mathbb{Z}_p)$$

while for a negative root α we get $x_\alpha(t) \in N^-(\mathbb{Z}_p)$.

Notation. If $\pi = \rho_\lambda$ is an irreducible representation then we simplify notation and set $x_\alpha^\lambda(t) = x_\alpha^{\rho_\lambda}(t)$, $\mathbf{G}_\lambda = \mathbf{G}_{\rho_\lambda}$, $\mu_\alpha^\lambda = \mu_\alpha^{\rho_\lambda}$, $h_\alpha^\lambda(t) = h_\alpha^{\rho_\lambda}(t)$ and $\mathbf{T}^\lambda = \mathbf{T}^{\rho_\lambda}$; we note that in Section 1.1 we already used the notation L_λ for L_{ρ_λ} and Γ_λ for Γ_{ρ_λ} .

1.3. Mod p reduction. Let $p \in \mathbb{N}$ be a prime element. We denote by $\mathbf{G}_{\pi,(p)}$ or by $\mathbf{G}_\pi/\mathbb{F}_p$ the mod p reduction of $\mathbf{G}_\pi/\mathbb{Z}_p$; i.e., $\mathbf{G}_{\pi,(p)}(\bar{\mathbb{F}}_p)$ is the set of zeros of $\mathbb{F}_p \otimes J \leq \mathbb{F}_p \otimes \mathbb{Z}_p[\mathbf{GL}_n]$ in $\mathbf{GL}_n(\bar{\mathbb{F}}_p)$. Hence, $\mathbf{G}_{\pi,(p)}$ is an affine variety defined over \mathbb{F}_p (a closed subgroup of $\mathbf{GL}_n/\mathbb{F}_p$). Moreover, since $\mathbf{G}_\pi/\mathbb{Z}_p$ has good reduction (see [Borel 1970, 3.4, p. 18]) we know that $\mathbb{F}_p \otimes J$ equals the ideal consisting of all $f \in \mathbb{F}_p \otimes \mathbb{Z}_p[\mathbf{GL}_n]$ which vanish on $\mathbf{G}_{\pi,(p)}(\bar{\mathbb{F}}_p)$, hence, $\mathbb{F}_p[\mathbf{G}_{\pi,(p)}] = \mathbb{F}_p \otimes \mathbb{Z}_p[\mathbf{G}_\pi]$. The mod p reduction $\mathbf{G}_{\pi,(p)}$ is a semisimple group defined over \mathbb{F}_p (ibid., 4.3, p. 21/22). Analogously, the mod p reductions $N_{(p)}, \mathbf{B}_{(p)}, N_{(p)}^-, \mathbf{B}_{(p)}^-, \dots$ are defined.

We denote by

$$\wp : \text{Aut}(L_\pi(\mathbb{Z}_p)) \stackrel{\mathcal{B}}{=} \mathbf{GL}_n(\mathbb{Z}_p) \rightarrow \mathbf{GL}_n(\mathbb{F}_p) \stackrel{\bar{\mathcal{B}}}{=} \text{Aut}(L_\pi(\mathbb{F}_p))$$

the mod p reduction map which sends (x_{ij}) to $(x_{ij} \pmod p)$ ($\bar{\mathcal{B}}$ is the basis of $L_\pi(\mathbb{F}_p) = \mathbb{F}_p \otimes L_\pi(\mathbb{Z}_p)$ induced by the basis \mathcal{B} of $L_\pi(\mathbb{Z}_p)$). If $(x_{ij}) \in \mathbf{G}_\pi(\mathbb{Z}_p) \leq \mathbf{GL}_n(\mathbb{Z}_p)$ then $\wp(x)$ obviously is contained in $\mathbf{G}_\pi(\mathbb{F}_p)$, hence, \wp induces a map $\mathbf{G}_\pi(\mathbb{Z}_p) \rightarrow \mathbf{G}_\pi(\mathbb{F}_p)$.

The Iwahori subgroup \mathcal{I} of $\mathbf{G}_\pi(\mathbb{Z}_p)$ then is defined as the set of all $k \in \mathbf{G}_\pi(\mathbb{Z}_p)$ such that $\wp(k) \in \mathbf{B}^-(\mathbb{F}_p)$, i.e., $\mathcal{I} = \wp^{-1}(\mathbf{B}^-(\mathbb{F}_p))$.

1.4. Irreducible representations of $\mathbf{G}_\pi/\mathbb{Z}_p$. The group \mathbf{G}_π is a semisimple connected \mathbb{Q}_p -split group with \mathbb{Q}_p -split maximal torus $\mathbf{T} = \mathbf{T}^\pi$. Let $\lambda^\circ \in X(\mathbf{T})$ be a dominant weight. We set $\lambda = d\lambda^\circ \in \mathfrak{h}^*$. As in Section 1.2 the choice of a \mathbb{Z}_p -basis \mathcal{B} of the $\mathcal{U}_{\mathbb{Z}_p}$ -invariant lattice $L_\lambda(\mathbb{Z}_p) (= L_{\rho_\lambda}(\mathbb{Z}_p))$ yields an identification $\text{Aut}(L_\lambda(\bar{\mathbb{Q}}_p)) \stackrel{\mathcal{B}}{=} \mathbf{GL}_m(\bar{\mathbb{Q}}_p)$ (where $m = \dim(L_\lambda)$).

If $\Gamma_\pi \supseteq \Gamma_\lambda$ we define a representation of algebraic groups

$$\rho_{\lambda^\circ} : \mathbf{G}_\pi(\bar{\mathbb{Q}}_p) \rightarrow \text{Aut}(L_\lambda(\bar{\mathbb{Q}}_p)) = \mathbf{GL}_m(\bar{\mathbb{Q}}_p)$$

by mapping $x_\alpha^\pi(t)$ to $x_\alpha^\lambda(t)$, $\alpha \in \Phi$, $t \in \bar{\mathbb{Q}}_p$ (see [Borel 1970, 3.2(4), p. 14 and 3.3(2), p. 15], where ρ_{λ° is denoted by $\lambda_{\rho_\lambda, \pi}$). We note that ρ_{λ° has image $\mathbf{G}_\lambda(\bar{\mathbb{Q}}_p)$ (which equals $\mathbf{G}_{\rho_\lambda}(\bar{\mathbb{Q}}_p)$).

Lemma. *The map ρ_{λ° induces a map of tori*

$$\rho_{\lambda^\circ} : \mathbf{T}^\pi(\bar{\mathbb{Q}}_p) \rightarrow \mathbf{T}^\lambda(\bar{\mathbb{Q}}_p)$$

which maps $h_\alpha^\pi(t) \mapsto h_\alpha^\lambda(t)$ for all $\alpha \in \Delta$ and $t \in \bar{\mathbb{Q}}_p^*$.

Proof. We first claim that for each $\alpha \in \Phi$ the diagram

$$\begin{array}{ccc} \mathbf{G}_\pi(\bar{\mathbb{Q}}_p) & \xrightarrow{\rho_{\lambda^\circ}} & \mathbf{G}_\lambda(\bar{\mathbb{Q}}_p) \\ & \swarrow \mu_\alpha^\pi & \nearrow \mu_\alpha^\lambda \\ & \mathbf{SL}_2(\bar{\mathbb{Q}}_p) & \end{array}$$

commutes. It is sufficient to show commutativity of the diagram for all $\begin{pmatrix} 1 & t \\ & 1 \end{pmatrix}$ and $\begin{pmatrix} 1 & \\ t & 1 \end{pmatrix}$ with $t \in \bar{\mathbb{Q}}_p$ because these elements generate $\mathbf{SL}_2(\bar{\mathbb{Q}}_p)$. But

$$\rho_{\lambda^\circ}(\mu_\alpha^\pi(\begin{pmatrix} 1 & t \\ & 1 \end{pmatrix})) = \rho_{\lambda^\circ}(x_\alpha^\pi(t)) = x_\alpha^\lambda(t) = \mu_\alpha^\lambda(\begin{pmatrix} 1 & t \\ & 1 \end{pmatrix}),$$

and analogously for the lower unipotent matrices. Hence, the diagram commutes and we obtain

$$\rho_{\lambda^\circ}(h_\alpha^\pi(t)) = \rho_{\lambda^\circ}(\mu_\alpha^\pi(\begin{pmatrix} t & \\ & t^{-1} \end{pmatrix})) = \mu_\alpha^\lambda(\begin{pmatrix} t & \\ & t^{-1} \end{pmatrix}) = h_\alpha^\lambda(t).$$

Since $\mathbf{T}^\pi(\bar{\mathbb{Q}}_p)$ is generated by the $h_\alpha^\pi(t)$, where $\alpha \in \Delta$ and $t \in \bar{\mathbb{Q}}_p^*$, this implies that $\rho_{\lambda^\circ}(\mathbf{T}^\pi(\bar{\mathbb{Q}}_p)) \subseteq \mathbf{T}^\lambda(\bar{\mathbb{Q}}_p)$ and the lemma is proven. \square

The lemma implies that for all $t \in T^\pi(\bar{\mathbb{Q}}_p)$ and any weight vector $v_\mu \in L_\lambda(\bar{\mathbb{Q}}_p, \mu)$, $\mu \in P_\lambda$, (i.e., v_μ has weight μ w.r.t. \mathfrak{h}) we have

$$(4) \quad \rho_{\lambda^\circ}(t)(v_\mu) = \mu^\circ(t)v_\mu.$$

In fact since $T^\pi(\bar{\mathbb{Q}}_p)$ is generated by the $h_\alpha^\pi(s)$ we may assume that $t = h_\alpha^\pi(s)$ for some $\alpha \in \Delta$ and $s \in \bar{\mathbb{Q}}_p^*$. Using the lemma and the equation in [Borel 1970, 3.2(1), p. 13], we obtain

$$\rho_{\lambda^\circ}(t)(v_\mu) = \rho_{\lambda^\circ}(h_\alpha^\pi(s))(v_\mu) = h_\alpha^\lambda(s)v_\mu = s^{\mu(h_\alpha)}v_\mu = \mu^\circ(t)v_\mu.$$

The following result seems to be well known. Since we could not find a direct reference we add a proof.

- Proposition.** 1. *The $G_\pi(\bar{\mathbb{Q}}_p)$ -module $L_\lambda(\bar{\mathbb{Q}}_p)$ contains a vector v_{λ° which is invariant under $N^\pi(\bar{\mathbb{Q}}_p)$ and satisfies $tv_{\lambda^\circ} = \lambda^\circ(t)v_{\lambda^\circ}$ for all $t \in T^\pi(\bar{\mathbb{Q}}_p)$.*
2. *The representation ρ_{λ° is the irreducible representation of $G_\pi(\bar{\mathbb{Q}}_p)$ of highest weight λ° .*

Proof. 1. We choose for v_{λ° the highest weight vector $v_\lambda \in L_\lambda(\mathbb{Z})$, which we selected in Section 1.1. Since $x_\alpha v_{\lambda^\circ} (= \rho_\lambda(x_\alpha)v_{\lambda^\circ})$ vanishes for all $\alpha \in \Phi^+$ we obtain

$$\rho_{\lambda^\circ}(x_\alpha^\pi(t))v_{\lambda^\circ} = x_\alpha^\lambda(t)v_{\lambda^\circ} = v_{\lambda^\circ} + t\rho_\lambda(x_\alpha)(v_{\lambda^\circ}) + \dots = v_{\lambda^\circ}.$$

Since $N^\pi(\bar{\mathbb{Q}}_p)$ is generated by the $x_\alpha^\pi(t)$ with $\alpha \in \Phi^+$, $t \in \bar{\mathbb{Q}}_p$, this yields the first claim about v_{λ° . The second claim is immediate by equation (4) since $v_{\lambda^\circ} = v_\lambda$ has \mathfrak{h} -weight λ .

2. For the moment we denote by $(\sigma_{\mu^\circ}, \Sigma_{\mu^\circ})$ the irreducible representation of $G_\pi(\bar{\mathbb{Q}}_p)$ of highest weight $\mu^\circ \in X(T^\pi)$. The derived representation of σ_{μ° is $(\rho_\mu, L_\mu(\bar{\mathbb{Q}}_p))$, where $\mu = d\mu^\circ$; hence, $\dim \Sigma_{\mu^\circ} = \dim L_\mu(\bar{\mathbb{Q}}_p)$. Since G_π is semisimple and since we are in characteristic 0 any representation of $G_\pi(\bar{\mathbb{Q}}_p)$ is semisimple, hence, we can write

$$\rho_{\lambda^\circ} = \bigoplus_{i=1}^r \Sigma_{\mu_i^\circ}.$$

Any representation $\Sigma_{\mu_i^\circ}$ contains a unique (up to scalars) nontrivial vector $v_{\mu_i^\circ}$ invariant under $N^\pi(\bar{\mathbb{Q}}_p)$. This vector $v_{\mu_i^\circ}$ then satisfies $tv_{\mu_i^\circ} = \mu_i^\circ(t)v_{\mu_i^\circ}$, $t \in T^\pi(\bar{\mathbb{Q}}_p)$ (i.e., $\bar{\mathbb{Q}}_p v_{\mu_i^\circ}$ is the unique line which is stable under $B(\bar{\mathbb{Q}}_p)$). The vector v_{λ° decomposes as

$$v_{\lambda^\circ} = \sum_{i=1}^r v_i,$$

where $v_i \in \Sigma_{\mu_i^\circ}$ and at least one vector v_j does not vanish. Since v_{λ° is invariant under $N(\bar{\mathbb{Q}}_p)$ by part 1, we obtain $\sum_i n v_i = \sum_i v_i$ for any $n \in N(\bar{\mathbb{Q}}_p)$, hence, $n v_i = v_i$

for all i and all $n \in N(\bar{\mathbb{Q}}_p)$. Thus, $v_i = c_i v_{\mu_i^\circ}$ for some $c_i \in \bar{\mathbb{Q}}_p$ by the uniqueness of $v_{\mu_i^\circ}$; in particular, v_i has weight μ_i° w.r.t. $T^\pi(\bar{\mathbb{Q}}_p)$. On the other hand, since $tv_{\lambda^\circ} = \lambda^\circ(t)v_{\lambda^\circ}$ by part 1, we obtain $\sum_i tv_i = \sum_i \lambda^\circ(t)v_i$; hence, $tv_i = \lambda^\circ(t)v_i$ for all i and $t \in T^\pi(\bar{\mathbb{Q}}_p)$. Since $v_j \neq 0$ we deduce that $\mu_j^\circ = \lambda^\circ$. The representation ρ_{λ° therefore decomposes as a direct sum $\rho_{\lambda^\circ} = \Sigma_{\lambda^\circ} \oplus C$. Since ρ_{λ° is a representation on the space $L_\lambda(\bar{\mathbb{Q}}_p)$ we know that $\dim \rho_{\lambda^\circ} = \dim L_\lambda(\bar{\mathbb{Q}}_p) = \dim \Sigma_{\lambda^\circ}$. This implies that $C = 0$, hence, $\rho_{\lambda^\circ} = \Sigma_{\lambda^\circ}$ is the irreducible representation of $G_\pi(\bar{\mathbb{Q}}_p)$ of highest weight λ° . Thus, the proof is complete. \square

From [Borel 1970, 3.5, p. 19], we know that the morphism ρ_{λ° is defined over \mathbb{Z}_p , i.e., it is associated to a morphism of \mathbb{Z}_p -group schemes $\rho_{\lambda^\circ} : G_\pi/\mathbb{Z}_p \rightarrow G_\lambda/\mathbb{Z}_p$. Since G_λ is a closed subscheme of $\mathbf{Aut}(L_\lambda) = \mathbf{GL}_m/\mathbb{Z}_p$ we obtain that the representation ρ_{λ° is defined over \mathbb{Z}_p , i.e.,

$$\rho_{\lambda^\circ} : G_\pi/\mathbb{Z}_p \rightarrow \mathbf{Aut}(L_\lambda) = \mathbf{GL}_m/\mathbb{Z}_p$$

In particular, $L_\lambda(R)$ is a $G_\pi(R)$ -module for all \mathbb{Z}_p -algebras R . Using equation (4) we deduce that $T^\pi(\mathbb{Z}_p)$ leaves $L_\lambda(\mathbb{Z}_p, \mu) = L_\lambda(\mathbb{Z}_p) \cap L_\lambda(\bar{\mathbb{Q}}_p, \mu)$ invariant and acts via the character μ° .

1.5. The level subgroup $K_*(p, \sigma)$. From now on we fix a prime element $p \in \mathbb{N}$. For any $\sigma \in \mathbb{N}$ we define the level subgroup

$$K_*(\sigma) = K_*(p, \sigma) = K_*^\pi(p, \sigma) \leq G_\pi(\mathbb{Z}_p)$$

as the subgroup generated by the following elements: all $x_\alpha(t_\alpha)$ with $\alpha \in \Phi^-$ and $t_\alpha \in \mathbb{Z}_p$, all $x_\alpha(t_\alpha)$ with $\alpha \in \Phi^+$ and $t_\alpha \in p^{\lceil \frac{1}{\sigma} \text{ht}(\alpha) \rceil} \mathbb{Z}_p$ and all $h_\alpha^\pi(t_\alpha)$ with $\alpha \in \Delta$ and $t_\alpha \in \mathbb{Z}_p^*$. We note that the equations at the end of Section 1.2 imply that $K_*(\sigma) \leq G_\pi(\mathbb{Z}_p)$ and even that $K_*(\sigma) \leq \mathcal{I}$, because $\wp(x_\alpha(t))$ is the identity in $\mathbf{Aut}(L_\pi(\mathbb{F}_p))$ if $t \in p\mathbb{Z}_p$. If $\sigma \geq \max_{\alpha \in \Phi^+} \text{ht}(\alpha)$ we conclude that $K_*(\sigma)$ equals

$$\langle x_\alpha(t_\alpha), t_\alpha \in p\mathbb{Z}_p \text{ if } \alpha > 0 \text{ and } t_\alpha \in \mathbb{Z}_p \text{ if } \alpha < 0, h_\alpha(t_\alpha), \alpha \in \Delta, t_\alpha \in \mathbb{Z}_p^* \rangle = \mathcal{I};$$

the latter equality follows from [Iwahori and Matsumoto 1965, p. 259] (the Iwahori subgroup is denoted by B there).

We define the subgroups $N_{\mathbb{Z}_p} = N_{\mathbb{Z}_p}^\pi := \langle x_\alpha(t_\alpha), \alpha > 0, t_\alpha \in \mathbb{Z}_p \rangle \subseteq N(\mathbb{Z}_p)$, $N_{p\mathbb{Z}_p} = N_{p\mathbb{Z}_p}^\pi := \langle x_\alpha(t_\alpha), \alpha > 0, t_\alpha \in p\mathbb{Z}_p \rangle$, $N_{\mathbb{Z}_p}^- = N_{\mathbb{Z}_p}^{-,\pi} := \langle x_\alpha(t_\alpha), \alpha < 0, t_\alpha \in \mathbb{Z}_p \rangle$ and $T_{\mathbb{Z}_p} = T_{\mathbb{Z}_p}^\pi := \langle h_\alpha^\pi(t_\alpha), \alpha \in \Delta, t_\alpha \in \mathbb{Z}_p^* \rangle$ of $G_\pi(\mathbb{Z}_p)$. The Iwahori subgroup then satisfies the decomposition

$$\mathcal{I} = N_{p\mathbb{Z}_p} T_{\mathbb{Z}_p} N_{\mathbb{Z}_p}^-$$

(see [Iwahori and Matsumoto 1965, Theorem 2.5, p. 263]; note that $T_{\mathbb{Z}_p} = \mathfrak{h}_\mathcal{D}$). We note the following consequences.

1. Assume $h \in \mathbf{T}(\mathbb{Q}_p)^{++}$; i.e., $v_p(\alpha(h)) > 0$ for all simple roots $\alpha \in \Delta$. Then, for all $e, f \in \mathbb{N}_0$ we have

$$(5) \quad \mathcal{I}h^e \mathcal{I}h^f \mathcal{I} = \mathcal{I}h^{e+f} \mathcal{I}.$$

2. Assume $t, t' \in \mathbf{T}(\mathbb{Q}_p)^{++}$. Then

$$(6) \quad \mathcal{I}t\mathcal{I} = \mathcal{I}t'\mathcal{I} \iff T_{\mathbb{Z}_p}tT_{\mathbb{Z}_p} = T_{\mathbb{Z}_p}t'T_{\mathbb{Z}_p}.$$

Proof of equation (6). The leftward implication is trivial. To prove the reverse implication we note that $t \in \mathcal{I}t'\mathcal{I}$ implies that there are $k^+, m^+ \in N_{p\mathbb{Z}_p}, k^\circ, m^\circ \in T_{\mathbb{Z}_p}$ and $k^-, m^- \in N_{\mathbb{Z}_p}^-$ such that $tk^+k^\circ k^- = m^+m^\circ m^-t'$. Since $\text{Ad}(t)(x_\alpha(t_\alpha)) = x_\alpha(\alpha(t)t_\alpha)$ the element t normalizes $N_{p\mathbb{Z}_p}$ and $(t')^{-1}$ normalizes $N_{\mathbb{Z}_p}^-$, hence, we obtain $\tilde{k}^+tk^\circ k^- = m^+m^\circ t'\tilde{m}^-$ with $\tilde{k}^+ \in N_{p\mathbb{Z}_p}, \tilde{m}^- \in N_{\mathbb{Z}_p}^-$. Equivalently,

$$(7) \quad (m^+)^{-1}\tilde{k}^+tk^\circ = m^\circ t'\tilde{m}^-(k^-)^{-1}.$$

The left-hand side is contained in $\mathbf{B}(\mathbb{Q}_p)$ and the right-hand side is contained in $\mathbf{B}^-(\mathbb{Q}_p)$, whose intersection is $\mathbf{T}(\mathbb{Q}_p)$. Hence, $(m^+)^{-1}\tilde{k}^+ \in \mathbf{T}(\mathbb{Q}_p) \cap N_{p\mathbb{Z}_p} = \{1\}$ (note that $N_{p\mathbb{Z}_p} \subseteq N_{\mathbb{Z}_p} \subseteq \mathbf{N}(\mathbb{Z}_p)$) and, similarly, $\tilde{m}^-(k^-)^{-1} \in \mathbf{T}(\mathbb{Q}_p) \cap N_{\mathbb{Z}_p}^- = \{1\}$. Equation (7) thus implies that $tk^\circ = m^\circ t'$, which proves the claim. \square

2. Hecke algebra and cohomology

2.1. Reductive algebraic groups. From now on, $\tilde{\mathbf{G}}$ denotes a connected reductive algebraic group defined over \mathbb{Q} . Since $\tilde{\mathbf{G}}$ is defined over \mathbb{Q} it contains a maximal torus which is defined over \mathbb{Q} and we assume that $\tilde{\mathbf{G}}$ contains a maximal torus $\tilde{\mathbf{T}}$ which is defined over \mathbb{Q} and split over \mathbb{Q}_p (hence, $\tilde{\mathbf{G}}$ is \mathbb{Q}_p -split). This assumption is in particular satisfied if $\tilde{\mathbf{G}}$ is \mathbb{Q} -split. We denote by $\mathbf{G} = \tilde{\mathbf{G}}^{\text{der}}$ the derived group and by $\tilde{\mathbf{Z}}$ the center of $\tilde{\mathbf{G}}$; hence, $\tilde{\mathbf{G}} = (\mathbf{G} \times \tilde{\mathbf{Z}})/\mathbf{Z}$ as algebraic groups over \mathbb{Q} , where \mathbf{Z} is the center of \mathbf{G} (embedded via $z \mapsto (z, z^{-1})$). We denote by $\text{Lie}(\mathbf{G})$ the Lie algebra of \mathbf{G} . We use the notations introduced in Section 1.1 for the complex Lie algebra $\mathfrak{g} = \text{Lie}(\mathbf{G}) \otimes_{\mathbb{Q}} \mathbb{C}$; e.g., \mathfrak{h} is a Cartan subalgebra in \mathfrak{g} , $\Phi = \Phi(\mathfrak{g}, \mathfrak{h})$ the set of roots and Δ a choice of a basis of Φ . Since \mathbf{G} is a \mathbb{Q}_p -split, semisimple algebraic group, there is a finite dimensional complex representation π of \mathfrak{g} such that

$$\mathbf{G}/\mathbb{Q}_p \cong \mathbf{G}_\pi/\mathbb{Q}_p$$

as \mathbb{Q}_p -groups, where $\mathbf{G}_\pi/\mathbb{Z}_p$ is the Chevalley group attached to π ; see Section 1.2. In the following we may assume that $\mathbf{G}/\mathbb{Q}_p = \mathbf{G}_\pi/\mathbb{Q}_p$.

The \mathbb{Q}_p -structure on $\tilde{\mathbf{G}}$. We denote by \mathbf{T} the \mathbb{Q}_p -split maximal torus and by \mathbf{N}, \mathbf{N}^- the maximal unipotent subgroups in $\mathbf{G}/\mathbb{Q}_p = \mathbf{G}_\pi/\mathbb{Q}_p$ defined in Section 1.2. The

subgroups N, N^- remain maximal unipotent in \tilde{G} and T is a \mathbb{Q}_p -defined and \mathbb{Q}_p -split torus in \tilde{G} . We denote by \tilde{B}/\mathbb{Q}_p the Borel subgroup in \tilde{G} containing \tilde{T} which corresponds to Δ . The torus \tilde{T} decomposes $\tilde{T} = (T' \times \tilde{Z})/Z$ over \mathbb{Q}_p , where T'/\mathbb{Q}_p is a \mathbb{Q}_p -split maximal torus in G/\mathbb{Q}_p . Since any two \mathbb{Q}_p -split maximal tori in G/\mathbb{Q}_p are conjugate by an element $x \in G(\mathbb{Q}_p)$ (see [Springer 1981, 15.2.6 Theorem, p. 256]) we may assume after composing the isomorphism $G/\mathbb{Q}_p \cong G_\pi/\mathbb{Q}_p$ with conjugation by x that $T = T'$, hence, $\tilde{T} = (T \times \tilde{Z})/Z$ as algebraic groups over \mathbb{Q}_p . We denote by $X(\tilde{T})$ (resp. $X_*(\tilde{T})$) the (additively written) group of \mathbb{Q}_p -characters (resp. \mathbb{Q}_p -cocharacters) of \tilde{T}/\mathbb{Q}_p and by $\langle \cdot, \cdot \rangle : X(\tilde{T}) \times X_*(\tilde{T}) \rightarrow \mathbb{Z}$ the canonical pairing defined by $\chi \circ \eta(x) = x^{\langle \chi, \eta \rangle}$ for all $x \in \mathbb{G}_m(\bar{\mathbb{Q}}_p)$. We recall that by α we denote a root in $\Phi \subseteq \mathfrak{h}^*$ and also its exponential in $X(T)$ (i.e., we write α for α° ; see Section 1.2). Any root $\alpha \in X(T)$ vanishes on the center Z of G , hence, it extends to a character on $\tilde{T} = (T \times \tilde{Z})/Z$ by setting it equal to 1 on \tilde{Z} ; we denote this extension of the root again by α ; hence, $\alpha(t) = \alpha(t^\circ) (= \alpha^\circ(t^\circ))$ if $t = t^\circ z \in T(\bar{\mathbb{Q}}_p)\tilde{Z}(\bar{\mathbb{Q}}_p)$. We denote by $\alpha^\vee \in X_*(T) \subseteq X_*(\tilde{T})$ the coroot corresponding to α ; explicitly, $\alpha^\vee(t) = h_\alpha(t)$, $t \in \mathbb{G}_m(\bar{\mathbb{Q}}_p)$. Any character $\tilde{\lambda} \in X(\tilde{T})$ is of the form $\tilde{\lambda} = \lambda^\circ \otimes \kappa$, where $\kappa = \tilde{\lambda}|_{\tilde{Z}} \in X(\tilde{Z})$ and $\lambda^\circ = \tilde{\lambda}|_T \in X(T)$ satisfy $\lambda^\circ|_Z = \kappa|_Z$. We note that λ° corresponds to a weight $\lambda \in \Gamma_\pi$, i.e., $\lambda = d\lambda^\circ$; see Section 1.2. We call $\tilde{\lambda} \in X(\tilde{T})$ dominant if

$$\langle \tilde{\lambda}, \alpha^\vee \rangle = \langle \lambda^\circ, \alpha^\vee \rangle = \lambda(h_\alpha) \geq 0$$

for all $\alpha \in \Delta$. We denote by $\tilde{T}(\mathbb{Q}_p)^+$ (resp. $\tilde{T}(\mathbb{Q}_p)^{++}$, $\tilde{T}(\mathbb{Q}_p)^{--}$) the set of all elements $t \in \tilde{T}(\mathbb{Q}_p)$ such that $v_p(\alpha(t)) \geq 0$ (resp. $v_p(\alpha(t)) > 0$, $v_p(\alpha(t)) < 0$) for all $\alpha \in \Delta$ and by $X(\tilde{T})^{\text{dom}}$ the set of dominant characters.

The \mathbb{Z}_p -structure on \tilde{G} . We recall that the derived group $G/\mathbb{Q}_p = G_\pi/\mathbb{Q}_p$ has a \mathbb{Z}_p -structure; see Section 1.2. In Section 1.5 we defined the level subgroup $K_*(\sigma) = K_*^\pi(p, \sigma)$ which is a subgroup of $G(\mathbb{Z}_p)$. We define a \mathbb{Z}_p -structure on \tilde{Z} by selecting as a \mathbb{Z}_p -form of $\mathbb{Q}_p[\tilde{Z}]$ the algebra $\mathbb{Z}_p[\text{res}_{\tilde{T}/\tilde{Z}} X(\tilde{T})] = \mathbb{Z}_p[X(\tilde{T})/X'(\tilde{T})]$, where $X'(\tilde{T}) = X(\tilde{T}) \cap \sum_{\alpha \in \Phi} \mathbb{Q}\alpha$. It follows that for any $\tilde{\lambda} \in X(\tilde{T})$

$$(8) \quad \tilde{\lambda}|_{\tilde{Z}} : \tilde{Z} \rightarrow \mathbb{G}_m$$

is defined over \mathbb{Z}_p . The \mathbb{Z}_p -structures on G and \tilde{Z} yield a \mathbb{Z}_p -structure on \tilde{G} .

2.2. From now on, we fix a prime $p \in \mathbb{N}$ and we define the subgroup

$$\tilde{\mathcal{I}} := \langle \mathcal{I}, \tilde{Z}(\mathbb{Z}_p) \rangle \leq \tilde{G}(\mathbb{Z}_p).$$

We select an arithmetic subgroup $\Gamma \leq \tilde{G}(\mathbb{Q})$ satisfying $\Gamma \leq \tilde{\mathcal{I}}$.

2.3. The Hecke algebra. Also, from now on, we let h be an element in $\tilde{T}(\mathbb{Q})^{++}$; i.e., $h \in \tilde{T}(\mathbb{Q})$ and $v_p(\alpha(h)) > 0$ for all $\alpha \in \Delta$. We denote by $\mathcal{K} = \mathcal{K}_h = \langle h, \tilde{\mathcal{I}} \rangle_{\text{semigrp}} \leq$

$\tilde{\mathbf{G}}(\mathbb{Q}_p)$ the (sub)semigroup of $\tilde{\mathbf{G}}(\mathbb{Q}_p)$ which is generated by h and $\tilde{\mathbf{T}}$ and we set

$$\Delta = \Delta_h = \{g \in \tilde{\mathbf{G}}(\mathbb{Q}) : g \in \mathcal{K}\}.$$

Thus, $\Delta \leq \tilde{\mathbf{G}}(\mathbb{Q})$ is a subsemigroup containing Γ and we denote by

$$\mathcal{H} = \mathcal{H}_h = \mathcal{H}(\Gamma \backslash \Delta / \Gamma)$$

the Hecke algebra attached to the pair (Δ, Γ) . Thus, \mathcal{H} is a \mathbb{Z} -algebra which is a free \mathbb{Z} -module with basis $\{\Gamma \zeta \Gamma, \zeta \in \Delta\}$. For any \mathbb{Z} -algebra R we set $\mathcal{H}_R = \mathcal{H} \otimes R$ and we put

$$T_\zeta = \Gamma \zeta \Gamma \in \mathcal{H} \quad (\zeta \in \Delta).$$

2.4. Irreducible representations of $\tilde{\mathbf{G}}$. Let $\tilde{\lambda} \in X(\tilde{\mathbf{T}})$ be dominant. We set $\lambda^\circ = \tilde{\lambda}|_T \in X(T)$ and $\lambda = d\lambda^\circ \in \mathfrak{h}^*$. Since $X(T) \cong \Gamma_\pi$ (see Section 1.2) we know that $\lambda \in \Gamma_\pi$ and we let $\rho_{\lambda^\circ} : \mathbf{G}_\pi / \mathbb{Z}_p \rightarrow \mathbf{Aut}(L_\lambda)$ be the irreducible representation of $\mathbf{G} / \mathbb{Z}_p = \mathbf{G}_\pi / \mathbb{Z}_p$ of highest weight λ° ; see Section 1.4. The morphism $\rho_{\lambda^\circ} \otimes \tilde{\lambda}|_{\tilde{\mathbf{Z}}} : \mathbf{G} \times \tilde{\mathbf{Z}} \rightarrow \mathbf{Aut}(L_\lambda)$, given by sending $(g, z) \in \mathbf{G}(R) \times \tilde{\mathbf{Z}}(R)$ to $\tilde{\lambda}(z)\rho_{\lambda^\circ}(g) \in \mathbf{Aut}(L_\lambda(R))$, R any \mathbb{Z}_p -algebra, is defined over \mathbb{Z}_p (see equation (8)) and factorizes over \mathbf{Z} , hence, we obtain a representation

$$\rho_{\tilde{\lambda}} : \tilde{\mathbf{G}} = (\mathbf{G} \times \tilde{\mathbf{Z}}) / \mathbf{Z} \rightarrow \mathbf{Aut}(L_\lambda)$$

of \mathbb{Z}_p -groups (group schemes). The representation $(\rho_{\tilde{\lambda}}, L_\lambda)$ is irreducible of highest weight $\tilde{\lambda}$ and $\rho_{\lambda^\circ} = \rho_{\tilde{\lambda}}|_{\mathbf{G}}$. In particular, for any \mathbb{Z}_p -algebra R the $\mathbf{G}(R)$ -module $L_\lambda(R)$ also is a $\tilde{\mathbf{G}}(R)$ -module and we write $L_{\tilde{\lambda}}(R)$ for $L_\lambda(R)$ if we view it as $\tilde{\mathbf{G}}(R)$ -module. Thus, $L_{\tilde{\lambda}}(R)$ and $L_\lambda(R)$ are isomorphic as $\mathbf{G}(R)$ -modules, but on $L_{\tilde{\lambda}}(R)$ we have an action of $\tilde{\mathbf{Z}}(R)$ via $\tilde{\lambda}|_{\tilde{\mathbf{Z}}}$ and, hence, an action of $\tilde{\mathbf{T}}(R)$. Similarly, we obtain a representation

$$\tilde{\mathbf{T}} = (\mathbf{T} \times \tilde{\mathbf{Z}}) / \mathbf{Z} \rightarrow \mathbf{Aut}(L_\lambda(\mu)), \quad \mu \in P_\lambda,$$

by sending $(t, z) \in \mathbf{T}(R) \times \tilde{\mathbf{Z}}(R)$ to $\tilde{\lambda}(z)\rho_{\lambda^\circ}(t) \in \mathbf{Aut}(L_\lambda(\mu, R))$, R any \mathbb{Z}_p -algebra (note that $L_\lambda(\mathbb{Z}_p, \mu)$ is a \mathbb{Z}_p -module, hence, $\mathbf{Aut}(L_\lambda(\mu))$ is a \mathbb{Z}_p -group). If we view the weight space $L_\lambda(R, \mu)$ as $\tilde{\mathbf{T}}(R)$ -module we write it as $L_{\tilde{\lambda}}(R, \mu)$. Thus, $L_{\tilde{\lambda}}(R, \mu) = L_\lambda(R, \mu)$ as abelian groups and also as $\mathbf{T}(R)$ -modules but on $L_{\tilde{\lambda}}(R, \mu)$ the torus $\tilde{\mathbf{T}}(R)$ acts via the character $\tilde{\mu} := \mu^\circ \otimes \tilde{\lambda}|_{\tilde{\mathbf{Z}}}$ of $\tilde{\mathbf{T}}$ (see Section 1.4 and equation (4) in particular). The weight decomposition of $L_{\tilde{\lambda}}(R)$ w.r.t. $\tilde{\mathbf{T}}(R)$ then reads

$$(9) \quad L_{\tilde{\lambda}}(R) = \bigoplus_{\mu \leq \lambda} L_{\tilde{\lambda}}(R, \mu),$$

where $L_{\tilde{\lambda}}(R, \mu)$ is the weight $\tilde{\mu}$ -subspace of $L_{\tilde{\lambda}}(R)$ w.r.t. $\tilde{\mathbf{T}}(R)$.

2.5. Splitting field. Since the maximal torus \tilde{T}/\mathbb{Q} is assumed to be split over \mathbb{Q}_p there is a subfield $F \subseteq \mathbb{Q}_p$ which is a finite extension of \mathbb{Q} such that \tilde{T}/F is split. In particular, \tilde{G} is F -split and $\tilde{\lambda} \in X(\tilde{T})$ and the irreducible highest weight representation $(\rho_{\tilde{\lambda}}, L_{\tilde{\lambda}})$ are defined over F ; hence, $L_{\tilde{\lambda}}(F)$ is defined and is a $\tilde{G}(\mathbb{Q})$ -module.

We fix an algebraic closure $\bar{\mathbb{Q}}_p$ of \mathbb{Q}_p with valuation v_p normalized by $v_p(p) = 1$. Since $F \subseteq \mathbb{Q}_p \subseteq \bar{\mathbb{Q}}_p$ this induces a p -adic valuation v_p on F and we obtain $F_{v_p} = \mathbb{Q}_p$. We also fix an embedding $F \subseteq \mathbb{C}$. We extend the embeddings of F to embeddings of its algebraic closure $\bar{F} \subseteq \bar{\mathbb{Q}}_p$ and $\bar{F} \subseteq \mathbb{C}$; hence, we may view F as a subfield of \mathbb{Q}_p and of \mathbb{C} and \bar{F} as a subfield of \mathbb{C} and of $\bar{\mathbb{Q}}_p$.

2.6. Cohomology with coefficients $L_{\tilde{\lambda}}$. We denote by $\Delta^{-1} \leq \tilde{G}(\mathbb{Q})$ the sub semi-group consisting of the inverses of elements in Δ . The representation space $L_{\tilde{\lambda}}(F)$ in particular is a Δ^{-1} -module, hence, the Hecke algebra \mathcal{H} acts on cohomology $H^i(\Gamma, L_{\tilde{\lambda}}(F))$. For later use we recall the definition of this action. Let $T_\zeta = \Gamma \zeta \Gamma \in \mathcal{H}$ ($\zeta \in \Delta$). We select a system of representatives $\gamma_1, \dots, \gamma_r$ for $(\zeta^{-1} \Gamma \zeta \cap \Gamma) \backslash \Gamma$, hence,

$$T_\zeta = \bigcup_{i=1, \dots, r} \Gamma \zeta \gamma_i.$$

Thus, for any $\eta \in \Gamma$ and any index i satisfying $1 \leq i \leq r$ there is an index $\eta(i)$ such that

$$\Gamma \zeta \gamma_i \eta = \Gamma \zeta \gamma_{\eta(i)}.$$

In particular, there are $\rho_i(\eta) \in \Gamma$, $i = 1, \dots, r$, such that $\zeta \gamma_i \eta = \rho_i(\eta) \zeta \gamma_{\eta(i)}$. Let now $c \in C^d(\Gamma, L_{\tilde{\lambda}}(F))$ be any cochain; we then define $T_\zeta(c)$ as the cochain $c' \in C^d(\Gamma, L_{\tilde{\lambda}}(F))$, which is given by

$$(10) \quad c'(\eta_0, \dots, \eta_d) = \sum_{1 \leq i \leq r} (\zeta \gamma_i)^{-1} c(\rho_i(\eta_0), \dots, \rho_i(\eta_d)).$$

Since T_ζ commutes with the coboundary operator, T_ζ acts on cohomology with coefficients in $L_{\tilde{\lambda}}(F)$, i.e., T_ζ defines an element in $\text{End}(H^i(\Gamma, L_{\tilde{\lambda}}(F)))$ which does not depend on the choice of the representatives $\gamma_1, \dots, \gamma_r$ (see [Kuga et al. 1981, p. 227]). We note that this also yields \mathcal{H} -module structures on $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))$ and $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{C}))$. We denote by

$$H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}}$$

the image of the canonical mapping $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p)) \rightarrow H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))$; this defines a lattice in $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))$.

Cuspidal cohomology. We select a maximal compact open subgroup $\tilde{K}_\infty \leq \tilde{G}(\mathbb{R})$. We denote by $A_{\tilde{G}}$ the connected component of the real points of a maximal \mathbb{Q} -split

torus $A_{\tilde{G}}$ in the center of \tilde{G} and we set $X = \tilde{G}(\mathbb{R})/\tilde{K}_{\infty}A_{\tilde{G}}$. The cuspidal cohomology $H_{\text{cusp}}^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))$ is a subspace of full cohomology $H^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))$. We note that if the highest weight $\tilde{\lambda}$ is regular and \tilde{G} has discrete series then there are isomorphisms

$$H_{(2)}^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C})) = H_!^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C})) = H_{\text{cusp}}^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))$$

and cuspidal cohomology vanishes in all degrees except for the middle degree $d = d_{\tilde{G}} = \frac{1}{2} \dim X$.

Weighted cohomology. For use in Section 4.14 involving the Goresky–MacPherson trace formula we briefly recall the relation between cuspidal cohomology and weighted cohomology. We denote by $W^{\nu} H^i(\overline{\Gamma \backslash X}, L_{\tilde{\lambda}}(F))$ the weighted cohomology groups of Γ (see [Goresky et al. 1994]). If ν is the middle weight profile and \tilde{G} has discrete series then [Nair 1999, Corollary B, p. 3] (see also Section 5.1 there) implies that there is an isomorphism

$$W^{\nu} H^i(\overline{\Gamma \backslash X}, L_{\tilde{\lambda}}(\mathbb{C})) = H_{(2)}^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C})).$$

(in the hermitian case this also follows from the Zucker conjecture, which was proven by Saper and Stern, and independently by Looijenga). Thus, if in addition the highest weight $\tilde{\lambda} \in X(\tilde{T})$ is regular then there is a canonical isomorphism of Hecke modules

$$W^{\nu} H^i(\overline{\Gamma \backslash X}, L_{\tilde{\lambda}}(\mathbb{C})) = H_{\text{cusp}}^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))$$

where the cohomology groups are nonvanishing only if $i = d$; in particular, we obtain

$$(-1)^d \text{tr}(T | H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))) = \text{Lef}(T | W^{\nu} H^{\bullet}(\overline{\Gamma \backslash X}, L_{\tilde{\lambda}}(\mathbb{C})))$$

where $T \in \mathcal{H}$ is a Hecke operator.

We mention that this implies an F -structure on cuspidal cohomology: we denote by $H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}}(F))$ the image of $W^{\nu} H^d(\overline{\Gamma \backslash X}, L_{\tilde{\lambda}}(F))$ in $H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))$ and obtain

$$H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C})) = H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}}(F)) \otimes \mathbb{C}.$$

2.7. Normalization of Hecke operators. We want to normalize the Hecke operators so that they act on cohomology with p -adically integral coefficients. We recall the following diagram of inclusions:

$$\begin{array}{ccc} \tilde{\mathcal{I}} & \subseteq & \mathcal{K} \\ \cup & & \cup \\ \Gamma & \subseteq & \Delta. \end{array}$$

Lemma. 1. $\mathcal{K} = \bigcup_{e \in \mathbb{N}_0} \tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$, i.e., any element $g \in \mathcal{K}$ can be written $g = k_1h^ek_2$ with $k_1, k_2 \in \tilde{\mathcal{I}}$ and $e \in \mathbb{N}_0$.

2. If $\tilde{\mathcal{I}}h^e\tilde{\mathcal{I}} = \tilde{\mathcal{I}}h^f\tilde{\mathcal{I}}$, $e, f \in \mathbb{N}_0$, then $e = f$.

3. Let $\tilde{\lambda} \in X(\tilde{\mathbf{T}})$ be a dominant weight. The mapping

$$\hat{\lambda} : \mathcal{K} \rightarrow F^*, \quad k_1h^ek_2 \mapsto \tilde{\lambda}(h^e) \quad (k_1, k_2 \in \tilde{\mathcal{I}}, e \in \mathbb{N}_0)$$

is a well defined morphism of semigroups.

Proof. Equations (5) and (6) in Section 1.5 remain valid with the same proof if \mathcal{I} is replaced by $\tilde{\mathcal{I}}$, $T_{\mathbb{Z}_p}$ is replaced by $\tilde{T}_{\mathbb{Z}_p} = T_{\mathbb{Z}_p} \tilde{\mathbf{G}}(\mathbb{Z}_p)$ and if $h, t, t' \in \tilde{\mathbf{T}}(\mathbb{Q}_p)$. Conclusion 1 is then immediate by equation (5). As for 2 we note that equation (6) implies $h^e = \delta h^f$, where $\delta \in \tilde{T}_{\mathbb{Z}_p} \subseteq \tilde{\mathbf{T}}(\mathbb{Z}_p)$. Applying an arbitrary simple root α and taking p -adic values yields $e v_p(\alpha(h)) = v_p(\alpha(\delta)) + f v_p(\alpha(h))$ and since $\alpha(\delta) \in \mathbb{Z}_p^*$ and $v_p(\alpha(h)) > 0$ we deduce that $e = f$. As for 3 we remark that parts 1 and 2 show that $\hat{\lambda}$ is well defined and equation (5) in Section 1.5 implies that $\hat{\lambda}$ is a morphism of semigroups. Thus, the lemma is proven. \square

Let $\tilde{\lambda} \in X(\tilde{\mathbf{T}})$ be a dominant weight. By restriction, $\hat{\lambda}$ induces a mapping $\hat{\lambda} : \Delta \rightarrow F^*$. For any F -algebra R we define an R -linear mapping

$$\mathcal{H}_R \rightarrow \mathcal{H}_R$$

by sending $\Gamma\zeta\Gamma \mapsto \hat{\lambda}(\zeta)\Gamma\zeta\Gamma$, $\zeta \in \Delta$; note that $\{\Gamma\zeta\Gamma, \zeta \in \Delta\}$ is a basis for \mathcal{H}_R and that the assignment is well defined since $\hat{\lambda}$ vanishes on $\tilde{\mathcal{I}}$ by definition and, hence, vanishes on $\Gamma \subseteq \tilde{\mathcal{I}}$. We denote the image of $T \in \mathcal{H}_R$ under the above mapping by $T_{\tilde{\lambda}} \in \mathcal{H}_R$ and we call $T_{\tilde{\lambda}}$ the $\tilde{\lambda}$ -normalization of T . In particular, if $\zeta \in \Delta$ with $\zeta \in \tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$ then

$$(T_{\tilde{\lambda}})_{\zeta} = \hat{\lambda}(\zeta)T_{\zeta} = \tilde{\lambda}(h^e)T_{\zeta}.$$

The normalization $T_{\tilde{\lambda}}$ of any $T \in \mathcal{H}_{\mathbb{Q}_p}$ leaves $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))$ invariant and we want to show that $T_{\tilde{\lambda}}$ leaves cohomology with integral coefficients $L_{\tilde{\lambda}}(\mathbb{Z}_p)$ invariant. To this end we first show that the “normalization $\hat{\lambda}(g)g^{-1}$ ” of any $g \in \mathcal{K}$ leaves the lattice $L_{\tilde{\lambda}}(\mathbb{Z}_p)$ invariant.

2.8. Lemma. For all $g \in \mathcal{K}$ and $v \in L_{\tilde{\lambda}}(\mathbb{Z}_p)$ we have $\hat{\lambda}(g)g^{-1}v \in L_{\tilde{\lambda}}(\mathbb{Z}_p)$.

Proof. Any $g \in \mathcal{K}$ has the form $g = k_1h^ek_2$ with $k_1, k_2 \in \tilde{\mathcal{I}}$. Since $\hat{\lambda}(g) = \tilde{\lambda}(h^e)$ and since $\tilde{\mathcal{I}} \subseteq \tilde{\mathbf{G}}(\mathbb{Z}_p)$ leaves $L_{\tilde{\lambda}}(\mathbb{Z}_p)$ invariant it is sufficient to show that $\tilde{\lambda}(h^e)h^{-e}$ leaves $L_{\tilde{\lambda}}(\mathbb{Z}_p)$ invariant. Since, as we saw in equation (9), we further have

$$L_{\tilde{\lambda}}(\mathbb{Z}_p) = \bigoplus_{\mu \leq \tilde{\lambda}} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu),$$

it is sufficient to show that $\tilde{\lambda}(h^e)h^{-e}v_{\mu} \in L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu)$ for any weight vector v_{μ}

in $L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu)$. Equation (9) implies that $\tilde{\lambda}(h^e)h^{-e}v_\mu = \tilde{\lambda}(h^e)\tilde{\mu}(h^{-e})v_\mu$. We write $\mu = \lambda - \nu$ where $\nu = \sum_{\alpha \in \Delta} n_\alpha \alpha$ with $n_\alpha \in \mathbb{N}_0$ and $h = tz$ with $t \in \mathbf{T}(\bar{\mathbb{Q}}_p)$, $z \in \tilde{\mathbf{Z}}(\bar{\mathbb{Q}}_p)$ and obtain

$$\begin{aligned} \tilde{\lambda}(h^e)\tilde{\mu}(h^{-e}) &= \lambda^\circ(t^e)\tilde{\lambda}(z^e)\mu^\circ(t^{-e})\tilde{\lambda}(z^{-e}) = \lambda^\circ(t^e)\mu^\circ(t^{-e}) \\ &= \lambda^\circ(t^e)\lambda^\circ(t^{-e})\nu^\circ(t^e) = \prod_{\alpha \in \Delta} \alpha(t)^{en_\alpha}. \end{aligned}$$

Since $v_p(\alpha(t)) = v_p(\alpha(h)) \geq 1$ for all simple roots α we deduce that

$$v_p\left(\prod_{\alpha \in \Delta} \alpha(t)^{en_\alpha}\right) \geq e \sum_{\alpha \in \Delta} n_\alpha = e \operatorname{ht}(\nu) \geq 0.$$

Thus, taking into account that $\operatorname{ht}(\nu) = \operatorname{ht}_\lambda(\mu)$ we obtain

$$(11) \quad \tilde{\lambda}(h^e)h^{-e}v_\mu \in p^{e \operatorname{ht}_\lambda(\mu)} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) \subseteq L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu),$$

which implies that $\tilde{\lambda}(h^e)h^{-e}$ leaves $L_{\tilde{\lambda}}(\mathbb{Z}_p)$ invariant. The lemma is proven. \square

- 2.9. Corollary.** 1. For any $T \in \mathcal{H}_{\mathbb{Z}_p}$ the normalized operator $T_{\tilde{\lambda}}$ leaves the group $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))$ invariant. In particular, $T_{\tilde{\lambda}}$ acts on integral cohomology $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}}$.
2. For any $T \in \mathcal{H}$ the eigenvalues of $T_{\tilde{\lambda}} \in \mathcal{H}_F$ on $H^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))$ and on $H^i_{\text{cusp}}(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))$ are algebraic over F (note that $F \subseteq \mathbb{C}$) and are contained in $\mathcal{O}_{\bar{\mathbb{Q}}_p}$ (note that $\bar{F} \subseteq \bar{\mathbb{Q}}_p$).

Proof. 1. We may assume that $T = T_\zeta$ for some $\zeta \in \Delta$. The claim then follows directly from the definition of the action of T_ζ on cohomology given in equation (10) and the lemma just proved (note that $\zeta \in \Delta \subseteq \mathcal{K}$ and that $\gamma_i^{-1} \in \Gamma \subseteq \tilde{\mathbf{G}}(\mathbb{Z}_p)$ leaves $L_{\tilde{\lambda}}(\mathbb{Z}_p)$ invariant).

2. The cuspidal cohomology $H^i_{\text{cusp}}(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C})) \subseteq H^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))$ is a Hecke submodule of full cohomology, hence, the eigenvectors and (complex) eigenvalues of $T_{\tilde{\lambda}}$ on $H^i_{\text{cusp}}(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))$ are contained in the set of eigenvectors and eigenvalues of $T_{\tilde{\lambda}}$ on $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{C}))$. Since $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{C})) = H^i(\Gamma, L_{\tilde{\lambda}}(F)) \otimes \mathbb{C}$ all eigenvalues of $T_{\tilde{\lambda}}$ on $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{C}))$ are algebraic over F and, hence, they already appear as eigenvalues of $T_{\tilde{\lambda}}$ on $H^i(\Gamma, L_{\tilde{\lambda}}(\bar{F}))$. In particular, they also appear as eigenvalues of $T_{\tilde{\lambda}}$ on $H^i(\Gamma, L_{\tilde{\lambda}}(\bar{\mathbb{Q}}_p))$ where the latter contains the $T_{\tilde{\lambda}}$ -invariant \mathbb{Z}_p -lattice $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}}$. Thus, all eigenvalues of $T_{\tilde{\lambda}}$ on $H^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))$ are algebraic over F and contained in the integer ring $\mathcal{O}_{\bar{\mathbb{Q}}_p}$ after embedding \bar{F} in $\bar{\mathbb{Q}}_p$. This completes the proof. \square

The diagram of inclusions on the next page recapitulates the objects appearing in the proof above and groups them together for easy lookup as they come up later in the discussion.

$$\begin{aligned}
 H_{\text{cusp}}^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C})) &\subseteq H^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C})) \\
 &\cup \\
 &H^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\bar{\mathbb{F}})) \subseteq H^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\bar{\mathbb{Q}}_p)) \\
 &\cup \\
 &H^i(\Gamma \backslash X, L_{\tilde{\lambda}}(F)) \subseteq H^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{Q}_p)) \\
 &\cup \\
 &H^i(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}}
 \end{aligned}$$

2.10. Mod p reduction of irreducible representations. We denote by T_m the maximal torus in \mathbf{GL}_m consisting of diagonal matrices, by \mathbf{B}_m^- the Borel subgroup in \mathbf{GL}_m consisting of all lower triangular matrices and by $\wp : \mathbf{GL}_m(\mathbb{Z}_p) \rightarrow \mathbf{GL}_m(\mathbb{F}_p)$ the mod p reduction map for \mathbf{GL}_m . Since $\tilde{\mathbf{G}}/\mathbb{Z}_p$ is smooth the reduction map $\wp : \tilde{\mathbf{G}}(\mathbb{Z}_p) \rightarrow \tilde{\mathbf{G}}(\mathbb{F}_p)$ is surjective. We let $\tilde{\lambda} \in X(\tilde{\mathbf{T}})$ be a dominant weight and as before we set $\lambda = d(\tilde{\lambda}|_T) \in \mathfrak{h}^*$. We define the mod p reduction

$$\bar{\rho}_{\tilde{\lambda}} : \tilde{\mathbf{G}}(\mathbb{F}_p) \rightarrow \mathbf{GL}_m(\mathbb{F}_p)$$

of the representation $\rho_{\tilde{\lambda}} : \tilde{\mathbf{G}}/\mathbb{Z}_p \rightarrow \mathbf{GL}_m/\mathbb{Z}_p$ by $\bar{\rho}_{\tilde{\lambda}}(\bar{g}) = \wp(\rho_{\tilde{\lambda}}(g))$, where $g \in \tilde{\mathbf{G}}(\mathbb{Z}_p)$ satisfies $\wp(g) = \bar{g}$. We denote by $\mathcal{J}_m \subseteq \mathbf{GL}_m(\mathbb{Z}_p)$ the Iwahori subgroup consisting of all elements $g \in \mathbf{GL}_m(\mathbb{Z}_p)$ such that $\wp(g) \in \mathbf{B}_m^-(\mathbb{F}_p)$.

Lemma.

$$\rho_{\tilde{\lambda}}(\tilde{\mathcal{I}}) \subseteq \mathcal{J}_m.$$

Proof. In Section 1.4 we selected a \mathbb{Z}_p -basis \mathcal{B} of $L_{\lambda}(\mathbb{Z}_p)$ to identify $\text{Aut}(L_{\lambda}(\mathbb{Z}_p)) = \mathbf{GL}_m(\mathbb{Z}_p)$. Since $L_{\lambda}(\mathbb{Z}_p) = \bigoplus_{\mu \in P_{\lambda}} L_{\lambda}(\mathbb{Z}_p, \mu)$ — see Section 1.1 — we may choose a basis \mathcal{B} consisting of weight vectors w.r.t. \mathfrak{h} . We order \mathcal{B} so that, if $v_{\mu}, v_{\mu'} \in \mathcal{B}$ are vectors of respective weights $\mu, \mu' \in \mathfrak{h}^*$, then $\text{ht}_{\lambda}(\mu) < \text{ht}_{\lambda}(\mu')$ implies that $v_{\mu} < v_{\mu'}$. We consider the image $\rho_{\tilde{\lambda}}(x_{\alpha}^{\pi}(t_{\alpha})) = \rho_{\lambda \circ (x_{\alpha}^{\pi}(t_{\alpha}))} = x_{\alpha}^{\lambda}(t_{\alpha}) \in \text{Aut}(L_{\lambda}(\bar{\mathbb{Q}}_p))$, where $\alpha \in \Phi^-$ ($t_{\alpha} \in \bar{\mathbb{Q}}_p$). Let $v_{\mu} \in \mathcal{B}$ be a basis vector of weight μ . Since $x_{\alpha}^{\lambda}(t_{\alpha})v_{\mu} = v_{\mu} + t_{\alpha}x_{\alpha}v_{\mu} + \frac{1}{2}t_{\alpha}^2x_{\alpha}^2v_{\mu} + \dots$, we see that $x_{\alpha}^{\lambda}(t_{\alpha})v_{\mu}$ is a sum of vectors of weights $\mu, \mu + \alpha, \mu + 2\alpha, \dots$, which are of strictly increasing relative height (since $\alpha < 0$). Hence, $\rho_{\tilde{\lambda}}(x_{\alpha}^{\pi}(t_{\alpha}))$ has lower triangular form w.r.t. \mathcal{B} , i.e., $\rho_{\tilde{\lambda}}(x_{\alpha}^{\pi}(t_{\alpha})) \in \mathbf{B}_m^-(\bar{\mathbb{Q}}_p)$. This shows that $\rho_{\tilde{\lambda}}(N^-(\bar{\mathbb{Q}}_p)) \subseteq \mathbf{B}_m^-(\bar{\mathbb{Q}}_p)$. Since $T(\bar{\mathbb{Q}}_p)$ preserves weight spaces by equation (4) in Section 1.4 we find quite analogous that $\rho_{\tilde{\lambda}}(T(\bar{\mathbb{Q}}_p)) \subseteq T_m(\bar{\mathbb{Q}}_p)$, hence, $\rho_{\tilde{\lambda}}(\mathbf{B}^-(\bar{\mathbb{Q}}_p)) \subseteq \mathbf{B}_m^-(\bar{\mathbb{Q}}_p)$. Thus, we obtain $\rho_{\tilde{\lambda}}(\mathbf{B}^-(\mathbb{Z}_p)) \subseteq \mathbf{B}_m^-(\bar{\mathbb{Q}}_p) \cap \mathbf{GL}_m(\mathbb{Z}_p) = \mathbf{B}_m^-(\mathbb{Z}_p)$ and, hence,

$$\bar{\rho}_{\tilde{\lambda}}(\mathbf{B}^-(\mathbb{F}_p)) \subseteq \mathbf{B}_m^-(\mathbb{F}_p).$$

We obtain $\wp(\rho_{\tilde{\lambda}}(\mathcal{I})) = \bar{\rho}_{\tilde{\lambda}}(\wp(\mathcal{I})) \subseteq \bar{\rho}_{\tilde{\lambda}}(\mathbf{B}^-(\mathbb{F}_p)) \subseteq \mathbf{B}_m^-(\mathbb{F}_p)$ and since $\rho_{\tilde{\lambda}}(\mathcal{I}) \subseteq \rho_{\tilde{\lambda}}(\tilde{\mathbf{G}}(\mathbb{Z}_p)) \subseteq \mathbf{GL}_m(\mathbb{Z}_p)$ we deduce that $\rho_{\tilde{\lambda}}(\mathcal{I}) \subseteq \mathcal{J}_m$. Equation (8) implies that $\rho_{\tilde{\lambda}}(z) = \tilde{\lambda}(z)\mathbf{1}_{\mathbf{GL}_m} \in \mathcal{J}_m$ for all $z \in \tilde{\mathbf{Z}}(\mathbb{Z}_p)$, hence, we finally obtain $\rho_{\tilde{\lambda}}(\tilde{\mathcal{I}}) \subseteq \mathcal{J}_m$ and the lemma is proven. \square

3. Boundedness of dimension of slope subspaces

3.1. We keep the assumptions from the previous sections. In particular, \tilde{G}/\mathbb{Q} is a connected reductive group containing a \mathbb{Q}_p -split maximal torus \tilde{T}/\mathbb{Q} and $\Gamma \leq \tilde{G}(\mathbb{Q})$ is an arithmetic subgroup such that $\Gamma \subseteq \tilde{I}$. We will obtain bounds for the dimension of the slope subspaces of $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))$; see 3.10 Corollary. This extends the main result in [Mahnkopf 2014], since (i) we allow Γ to be an arbitrary subgroup in \tilde{I} (i.e., we do not assume that Γ is contained in the smaller group $K_*(p) < \tilde{I}$ defined in [Mahnkopf 2013; 2014]); (ii) we do not assume that $\Gamma \leq \mathbf{G}(\mathbb{Q})$ where \mathbf{G} is the derived group of \tilde{G} ; (iii) we obtain stronger bounds for the dimension of the slope subspaces than those in [Mahnkopf 2014]. The proof follows the one in that paper. To deal with arithmetic subgroups Γ which are only contained in \tilde{I} we have to generalize the notion of truncation of an irreducible representation introduced in [Mahnkopf 2013; 2014] (see Section 3.3).

3.2. We note the following corrections to the works just cited.

1. In [Mahnkopf 2014] we considered a connected reductive group \tilde{G} which is defined over a number field F with F_p -split maximal torus \tilde{T} (Section 1.4 there). As in the present article, we have to assume that \tilde{T} is defined over F (and split over F_p); thus, the F -points $\tilde{T}(F)$ are defined and we may select $h \in \tilde{T}(F)$ as done in Section 1.6 of [Mahnkopf 2014].

2. Let \mathbf{G} denote the derived group of the connected reductive group \tilde{G} which we considered in [Mahnkopf 2013; 2014]. Hence, \mathbf{G} is a semisimple group and in those two papers we assumed that it is isomorphic over a splitting field to a Chevalley group \mathbf{G}_{λ_0} for an irreducible representation ρ_{λ_0} of the Lie algebra $\mathfrak{g} = \text{Lie}(\mathbf{G}) \otimes \mathbb{C}$. In general, \mathbf{G} over its splitting field only is isomorphic to a Chevalley group \mathbf{G}_{π} for a semisimple representation π of \mathfrak{g} (if one restricts to irreducible representations π one does not obtain all covering groups of the adjoint group with Lie algebra \mathfrak{g}). Since in the cited papers we did not make use of the irreducibility of the representation ρ_{λ_0} the results also hold if we consider a Chevalley group \mathbf{G}_{π} which is attached to a semisimple representation π of \mathfrak{g} .

3. In the summation formula (7) in Section 2.5.3 of [Mahnkopf 2014], the Bernoulli number $B_s(0)$ has to be replaced by $B_s(1)$ (note that $B_s(1) = B_s(0)$ for all $s > 1$ but $B_1(1) = -B_1(0) = \frac{1}{2}$).

3.3. Truncations with slope parameter. For the moment we let $\sigma \in \mathbb{N}$ be any natural number and we consider the subgroup

$$\tilde{K}_*(\sigma) = \tilde{K}_*(p, \sigma) = \langle K_*(p, \sigma), \tilde{Z}(\mathbb{Z}_p) \rangle \leq \tilde{G}(\mathbb{Z}_p).$$

Thus, if $\sigma \geq \max_{\alpha \in \Phi^+} \text{ht}(\alpha)$ then $\tilde{K}_*(\sigma) = \tilde{\mathcal{L}}$. For any $r \in \mathbb{N}_0$ we define the \mathbb{Z}_p -submodule

$$(12) \quad L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma) := \bigoplus_{\substack{\mu \leq \lambda \\ 0 \leq \text{ht}_{\tilde{\lambda}}(\mu) \leq r\sigma}} p^{r - \lceil \frac{1}{\sigma} \text{ht}_{\tilde{\lambda}}(\mu) \rceil} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) \oplus \bigoplus_{\substack{\mu \leq \lambda \\ \text{ht}_{\tilde{\lambda}}(\mu) > r\sigma}} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu)$$

of $L_{\tilde{\lambda}}(\mathbb{Z}_p)$.

Lemma. *The \mathbb{Z}_p -module $L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)$ of $L_{\tilde{\lambda}}(\mathbb{Z}_p)$ is $\tilde{K}_*(\sigma)$ -invariant.*

Proof. In view of the definition of $\tilde{K}_*(\sigma)$ we have to show that the three types of generators of $\tilde{K}_*(\sigma)$ map any of the weight subspaces of $L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)$ (see equation (12)) to $L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)$. Since $t_{\alpha}^n x_{\alpha}^n / n!$, where $t_{\alpha} \in \mathbb{Z}_p$ and $\alpha \in \Phi$, maps $L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu)$ to $L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu + n\alpha)$, it is immediate that any $t_{\alpha}^n x_{\alpha}^n / n!$ with $\alpha < 0$ and $t_{\alpha} \in \mathbb{Z}_p$ maps any weight subspace $p^m L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu)$ contained in $L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)$ to $p^m L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu + n\alpha)$, which is contained in $L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)$ because $\text{ht}_{\tilde{\lambda}}(\mu + n\alpha) \geq \text{ht}_{\tilde{\lambda}}(\mu)$. Hence, any generator $x_{\alpha}(t_{\alpha})$, $\alpha \in \Phi^-$, $t_{\alpha} \in \mathbb{Z}_p$, leaves $L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)$ invariant. We look at generators $x_{\alpha}(t_{\alpha})$ where $\alpha \in \Phi^+$ —hence, $t_{\alpha} \in p^{\lceil \frac{1}{\sigma} \text{ht}(\alpha) \rceil} \mathbb{Z}_p$. Using the inequalities $\lceil x \rceil - \lceil y \rceil \leq \lceil x - y \rceil$ and $n \lceil x \rceil \geq \lceil nx \rceil$, $x, y \in \mathbb{R}$, $n \in \mathbb{N}$, we find for all $n \in \mathbb{N}$ and all weights $\mu \leq \lambda$ with $\text{ht}_{\tilde{\lambda}}(\mu) \leq r\sigma$:

$$\begin{aligned} t_{\alpha}^n \frac{x_{\alpha}^n}{n!} p^{r - \lceil \frac{1}{\sigma} \text{ht}_{\tilde{\lambda}}(\mu) \rceil} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) &\subseteq p^{n \lceil \frac{1}{\sigma} \text{ht}(\alpha) \rceil} p^{r - \lceil \frac{1}{\sigma} \text{ht}_{\tilde{\lambda}}(\mu) \rceil} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu + n\alpha) \\ &\subseteq p^{r - \lceil \frac{1}{\sigma} \text{ht}_{\tilde{\lambda}}(\mu) \rceil + \lceil n \frac{1}{\sigma} \text{ht}(\alpha) \rceil} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu + n\alpha) \\ &\subseteq p^{r - (\lceil \frac{1}{\sigma} \text{ht}_{\tilde{\lambda}}(\mu) \rceil - n \frac{1}{\sigma} \text{ht}(\alpha))} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu + n\alpha) \\ &= p^{r - (\lceil \frac{1}{\sigma} \text{ht}_{\tilde{\lambda}}(\mu + n\alpha) \rceil)} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu + n\alpha) \\ &\subseteq L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma); \end{aligned}$$

for the last inclusion note that $\text{ht}_{\tilde{\lambda}}(\mu + n\alpha) \leq \text{ht}_{\tilde{\lambda}}(\mu) \leq r\sigma$ (we remark that if $\mu + n\alpha$ is not $\leq \lambda$ then $t_{\alpha}^n (x_{\alpha}^n / n!) p^{r - \lceil \frac{1}{\sigma} \text{ht}_{\tilde{\lambda}}(\mu) \rceil} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) = 0$). For weights $\mu \leq \lambda$ with $\text{ht}_{\tilde{\lambda}}(\mu) > r\sigma$ we find

$$t_{\alpha}^n \frac{x_{\alpha}^n}{n!} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) \subseteq p^{n \lceil \frac{1}{\sigma} \text{ht}(\alpha) \rceil} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu + n\alpha) \subseteq p^{\lceil \frac{n}{\sigma} \text{ht}(\alpha) \rceil} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu + n\alpha).$$

Since $\lceil \frac{n}{\sigma} \text{ht}(\alpha) \rceil \geq 0$ this shows that $t_{\alpha}^n (x_{\alpha}^n / n!) L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) \subseteq L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)$ if $\text{ht}_{\tilde{\lambda}}(\mu + n\alpha) > r\sigma$. If $\text{ht}_{\tilde{\lambda}}(\mu + n\alpha) \leq r\sigma$ we note that

$$r - \lceil \frac{1}{\sigma} \text{ht}_{\tilde{\lambda}}(\mu + n\alpha) \rceil \leq r - \frac{1}{\sigma} (\text{ht}_{\tilde{\lambda}}(\mu) - n \text{ht}(\alpha)) \leq \frac{n}{\sigma} \text{ht}(\alpha),$$

which shows that again $t_{\alpha}^n (x_{\alpha}^n / n!) L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) \subseteq L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)$. Hence, the generators $x_{\alpha}(t_{\alpha})$, $\alpha \in \Phi^+$, $t_{\alpha} \in p^{\lceil \frac{1}{\sigma} \text{ht}(\alpha) \rceil} \mathbb{Z}_p$, also leave $L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)$ invariant. Finally if $t \in \tilde{\mathcal{T}}(\mathbb{Z}_p)$ then t leaves all weight spaces $L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu)$ invariant because the representation

$\rho_{\tilde{\lambda}}$ is defined over \mathbb{Z}_p . Hence, $L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)$ is invariant under all generators of $\tilde{K}_*(\sigma)$ and the lemma is proven. \square

Definition. The quotient

$$L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma) = \frac{L_{\tilde{\lambda}}(\mathbb{Z}_p)}{L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)} = \bigoplus_{\substack{\mu \leq \lambda \\ \text{ht}_\lambda(\mu) \leq r\sigma}} \frac{L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu)}{p^{r - \lceil \frac{1}{\sigma} \text{ht}_\lambda(\mu) \rceil} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu)}$$

is called the truncation of $L_{\tilde{\lambda}}(\mathbb{Z}_p)$ of height $r \in \mathbb{N}_0$ and slope $\frac{1}{\sigma}$, $\sigma \in \mathbb{N}$.

Remark. The lemma just proved implies that $L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma)$ is a $\tilde{K}_*(\sigma)$ -module. Therefore, if $\sigma \geq \max_{\alpha \in \Phi^+} \text{ht}(\alpha)$, i.e., $\tilde{K}_*(\sigma) = \tilde{\mathcal{I}}$, then $L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma)$ is a Γ -module (recall that $\Gamma \leq \tilde{\mathcal{I}}$).

3.4. Lemma. For any dominant and integral weight λ and any $r \in \mathbb{N}$, $\sigma \in \mathbb{N}$ there is an embedding (of \mathbb{Z}_p -modules)

$$L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma) \leq \bigoplus_{h=0}^r \left(\frac{\mathbb{Z}_p}{p^{r-h}\mathbb{Z}_p} \right)^{M_{\sigma,h}},$$

where $M_{\sigma,h} = \sigma(\sigma h + 1)^{s-1}$ ($s = |\Phi^+|$).

Proof. We write

$$L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma) = \bigoplus_{\substack{h \in \mathbb{N}_0 \\ 0 \leq h \leq r\sigma}} \bigoplus_{\substack{\mu \leq \lambda \\ \text{ht}_\lambda(\mu) = h}} \frac{\mathbb{Z}_p}{p^{r - \lceil \frac{1}{\sigma} h \rceil} \mathbb{Z}_p} \otimes L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu).$$

In the proof of 2.2 Lemma in [Mahnkopf 2014] we have seen that

$$\dim_{\mathbb{Z}_p} \bigoplus_{\substack{\mu \leq \lambda \\ \text{ht}_\lambda(\mu) = h}} L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) \leq N_h,$$

where N_h is the number of tuples $\mathbf{n} = (n_1, \dots, n_s) \in \mathbb{N}_0^s$ such that

$$\sum_{i=1}^s n_i \text{ht}(\alpha_i) = h$$

(recall that $\Phi^+ = \{\alpha_1, \dots, \alpha_s\}$). Hence,

$$(13) \quad L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma) \leq \bigoplus_{\substack{h \in \mathbb{N}_0 \\ 0 \leq h \leq r\sigma}} \left(\frac{\mathbb{Z}_p}{p^{r - \lceil \frac{1}{\sigma} h \rceil} \mathbb{Z}_p} \right)^{N_h}.$$

We select an $a \in \mathbb{N}$. The terms of the form $r - \lceil \frac{1}{\sigma} h \rceil$, $h \in \mathbb{N}_0$, which equal $r - a$ are then precisely those with $h = \sigma a - \sigma + 1, \sigma a - \sigma + 2, \dots, \sigma a$. Thus, the term

$\frac{\mathbb{Z}_p}{p^{r-a}\mathbb{Z}_p}$ appears with multiplicity

$$N_{\sigma a - \sigma + 1} + N_{\sigma a - \sigma + 2} + \dots + N_{\sigma a}$$

in the right-hand side of equation (13). It is easy to see that $N_h \leq (h + 1)^{s-1}$, which implies that

$$N_{\sigma a - \sigma + 1} + N_{\sigma a - \sigma + 2} + \dots + N_{\sigma a} \leq \sigma(\sigma a + 1)^{s-1}.$$

Thus, we obtain, as desired,

$$L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma) \leq \bigoplus_{\substack{a \in \mathbb{N}_0 \\ 0 \leq a \leq r}} \left(\frac{\mathbb{Z}_p}{p^{r-a}\mathbb{Z}_p} \right)^{M_{\sigma,a}}. \quad \square$$

3.5. From now on we set

$$\sigma = \max_{\alpha \in \Phi^+} \text{ht}(\alpha).$$

Hence, σ only depends on \tilde{G} and $\tilde{K}_*(\sigma) = \tilde{\mathcal{I}}$. In particular, $L(\mathbb{Z}_p, r, \sigma)$ is a $\tilde{\mathcal{I}}$ and, hence, a Γ -module. The inclusion $i : L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma) \subseteq L_{\tilde{\lambda}}(\mathbb{Z}_p)$ induces a mapping

$$i^* : H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)) \rightarrow H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p)).$$

We recall that h is an element in $\tilde{T}(\mathbb{Q})^{++}$ (see Section 2.3).

Lemma. 1. *The mapping i^* induces an injection*

$$i^* : H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))^{\text{TF}} \hookrightarrow H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}},$$

where superscript *TF* denotes the maximal torsion-free quotient. In particular, we may identify $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))^{\text{TF}}$ with its image in $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}}$ under i^* .

2. *Let $\zeta \in \Delta$; hence, $\zeta \in \tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$ for some $e \in \mathbb{N}_0$ and we assume that $e \in \mathbb{N}$. Then the Hecke operator $(T_\zeta)_{\tilde{\lambda}}$ induces an operator on $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}}$ and we obtain*

$$(T_\zeta)_{\tilde{\lambda}}(H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))^{\text{TF}}) \subseteq p^r H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}}.$$

Proof. 1. The exact sequence

$$0 \rightarrow L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma) \xrightarrow{i} L_{\tilde{\lambda}}(\mathbb{Z}_p) \xrightarrow{\pi} L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma) \rightarrow 0$$

yields an exact sequence

$$H^{i-1}(\Gamma, L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma)) \rightarrow H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)) \xrightarrow{i^*} H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p)) \xrightarrow{\pi^*} H^i(\Gamma, L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma)).$$

Since $H^i(\Gamma, L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma))$ is a finite abelian group we further obtain an exact sequence

$$(14) \quad 0 \rightarrow H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))^{\text{TF}} \xrightarrow{i^*} H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}} \xrightarrow{\pi^*} Q \rightarrow 0,$$

where Q is a certain subquotient of $H^i(\Gamma, L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma))$. Thus, i^* is injective.

2. The first claim follows from 2.9 Corollary. In equation (11) in Section 2.8 we have seen that for any $v_\mu \in L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu)$

$$\tilde{\lambda}(h^e)h^{-e}v_\mu \in p^{e \text{ht}_\lambda(\mu)}L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu).$$

Hence, for all weights $\mu \leq \lambda$ satisfying $\text{ht}_\lambda(\mu) \leq r\sigma$ we obtain

$$\begin{aligned} \tilde{\lambda}(h^e)h^{-e}p^{r-\lceil \frac{1}{\sigma} \text{ht}_\lambda(\mu) \rceil}L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) &\subseteq p^{r-\lceil \frac{1}{\sigma} \text{ht}_\lambda(\mu) \rceil + e \text{ht}_\lambda(\mu)}L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) \\ &\subseteq p^{r+(e-1)\text{ht}_\lambda(\mu)}L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) \subseteq p^rL_{\tilde{\lambda}}(\mathbb{Z}_p, \mu), \end{aligned}$$

and for all weights $\mu \leq \lambda$ satisfying $\text{ht}_\lambda(\mu) > r\sigma (\geq r)$ we obtain

$$\tilde{\lambda}(h^e)h^{-e}L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) \subseteq p^{e \text{ht}_\lambda(\mu)}L_{\tilde{\lambda}}(\mathbb{Z}_p, \mu) \subseteq p^rL_{\tilde{\lambda}}(\mathbb{Z}_p, \mu).$$

Hence, we obtain $\tilde{\lambda}(h^e)h^{-e}L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma) \subseteq p^rL_{\tilde{\lambda}}(\mathbb{Z}_p)$. Since $\zeta \in \tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$ with $e \geq 1$ and $\tilde{\mathcal{I}}$ leaves $L_{\tilde{\lambda}}(\mathbb{Z}_p)$ and $L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)$ invariant (see 3.3.1 Lemma) we obtain

$$\tilde{\lambda}(h^e)\zeta^{-1}L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma) \subseteq p^rL_{\tilde{\lambda}}(\mathbb{Z}_p)$$

which yields

$$(T_\zeta)_{\tilde{\lambda}}(C^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))) \subseteq p^rC^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p)) (\subseteq C^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p)))$$

(here, we view $C^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))$ as embedded in $C^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))$ via i^*). The last equation implies the claim. \square

We note that $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}} \cong H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}}$.

3.6. We select a resolution of the trivial Γ -module \mathbb{Z} ,

$$0 \rightarrow M_d \rightarrow \dots \rightarrow M_1 \rightarrow M_0 \rightarrow \mathbb{Z} \rightarrow 0,$$

where M_i is a free $\mathbb{Z}\Gamma$ -module of finite rank (see [Brown 1982, p. 199]; note that Γ is of type FL; see p. 218 in the same work). The groups $H^i(\Gamma, L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma))$ then may be computed as the cohomology of the complex

$$0 \rightarrow \text{Hom}_{\mathbb{Z}_p\Gamma}(M_{0,p}, L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma)) \rightarrow \dots \rightarrow \text{Hom}_{\mathbb{Z}_p\Gamma}(M_{d,p}, L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma)) \rightarrow 0$$

where $M_{i,p} = \mathbb{Z}_p \otimes M_i$. We set

$$g_i = g_{i,\Gamma} = \text{rk}_{\mathbb{Z}\Gamma} M_i.$$

Thus, g_i depends on i and the arithmetic group Γ .

3.7. Borel–Serre compactification. Following [Borel and Serre 1973] (see also [Brown 1982, pp. 14 and 218]) we can construct a finite free resolution $(M_i)_{d \geq i \geq 0}$ of \mathbb{Z} as follows. We denote by $Y = \Gamma \backslash \tilde{X}$ the Borel–Serre compactification of the locally symmetric space $\Gamma \backslash X$ attached to \tilde{G}/\mathbb{Q} . By the work of Borel and Serre Y is a compact $K(\Gamma, 1)$ space; hence, there is a *finite* CW complex \mathcal{Z} having the same homotopy type as Y . The universal cover \tilde{Y} of Y inherits a structure of CW complex $\tilde{\mathcal{Z}}$ from \mathcal{Z} and the cellular complex $\tilde{C}_\bullet = (\tilde{C}_i)_{d \geq i \geq 0}$ ($d = \dim X$) attached to $\tilde{\mathcal{Z}}$ is a complex consisting of free $\mathbb{Z}\Gamma$ -modules \tilde{C}_i . The module \tilde{C}_i has a natural $\mathbb{Z}\Gamma$ -basis which is in bijection with the set of i -cells of \mathcal{Z} (see [Brown 1982, p. 15]); hence, \tilde{C}_\bullet is a finite complex consisting of free, finitely generated $\mathbb{Z}\Gamma$ -modules. Since \tilde{Y} is contractible, its cohomology vanishes except in degree 0; hence, the complex

$$0 \rightarrow \tilde{C}_d \rightarrow \dots \rightarrow \tilde{C}_0 \rightarrow \mathbb{Z} \rightarrow 0$$

is exact and thus a resolution of \mathbb{Z} . In particular, we may select $M_i = \tilde{C}_i$ and since $\text{rk}_{\mathbb{Z}\Gamma} \tilde{C}_i$ equals the number of i -cells of \mathcal{Z} this shows that we can take for g_i the number of i -cells of a CW complex \mathcal{Z} which has the same homotopy type as $\Gamma \backslash \tilde{X}$.

3.8. Slope subspaces. We select a Hecke operator $T \in \mathcal{H}_{\mathbb{Z}_p}$. Let $\tilde{\lambda} \in X(\tilde{T})^{\text{dom}}$ and let E/\mathbb{Q}_p be an extension which is contained in $\bar{\mathbb{Q}}_p$. For any $\beta \in \mathbb{Q}_{\geq 0}$ we denote by

$$H^i(\Gamma, L_{\tilde{\lambda}}(E))^\beta = p_\beta(T_{\tilde{\lambda}})H^i(\Gamma, L_{\tilde{\lambda}}(E))$$

the slope β subspace of $H^i(\Gamma, L_{\tilde{\lambda}}(E))$ w.r.t. to the (normalized) Hecke operator $T_{\tilde{\lambda}}$. Here, $p(X) \in \mathbb{Z}_p[X]$ is the characteristic polynomial of $T_{\tilde{\lambda}}$ acting on $H^i(\Gamma, L_{\tilde{\lambda}}(E))$ and $p_\beta(X) = \prod_{\mu, v_p(\mu) \neq \beta} (X - \mu) \in \mathbb{Z}_p[X]$, where μ runs over all roots of $p(X)$ whose p -adic value is different from β . Thus, we obtain $H^i(\Gamma, L_{\tilde{\lambda}}(\bar{\mathbb{Q}}_p))^\beta = \bigoplus_{\mu \in \mathcal{O}_{\bar{\mathbb{Q}}_p}, v_p(\mu) = \beta} H^i(\Gamma, L_{\tilde{\lambda}}(\bar{\mathbb{Q}}_p))(\mu)$ where $H^i(\Gamma, L_{\tilde{\lambda}}(\bar{\mathbb{Q}}_p))(\mu)$ is the generalized eigenspace attached to the eigenvalue μ . We set

$$H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{\leq \beta} = \bigoplus_{0 \leq \gamma \leq \beta} H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))^\gamma$$

and we denote by $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{< \infty} = \bigoplus_{0 \leq \gamma < \infty} H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))^\gamma$ the finite slope subspace.

3.9. An estimate for the Newton polygon. We denote by

$$\mathcal{H}_{\mathbb{Z}_p}^{\text{reg}} \subseteq \mathcal{H}_{\mathbb{Z}_p}$$

the set of all Hecke operators $T = \sum_\zeta c_\zeta T_\zeta \in \mathcal{H}_{\mathbb{Z}_p}$ ($\zeta \in \Delta$, $c_\zeta \in \mathbb{Z}_p$) where $\zeta \in \tilde{\mathcal{I}}h^{e_\zeta} \tilde{\mathcal{I}}$ with $e_\zeta \geq 1$ for all ζ with $c_\zeta \neq 0$. We let $T \in \mathcal{H}_{\mathbb{Z}_p}^{\text{reg}}$. We set $t' = t'(\tilde{\lambda}, i) =$

$\dim H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{<\infty}$ and we denote by $p'(X) = \sum_{i=0}^{t'} a_i X^{t'-i} \in \mathbb{Z}_p[X]$ the characteristic polynomial and by

$$\mathcal{N}^{<\infty} = \mathcal{N}_{\tilde{\lambda}, i}^{<\infty} : [0, t'] \rightarrow \mathbb{R}_{\geq 0}$$

the Newton polygon of $T_{\tilde{\lambda}}$ acting on $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{<\infty}$ which contains the $T_{\tilde{\lambda}}$ -invariant lattice $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}}^{<\infty} = H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{<\infty} \cap H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}}$. Thus, $\mathcal{N}^{<\infty}$ is the lower convex hull of the points $(i, v_p(a_i))$, $i = 0, \dots, t'$, where we omit all points with $a_i = 0$ (note that $p'(0) \neq 0$, hence, $a_{t'} \neq 0$). We recall that $g_i = \text{rk}_{\mathbb{Z}\Gamma} M_i$ (see Sections 3.6 and 3.7) and that $B_s \in \mathbb{Q}[X]$ denotes the s -th Bernoulli polynomial.

Theorem. *For all dominant weights $\tilde{\lambda} \in X(\tilde{T})$ and all $i \in \mathbb{N}_0$ the Newton polygon $\mathcal{N}^{<\infty} = \mathcal{N}_{\tilde{\lambda}, i}^{<\infty}$ lies above the restriction to $[0, t']$ of the piecewise linear function $f_{\infty}^* = f_{i, \infty}^* : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ which connects the points $(0, 0)$ and*

$$P_j = \left(g_i \sigma \frac{B_s(\sigma(j+1)+1) - B_s(1)}{s}, g_i \sigma^{s+1} \frac{B_{s+1}(j+1) - B_{s+1}(1)}{s+1} \right),$$

where $j = 0, 1, 2, \dots$

Proof. We proceed in steps.

3.9.1. We let $\tilde{\lambda} \in X(\tilde{T})^{\text{dom}}$ and set $\lambda = d\tilde{\lambda}|_T \in \mathfrak{h}^*$. Moreover, we select a natural number $r \in \mathbb{N}$. Since $\sigma = \max_{\alpha \in \Phi^+} \text{ht}(\alpha)$ the \mathbb{Z}_p -module $L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma)$ is a Γ -module and by 3.4 Lemma we know that

$$L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma) \leq \bigoplus_{h=0}^r (\mathbb{Z}_p/p^{r-h}\mathbb{Z}_p)^{M_{\sigma, h}},$$

which implies that

$$(15) \quad \text{Hom}_{\mathbb{Z}_p\Gamma}(M_{i, p}, L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma)) \leq \bigoplus_{h=0}^r (\mathbb{Z}_p/p^{r-h}\mathbb{Z}_p)^{g_i M_{\sigma, h}}.$$

We denote by $(p^{a_l})_l$, $a_1 \geq a_2 \geq \dots \geq a_n > 0$, the sequence of elementary divisors of the right-hand side of equation (15), i.e.,

$$(16) \quad (p^{a_l})_{l=1, \dots, n} = (p^r, \dots, p^r, p^{r-1}, \dots, p^{r-1}, \dots, p, \dots, p),$$

where p^{r-h} appears $g_i M_{\sigma, h}$ -times. From 3.5 Lemma it follows that there is a natural embedding of $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))^{\text{TF}}$ in $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}}$ and the exact sequence in equation (14) shows that

$$(17) \quad \frac{H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}}}{H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))^{\text{TF}}} \text{ is a subquotient of } H^i(\Gamma, L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma)).$$

We denote by t the rank of $H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}}$ and by $(p^{b_l})_l, b_1 \geq b_2 \geq \dots \geq b_m > 0$ ($m \leq t$) the sequence of elementary divisors of the quotient on the left in (17). Equation (17) implies that this quotient is a subquotient of the Hom space on the left-hand side of equation (15); hence, it is a subquotient of the right-hand side of (15) and equation (16) yields $m \leq n$ and

$$(18) \quad b_1 \leq a_1, b_2 \leq a_2, \dots, b_m \leq a_m.$$

We set $b_l = 0$ for $m < l \leq t$ and $a_l = 0$ for $n < l \leq t$ (if $n < t$); hence, $b_i \leq a_i$ for $i = 1, \dots, t$.

3.9.2. Using the results so far we can give a lower bound for $\mathcal{N}^{<\infty}$. Equations (16) and (18) imply that the b_l are all smaller than or equal to r . Moreover, since $T = \sum_{\zeta} c_{\zeta} T_{\zeta}$ where $c_{\zeta} \in \mathbb{Z}_p$ and $\zeta \in \tilde{\mathcal{I}}h^{e_{\zeta}}\tilde{\mathcal{I}}$ with $e_{\zeta} \geq 1$ if $c_{\zeta} \neq 0$ 3.5 Lemma implies that

$$T_{\tilde{\lambda}}(H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))^{\text{TF}}) \subseteq p^r H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}}.$$

Thus, we may apply Lemma 1 in [Buzzard 2001], as recalled in Section 1.7 of [Mahnkopf 2014], to the pair $L = H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}}$ and $K = H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))^{\text{TF}}$ and the operator $\xi = T_{\tilde{\lambda}}$. More precisely, we denote by $f_b = f_{b,r} : [0, t'] \rightarrow \mathbb{R}_{\geq 0}$ the piecewise linear function attached to the sequence (b_1, \dots, b_t) ; i.e., f_b is the piecewise linear function joining the points $(j, C(j))$, $j = 0, \dots, t'$, where $C(j) = \sum_{l=1}^j (r - b_l)$. Then (1.7) Lemma in [Mahnkopf 2014] states that the Newton polygon $\mathcal{N}^{<\infty}$ of $T_{\tilde{\lambda}}$ acting on

$$(H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}} \otimes \mathbb{Q}_p)^{<\infty} = H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{<\infty}$$

is bounded from below by the graph of f_b .

3.9.3. We further estimate the function f_b . Equation (18) implies that f_b lies above the piecewise linear function $f_{a,r} : [0, t'] \rightarrow \mathbb{R}_{\geq 0}$ attached to the sequence $(a_1, \dots, a_{t'})$, i.e., $f_{a,r}$ joins the points $(j, A(j))$, $j = 0, \dots, t'$, where $A(j) = \sum_{l=1}^j r - a_l$ (the $A(j)$'s are equal to or smaller than the $C(j)$'s). Thus, we have

$$(19) \quad \mathcal{N}^{<\infty} \geq f_{b,r} \geq f_{a,r}$$

and this inequality holds for all $r \in \mathbb{N}$ since r was chosen arbitrarily. Using equation (16) it is not difficult to see that the function $f_{a,r}$ is the restriction to $[0, t']$ of the piecewise linear function on $\mathbb{R}_{\geq 0}$ which starts in $(0, 0)$ and has slope j for $g_i \sum_{h=0}^{j-1} M_{\sigma,h} \leq x \leq g_i \sum_{h=0}^j M_{\sigma,h}$, $j = 0, \dots, r - 1$, and slope r for $x \geq g_i \sum_{h=0}^{r-1} M_{\sigma,h}$. Since $M_{\sigma,h} = \sigma(\sigma h + 1)^{s-1}$, $h \geq 0$, we see that the function $f_{a,r}$ may be equivalently described as the piecewise linear function $f_{a,r} : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$

which starts in $(0, 0)$, has slope j for

$$(20) \quad g_i \sigma \sum_{h=0}^{j-1} (\sigma h + 1)^{s-1} \leq x \leq g_i \sigma \sum_{h=0}^j (\sigma h + 1)^{s-1}, \quad j = 0, \dots, r-1,$$

and slope r for $x \geq g_i \sigma \sum_{h=0}^{r-1} (\sigma h + 1)^{s-1}$. We set

$$x_s(j) = g_i \sigma \frac{B_s(\sigma(j+1)+1) - B_s(1)}{s}, \quad j = 0, 1, 2, \dots,$$

and we denote by $f_\infty : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ the piecewise linear function which starts in $(0, 0)$ and has slope 0 in the interval

$$0 \leq x \leq x_s(0)$$

and slope j in the interval

$$x_s(j-1) \leq x \leq x_s(j), \quad j = 1, 2, \dots$$

The function f_∞ like the function $f_{a,r}$ is monotonely increasing. Taking into account that for all $j \in \mathbb{N}_0$

$$(21) \quad \sum_{h=0}^{\sigma j - 1} (h + 1)^{s-1} = \sum_{h=1}^{\sigma j} h^{s-1} = \begin{cases} \frac{B_s(\sigma j + 1) - B_s(1)}{s} & \text{if } j \geq 1, \\ 0 & \text{if } j = 0, \end{cases}$$

we deduce that

$$x_s(0) = g_i \sigma \frac{B_s(\sigma + 1) - B_s(1)}{s} \geq g_i \sigma 1^{s-1} = g_i \sigma$$

and for all $j \in \mathbb{N}$

$$\begin{aligned} x_s(j) - x_s(j-1) &= g_i \sigma \sum_{h=\sigma j}^{\sigma(j+1)-1} (h + 1)^{s-1} \\ &\geq g_i \sigma (\sigma j + 1)^{s-1} = g_i \sigma \left(\sum_{h=0}^j (\sigma h + 1)^{s-1} - \sum_{h=0}^{j-1} (\sigma h + 1)^{s-1} \right). \end{aligned}$$

Thus, equation (20) implies that the segments of slope $0, \dots, r-1$ of f_∞ are longer than those of $f_{a,r}$, hence, $f_{a,r}(x) \geq f_\infty(x)$ for all $x \in [0, x_s(r-1)]$. This implies by equation (19) that

$$\mathcal{N}^{<\infty}(x) \geq f_\infty(x)$$

for all $x \in [0, x_s(r-1)]$. Since equation (19) holds for arbitrarily large $r \in \mathbb{N}$, and since $x_s(r-1) \rightarrow \infty$ for $r \rightarrow \infty$ by equation (21), we finally obtain

$$(22) \quad \mathcal{N}^{<\infty}(x) \geq f_\infty(x), \quad x \in [0, \infty).$$

3.9.4. We show that $f_\infty \geq f_\infty^*$. In view of equation (22) this completes the proof. By definition f_∞ is the piecewise linear function joining the points

$$(0, 0), \quad Q_j = \left(x_s(j), \sum_{h=1}^j h(x_s(h) - x_s(h-1)) \right), \quad j = 0, 1, 2, 3, \dots$$

We obtain the following estimate for the second coordinate $y_s(j)$ of Q_j , i.e., for the value $f_\infty(x_s(j))$:

$$y_s(0) = 0 = f_\infty^*(x_s(0))$$

and if $j \geq 1$ then equation (21) implies that

$$\begin{aligned} y_s(j) &= \sum_{h=1}^j h(x_s(h) - x_s(h-1)) \\ &= \sum_{h=1}^j h g_i \sigma \left(\frac{B_s(\sigma(h+1)+1) - B_s(1)}{s} - \frac{B_s(\sigma h+1) - B_s(1)}{s} \right) \\ &\stackrel{(21)}{=} g_i \sigma \sum_{h=1}^j h \sum_{k=\sigma h}^{\sigma(h+1)-1} (k+1)^{s-1} \geq g_i \sigma \sum_{h=1}^j h \sum_{k=\sigma h}^{\sigma h+\sigma-1} k^{s-1} \\ &\geq g_i \sigma \sum_{h=1}^j h \sigma (\sigma h)^{s-1} = g_i \sigma^{s+1} \sum_{h=1}^j h^s \\ &\stackrel{(21)}{=} g_i \sigma^{s+1} \frac{B_{s+1}(j+1) - B_{s+1}(1)}{s+1} = f_\infty^*(x_s(j)). \end{aligned}$$

Thus, $f_\infty \geq f_\infty^*$ and the theorem is proven. □

3.10. A bound for the dimension of slope subspaces. We recall that $s = |\Phi^+|$, $\sigma = \max_{\alpha \in \Phi^+} \text{ht}(\alpha)$ and g_i is the number of i -cells in a cell complex \tilde{Z} which is homotopy equivalent to $\Gamma \backslash \tilde{X}$.

Corollary. For all $\beta \in \mathbb{Q}_{\geq 0}$, all dominant weights $\tilde{\lambda} \in X(\tilde{T})$, all i and all Hecke operators $T \in \mathcal{H}_{\mathbb{Z}_p}^{\text{reg}}$ we have

$$\dim H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{\leq \beta} \leq m\beta^s + n;$$

here, $m = m_\Gamma = 12(g_i/s)\sigma^{s+1} \in \mathbb{Q}_{\geq 0}$ and $n = n_\Gamma \in \mathbb{N}$ is an integer which also only depends on g_i, σ, s (see (26) below); in particular, m and n only depend on Γ (and so on \tilde{G} and p) and i , but not on $\tilde{\lambda}, h$ and T .

Proof. Let $h : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$ be any function such that $f_\infty^*(x) \geq h(x)$ for all $x \geq 0$ and let $(d(\epsilon), y)$ with $d(\epsilon) > 0$ be an intersection point of h and the function

$w_\epsilon : x \mapsto (\beta + \epsilon)x$ ($\epsilon > 0$). Since

$$(\beta + \epsilon)x > \mathcal{N}^{<\infty}(x) \geq h(x)$$

for all $x \in [0, \dim H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{\leq \beta}]$ by 3.9 Theorem we deduce that $d(\epsilon) \geq \dim H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{\leq \beta}$; hence, we obtain an upper bound for the dimension of the slope $\leq \beta$ -subspace. We explicitly define a lower bound h for f_∞^* as follows. Since B_s is a polynomial of degree s and leading coefficient 1 there is a natural number $M = M(\sigma, s) \in \mathbb{N}$ such that

$$(23) \quad x_s(j) = g_i \sigma \frac{B_s(\sigma(j+1) + 1) - B_s(1)}{s} \leq 2^{\frac{1}{s+1}} \frac{g_i \sigma^{s+1}}{s} j^s$$

and

$$(24) \quad y_s(j) := g_i \sigma^{s+1} \frac{B_{s+1}(j+1) - B_{s+1}(1)}{s+1} \geq 2^{-\frac{1}{s}} \frac{g_i \sigma^{s+1}}{s+1} j^{s+1}$$

for all $j \geq M$. We define the function

$$h : [x_s(M), \infty) \rightarrow \mathbb{R}_{\geq 0}, \quad x \mapsto cx^{\frac{s+1}{s}},$$

where $c = 4^{-\frac{1}{s}} g_i^{-\frac{1}{s}} s^{\frac{s+1}{s}} \frac{1}{(s+1)} \sigma^{-\frac{s+1}{s}}$. We note that $x_s(M) \geq 0$ by equation (21). We then obtain for all $j \geq M$

$$\begin{aligned} h(x_s(j)) &= h\left(g_i \sigma \frac{B_s(\sigma(j+1) + 1) - B_s(1)}{s}\right) \stackrel{(23)}{\leq} c \left(2^{\frac{1}{s+1}} \frac{g_i \sigma^{s+1}}{s} j^s\right)^{\frac{s+1}{s}} \\ &= c 2^{\frac{1}{s}} \left(\frac{g_i}{s}\right)^{\frac{s+1}{s}} \sigma^{\frac{(s+1)^2}{s}} j^{s+1} \\ &= 2^{-\frac{1}{s}} \frac{g_i \sigma^{s+1}}{s+1} j^{s+1} \stackrel{(24)}{\leq} y_s(j). \end{aligned}$$

Since f_∞^* is the piecewise linear function connecting the points $P_j = (x_s(j), y_s(j))$, $j \in \mathbb{N}_0$ and $(0, 0)$, and since h passes below the points P_j , $j \geq M$, and is convex this implies that $h(x) \leq f_\infty^*(x)$ for all $x \geq x_s(M)$. We extend h to a function $h : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$ by setting $h(x) = f_\infty^*(x)$ for $x \in [0, x_s(M)]$ and $h(x) = cx^{\frac{s+1}{s}}$ if $x > x_s(M)$, hence, $f_\infty^*(x) \geq h(x)$ for all $x \in [0, \infty)$. As in the proof of 3.3 Corollary in [Mahnkopf 2014] we see that for all $\epsilon > 0$ the functions h and $x \mapsto (\beta + \epsilon)x$ always intersect in a point $(d(\epsilon), y)$ with $d(\epsilon) > 0$ and this point satisfies $d(\epsilon) \leq \max((\frac{\beta + \epsilon}{c})^s, x_s(M))$. Since $(\beta/c)^s, x_s(M) \geq 0$ are positive we obtain

$$\dim H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{\leq \beta} \leq \max((\beta/c)^s, x_s(M)) \leq (\beta/c)^s + x_s(M).$$

Since further

$$(25) \quad c^{-s} = 4g_i s^{-(s+1)} (s+1)^s \sigma^{s+1} = 4g_i s^{-1} \left(1 + \frac{1}{s}\right)^s \sigma^{s+1} \leq 4 \frac{g_i}{s} \sigma^{s+1} \exp(1)$$

and $\exp(1) \leq 3$ the claim of the corollary holds with $m = 12(g_i/s)\sigma^{s+1}$ and $n = x_s(M)$. The claim still holds if we replace n with any larger number and since equation (23) implies that $x_s(M) \leq 2^{\frac{1}{s+1}}(g_i\sigma^{s+1}/s)M^s$ the corollary in particular holds with

$$(26) \quad n = \left\lceil 2^{\frac{1}{s+1}} \frac{g_i \sigma^{s+1}}{s} M^s \right\rceil + 1 \in \mathbb{N}. \quad \square$$

4. Mod p^n reduction of traces of Hecke operators

4.1. In this section we will prove congruences between traces of powers of normalized Hecke operators on cuspidal cohomology for varying weight $\tilde{\lambda}$. Our main tool will be a comparison of Bewersdorff’s elementary trace formula for pairs $\tilde{\lambda}, \tilde{\lambda}'$ of congruent weights. The equality of mod p^n reductions of geometric sides follows from p -adic properties of the diagonalization of elements in $\tilde{\mathcal{I}}h^e\tilde{\mathcal{I}} \subseteq \tilde{\mathcal{G}}(\mathbb{Q}_p)$; see 4.4 Proposition (note that the Hecke operator $\Gamma\zeta\Gamma$, $\zeta \in \Delta$, is contained in $\tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$ for some $e \in \mathbb{N}_0$). In particular, the comparison is elementary and does not make use of advanced methods such as rigid analytic geometry or p -adic Banach space methods such as overconvergent cohomology. Using an adelic setting we prove analogous congruences on the Eisenstein part of cohomology and subtracting from full cohomology we obtain congruences on cuspidal cohomology (Sections 4.11–4.13).

In Section 4.14 we compare two Goresky–MacPherson trace formulas for two congruent weights. Equality of mod p^n reductions of the geometric sides again follows from the same diagonalization of elements in $\tilde{\mathcal{I}}h^e\tilde{\mathcal{I}} \subseteq \tilde{\mathcal{G}}(\mathbb{Q}_p)$ but now applied for all Levi subgroups \tilde{M} of \mathbb{Q} -parabolic subgroups of $\tilde{\mathcal{G}}$. This yields congruences on weighted cohomology groups and also has an application to a more explicit version of the Gouvêa–Mazur conjecture for symplectic groups of rank 2 (see Section 5.8).

As before, $\tilde{\mathcal{G}}/\mathbb{Q}$ is a connected reductive group containing a \mathbb{Q}_p -split maximal torus \tilde{T}/\mathbb{Q} and $\Gamma \subseteq \tilde{\mathcal{G}}(\mathbb{Q})$ is an arithmetic subgroup satisfying $\Gamma \subseteq \tilde{\mathcal{I}}$.

4.2. The fixed point principle of Bewersdorff. As in Section 2.6 we denote by $X = \tilde{\mathcal{G}}(\mathbb{R})/\tilde{K}_\infty A_{\tilde{\mathcal{G}}}$ the symmetric space attached to $\tilde{\mathcal{G}}$. The Γ -module $L_{\tilde{\lambda}}(\mathbb{Q}_p)$ defines a locally constant sheaf on the locally symmetric space $\Gamma \backslash X$ and its Borel–Serre compactification $\Gamma \backslash \tilde{X}$, which we will also denote by $L_{\tilde{\lambda}}(\mathbb{Q}_p)$, and the Hecke algebra \mathcal{H} acts on the cohomology groups

$$H^i(\Gamma, L_{\tilde{\lambda}}(\mathbb{Q}_p)) \cong H^i(\Gamma \backslash \tilde{X}, L_{\tilde{\lambda}}(\mathbb{Q}_p)).$$

We recall that in Section 2.5 we selected a finite extension F/\mathbb{Q} which splits $\tilde{\mathcal{G}}$ and which embeds in \mathbb{C}, \mathbb{Q}_p ; in particular, $(\rho_{\tilde{\lambda}}, L_{\tilde{\lambda}})$ is defined over F . We write

$\Gamma\zeta\Gamma/\sim_\Gamma$ for the set of Γ -conjugacy classes contained in the double coset $\Gamma\zeta\Gamma$, $\zeta \in \Delta$, and $[\xi]_\Gamma$ denotes the Γ -conjugacy class of $\xi \in \tilde{\mathbf{G}}(\mathbb{Q})$. We now borrow from [Bewersdorff 1985, Satz 2.6] a simple and elementary formula for the Lefschetz number of Hecke correspondences on full cohomology:

Theorem (Bewersdorff). *Let $\Gamma\zeta\Gamma \in \mathcal{H}$ (i.e., $\zeta \in \Delta$). There are rational integers $c_{[\xi]_\Gamma} \in \mathbb{Z}$, $[\xi]_\Gamma \in \Gamma\zeta\Gamma/\sim_\Gamma$, such that for all dominant weights $\tilde{\lambda} \in X(\tilde{\mathbf{T}})$*

$$(27) \quad \text{Lef}(\Gamma\zeta\Gamma|H^\bullet(\Gamma\backslash X, L_{\tilde{\lambda}}(F))) = \sum_{[\xi]_\Gamma \in \Gamma\zeta\Gamma/\sim_\Gamma} c_{[\xi]_\Gamma} \text{tr}(\xi^{-1}|L_{\tilde{\lambda}}(F))$$

and $c_{[\xi]_\Gamma}$ vanishes if $\xi x \neq x$ for all $x \in \tilde{X}$.

Remark. 1. The integers $c_{[\xi]_\Gamma}$ do not depend on the weight $\tilde{\lambda}$.

2. The trace formula (27) is of an elementary nature. Apart from the existence of a nice compactification of the locally symmetric space $\Gamma\backslash X$ (the Borel–Serre compactification) its proof is a direct application of the Lefschetz fixed point principle which is a general and basic principle of algebraic topology.

4.3. The following lemma will be applied in the proof of 4.4 Proposition, where representations $\tilde{\mathbf{G}}/\mathbb{Q}_p \hookrightarrow \mathbf{GL}_m/\mathbb{Q}_p$ of $\tilde{\mathbf{G}}$ as matrix group are used.

Lemma. *Let $\beta = (\beta_{ij}) \in \mathcal{J}_m$ and let $t = \text{diag}(t_1, \dots, t_m) \in \mathbf{T}_m(\mathbb{Q}_p)$ with $v_p(t_1) > v_p(t_i)$ for all $i = 2, \dots, m$. Then the characteristic polynomial $\text{ch}_{\beta t}$ of $\beta t \in \mathbf{GL}_m(\mathbb{Q}_p)$ has m roots t'_1, t'_2, \dots, t'_m in $\bar{\mathbb{Q}}_p$ (roots appearing several times according their multiplicity) such that $v_p(t'_1) > v_p(t'_i)$ for all $i = 2, \dots, m$ (in particular, t'_1 has multiplicity 1) and*

$$v_p(t'_1) = v_p(t_1) \quad \text{and} \quad t'_1 \equiv \beta_{11}t_1 \pmod{p^{v_p(t_1)+1}\mathcal{O}_{\bar{\mathbb{Q}}_p}}.$$

Proof. We put $[a, b] = \{a, a + 1, a + 2, \dots, b\}$ ($a, b \in \mathbb{N}$, $a \leq b$) and we denote by S_M the symmetric group on the set M . We write the characteristic polynomial of βt as $\text{ch}_{\beta t}(X) = (-1)^m X^m + (-1)^{m-1} c_1 X^{m-1} + \dots + c_{m-1} X + c_m$, $c_i \in \mathbb{Q}_p$ (i.e., $c_0 = 1$). The Leibniz formula

$$\text{ch}_{\beta t}(X) = \det(\beta t - X\mathbf{1}) = \sum_{\pi \in S_{[1,m]}} \text{sgn}(\pi) \prod_{i=1}^m (\beta_{\pi(i),i} t_i - \delta_{\pi(i),i} X)$$

yields

$$c_i = \sum_{\substack{T \subseteq [1,m] \\ |T|=i}} \sum_{\pi \in S_T} c_{T,\pi}$$

for all $i = 1, \dots, m$, where

$$c_{T,\pi} = \text{sgn}(\pi) \prod_{h \in T} \beta_{\pi(h),h} t_h \in \mathbb{Q}_p.$$

Since

$$(28) \quad v_p(\beta_{g,h}t_h) \begin{cases} \geq v_p(t_h) + 1 & \text{if } g < h, \\ = v_p(t_h) & \text{if } g = h, \\ \geq v_p(t_h) & \text{if } g > h, \end{cases}$$

we obtain $v_p(c_{T,\pi}) \geq v_p(c_{T,\text{id}}) + 1$ for all $T \subseteq [1, m]$ and all $\pi \in S_T, \pi \neq \text{id}$. Hence,

$c_m = c_{[1,m],\text{id}} + \text{terms with } p\text{-adic value equal to or greater than } v_p(c_{[1,m],\text{id}}) + 1,$

which implies that

$$(29) \quad c_m \equiv \prod_{h=1}^m \beta_{hh}t_h \pmod{p^{v_p(c_m)+1}\mathbb{Z}_p} \quad \text{and} \quad v_p(c_m) = v_p\left(\prod_{h=1}^m t_h\right).$$

Since, moreover, $v_p(t_1) \geq v_p(t_i) + 1$ for all $i = 2, \dots, m$ we obtain from equation (28) that

- for any subset $T \subset [1, m], T \neq [2, m]$, of cardinality $m - 1$ and any $\pi \in S_T$ we have $v_p(c_{T,\pi}) \geq v_p(c_{T,\text{id}}) \geq v_p(c_{[2,m],\text{id}}) + 1$
- for any $\pi \in S_{[2,m]}, \pi \neq \text{id}$, we have $v_p(c_{[2,m],\pi}) \geq v_p(c_{[2,m],\text{id}}) + 1$.

Hence,

$c_{m-1} = c_{[2,m],\text{id}} + \text{terms with } p\text{-adic value equal to or greater than } v_p(c_{[2,m],\text{id}}) + 1,$

which implies that

$$(30) \quad c_{m-1} \equiv \prod_{h=2}^m \beta_{hh}t_h \pmod{p^{v_p(c_{m-1})+1}\mathbb{Z}_p} \quad \text{and} \quad v_p(c_{m-1}) = v_p\left(\prod_{h=2}^m t_h\right).$$

In particular,

$$(31) \quad v_p(c_m) - v_p(c_{m-1}) = v_p(t_1).$$

Finally, for $i = 1, \dots, m - 1$ we denote by $T_{i,\min} \subset [1, m]$ a subset of cardinality i such that $v_p\left(\prod_{h \in T_{i,\min}} t_h\right) \leq v_p\left(\prod_{h \in T} t_h\right)$ for all subsets $T \subseteq [1, m]$ of cardinality i (thus, $T_{m-1,\min} = [2, m]$). As above equation (28) implies

$$(32) \quad v_p(c_i) \geq v_p\left(\prod_{h \in T_{i,\min}} t_h\right) = \sum_{h \in T_{i,\min}} v_p(t_h)$$

for all $i = 1, \dots, m - 1$. Since $v_p(t_1) > v_p(t_h)$ for all $h = 2, \dots, m$ we obtain $T_{i,\min} \subseteq [2, m]$ and equations (30) and (32) imply that

$$v_p(c_{m-1}) - v_p(c_i) \leq \sum_{h \in [2,m] - T_{i,\min}} v_p(t_h) \leq (m - 1 - i)r$$

for all $i = 1, \dots, m - 1$, where $r = \max_{h=2}^m v_p(t_h)$. Equivalently,

$$(33) \quad v_p(c_i) \geq v_p(c_{m-1}) - (m - 1 - i)r$$

for all $i = 1, \dots, m - 1$. Since $(m - 1)r \geq \sum_{h=2}^m v_p(t_h) = v_p(c_{m-1})$ in view of equation (30), we see that equation (33) also holds for $i = 0$ ($v_p(c_0) = 0$). Thus:

- The line connecting the points $(m - 1, v_p(c_{m-1}))$ and $(m, v_p(c_m))$ has slope $v_p(t_1)$; see equation (31).
- All points $(i, v_p(c_i))$ with $0 \leq i \leq m - 1$ lie on or above the line g , which has slope r and passes through $(m - 1, v_p(c_{m-1}))$; see equation (33).

Since r is strictly smaller than $v_p(t_1)$ this shows that the Newton polygon \mathcal{N} of $\text{ch}_{\beta t}$ has the segment connecting $(m, v_p(c_m))$ and $(m - 1, v_p(c_{m-1}))$ as one of its sides while all other segments have slope less than or equal to r . We deduce that there is precisely one root $t'_1 \in \bar{\mathbb{Q}}_p$ of $\text{ch}_{\beta t}$ (counted with multiplicity) such that

$$v_p(t'_1) = v_p(t_1),$$

while all remaining roots $t'_h \in \bar{\mathbb{Q}}_p$ of $\text{ch}_{\beta t}$, $h = 2, \dots, m$, have p -adic value smaller than or equal to r (in particular t'_1 appears with multiplicity 1). Since $r = \max_{h=2}^m v_p(t_h) < v_p(t_1)$ we obtain $v_p(t'_1) \geq v_p(t'_h) + 1$ for all $h = 2, \dots, m$. This implies that

$$c_m = \prod_{h=1}^m t'_h \quad \text{and} \quad c_{m-1} \equiv \prod_{h=2}^m t'_h \pmod{p^{v_p(c_{m-1})+1} \mathcal{O}_{\bar{\mathbb{Q}}_p}}$$

(note that $\text{ch}_{\beta t}(X) = \prod_{h=1}^m (t'_h - X)$ because both sides have leading coefficient $(-1)^m$). Together with equations (29) and (30) we obtain

$$\prod_{h=1}^m \beta_{hh} t_h \equiv \prod_{h=1}^m t'_h \pmod{p^{v_p(c_m)+1} \mathbb{Z}_p}$$

and

$$\prod_{h=2}^m \beta_{hh} t_h \equiv \prod_{h=2}^m t'_h \pmod{p^{v_p(c_{m-1})+1} \mathcal{O}_{\bar{\mathbb{Q}}_p}}.$$

Since $v_p(c_m) - v_p(c_{m-1}) = v_p(t_1)$ the above two equations imply that $\beta_{11} t_1 \equiv t'_1 \pmod{p^{v_p(t_1)+1} \mathcal{O}_{\bar{\mathbb{Q}}_p}}$ and the proof of the lemma is complete. □

4.4. The following proposition is an extension of 4.3 Lemma to closed subgroups of \mathbf{GL}_m and is used in the proof of 4.7 Proposition. We denote by $W_{\tilde{\mathbf{G}}}$ the Weyl group of $\tilde{\mathbf{T}}/F \leq \tilde{\mathbf{G}}/F$.

Proposition. *Let $t \in \tilde{T}(\mathbb{Q}_p)^{++}$ and let $x \in \tilde{\mathcal{I}}t\tilde{\mathcal{I}}$. Then the semisimple part x_s of $x \in \tilde{\mathbf{G}}(\mathbb{Q}_p)$ is $\tilde{\mathbf{G}}(\bar{\mathbb{Q}}_p)$ -conjugate to a uniquely determined element $t' = t'_x \in \tilde{T}(\bar{\mathbb{Q}}_p)^{++}$. The element t' satisfies*

$$v_p(\tilde{\lambda}(t')) = v_p(\tilde{\lambda}(t)) \quad \text{and} \quad \tilde{\lambda}(t') \equiv \epsilon \tilde{\lambda}(t) \pmod{p^{v_p(\tilde{\lambda}(t))+1} \mathcal{O}_{\bar{\mathbb{Q}}_p}}$$

for all $\tilde{\lambda} \in X(\tilde{T})$, where $\epsilon = \epsilon_{\tilde{\lambda}, x} \in \mathbb{Z}_p^*$ is a p -adic unit in \mathbb{Z}_p .

Proof. Conjugating x by an element in $\tilde{\mathcal{I}} \subseteq \tilde{\mathbf{G}}(\mathbb{Q}_p)$ we may assume that $x = \beta t$ with $\beta \in \tilde{\mathcal{I}}$. Since \mathbb{Q}_p is a perfect field we know that the semisimple part $(\beta t)_s$ of $\beta t \in \tilde{\mathbf{G}}(\mathbb{Q}_p)$ also is contained in $\tilde{\mathbf{G}}(\mathbb{Q}_p)$ (see [Sp 1], 12.1.7 (c), p. 211). By 6.4.5 Theorem (ii) in [Sp 1], p. 109, $(\beta t)_s$ is contained in $\tilde{\mathbf{S}}(\bar{\mathbb{Q}}_p)$, where $\tilde{\mathbf{S}} = \tilde{\mathbf{S}}_{\beta t}$ is a maximal $\bar{\mathbb{Q}}_p$ -torus in $\tilde{\mathbf{G}}/\bar{\mathbb{Q}}_p$. Since all maximal tori in $\tilde{\mathbf{G}}/\bar{\mathbb{Q}}_p$ are conjugate over $\bar{\mathbb{Q}}_p$, there is $g \in \tilde{\mathbf{G}}(\bar{\mathbb{Q}}_p)$ such that ${}^s\tilde{T} := g\tilde{T}g^{-1} = \tilde{\mathbf{S}}$; in particular,

$$t' := g^{-1}(\beta t)_s g \in \tilde{T}(\bar{\mathbb{Q}}_p).$$

Conjugating further by some $w \in W_{\tilde{\mathbf{G}}}$ we may assume that $t' \in \tilde{T}(\bar{\mathbb{Q}}_p)^+$. Thus, t' is $\tilde{\mathbf{G}}(\bar{\mathbb{Q}}_p)$ -conjugate to $x_s = (\beta t)_s$ and we will show that it satisfies the conditions of the Proposition. To this end we let $\tilde{\lambda} \in X(\tilde{T})$ and we first assume that $\tilde{\lambda}$ is dominant. We denote by $(\rho_{\tilde{\lambda}}, L_{\tilde{\lambda}})$ the irreducible representation of $\tilde{\mathbf{G}}/\mathbb{Z}_p$ of highest weight $\tilde{\lambda}$ (see Section 1.4 and 2.4). We select a basis (v_1, v_2, \dots, v_m) of $L_{\tilde{\lambda}}(\mathbb{Q}_p)$ consisting of weight vectors w.r.t. \mathfrak{h} as in the proof of 2.10 Lemma, i.e., if μ_i denotes the weight of v_i then $\text{ht}_{\tilde{\lambda}}(\mu_i) > \text{ht}_{\tilde{\lambda}}(\mu_j)$ implies $i > j$. We note that v_i has weight $\tilde{\mu}_i = \mu_i \circ \tilde{\lambda}|_{\tilde{\mathcal{Z}}}$ w.r.t. $\tilde{T}(\bar{\mathbb{Q}}_p)$ (see equation (9) in Section 2.4). In particular, v_1 is the highest weight vector, i.e., $\tilde{\mu}_1 = \tilde{\lambda}$. The above choice of a basis of $L_{\tilde{\lambda}}$ yields a matrix representation

$$\rho_{\tilde{\lambda}} : \tilde{\mathbf{G}}(\mathbb{Q}_p) \rightarrow \mathbf{GL}_m(\mathbb{Q}_p)$$

and 2.10 Lemma implies that

$$\rho_{\tilde{\lambda}}(\tilde{\mathcal{I}}) \subseteq \mathcal{I}_m.$$

Moreover, we obtain

$$\rho_{\tilde{\lambda}}(t) = \text{diag}(\tilde{\mu}_1(t), \tilde{\mu}_2(t), \dots, \tilde{\mu}_m(t)) \in \mathbf{T}_m(\mathbb{Q}_p)$$

and

$$\rho_{\tilde{\lambda}}(t') = \text{diag}(\tilde{\mu}_1(t'), \tilde{\mu}_2(t'), \dots, \tilde{\mu}_m(t')) \in \mathbf{T}_m(\bar{\mathbb{Q}}_p).$$

Any weight μ_i , $i \geq 2$, has the form $\mu_i = \lambda - \sum_{\alpha \in \Delta} n_{\alpha} \alpha$ where not all $n_{\alpha} \in \mathbb{N}_0$ are equal to zero and since $t \in \tilde{T}(\mathbb{Q}_p)^{++}$ (i.e., $v_p(\alpha(t)) > 0$ for all $\alpha \in \Delta$) we obtain

$$(34) \quad v_p(\tilde{\mu}_i(t)) = v_p(\tilde{\lambda}(t)) - \sum_{\alpha \in \Delta} n_{\alpha} v_p(\alpha(t)) < v_p(\tilde{\lambda}(t))$$

for all weights $\tilde{\mu}_i$, $i = 2, \dots, m$. Analogously, since $t' \in \tilde{T}(\bar{\mathbb{Q}}_p)^+$ we obtain

$$(35) \quad v_p(\tilde{\mu}_i(t')) = v_p(\tilde{\lambda}(t')) - \sum_{\alpha \in \Delta} n_\alpha v_p(\alpha(t')) \leq v_p(\tilde{\lambda}(t'))$$

for all weights $\tilde{\mu}_i, i = 2, \dots, m$. Since $\rho_{\tilde{\lambda}}(\beta) \in \rho_{\tilde{\lambda}}(\tilde{X}) \subseteq \mathcal{J}_m$ and $\rho_{\tilde{\lambda}}(t) \in \mathbf{T}_m(\mathbb{Q}_p)$ where the first entry of $\rho_{\tilde{\lambda}}(t)$ has p -adic value strictly bigger than the remaining entries (see equation (34)) we may apply 4.3 Lemma to $\rho_{\tilde{\lambda}}(\beta t) = \rho_{\tilde{\lambda}}(\beta)\rho_{\tilde{\lambda}}(t) \in \mathbf{GL}_m(\mathbb{Q}_p)$; we obtain that $\rho_{\tilde{\lambda}}(\beta t)$ has an eigenvalue $t'_1 \in \bar{\mathbb{Q}}_p$ of multiplicity 1 whose p -adic value is strictly larger than the p -adic values of the $m-1$ remaining eigenvalues of $\rho_{\tilde{\lambda}}(\beta t)$ and which satisfies

$$(36) \quad v_p(\tilde{\lambda}(t)) = v_p(t'_1) \quad \text{and} \quad \tilde{\lambda}(t) \equiv \epsilon t'_1 \pmod{p^{v_p(\tilde{\lambda}(t))+1} \mathcal{O}_{\bar{\mathbb{Q}}_p}}$$

for some $\epsilon = \epsilon_{\tilde{\lambda}, \beta} \in \mathbb{Z}_p^*$. Since the matrix $\rho_{\tilde{\lambda}}(t')$ has the same eigenvalues as $\rho_{\tilde{\lambda}}((\beta t)_s) = (\rho_{\tilde{\lambda}}(\beta t))_s$ it has the same eigenvalues as $\rho_{\tilde{\lambda}}(\beta t)$. In particular, $\rho_{\tilde{\lambda}}(t')$ has the eigenvalue t'_1 with multiplicity 1 whose p -adic value is strictly bigger than the p -adic values of the $m-1$ remaining eigenvalues of $\rho_{\tilde{\lambda}}(t')$. Thus, equation (35) shows that $t'_1 = \tilde{\lambda}(t')$ and equation (36) yields

$$(37) \quad v_p(\tilde{\lambda}(t)) = v_p(\tilde{\lambda}(t')) \quad \text{and} \quad \tilde{\lambda}(t) \equiv \epsilon \tilde{\lambda}(t') \pmod{p^{v_p(\tilde{\lambda}(t))+1} \mathcal{O}_{\bar{\mathbb{Q}}_p}}.$$

We recall that we have proven (37) under the assumption that $\tilde{\lambda}$ is dominant. For the general case we will need the following consequence of (37). Let $\tilde{\lambda}$ be dominant and write $\tilde{\lambda} = \tilde{\lambda}|_{\tilde{Z}} \lambda^\circ$ where $\lambda^\circ = \tilde{\lambda}|_T$. We write $t = t^\circ z$ where $t^\circ \in \mathbf{T}(\bar{\mathbb{Q}}_p), z \in \tilde{Z}(\bar{\mathbb{Q}}_p)$, hence, $t' = t'^\circ z$ where $t'^\circ = g^{-1}(\beta t^\circ)_s g$. Equation (37) then implies that

$$(38) \quad 1 \equiv \epsilon \frac{\lambda^\circ(t'^\circ)}{\lambda^\circ(t^\circ)} \pmod{p \mathcal{O}_{\bar{\mathbb{Q}}_p}}.$$

Since $\mathbf{T} \subseteq \tilde{\mathbf{T}}$ is a closed subset the restriction map $X(\tilde{\mathbf{T}}) \rightarrow X(\mathbf{T})$ is surjective, hence, any dominant character $\lambda^\circ \in X(\mathbf{T})$ is the restriction of a dominant character $\tilde{\lambda} \in X(\tilde{\mathbf{T}})$ and we deduce that equation (38) holds for all dominant $\lambda^\circ \in X(\mathbf{T})$.

We now let $\tilde{\lambda} \in X(\tilde{\mathbf{T}})$ be arbitrary and we show that equation (37) still holds. We write $\tilde{\lambda} = \tilde{\lambda}|_{\tilde{Z}} \lambda^\circ$ where $\lambda^\circ = \tilde{\lambda}|_T$. Applying 4.4.1 Lemma below to $\lambda = d \lambda^\circ \in \Gamma_\pi$ we can write $\lambda^\circ = \prod_i (\mu_i^\circ)^{n_i}$, where $\mu_i^\circ \in X(\mathbf{T})$ is dominant and $n_i \in \mathbb{Z}$. Equation (38) implies that

$$1 \equiv \prod_i \epsilon_i^{n_i} \frac{\prod_i \mu_i^\circ(t'^\circ)^{n_i}}{\prod_i \mu_i^\circ(t^\circ)^{n_i}} \pmod{p \mathcal{O}_{\bar{\mathbb{Q}}_p}}$$

with certain $\epsilon_i \in \mathbb{Z}_p^*$. Multiplying this by $\tilde{\lambda}(t) = \tilde{\lambda}(z) \lambda^\circ(t^\circ)$ we finally obtain

$$\tilde{\lambda}(t) \equiv \prod_i \epsilon_i^{n_i} \tilde{\lambda}(t') \pmod{p^{v_p(\tilde{\lambda}(t))+1} \mathcal{O}_{\bar{\mathbb{Q}}_p}}.$$

The last equation also implies $v_p(\tilde{\lambda}(t)) = v_p(\tilde{\lambda}(t'))$. Since t was strictly dominant this shows in particular that $t' \in \tilde{\mathbf{T}}(\bar{\mathbb{Q}}_p)^{++}$.

It only remains to prove uniqueness of t' . Let $t'' \in \tilde{T}(\bar{\mathbb{Q}}_p)^{++}$ be another element which is $\tilde{G}(\bar{\mathbb{Q}}_p)$ -conjugate to $x_s = (\beta t)_s$. Hence, $t'' = gt'g^{-1}$ for some $g \in \tilde{G}(\bar{\mathbb{Q}}_p)$. Since t', t'' are regular we know that $\tilde{T}(\bar{\mathbb{Q}}_p)$ is the centralizer of t' and of t'' . This implies $g\tilde{T}(\bar{\mathbb{Q}}_p)g^{-1} = \tilde{T}(\bar{\mathbb{Q}}_p)$, hence, g yields an element in $W_{\tilde{G}}$. Since t', t'' are both strictly dominant g is the unit element in $W_{\tilde{G}}$, i.e., $g \in \mathcal{C}(\tilde{T}(\bar{\mathbb{Q}}_p))$ (centralizer of $\tilde{T}(\bar{\mathbb{Q}}_p)$ in $\tilde{G}(\bar{\mathbb{Q}}_p)$), hence, $t'' = t'$. Thus, the proposition is proven. \square

4.4.1. Lemma. *The lattice $\Gamma_\pi \subseteq \mathfrak{h}^*$ is generated by dominant weights.*

Proof. We write the representation π of \mathfrak{g} defining $G = G_\pi$ as $\pi = \bigoplus_{i=1}^n \rho_{\lambda_i}$ with dominant weights $\lambda_i \in \Gamma_\pi$. Since $P_\pi = \bigcup_i P_{\lambda_i}$ we obtain

$$\Gamma_\pi = \langle P_\pi \rangle = \langle \bigcup_i P_{\lambda_i} \rangle \subseteq \langle \bigcup_i \lambda_i + \Gamma_{\text{ad}} \rangle \subseteq \Gamma_\pi;$$

hence, $\Gamma_\pi = \langle \bigcup_i \lambda_i + \Gamma_{\text{ad}} \rangle$ (for the last inclusion note that π is faithful; hence, $\Gamma_{\text{ad}} \subseteq \Gamma_\pi$). We select a weight $\gamma \in \Gamma_{\text{ad}}$ which for the moment is arbitrary and we put $\mu_i = \lambda_i + \gamma, i = 1, \dots, n$. Since Γ_{ad} is generated by the simple roots we obtain

$$\Gamma_\pi = \langle \bigcup_i \mu_i + \Gamma_{\text{ad}} \rangle = \langle \mu_i, \mu_i - \alpha, i = 1, \dots, n, \alpha \in \Delta \rangle.$$

If we choose the weight $\gamma \in \Gamma_{\text{ad}}$ dominant and sufficiently regular, i.e., $\langle \gamma, h_\beta \rangle > 0$ is positive and sufficiently large for all $\beta \in \Delta$ then μ_i and $\mu_i - \alpha$ are dominant for all $i = 1, \dots, n$ and all $\alpha \in \Delta$. This implies that Γ_π is generated by dominant weights. \square

4.5. We denote by ${}^w\chi$ or sometimes by $w\chi, \chi \in X(\tilde{T}), w \in W_{\tilde{G}}$, the character sending t to $\chi(w^{-1}tw)$. We write $\rho = \rho_{\tilde{G}} \in \Gamma_{\text{sc}}$ for the half sum of the positive roots and we put

$$w \cdot \tilde{\lambda} = w(\tilde{\lambda} + \rho^\circ) - \rho^\circ \in X(\tilde{T}) \quad \text{for } \tilde{\lambda} \in X(\tilde{T}),$$

with $\rho^\circ = \frac{1}{2} \sum_{\alpha \in \Phi^+} \alpha \in X(\tilde{T}) \otimes \mathbb{Q}$, where $\alpha = \alpha^\circ \in X(\tilde{T})$ is the exponential of the root α ; see Section 1.2.

Lemma. *Let $\tilde{\lambda} \in X(\tilde{T})$ be a dominant weight. For any $w \in W_{\tilde{G}}, w \neq 1$, and any $t \in \tilde{T}(\bar{\mathbb{Q}}_p)$ we have $w\tilde{\lambda}(t) = \tilde{\lambda}(t) (\sum_{\alpha \in \Delta} -b_\alpha \alpha)(t)$, where $b_\alpha \in \mathbb{N}_0$ and*

$$b_{\alpha_0} \geq \frac{\langle \tilde{\lambda}, \alpha_0^\vee \rangle}{2}$$

for at least one root $\alpha_0 \in \Delta$. Also, if $w \neq 1$ we have $w \cdot \tilde{\lambda}(t) = \tilde{\lambda}(t) (\sum_{\alpha \in \Delta} -b_\alpha \alpha)(t)$, where $b_\alpha \in \mathbb{N}_0$ and

$$b_{\alpha_0} \geq \frac{\langle \tilde{\lambda}, \alpha_0^\vee \rangle}{2}$$

for at least one root $\alpha_0 \in \Delta$.

Proof. We prove the claim about $w \cdot \tilde{\lambda}$. We write $\tilde{\lambda} = \tilde{\lambda}|_{\tilde{Z}} \lambda^\circ$, where $\lambda^\circ = \tilde{\lambda}|_T$ and $t = zt^\circ \in \tilde{T}(\bar{\mathbb{Q}}_p)$, where $z \in \tilde{Z}(\bar{\mathbb{Q}}_p)$ and $t^\circ \in T(\bar{\mathbb{Q}}_p)$. Since $w\rho^\circ - \rho^\circ \in \sum_{\alpha \in \Delta} \mathbb{Z}\alpha$ we obtain $w \cdot \tilde{\lambda}(z) = (w\tilde{\lambda} + (w\rho^\circ - \rho^\circ))(z) = \tilde{\lambda}(z)$. Since $\tilde{\lambda} = \lambda^\circ$ on T we obtain $w \cdot \tilde{\lambda}(t^\circ) = w \cdot \lambda^\circ(t^\circ)$. To determine $w \cdot \lambda^\circ(t^\circ)$ we set as before $\lambda = d\lambda^\circ \in \Gamma_\pi$, i.e., λ° corresponds to λ under the isomorphism $(\cdot)^\circ : \Gamma_\pi \rightarrow X(T)$ (see Section 1.2). The Weyl group $W_{\tilde{G}}$ acts on $\Gamma_\pi \subseteq \mathfrak{h}^*$ via $\lambda \mapsto \lambda \circ \text{Ad}(w^{-1})$, $w \in W_{\tilde{G}}$, and since $(\cdot)^\circ$ is equivariant w.r.t. the action of W_G (see [B], Section 3.3, Remarks (1), p. 16) we obtain $w \cdot \lambda^\circ = (w \cdot \lambda)^\circ$, where $w \cdot \lambda := w(\lambda + \rho) - \rho$. Since $w\lambda$ is a weight of the irreducible \mathfrak{g} -module of highest weight λ we know that

$$w\lambda = \lambda - \sum_{\alpha \in \Delta} c_\alpha \alpha$$

for certain $c_\alpha \in \mathbb{N}_0$. Since λ is a dominant element in the weight lattice Γ_{sc} we may write $\lambda = \sum_{\alpha \in \Delta} d_\alpha \omega_\alpha$, where ω_α , $\alpha \in \Delta$, are the fundamental weights and $d_\alpha \in \mathbb{N}_0$. On the other hand, $w \neq 1$ implies that $w\lambda$ is not contained in the Weyl chamber corresponding to the basis Δ , hence, $\langle w\lambda, h_{\alpha_0} \rangle \leq 0$ for some root $\alpha_0 \in \Delta$. We obtain

$$0 \geq \langle w\lambda, h_{\alpha_0} \rangle = \left\langle \sum_{\alpha \in \Delta} d_\alpha \omega_\alpha - \sum_{\alpha \in \Delta} c_\alpha \alpha, h_{\alpha_0} \right\rangle = d_{\alpha_0} - \sum_{\alpha \in \Delta} c_\alpha \langle \alpha, h_{\alpha_0} \rangle.$$

Since $\langle \alpha, h_{\alpha_0} \rangle = 2$ if $\alpha = \alpha_0$ and $\langle \alpha, h_{\alpha_0} \rangle \leq 0$ if $\alpha \neq \alpha_0$ this yields $0 \geq d_{\alpha_0} - 2c_{\alpha_0}$. Thus,

$$c_{\alpha_0} \geq \frac{1}{2}d_{\alpha_0} = \frac{1}{2}\langle \lambda, h_{\alpha_0} \rangle = \frac{1}{2}\langle \lambda^\circ, \alpha_0^\vee \rangle = \frac{1}{2}\langle \tilde{\lambda}, \alpha_0^\vee \rangle.$$

Altogether we obtain

$$\begin{aligned} (w \cdot \tilde{\lambda})(t) &= \tilde{\lambda}(z) (w \cdot \lambda^\circ)(t^\circ) = \tilde{\lambda}(z) (w \cdot \lambda)^\circ(t^\circ) = \tilde{\lambda}(z) (w\lambda + w\rho - \rho)^\circ(t^\circ) \\ &= \tilde{\lambda}(z) \left(\lambda + \sum_{\alpha \in \Delta} -c_\alpha \alpha + w\rho - \rho \right)^\circ(t^\circ) = \tilde{\lambda}(t) \left(\sum_{\alpha \in \Delta} -c_\alpha \alpha + w\rho - \rho \right)^\circ(t^\circ). \end{aligned}$$

Since $w\rho - \rho \in \mathbb{Z}\Phi$ is a sum of negative roots, this shows that $w \cdot \tilde{\lambda}(t)$ has the claimed form (note that $\alpha(t) = \alpha(t^\circ)$ since $\alpha = \alpha^\circ$ vanishes on $\tilde{Z}(\bar{\mathbb{Q}}_p)$). The claim about $w\tilde{\lambda}$ follows analogously. \square

4.6. Notation. We recall that in Section 2.3 we selected an element $h \in \tilde{T}(\mathbb{Q})^{++}$. We set

$$\kappa_1 = \kappa_{1, \tilde{G}}(h) = \sum_{\alpha \in \Phi_{\tilde{G}}^+} v_p(\alpha(h)) \in \mathbb{N}.$$

Thus, κ_1 depends on \tilde{G} and h . Since $\rho - {}^w\rho$, $w \in W_{\tilde{G}}$, is a sum of certain positive roots all of which occur with multiplicity 1 we obtain

$$v_p((\rho^\circ - {}^w\rho^\circ)(h^e)) \leq e\kappa_1.$$

We write $2\rho = 2\rho_{\tilde{G}} = \sum_{\beta \in \Delta_{\tilde{G}}} m_{\beta} \beta$, where $m_{\beta} \in \mathbb{N}_0$ for all $\beta \in \Delta_{\tilde{G}}$, and we set

$$\kappa_2 = \kappa_{2, \tilde{G}} = \max_{\beta \in \Delta_{\tilde{G}}} m_{\beta} \in \mathbb{N}_0;$$

i.e., κ_2 is the maximum multiplicity with which a simple root can occur in $2\rho_{\tilde{G}}$ and only depends on \tilde{G} . Since $\rho - {}^w\rho$ is a sum of certain positive roots each of which occurs with multiplicity 1 we can write $\rho - {}^w\rho = \sum_{\beta \in \Delta_{\tilde{G}}} n_{\beta} \beta$ where $n_{\beta} \in \mathbb{N}_0$ and $n_{\beta} \leq m_{\beta}$ for all $\beta \in \Delta_{\tilde{G}}$. Since $\langle \beta, \alpha^{\vee} \rangle$, $\alpha, \beta \in \Delta_{\tilde{G}}$, equals 2 if $\alpha = \beta$ and is ≤ 0 otherwise we obtain for all $\alpha \in \Delta_{\tilde{G}}$

$$\langle \rho - {}^w\rho, \alpha^{\vee} \rangle \leq 2n_{\alpha} \leq 2m_{\alpha} \leq 2\kappa_2.$$

Let $\tilde{P}/\mathbb{Q}_p \leq \tilde{G}/\mathbb{Q}_p$ be a standard parabolic subgroup (i.e., $\tilde{P}/\mathbb{Q}_p \supseteq \tilde{B}/\mathbb{Q}_p$) with Levi decomposition $\tilde{P} = \tilde{M}\tilde{N}$. The Levi subgroup \tilde{M} contains the \mathbb{Q}_p -split maximal torus \tilde{T} and we denote by $W^{\tilde{P}}$ the set of Kostant representatives for the quotient of Weyl groups $W_{\tilde{M}} \backslash W_{\tilde{G}}$. The intersection $\tilde{B} \cap \tilde{M} \leq \tilde{M}/\mathbb{Q}_p$ is a Borel subgroup in \tilde{M}/\mathbb{Q}_p and for any dominant (w.r.t. to $\tilde{B} \cap \tilde{M}$) weight $\tilde{\lambda} \in X(\tilde{T})$ we then denote by $\rho_{\tilde{\lambda}}^{\tilde{M}} : \tilde{M} \rightarrow \mathbf{Aut}(L_{\tilde{\lambda}}^{\tilde{M}})$ the irreducible representation of \tilde{M}/\mathbb{Q}_p of highest weight $\tilde{\lambda}$ (see Section 2.4). Any $p \in \tilde{P}(\bar{\mathbb{Q}}_p)$ can be written $p = \bar{p}u$, $\bar{p} \in \tilde{M}(\bar{\mathbb{Q}}_p)$, $u \in \tilde{N}(\bar{\mathbb{Q}}_p)$ and we denote by $v_{\bar{p}} : \tilde{P}/\mathbb{Q}_p \rightarrow \tilde{M}/\mathbb{Q}_p$, $p \mapsto \bar{p}$, the morphism to the Levi subgroup.

In the next proposition we will use the following notation: for $c \in \mathbb{Q}$ and $x, y \in \bar{\mathbb{Q}}_p$ we write $x \equiv y \pmod{p^c \mathcal{O}_{\bar{\mathbb{Q}}_p}}$ to denote that $v_p(x - y) \geq c$. Thus, in case $c \in \mathbb{Z}$ the term “ $x \equiv y \pmod{p^c \mathcal{O}_{\bar{\mathbb{Q}}_p}}$ ” has two meanings that coincide.

4.7. Proposition. *Let $C \in \mathbb{Q}_{>0}$ and assume that $\tilde{\lambda} \in X(\tilde{T})^{\text{dom}}$ satisfies*

$$\langle \tilde{\lambda}, \alpha^{\vee} \rangle > 2C$$

for all $\alpha \in \Delta_{\tilde{G}}$. Select a standard parabolic subgroup $\tilde{P}/\mathbb{Q}_p \leq \tilde{G}/\mathbb{Q}_p$ with Levi decomposition $\tilde{P} = \tilde{M}\tilde{N}$ and let $\xi \in \tilde{M}(\mathbb{Q}_p)$. Assume that there is $u \in \tilde{N}(\mathbb{Q}_p)$ such that $\xi u \in \tilde{P}(\mathbb{Q}_p)$ is $\tilde{G}(\mathbb{Q}_p)$ -conjugate to an element in $\tilde{I}h^e\tilde{I}$, $e \in \mathbb{N}$. We denote by $t = t_{\xi u}$ the unique element in $\tilde{T}(\bar{\mathbb{Q}}_p)^{++}$ which is $\tilde{G}(\bar{\mathbb{Q}}_p)$ -conjugate to $(\xi u)_s$ (see 4.4 Proposition; note that ξu is conjugate to an element $x \in \tilde{I}h^e\tilde{I}$). Then there is an element $s = s_{\xi u} \in W^{\tilde{P}}$ such that for all $w \in W^{\tilde{P}}$ the following congruence holds:

$$\tilde{\lambda}(h^e) \text{tr}(\xi^{-1} | L_{w, \tilde{\lambda}}^{\tilde{M}}(\mathbb{Q}_p)) \equiv \begin{cases} \varepsilon_{\bar{p}} \tilde{\lambda}(h^e t^{-1}) \pmod{p^{(C-\kappa_1-\kappa_2)e} \mathcal{O}_{\bar{\mathbb{Q}}_p}} & \text{if } w = s, \\ 0 \pmod{p^{(C-\kappa_1-\kappa_2)e} \mathcal{O}_{\bar{\mathbb{Q}}_p}} & \text{if } w \neq s. \end{cases}$$

Here, $\varepsilon_{\bar{p}} = \varepsilon_{\bar{p}, \xi u}$ is an element in $\bar{\mathbb{Q}}_p$ which does not depend on $\tilde{\lambda}$ and satisfies

$$v_p(\varepsilon_{\bar{p}, \xi u}) \geq -e\kappa_1.$$

Proof. By 12.1.7(c) (p. 211) of [Springer 1981], the semisimple part $(\xi u)_s$ is contained in $\tilde{P}(\mathbb{Q}_p)$, and Theorem 6.4.5(ii) (p. 109) of the same reference shows that it is contained in a maximal $\bar{\mathbb{Q}}_p$ -torus $\tilde{S} = \tilde{S}_{\xi u}$ in $\tilde{P}/\bar{\mathbb{Q}}_p$. Since all maximal tori in $\tilde{P}/\bar{\mathbb{Q}}_p$ are conjugate there is $y \in \tilde{P}(\bar{\mathbb{Q}}_p)$ such that $\tilde{S} = {}^y\tilde{T}$. In particular there is $t' \in \tilde{T}(\bar{\mathbb{Q}}_p)$ such that

$$(\xi u)_s = {}^y t'.$$

Modifying y by an element in $W_{\tilde{M}}$ we may even assume that t' is dominant w.r.t. $\Delta_{\tilde{M}}$, i.e., $v_p(\alpha(t')) \geq 0$ for all $\alpha \in \Delta_{\tilde{M}}$.

Since ξu is conjugate to an element $x \in \tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$ the semisimple part $(\xi u)_s$ is $\tilde{G}(\bar{\mathbb{Q}}_p)$ -conjugate to an element $t = t_{\xi u} \in \tilde{T}(\bar{\mathbb{Q}}_p)^{++}$ satisfying

$$(39) \quad v_p(\chi(t)) = v_p(\chi(h^e)) \quad \text{and} \quad \chi(t) \equiv \epsilon \chi(h^e) \pmod{p^{v_p(\chi(t))+1} \mathcal{O}_{\bar{\mathbb{Q}}_p}}$$

for all $\chi \in X(\tilde{T})$ where $\epsilon = \epsilon_\chi \in \mathbb{Z}_p^*$ (see 4.4 Proposition). Since t' also is $\tilde{G}(\bar{\mathbb{Q}}_p)$ -conjugate to $(\xi u)_s$ we find that $t, t' \in \tilde{T}(\bar{\mathbb{Q}}_p)$ are conjugate by an element $s = s_{\xi u} \in \tilde{G}(\bar{\mathbb{Q}}_p)$:

$$t' = {}^s t.$$

Since t is regular, t' also is regular, and it follows that $\tilde{T} = C(t)^0 = C(t')^0$. This implies ${}^s\tilde{T} = \tilde{T}$; hence, s is contained in the normalizer of \tilde{T} and we therefore can select $s \in W_{\tilde{G}}$ (i.e., s is representative of an element in $W_{\tilde{G}}$). Since t' is dominant w.r.t. $\Delta_{\tilde{M}}$ and t is strictly dominant w.r.t. $\Delta_{\tilde{G}}$ we see that s^{-1} maps $\Phi_{\tilde{M}}^+$ to $\Phi_{\tilde{G}}^+$, which implies $s \in W_{\tilde{P}}$. Denote by $L_{w, \tilde{\lambda}}^{\tilde{P}}$ the extension of the representation $L_{w, \tilde{\lambda}}^{\tilde{M}}$ to \tilde{P}/\mathbb{Q}_p via the morphism $v_{\tilde{P}} : \tilde{P} \rightarrow \tilde{M}$. We have obtained

$$\begin{aligned} \text{tr}(\xi^{-1} | L_{w, \tilde{\lambda}}^{\tilde{M}}(\mathbb{Q}_p)) &= \text{tr}(\xi^{-1} | L_{w, \tilde{\lambda}}^{\tilde{P}}(\mathbb{Q}_p)) = \text{tr}((\xi u)^{-1} | L_{w, \tilde{\lambda}}^{\tilde{P}}(\mathbb{Q}_p)) \\ &= \text{tr}((\xi u)_s^{-1} | L_{w, \tilde{\lambda}}^{\tilde{P}}(\mathbb{Q}_p)) = \text{tr}((t')^{-1} | L_{w, \tilde{\lambda}}^{\tilde{M}}(\mathbb{Q}_p)) \\ &= \text{tr}({}^s t^{-1} | L_{w, \tilde{\lambda}}^{\tilde{M}}(\mathbb{Q}_p)). \end{aligned}$$

The Weyl character formula (see [Popov and Vinberg 1994, I.4.6.4 Theorem, p. 45] or [Jantzen 2003, II.5.10 Proposition, p. 223]) then yields

$$(40) \quad \tilde{\lambda}(h^e) \text{tr}(\xi^{-1} | L_{w, \tilde{\lambda}}^{\tilde{M}}(\mathbb{Q}_p)) = \tilde{\lambda}(h^e) \frac{\sum_{v \in W_{\tilde{M}}} (-1)^{\ell(v)} v \cdot \tilde{M}(w \cdot \tilde{\lambda})({}^s t^{-1})}{\prod_{\alpha \in \Phi_{\tilde{M}}^+} (1 - \alpha^{-1}(\underbrace{{}^s t^{-1}}_{= t'^{-1}}))}.$$

Here, we use the notation $v \cdot_{\tilde{M}} \tilde{\mu} = v(\tilde{\mu} + \rho_{\tilde{M}}^\circ) - \rho_{\tilde{M}}^\circ$. We denote the denominator appearing on the right-hand side of (40) by

$$N(t') = N_{\tilde{M}}(t').$$

For all $\alpha \in \Phi_{\tilde{G}}$ equation (39) implies that $v_p(\alpha(t)) \geq e$ or $v_p(\alpha(t)) \leq -e$. Since $\alpha({}^s t) = (s^{-1}\alpha)(t)$ the same then is true of $t' = {}^s t$, i.e., $v_p(\alpha(t')) \geq e$ or $v_p(\alpha(t')) \leq -e$ for all $\alpha \in \Phi_{\tilde{G}}$. Since $v_p(\alpha(t')) \geq 0$ for all $\alpha \in \Phi_{\tilde{M}}^+$ we obtain

$$(41) \quad v_p(\alpha(t')) \geq e \quad \text{for all } \alpha \in \Phi_{\tilde{M}}^+.$$

Hence,

$$v_p(N(t')) = v_p\left(\prod_{\alpha \in \Phi_{\tilde{M}}^+} (1 - \alpha(t'))\right) = 0,$$

i.e., $N(t')$ is a p -adic unit in $\mathcal{O}_{\tilde{\mathbb{Q}}_p}$; in particular, $N(t') \neq 0$.

We look at the individual summands indexed by $v \in W_{\tilde{M}}$ which appear in the numerator in equation (40) and distinguish cases.

4.7.1. We first assume that $v \neq 1$. Using 4.5 Lemma with \tilde{M} in place of \tilde{G} and the definition of $w \cdot \tilde{\lambda}$ we can write

$$(42) \quad \begin{aligned} \tilde{\lambda}(h^e) v \cdot_{\tilde{M}} (w \cdot \tilde{\lambda})({}^s t^{-1}) &= \tilde{\lambda}(h^e) (w \cdot \tilde{\lambda})({}^s t^{-1}) \left(\sum_{\alpha \in \Delta_{\tilde{M}}} -b_{v,\alpha} \alpha \right) ({}^s t^{-1}) \\ &= \tilde{\lambda}(h^e) (s^{-1} w \tilde{\lambda})(t^{-1}) (w \rho^\circ - \rho^\circ) ({}^s t^{-1}) \left(\sum_{\alpha \in \Delta_{\tilde{M}}} b_{v,\alpha} \alpha \right) ({}^s t) \end{aligned}$$

with $b_{v,\alpha} \in \mathbb{N}_0$ and $b_{v,\alpha_v} \geq \frac{1}{2} \langle w \cdot \tilde{\lambda}, \alpha_v^\vee \rangle$ for (at least) one root $\alpha_v \in \Delta_{\tilde{M}}$ (note that $w \cdot \tilde{\lambda}$ is dominant for $\Delta_{\tilde{M}}$ since $w \in W^{\tilde{P}}$). Since $w \in W^{\tilde{P}}$ we know that $w^{-1} \alpha_v \in \Phi_{\tilde{G}}^+$, hence, we obtain

$$b_{v,\alpha_v} \geq \frac{\langle w \cdot \tilde{\lambda}, \alpha_v^\vee \rangle}{2} = \frac{\langle \tilde{\lambda}, w^{-1} \alpha_v^\vee \rangle}{2} + \frac{\langle w \rho^\circ - \rho^\circ, \alpha_v^\vee \rangle}{2} \geq C - \kappa_2.$$

Equation (41) implies that

$$(43) \quad v_p\left(\left(\sum_{\alpha \in \Delta_{\tilde{M}}} b_{v,\alpha} \alpha\right)({}^s t)\right) \geq (C - \kappa_2)e.$$

Since $w\rho - \rho$ is a sum of certain negative roots all appearing with multiplicity 1 and since t is strictly dominant we obtain using equation (39)

$$(44) \quad v_p((w\rho^\circ - \rho^\circ)({}^s t^{-1})) \geq - \sum_{\alpha \in \Phi_{\tilde{G}}^+} v_p(\alpha(t)) = - \sum_{\alpha \in \Phi_{\tilde{G}}^+} v_p(\alpha(h^e)) = -e\kappa_1.$$

If $s^{-1}w \neq 1$ then 4.5 Lemma yields

$$\tilde{\lambda}(h^e) (s^{-1} w \tilde{\lambda})(t^{-1}) = \tilde{\lambda}(h^e t^{-1}) \left(\sum_{\alpha \in \Delta_{\tilde{G}}} -c_{w,\alpha} \alpha \right) (t^{-1}),$$

where $c_{w,\alpha} \in \mathbb{N}_0$ and $c_{w,\alpha_w} \geq \frac{1}{2} \langle \tilde{\lambda}, \alpha_w^\vee \rangle \geq C$ for (at least) one root $\alpha_w \in \Delta_{\tilde{G}}$. It follows

from (39) that $\tilde{\lambda}(h^e)\tilde{\lambda}(t^{-1})$ is a p -adic unit and that $v_p(\alpha(t)) = v_p(\alpha(h^e)) \geq e$ for all $\alpha \in \Delta_{\tilde{G}}$; hence,

$$(45) \quad v_p(\tilde{\lambda}(h^e)(s^{-1}w\tilde{\lambda})(t^{-1})) = v_p\left(\left(\sum_{\alpha \in \Delta_{\tilde{G}}} c_{w,\alpha}\alpha\right)(t)\right) \geq Ce (> 0).$$

If $s = w$ then $\tilde{\lambda}(h^e)(s^{-1}w\tilde{\lambda})(t^{-1}) = \tilde{\lambda}(h^e t^{-1})$ is a p -adic integer. Thus, if $v \neq 1$ equations (42), (43), (44), (45) yield

$$(46) \quad v_p(\tilde{\lambda}(h^e) v \cdot_{\tilde{M}} (w \cdot \tilde{\lambda})(s t^{-1})) \geq (C - \kappa_1 - \kappa_2)e$$

for all $w \in W^{\tilde{P}}$. Hence, modulo $p^{(C-\kappa_1-\kappa_2)e} \mathcal{O}_{\tilde{\mathbb{Q}}_p}$ we may neglect all summands with $v \neq 1$.

4.7.2. We assume $v = 1$. If $w \neq s$ equations (42), (44), (45) yield

$$(47) \quad v_p(\tilde{\lambda}(h^e) v \cdot_{\tilde{M}} (w \cdot \tilde{\lambda})(s t^{-1})) = v_p(\tilde{\lambda}(h^e)(s^{-1}w\tilde{\lambda})(t^{-1})(w\rho^\circ - \rho^\circ)(s t^{-1})) \geq (C - \kappa_1)e.$$

If $w = s$ we obtain as above

$$(48) \quad \tilde{\lambda}(h^e) v \cdot_{\tilde{M}} (w \cdot \tilde{\lambda})(s t^{-1}) = \tilde{\lambda}(h^e)\tilde{\lambda}(t^{-1})(s\rho^\circ - \rho^\circ)(s t^{-1}).$$

4.7.3. Taking into account that $C - \kappa_1$ is bigger than or equal to $C - \kappa_1 - \kappa_2$, equations (40) and (46), (47), (48) now yield (note that $N(t')$ is a p -adic unit)

$$\begin{aligned} &\tilde{\lambda}(h^e) \operatorname{tr}(\xi^{-1} | L_{w \cdot \tilde{\lambda}}^{\tilde{M}}(\mathbb{Q}_p)) \\ &\equiv \begin{cases} \frac{\tilde{\lambda}(h^e)\tilde{\lambda}(t^{-1})(s\rho^\circ - \rho^\circ)(s t^{-1})}{N(t')} \pmod{p^{(C-\kappa_1-\kappa_2)e} \mathcal{O}_{\tilde{\mathbb{Q}}_p}} & \text{if } w = s, \\ 0 \pmod{p^{(C-\kappa_1-\kappa_2)e} \mathcal{O}_{\tilde{\mathbb{Q}}_p}} & \text{if } w \neq s. \end{cases} \end{aligned}$$

We put $\varepsilon_{\tilde{P}} = \varepsilon_{\tilde{P}, \xi u} = \frac{(s\rho^\circ - \rho^\circ)(s t^{-1})}{N(t')} \in \tilde{\mathbb{Q}}_p$. Since $N(t')$ is a p -adic unit, equation (44) shows that

$$v_p(\varepsilon_{\tilde{P}, \xi u}) \geq -e\kappa_1.$$

This completes the proof of the proposition. □

4.8. We look at the special case $\tilde{P} = \tilde{G}$ in 4.7 Proposition which is sufficient for application to Bewersdorff’s trace formula. (The general case will be needed in application to the Goresky–MacPherson trace formula which involves contributions from parabolic subgroups of \tilde{G} as well; see Section 4.14). In this case, $\xi \in \tilde{G}(\mathbb{Q}_p)$, $u = 1$, $W^{\tilde{P}} = 1$ and $s = w = 1$; hence, $t = t_\xi = t'$. In particular, $\varepsilon_{\tilde{P}} = \varepsilon_{\tilde{G}, 1, \xi} = 1/N(t_\xi)$ is a p -adic integer. Moreover, we can choose $\kappa_1 = \kappa_2 = 0$ since then the equations involving ρ and κ_1, κ_2 in Section 4.6 still hold (note that $w = 1$; see also equation (44) and the equation following (42)). We thus obtain:

Corollary. *Let $C \in \mathbb{Q}_{>0}$ and assume that $\tilde{\lambda} \in X(\tilde{T})^{\text{dom}}$ satisfies*

$$\langle \tilde{\lambda}, \alpha^\vee \rangle > 2C$$

for all $\alpha \in \Delta$. Then for any $\xi \in \tilde{G}(\mathbb{Q}_p)$ which is $\tilde{G}(\mathbb{Q}_p)$ -conjugate to an element in $\tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$, $e \in \mathbb{N}$, the following congruence holds:

$$\tilde{\lambda}(h^e) \text{tr}(\xi^{-1} | L_{\tilde{\lambda}}(\mathbb{Q}_p)) \equiv \frac{\tilde{\lambda}(h^e t_\xi^{-1})}{N(t_\xi)} \pmod{p^{Ce} \mathcal{O}_{\tilde{\mathbb{Q}}_p}}.$$

Here, $t_\xi \in \tilde{T}(\tilde{\mathbb{Q}}_p)^{++}$ denotes the unique element which is $\tilde{G}(\tilde{\mathbb{Q}}_p)$ -conjugate to ξ_s (see 4.4 Proposition) and

$$N(t_\xi) = \prod_{\alpha \in \Phi_G^+} (1 - \alpha(t_\xi))$$

is a p -adic unit in $\mathcal{O}_{\tilde{\mathbb{Q}}_p}$.

4.9. Proposition. *Let $C \in \mathbb{Q}_{>0}$ and assume that $\tilde{\lambda} \in X(\tilde{T})^{\text{dom}}$ satisfies*

$$\langle \tilde{\lambda}, \alpha^\vee \rangle > 2C$$

for all $\alpha \in \Delta_{\tilde{G}}$. Let ζ be contained in the semigroup Δ , so $\zeta \in \tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$ for some $e \in \mathbb{N}_0$ (by 2.7 Lemma); we assume that $e \in \mathbb{N}$. Then the Lefschetz number $\text{Lef}((\Gamma\zeta\Gamma)_{\tilde{\lambda}} | H^\bullet(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{Q}_p)))$ is contained in \mathbb{Z}_p and the following congruence holds:

$$\text{Lef}((\Gamma\zeta\Gamma)_{\tilde{\lambda}} | H^\bullet(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{Q}_p))) \equiv \sum_{[\xi]_\Gamma \in \Gamma\zeta\Gamma/\sim_\Gamma} c_{[\xi]_\Gamma} \frac{\tilde{\lambda}(h^e t_\xi^{-1})}{N(t_\xi)} \pmod{p^{Ce} \mathcal{O}_{\tilde{\mathbb{Q}}_p}}.$$

Proof. By 2.9 Corollary, the Lefschetz numbers of normalized Hecke operators are contained in \mathbb{Z}_p . Since $\zeta \in \tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$ we know that $(\Gamma\zeta\Gamma)_{\tilde{\lambda}} = \tilde{\lambda}(h^e)\Gamma\zeta\Gamma$. On the other hand, a representative ξ of a Γ -conjugacy class contained in $\Gamma\zeta\Gamma$ is contained in $\tilde{G}(\mathbb{Q})$ and in $\tilde{\mathcal{I}}\zeta\tilde{\mathcal{I}} = \tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$. The second claim thus follows from Bewersdorff's trace formula (see 4.2 Theorem) and 4.8 Corollary (note that $F \subseteq \mathbb{Q}_p$). \square

4.10. Proposition. *Let $C \in \mathbb{Q}_{>0}$ and let the dominant weights $\tilde{\lambda}, \tilde{\lambda}' \in X(\tilde{T})$ satisfy*

- $\langle \tilde{\lambda}, \alpha^\vee \rangle > 2C$ and $\langle \tilde{\lambda}', \alpha^\vee \rangle > 2C$ for all $\alpha \in \Delta_{\tilde{G}}$;
- $\tilde{\lambda} \equiv \tilde{\lambda}' \pmod{(p-1)p^{m-1}X(\tilde{T})}$ ($m \in \mathbb{N}$).

Let ζ be contained in the semigroup Δ , so $\zeta \in \tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$ for some $e \in \mathbb{N}_0$ which we assume positive. Then the Lefschetz number $\text{Lef}((\Gamma\zeta\Gamma)_{\tilde{\lambda}} | H^\bullet(\Gamma \backslash X, L_{\tilde{\lambda}}(F)))$ is contained in \mathbb{Z}_p (note that $F \subseteq \mathbb{Q}_p$) and the following congruence holds:

$$\begin{aligned} &\text{Lef}((\Gamma\zeta\Gamma)_{\tilde{\lambda}} | H^\bullet(\Gamma \backslash X, L_{\tilde{\lambda}}(F))) \\ &\equiv \text{Lef}((\Gamma\zeta\Gamma)_{\tilde{\lambda}'} | H^\bullet(\Gamma \backslash X, L_{\tilde{\lambda}'}(F))) \pmod{p^{\lceil \min(m, Ce) \rceil} \mathbb{Z}_p}. \end{aligned}$$

Proof. Since $\text{Lef}((\Gamma\zeta\Gamma)_{\tilde{\lambda}}|H^\bullet(\Gamma\backslash X, L_{\tilde{\lambda}}(F))) = \text{Lef}((\Gamma\zeta\Gamma)_{\tilde{\lambda}}|H^\bullet(\Gamma\backslash X, L_{\tilde{\lambda}}(\mathbb{Q}_p)))$, in order to prove the proposition we may consider Lefschetz numbers over \mathbb{Q}_p . Integrality of the Lefschetz number then follows from 4.9 Proposition. We let $[\xi]_\Gamma$ be any Γ -conjugacy class contained in $\Gamma\zeta\Gamma$. Hence, $\xi \in \tilde{T}h^e\tilde{T}$ and we denote by $t_\xi \in \tilde{T}(\bar{\mathbb{Q}}_p)^{++}$ the element which is $\tilde{G}(\bar{\mathbb{Q}}_p)$ -conjugate to ξ_s (see 4.4 Proposition). Since $\tilde{\lambda} \equiv \tilde{\lambda}' \pmod{(p-1)p^{m-1}X(\tilde{T})}$ there is $\chi \in X(\tilde{T})$ such that $\tilde{\lambda} - \tilde{\lambda}' = (p-1)p^{m-1}\chi$. Taking into account that $\chi(h^e t_\xi^{-1}) \equiv \epsilon \pmod{p\mathcal{O}_{\bar{\mathbb{Q}}_p}}$ by 4.4 Proposition, where $\epsilon = \epsilon_\chi \in \mathbb{Z}_p^*$, we therefore obtain

$$\frac{\tilde{\lambda}(h^e t_\xi^{-1})}{\tilde{\lambda}'(h^e t_\xi^{-1})} = \chi(h^e t_\xi^{-1})^{(p-1)p^{m-1}} \in 1 + p^m \mathcal{O}_{\bar{\mathbb{Q}}_p}.$$

Since also $\tilde{\lambda}'(h^e t_\xi^{-1})$ is a p -adic unit by the same proposition, this implies

$$\tilde{\lambda}(h^e t_\xi^{-1}) \equiv \tilde{\lambda}'(h^e t_\xi^{-1}) \pmod{p^m \mathcal{O}_{\bar{\mathbb{Q}}_p}}.$$

The claim now follows from 4.9 Proposition taking into account that the Lefschetz numbers are contained in \mathbb{Q}_p , hence, their p -adic valuations are integers and that $c_{[\xi]_\Gamma} \in \mathbb{Z}$ and $N(t_\xi)$ is a p -adic unit. Thus, the proof is complete. \square

Remark. The proposition also holds trivially for $e = 0$ since both sides of the congruence are integers by 2.9 Corollary.

4.11. Adelic formulation. Using adelic formulation in Section 4.13 we will prove congruences between traces of Hecke operators on Eisenstein cohomology and, hence, on cuspidal cohomology. In this section we therefore reformulate 3.10 Corollary and 4.10 Proposition in adelic language.

We denote by \mathbb{A} (resp. \mathbb{A}_f) the ring of adèles (resp. of finite adèles) of \mathbb{Q} . For any compact open subgroup $\tilde{K} \leq \tilde{G}(\mathbb{A}_f)$ we set $S_{\tilde{K}} = \tilde{G}(\mathbb{Q}) \backslash \tilde{G}(\mathbb{A}) / \tilde{K} \tilde{K}_\infty A_{\tilde{G}}$. We assume that \tilde{G}/\mathbb{Q} satisfies strong approximation; in particular, $\tilde{G}(\mathbb{A})$ is a finite disjoint union $\tilde{G}(\mathbb{A}) = \bigcup_{i=1}^t \tilde{G}(\mathbb{Q}) g_i \tilde{G}(\mathbb{R}) \tilde{K}$, $g_i \in \tilde{G}(\mathbb{A}_f)$, and we obtain

$$S_{\tilde{K}} = \bigcup_{i=1}^t \Gamma_i \backslash X$$

where

$$\Gamma_i = \tilde{G}(\mathbb{Q}) \cap g_i \tilde{K} g_i^{-1}.$$

We assume that we can choose a system of double coset representatives g_i as above which is contained in $\tilde{G}(\mathbb{A}_f)^{(p)}$, where $\tilde{G}(\mathbb{A}_f)^{(p)} \leq \tilde{G}(\mathbb{A}_f)$ is the subgroup consisting of elements whose p -component equals 1 (e.g., \tilde{G} satisfies weak approximation at p ; note that weak approximation holds for almost all primes p).

We fix a compact open subgroup $\tilde{K} = \prod_{\ell \neq \infty} \tilde{K}_\ell \leq \tilde{G}(\mathbb{A}_f)$ such that $\tilde{K}_p = \tilde{I}$ and we set $\tilde{K}^{(p)} = \prod_{\ell \neq p, \infty} \tilde{K}_\ell$. Since the p -component of g_i is equal to 1 any of the arithmetic subgroups Γ_i is contained in \tilde{I} .

4.11.1. Hecke algebra. We fix a Haar measure $dg = \otimes_{\ell \neq \infty} dg_\ell$ on $\tilde{G}(\mathbb{A}_f)$ and we denote by $\mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{A}_f))$ the Hecke algebra consisting of compactly supported smooth \mathbb{Q} -valued functions on $\tilde{G}(\mathbb{A}_f)$. We have the decomposition $\mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{A}_f)) = \mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{Q}_p)) \otimes \mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{A}_f)^{(p)})$, where the two factors are the Hecke algebras consisting respectively of compactly supported smooth \mathbb{Q} -valued functions on $\tilde{G}(\mathbb{Q}_p)$ and on $\tilde{G}(\mathbb{A}_f)^{(p)}$. Let the subalgebra consisting of \tilde{K} - (resp. \tilde{I} - or $\tilde{K}^{(p)}$ -) bi-invariant functions be denoted by $\mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{A}_f) // \tilde{K}) \leq \mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{A}_f))$ (resp. $\mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{Q}_p) // \tilde{I}) \leq \mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{Q}_p))$ or $\mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{A}_f)^{(p)} // \tilde{K}^{(p)}) \leq \mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{A}_f)^{(p)})$). Let

$$[\tilde{K}x\tilde{K}] = \frac{1}{\text{vol}(\tilde{K})} \mathbf{1}_{\tilde{K}x\tilde{K}}, \quad x \in \tilde{G}(\mathbb{A}_f),$$

where $\mathbf{1}_X$ is the characteristic function of the set X , and define likewise $[\tilde{I}x\tilde{I}]$ for $x \in \tilde{G}(\mathbb{Q}_p)$ and $[\tilde{K}^{(p)}x\tilde{K}^{(p)}]$ for $x \in \tilde{G}(\mathbb{A}_f)^{(p)}$, by replacing \tilde{K} with \tilde{I} and with $\tilde{K}^{(p)}$. The elements $[\tilde{K}x\tilde{K}]$ (resp. $[\tilde{I}x\tilde{I}]$ or $[\tilde{K}^{(p)}x\tilde{K}^{(p)}]$) form a \mathbb{Q} -basis of $\mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{A}_f) // \tilde{K})$ (resp. $\mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{Q}_p) // \tilde{I})$ or $\mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{A}_f)^{(p)} // \tilde{K}^{(p)})$); their \mathbb{Z} -spans define \mathbb{Z} -structures $\mathcal{C}_0(\tilde{G}(\mathbb{A}_f) // \tilde{K})$ (resp. $\mathcal{C}_0(\tilde{G}(\mathbb{Q}_p) // \tilde{I})$ or $\mathcal{C}_0(\tilde{G}(\mathbb{A}_f)^{(p)} // \tilde{K}^{(p)})$) in the respective Hecke algebras. Thus, the \mathbb{Z} -structure on $\mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{A}_f) // \tilde{K})$ is given as the subspace of $\text{vol}(\tilde{K})^{-1} \cdot \mathbb{Z}$ -valued functions and analogously for the other two Hecke algebras.

In Section 2.3 we selected an element $h \in \tilde{T}(\mathbb{Q})^{++}$ and we now denote by

$$\mathcal{C}_0(\tilde{G}(\mathbb{Q}_p) // \tilde{I})_h$$

the \mathbb{Z} -subalgebra of $\mathcal{C}_0(\tilde{G}(\mathbb{Q}_p) // \tilde{I})$ generated by $[\tilde{I}h^{-1}\tilde{I}]$ and we set

$$\mathcal{C}_0(\tilde{G}(\mathbb{A}_f) // \tilde{K})_h = \mathcal{C}_0(\tilde{G}(\mathbb{Q}_p) // \tilde{I})_h \otimes \mathcal{C}_0(\tilde{G}(\mathbb{A}_f)^{(p)} // \tilde{K}^{(p)}).$$

Since $[\tilde{I}h^{-1}\tilde{I}]^e = [\tilde{I}h^{-e}\tilde{I}]$ the algebra $\mathcal{C}_0(\tilde{G}(\mathbb{Q}_p) // \tilde{I})_h$ is the \mathbb{Z} -span of $[\tilde{I}h^{-e}\tilde{I}]$, $e \in \mathbb{N}_0$, hence, $\mathcal{C}_0(\tilde{G}(\mathbb{A}_f) // \tilde{K})_h$ is the \mathbb{Z} -span of

$$[\tilde{K}[h]_p^{-e}r\tilde{K}] = [\tilde{I}h^{-e}\tilde{I}] \otimes [\tilde{K}^{(p)}r\tilde{K}^{(p)}], \quad e \in \mathbb{N}_0, r \in \tilde{G}(\mathbb{A}_f)^{(p)};$$

here, $[h]_p \in \tilde{G}(\mathbb{A}_f)$ is the element with h in the p -component and all remaining components equal to 1.

For any \mathbb{Z} -algebra R we put $\mathcal{C}_{0, R}(\tilde{G}(\mathbb{A}_f) // \tilde{K})_? = \mathcal{C}_0(\tilde{G}(\mathbb{A}_f) // \tilde{K})_? \otimes R$ where $? = \text{blank}, h$. We define the $\tilde{\lambda}$ -normalization of $[\tilde{K}[h]_p^{-e}r\tilde{K}]$ as

$$[\tilde{K}[h]_p^{-e}r\tilde{K}]_{\tilde{\lambda}} = \tilde{\lambda}(h)^e [\tilde{K}[h]_p^{-e}r\tilde{K}] \in \mathcal{C}_{0, F}(\tilde{G}(\mathbb{A}_f) // \tilde{K})_h$$

(note that $\tilde{\lambda}(h)^e \in F$; see Section 2.5) and we extend linearly to $\mathcal{C}_{0,F}(\tilde{\mathbf{G}}(\mathbb{A}_f)//\tilde{\mathbf{K}})_h$. Since $[\tilde{\mathcal{I}}h^{-e}\tilde{\mathcal{I}}][\tilde{\mathcal{I}}h^{-f}\tilde{\mathcal{I}}] = [\tilde{\mathcal{I}}h^{-(e+f)}\tilde{\mathcal{I}}]$ we obtain that the assignment

$$\mathcal{C}_{0,F}(\tilde{\mathbf{G}}(\mathbb{A}_f)//\tilde{\mathbf{K}})_h \rightarrow \mathcal{C}_{0,F}(\tilde{\mathbf{G}}(\mathbb{A}_f)//\tilde{\mathbf{K}})_h, \quad \mathbb{T} \mapsto \mathbb{T}_{\tilde{\lambda}},$$

is a morphism of F -algebras.

4.11.2. Cohomology. The algebra $\mathcal{C}_0(\tilde{\mathbf{G}}(\mathbb{A}_f)//\tilde{\mathbf{K}})$, and hence also $\mathcal{C}_0(\tilde{\mathbf{G}}(\mathbb{A}_f)//\tilde{\mathbf{K}})_h$, acts on $H^n(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(F)) = \bigoplus_i H^n(\Gamma_i \backslash X, L_{\tilde{\lambda}}(F))$. We define the integral cohomology $H^n(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}}$ as the image of $H^n(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{Z}_p))$ in $H^n(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{Q}_p))$; hence, $H^n(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}} = \bigoplus_i H^n(\Gamma_i \backslash X, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}}$. Consider $e \in \mathbb{N}_0$ and $r \in \tilde{\mathbf{G}}(\mathbb{A}_f)^{(p)}$; we write $g_i[h]_p^{-e}r = \zeta_i g_{j(i)}k$ with $\zeta_i \in \tilde{\mathbf{G}}(\mathbb{Q})$, $k = (k_\ell)_\ell \in \tilde{\mathbf{K}}$ and obtain

$$[\tilde{\mathbf{K}}[h]_p^{-e}r\tilde{\mathbf{K}}] = \bigoplus_i \Gamma_i \zeta_i^{-1} \Gamma_{j(i)}.$$

Looking at the p -component and recalling that $g_i \in \tilde{\mathbf{G}}(\mathbb{A}_f)$ has trivial p -component we find $h^{-e} = \zeta_i k_p$. Hence, $\zeta_i^{-1} = k_p h^e$ is contained in $\tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$ and, thus, in $\Delta = \Delta_h$ and we deduce that the normalization $\tilde{\lambda}(h^e)\Gamma_i\zeta_i^{-1}\Gamma_{j(i)}$ maps $H^n(\Gamma_i \backslash X, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}}$ to $H^n(\Gamma_{j(i)} \backslash X, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}}$. Hence, $[\tilde{\mathbf{K}}[h]_p^{-e}r\tilde{\mathbf{K}}]_{\tilde{\lambda}}$ and, thus, any $\mathbb{T}_{\tilde{\lambda}}$ with \mathbb{T} in $\mathcal{C}_{0,\mathbb{Z}_p}(\tilde{\mathbf{G}}(\mathbb{A}_f)//\tilde{\mathbf{K}})_h$ leaves $H^n(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{Z}_p))_{\text{int}}$ invariant. In particular, the Lefschetz number $\text{Lef}(\mathbb{T}_{\tilde{\lambda}}|H^\bullet(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(F)))$, $\mathbb{T} \in \mathcal{C}_0(\tilde{\mathbf{G}}(\mathbb{A}_f)//\tilde{\mathbf{K}})_h$, is contained in F and in \mathbb{Z}_p (note that $F \subseteq \mathbb{Q}_p$). Moreover, as in the proof of 2.9 Corollary we see that the eigenvalues of $\mathbb{T}_{\tilde{\lambda}}$, $\mathbb{T} \in \mathcal{C}_0(\tilde{\mathbf{G}}(\mathbb{A}_f)//\tilde{\mathbf{K}})_h$, on $H^n_{\text{cusp}}(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{C})) \subseteq H^n(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(\mathbb{C}))$ are algebraic over F (note that $F \subseteq \mathbb{C}$) and integral over \mathbb{Z}_p , hence, they are contained in $\mathcal{O}_{\bar{\mathbb{Q}}_p}$ (note that $\bar{F} \subseteq \bar{\mathbb{Q}}_p$).

4.11.3. We denote by

$$\mathcal{C}_0(\tilde{\mathbf{G}}(\mathbb{A}_f)//\tilde{\mathbf{K}})_h^{\text{reg}} \subseteq \mathcal{C}_0(\tilde{\mathbf{G}}(\mathbb{A}_f)//\tilde{\mathbf{K}})_h$$

the \mathbb{Z} -submodule generated by all Hecke operators $[\tilde{\mathbf{K}}[h]_p^{-e}r\tilde{\mathbf{K}}]$ with $r \in \tilde{\mathbf{G}}(\mathbb{A}_f)^{(p)}$ and $e \in \mathbb{N}$ (i.e., $e \geq 1$). Keeping in mind that $[\tilde{\mathcal{I}}h^{-e}\tilde{\mathcal{I}}][\tilde{\mathcal{I}}h^{-f}\tilde{\mathcal{I}}] = [\tilde{\mathcal{I}}h^{-(e+f)}\tilde{\mathcal{I}}]$ we find that $\mathcal{C}_0(\tilde{\mathbf{G}}(\mathbb{A}_f)//\tilde{\mathbf{K}})_h^{\text{reg}}$ is an ideal in $\mathcal{C}_0(\tilde{\mathbf{G}}(\mathbb{A}_f)//\tilde{\mathbf{K}})_h$.

Proposition. *Let $C \in \mathbb{Q}_{>0}$. If the dominant weights $\tilde{\lambda}, \tilde{\lambda}' \in X(\tilde{\mathbf{T}})$ satisfy*

- $\langle \tilde{\lambda}, \alpha^\vee \rangle > 2C$ and $\langle \tilde{\lambda}', \alpha^\vee \rangle > 2C$ for all $\alpha \in \Delta_{\tilde{\mathbf{G}}}$,
- $\tilde{\lambda} \equiv \tilde{\lambda}' \pmod{(p-1)p^{m-1}X(\tilde{\mathbf{T}})}$ ($m \in \mathbb{N}$),

then for all $e \in \mathbb{N}$ and $r \in \tilde{\mathbf{G}}(\mathbb{A}_f)^{(p)}$ the Lefschetz number of $[\tilde{\mathbf{K}}[h]_p^{-e}r\tilde{\mathbf{K}}]_{\tilde{\lambda}}$ on $H^\bullet(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(F))$ is contained in \mathbb{Z}_p and the following congruence holds:

$$\text{Lef}([\tilde{\mathbf{K}}[h]_p^{-e}r\tilde{\mathbf{K}}]_{\tilde{\lambda}}|H^\bullet(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}}(F))) \equiv \text{Lef}([\tilde{\mathbf{K}}[h]_p^{-e}r\tilde{\mathbf{K}}]_{\tilde{\lambda}'}|H^\bullet(S_{\tilde{\mathbf{K}}}, L_{\tilde{\lambda}'}(F))) \pmod{p^{\lceil \min(m, Ce) \rceil} \mathbb{Z}_p}.$$

Proof. We have

$$\text{Lef}([\tilde{K}[h]_p^{-e} r \tilde{K}]_{\tilde{\lambda}} | H^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(F))) = \sum_{i=j(i)} \tilde{\lambda}(h^e) \text{Lef}(\Gamma_i \zeta_i^{-1} \Gamma_i | H^\bullet(\Gamma_i \backslash X, L_{\tilde{\lambda}}(F)))$$

where $\zeta_i^{-1} \in \Delta_h$ and $\zeta_i^{-1} \in \tilde{\mathcal{I}} h^e \tilde{\mathcal{I}}$. Hence, $\tilde{\lambda}(h^e) = \hat{\lambda}(\zeta_i^{-1})$ and the claim follows from 4.10 Proposition. \square

4.11.4. Slope subspaces. We select a Hecke operator $\mathbb{T} \in \mathcal{C}_{0, \mathbb{Z}_p}(\tilde{\mathbf{G}}(\mathbb{A}_f)/\tilde{K})_h^{\text{reg}}$ and we denote by $H^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Q}_p))^\beta$ the slope β subspace of $H^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Q}_p))$ w.r.t. $\mathbb{T}_{\tilde{\lambda}}$. We also denote by $\mathbf{g}_i = \mathbf{g}_{i, \tilde{K}}$ the number of i -cells in a cell complex which is homotopy equivalent to the Borel–Serre compactification $\bar{S}_{\tilde{K}}$ of $S_{\tilde{K}}$.

Theorem. For all $\beta \in \mathbb{Q}_{\geq 0}$, all dominant weights $\tilde{\lambda} \in X(\tilde{\mathbf{T}})$, all i and all $\mathbb{T} \in \mathcal{C}_{0, \mathbb{Z}_p}(\tilde{\mathbf{G}}(\mathbb{A}_f)/\tilde{K})_h^{\text{reg}}$ we have

$$\dim H^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{\leq \beta} \leq \mathfrak{m} \beta^s + \mathfrak{n};$$

here, $\mathfrak{m} = \mathfrak{m}_{\tilde{K}} = 12 \frac{\mathbf{g}_i}{s} \sigma^{s+1} \in \mathbb{Q}_{\geq 0}$ and $\mathfrak{n} = \mathfrak{n}_{\tilde{K}} \in \mathbb{N}$ is an integer which also only depends on \mathbf{g}_i, σ, s (see (50) below); in particular, \mathfrak{m} and \mathfrak{n} only depend on \tilde{K} (and, hence, on $\tilde{\mathbf{G}}$ and p) and i , i.e., they do not depend on $\tilde{\lambda}, h$ and \mathbb{T} .

The proof follows those of 3.9 Theorem and 3.10 Corollary. More precisely, for any $r \in \mathbb{N}_0$ we define the \mathbb{Z}_p -submodule

$$H^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma)) = \bigoplus_{j=1}^t H^i(\Gamma_j, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))$$

of $H^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Z}_p)) = \bigoplus_{j=1}^t H^i(\Gamma_j, L_{\tilde{\lambda}}(\mathbb{Z}_p))$ (note that $\Gamma_j \subseteq \tilde{\mathcal{I}}$). Using the decomposition $[\tilde{K}[h]_p^{-e} r \tilde{K}] = \bigoplus_i \Gamma_i \zeta_i^{-1} \Gamma_{j(i)}$ where $\zeta_i^{-1} \in \tilde{\mathcal{I}} h^e \tilde{\mathcal{I}}$ and following the proof in 3.5 Lemma we see that the submodule $H^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))$ satisfies the following properties.

- $H^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}}$ is $\mathbb{T}_{\tilde{\lambda}}$ -invariant.
- $\mathbb{T}_{\tilde{\lambda}} H^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))^{\text{TF}} \subseteq p^r H^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}}$.

We denote by $(p^{b_l})_l$ the elementary divisors of the quotient

$$(49) \quad \frac{H^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Z}_p))^{\text{TF}}}{H^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Z}_p, r, \sigma))^{\text{TF}}}$$

in decreasing order, i.e., $b_1 \geq b_2 \geq b_3 \geq \dots$. As in Section 3.9.1 we see that (49) is a subquotient of $\bigoplus_j H^i(\Gamma_j, L_{\tilde{\lambda}}^{[r]}(\mathbb{Z}_p, \sigma))$, which is a subquotient of

$$\bigoplus_j \bigoplus_{h=0}^r \left(\frac{\mathbb{Z}_p}{p^{r-h} \mathbb{Z}_p} \right)^{\mathbf{g}_{i, \Gamma_j} M_{\sigma, h}} = \bigoplus_{h=0}^r \left(\frac{\mathbb{Z}_p}{p^{r-h} \mathbb{Z}_p} \right)^{\mathbf{g}_i M_{\sigma, h}}.$$

We denote by $(p^{a_i})_l$ the elementary divisors of the latter sum in decreasing order, i.e., $a_1 \geq a_2 \geq \dots$ (the elementary divisor p^{r-h} appears $\mathbf{g}_i M_{\sigma, h}$ -times). We obtain $b_i \leq a_i \leq r$ for all i and following the arguments in Section 3.9.2 - 3.9.4 with \mathbf{g}_i in place of g_i we obtain that the piecewise linear function defined in 3.9 Theorem but with g_i replaced by \mathbf{g}_i is a lower bound for the Newton polygon of $\mathbb{T}_{\tilde{\lambda}}$ acting on $H^i(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{<\infty}$. Following the proof of 3.10 Corollary we find the bound for the dimension of the slope β subspace as in the statement of the theorem where we can choose

$$(50) \quad \mathbf{n} = \left\lceil 2^{\frac{1}{s+1}} \frac{\mathbf{g}_i \sigma^{s+1}}{s} M^s \right\rceil + 1 \in \mathbb{N};$$

here, M is defined in equations (23) and (24) (note that the definition of M does not depend on g_i). This finishes the proof of the theorem.

4.12. Induced representations. We look at traces of Hecke operators on induced representations. This will be applied in the next section where we consider Hecke operators on Eisenstein cohomology.

4.12.1. We select a \mathbb{Q} -parabolic subgroup $\tilde{Q} \leq \tilde{G}$ with Levi decomposition $\tilde{Q} = \tilde{M}\tilde{N}$ and a representation π of $\tilde{M}(\mathbb{A}_f)$. By $\text{Ind}_{\tilde{Q}(\mathbb{A}_f)}^{\tilde{G}(\mathbb{A}_f)} \pi$ we understand the non-unitarily induced representation. We select a maximal compact open subgroup $\tilde{K}_0 = \prod_{\ell \neq \infty} \tilde{K}_{0, \ell} \leq \tilde{G}(\mathbb{A}_f)$ and we may assume that $\tilde{K} \leq \tilde{K}_0$. We set

$$\tilde{K}^{\tilde{M}} = \prod_{\ell \neq \infty} \tilde{K}_{\ell}^{\tilde{M}} \quad \text{and} \quad \tilde{K}^{(p), \tilde{M}} = \prod_{\ell \neq p, \infty} \tilde{K}_{\ell}^{\tilde{M}},$$

where $\tilde{K}_{\ell}^{\tilde{M}} = \tilde{K}_{\ell} \cap \tilde{M}(\mathbb{Q}_{\ell})$, and we use the same definition with \tilde{K} replaced by \tilde{K}_0 . We also set $\tilde{K}^{\tilde{N}} = \tilde{K} \cap \tilde{N}(\mathbb{A}_f)$ and $\tilde{K}^{(p), \tilde{N}} = \tilde{K}^{(p)} \cap \tilde{N}(\mathbb{A}_f)^{(p)}$.

Let $f \in \mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{A}_f))$; we define the constant term $f_{\tilde{M}} \in \mathcal{C}_{0, \mathbb{Q}}(\tilde{M}(\mathbb{A}_f))$ by

$$f_{\tilde{M}}(x) = \int_{\tilde{K}_0} \int_{\tilde{N}(\mathbb{A}_f)} f(k^{-1}xnk) \, dn \, dk.$$

Here and below we normalize Haar measures on $\tilde{G}(\mathbb{A}_f)$ (and, hence, on \tilde{K}_0), $\tilde{M}(\mathbb{A}_f)$ and $\tilde{N}(\mathbb{A}_f)$ so that $\text{vol}(\tilde{K}) = 1$, $\text{vol}(\tilde{K}^{\tilde{M}}) = 1$, $\text{vol}(\tilde{K}^{\tilde{N}}) = 1$. With these conventions we have

$$\text{tr}(f | \text{Ind}_{\tilde{Q}(\mathbb{A}_f)}^{\tilde{G}(\mathbb{A}_f)} \pi) = \text{tr}(f_{\tilde{M}} | \pi).$$

If $f^{(p)} \in \mathcal{C}_{0, \mathbb{Q}}(\tilde{G}(\mathbb{A}_f)^{(p)})$ we define $f_{\tilde{M}}^{(p)} \in \mathcal{C}_{0, \mathbb{Q}}(\tilde{M}(\mathbb{A}_f)^{(p)})$ by replacing f with $f^{(p)}$, \tilde{K}_0 with $\tilde{K}_0^{(p)} = \prod_{\ell \neq p, \infty} \tilde{K}_{0, \ell}$ and $\tilde{N}(\mathbb{A}_f)$ with $\tilde{N}(\mathbb{A}_f)^{(p)}$; an analogous identity for the trace of $f^{(p)}$ on representations induced from $\tilde{M}(\mathbb{A}_f)^{(p)}$ then holds. We will also need the following classical identity which holds for a representation π_p of $\tilde{M}(\mathbb{Q}_p)$. Assume that \tilde{Q}/F contains \tilde{B}^-/F , i.e., the unipotent radical of \tilde{Q}/F

is generated by root subgroups attached to negative roots (compare the definition of \mathcal{I} in Section 1.3); then

$$(51) \quad \text{tr}([\tilde{\mathcal{I}}h^{-e}\tilde{\mathcal{I}}](\text{Ind}_{\tilde{\mathcal{Q}}(\mathbb{Q}_p)}^{\tilde{\mathcal{G}}(\mathbb{Q}_p)}\pi_p)^{\tilde{\mathcal{I}}}) = \sum_{v \in W^{\tilde{\mathcal{Q}}}} c_{v,h^{-e}} \text{tr}([\tilde{\mathcal{I}}^{\tilde{M}}vh^{-e}v^{-1}\tilde{\mathcal{I}}^{\tilde{M}}]|\pi_p^{\tilde{\mathcal{I}}^{\tilde{M}}}),$$

where $c_{v,h^{-e}} \in \mathbb{N}$ and $\tilde{\mathcal{I}}^{\tilde{M}} = \tilde{\mathcal{I}} \cap \tilde{\mathcal{M}}(\mathbb{Q}_p)$ (see [Urban 2011, p. 1751], for example). We obtain an analogous formula when $\tilde{\mathcal{Q}}$ is standard parabolic, i.e., $\tilde{\mathcal{B}}/F \leq \tilde{\mathcal{Q}}/F$. To this end we denote by $a \in W_{\tilde{\mathcal{G}}}$ and $b \in W_{\tilde{\mathcal{M}}} = W(\tilde{\mathcal{T}}/\mathbb{Z}_p, \tilde{\mathcal{M}}/\mathbb{Z}_p)$ the longest elements; i.e., a maps $\Phi_{\tilde{\mathcal{G}}}^+$ to $\Phi_{\tilde{\mathcal{G}}}^-$ and b maps $\Phi_{\tilde{\mathcal{M}}}^+$ to $\Phi_{\tilde{\mathcal{M}}}^-$. Using (51) but with positivity defined by $-\Delta_{\tilde{\mathcal{G}}}$ — that is, $\tilde{\mathcal{B}}/F \leq \tilde{\mathcal{Q}}/F$, $\tilde{\mathcal{I}} \equiv \tilde{\mathcal{B}}(\mathbb{F}_p) \pmod{p}$ and $h \in \tilde{\mathcal{T}}(\mathbb{Q})^{-}$ — we obtain

$$\begin{aligned} \text{tr}([\tilde{\mathcal{I}}h^{-e}\tilde{\mathcal{I}}](\text{Ind}_{\tilde{\mathcal{Q}}(\mathbb{Q}_p)}^{\tilde{\mathcal{G}}(\mathbb{Q}_p)}\pi_p)^{\tilde{\mathcal{I}}}) &= \text{tr}([{}^a\tilde{\mathcal{I}}({}^ah^{-e}{}^a\tilde{\mathcal{I}})](\text{Ind}_{\tilde{\mathcal{Q}}(\mathbb{Q}_p)}^{\tilde{\mathcal{G}}(\mathbb{Q}_p)}\pi_p)^{{}^a\tilde{\mathcal{I}}}) \\ &= \sum_{v \in W^{\tilde{\mathcal{Q}}}} c_{v,h^{-e}} \text{tr}([({}^a\tilde{\mathcal{I}})^{\tilde{M}}v({}^ah^{-e})v^{-1}({}^a\tilde{\mathcal{I}})^{\tilde{M}}]|\pi_p^{({}^a\tilde{\mathcal{I}})^{\tilde{M}}}) \\ &= \sum_{v \in W^{\tilde{\mathcal{Q}}}} c_{v,h^{-e}} \text{tr}([{}^b({}^a\tilde{\mathcal{I}})^{\tilde{M}}b v({}^ah^{-e})v^{-1}b^{-1}({}^a\tilde{\mathcal{I}})^{\tilde{M}}]|\pi_p^{b({}^a\tilde{\mathcal{I}})^{\tilde{M}}}) \\ &= \sum_{v \in W^{\tilde{\mathcal{Q}}}} c_{v,h^{-e}} \text{tr}([\tilde{\mathcal{I}}^{\tilde{M}}({}^{bva}h^{-e})\tilde{\mathcal{I}}^{\tilde{M}}]|\pi_p^{\tilde{\mathcal{I}}^{\tilde{M}}}), \end{aligned}$$

where $c_v = c_{v,h^{-e}} \in \mathbb{Z}$ are certain integers.

4.12.2. We look more closely at the constant term $f_{\tilde{\mathcal{M}}}^{(p)}$ of $f^{(p)} \in \mathcal{C}_0(\tilde{\mathcal{G}}(\mathbb{A}_f)^{(p)}/\tilde{\mathcal{K}}^{(p)})$, i.e., $f^{(p)}$ is \mathbb{Z} -valued and $\tilde{\mathcal{K}}^{(p)}$ bi-invariant (note that $\text{vol}(\tilde{\mathcal{K}}^{(p)}) = 1$ by our normalizations). For simplicity we assume that $\tilde{\mathcal{K}}^{(p)} \leq \tilde{\mathcal{K}}_0^{(p)}$ is a normal subgroup. The definition yields

$$\begin{aligned} f_{\tilde{\mathcal{M}}}^{(p)}(x) &= \sum_{k \in \tilde{\mathcal{K}}_0^{(p)}/\tilde{\mathcal{K}}^{(p)}} \int_{\tilde{\mathcal{N}}(\mathbb{A}_f)^{(p)}} f^{(p)}(k^{-1}xnk) \, dn \\ &= \text{vol}(\tilde{\mathcal{K}}^{(p),\tilde{\mathcal{N}}}) \sum_{k \in \tilde{\mathcal{K}}_0^{(p)}/\tilde{\mathcal{K}}^{(p)}} \int_{\tilde{\mathcal{N}}(\mathbb{A}_f)^{(p)}/\tilde{\mathcal{K}}^{(p),\tilde{\mathcal{N}}}} f^{(p)}(k^{-1}xnk) \, dn, \end{aligned}$$

The first of these equalities shows that $f_{\tilde{\mathcal{M}}}^{(p)}$ is $\tilde{\mathcal{K}}^{(p),\tilde{\mathcal{M}}}$ bi-invariant (the modulus $\delta_{\tilde{\mathcal{Q}}(\mathbb{A}_f)^{(p)}}(m)$ vanishes for $m \in \tilde{\mathcal{K}}^{(p),\tilde{\mathcal{M}}}$ because $\tilde{\mathcal{K}}^{(p),\tilde{\mathcal{M}}}$ is contained in the compact group $\tilde{\mathcal{K}}^{(p)} \cap \tilde{\mathcal{M}}(\mathbb{A}_f)^{(p)}$). Since $\text{vol}(\tilde{\mathcal{K}}^{(p),\tilde{\mathcal{N}}}) = \text{vol}(\tilde{\mathcal{K}}^{(p),\tilde{\mathcal{M}}}) = 1$ the second equality implies

$$(52) \quad f_{\tilde{\mathcal{M}}}^{(p)} \in \mathcal{C}_0(\tilde{\mathcal{M}}(\mathbb{A}_f)^{(p)}/\tilde{\mathcal{K}}^{(p),\tilde{\mathcal{M}}}).$$

4.13. Eisenstein cohomology.

4.13.1. We assume that the highest weight $\tilde{\lambda} \in X(\tilde{\mathcal{T}})$ is dominant and regular. The

full cohomology then decomposes as

$$H^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C})) = H_{\text{cusp}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C})) \oplus H_{\text{Eis}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C})),$$

where

$$(53) \quad H_{\text{Eis}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C})) = \bigoplus_{\tilde{M} \neq \tilde{G}} \bigoplus_w (\text{Ind}_{\tilde{Q}(\mathbb{A}_f)}^{\tilde{G}(\mathbb{A}_f)} H_{\text{cusp}}^{\bullet-\ell(w)}(S^{\tilde{M}}, L_{w \cdot \tilde{\lambda}}(\mathbb{C})))^{\tilde{K}};$$

here, \tilde{M} runs over a system of representatives of $\tilde{G}(\mathbb{Q})$ -conjugacy classes of proper \mathbb{Q} -Levi subgroups of \tilde{G} , i.e., there is a (standard) \mathbb{Q} -parabolic subgroup $\tilde{Q} \leq \tilde{G}$ with Levi decomposition

$$\tilde{Q} = \tilde{M}\tilde{N}, \quad S^{\tilde{M}} = \tilde{M}(\mathbb{Q}) \backslash \tilde{M}(\mathbb{A}) / A_{\tilde{Q}} \tilde{K}_{\infty}^{\tilde{M}},$$

where $\tilde{K}_{\infty}^{\tilde{M}} = \tilde{K}_{\infty} \cap \tilde{M}(\mathbb{R})$, is the locally symmetric space attached to \tilde{M} , $A_{\tilde{Q}}$ is the connected component of the real points of a maximal \mathbb{Q} -split torus $A_{\tilde{Q}}$ in the center of \tilde{M} and w runs over those elements in $W^{\tilde{Q}}$ which satisfy the condition that $-w(\tilde{\lambda} + \rho^\circ)|_{A_{\tilde{Q}}}$ is nonnegative, i.e., $\langle -\text{Re}(w(\tilde{\lambda} + \rho^\circ)), \alpha^\vee \rangle \geq 0$ for all roots α of $A_{\tilde{Q}}$ acting on $\text{Lie}(\tilde{N}) \otimes \mathbb{R}$ (see [Franke 1998, Theorem 19 II, p. 257] and [Schwermer 1994, Proof of 6.3 Theorem, p. 505]).

4.13.2. Theorem. *Let $C \in \mathbb{Q}_{>0}$ and suppose the dominant weights $\tilde{\lambda}, \tilde{\lambda}' \in X(\tilde{T})$ satisfy*

- $\langle \tilde{\lambda}, \alpha^\vee \rangle > 2C$ and $\langle \tilde{\lambda}', \alpha^\vee \rangle > 2C$ for all $\alpha \in \Delta_{\tilde{G}}$,
- $\tilde{\lambda} \equiv \tilde{\lambda}' \pmod{(p-1)p^{m-1}X(\tilde{T})}$ ($m \in \mathbb{N}$).

Then, for all $e \in \mathbb{N}$ and $r \in \tilde{G}(\mathbb{A}_f)^{(p)}$, the Lefschetz number of $[\tilde{K}[h]_p^{-e} r \tilde{K}]_{\tilde{\lambda}}$ on $H_{\text{cusp}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))$ is contained in F and the following congruence holds (note that $F \subseteq \mathbb{Q}_p$):

$$\text{Lef}([\tilde{K}[h]_p^{-e} r \tilde{K}]_{\tilde{\lambda}} | H_{\text{cusp}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))) \equiv \text{Lef}([\tilde{K}[h]_p^{-e} r \tilde{K}]_{\tilde{\lambda}'} | H_{\text{cusp}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}'}(\mathbb{C}))) \pmod{p^\dagger \mathbb{Z}_p}.$$

Here, $\dagger = \lceil \min(m, e(C - \kappa_2 \text{rk}(\tilde{G}))) \rceil - e\kappa_1 \text{rk}(\tilde{G})$ with $\kappa_i = \kappa_{i, \tilde{G}}$ and $\text{rk}(\tilde{G}) = \text{rk}_{\mathbb{Q}}(\tilde{G})$ is the \mathbb{Q} -rank of \tilde{G} .

Proof. We use induction on the \mathbb{Q} -rank of \tilde{G} . If $\text{rk}(\tilde{G}) = 0$ then

$$H_{\text{cusp}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C})) = H^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}));$$

hence, 4.11.3 Proposition implies the claim. We assume $\text{rk}(\tilde{G}) > 0$. Equation (53)

yields

$$\begin{aligned}
 (54) \quad & \text{Lef}([\tilde{K}[h]_p^{-e} r \tilde{K}] | H_{\text{Eis}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))) \\
 &= \sum_{\tilde{M} \neq \tilde{G}} \sum_w \sum_i (-1)^{i+\ell(w)} \text{tr}([\tilde{K}[h]_p^{-e} r \tilde{K}] | (\text{Ind}_{\tilde{Q}(\mathbb{A}_f)}^{\tilde{G}(\mathbb{A}_f)} H_{\text{cusp}}^i(S^{\tilde{M}}, L_{w \cdot \tilde{\lambda}}(\mathbb{C})))^{\tilde{K}}) \\
 &= \sum_{\tilde{M} \neq \tilde{G}} \sum_w (-1)^{\ell(w)} \text{Lef}([\tilde{K}[h]_p^{-e} r \tilde{K}] | \bigoplus_i (\text{Ind}_{\tilde{Q}(\mathbb{A}_f)}^{\tilde{G}(\mathbb{A}_f)} \pi_{w \cdot \tilde{\lambda}}^i)^{\tilde{K}}).
 \end{aligned}$$

Here, we have set

$$\pi_{w \cdot \tilde{\lambda}}^i = H_{\text{cusp}}^i(S^{\tilde{M}}, L_{w \cdot \tilde{\lambda}}(\mathbb{C}));$$

this is a module under the Hecke algebra attached to $\tilde{M}(\mathbb{A}_f)$ and we have

$$H_{\text{cusp}}^i(S^{\tilde{M}}, L_{w \cdot \tilde{\lambda}}(\mathbb{C}))^{\tilde{H}} = H_{\text{cusp}}^i(S^{\tilde{M}}/\tilde{H}, L_{w \cdot \tilde{\lambda}}(\mathbb{C})),$$

with $\tilde{H} \leq \tilde{M}(\mathbb{A}_f)$ compact open. We select a proper \mathbb{Q} -parabolic subgroup $\tilde{Q} = \tilde{M}\tilde{N}$ of \tilde{G} and an element $w \in W^{\tilde{Q}}$ as in equation (53). We denote by $\Phi_{\tilde{M}}$ the set of roots of \tilde{T}/F acting on $\text{Lie}(\tilde{M}/F)$ and we set $\Delta_{\tilde{M}} = \Phi_{\tilde{M}} \cap \Delta_{\tilde{G}}$. The set $\Delta_{\tilde{M}}$ is the basis for the root system $\Phi_{\tilde{M}}$ corresponding to the Borel subgroup $\tilde{B}^{\tilde{M}}/F = \tilde{B}/F \cap \tilde{M}/F$ of \tilde{M}/F ; in particular, this determines the set of positive roots $\Phi_{\tilde{M}}^+$. The subgroup $\tilde{\mathcal{I}}^{\tilde{M}} = \tilde{\mathcal{I}} \cap \tilde{M}(\mathbb{Q}_p)$ is a Iwahori subgroup in $\tilde{M}(\mathbb{Z}_p)$, i.e., $\tilde{\mathcal{I}}^{\tilde{M}} \pmod{p}$ is contained in the Borel subgroup $(\tilde{B}^- \cap \tilde{M})(\mathbb{F}_p) \leq \tilde{M}(\mathbb{F}_p)$. The identities in Section 4.12.1 for the traces of induced representations and equation (52) yield

$$\begin{aligned}
 (55) \quad & \text{Lef}([\tilde{K}[h]_p^{-e} r \tilde{K}] | \bigoplus_i (\text{Ind}_{\tilde{Q}(\mathbb{A}_f)}^{\tilde{G}(\mathbb{A}_f)} \pi_{w \cdot \tilde{\lambda}}^i)^{\tilde{K}}) \\
 &= \text{Lef}([\tilde{\mathcal{I}}h^{-e} \tilde{\mathcal{I}}] \otimes [\tilde{K}^{(p)} r \tilde{K}^{(p)}] | \bigoplus_i \text{Ind}_{\tilde{Q}(\mathbb{A}_f)}^{\tilde{G}(\mathbb{A}_f)} \pi_{w \cdot \tilde{\lambda}}^i) \\
 &= \sum_{v \in W^{\tilde{Q}}} c_v \text{Lef}([\tilde{\mathcal{I}}^{\tilde{M}}(bva h^{-e}) \tilde{\mathcal{I}}^{\tilde{M}}] \otimes [\tilde{K}^{(p)} r \tilde{K}^{(p)}]_{\tilde{M}} | \bigoplus_i \pi_{w \cdot \tilde{\lambda}}^i) \\
 &= \sum_{v \in W^{\tilde{Q}}} c_v \text{Lef}([\tilde{\mathcal{I}}^{\tilde{M}}(bva h^{-e}) \tilde{\mathcal{I}}^{\tilde{M}}] \otimes [\tilde{K}^{(p)} r \tilde{K}^{(p)}]_{\tilde{M}} | \bigoplus_i (\pi_{w \cdot \tilde{\lambda}}^i)^{\tilde{K}_{\tilde{M}}}),
 \end{aligned}$$

where $c_v = c_{v, h^{-e}} \in \mathbb{Z}$, a (resp. b) is the longest element in the Weyl group $W_{\tilde{G}}$ (resp. $W_{\tilde{M}}$) and $\tilde{K}^{\tilde{M}} = \tilde{\mathcal{I}}^{\tilde{M}} \times \tilde{K}^{(p), \tilde{M}}$. We select an element $v \in W^{\tilde{Q}}$. Since $v^{-1} \Phi_{\tilde{M}}^+ \subseteq \Phi_{\tilde{G}}^+$ and ${}^a h \in \tilde{T}(\mathbb{Q}_p)^{-}$ we obtain for all $\alpha \in \Phi_{\tilde{M}}^+$ that $v_p(\alpha(bva h)) = v_p((v^{-1} b^{-1} \alpha)({}^a h)) > 0$; hence, $bva h$ is regular dominant w.r.t. $\Phi_{\tilde{M}}^+$. Thus, using equation (52) we obtain that

$$(56) \quad [\tilde{\mathcal{I}}^{\tilde{M}}(bva h^{-e}) \tilde{\mathcal{I}}^{\tilde{M}}] \otimes [\tilde{K}^{(p)} r \tilde{K}^{(p)}]_{\tilde{M}} \in \mathcal{C}_0(\tilde{M}(\mathbb{A}_f) // \tilde{K}^{\tilde{M}})_{bva h}$$

is contained in the integral Hecke algebra attached to \tilde{M} and the dominant regular element $bva h \in \tilde{T}(F)$. Now, in Section 4.6 we have seen that $\langle w\rho - \rho, \alpha^\vee \rangle \geq -2\kappa_2$

for all $\alpha \in \Delta_{\tilde{G}}$ ($\rho = \rho_{\tilde{G}}$); hence, we obtain for all $\alpha \in \Delta_{\tilde{M}}$

$$\langle w \cdot \tilde{\lambda}, \alpha^\vee \rangle = \langle \tilde{\lambda}, w^{-1} \alpha^\vee \rangle + \langle w \rho^\circ - \rho^\circ, \alpha^\vee \rangle \geq 2C - 2\kappa_2.$$

Since also $w \cdot \tilde{\lambda} \equiv w \cdot \tilde{\lambda}' \pmod{(p-1)p^m X(\tilde{T})}$ the induction hypotheses for the group \tilde{M} which has strictly smaller rank than \tilde{G} (since $\tilde{M} \neq \tilde{G}$) is satisfied and we obtain that the following congruence holds:

$$(57) \quad (w \cdot \tilde{\lambda})(^{bva}h^e) \text{Lef}([\tilde{\mathcal{I}}^{\tilde{M}}(^{bva}h^{-e})\tilde{\mathcal{I}}^{\tilde{M}}] \otimes [\tilde{K}^{(p)}r\tilde{K}^{(p)}]_{\tilde{M}} | \bigoplus_i (\pi_{w \cdot \tilde{\lambda}}^i)^{\tilde{K}^{\tilde{M}}}) \\ \equiv (w \cdot \tilde{\lambda}')(^{bva}h^e) \text{Lef}([\tilde{\mathcal{I}}^{\tilde{M}}(^{bva}h^{-e})\tilde{\mathcal{I}}^{\tilde{M}}] \otimes [\tilde{K}^{(p)}r\tilde{K}^{(p)}]_{\tilde{M}} | \bigoplus_i (\pi_{w \cdot \tilde{\lambda}'}^i)^{\tilde{K}^{\tilde{M}}}) \\ \pmod{p^\star \mathbb{Z}_p},$$

where $\star = \lceil \min(m, e(C - \kappa_2 - \kappa_{2, \tilde{M}} \text{rk}(\tilde{M}))) \rceil - e\kappa_{1, \tilde{M}} \text{rk}(\tilde{M})$; moreover, the above Lefschetz numbers are contained in F . Equations (54) and (55) thus imply that

$$\tilde{\lambda}(h^e) \text{Lef}([\tilde{K}[h]_p^{-e}r\tilde{K}] | H_{\text{Eis}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))) \in F$$

(note that $\tilde{\lambda}$ and all $\alpha \in \Phi_{\tilde{G}}$ are defined over F , $h \in \tilde{T}(\mathbb{Q})$ and a, b, v, w normalize $\tilde{T}(F)$) and since full cohomology is defined over F we obtain that

$$\text{Lef}([\tilde{K}[h]_p^{-e}r\tilde{K}]_{\tilde{\lambda}} | H_{\text{cusp}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))) \in F.$$

It remains to prove the congruences. Since $\tilde{\lambda} - (bva)^{-1}(w \cdot \tilde{\lambda}) = \tilde{\lambda} - ((bva)^{-1}w) \cdot \tilde{\lambda} + (bva)^{-1}\rho^\circ - \rho^\circ$ we obtain using 4.5 Lemma (note that $h \in \tilde{T}(\mathbb{Q}_p)^{++}$) and $v_p(((bva)^{-1}\rho^\circ - \rho^\circ)(h^e)) \geq -e\kappa_1$ (see Section 4.6) that

$$(58) \quad \frac{\tilde{\lambda}(h^e)}{(bva)^{-1}(w \cdot \tilde{\lambda})(h^e)} \begin{cases} \text{has } p\text{-adic value} \geq Ce - e\kappa_1 & \text{if } bva \neq w, \\ \text{equals } ((bva)^{-1}\rho^\circ - \rho^\circ)(h^e) & \text{if } bva = w, \end{cases}$$

and the same holds for $\tilde{\lambda}'$. Thus, if $bva = w$, we obtain from equation (57), after multiplying by $((bva)^{-1}\rho^\circ - \rho^\circ)(h^e)$,

$$\tilde{\lambda}(h^e) \text{Lef}([\tilde{\mathcal{I}}^{\tilde{M}}(^{bva}h^{-e})\tilde{\mathcal{I}}^{\tilde{M}}] \otimes [\tilde{K}^{(p)}r\tilde{K}^{(p)}]_{\tilde{M}} | \bigoplus_i (\pi_{w \cdot \tilde{\lambda}}^i)^{\tilde{K}^{\tilde{M}}}) \\ \equiv \tilde{\lambda}'(h^e) \text{Lef}([\tilde{\mathcal{I}}^{\tilde{M}}(^{bva}h^{-e})\tilde{\mathcal{I}}^{\tilde{M}}] \otimes [\tilde{K}^{(p)}r\tilde{K}^{(p)}]_{\tilde{M}} | \bigoplus_i (\pi_{w \cdot \tilde{\lambda}'}^i)^{\tilde{K}^{\tilde{M}}}) \pmod{p^{\star - e\kappa_1} \mathbb{Z}_p}.$$

We look at the case $bva \neq w$. Since

$$[\tilde{K}^{(p)}r\tilde{K}^{(p)}]_{\tilde{M}} = \sum_{s \in \tilde{M}(\mathbb{A}_f)^{(p)}} z_s [\tilde{K}^{(p), \tilde{M}}_s \tilde{K}^{(p), \tilde{M}}]$$

with $z_s \in \mathbb{Z}$, by equation (52), and since the trace of the $w \cdot \tilde{\lambda}$ -normalization of

$$[\tilde{\mathcal{I}}^{\tilde{M}}(^{bva}h^{-e})\tilde{\mathcal{I}}^{\tilde{M}}] \otimes [\tilde{K}^{(p), \tilde{M}}_s \tilde{K}^{(p), \tilde{M}}] \in C_0(\tilde{M}(\mathbb{A}_f) // \tilde{K}^{\tilde{M}})_{bva_h}$$

on $H_{\text{cusp}}^i(S^{\tilde{M}}, L_{w \cdot \tilde{\lambda}}(\mathbb{C}))$ is contained in $\mathcal{O}_{\mathbb{Q}_p}$ (see Section 4.11.2) we obtain that the Lefschetz number of $(w \cdot \tilde{\lambda})(^{bva}h^e) [\tilde{\mathcal{I}}^{\tilde{M}}(^{bva}h^{-e})\tilde{\mathcal{I}}^{\tilde{M}}] \otimes [\tilde{K}^{(p)}r\tilde{K}^{(p)}]_{\tilde{M}}$ on cuspidal

cohomology $\bigoplus_i (\pi_{w,\tilde{\lambda}}^i)^{\tilde{K}_{\tilde{M}}}$ is p -adically integral; thus using equation (58) we obtain

$$\begin{aligned} & \tilde{\lambda}(h^e) \operatorname{Lef}([\tilde{\mathcal{I}}^{\tilde{M}}(bva h^{-e})\tilde{\mathcal{I}}^{\tilde{M}}] \otimes [\tilde{K}^{(p)}r\tilde{K}^{(p)}]_{\tilde{M}} | \bigoplus_i (\pi_{w,\tilde{\lambda}}^i)^{\tilde{K}_{\tilde{M}}}) \\ & \equiv \tilde{\lambda}'(h^e) \operatorname{Lef}([\tilde{\mathcal{I}}^{\tilde{M}}(bva h^{-e})\tilde{\mathcal{I}}^{\tilde{M}}] \otimes [\tilde{K}^{(p)}r\tilde{K}^{(p)}]_{\tilde{M}} | \bigoplus_i (\pi_{w,\tilde{\lambda}'}^i)^{\tilde{K}_{\tilde{M}}}) \pmod{p^{Ce-e\kappa_1}\mathbb{Z}_p}. \end{aligned}$$

Since $Ce - e\kappa_1 \geq \star - e\kappa_1$, in both cases the congruence holds modulo $p^{\star - e\kappa_1}\mathbb{Z}_p$. Using equations (54) and (55) we obtain

$$\begin{aligned} & \operatorname{Lef}([\tilde{K}[h]_p^{-e}r\tilde{K}]_{\tilde{\lambda}} | H_{\text{Eis}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))) \\ & \equiv \operatorname{Lef}([\tilde{K}[h]_p^{-e}r\tilde{K}]_{\tilde{\lambda}'} | H_{\text{Eis}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}'}(\mathbb{C}))) \pmod{p^{\star - e\kappa_1}\mathbb{Z}_p}. \end{aligned}$$

The rank of \tilde{M} is strictly smaller than the rank of \tilde{G} and $\kappa_{i,\tilde{M}} \leq \kappa_i$, as follows from Section 4.6; hence, $\kappa_i + \kappa_{i,\tilde{M}}\operatorname{rk}(\tilde{M}) \leq \kappa_i\operatorname{rk}(\tilde{G})$ which yields

$$\star - e\kappa_1 \geq \lceil \min(m, e(C - \kappa_2\operatorname{rk}(\tilde{G}))) \rceil - \operatorname{rk}(\tilde{G})e\kappa_1.$$

Together with 4.11.3 Proposition this implies the claim about congruences for the Lefschetz numbers of $[\tilde{K}[h]_p^{-e}r\tilde{K}]_{\tilde{\lambda}}$ on cuspidal cohomology for $S_{\tilde{K}}$. Thus, the theorem is proven. \square

4.13.3. Remark. Section 4.11.2 implies that the Lefschetz number

$$\operatorname{Lef}(\mathbb{T}_{\tilde{\lambda}} | H_{\text{cusp}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))), \quad \mathbb{T} \in \mathcal{C}_0(\tilde{G}(\mathbb{A}_f) // \tilde{K})_h^{\text{reg}},$$

is contained in $\mathcal{O}_{\tilde{\mathbb{Q}}_p}$. Thus, 4.13.2 Theorem implies that $\operatorname{Lef}(\mathbb{T}_{\tilde{\lambda}} | H_{\text{cusp}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C})))$ is contained in $F \cap \mathcal{O}_{\tilde{\mathbb{Q}}_p} \subseteq \mathbb{Z}_p$.

4.14. Weighted cohomology. In this section we compare two Goresky–MacPherson trace formulas for two different weights $\tilde{\lambda}$ and $\tilde{\lambda}'$. This is analogous to the comparison of Bewersdorff’s trace formula in previous sections and relies on the same diagonalization of elements in $\tilde{\mathcal{I}}h^e\tilde{\mathcal{I}}$ (see 4.4 Proposition) but now applied to all Levi subgroups \tilde{M} of parabolic subgroups in \tilde{G}/\mathbb{Q} (see 4.7 Proposition). As a result we obtain congruences for Hecke operators on weighted cohomology for varying weight $\tilde{\lambda}$.

Here, we will work again in a classical, non-adelic setting (see Section 2.2 and 2.3); e.g., $\Gamma \leq \tilde{G}(\mathbb{Q})$ is an arithmetic subgroup contained in $\tilde{\mathcal{I}}$, $h \in \tilde{T}(\mathbb{Q})^{++}$ and we will consider Hecke operators $\Gamma\zeta\Gamma$ where $\zeta \in \Delta = \Delta_h$.

4.14.1. The trace formula of Goresky–MacPherson. We select a minimal \mathbb{Q} -parabolic subgroup \tilde{P}_0 in \tilde{G}/\mathbb{Q} with Levi decomposition $\tilde{P}_0 = \tilde{M}_0\tilde{N}_0$ and we denote by A_0 a maximal \mathbb{Q} -split torus in the center of the Levi subgroup \tilde{M}_0 . We may assume that $\tilde{B} \subseteq \tilde{P}_0/F$ and $\tilde{T} \supseteq A_0/F$. Let $\tilde{P} = \tilde{M}\tilde{N}$ be a \mathbb{Q} -parabolic subgroup in \tilde{G} . We denote by $A_{\tilde{P}}$ a maximal \mathbb{Q} -split torus in the center of \tilde{M} and we write

$\Delta_{\tilde{P}} = \{\alpha_1, \dots, \alpha_m\} \subset X(A_{\tilde{P}}/A_{\tilde{G}})$ for the set of simple roots of $A_{\tilde{P}}$ occurring in $\text{Lie}(\tilde{N})$ and

$$\{t_{\alpha_1}, \dots, t_{\alpha_m}\} \subset X_*(A_{\tilde{P}}/A_{\tilde{G}}) \otimes \mathbb{Q}$$

for the basis dual to $\Delta_{\tilde{P}}$. We select $h \in \tilde{T}(\mathbb{Q})^{++}$ and we let $\zeta \in \Delta = \Delta_h$. The double coset $\Gamma\zeta\Gamma$ induces an operator on the weighted cohomology groups $W^\nu H^i(\overline{\Gamma\backslash X}, L_{\tilde{\lambda}}(\mathbb{C}))$ with weight profile $\nu \in X(A_0) \otimes \mathbb{Q}$. Goresky and MacPherson computed the Lefschetz number of $\Gamma\zeta\Gamma$ acting on weighted cohomology:

Theorem [Goresky and MacPherson 2003, 1.4 Theorem].

$$\begin{aligned} \text{Lef}(\Gamma\zeta\Gamma | W^\nu H^*(\overline{\Gamma\backslash X}, L_{\tilde{\lambda}}(\mathbb{C}))) \\ = \sum_{\{\tilde{P}\}} \sum_i \sum_{\{\xi\}} \Xi_{\tilde{P}, \xi} \sum_{\substack{w \in W^{\tilde{P}} \\ I_\nu(w, \tilde{\lambda}) = \Delta_{\tilde{P}}^+(\xi)}} (-1)^{\ell(w)} \text{tr}(\xi^{-1} | L_{w \cdot \tilde{\lambda}}^{\tilde{M}}(F)). \end{aligned}$$

In this formula, \tilde{P} runs over a choice of representatives for the Γ -conjugacy classes of \mathbb{Q} -parabolic subgroups of \tilde{G} . For each such \tilde{P} we write $\Gamma\zeta\Gamma \cap \tilde{P}(\mathbb{Q}) = \coprod_i \Gamma_{\tilde{P}} \zeta_i \Gamma_{\tilde{P}}$ where $\Gamma_{\tilde{P}} = \Gamma \cap \tilde{P}(\mathbb{Q})$ and $\zeta_i \in \tilde{P}(\mathbb{Q})$. The second sum runs over these finitely many double cosets. We set $\Gamma_{\tilde{M}} = \nu_{\tilde{P}}(\Gamma_{\tilde{P}}) \subseteq \tilde{M}(\mathbb{Q})$ and $\zeta_i = \nu_{\tilde{P}}(\zeta_i) \in \tilde{M}(\mathbb{Q})$; here $\tilde{P} = \tilde{M}\tilde{N}$ is the Levi decomposition and $\nu_{\tilde{P}} : \tilde{P} \rightarrow \tilde{M}$ is the canonical mapping. The third sum is over a set of representatives ξ of the $\Gamma_{\tilde{M}}$ -conjugacy classes of elliptic (modulo $A_{\tilde{P}}(\mathbb{R})$) elements in $\Gamma_{\tilde{M}} \zeta_i \Gamma_{\tilde{M}} \subseteq \tilde{M}(\mathbb{Q})$. (The numbers $\Xi_{\tilde{P}, \xi}$ are explained in [Goresky and MacPherson 2003, 1.4]; we only need to know that they are contained in \mathbb{Z} and do not depend on the weight $\tilde{\lambda}$.) Moreover,

$$\Delta_{\tilde{P}}^+(\xi) = \{\alpha \in \Delta_{\tilde{P}} : \alpha(a_\xi) < 1\},$$

where a_ξ is the projection of ξ to the identity component $A_{\tilde{P}}$ of $A_{\tilde{P}}(\mathbb{R})$ and $I_\nu(w, \tilde{\lambda})$ is given as

$$I_\nu(w, \tilde{\lambda}) = \{\alpha_i \in \Delta_{\tilde{P}} : \langle w(\tilde{\lambda} + \rho) - \rho - \nu, t_{\alpha_i} \rangle < 0\},$$

where $\rho = \rho_{\tilde{G}}$. Finally, since $\xi \in \tilde{M}(\mathbb{Q})$ and $L_{w \cdot \tilde{\lambda}}^{\tilde{M}}$ is defined over F , the trace may be computed on F -points of $L_{w \cdot \tilde{\lambda}}^{\tilde{M}}$. We note that to make sense of the trace of $\xi^{-1} \in \tilde{M}(\mathbb{Q})$ on $L_{\tilde{\lambda}}^{\tilde{M}}(F)$ ($L_{\tilde{\lambda}}^{\tilde{M}}$ was defined for standard parabolic subgroups) as well as of the definition of $I_\nu(w, \tilde{\lambda})$ ($\tilde{\lambda}, \rho$ are characters of \tilde{T}) we have to conjugate \tilde{P} , i.e., we select $x \in \tilde{G}(F)$ such that ${}^x\tilde{P}/F$ is standard parabolic.

4.14.2. We select a \mathbb{Q} -parabolic subgroup \tilde{P} in \tilde{G} with Levi decomposition $\tilde{P} = \tilde{M}\tilde{N}$. For any $w \in W^{\tilde{P}}$ and any $\alpha_i \in \Delta_{\tilde{P}}$ the assignment $L_{w,i} : X(\tilde{T}) \otimes \mathbb{Q} \rightarrow \mathbb{Q}$ taking $\tilde{\lambda}$ to $\langle w\tilde{\lambda}, t_{\alpha_i} \rangle$ is linear and we denote by $H_{w,i}^+$ (resp. $H_{w,i}^-$) the half space consisting of all $\tilde{\lambda} \in X(\tilde{T})$ such that $\langle w(\tilde{\lambda} + \rho) - \rho - \nu, t_{\alpha_i} \rangle$ is positive (resp. negative). For

any $\underline{\epsilon} = (\epsilon_{w,i}) \in \{\pm\}^{W^{\tilde{P}} \times [1, \dots, m]}$ we set

$$C(\underline{\epsilon}) = X(\tilde{T})^{\text{dom}} \cap \bigcap_{w,i} H_{w,i}^{\epsilon(w,i)}.$$

Thus, $C(\underline{\epsilon})$ is an intersection of half spaces which may be empty. For all $w \in W^{\tilde{P}}$ and $i = 1, \dots, m$ the values $\langle w(\tilde{\lambda} + \rho) - \rho - \nu, t_{\alpha_i} \rangle$ and $\langle w(\tilde{\lambda}' + \rho) - \rho - \nu, t_{\alpha_i} \rangle$, where $\tilde{\lambda}, \tilde{\lambda}' \in C(\underline{\epsilon})$ have the same sign, hence, for all $w \in W^{\tilde{P}}$ we obtain $I_\nu(w, \tilde{\lambda}) = I_\nu(w, \tilde{\lambda}')$, i.e., $I_\nu(w, \tilde{\lambda})$ does not depend on $\tilde{\lambda} \in C(\underline{\epsilon})$.

Theorem. *Let $C \in \mathbb{Q}_{>0}$ and $\underline{\epsilon} \in \{\pm\}^{W^{\tilde{P}} \times [1, \dots, m]}$. Let $\tilde{\lambda}, \tilde{\lambda}' \in C(\underline{\epsilon})$ be (dominant) weights satisfying*

- $\langle \tilde{\lambda}, \alpha^\vee \rangle > 2C$ and $\langle \tilde{\lambda}', \alpha^\vee \rangle > 2C$ for all $\alpha \in \Delta_{\tilde{G}}$,
- $\tilde{\lambda} \equiv \tilde{\lambda}' \pmod{(p-1)p^{m-1}X(\tilde{T})}$ ($m \in \mathbb{N}$).

Let ζ be contained in the semigroup Δ_h , hence, $\zeta \in \tilde{I}h^e\tilde{I}$ for some $e \in \mathbb{N}_0$ and we assume that $e \in \mathbb{N}$. Then the Lefschetz number $\text{Lef}((\Gamma\zeta\Gamma)_{\tilde{\lambda}}|W^\nu H^\bullet(\overline{\Gamma\backslash X}, L_{\tilde{\lambda}}(F)))$ is contained in F and the following congruence holds:

$$\begin{aligned} \text{Lef}((\Gamma\zeta\Gamma)_{\tilde{\lambda}}|W^\nu H^\bullet(\overline{\Gamma\backslash X}, L_{\tilde{\lambda}}(F))) &\equiv \text{Lef}((\Gamma\zeta\Gamma)_{\tilde{\lambda}'}|W^\nu H^\bullet(\overline{\Gamma\backslash X}, L_{\tilde{\lambda}'}(F))) \\ &\pmod{p^{\lceil \min((C-\kappa_1-\kappa_2)e, m-e\kappa_1) \rceil} \mathbb{Z}_p}. \end{aligned}$$

Proof. We look at the Goresky–MacPherson trace formula. Since $\Gamma\zeta\Gamma \cap \tilde{P}(\mathbb{Q}) \supseteq \Gamma_{\tilde{P}}\zeta_i\Gamma_{\tilde{P}}$ we obtain $\zeta_i \in \Gamma\zeta\Gamma \cap \tilde{P}(\mathbb{Q})$, hence, we can write $\zeta_i = \tilde{\zeta}_i u$ where $\tilde{\zeta}_i \in \tilde{M}(\mathbb{Q})$ and $u \in \tilde{N}(\mathbb{Q})$. Let $\xi \in \Gamma_{\tilde{M}}\tilde{\zeta}_i\Gamma_{\tilde{M}}$ be a representative of a $\Gamma_{\tilde{M}}$ -conjugacy class. We may assume that $\xi = \gamma_M \tilde{\zeta}_i$ for some $\gamma_M \in \Gamma_{\tilde{M}}$, i.e., $\gamma_M = \nu_{\tilde{P}}(\gamma_P)$ for some $\gamma_P \in \Gamma_{\tilde{P}}$. Hence, we can write $\gamma_P = \gamma_N \gamma_M$, where $\gamma_N \in \tilde{N}(\mathbb{Q})$, and obtain

$$\gamma_N \xi u = \gamma_N \gamma_M \tilde{\zeta}_i u = \gamma_P \zeta_i \in \Gamma_{\tilde{P}}\zeta_i \subseteq \Gamma\zeta\Gamma \subseteq \tilde{I}h^e\tilde{I}.$$

Since $\tilde{M}(\mathbb{Q})$ normalizes $\tilde{N}(\mathbb{Q})$ we can write $\gamma_N \xi u = \xi u'$ with $u' \in \tilde{N}(\mathbb{Q})$. Thus, $\xi u' \in \tilde{I}h^e\tilde{I}$ and we may apply 4.7 Proposition to compute $\text{tr}(\xi^{-1}|L_{w,\tilde{\lambda}}^{\tilde{M}}(\mathbb{Q}_p))$. If we put the result in the trace formula of Goresky–MacPherson we obtain

$$\begin{aligned} (59) \quad &\text{Lef}((\Gamma\zeta\Gamma)_{\tilde{\lambda}}|W^\nu H^\bullet(\overline{\Gamma\backslash X}, L_{\tilde{\lambda}}(F))) \\ &\equiv \sum_{\{\tilde{P}\}} \sum_{i\xi} \sum_{\{\xi\}} \Xi_{\tilde{P},\xi} \delta_s (-1)^{\ell(s)} \epsilon_{\tilde{P},\xi u'} \tilde{\lambda}(h^e t^{-1}) \pmod{p^{(C-\kappa_1-\kappa_2)e} \mathcal{O}_{\tilde{\mathbb{Q}}_p}}, \end{aligned}$$

where $s = s_{\xi u'} \in W^{\tilde{P}}$, $t = t_{\xi u'} \in \tilde{T}(\tilde{\mathbb{Q}}_p)^{++}$ is the unique element which is $\tilde{G}(\tilde{\mathbb{Q}}_p)$ -conjugate to $(\xi u')_s$, $\epsilon_{\tilde{P},\xi u'} \in \tilde{P}$ has p -adic value $\geq -e\kappa_1$ and $\delta_s = 1$ if $I_\nu(s, \tilde{\lambda}) = \Delta_{\tilde{P}}^+(\xi)$ and vanishes otherwise. We note that δ_s does not depend on $\tilde{\lambda}$ since we assume that $\tilde{\lambda}$ is contained in $C(\underline{\epsilon})$. By our assumption there is $\chi \in X(\tilde{T})$ such that $\tilde{\lambda} - \tilde{\lambda}' = (p-1)p^{m-1}\chi$. Since $\chi(h^e t^{-1}) \equiv \epsilon \pmod{p\mathcal{O}_{\tilde{\mathbb{Q}}_p}}$ where $\epsilon \in \mathbb{Z}_p^*$ (see 4.4 Proposition) this implies $\tilde{\lambda}(h^e t^{-1}) \equiv \tilde{\lambda}'(h^e t^{-1}) \pmod{p^m \mathcal{O}_{\tilde{\mathbb{Q}}_p}}$ (see the proof of

4.10 Proposition) and we find that equation (59) (which also holds for $\tilde{\lambda}'$) implies that the Lefschetz numbers in the Theorem are congruent to each other modulo $p^{\lceil \min((C-\kappa_1-\kappa_2)e, m-e\kappa_1) \rceil} \mathbb{Z}_p$ (note that the Lefschetz numbers are contained in F). Thus the proof is complete. \square

4.14.3. The case C_2 . We assume that \tilde{G}/\mathbb{Q} is connected and \mathbb{Q} -split with root system of type C_2 , hence, there are two simple roots α_1, α_2 . We also assume that $\nu = -\rho$, i.e., ν is the middle weight profile. Thus, if $\tilde{\lambda}$ is regular then $W^\nu H^d(\overline{\Gamma \backslash X}, L_{\tilde{\lambda}}(\mathbb{C})) = H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))$ and both these cohomology groups vanish for all degrees $i \neq d$. Let $\tilde{\lambda} \in X(\tilde{T})^{\text{dom}}$. Hence, $\tilde{\lambda} + \rho \in X(\tilde{T})^{\text{dom}}$ is strictly dominant and for any $w \in W_{\tilde{G}}$ we write $w(\tilde{\lambda} + \rho) = a_1\alpha_1 + a_2\alpha_2$ with $a_i = a_{i,w,\tilde{\lambda}} \in \mathbb{Q}$. For the root system of type C_2 it happens that the sign of a_1 as well as the sign of a_2 does not depend on $\tilde{\lambda} \in X(\tilde{T})^{\text{dom}}$, i.e., $\text{sign}(a_{i,w,\tilde{\lambda}}) = \text{sign}(a_{i,w,\tilde{\lambda}'})$ for any $\tilde{\lambda}, \tilde{\lambda}' \in X(\tilde{T})^{\text{dom}}$ and all $w \in W_{\tilde{G}}$ and all $i = 1, 2$. Thus, if $\tilde{P} = \tilde{B}$, hence, $A_{\tilde{P}} = \tilde{T}/\mathbb{Q}$ and if $\{t_{\alpha_1}, t_{\alpha_2}\}$ denotes the basis of $X_*(\tilde{T}) \otimes \mathbb{Q}$ dual to $\{\alpha_1, \alpha_2\}$ then we obtain that the sign of $\langle w(\tilde{\lambda} + \rho), t_{\alpha_i} \rangle = a_i$ does not depend on $\tilde{\lambda} \in X(\tilde{T})^{\text{dom}}$. Similarly if $\tilde{P} = \tilde{P}_{\alpha_i}$, hence, $A_{\tilde{P}} = \ker \alpha_j, j \neq i$, and if $\{t_{\alpha_i}\}$ denotes the basis of $X_*(A_{\tilde{P}}) \otimes \mathbb{Q}$ dual to $\{\alpha_i\}$ then we obtain that the sign of $\langle w(\tilde{\lambda} + \rho)|_{A_{\tilde{P}}}, t_{\alpha_i} \rangle = a_i$ does not depend on $\tilde{\lambda} \in X(\tilde{T})^{\text{dom}}$. This shows that $I_\nu(w, \tilde{\lambda})$ does not depend on $\tilde{\lambda} \in X(\tilde{T})^{\text{dom}}$ (if $\tilde{P} = \tilde{G}$ then $I_\nu(w, \tilde{\lambda})$ is empty). 4.14.2 Theorem therefore holds with $X(\tilde{T})^{\text{dom}}$ in place of $C(\epsilon)$:

Corollary. Assume \tilde{G}/\mathbb{Q} is connected and \mathbb{Q} -split with root system C_2 . Let $C \in \mathbb{Q}_{>0}$ and let $\tilde{\lambda}, \tilde{\lambda}' \in X(\tilde{T})^{\text{dom}}$ be weights satisfying

- $\langle \tilde{\lambda}, \alpha^\vee \rangle > 2C$ and $\langle \tilde{\lambda}', \alpha^\vee \rangle > 2C$ for all $\alpha \in \Delta_{\tilde{G}}$, and
- $\tilde{\lambda} \equiv \tilde{\lambda}' \pmod{(p-1)p^{m-1}X(\tilde{T})}$ ($m \in \mathbb{N}$).

Then $\text{tr}((\Gamma \zeta \Gamma)_{\tilde{\lambda}} | H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C})))$ is contained in F and

$$\text{tr}((\Gamma \zeta \Gamma)_{\tilde{\lambda}} | H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))) \equiv \text{tr}((\Gamma \zeta \Gamma)_{\tilde{\lambda}'} | H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}'}(\mathbb{C}))) \pmod{p^{\lceil \min((C-\kappa_1-\kappa_2)e, m-e\kappa_1) \rceil} \mathbb{Z}_p}.$$

Proof. Use the equality

$$\text{tr}((\Gamma \zeta \Gamma)_{\tilde{\lambda}} | H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))) = (-1)^d \text{Lef}((\Gamma \zeta \Gamma)_{\tilde{\lambda}} | W^\nu H^\bullet(\overline{\Gamma \backslash X}, L_{\tilde{\lambda}}(\mathbb{C}))). \quad \square$$

Remark. The \mathbb{Q} -rank of \tilde{G} does not appear in the modulus of these congruences.

5. Local constancy of dimension of slope subspaces

5.1. Slope subspaces of cuspidal cohomology. As before we let \tilde{G}/\mathbb{Q} be a connected reductive group with \mathbb{Q}_p -split maximal torus \tilde{T}/\mathbb{Q} and from now on we assume in addition that \tilde{G}/\mathbb{Q} has discrete series. We denote by ℓ the \mathbb{Q} -rank of

\tilde{G} and we keep the notations from Section 4.11. In particular, $\tilde{K} \leq \tilde{G}(\mathbb{A}_f)$ is a compact open subgroup with p -component $\tilde{K}_p = \tilde{I}$ and $S_{\tilde{K}}$ is the adelic locally symmetric space of level \tilde{K} . We select elements $h \in \tilde{T}(\mathbb{Q})^{++}$ and $r \in \tilde{G}(\mathbb{A}_f)^{(p)}$ and we set

$$\mathbb{T} = [\tilde{K}[h]_p^{-1}r\tilde{K}] \in \mathcal{C}_0(\tilde{G}(\mathbb{A}_f)//\tilde{K})_h^{\text{reg}}.$$

As before we denote by $\mathbb{T}_{\tilde{\lambda}}$ the $\tilde{\lambda}$ -normalization of \mathbb{T} . The (normalized) operator $\mathbb{T}_{\tilde{\lambda}}$ acts on

$$H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C}) = H_{\text{cusp}}^d(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))$$

where $d = d_{\tilde{G}}$ is the middle degree. For $\beta \in \mathbb{Q}_{\geq 0}$ we denote by

$$H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})^\beta = \bigoplus_{\substack{\mu \in \bar{F} \hookrightarrow \bar{\mathbb{Q}}_p \\ v_p(\mu) = \beta}} H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})(\mu),$$

the slope β subspace of $H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})$ w.r.t. $\mathbb{T}_{\tilde{\lambda}}$; here, $H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})(\mu) = H_{\tilde{\lambda}, \text{cusp}, \mathbb{T}}^d(\mathbb{C})(\mu)$ is the generalized eigenspace attached to $\mathbb{T}_{\tilde{\lambda}}$ and the eigenvalue μ and we remark that the eigenvalues μ of $\mathbb{T}_{\tilde{\lambda}}$ on $H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})$ are contained in $\bar{F}(\subseteq \mathbb{C})$ and are p -adically integral, i.e., contained in $\mathcal{O}_{\bar{\mathbb{Q}}_p}$ (note that $\bar{F} \subseteq \bar{\mathbb{Q}}_p$; see Section 4.11.2). We set $H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})^{\leq \beta} = \bigoplus_{0 \leq \gamma \leq \beta} H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})^\gamma$. Since

$$\begin{aligned} \dim H_{\text{cusp}}^d(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))^{\leq \beta} &= \dim \bigoplus_{\substack{\mu \in \bar{F} \hookrightarrow \bar{\mathbb{Q}}_p \\ 0 \leq v_p(\mu) \leq \beta}} H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})(\mu) \\ &\leq \dim \bigoplus_{\substack{\mu \in \bar{F} \\ 0 \leq v_p(\mu) \leq \beta}} H^d(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))(\mu) \\ &= \dim \bigoplus_{\substack{\mu \in \bar{F} \\ 0 \leq v_p(\mu) \leq \beta}} H^d(S_{\tilde{K}}, L_{\tilde{\lambda}}(\bar{F}))(\mu) \\ &= \dim \bigoplus_{\substack{\mu \in \bar{F} \hookrightarrow \bar{\mathbb{Q}}_p \\ 0 \leq v_p(\mu) \leq \beta}} H^d(S_{\tilde{K}}, L_{\tilde{\lambda}}(\bar{\mathbb{Q}}_p))(\mu) \\ &= \dim H^d(S_{\tilde{K}}, L_{\tilde{\lambda}}(\bar{\mathbb{Q}}_p))^{\leq \beta} \\ &= \dim H^d(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{Q}_p))^{\leq \beta}, \end{aligned}$$

we obtain the following bound from 4.11.4 Theorem. Recall that $s = |\Phi_{\tilde{G}}^+|$, $\sigma = \max_{\alpha \in \Phi_{\tilde{G}}^+} \text{ht}(\alpha)$ and we denote by $\mathbf{g} = \mathbf{g}_{\tilde{K}}$ the number of d cells in a cell complex

which is homotopy equivalent to the Borel–Serre compactification $\bar{S}_{\tilde{K}}$. We set

$$m = m_{\tilde{K}} = 12 \frac{g}{s} \sigma^{s+1} \in \mathbb{Q}_{\geq 0} \quad \text{and} \quad n = n_{\tilde{K}} = \left\lceil \frac{1}{2^{s+1}} \frac{g \sigma^{s+1}}{s} M^s \right\rceil + 1 \in \mathbb{N},$$

where $M = M(\sigma, s) \in \mathbb{N}$ is defined in equations (23) and (24); then

$$(60) \quad \dim H_{\text{cusp}}^d(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))^{\leq \beta} \leq B(\beta) := m\beta^s + n$$

for all dominant $\tilde{\lambda} \in X(\tilde{T})$, all $\beta \in \mathbb{Q}_{\geq 0}$ and all $h \in \tilde{T}(\mathbb{Q})^{++}$ and $r \in \tilde{G}(\mathbb{A}_f)^{(p)}$.

5.2. We want to consider the function

$$d(\cdot, \cdot) : \mathbb{Q}_{\geq 0} \times X(\tilde{T})^{\text{dom}} \rightarrow \mathbb{N}_0, \quad (\beta, \tilde{\lambda}) \mapsto \dim H_{\text{cusp}}^d(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))^\beta;$$

i.e., we want to understand how the dimension of the slope subspace varies as a function of the weight. To this end, for any $\beta \in \mathbb{Q}_{\geq 0}$ we set

$$(61) \quad \begin{aligned} m_1(\beta) &= \left(\beta + \frac{1}{p-1} + (\kappa_1 + \kappa_2)\ell\right)B(\beta) + 1 \\ &= m\beta^{s+1} + m\left(\frac{1}{p-1} + (\kappa_1 + \kappa_2)\ell\right)\beta^s + n\beta + n\left(\frac{1}{p-1} + (\kappa_1 + \kappa_2)\ell\right) + 1 \\ &\in \mathbb{Q}_{>0}. \end{aligned}$$

and

$$(62) \quad \begin{aligned} m_2(\beta) &= \left(\beta + \frac{1}{p-1} + \kappa_1\ell\right)B(\beta) + 1 \\ &= m\beta^{s+1} + m\left(\frac{1}{p-1} + \kappa_1\ell\right)\beta^s + n\beta + n\left(\frac{1}{p-1} + \kappa_1\ell\right) + 1 \in \mathbb{Q}_{>0}. \end{aligned}$$

Thus, $m_1(\beta), m_2(\beta) \in \mathbb{Q}[\beta]$ are polynomials in β with positive coefficients, degree $s + 1$ and leading term $m = 12 \frac{g}{s} \sigma^{s+1}$ which depend on \tilde{K} (and, hence, on \tilde{G} and p) and on h (since $\kappa_1 = \kappa_1(h)$) but do not depend on $\tilde{\lambda} \in X(\tilde{T})$.

Theorem. *Assume that \tilde{G} has discrete series. Let $\beta \in \mathbb{Q}_{\geq 0}$ and assume the dominant weights $\tilde{\lambda}, \tilde{\lambda}' \in X(\tilde{T})$ satisfy*

- $\langle \tilde{\lambda}, \alpha^\vee \rangle > 2m_1(\beta)$ and $\langle \tilde{\lambda}', \alpha^\vee \rangle > 2m_1(\beta)$ for all $\alpha \in \Delta_{\tilde{G}}$,
- $\tilde{\lambda} \equiv \tilde{\lambda}' \pmod{(p-1)p^{m-1}X(\tilde{T})}$ with $m \geq m_2(\beta)$ ($m \in \mathbb{N}$).

Then

$$\dim H_{\text{cusp}}^d(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))^\gamma = \dim H_{\text{cusp}}^d(S_{\tilde{K}}, L_{\tilde{\lambda}'}(\mathbb{C}))^\gamma \quad \text{for all } 0 \leq \gamma \leq \beta.$$

5.3. Characteristic polynomial. The proof of the preceding theorem will be given in Section 5.6. To prepare it we consider the characteristic polynomial. We denote by $\text{ch}_{\tilde{\lambda}}(X) = \det(X\mathbf{1} - \mathbb{T}_{\tilde{\lambda}}) = \sum_{i=0}^m (-1)^i a_{i,\tilde{\lambda}} X^{m-i} \in \mathbb{C}[X]$, where $m = m_{\tilde{\lambda}} = \dim H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})$, the characteristic polynomial of $\mathbb{T}_{\tilde{\lambda}}$ acting on $H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})$. We set $a_{i,\tilde{\lambda}} = 0$ if $i > m_{\tilde{\lambda}}$. This definition of the characteristic polynomial differs from the one used in the proof of 4.3 Lemma by a factor $(-1)^m$, but we can refer directly to

[Koecher 1983, 3.4.6 Satz, p. 117], where this definition is used and which yields the following inductive formula for the coefficients of $\text{ch}_{\tilde{\lambda}}(X)$: $a_{0,\tilde{\lambda}} = 1$ and

$$(63) \quad ia_{i,\tilde{\lambda}} = \sum_{e=1}^i (-1)^{e+1} \text{tr } \mathbb{T}_{\tilde{\lambda}}^e a_{i-e,\tilde{\lambda}}, \quad i = 1, 2, \dots,$$

where we have set

$$\text{tr } \mathbb{T}_{\tilde{\lambda}}^e = \text{tr} ((\mathbb{T}_{\tilde{\lambda}})^e | H_{\tilde{\lambda},\text{cusp}}^d(\mathbb{C})).$$

Since $[\tilde{\mathcal{I}}h^{-e}\tilde{\mathcal{I}}][\tilde{\mathcal{I}}h^{-f}\tilde{\mathcal{I}}] = [\tilde{\mathcal{I}}h^{-(e+f)}\tilde{\mathcal{I}}]$ we obtain that $\mathbb{T}^e = \sum_s c_s [\tilde{K}[h]_p^{-e} s \tilde{K}]$ for all $e \geq 1$, where s runs over $\tilde{\mathbf{G}}(\mathbb{A}_f)^{(p)}$ and $c_s \in \mathbb{Z}$. Hence $(\mathbb{T}_{\tilde{\lambda}})^e = \sum_s c_s [\tilde{K}[h]_p^{-e} s \tilde{K}]_{\tilde{\lambda}}$. Since $\tilde{\mathbf{G}}$ has discrete series we therefore obtain if the highest weight $\tilde{\lambda}$ is regular

$$(64) \quad \begin{aligned} \text{tr } \mathbb{T}_{\tilde{\lambda}}^e &= \sum_s c_s \text{tr} ([\tilde{K}[h]_p^{-e} s \tilde{K}]_{\tilde{\lambda}} | H_{\tilde{\lambda},\text{cusp}}^d(\mathbb{C})) \\ &= \sum_s c_s (-1)^d \text{Lef}([\tilde{K}[h]_p^{-e} s \tilde{K}]_{\tilde{\lambda}} | H_{\text{cusp}}^{\bullet}(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))). \end{aligned}$$

Hence, equation (63) and 4.13.2 Theorem yield $a_{i,\tilde{\lambda}} \in F$ for all i , i.e.,

$$\text{ch}_{\tilde{\lambda}}(X) \in F[X].$$

In particular, the roots of $\text{ch}_{\tilde{\lambda}}$, which are the eigenvalues of $\mathbb{T}_{\tilde{\lambda}}$ on $H_{\tilde{\lambda},\text{cusp}}^d(\mathbb{C})$, are algebraic over F and in Section 4.11.2 we have seen that after embedding $\bar{F} \subseteq \bar{\mathbb{Q}}_p$ they are contained in $\mathcal{O}_{\bar{\mathbb{Q}}_p}$. Hence, $a_{i,\tilde{\lambda}} \in \mathcal{O}_{\bar{\mathbb{Q}}_p}$, i.e., the coefficients are p -adically integral which implies that $a_{i,\tilde{\lambda}} \in \mathbb{Z}_p$, $i = 0, \dots, m$; in particular,

$$\text{ch}_{\tilde{\lambda}}(X) \in \mathbb{Z}_p[X].$$

5.4. Proposition. *Let $\beta \in \mathbb{Q}_{\geq 0}$. Assume that the weights $\tilde{\lambda}, \tilde{\lambda}' \in X(\tilde{\mathbf{T}})^{\text{dom}}$ satisfy the two assumptions of 5.2 Theorem. Then for all $i = 0, 1, 2, 3, \dots, B(\beta)$ the following congruence holds:*

$$a_{i,\tilde{\lambda}} \equiv a_{i,\tilde{\lambda}'} \pmod{p^{\lceil \beta B(\beta) + 1 \rceil} \mathbb{Z}_p}.$$

Proof. We will prove that for all $i = 0, 1, \dots, B(\beta)$ the congruence

$$a_{i,\tilde{\lambda}} \equiv a_{i,\tilde{\lambda}'} \pmod{p^{\lceil \beta B(\beta) + \frac{B(\beta)}{p-1} + 1 \rceil - v_p(i!)} \mathbb{Z}_p}$$

holds. Since $v_p(i!) \leq i/(p-1) \leq B(\beta)/(p-1)$ these congruences imply that the congruences of the Proposition hold. The congruences hold trivially for $i = 0$ ($a_{0,\tilde{\lambda}} = a_{0,\tilde{\lambda}'} = 1$). To prove them for $i = 1, 2, 3, \dots, B(\beta)$ we use equation (63). First, for all $e \in \mathbb{N}$ with $1 \leq e \leq B(\beta)$ we have

$$m - e\kappa_1 \ell \geq m_2(\beta) - e\kappa_1 \ell \geq \beta B(\beta) + \frac{B(\beta)}{p-1} + 1$$

and

$$\begin{aligned} \lceil e(\mathbf{m}_1(\beta) - \kappa_2 \ell) \rceil - e\kappa_1 \ell &\geq \mathbf{m}_1(\beta) - e\kappa_2 \ell - e\kappa_1 \ell \\ &= \left(\beta + \frac{1}{p-1}\right)B(\beta) + (\kappa_1 + \kappa_2)\ell B(\beta) + 1 - e\kappa_2 \ell - e\kappa_1 \ell \\ &\geq \beta B(\beta) + \frac{B(\beta)}{p-1} + 1. \end{aligned}$$

Hence, 4.13.2 Theorem yields for all $1 \leq e \leq B(\beta)$ and all $s \in \tilde{\mathbf{G}}(\mathbb{A}_f)^{(p)}$ that

$$\begin{aligned} \text{Lef}([\tilde{K}[h]_p^{-e} s \tilde{K}]_{\tilde{\lambda}} | H_{\text{cusp}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}}(\mathbb{C}))) &\equiv \text{Lef}([\tilde{K}[h]_p^{-e} s \tilde{K}]_{\tilde{\lambda}'} | H_{\text{cusp}}^\bullet(S_{\tilde{K}}, L_{\tilde{\lambda}'}(\mathbb{C}))) \\ &\pmod{p^{\lceil \beta B(\beta) + \frac{B(\beta)}{p-1} + 1 \rceil} \mathbb{Z}_p}. \end{aligned}$$

Together with equation (64) this implies

$$(65) \quad \text{tr } \mathbb{T}_{\tilde{\lambda}}^e \equiv \text{tr } \mathbb{T}_{\tilde{\lambda}'}^e \pmod{p^{\lceil \beta B(\beta) + \frac{B(\beta)}{p-1} + 1 \rceil} \mathbb{Z}_p}$$

for all $1 \leq e \leq B(\beta)$. Since $a_{1,?} = \text{tr } \mathbb{T}_?$, equation (65) implies

$$a_{1,\tilde{\lambda}} \equiv a_{1,\tilde{\lambda}'} \pmod{p^{\lceil \beta B(\beta) + \frac{B(\beta)}{p-1} + 1 \rceil} \mathbb{Z}_p},$$

which is the claim for $i = 1$. We now let $i \leq B(\beta)$ be arbitrary and assume that the claim holds for $0, 1, 2, \dots, i - 1$. The recursive relation in equation (63)

$$ia_{i,?} = \sum_{e=1}^i (-1)^{e+1} \text{tr } \mathbb{T}_?^e a_{i-e,?}$$

together with the induction assumption and equation (65) yields

$$ia_{i,\tilde{\lambda}} \equiv ia_{i,\tilde{\lambda}'} \pmod{p^{\lceil \beta B(\beta) + \frac{B(\beta)}{p-1} + 1 \rceil - v_p((i-1)!) } \mathbb{Z}_p}$$

from which the claim for i is immediate. □

5.5. Newton polygon. The *Newton polygon* $\mathcal{N}_{\tilde{\lambda}}$ of $\text{ch}_{\tilde{\lambda}} \in F[X] \subseteq \mathbb{Q}_p[X]$ is the lower convex hull of the points $(0, v_p(a_{0,\tilde{\lambda}})), \dots, (m, v_p(a_{m,\tilde{\lambda}}))$, where we omit from this list all points $(i, v_p(a_{i,\tilde{\lambda}}))$ with $v_p(a_{i,\tilde{\lambda}}) = \infty$ (i.e., $a_{i,\tilde{\lambda}} = 0$). Thus, if $a_{m-k+1,\tilde{\lambda}} = \dots = a_{m,\tilde{\lambda}} = 0$ and $a_{m-k,\tilde{\lambda}} \neq 0$ (i.e., if 0 occurs in $\text{ch}_{\tilde{\lambda}}(X)$ with multiplicity $k = \text{ord}_X(\text{ch}_{\tilde{\lambda}})$), we omit the points $(m-k+1, v_p(a_{m-k+1,\tilde{\lambda}})), \dots, (m, v_p(a_{m,\tilde{\lambda}}))$. In particular, $\mathcal{N}_{\tilde{\lambda}}$ represents a piecewise linear function on the interval $[0, n_{\tilde{\lambda}}]$ which starts at the point $(0, 0)$ corresponding to the leading term $a_{0,\tilde{\lambda}} = 1$ of $\text{ch}_{\tilde{\lambda}}$; here,

$$n_{\tilde{\lambda}} = m - k = \dim H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C}) - \text{ord}_X(\text{ch}_{\tilde{\lambda}}) = \dim \bigoplus_{\gamma \in \mathbb{Q}_{\geq 0}} H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})^\gamma$$

is the dimension of the finite slope subspace $H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})^{<\infty}$ of $H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})$. We have $a_{i,\tilde{\lambda}} = 0$ for all $i > n_{\tilde{\lambda}}$ (note that $a_{i,\tilde{\lambda}} = 0$ for all $i > m_{\tilde{\lambda}}$). Since $\text{ch}_{\tilde{\lambda}}/X^k$, for

$k = \text{ord}_X(\text{ch}_{\tilde{\lambda}})$, also has $\mathcal{N}_{\tilde{\lambda}}$ as its Newton polygon but has nonvanishing constant term we deduce that if the segment S_γ of (necessarily finite) slope $\gamma \in \mathbb{Q}_{\geq 0}$ of $\mathcal{N}_{\tilde{\lambda}}$ has length s_γ (if projected to the x -axis) then $\text{ch}_{\tilde{\lambda}}/X^k$ and, hence, $\text{ch}_{\tilde{\lambda}}$ has precisely s_γ many roots (counted with their multiplicities) in $\bar{F}(\subseteq \bar{\mathbb{Q}}_p)$ of p -adic value equal to γ .

5.6. Proof of 5.2 Theorem. We denote by \mathcal{S} the finite set consisting of all $\gamma \in \mathbb{Q}_{\geq 0}$ such that $\gamma \leq \beta$ and the segment S_γ of slope γ of the Newton polygon $\mathcal{N}_{\tilde{\lambda}}$ or the segment S'_γ of slope γ of $\mathcal{N}_{\tilde{\lambda}'}$ has strictly positive length (i.e., $H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})^\gamma \neq 0$ or $H_{\tilde{\lambda}', \text{cusp}}^d(\mathbb{C})^\gamma \neq 0$). We have to show that

$$\dim H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})^\gamma = \dim H_{\tilde{\lambda}', \text{cusp}}^d(\mathbb{C})^\gamma \quad \text{for all } \gamma \in \mathcal{S}.$$

Since $\dim H_{\tilde{\lambda}, \text{cusp}}^d(\mathbb{C})^\gamma$ equals the number of roots $\mu \in \bar{F}$ of $\text{ch}_{\tilde{\lambda}}$ having p -adic value γ this is equivalent to showing that for all $\gamma \in \mathcal{S}$ the corresponding segments S_γ and S'_γ have the same length (length 0 if the slope γ subspace is trivial). We assume this is not the case and we denote by $\gamma \in \mathbb{Q}_{\geq 0}$ the smallest number in \mathcal{S} such that S_γ and S'_γ have different length; without loss of generality we may assume that S'_γ is (strictly) longer than S_γ . For all $\gamma^* \in \mathcal{S}$ with $0 \leq \gamma^* < \gamma$ the segments S_{γ^*} and S'_{γ^*} have the same length, hence, S_γ and S'_γ have the same initial point (note that $\mathcal{N}_{\tilde{\lambda}}, \mathcal{N}_{\tilde{\lambda}'}$ both start in $(0, 0)$). We denote by $P' = (x', y')$ the endpoint of S'_γ . Hence, (x', y') is a vertex of $\mathcal{N}_{\tilde{\lambda}'}$ which implies that $x' \in \mathbb{N}$ and $y' = v_p(a_{x', \tilde{\lambda}'})$. We also know that $x' \leq B(\beta)$ because $x' = \dim H_{\tilde{\lambda}', \text{cusp}}^d(\mathbb{C})^{\leq \gamma} \leq \dim H_{\tilde{\lambda}', \text{cusp}}^d(\mathbb{C})^{\leq \beta} \leq B(\beta)$. Since $x' \leq B(\beta)$ and since all segments of $\mathcal{N}_{\tilde{\lambda}'}$ which are left to x' have slopes less than or equal to γ we deduce that

$$(66) \quad v_p(a_{x', \tilde{\lambda}'}) = y' \leq \gamma B(\beta) \leq \beta B(\beta).$$

Together with 5.4 Proposition this implies that

$$(67) \quad v_p(a_{x', \tilde{\lambda}}) = v_p(a_{x', \tilde{\lambda}'}).$$

We distinguish cases.

Case 1: $\mathcal{N}_{\tilde{\lambda}}$ is defined at x' (i.e., $x' \leq n_{\tilde{\lambda}}$). Since $\mathcal{N}_{\tilde{\lambda}}$ is convex and S_γ is strictly shorter than S'_γ in this case we know that $\mathcal{N}_{\tilde{\lambda}}(x')$ lies strictly above $\mathcal{N}_{\tilde{\lambda}'}(x') = y'$, hence, we obtain

$$v_p(a_{x', \tilde{\lambda}}) \geq \mathcal{N}_{\tilde{\lambda}}(x') > \mathcal{N}_{\tilde{\lambda}'}(x') = y' = v_p(a_{x', \tilde{\lambda}'}),$$

i.e., we obtain a contradiction to equation (67).

Case 2: $\mathcal{N}_{\tilde{\lambda}}$ is not defined at x' (i.e., $x' > n_{\tilde{\lambda}}$). In this case we know that $a_{x', \tilde{\lambda}} = 0$. Since $v_p(a_{x', \tilde{\lambda}'})$ is finite by equation (66), again, this contradicts equation (67).

Thus, the assumption does not hold and we have shown that all segments of slope $\leq \beta$ in $\mathcal{N}_{\tilde{\lambda}}$ and $\mathcal{N}_{\tilde{\lambda}'}$ have the same length which finishes the proof.

5.7. Remark. The theorem in particular implies that for any $\beta \in \mathbb{Q}_{\geq 0}$ the function $\tilde{\lambda} \mapsto d(\beta, \tilde{\lambda})$ is constant on cosets modulo $(p - 1)p^{\lceil m_2(\beta) \rceil - 1} X(\tilde{T})$.

5.8. Example: C_2/\mathbb{Q} . We look at a special case and assume that \tilde{G}/\mathbb{Q} is connected and \mathbb{Q} -split with root system of type C_2 . As in Section 4.14 we use a non-adelic setting, i.e., $\Gamma \leq \tilde{G}(\mathbb{Q})$ is an arithmetic subgroup contained in \tilde{T} , $h \in \tilde{T}(\mathbb{Q})^{++}$ and we set $T = \Gamma \zeta \Gamma \in \mathcal{H}$ where $\zeta \in \Delta_h$. We assume $\zeta \in \tilde{T}h\tilde{T}$. All eigenvalues of $T_{\tilde{\lambda}}$ on $H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))$ are algebraic over F and p -adically integral (see 2.9 Corollary) and we define the slope β subspace $H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))^\beta$ as the sum of the generalized $T_{\tilde{\lambda}}$ -eigenspaces attached to eigenvalues of p -adic value β . As in Section 5.1 we obtain that 3.10 Corollary implies

$$\dim H_{\text{cusp}}^d(\Gamma \backslash X, L_{\tilde{\lambda}}(\mathbb{C}))^{\leq \beta} \leq B_\Gamma(\beta),$$

where $B_\Gamma(\beta) = \mathfrak{m}_\Gamma \beta^s + \mathfrak{n}_\Gamma$. We define the polynomials

$$\begin{aligned} m_1(\beta) &= \beta B_\Gamma(\beta) + \frac{B_\Gamma(\beta)}{p-1} + 1 + B_\Gamma(\beta)(\kappa_1 + \kappa_2) \\ &= \mathfrak{m}_\Gamma \beta^{s+1} + \left(\frac{\mathfrak{m}_\Gamma}{p-1} + \mathfrak{m}_\Gamma(\kappa_1 + \kappa_2) \right) \beta^s + \mathfrak{n}_\Gamma \beta + \frac{\mathfrak{n}_\Gamma}{p-1} + \mathfrak{n}_\Gamma(\kappa_1 + \kappa_2) + 1 \end{aligned}$$

and

$$\begin{aligned} m_2(\beta) &= \beta B_\Gamma(\beta) + \frac{B_\Gamma(\beta)}{p-1} + 1 + B_\Gamma(\beta)\kappa_1 \\ &= \mathfrak{m}_\Gamma \beta^{s+1} + \left(\frac{\mathfrak{m}_\Gamma}{p-1} + \mathfrak{m}_\Gamma \kappa_1 \right) \beta^s + \mathfrak{n}_\Gamma \beta + \frac{\mathfrak{n}_\Gamma}{p-1} + \mathfrak{n}_\Gamma \kappa_1 + 1. \end{aligned}$$

Following the arguments in the previous subsections but using the congruences for the traces of Hecke operators in 4.14.3 Corollary instead of those in 4.13.2 Theorem we obtain this:

Theorem. *If \tilde{G} is connected and \mathbb{Q} -split with root system of type C_2 then 5.2 Theorem holds with m_1, m_2 as defined above.*

We want to determine the polynomial m_2 more explicitly. Since \tilde{G}/\mathbb{Q} is of type C_2 there are two simple roots $\Delta = \{\alpha_1, \alpha_2\}$, the positive roots are $\Phi^+ = \{\alpha_1, \alpha_2, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2\}$, hence, $s = 4$ and $2\rho_{\tilde{G}} = 4\alpha_1 + 3\alpha_2$. We denote by $g = g_d$ the number of d cells in a cell complex which is homotopy equivalent to the Borel–Serre compactification of $\Gamma \backslash X$. We assume that $h \in \tilde{T}(\mathbb{Q})^{++}$ satisfies $v_p(\alpha_1(h)) = v_p(\alpha_2(h)) = 1$. We then obtain

$$\kappa_1 = \sum_{\alpha > 0} v_p(\alpha(h)) = 7, \quad \kappa_2 = \max(m_{\alpha_1}, m_{\alpha_2}) = 4, \quad \sigma = \max_{\alpha > 0} \text{ht}(\alpha) = 3,$$

and hence

$$m_{\Gamma} = \frac{12g\sigma^{s+1}}{s} = \frac{12g3^5}{4} = 729g.$$

To determine n_{Γ} we have to find $M \in \mathbb{N}$ such that equations (23) and (24) hold. A numerical evaluation shows that we may choose $M = 34$; hence, equation (26) yields

$$n_{\Gamma} = \left\lceil 2^{1/(s+1)} \frac{g\sigma^{s+1} M^s}{s} \right\rceil + 1 = \left\lceil \frac{2^{1/5} g 3^5 1336336}{4} \right\rceil + 1 \leq 93254104g + 1.$$

This yields the bound

$$\begin{aligned} m_2(X) &\leq 729gX^5 + 729g\left(\frac{1}{p-1} + 7\right)X^4 + (93254104g + 1)X \\ &\quad + (93254104g + 1)\left(\frac{1}{p-1} + 7\right) + 1 \\ &\leq 729gX^5 + 5832gX^4 + (93254104g + 1)X + 746032832g + 9. \end{aligned}$$

We note that since p is in the level of Γ the number $g = g_{\Gamma}$ depends on the prime p .

Acknowledgement

I am grateful to A. Nair for pointing out an error in an earlier version and for enlightening discussions concerning the work of Goresky–Kottwitz–MacPherson.

References

- [Andreatta et al. 2015] F. Andreatta, A. Iovita, and V. Pilloni, “ p -adic families of Siegel modular cuspforms”, *Ann. of Math. (2)* **181**:2 (2015), 623–697. MR Zbl
- [Ash and Stevens 2008] A. Ash and G. Stevens, “ p -adic deformations of arithmetic cohomology”, preprint, 2008, <https://www2.bc.edu/~ashav/Papers/Ash-Stevens-Oct-07-DRAFT-copy.pdf>.
- [Bewersdorff 1985] J. Bewersdorff, *Eine Lefschetzsche Fixpunktformel für Hecke-Operatoren*, Dissertation (Bonner Mathematische Schriften **164**), Universität Bonn, 1985. MR Zbl
- [Borel 1970] A. Borel, “Properties and linear representations of Chevalley groups”, pp. 1–55 in *Seminar on algebraic groups and related finite groups* (Princeton, NJ, 1968/69), Lecture Notes in Mathematics **131**, Springer, Berlin, 1970. MR Zbl
- [Borel and Serre 1973] A. Borel and J.-P. Serre, “Corners and arithmetic groups”, *Comment. Math. Helv.* **48** (1973), 436–491. MR Zbl
- [Brown 1982] K. S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics **87**, Springer, New York, 1982. MR Zbl
- [Buzzard 1998] K. Buzzard, “ p -adic modular forms on definite quaternion algebras”, preprint, 1998, <http://wwwf.imperial.ac.uk/~buzzard/maths/research/notes/definite.dvi>.
- [Buzzard 2001] K. Buzzard, “Families of modular forms”, *J. Théor. Nombres Bordeaux* **13**:1 (2001), 43–52. MR Zbl
- [Buzzard and Calegari 2004] K. Buzzard and F. Calegari, “A counterexample to the Gouvêa–Mazur conjecture”, *C. R. Math. Acad. Sci. Paris* **338**:10 (2004), 751–753. MR Zbl

- [Coleman 1997] R. F. Coleman, “ p -adic Banach spaces and families of modular forms”, *Invent. Math.* **127**:3 (1997), 417–479. MR Zbl
- [Franke 1998] J. Franke, “Harmonic analysis in weighted L_2 -spaces”, *Ann. Sci. École Norm. Sup.* (4) **31**:2 (1998), 181–279. MR Zbl
- [Goresky and MacPherson 2003] M. Goresky and R. MacPherson, “The topological trace formula”, *J. Reine Angew. Math.* **560** (2003), 77–150. MR Zbl
- [Goresky et al. 1994] M. Goresky, G. Harder, and R. MacPherson, “Weighted cohomology”, *Invent. Math.* **116**:1-3 (1994), 139–213. MR Zbl
- [Goresky et al. 1997] M. Goresky, R. Kottwitz, and R. MacPherson, “Discrete series characters and the Lefschetz formula for Hecke operators”, *Duke Math. J.* **89**:3 (1997), 477–554. MR Zbl
- [Gouvêa and Mazur 1992] F. Gouvêa and B. Mazur, “Families of modular eigenforms”, *Math. Comp.* **58**:198 (1992), 793–805. MR Zbl
- [Hida 1988] H. Hida, “On p -adic Hecke algebras for GL_2 over totally real fields”, *Ann. of Math.* (2) **128**:2 (1988), 295–384. MR Zbl
- [Hida 1993] H. Hida, *Elementary theory of L -functions and Eisenstein series*, London Mathematical Society Student Texts **26**, Cambridge University Press, 1993. MR Zbl
- [Humphreys 1972] J. E. Humphreys, *Introduction to Lie algebras and representation theory*, Graduate Texts in Mathematics **9**, Springer, New York, 1972. MR Zbl
- [Iwahori and Matsumoto 1965] N. Iwahori and H. Matsumoto, “On some Bruhat decomposition and the structure of the Hecke rings of p -adic Chevalley groups”, *Inst. Hautes Études Sci. Publ. Math.* **1965**:25 (1965), 5–48. MR Zbl
- [Jantzen 2003] J. C. Jantzen, *Representations of algebraic groups*, 2nd ed., Mathematical Surveys and Monographs **107**, American Mathematical Society, Providence, RI, 2003. MR Zbl
- [Koecher 1983] M. Koecher, *Lineare Algebra und analytische Geometrie*, Grundwissen Mathematik **2**, Springer, Berlin, 1983. MR Zbl
- [Koike 1975] M. Koike, “On some p -adic properties of the Eichler–Selberg trace formula”, *Nagoya Math. J.* **56** (1975), 45–52. MR Zbl
- [Koike 1976] M. Koike, “On p -adic properties of the Eichler–Selberg trace formula, II”, *Nagoya Math. J.* **64** (1976), 87–96. MR Zbl
- [Kuga et al. 1981] M. Kuga, W. Parry, and C. H. Sah, “Group cohomology and Hecke operators”, pp. 223–266 in *Manifolds and Lie groups* (Notre Dame, IN, 1980), Progr. Math. **14**, Birkhäuser, Boston, 1981. MR Zbl
- [Mahnkopf 2013] J. Mahnkopf, “On truncation of irreducible representations of Chevalley groups”, *J. Number Theory* **133**:9 (2013), 3149–3182. MR Zbl
- [Mahnkopf 2014] J. Mahnkopf, “On truncation of irreducible representations of Chevalley groups, II”, *J. Number Theory* **143** (2014), 46–61. MR Zbl
- [Nair 1999] A. Nair, “Weighted cohomology of arithmetic groups”, *Ann. of Math.* (2) **150**:1 (1999), 1–31. MR Zbl
- [Pande 2009] A. Pande, “Local constancy of dimensions of Hecke eigenspaces of automorphic forms”, *J. Number Theory* **129**:1 (2009), 15–27. MR
- [Popov and Vinberg 1994] V. L. Popov and E. B. Vinberg, *Invariant theory*, edited by I. R. Shafarevich and A. N. Parshin, Encyclopaedia of Mathematical Sciences **55**, Springer, Berlin, 1994. MR Zbl
- [Schwermer 1994] J. Schwermer, “Eisenstein series and cohomology of arithmetic groups: the generic case”, *Invent. Math.* **116**:1-3 (1994), 481–511. MR Zbl

- [Spallone 2009] S. Spallone, “Stable discrete series characters at singular elements”, *Canad. J. Math.* **61**:6 (2009), 1375–1382. MR Zbl
- [Springer 1981] T. A. Springer, *Linear algebraic groups*, Birkhäuser, Boston, 1981. MR Zbl
- [Urban 2011] E. Urban, “Eigenvarieties for reductive groups”, *Ann. of Math. (2)* **174**:3 (2011), 1685–1784. MR Zbl
- [Wan 1998] D. Wan, “Dimension variation of classical and p -adic modular forms”, *Invent. Math.* **133**:2 (1998), 449–463. MR Zbl

Received March 27, 2015. Revised May 20, 2016.

JOACHIM MAHNKOPF
FAKULTÄT FÜR MATHEMATIK
UNIVERSITÄT WIEN
OSKAR-MORGENSTERN-PLATZ 1
1190 WIEN
AUSTRIA
joachim.mahnkopf@univie.ac.at

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 289 No. 2 August 2017

Regular representations of completely bounded maps	257
B. V. RAJARAMA BHAT, NIRUPAMA MALLICK and K. SUMESH	
Ball convex bodies in Minkowski spaces	287
THOMAS JAHN, HORST MARTINI and CHRISTIAN RICHTER	
Local constancy of dimension of slope subspaces of automorphic forms	317
JOACHIM MAHNKOPF	
Weakening idempotency in K -theory	381
VLADIMIR MANUILOV	
On Langlands quotients of the generalized principal series isomorphic to their Aubert duals	395
IVAN MATIĆ	
Exact Lagrangian fillings of Legendrian $(2, n)$ torus links	417
YU PAN	
Elementary calculation of the cohomology rings of real Grassmann manifolds	443
RUSTAM SADYKOV	
Cluster tilting modules and noncommutative projective schemes	449
KENTA UEYAMA	
Concentration for a biharmonic Schrödinger equation	469
DONG WANG	
Global existence of smooth solutions to exponential wave maps in FLRW spacetimes	489
CHANG-HUA WEI and NING-AN LAI	



0030-8730(201708)289:2;1-O