

*Pacific
Journal of
Mathematics*

**ON LANGLANDS QUOTIENTS
OF THE GENERALIZED PRINCIPAL SERIES
ISOMORPHIC TO THEIR AUBERT DUALS**

IVAN MATIĆ

ON LANGLANDS QUOTIENTS OF THE GENERALIZED PRINCIPAL SERIES ISOMORPHIC TO THEIR AUBERT DUALS

IVAN MATIĆ

We determine under which conditions is the Langlands quotient of an induced representation of the form $\delta \rtimes \sigma$, where δ is an irreducible essentially square-integrable representation of a general linear group and σ is a discrete series representation of the classical p -adic group, isomorphic to its Aubert dual.

1. Introduction

Let F denote a nonarchimedean local field and let G_n stand for the symplectic or (full) orthogonal group having split rank n . The involution on the Grothendieck group of the smooth finite-length representations of a reductive group has been intensively studied by many authors, and we use an involution defined for general reductive p -adic groups in [Aubert 1995] and [Schneider and Stuhler 1997]. This involution is known as the Aubert involution and the image of a representation under this involution is called the Aubert dual of a representation. In this paper we regard the Aubert dual of an admissible finite-length representation as a genuine representation, taking the $+$ or $-$ sign in such way that we obtain the positive element in the appropriate Grothendieck group.

The Aubert involution has a number of prominent applications in the representation theory of classical p -adic groups, and one would also like to gain a deeper knowledge on the explicit structure of the Aubert duals of irreducible representations.

In our previous work [Matić 2016a; 2017], we obtained an explicit description of the Aubert duals of certain classes of discrete series representations of G_n , and in this paper we use developed methods to identify certain classes of irreducible representations which are fixed by the Aubert involution, i.e., which are isomorphic to their Aubert duals. We tackle this problem for the Langlands quotients of the generalized principal series of the group G_n . We note that the generalized principal series is an induced representation of the form $\delta \rtimes \sigma$, obtained by the parabolic

MSC2010: primary 22E35; secondary 11F70, 22E50.

Keywords: discrete series, classical p -adic groups, Aubert involution.

induction with respect to the maximal parabolic subgroup, where the inducing representation $\delta \otimes \sigma$ has an irreducible essentially square-integrable representation on the general linear group part and an irreducible square-integrable representation on the classical group part. If $\nu^x \delta$ is unitarizable for $x < 0$, where $\nu = |\det|_F$, then the generalized principal series $\delta \rtimes \sigma$ has a unique irreducible (Langlands) quotient, which is also isomorphic to the unique irreducible subrepresentation of $\tilde{\delta} \rtimes \sigma$. Such irreducible nontempered representations can be observed as the first step in the Langlands classification of the nonunitary dual of G_n .

To obtain the necessary conditions under which the Langlands quotient of the generalized principal series is isomorphic to its Aubert dual, we use the Jacquet modules method and some elementary properties of the Aubert involution, together with descriptions of the Jacquet modules of certain discrete series representations, obtained in [Matić 2013; 2016c].

Afterwards, we explicitly determine the Aubert duals of Langlands quotients satisfying the obtained necessary conditions, using methods introduced in [Matić 2017], and further developed in [Matić 2016a]. Perhaps a bit surprisingly, an important role in such a procedure is, in the considered case, played by the composition factors of the generalized principal series $\delta \rtimes \sigma$ with a strongly positive σ , obtained in [Muić 2004] and [Matić 2016b, Proposition 3.2]. Such a description of the composition factors enables us to control the Jacquet modules of the investigated nontempered representations, similarly to in [Matić 2015].

We summarize our main results in the following theorem.

Theorem 1.1. *The Langlands quotient of the generalized principal series*

$$\delta([\nu^x \rho, \nu^y \rho]) \rtimes \sigma, \quad x + y > 0,$$

is isomorphic to its Aubert dual if and only if one of the following holds:

- (1) *The discrete series representation σ is cuspidal, $x = y$, and $\nu^x \rho \rtimes \sigma$ is irreducible.*
- (2) *The discrete series representation σ is cuspidal, $x = 0$, $y = 1$, and $\rho \rtimes \sigma$ reduces.*
- (3) *The cuspidal representation ρ is self-contragredient, the induced representation $\nu^\alpha \rho \rtimes \sigma_{\text{cusp}}$ reduces for $\alpha > 0$ (here σ_{cusp} stands for the partial cuspidal support of σ), $y = \alpha + 1$, and one of the following holds:*
 - (i) *x is a half-integer, $\frac{3}{2} \leq x \leq \alpha$, and σ is the unique irreducible subrepresentation of the induced representation*

$$\nu^x \rho \times \nu^{x+1} \rho \times \cdots \times \nu^\alpha \rho \rtimes \sigma_{\text{cusp}}.$$

- (ii) x is a positive integer, $x \leq \alpha$, and σ is the unique irreducible subrepresentation of the induced representation

$$v^x \rho \times v^{x+1} \rho \times \cdots \times v^\alpha \rho \rtimes \sigma_{\text{cusp}}.$$

- (iii) $x = 0$ and σ is the unique irreducible subrepresentation of the induced representation

$$v \rho \times v^2 \rho \times \cdots \times v^\alpha \rho \rtimes \sigma_{\text{cusp}}.$$

We now describe the contents of the paper in more detail. In [Section 2](#) we set up the notation and terminology, and prove some technical results which will be helpful in our investigation. In [Section 3](#) we state and prove our main results, using a case-by-case consideration.

2. Notation and preliminaries

Let F denote a nonarchimedean local field of characteristic $\neq 2$.

Let us first recall the definition of the Aubert involution and its basic properties.

For a connected reductive p -adic group G defined over F , let Σ denote the set of roots of G with respect to a fixed minimal parabolic subgroup and let Δ stand for a basis of Σ . For $\Theta \subseteq \Delta$, we let P_Θ be the standard parabolic subgroup of G corresponding to Θ and M_Θ the standard Levi factor of G corresponding to Θ .

For a parabolic subgroup P of G with the Levi factor M and a representation σ of M , we denote by $i_M(\sigma)$ a normalized parabolically induced representation of G_n induced from σ . For an admissible finite-length representation σ of G , the normalized Jacquet module of σ with respect to the standard parabolic subgroup having Levi factor equal to M will be denoted by $r_M(\sigma)$. We recall the following definition and results from [\[Aubert 1995\]](#):

Theorem 2.1. *Define the operator on the Grothendieck group of admissible representations of G of finite length by*

$$D_G = \sum_{\Theta \subseteq \Delta} (-1)^{|\Theta|} i_{M_\Theta} \circ r_{M_\Theta}.$$

The operator D_G has the following properties:

- (1) D_G is an involution.
- (2) D_G takes irreducible representations to irreducible ones.
- (3) If σ is an irreducible cuspidal representation, then $D_G(\sigma) = (-1)^{|\Delta|} \sigma$.
- (4) For a standard Levi subgroup $M = M_\Theta$, we have

$$D_G \circ i_M = i_M \circ D_M.$$

(5) For a standard Levi subgroup $M = M_\Theta$, we have

$$r_M \circ D_G = \text{Ad}(w) \circ D_{w^{-1}(M)} \circ r_{w^{-1}(M)},$$

where w is the longest element of the set $\{w \in W : w^{-1}(\Theta) > 0\}$.

Let us now describe the groups that we consider. We look at the usual towers of orthogonal or symplectic groups $G_n = G(V_n)$ that are the groups of isometries of F -spaces $(V_n, (\cdot, \cdot))$, $n \geq 0$, where the form (\cdot, \cdot) is nondegenerate and it is skew-symmetric if the tower is symplectic and symmetric otherwise. The set of standard parabolic subgroups will be fixed in a usual way. Then the Levi factors of standard parabolic subgroups have the form

$$M \cong \text{GL}(n_1, F) \times \text{GL}(n_2, F) \times \cdots \times \text{GL}(n_k, F) \times G_{n'}.$$

If δ_i is a representation of $\text{GL}(n_i, F)$ for $i = 1, 2, \dots, k$, and τ a representation of $G_{n'}$, the induced representation $i_M(\delta_1 \otimes \delta_2 \otimes \cdots \otimes \delta_k \otimes \tau)$ will be denoted by $\delta_1 \times \delta_2 \times \cdots \times \delta_k \rtimes \tau$. We use a similar notation to denote a parabolically induced representation of $\text{GL}(m, F)$.

If π is an irreducible representation of G_n , we denote by $\hat{\pi}$ the representation $\pm D_{G_n}(\pi)$, taking the sign $+$ or $-$ such that $\hat{\pi}$ is a positive element in the Grothendieck group of finite-length admissible representations of G_n . We call $\hat{\pi}$ the Aubert dual of π .

By $\text{Irr}(G_n)$ we denote the set of all irreducible admissible representations of G_n . Furthermore, let $R(G_n)$ denote the Grothendieck group of admissible representations of finite length of G_n and define $R(G) = \bigoplus_{n \geq 0} R(G_n)$. Similarly, let $\text{Irr}(\text{GL}(n, F))$ denote the set of all irreducible admissible representations of $\text{GL}(n, F)$, let $R(\text{GL}(n, F))$ denote the Grothendieck group of admissible representations of finite length of $\text{GL}(n, F)$ and define $R(\text{GL}) = \bigoplus_{n \geq 0} R(\text{GL}(n, F))$.

The generalized principal series are the induced representations of the form $\delta \rtimes \sigma$, where $\delta \in R(\text{GL})$ is an irreducible essentially square-integrable representation and $\sigma \in R(G)$ is a discrete series representation.

There is a unique $e(\delta) \in \mathbb{R}$ such that $v^{-e(\delta)}\delta$ is unitarizable, where $v = |\text{det}|_F$. If $e(\delta) > 0$, the generalized principal series $\delta \rtimes \sigma$ has a unique irreducible (Langlands) quotient, which is also the unique irreducible subrepresentation of $\tilde{\delta} \rtimes \sigma$, where $\tilde{\delta}$ denotes the contragredient of δ .

By the results of [Zelevinsky 1980], such a representation δ is attached to the segment and we write $\delta = \delta([v^a\rho, v^b\rho])$, where $a, b \in \mathbb{R}$ are such that $b - a$ is a nonnegative integer and $\rho \in \text{Irr}(\text{GL}(n, F))$ is a unitary cuspidal representation. We recall that $\delta([v^a\rho, v^b\rho])$ is the unique irreducible subrepresentation of the induced representation $v^b\rho \times v^{b-1}\rho \times \cdots \times v^a\rho$.

For our Jacquet module considerations it is more convenient to use the subrepresentation version of the Langlands classification and write a nontempered irreducible representation π of G_n as the unique irreducible (Langlands) subrepresentation of the induced representation of the form $\delta_1 \times \delta_2 \times \cdots \times \delta_k \rtimes \tau$, where $\tau \in \text{Irr}(G_m)$ is a tempered representation, $\delta_i \in \text{Irr}(\text{GL}(n_i, F))$ is an essentially square-integrable representation attached to the segment $[v^{a_i}\rho_i, v^{b_i}\rho_i]$ for $i = 1, 2, \dots, k$, and $a_1 + b_1 \leq a_2 + b_2 \leq \cdots \leq a_k + b_k < 0$. In this case, we write $\pi = L(\delta_1 \times \delta_2 \times \cdots \times \delta_k \rtimes \tau)$.

For $\sigma \in \text{Irr}(G_n)$ and $1 \leq k \leq n$, denote by $r_{(k)}(\sigma)$ the normalized Jacquet module of σ with respect to the parabolic subgroup with Levi factor $\text{GL}(k, F) \times G_{n-k}$. Identify $r_{(k)}(\sigma)$ with its semisimplification in $R(\text{GL}(k, F)) \otimes R(G_{n-k})$ and consider

$$\mu^*(\sigma) = 1 \otimes \sigma + \sum_{k=1}^n r_{(k)}(\sigma) \in R(\text{GL}) \otimes R(G).$$

The following result, derived in [Tadić 1995], presents the crucial structural formula for our calculations of Jacquet modules.

Theorem 2.2. *Let ρ be an irreducible cuspidal representation of $\text{GL}(m, F)$ and $k, l \in \mathbb{R}$ such that $k + l \in \mathbb{Z}_{\geq 0}$. Let σ be an admissible representation of G_n of finite length. Write $\mu^*(\sigma) = \sum_{\tau, \sigma'} \tau \otimes \sigma'$. Then we have*

$$\begin{aligned} & \mu^*(\delta([v^{-k}\rho, v^l\rho]) \rtimes \sigma) \\ &= \sum_{i=-k-1}^l \sum_{j=i}^l \sum_{\tau, \sigma'} \delta([v^{-i}\tilde{\rho}, v^k\tilde{\rho}]) \times \delta([v^{j+1}\rho, v^l\rho]) \times \tau \otimes \delta([v^{i+1}\rho, v^j\rho]) \rtimes \sigma'. \end{aligned}$$

We omit $\delta([v^x\rho, v^y\rho])$ if $x > y$.

We note the following direct consequence of the previous theorem and of the Casselman square-integrability criterion:

Corollary 2.3. *Let ρ denote an irreducible self-contragredient cuspidal representation of $\text{GL}(m, F)$ and $k, l \in \mathbb{R}$ such that $k + l \in \mathbb{Z}_{\geq 0}$ and $k > 0$. If $\sigma \in \text{Irr}(G_n)$ is a discrete series representation, then $\mu^*(\delta([v^{-k}\rho, v^l\rho]) \rtimes \sigma)$ contains an irreducible constituent of the form $v^r\rho' \otimes \pi$, where $r \leq 0$ and ρ' is cuspidal, if and only if $l \leq 0$. Furthermore, if $l \leq 0$ and $\mu^*(\delta([v^{-k}\rho, v^l\rho]) \rtimes \sigma)$ contains an irreducible constituent of the form $v^r\rho' \otimes \pi$, where $r \leq 0$ and ρ' is cuspidal, then $r = l$ and $\rho' \cong \rho$.*

The following technical result will be used several times in the paper:

Lemma 2.4. *Suppose that $\pi \in \text{Irr}(G_n)$ is a subrepresentation of an induced representation of the form $v^{a_1}\rho_1 \times v^{a_2}\rho_2 \times \cdots \times v^{a_k}\rho_k \rtimes \pi_1$, where $\rho_i \in \text{Irr}(\text{GL}(m_i, F))$ is a unitary cuspidal self-contragredient representation for $i = 1, 2, \dots, k$, and π_1 is an admissible representation of finite length. Then the Jacquet module of $\hat{\pi}$ with*

respect to an appropriate parabolic subgroup contains an irreducible representation of the form $v^{-a_1}\rho_1 \otimes v^{-a_2}\rho_2 \otimes \cdots \otimes v^{-a_k}\rho_k \otimes \pi_2$.

Proof. Frobenius reciprocity and transitivity of Jacquet modules imply that there is an irreducible cuspidal representation π' such that the Jacquet module of π with respect to an appropriate parabolic subgroup contains the irreducible cuspidal representation $v^{a_1}\rho_1 \otimes v^{a_2}\rho_2 \otimes \cdots \otimes v^{a_k}\rho_k \otimes \pi'$. Using [Theorem 2.1](#), we obtain the claim of the lemma. \square

We note that one can also deduce that the representation π_2 in the previous lemma is in fact isomorphic to $\widehat{\pi}_1$. However, we will not use this in the sequel.

We will now recall the Mœglin–Tadić classification of discrete series for groups that we consider. Every discrete series representation of G_n is uniquely determined by three invariants: the partial cuspidal support, the Jordan block and the ϵ function.

The partial cuspidal support of a discrete series $\sigma \in \text{Irr}(G_n)$ is an irreducible cuspidal representation σ_{cusp} of some G_m such that there is an irreducible admissible representation π of $\text{GL}(n-m, F)$ such that σ is a subrepresentation of $\pi \rtimes \sigma_{\text{cusp}}$.

The Jordan block of σ , denoted by $\text{Jord}(\sigma)$, is the set of all pairs (c, ρ) where ρ is an irreducible cuspidal self-contragredient representation of some $\text{GL}(n_\rho, F)$ and $c > 0$ is an integer such that the following two conditions are satisfied:

- (1) c is even if and only if $L(s, \rho, r)$ has a pole at $s = 0$. The local L -function $L(s, \rho, r)$ is the one defined by Shahidi [[1990](#); [1992](#)], where $r = \bigwedge^2 \mathbb{C}^{n_\rho}$ is the exterior-square representation of the standard representation on \mathbb{C}^{n_ρ} of $\text{GL}(n_\rho, \mathbb{C})$ if G_n is a symplectic or even-orthogonal group, and $r = \text{Sym}^2 \mathbb{C}^{n_\rho}$ is the symmetric-square representation of the standard representation on \mathbb{C}^{n_ρ} of $\text{GL}(n_\rho, \mathbb{C})$ if G_n is an odd-orthogonal group.
- (2) The induced representation $\delta([v^{-(c-1)/2}\rho, v^{(c-1)/2}\rho]) \rtimes \sigma$ is irreducible.

To explain the notion of the ϵ function, we will first define Jordan triples. These are triples of the form $(\text{Jord}, \sigma', \epsilon)$, where

- σ' is an irreducible cuspidal representation of some G_n .
- Jord is the finite (possibly empty) set of ordered pairs (c, ρ) , where $\rho \in \text{Irr}(\text{GL}(n_\rho, F))$ is a self-contragredient cuspidal representation, and c is a positive integer which is even if and only if $L(s, \rho, r)$ has a pole at $s = 0$ (for the local L function as above). For an irreducible self-contragredient cuspidal representation ρ of $\text{GL}(n_\rho, F)$ we write $\text{Jord}_\rho = \{c : (c, \rho) \in \text{Jord}\}$. If $\text{Jord}_\rho \neq \emptyset$ and $c \in \text{Jord}_\rho$, we put $c_- = \max\{d \in \text{Jord}_\rho : d < c\}$, if it exists.
- ϵ is the function defined on a subset of $\text{Jord} \cup (\text{Jord} \times \text{Jord})$ and attains the values 1 and -1 . If $(c, \rho) \in \text{Jord}$, then $\epsilon(c, \rho)$ is not defined if and only if c is odd and $(c', \rho) \in \text{Jord}(\sigma')$ for some positive integer c' . Next, ϵ is defined on a pair $((c, \rho), (c', \rho')) \in \text{Jord} \times \text{Jord}$ if and only if $\rho \cong \rho'$ and $c \neq c'$.

Suppose that, for the Jordan triple $(\text{Jord}, \sigma', \epsilon)$, there is a $(c, \rho) \in \text{Jord}$ such that $\epsilon((c_-, \rho), (c, \rho)) = 1$. If we put $\text{Jord}' = \text{Jord} \setminus \{(c_-, \rho), (c, \rho)\}$ and consider the restriction ϵ' of ϵ to $\text{Jord}' \cup (\text{Jord}' \times \text{Jord}')$, we obtain a new Jordan triple $(\text{Jord}', \sigma', \epsilon')$, and we say that such Jordan triple is subordinated to $(\text{Jord}, \sigma', \epsilon)$.

We say that the Jordan triple $(\text{Jord}, \sigma', \epsilon)$ is a triple of alternated type if

$$\epsilon((c_-, \rho), (c, \rho)) = -1$$

whenever c_- is defined and there is an increasing bijection $\phi_\rho : \text{Jord}_\rho \rightarrow \text{Jord}'_\rho(\sigma')$, where $\text{Jord}'_\rho(\sigma')$ equals $\text{Jord}_\rho(\sigma') \cup \{0\}$ if a is even and $\epsilon(\min \text{Jord}_\rho, \rho) = 1$, and $\text{Jord}'_\rho(\sigma')$ equals $\text{Jord}_\rho(\sigma')$ otherwise.

The Jordan triple $(\text{Jord}, \sigma', \epsilon)$ dominates the Jordan triple $(\text{Jord}', \sigma', \epsilon')$ if there is a sequence of Jordan triples $(\text{Jord}_i, \sigma', \epsilon_i)$, $0 \leq i \leq k$, such that $(\text{Jord}_0, \sigma', \epsilon_0) = (\text{Jord}, \sigma', \epsilon)$, $(\text{Jord}_k, \sigma', \epsilon_k) = (\text{Jord}', \sigma', \epsilon')$, and $(\text{Jord}_i, \sigma', \epsilon_i)$ is subordinated to $(\text{Jord}_{i-1}, \sigma', \epsilon_{i-1})$ for $i \in \{1, 2, \dots, k\}$. The Jordan triple $(\text{Jord}, \sigma', \epsilon)$ is called admissible if it dominates a triple of alternated type.

The classification given in [Mœglin 2002] and [Mœglin and Tadić 2002] states that there is a one-to-one correspondence between the set of all discrete series in $\text{Irr}(G)$ and the set of all admissible triples $(\text{Jord}, \sigma', \epsilon)$ given by $\sigma = \sigma_{(\text{Jord}, \sigma', \epsilon)}$, such that $\sigma_{\text{cusps}} = \sigma'$ and $\text{Jord}(\sigma) = \text{Jord}$. Furthermore, if $(c, \rho) \in \text{Jord}$ is such that $\epsilon((c_-, \rho), (c, \rho)) = 1$, we set $\text{Jord}' = \text{Jord} \setminus \{(c_-, \rho), (c, \rho)\}$ and consider the restriction ϵ' of ϵ to $\text{Jord}' \cup (\text{Jord}' \times \text{Jord}')$. Then $(\text{Jord}', \sigma', \epsilon')$ is an admissible triple and σ is a subrepresentation of $\delta([v^{-(c_- - 1)/2}\rho, v^{(c_- - 1)/2}\rho]) \rtimes \sigma_{(\text{Jord}', \sigma', \epsilon')}$.

An irreducible representation $\sigma \in R(G)$ is called strongly positive if for every embedding

$$\sigma \hookrightarrow v^{s_1}\rho_1 \times v^{s_2}\rho_2 \times \cdots \times v^{s_k}\rho_k \rtimes \sigma_{\text{cusps}},$$

where $\rho_i \in R$, $i = 1, 2, \dots, k$, are irreducible cuspidal unitary representations and $\sigma_{\text{cusps}} \in R(G)$ is an irreducible cuspidal representation, we have $s_i > 0$ for $i = 1, 2, \dots, k$.

It was shown in [Mœglin 2002, Proposition 5.3] and [Mœglin and Tadić 2002, Proposition 7.1] that triples of alternated type correspond to strongly positive discrete series. Let us recall an inductive description of the noncuspidal strongly positive discrete series, obtained in [Matić 2011, Theorem 5.1], which also holds in the classical group case.

Proposition 2.5. *Suppose that $\sigma_{\text{sp}} \in R(G)$ is an irreducible strongly positive representation and let $\rho \in \text{Irr}(\text{GL}(m, F))$ denote an irreducible cuspidal representation such that some twist of ρ appears in the cuspidal support of σ_{sp} . We denote by σ_{cusps} the partial cuspidal support of σ_{sp} . Then there exist unique $a, b \in \mathbb{R}$ such that $a > 0$, $b > 0$, $b - a$ is a nonnegative integer, and a unique irreducible strongly positive representation $\sigma_{\text{sp}}^{(1)}$ without $v^a\rho$ in the cuspidal support, with the property that σ_{sp}*

is the unique irreducible subrepresentation of $\delta([v^a\rho, v^b\rho]) \rtimes \sigma_{\text{sp}}^{(1)}$. Furthermore, there is a nonnegative integer l such that $\alpha := a + l > 0$ and $v^\alpha\rho \rtimes \sigma_{\text{cusp}}$ reduces. If $l = 0$ there are no twists of ρ appearing in the cuspidal support of $\sigma_{\text{sp}}^{(1)}$, and if $l > 0$ there exist a unique $b' > b$ and a unique strongly positive discrete series $\sigma_{\text{sp}}^{(2)}$, which contains neither $v^a\rho$ nor $v^{a+1}\rho$ in its cuspidal support, such that $\sigma_{\text{sp}}^{(1)}$ can be written as the unique irreducible subrepresentation of $\delta([v^{a+1}\rho, v^{b'}\rho]) \rtimes \sigma_{\text{sp}}^{(2)}$.

We say a representation $\sigma \in \text{Irr}(G_n)$ belongs to the set $D(\rho_1, \dots, \rho_m; \sigma_{\text{cusp}})$ if every element of the cuspidal support of σ belongs to the set $\{v^x\rho_1, \dots, v^x\rho_m, \sigma_{\text{cusp}} : x \in \mathbb{R}\}$, where ρ_1, \dots, ρ_m are mutually nonisomorphic irreducible cuspidal representations of general linear groups and σ_{cusp} is an irreducible cuspidal representation of $G_{n'}$ for some $n' \leq n$.

We note that for a self-contragredient cuspidal $\rho \in \text{Irr}(\text{GL}(m, F))$ and a cuspidal $\sigma_{\text{cusp}} \in \text{Irr}(G_n)$, there is a unique nonnegative α such that the induced representation $v^\alpha\rho \rtimes \sigma_{\text{cusp}}$ reduces, and it follows from [Arthur 2013] and [Mœglin 2014, Théorème 3.1.1] that α is a half-integer.

Directly from the previous proposition we obtain

Proposition 2.6. *Let $\sigma_{\text{sp}} \in \text{Irr}(G_n)$ denote a strongly positive representation and suppose that $\sigma_{\text{sp}} \in D(\rho; \sigma_{\text{cusp}})$ for an irreducible cuspidal self-contragredient representation ρ . Let α stand for the unique nonnegative half-integer such that $v^\alpha\rho \rtimes \sigma_{\text{cusp}}$ reduces, and let $k = \lceil \alpha \rceil$, the smallest integer which is not smaller than α . If $k = 0$, then $\sigma_{\text{sp}} \cong \sigma_{\text{cusp}}$. Otherwise, there exists a unique k -tuple (a_1, a_2, \dots, a_k) such that $a_i - \alpha \in \mathbb{Z}$ for $i = 1, 2, \dots, k$, $-1 < a_1 < a_2 < \dots < a_k$, and σ_{sp} is the unique irreducible subrepresentation of the induced representation*

$$\delta([v^{\alpha-k+1}\rho, v^{a_1}\rho]) \times \delta([v^{\alpha-k+2}\rho, v^{a_2}\rho]) \times \dots \times \delta([v^\alpha\rho, v^{a_k}\rho]) \rtimes \sigma_{\text{cusp}}.$$

3. Langlands quotients fixed by the Aubert involution

In this section we describe all Langlands quotients of the generalized principal series $\delta \rtimes \sigma$ which are isomorphic to their Aubert duals, using case-by-case considerations. We write $\delta = \delta([v^x\rho, v^y\rho])$, for x, y such that $x + y > 0$. The induced representation $\delta \rtimes \sigma$ then contains a unique irreducible (Langlands) quotient, which is also the unique irreducible subrepresentation of the induced representation $\delta([v^{-y}\tilde{\rho}, v^{-x}\tilde{\rho}]) \rtimes \sigma$, and in what follows will be denoted by π , i.e., let $\pi = L(\tilde{\delta} \rtimes \sigma)$.

Lemma 3.1. *If π is isomorphic to $\hat{\pi}$, then $x \geq 0$.*

Proof. Since $x + y > 0$, we obviously have $y > 0$. Suppose that $x < 0$. From the embedding $\pi \hookrightarrow v^{-x}\tilde{\rho} \rtimes \delta([v^{-y}\tilde{\rho}, v^{-x-1}\tilde{\rho}]) \rtimes \sigma$ and the transitivity of Jacquet modules, in the same way as in the proof of Lemma 2.4 we obtain that $\mu^*(\hat{\pi})$ contains an irreducible constituent of the form $v^x\rho \otimes \pi'$. Since $y \neq x$, it follows

directly from the structural formula that $\mu^*(\delta([v^{-y}\tilde{\rho}, v^{-x}\tilde{\rho}]) \rtimes \sigma)$ does not contain such an irreducible constituent, so $\hat{\pi}$ is not isomorphic to π , a contradiction. \square

Let us first consider the case of cuspidal σ .

Proposition 3.2. *Suppose that $\sigma \in \text{Irr}(G_n)$ is a cuspidal representation. Then π is isomorphic to its Aubert dual if and only if one of the following holds:*

- (1) $x = y > 0$ and the induced representation $v^x\rho \rtimes \sigma$ is irreducible,
- (2) $(x, y) = (0, 1)$ and the induced representation $\rho \rtimes \sigma$ reduces.

Proof. We have already seen that if $\pi \cong \hat{\pi}$ then $x \geq 0$. In the same way as in the proof of the previous lemma we deduce that $\mu^*(\hat{\pi}) \geq v^{-x}\tilde{\rho} \otimes \pi'$, for some irreducible representation π' . From the structural formula we see that this is possible only if either $x = y$ or $(x, \tilde{\rho}) = (0, \rho)$.

Let us first consider the case $x = y$. Note that then we have $x > 0$. Furthermore, if $v^x\rho \rtimes \sigma$ reduces, it follows from [Muić 2004, Proposition 3.1(i)] that $\mu^*(\pi)$ does not contain an irreducible constituent of the form $v^{-x}\tilde{\rho} \otimes \pi'$. Consequently, if $\pi \cong \hat{\pi}$ and $x = y$, then $v^x\rho \rtimes \sigma$ is irreducible.

Conversely, if the induced representation $v^x\rho \rtimes \sigma$ is irreducible, then $\pi \cong v^x\rho \rtimes \sigma$ and from part (4) of Theorem 2.1 we have $\hat{\pi} \leq v^x\rho \rtimes \sigma \cong \pi$, so π is isomorphic to its Aubert dual.

Let us now assume that $x = 0$ and $\rho \cong \tilde{\rho}$. Let s denote a unique nonnegative half-integer such that $v^s\rho \rtimes \sigma$ reduces. We obviously have $y > 0$ and there are two possibilities to consider.

First, suppose that $y = s$ and let σ_{sp} stand for a unique irreducible subrepresentation of the induced representation $v\rho \times v^2\rho \times \cdots \times v^y\rho \rtimes \sigma$. It follows from [Matić 2011, Theorem 4.6] that σ_{sp} is a strongly positive discrete series and one can see directly from [Mœglin and Tadić 2002, Proposition 2.1] that the induced representation $\rho \rtimes \sigma_{\text{sp}}$ reduces. By [Tadić 2013, Section 4], there is a unique irreducible subrepresentation τ of $\rho \rtimes \sigma_{\text{sp}}$ such that $\mu^*(\tau)$ does not contain an irreducible constituent of the form $v\rho \otimes \pi'$. We note that τ is a tempered representation. Furthermore, if $\mu^*(\tau)$ contains an irreducible constituent of the form $v^a\rho \otimes \pi'_1$, then $a = 0$. Thus, if $\mu^*(\hat{\tau})$ contains an irreducible constituent of the form $v^a\rho \otimes \pi'_2$, then $a = 0$. Since τ is a subrepresentation of $\rho \times v\rho \times v^2\rho \times \cdots \times v^y\rho \rtimes \sigma$, using Lemma 2.4 we deduce that $\hat{\tau}$ is a subrepresentation of $\rho \times v^{-1}\rho \times v^{-2}\rho \times \cdots \times v^{-y}\rho \rtimes \sigma$ and, in the same way as in the proof of [Matić 2017, Lemma 3.4], we deduce that $\hat{\tau}$ is a subrepresentation of $\delta([v^{-y}\rho, \rho]) \rtimes \sigma$. Consequently, $\hat{\tau} \cong \pi$ and $\hat{\pi} \not\cong \pi$.

Now, suppose that $y \neq s$. If $y \geq 2$, we have the following embedding and isomorphism:

$$\pi \hookrightarrow \delta([v^{-y+1}\rho, \rho]) \times v^{-y}\rho \rtimes \sigma \cong v^y\rho \times \delta([v^{-y+1}\rho, \rho]) \rtimes \sigma.$$

Lemma 2.4 implies that $\mu^*(\hat{\pi})$ contains an irreducible constituent of the form $v^{-y}\rho \otimes \pi'$ and it follows directly from the structural formula that $\hat{\pi} \not\cong \pi$. Thus, we can assume that $y = 1$. If $s > 0$, then $s \neq y$ and [Muić 2004, Theorem 4.1(i)] imply that $\delta([v^{-1}\rho, \rho]) \rtimes \sigma$ is irreducible and $\pi \cong \delta([\rho, v\rho]) \rtimes \sigma$. Consequently, if $s > 0$ then $\mu^*(\hat{\pi})$ contains an irreducible constituent of the form $v^{-1}\rho \otimes \pi'$, and it follows directly from the structural formula that $\hat{\pi} \not\cong \pi$.

It remains to consider the case $s = 0$. According to [Muić 2004, Theorem 2.1], in $R(G)$ we have

$$\delta([\rho, v\rho]) \rtimes \sigma = \pi + \sigma_{\text{ds}}^{(1)} + \sigma_{\text{ds}}^{(2)},$$

where $\sigma_{\text{ds}}^{(1)}$ and $\sigma_{\text{ds}}^{(2)}$ are mutually nonisomorphic discrete series subrepresentations of $\delta([\rho, v\rho]) \rtimes \sigma$. Frobenius reciprocity implies that both $\mu^*(\sigma_{\text{ds}}^{(1)})$ and $\mu^*(\sigma_{\text{ds}}^{(2)})$ contain irreducible constituents of the form $v\rho \otimes \pi'$. It follows from the structural formula that only irreducible constituents of the form $v\rho \otimes \pi'$ appearing in $\mu^*(\delta([\rho, v\rho]) \rtimes \sigma)$ are $v\rho \otimes \tau_1$ and $v\rho \otimes \tau_{-1}$, where τ_1 and τ_{-1} are irreducible mutually nonisomorphic tempered representations such that in $R(G)$ we have $\rho \rtimes \sigma = \tau_1 + \tau_{-1}$. Furthermore, both $v\rho \otimes \tau_1$ and $v\rho \otimes \tau_{-1}$ appear in $\mu^*(\delta([\rho, v\rho]) \rtimes \sigma)$ with multiplicity one. Thus, $\mu^*(\pi)$ does not contain an irreducible constituent of the form $v\rho \otimes \pi'$ and, consequently, $\mu^*(\hat{\pi})$ does not contain an irreducible constituent of the form $v^{-1}\rho \otimes \pi''$. Since π is a subrepresentation of $\rho \times v\rho \rtimes \sigma$, using **Lemma 2.4** we obtain that $\hat{\pi}$ is a subrepresentation of $\rho \times v^{-1}\rho \rtimes \sigma$ and it follows that $\hat{\pi}$ is a unique irreducible subrepresentation of $\delta([v^{-1}\rho, \rho]) \rtimes \sigma$, i.e., $\pi \cong \hat{\pi}$. This completes the proof. \square

In the rest of this section we assume that σ is a noncuspidal discrete series representation, and let σ_{cusp} denote the partial cuspidal support of σ .

Lemma 3.3. *If π is isomorphic to $\hat{\pi}$, then $\sigma \in D(\rho; \sigma_{\text{cusp}})$. In particular, ρ is self-contragredient.*

Proof. Suppose that $\sigma \notin D(\rho; \sigma_{\text{cusp}})$. Then there is an embedding of the form $\sigma \hookrightarrow v^a\rho' \rtimes \sigma'$ such that $a > 0$, ρ' is an irreducible self-contragredient cuspidal representation which is not isomorphic to ρ , and σ' is irreducible. We have

$$\pi \hookrightarrow \tilde{\delta} \times v^a\rho' \rtimes \sigma' \cong v^a\rho' \times \tilde{\delta} \rtimes \sigma',$$

and **Lemma 2.4**, together with transitivity of Jacquet modules, implies that Jacquet module of $\hat{\pi}$ with respect to an appropriate parabolic subgroup contains an irreducible representation of the form $v^{-a}\rho' \otimes \sigma''$. Since σ is square-integrable, it follows that $\mu^*(\delta \rtimes \sigma)$ does not contain an irreducible constituent of the form $v^{-a}\rho' \otimes \sigma''$. Thus, π is not isomorphic to $\hat{\pi}$, a contradiction. \square

According to the previous lemma, in what follows we can assume that ρ is a self-contragredient representation and that $\sigma \in D(\rho; \sigma_{\text{cusp}})$. We denote by α a unique nonnegative half-integer s such that $v^s \rho \rtimes \sigma_{\text{cusp}}$ reduces.

The following result presents the crucial step towards our description.

Theorem 3.4. *If π is isomorphic to $\hat{\pi}$, then σ is a strongly positive discrete series. In particular, $\alpha > 0$.*

Proof. Suppose on the contrary that σ is not a strongly positive representation and let $(\text{Jord}(\sigma), \sigma_{\text{cusp}}, \epsilon_\sigma)$ denote the corresponding Jordan triple. Since $\sigma \in D(\rho; \sigma_{\text{cusp}})$, there is $c \in \text{Jord}_\rho(\sigma)$ such that c_- is defined and $\epsilon_\sigma((c_-, \rho), (c, \rho)) = 1$. Also, σ is a subrepresentation of an induced representation of the form

$$\delta([v^{-(c-1)/2} \rho, v^{(c-1)/2} \rho]) \rtimes \sigma'$$

for an appropriate discrete series σ' . Using [Mœglin and Tadić 2002, Lemma 3.2], we deduce that σ is a subrepresentation of an induced representation of the form $v^{(c-1)/2} \rho \rtimes \pi_1$, for some irreducible π_1 . Since $(c-1)/2 \geq 1$ and $-x \leq 0$, if $(c, x) \neq (3, 0)$ we obtain an embedding $\pi \hookrightarrow v^{(c-1)/2} \rho \times \delta([v^{-y} \rho, v^{-x} \rho]) \rtimes \pi_1$. Lemma 2.4 implies that $\mu^*(\hat{\pi})$ contains an irreducible constituent of the form $v^{-(c-1)/2} \rho \otimes \pi_2$, and Corollary 2.3 implies that $x = (c-1)/2$.

If $c > 3$, we also have $(c-3)/2 > -(c-1)/2 + 1$, which gives the following embeddings and isomorphisms:

$$\begin{aligned} \pi &\hookrightarrow \delta([v^{-y} \rho, v^{-\frac{c-1}{2}} \rho]) \times \delta([v^{-\frac{c-1}{2}} \rho, v^{\frac{c-1}{2}} \rho]) \rtimes \sigma' \\ &\hookrightarrow \delta([v^{-y} \rho, v^{-\frac{c-1}{2}} \rho]) \times v^{\frac{c-1}{2}} \rho \times v^{\frac{c-3}{2}} \rho \times \delta([v^{-\frac{c-1}{2}} \rho, v^{\frac{c-5}{2}} \rho]) \rtimes \sigma' \\ &\cong v^{\frac{c-1}{2}} \rho \times \delta([v^{-y} \rho, v^{-\frac{c-1}{2}} \rho]) \times v^{\frac{c-3}{2}} \rho \times \delta([v^{-\frac{c-1}{2}} \rho, v^{\frac{c-5}{2}} \rho]) \rtimes \sigma' \\ &\cong v^{\frac{c-1}{2}} \rho \times v^{\frac{c-3}{2}} \rho \times \delta([v^{-y} \rho, v^{-\frac{c-1}{2}} \rho]) \times \delta([v^{-\frac{c-1}{2}} \rho, v^{\frac{c-5}{2}} \rho]) \rtimes \sigma'. \end{aligned}$$

Since $\pi \cong \hat{\pi}$, Lemma 2.4 and the transitivity of Jacquet modules imply that the Jacquet module of π with respect to an appropriate parabolic subgroup contains an irreducible representation of the form $v^{-(c-1)/2} \rho \otimes v^{-(c-3)/2} \rho \otimes \pi_3$.

From $\pi \hookrightarrow \delta([v^{-y} \rho, v^{-(c-1)/2} \rho]) \rtimes \sigma$, using the structural formula recalled in Theorem 2.2, we obtain that $v^{-(c-3)/2} \rho \otimes \pi_3 \leq \mu^*(\delta([v^{-y} \rho, v^{-(c+1)/2} \rho]) \rtimes \sigma)$, which is impossible.

It remains to consider the case $c = 3$. This directly implies that $c_- = 1$ and $x \in \{0, 1\}$. In other words, σ is a subrepresentation of an induced representation of the form $\delta([\rho, v\rho]) \rtimes \sigma'$, where σ' is a discrete series such that 1 and 3 do not appear in $\text{Jord}_\rho(\sigma')$, and it follows that σ' is a strongly positive representation, since otherwise we can apply the same arguments as before to deduce that π is not isomorphic to $\hat{\pi}$.

Let us first assume that $\text{Jord}_\rho(\sigma') \neq \emptyset$. Then, as in [Matić 2011, Section 4] and [Mœglin and Tadić 2002, Proposition 2.1], we see that there is an $a \geq 2$ such that σ' is a subrepresentation of $v^a \rho \rtimes \sigma''$ for an appropriate strongly positive representation σ'' .

If $a > 2$, we have $\sigma \hookrightarrow v^a \rho \times \delta([\rho, v\rho]) \rtimes \sigma''$. Since $x \in \{0, 1\}$, in the same way as before we deduce that $\mu^*(\pi) \geq v^{-a} \rho \otimes \pi'$ for an irreducible representation π' , which is impossible.

If $a = 2$, the irreducible representation σ' is also a subrepresentation of an induced representation of the form $v^2 \rho \times v\rho \rtimes \sigma_1$. If $x = 0$, we have the following embeddings:

$$\begin{aligned} \pi &\hookrightarrow \delta([v^{-y}\rho, \rho]) \times \delta([\rho, v\rho]) \times v^2 \rho \times v\rho \rtimes \sigma_1 \\ &\hookrightarrow \rho \times v\rho \times v^2 \rho \times \delta([v^{-y}\rho, v^{-1}\rho]) \times \rho \times v\rho \rtimes \sigma_1 \\ &\hookrightarrow \rho \times v\rho \times v^2 \rho \times v^{-1}\rho \times \rho \times v\rho \times \delta([v^{-y}\rho, v^{-2}\rho]) \rtimes \sigma_1. \end{aligned}$$

Using Lemma 2.4 and the transitivity of Jacquet modules, we obtain that the Jacquet module of π with respect to an appropriate parabolic subgroup contains an irreducible representation of the form $\rho \otimes v^{-1}\rho \otimes v^{-2}\rho \otimes v\rho \otimes \rho \otimes v^{-1}\rho \otimes \pi_4$. Since σ is a discrete series representation, applying the structural formula several times, we deduce that $y \geq 2$ and that $v\rho \otimes \rho \otimes v^{-1}\rho \otimes \pi_4$ is contained in the Jacquet module of $\delta([v^{-y}\rho, v^{-3}\rho]) \rtimes \sigma$ with respect to an appropriate parabolic subgroup. This directly implies that the Jacquet module of σ with respect to an appropriate parabolic subgroup contains an irreducible representation of the form $v\rho \otimes \rho \otimes v^{-1}\rho \otimes \pi_5$, contradicting the square-integrability criterion. The case $x = 1$ can be handled in the same way.

Let us now assume that $\text{Jord}_\rho(\sigma') = \emptyset$. This implies that σ' is a cuspidal representation and $\rho \rtimes \sigma'$ reduces. As in the proof of Proposition 3.2, in $R(G)$ we have $\rho \rtimes \sigma' = \tau_1 + \tau_{-1}$ and there is a unique $i \in \{1, -1\}$ such that σ is the unique irreducible subrepresentation of $v\rho \rtimes \tau_i$ or, equivalently, such that $\mu^*(\sigma) \geq v\rho \otimes \tau_i$. It follows from [Matić 2016a, Theorem 5.1] that $\hat{\sigma} \cong L(v^{-1}\rho \rtimes \tau_{-i})$, and if an irreducible constituent of the form $v^z \rho \otimes \pi'$ appears in $\mu^*(\hat{\sigma})$, then $z = -1$.

Again, we comment only on the case $x = 0$, since the case $x = 1$ can be handled in the same way.

We have the following embeddings:

$$\begin{aligned} (1) \quad \pi &\hookrightarrow \delta([v^{-y}\rho, \rho]) \rtimes \sigma \\ &\hookrightarrow \rho \times v^{-1}\rho \times \cdots \times v^{-y}\rho \rtimes \sigma \\ &\hookrightarrow \rho \times v^{-1}\rho \times \cdots \times v^{-y}\rho \times v\rho \rtimes \tau_i. \end{aligned}$$

Frobenius reciprocity implies that the Jacquet module of π with respect to an appropriate parabolic subgroup contains the irreducible representation

$$(2) \quad \rho \otimes v^{-1}\rho \otimes \cdots \otimes v^{-y}\rho \otimes v\rho \otimes \tau_i.$$

Since $\pi \cong \hat{\pi}$, applying [Theorem 2.1](#), part (4) to the induced representation appearing in (1), we deduce that π is an irreducible subquotient of

$$(3) \quad \rho \times v^{-1}\rho \times \cdots \times v^{-y}\rho \rtimes L(v^{-1}\rho \rtimes \tau_{-i}).$$

Using a repeated application of the structural formula and $\tau_i \not\cong \tau_{-i}$, one can show that the irreducible representation (2) is not contained in the Jacquet module of the induced representation (3) with respect to an appropriate parabolic subgroup, a contradiction. This completes the proof. \square

We denote $\lceil \alpha \rceil$ by k , and let (a_1, a_2, \dots, a_k) denote a unique ordered k -tuple such that $a_i - \alpha \in \mathbb{Z}$ for $i = 1, 2, \dots, k$, $-1 < a_1 < a_2 < \cdots < a_k$, and such that σ is the unique irreducible subrepresentation of

$$\delta([v^{\alpha-k+1}\rho, v^{a_1}\rho]) \times \delta([v^{\alpha-k+2}\rho, v^{a_2}\rho]) \times \cdots \times \delta([v^\alpha\rho, v^{a_k}\rho]) \rtimes \sigma_{\text{cuspidal}}.$$

We note that such a k -tuple exists by [Proposition 2.6](#). Since σ is noncuspidal, there is an $i \in \{1, 2, \dots, k\}$ such that $a_i \geq \alpha - k + i$. Denote the minimal such i by i_{\min} .

Lemma 3.5. *If π is isomorphic to $\hat{\pi}$, then $a_{i_{\min}} = \alpha - k + i_{\min}$ and $a_{j+1} = a_j + 1$ for $j = i_{\min}, i_{\min} + 1, \dots, k - 1$.*

Proof. It follows from [[Matić 2013](#), Theorem 4.6], or from [[Matić and Tadić 2015](#), Section 8], that σ is a subrepresentation of an induced representation of the form $v^{a_{i_{\min}}}\rho \rtimes \sigma_{\text{sp}}$, where σ_{sp} is a strongly positive representation. If $-x \neq a_{i_{\min}} - 1$, we have an embedding $\pi \hookrightarrow v^{a_{i_{\min}}}\rho \times \delta([v^{-y}\rho, v^{-x}\rho]) \rtimes \sigma_{\text{sp}}$, and an application of [Corollary 2.3](#) and [Lemma 2.4](#) gives $x = a_{i_{\min}}$.

If $-x = a_{i_{\min}} - 1$, since $x \geq 0$ and $a_{i_{\min}} \geq \frac{1}{2}$, it follows that $x \in \{0, \frac{1}{2}\}$. Thus, if $-x = a_{i_{\min}} - 1$ then $i_{\min} = 1$ and $a_{i_{\min}} = \alpha - k + 1$.

Let us now assume that $-x \neq a_{i_{\min}} - 1$ and $a_{i_{\min}} \geq \alpha - k + i_{\min} + 1$. It follows that $a_{i_{\min}} \geq \frac{3}{2}$, so $x = a_{i_{\min}}$ and $-x < a_{i_{\min}} - 1$. There is a strongly positive representation σ'_{sp} such that σ is a subrepresentation of $v^{a_{i_{\min}}}\rho \times v^{a_{i_{\min}}-1}\rho \rtimes \sigma'_{\text{sp}}$, so we have an embedding $\pi \hookrightarrow v^{a_{i_{\min}}}\rho \times v^{a_{i_{\min}}-1}\rho \times \delta([v^{-y}\rho, v^{-x}\rho]) \rtimes \sigma'_{\text{sp}}$. Since $\pi \cong \hat{\pi}$, it follows that the Jacquet module of π with respect to an appropriate parabolic subgroup contains an irreducible representation of the form $v^{-a_{i_{\min}}}\rho \otimes v^{-a_{i_{\min}}+1}\rho \otimes \pi'$. From the structural formula we directly obtain that $\mu^*(\delta([v^{-y}\rho, v^{-a_{i_{\min}}-1}\rho]) \rtimes \sigma)$ contains $v^{-a_{i_{\min}}+1}\rho \otimes \pi'$, a contradiction. Consequently, $a_{i_{\min}} = \alpha - k + i_{\min}$.

Now we assume that there is a $j \in \{i_{\min}, i_{\min} + 1, \dots, k - 1\}$ such that $a_{j+1} \neq a_j + 1$. It follows from [[Matić 2011](#), Section 4] that $a_{j+1} \geq a_j + 2$ and π is a subrepresentation of an induced representation of the form $v^{a_{j+1}}\rho \rtimes \sigma'_{\text{sp}}$, for a strongly

positive representation σ'_{sp} . Since we obviously have that $a_{j+1} > -x + 1$, there is an embedding $\pi \hookrightarrow v^{a_{j+1}}\rho \times \delta([v^{-y}\rho, v^{-x}\rho]) \rtimes \sigma'_{\text{sp}}$. This gives $\mu^*(\pi) \geq v^{-a_{j+1}}\rho \otimes \pi'$ for some irreducible π' , contradicting [Corollary 2.3](#), since $x < a_{j+1}$. \square

In the following theorem we state our first main result.

Theorem 3.6. *Suppose that α is a half-integer. Then π is isomorphic to $\hat{\pi}$ if and only if $\alpha \geq \frac{3}{2}$, $a_{i_{\min}} \geq \frac{3}{2}$, $x = a_{i_{\min}}$, and $y = \alpha + 1$.*

[Theorem 3.6](#) follows from the following two propositions:

Proposition 3.7. *Suppose that α is a half-integer and $\pi \cong \hat{\pi}$. Then $\alpha \geq \frac{3}{2}$, $a_{i_{\min}} \geq \frac{3}{2}$, $x = a_{i_{\min}}$, and $y = \alpha + 1$.*

Proof. Let us first show that $x = a_{i_{\min}}$. We have an embedding $\sigma \hookrightarrow v^{a_{i_{\min}}}\rho \rtimes \sigma_{\text{sp}}$ for some strongly positive representation σ_{sp} . Since $x \geq 0$ and $a_{i_{\min}} > 0$, if $(x, a_{i_{\min}}) \neq (\frac{1}{2}, \frac{1}{2})$, we obtain an embedding $\pi \hookrightarrow v^{a_{i_{\min}}}\rho \times \delta([v^{-y}\rho, v^{-x}\rho]) \rtimes \sigma_{\text{sp}}$, which implies that $\mu^*(\pi)$ contains an irreducible constituent of the form $v^{-a_{i_{\min}}}\rho \otimes \pi_1$, since $\pi \cong \hat{\pi}$. This is possible only if $x = a_{i_{\min}}$. Thus, in any case we have $x = a_{i_{\min}}$.

Let us now prove that $a_{i_{\min}} \geq \frac{3}{2}$. Assume on the contrary that $a_{i_{\min}} = \frac{1}{2}$. Using [Lemma 3.5](#) and [[Muić 2004](#), Theorem 5.1], we obtain that $v^z\rho \rtimes \sigma$ is irreducible for $z \notin \{\frac{1}{2}, \alpha + 1\}$. If $y \notin \{\frac{1}{2}, \alpha + 1\}$, we have the following embedding and isomorphism:

$$\pi \hookrightarrow \delta([v^{-y+1}\rho, v^{-1/2}\rho]) \times v^{-y}\rho \rtimes \sigma \cong v^y\rho \times \delta([v^{-y+1}\rho, v^{-1/2}\rho]) \rtimes \sigma.$$

In the same way as before we conclude that $\mu^*(\pi)$ contains an irreducible constituent of the form $v^{-y}\rho \otimes \pi_1$, which is impossible unless $y = \frac{1}{2}$. Thus, $y \in \{\frac{1}{2}, \alpha + 1\}$.

It follows at once that π is a subrepresentation of $v^{-1/2}\rho \times \delta([v^{-y}\rho, v^{-3/2}\rho]) \rtimes \sigma$, and [Lemma 2.4](#), together with transitivity of Jacquet modules, shows that $\mu^*(\pi)$ contains an irreducible constituent of the form $v^{1/2}\rho \otimes \pi_1$. We will show that this is impossible, implying $a_{i_{\min}} \geq \frac{3}{2}$. Note that $\epsilon_\sigma(\rho, 2) = 1$, where $(\text{Jord}(\sigma), \sigma_{\text{cusp}}, \epsilon_\sigma)$ stands for the Jordan triple corresponding to σ , so we use [[Muić 2004](#), Theorem 5.1(ii)]. Only the case $y = \alpha + 1$ and $\alpha \geq \frac{3}{2}$ will be described in detail, since other cases can be obtained in the same way and the case $(y, \alpha) = (\frac{3}{2}, \frac{1}{2})$ is also, in the split case, discussed in [[Jantzen 1996](#)]. The following equality holds in $R(G)$:

$$\begin{aligned} \delta([v^{1/2}\rho, v^{\alpha+1}\rho]) \rtimes \sigma &= \pi + L(\delta([v^{-\alpha-1}\rho, v^{1/2}\rho]) \rtimes \sigma_{\text{sp}}^{(1)}) \\ &\quad + L(\delta([v^{-\alpha}\rho, v^{-1/2}\rho]) \rtimes \sigma_{\text{sp}}^{(2)}) \\ &\quad + L(\delta([v^{-\alpha}\rho, v^{1/2}\rho]) \rtimes \sigma_{\text{sp}}^{(3)}), \end{aligned}$$

where $\sigma_{\text{sp}}^{(1)}$, $\sigma_{\text{sp}}^{(2)}$, $\sigma_{\text{sp}}^{(3)}$ are the unique irreducible subrepresentations of

$$\begin{aligned} & v^{3/2}\rho \times v^{5/2}\rho \times \cdots \times v^\alpha\rho \rtimes \sigma_{\text{cusps}}, \\ & v^{1/2}\rho \times v^{3/2}\rho \times \cdots \times v^{\alpha-1}\rho \times \delta([v^\alpha\rho, v^{\alpha+1}\rho]) \rtimes \sigma_{\text{cusps}}, \\ & v^{3/2}\rho \times v^{5/2}\rho \times \cdots \times v^{\alpha-1}\rho \times \delta([v^\alpha\rho, v^{\alpha+1}\rho]) \rtimes \sigma_{\text{cusps}}, \end{aligned}$$

respectively. We note that $\sigma_{\text{sp}}^{(i)}$ is strongly positive for $i = 1, 2, 3$.

Using the structural formula, we obtain that if $v^{1/2}\rho \otimes \pi_1$ is an irreducible constituent of $\mu^*(\delta([v^{1/2}\rho, v^{\alpha+1}\rho]) \rtimes \sigma)$, then π_1 is an irreducible subquotient of $\delta([v^{1/2}\rho, v^{\alpha+1}\rho]) \rtimes \sigma_{\text{sp}}^{(1)}$. By [Muić 2004, Theorem 5.1(i)], in $R(G)$ we have:

$$\begin{aligned} \delta([v^{1/2}\rho, v^{\alpha+1}\rho]) \rtimes \sigma_{\text{sp}}^{(1)} &= L(\delta([v^{-\alpha-1}\rho, v^{-1/2}\rho]) \rtimes \sigma_{\text{sp}}^{(1)}) \\ &\quad + L(\delta([v^{-\alpha}\rho, v^{-1/2}\rho]) \rtimes \sigma_{\text{sp}}^{(3)}). \end{aligned}$$

Furthermore, both irreducible constituents $v^{1/2}\rho \otimes L(\delta([v^{-\alpha-1}\rho, v^{-1/2}\rho]) \rtimes \sigma_{\text{sp}}^{(1)})$ and $v^{1/2}\rho \otimes L(\delta([v^{-\alpha}\rho, v^{-1/2}\rho]) \rtimes \sigma_{\text{sp}}^{(3)})$ appear in $\mu^*(\delta([v^{1/2}\rho, v^{\alpha+1}\rho]) \rtimes \sigma)$ with multiplicity one, and Frobenius reciprocity implies that both

$$\mu^*(L(\delta([v^{-\alpha-1}\rho, v^{1/2}\rho]) \rtimes \sigma_{\text{sp}}^{(1)})) \quad \text{and} \quad \mu^*(L(\delta([v^{-\alpha}\rho, v^{1/2}\rho]) \rtimes \sigma_{\text{sp}}^{(3)}))$$

contain irreducible constituents of the form $v^{1/2}\rho \otimes \pi_1$, so $\mu^*(\pi)$ does not contain such an irreducible constituent.

Since $a_{i_{\min}} \geq \frac{3}{2}$, Lemma 3.5 implies that $\alpha \geq \frac{3}{2}$. From $y \geq a_{i_{\min}}$, using [Muić 2004, Proposition 3.1], we obtain that $v^y\rho \rtimes \sigma$ is irreducible if $y \neq \alpha + 1$. Suppose that $y \neq \alpha + 1$. Then we have the following embedding and isomorphism:

$$\pi \hookrightarrow \delta([v^{-y+1}\rho, v^{-x}\rho]) \times v^{-y}\rho \rtimes \sigma \cong v^y\rho \times \delta([v^{-y+1}\rho, v^{-x}\rho]) \rtimes \sigma.$$

In the same way as before we conclude that $\mu^*(\pi)$ contains an irreducible constituent of the form $v^{-y}\rho \otimes \pi_1$, which is impossible unless $x = y$. Thus, $y = a_{i_{\min}}$. It follows from Lemma 3.5 and [Muić 2004, Proposition 3.1] that $v^{-a_{i_{\min}}}\rho \rtimes \sigma \cong v^{a_{i_{\min}}}\rho \rtimes \sigma$, so π is an irreducible subrepresentation of $v^{a_{i_{\min}}}\rho \times v^{a_{i_{\min}}}\rho \rtimes \sigma_{\text{sp}}$. Consequently, Lemma 2.4 and the transitivity of Jacquet modules imply that the Jacquet module of π with respect to an appropriate parabolic subgroup contains an irreducible representation of the form $v^{-a_{i_{\min}}}\rho \otimes v^{-a_{i_{\min}}}\rho \otimes \pi_2$, and an easy application of Theorem 2.2 implies the Jacquet module of $\delta([v^{-y}\rho, v^{-x}\rho]) \rtimes \sigma$ with respect to an appropriate parabolic subgroup does not contain such a representation. Thus, $y = \alpha + 1$, and the proposition is proved. \square

Proposition 3.8. *Suppose that α is a half-integer, $\alpha \geq \frac{3}{2}$, $a_{i_{\min}} \geq \frac{3}{2}$, $x = a_{i_{\min}}$, and $y = \alpha + 1$. Then π is isomorphic to $\hat{\pi}$.*

Proof. Let us first prove that if an irreducible constituent of the form $v^z\rho \otimes \pi_1$, with $z \geq 0$, appears in $\mu^*(\pi)$, then $z = a_{i_{\min}}$. Using the structural formula and [Matić 2013,

Theorem 4.6], we deduce that if an irreducible constituent of the form $v^z\rho \otimes \pi_1$, with $z \geq 0$, appears in $\mu^*(\delta([v^{-\alpha-1}\rho, v^{-a_{i_{\min}}}\rho]) \rtimes \sigma)$, then $z \in \{a_{i_{\min}}, \alpha + 1\}$.

By [Muić 2004, Proposition 3.1(i)], in $R(G)$ we have

$$\delta([v^{a_{i_{\min}}}\rho, v^{\alpha+1}\rho]) \rtimes \sigma = \pi + L(\delta([v^{-\alpha}\rho, v^{-a_{i_{\min}}}\rho]) \rtimes \sigma_{\text{sp}}),$$

where σ_{sp} denotes the unique irreducible subrepresentation of

$$v^{a_{i_{\min}}}\rho \times v^{a_{i_{\min}}+1}\rho \times \cdots \times v^{\alpha-1}\rho \times \delta([v^{\alpha}\rho, v^{\alpha+1}\rho]) \rtimes \sigma_{\text{cusp}}.$$

We note that σ_{sp} is a strongly positive representation. Using [Matić 2013, Theorem 4.6], Frobenius reciprocity, and the transitivity of Jacquet modules, we obtain that $\mu^*(L(\delta([v^{-\alpha}\rho, v^{-a_{i_{\min}}}\rho]) \rtimes \sigma_{\text{sp}}))$ contains an irreducible constituent of the form $v^{\alpha+1}\rho \otimes \pi'$. The induced representation $\delta([v^{a_{i_{\min}}}\rho, v^{\alpha}\rho]) \rtimes \sigma$ is irreducible (by [Muić 2004, Proposition 3.1(ii)]), so the only such irreducible constituent appearing in $\mu^*(\delta([v^{a_{i_{\min}}}\rho, v^{\alpha+1}\rho]) \rtimes \sigma)$ is $v^{\alpha+1}\rho \otimes \delta([v^{a_{i_{\min}}}\rho, v^{\alpha}\rho]) \rtimes \sigma$, which appears there with multiplicity one. Therefore, there is no irreducible constituent of the form $v^{\alpha+1}\rho \otimes \pi_1$ appearing in $\mu^*(\pi)$. Furthermore, Lemma 2.4 implies that if an irreducible constituent of the form $v^z\rho \otimes \pi_1$ with $z \leq 0$ appears in $\mu^*(\hat{\pi})$, then $z = -a_{i_{\min}}$.

Since π is a subrepresentation of $\delta([v^{-\alpha-1}\rho, v^{-a_{i_{\min}}}\rho]) \rtimes \sigma$ and $a_{i_{\min}} \geq \frac{3}{2}$, we have the following embedding and isomorphisms:

$$\begin{aligned} \pi &\hookrightarrow v^{a_{i_{\min}}}\rho \times \cdots \times v^{\alpha}\rho \times v^{-a_{i_{\min}}}\rho \times \cdots \times v^{-\alpha}\rho \times v^{-\alpha-1}\rho \rtimes \sigma_{\text{cusp}} \\ &\cong v^{a_{i_{\min}}}\rho \times \cdots \times v^{\alpha}\rho \times v^{-a_{i_{\min}}}\rho \times \cdots \times v^{-\alpha}\rho \times v^{\alpha+1}\rho \rtimes \sigma_{\text{cusp}} \\ &\cong v^{a_{i_{\min}}}\rho \times \cdots \times v^{\alpha}\rho \times v^{\alpha+1}\rho \times v^{-a_{i_{\min}}}\rho \times \cdots \times v^{-\alpha}\rho \rtimes \sigma_{\text{cusp}}. \end{aligned}$$

Using Lemma 2.4, Theorem 2.1, and [Mœglin and Tadić 2002, Lemma 3.1], we obtain that $\hat{\pi}$ is a subrepresentation of the induced representation

$$v^{-a_{i_{\min}}}\rho \times \cdots \times v^{-\alpha-1}\rho \times v^{a_{i_{\min}}}\rho \times \cdots \times v^{\alpha}\rho \rtimes \sigma_{\text{cusp}}.$$

It follows from Lemma 3.2 of the same work that there exists an irreducible subquotient π_1 of $v^{-a_{i_{\min}}}\rho \times \cdots \times v^{-\alpha-1}\rho$ such that $\hat{\pi}$ is a subrepresentation of $\pi_1 \times v^{a_{i_{\min}}}\rho \times \cdots \times v^{\alpha}\rho \rtimes \sigma_{\text{cusp}}$. Since $\mu^*(\hat{\pi})$ does not contain an irreducible constituent of the form $v^z\rho \otimes \pi_1$ for $z \leq 0$ and $z \neq -a_{i_{\min}}$, we deduce that $\pi_1 \cong \delta([v^{-\alpha-1}\rho, v^{-a_{i_{\min}}}\rho])$.

By the same lemma, there is an irreducible representation π' such that $\hat{\pi}$ is a subrepresentation of $\delta([v^{-\alpha-1}\rho, v^{-a_{i_{\min}}}\rho]) \rtimes \pi'$ and, obviously, the cuspidal support of π' equals $\{v^{a_{i_{\min}}}\rho, v^{a_{i_{\min}}+1}\rho, \dots, v^{\alpha}\rho, \sigma_{\text{cusp}}\}$.

Let us first suppose that π' is a nontempered representation and write $\pi' \cong L(\delta_1 \times \delta_2 \times \cdots \times \delta_k \rtimes \tau)$, where $\delta_i \in \text{Irr}(\text{GL}(n_i, F))$ is an irreducible essentially square-integrable representation for $i = 1, 2, \dots, k$, $e(\delta_i) \leq e(\delta_{i+1}) < 0$ for $i = 1, 2, \dots, k-1$,

and $\tau \in \text{Irr}(G_{n'})$ is an irreducible tempered representation. Write $\delta_i = \delta([v^{a_i}\rho, v^{b_i}\rho])$. From $a_i + b_i < 0$ and from the description of the cuspidal support of π' follows that $b_i \leq -a_{i_{\min}}$, for $i = 1, 2, \dots, k$. It directly follows that π' is a subrepresentation of an induced representation of the form $v^{b_1}\rho \rtimes \pi''$ and, since $-\alpha \leq b_1$, we have the following embedding and isomorphism:

$$\hat{\pi} \hookrightarrow \delta([v^{-\alpha-1}\rho, v^{-a_{i_{\min}}}\rho]) \times v^{b_1}\rho \rtimes \pi'' \cong v^{b_1}\rho \times \delta([v^{-\alpha-1}\rho, v^{-a_{i_{\min}}}\rho]) \rtimes \pi'',$$

and it follows that $\mu^*(\hat{\pi})$ contains an irreducible constituent of the form $v^{b_1}\rho \otimes \pi_2$, which is impossible unless $b_1 = -a_{i_{\min}}$. If this is the case, we have an embedding

$$\hat{\pi} \hookrightarrow v^{-a_{i_{\min}}}\rho \times v^{-a_{i_{\min}}}\rho \times \delta([v^{-\alpha-1}\rho, v^{-a_{i_{\min}}-1}\rho]) \rtimes \pi'',$$

and [Lemma 2.4](#) and the transitivity of Jacquet modules imply that the Jacquet module of π with respect to an appropriate parabolic subgroup contains an irreducible representation of the form $v^{a_{i_{\min}}}\rho \otimes v^{a_{i_{\min}}}\rho \otimes \pi_2$. Using the structural formula, [\[Matić 2013, Theorem 4.6\]](#), and the fact that $\alpha + 1 > a_{i_{\min}}$, we deduce that a representation of the form $v^{a_{i_{\min}}}\rho \otimes v^{a_{i_{\min}}}\rho \otimes \pi_2$ does not appear in the Jacquet module of the induced representation $\delta([v^{-\alpha-1}\rho, v^{-a_{i_{\min}}}\rho]) \rtimes \sigma$ with respect to an appropriate parabolic subgroup, a contradiction.

Consequently, π' is a tempered representation and, using the description of its cuspidal support and [\[Matić 2012, Theorem 3.5\]](#), we conclude that π' is strongly positive. Since the strongly positive representation is completely determined by its cuspidal support ([\[Matić 2013, Lemma 3.5\]](#)), it follows at once that π' is isomorphic to σ . Thus, $\hat{\pi}$ is an irreducible subrepresentation of $\delta([v^{-\alpha-1}\rho, v^{-a_{i_{\min}}}\rho]) \rtimes \sigma$, leading to $\hat{\pi} \cong \pi$. This completes the proof. \square

Now we state our second main result.

Theorem 3.9. *Suppose that α is an integer. Then π is isomorphic to $\hat{\pi}$ if and only if $y = \alpha + 1$ and either $x = a_{i_{\min}}$ or $(a_{i_{\min}}, x) = (1, 0)$.*

[Theorem 3.9](#) follows from the following two propositions:

Proposition 3.10. *Suppose that α is an integer and $\pi \cong \hat{\pi}$. Then $y = \alpha + 1$ and either $x = a_{i_{\min}}$ or $(a_{i_{\min}}, x) = (1, 0)$.*

Proof. If $a_{i_{\min}} \geq 2$, in the same way as in the proof of [Proposition 3.7](#), we deduce that $(x, y) = (a_{i_{\min}}, \alpha + 1)$.

Let us now assume that $a_{i_{\min}} = 1$. Then σ is a subrepresentation of an induced representation of the form $v\rho \rtimes \sigma'$ and if $x > 0$ we have an embedding $\pi \hookrightarrow v\rho \times \delta([v^{-y}\rho, v^{-x}\rho]) \rtimes \sigma'$. In the same way as in the proof of [Proposition 3.7](#), we get that $x \in \{0, 1\}$. Note that $y > x$ if $x = 0$. Let us prove that $y = \alpha + 1$. Suppose, contrary to our assumption, that $y \neq \alpha + 1$. Since $y \geq x$, it follows from [\[Muić](#)

2004, Proposition 3.1] that $v^y\rho \rtimes \sigma$ is irreducible. We have

$$\pi \hookrightarrow \delta([v^{-y+1}\rho, v^{-x}\rho]) \times v^{-y}\rho \rtimes \sigma \cong \delta([v^{-y+1}\rho, v^{-x}\rho]) \times v^y\rho \rtimes \sigma,$$

and if $y \neq -x + 1$ one obtains a contradiction in the same way as in the proof of Proposition 3.7. Since $y \geq x$ and $x \geq 0$, we see that $y = -x + 1$ holds only if $(x, y) = (0, 1)$. In that case, we have $\pi \hookrightarrow \rho \times v\rho \times v\rho \rtimes \sigma'$. Using Lemma 2.4 and the transitivity of Jacquet modules we get that $r_M(\pi)$ contains an irreducible representation of the form $\rho \otimes v^{-1}\rho \otimes v^{-1}\rho \otimes \sigma''$, where M denotes the Levi factor of an appropriate parabolic subgroup. But, since σ is strongly positive, $r_M(\delta([v\rho, v\rho]) \rtimes \sigma)$ does not contain an irreducible representation of the form $\rho \otimes v^{-1}\rho \otimes v^{-1}\rho \otimes \sigma'$, a contradiction. This completes the proof. \square

Proposition 3.11. *Suppose that α is an integer. If $y = \alpha + 1$ and either $x = a_{i_{\min}}$ or $(a_{i_{\min}}, x) = (1, 0)$, then π is isomorphic to $\hat{\pi}$.*

Proof. First we suppose that $(x, y) = (a_{i_{\min}}, \alpha + 1)$. Let us prove that if $\mu^*(\pi)$ contains an irreducible constituent of the form $v^z\rho \otimes \pi'$, with $z \geq 0$, then $z = a_{i_{\min}}$. It follows from the structural formula that if $\mu^*(\delta([v^{a_{i_{\min}}}\rho, v^{\alpha+1}\rho]) \rtimes \sigma)$ contains an irreducible constituent of the form $v^z\rho \otimes \pi'$, with $z \geq 0$, then $z \in \{a_{i_{\min}}, \alpha + 1\}$. Also, if $z = \alpha + 1$, then π' is an irreducible subquotient of $\delta([v^{a_{i_{\min}}}\rho, v^\alpha\rho]) \rtimes \sigma$. We have $\delta([v^{a_{i_{\min}}}\rho, v^\alpha\rho]) \rtimes \sigma$ is irreducible and $v^{\alpha+1}\rho \otimes \delta([v^{a_{i_{\min}}}\rho, v^\alpha\rho]) \rtimes \sigma$ is contained in $\mu^*(\delta([v^{a_{i_{\min}}}\rho, v^{\alpha+1}\rho]) \rtimes \sigma)$ with multiplicity one by [Muić 2004, Proposition 3.1]. Using part (i) of the same proposition, we deduce that in $R(G)$ we have $\delta([v^{a_{i_{\min}}}\rho, v^{\alpha+1}\rho]) \rtimes \sigma = \pi + L(\delta([v^{-\alpha}\rho, v^{-a_{i_{\min}}}\rho]) \rtimes \sigma_{\text{sp}})$, where σ_{sp} denotes the unique irreducible subrepresentation of

$$v^{a_{i_{\min}}}\rho \times v^{a_{i_{\min}}+1}\rho \times \dots \times v^{\alpha-1}\rho \times \delta([v^\alpha\rho, v^{\alpha+1}\rho]) \rtimes \sigma_{\text{cusp}}.$$

We note that σ_{sp} is a strongly positive representation. It is now easy to conclude, using Frobenius reciprocity and the irreducibility of $v^x\rho \times v^{\alpha+1}\rho$ for $x < \alpha$, that $\mu^*(L(\delta([v^{-\alpha}\rho, v^{-a_{i_{\min}}}\rho]) \rtimes \sigma_{\text{sp}}))$ contains an irreducible constituent of the form $v^{\alpha+1}\rho \otimes \pi'$, so $\mu^*(\pi)$ does not contain such an irreducible constituent. Now, in the same way as in the proof of Proposition 3.8, we obtain that $\hat{\pi}$ is a subrepresentation of $\delta([v^{-\alpha-1}\rho, v^{-a_{i_{\min}}}\rho]) \rtimes \sigma$, i.e., $\pi \cong \hat{\pi}$.

Now we turn our attention to the case $(a_{i_{\min}}, x, y) = (1, 0, \alpha + 1)$. In this case, we have the following embedding and isomorphisms:

$$\begin{aligned} \pi &\hookrightarrow \rho \times v^{-1}\rho \times \dots \times v^{-\alpha-1}\rho \times v\rho \times v^2\rho \times \dots \times v^\alpha\rho \rtimes \sigma_{\text{cusp}} \\ &\cong \rho \times v\rho \times v^2\rho \times \dots \times v^\alpha\rho \times v^{-1}\rho \times \dots \times v^{-\alpha}\rho \times v^{-\alpha-1}\rho \rtimes \sigma_{\text{cusp}} \\ &\cong \rho \times v\rho \times v^2\rho \times \dots \times v^\alpha\rho \times v^{-1}\rho \times \dots \times v^{-\alpha}\rho \times v^{\alpha+1}\rho \rtimes \sigma_{\text{cusp}} \\ &\cong \rho \times v\rho \times v^2\rho \times \dots \times v^\alpha\rho \times v^{\alpha+1}\rho \times v^{-1}\rho \times \dots \times v^{-\alpha}\rho \rtimes \sigma_{\text{cusp}}. \end{aligned}$$

In the same way as before, we obtain that $\hat{\pi}$ is a subrepresentation of the induced representation

$$\rho \times v^{-1}\rho \times v^{-2}\rho \times \cdots \times v^{-\alpha}\rho \times v^{-\alpha-1}\rho \times v\rho \times \cdots \times v^{\alpha}\rho \rtimes \sigma_{\text{cusp}}.$$

We will show that if $\mu^*(\pi)$ contains an irreducible constituent of the form $v^z\rho \otimes \pi_1$, with $z \geq 0$, then $z = 0$. The rest of the proof then follows similarly to [Proposition 3.8](#).

Note that if an irreducible constituent of the form $v^z\rho \otimes \pi_1$, with $z \geq 0$, appears in $\mu^*(\delta([\rho, v^{\alpha+1}\rho]) \rtimes \sigma)$, then $z \in \{0, 1, \alpha + 1\}$. We will comment only on the case $\alpha \geq 2$, since the case $\alpha = 1$ can be handled in the same way but more easily, and in the split case it can also be obtained using [\[Jantzen 1996\]](#).

According to [\[Muić 2004, Theorem 4.1\(iv\)\]](#), in $R(G)$ we have

$$\begin{aligned} \delta([\rho, v^{\alpha+1}\rho]) \rtimes \sigma &= \pi + L(\delta([v^{-\alpha-1}\rho, v\rho]) \rtimes \sigma_{\text{sp}}^{(1)}) + L(\delta([v^{-\alpha}\rho, \rho]) \rtimes \sigma_{\text{sp}}^{(2)}) \\ &\quad + L(\delta([v^{-\alpha}\rho, v\rho]) \rtimes \sigma_{\text{sp}}^{(3)}), \end{aligned}$$

where $\sigma_{\text{sp}}^{(1)}, \sigma_{\text{sp}}^{(2)}, \sigma_{\text{sp}}^{(3)}$ are the unique irreducible subrepresentations of

$$\begin{aligned} v^2\rho \times \cdots \times v^{\alpha}\rho \rtimes \sigma_{\text{cusp}}, \quad v\rho \times \cdots \times v^{\alpha-1}\rho \times \delta([v^{\alpha}\rho, v^{\alpha+1}\rho]) \rtimes \sigma_{\text{cusp}}, \\ v^2\rho \times \cdots \times v^{\alpha-1}\rho \times \delta([v^{\alpha}\rho, v^{\alpha+1}\rho]) \rtimes \sigma_{\text{cusp}}, \end{aligned}$$

respectively. We note that $\sigma_{\text{sp}}^{(i)}$ is strongly positive for $i = 1, 2, 3$.

If $\mu^*(\delta([\rho, v^{\alpha+1}\rho]) \rtimes \sigma)$ contains an irreducible constituent of the form $v\rho \otimes \pi_1$, then π_1 is an irreducible subquotient of $\delta([\rho, v^{\alpha+1}\rho]) \rtimes \sigma_{\text{sp}}^{(1)}$. By [\[Muić 2004, Theorem 4.1\(ii\)\]](#), in $R(G)$ we have

$$\delta([\rho, v^{\alpha+1}\rho]) \rtimes \sigma_{\text{sp}}^{(1)} = L(\delta([v^{-\alpha-1}\rho, \rho]) \rtimes \sigma_{\text{sp}}^{(1)}) + L(\delta([v^{-\alpha}\rho, \rho]) \rtimes \sigma_{\text{sp}}^{(3)}).$$

Also, $v\rho \otimes L(\delta([v^{-\alpha-1}\rho, \rho]) \rtimes \sigma_{\text{sp}}^{(1)})$ and $v\rho \otimes L(\delta([v^{-\alpha}\rho, \rho]) \rtimes \sigma_{\text{sp}}^{(3)})$ appear in $\mu^*(\delta([\rho, v^{\alpha+1}\rho]) \rtimes \sigma)$ with multiplicity one and are obviously contained in $\mu^*(L(\delta([v^{-\alpha-1}\rho, v\rho]) \rtimes \sigma_{\text{sp}}^{(1)}))$ and in $\mu^*(L(\delta([v^{-\alpha}\rho, v\rho]) \rtimes \sigma_{\text{sp}}^{(3)}))$. Thus, there are no irreducible constituents of the form $v\rho \otimes \pi_1$ appearing in $\mu^*(\pi)$.

Similarly, if $\mu^*(\delta([\rho, v^{\alpha+1}\rho]) \rtimes \sigma)$ contains an irreducible constituent of the form $v^{\alpha+1}\rho \otimes \pi_1$, then π_1 is an irreducible subquotient of $\delta([\rho, v^{\alpha}\rho]) \rtimes \sigma$. By [\[Muić 2004, Theorem 4.1\(iii\)\]](#), in $R(G)$ we have

$$\delta([\rho, v^{\alpha}\rho]) \rtimes \sigma = L(\delta([v^{-\alpha}\rho, \rho]) \rtimes \sigma) + L(\delta([v^{-\alpha}\rho, v\rho]) \rtimes \sigma_{\text{sp}}^{(1)}).$$

Also, $v^{\alpha+1}\rho \otimes L(\delta([v^{-\alpha}\rho, \rho]) \rtimes \sigma_{\text{sp}})$ and $v^{\alpha+1}\rho \otimes L(\delta([v^{-\alpha}\rho, v\rho]) \rtimes \sigma_{\text{sp}}^{(1)})$ appear in $\mu^*(\delta([\rho, v^{\alpha+1}\rho]) \rtimes \sigma)$ with multiplicity one and obviously appear in $\mu^*(L(\delta([v^{-\alpha}\rho, \rho]) \rtimes \sigma_{\text{sp}}^{(2)}))$ and in $\mu^*(L(\delta([v^{-\alpha}\rho, v\rho]) \rtimes \sigma_{\text{sp}}^{(3)}))$. Thus, $\mu^*(\pi)$ does not contain irreducible constituents of the form $v^{\alpha+1}\rho \otimes \pi_1$. Consequently, $\mu^*(\pi)$ does not contain an irreducible constituent of the form $v^z\rho \otimes \pi_1$ with $z > 0$, and the proposition is proved. \square

Acknowledgements

The author would like to thank the referee for helping to improve the presentation style and for the help with [Proposition 3.2](#). This work has been supported by Croatian Science Foundation under the project 9364.

References

- [Arthur 2013] J. Arthur, *The endoscopic classification of representations: orthogonal and symplectic groups*, American Mathematical Society Colloquium Publications **61**, American Mathematical Society, Providence, RI, 2013. [MR](#) [Zbl](#)
- [Aubert 1995] A.-M. Aubert, “Dualité dans le groupe de Grothendieck de la catégorie des représentations lisses de longueur finie d’un groupe réductif p -adique”, *Trans. Amer. Math. Soc.* **347**:6 (1995), 2179–2189. Correction in **348**:11 (1996), 4687–4690. [MR](#) [Zbl](#)
- [Jantzen 1996] C. Jantzen, *Degenerate principal series for symplectic and odd-orthogonal groups*, Mem. Amer. Math. Soc. **590**, American Mathematical Society, Providence, RI, 1996. [MR](#) [Zbl](#)
- [Matić 2011] I. Matić, “Strongly positive representations of metaplectic groups”, *J. Algebra* **334** (2011), 255–274. [MR](#) [Zbl](#)
- [Matić 2012] I. Matić, “Theta lifts of strongly positive discrete series: the case of $(\widetilde{\mathrm{Sp}}(n), O(V))$ ”, *Pacific J. Math.* **259**:2 (2012), 445–471. [MR](#) [Zbl](#)
- [Matić 2013] I. Matić, “Jacquet modules of strongly positive representations of the metaplectic group $\widetilde{\mathrm{Sp}}(n)$ ”, *Trans. Amer. Math. Soc.* **365**:5 (2013), 2755–2778. [MR](#) [Zbl](#)
- [Matić 2015] I. Matić, “Strongly positive subquotients in a class of induced representations of classical p -adic groups”, *J. Algebra* **444** (2015), 504–526. [MR](#) [Zbl](#)
- [Matić 2016a] I. Matić, “Aubert duals of discrete series: the first inductive step”, preprint, 2016, Available at <http://tinyurl.com/aubertduals>.
- [Matić 2016b] I. Matić, “On discrete series subrepresentations of the generalized principal series”, *Glas. Mat. Ser. III* **51**:1 (2016), 125–152. [MR](#) [Zbl](#)
- [Matić 2016c] I. Matić, “On Jacquet modules of discrete series: the first inductive step”, *J. Lie Theory* **26**:1 (2016), 135–168. [MR](#) [Zbl](#)
- [Matić 2017] I. Matić, “Aubert duals of strongly positive discrete series and a class of unitarizable representations”, *Proc. Amer. Math. Soc.* **145**:8 (2017), 3561–3570.
- [Matić and Tadić 2015] I. Matić and M. Tadić, “On Jacquet modules of representations of segment type”, *Manuscripta Math.* **147**:3-4 (2015), 437–476. [MR](#) [Zbl](#)
- [Mœglin 2002] C. Mœglin, “Sur la classification des séries discrètes des groupes classiques p -adiques: paramètres de Langlands et exhaustivité”, *J. Eur. Math. Soc.* **4**:2 (2002), 143–200. [MR](#) [Zbl](#)
- [Mœglin 2014] C. Mœglin, “Paquets stables des séries discrètes accessibles par endoscopie tordue; leur paramètre de Langlands”, pp. 295–336 in *Automorphic forms and related geometry: assessing the legacy of I. I. Piatetski-Shapiro*, edited by J. W. Cogdell et al., Contemp. Math. **614**, American Mathematical Society, Providence, RI, 2014. [MR](#) [Zbl](#)
- [Mœglin and Tadić 2002] C. Mœglin and M. Tadić, “Construction of discrete series for classical p -adic groups”, *J. Amer. Math. Soc.* **15**:3 (2002), 715–786. [MR](#) [Zbl](#)
- [Muić 2004] G. Muić, “Composition series of generalized principal series: the case of strongly positive discrete series”, *Israel J. Math.* **140** (2004), 157–202. [MR](#) [Zbl](#)

- [Schneider and Stuhler 1997] P. Schneider and U. Stuhler, “Representation theory and sheaves on the Bruhat–Tits building”, *Inst. Hautes Études Sci. Publ. Math.* 85 (1997), 97–191. MR Zbl
- [Shahidi 1990] F. Shahidi, “A proof of Langlands’ conjecture on Plancherel measures; complementary series for p -adic groups”, *Ann. of Math. (2)* 132:2 (1990), 273–330. MR Zbl
- [Shahidi 1992] F. Shahidi, “Twisted endoscopy and reducibility of induced representations for p -adic groups”, *Duke Math. J.* 66:1 (1992), 1–41. MR Zbl
- [Tadić 1995] M. Tadić, “Structure arising from induction and Jacquet modules of representations of classical p -adic groups”, *J. Algebra* 177:1 (1995), 1–33. MR Zbl
- [Tadić 2013] M. Tadić, “On tempered and square integrable representations of classical p -adic groups”, *Sci. China Math.* 56:11 (2013), 2273–2313. MR Zbl
- [Zelevinsky 1980] A. V. Zelevinsky, “Induced representations of reductive p -adic groups, II: On irreducible representations of $GL(n)$ ”, *Ann. Sci. École Norm. Sup. (4)* 13:2 (1980), 165–210. MR Zbl

Received October 21, 2016. Revised February 10, 2017.

IVAN MATIĆ
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF OSIJEK
TRG LJUDEVITA GAJA 6
31000 OSIJEK
CROATIA
imatic@mathos.hr

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 289 No. 2 August 2017

Regular representations of completely bounded maps	257
B. V. RAJARAMA BHAT, NIRUPAMA MALLICK and K. SUMESH	
Ball convex bodies in Minkowski spaces	287
THOMAS JAHN, HORST MARTINI and CHRISTIAN RICHTER	
Local constancy of dimension of slope subspaces of automorphic forms	317
JOACHIM MAHNKOPF	
Weakening idempotency in K -theory	381
VLADIMIR MANUILOV	
On Langlands quotients of the generalized principal series isomorphic to their Aubert duals	395
IVAN MATIĆ	
Exact Lagrangian fillings of Legendrian $(2, n)$ torus links	417
YU PAN	
Elementary calculation of the cohomology rings of real Grassmann manifolds	443
RUSTAM SADYKOV	
Cluster tilting modules and noncommutative projective schemes	449
KENTA UHEYAMA	
Concentration for a biharmonic Schrödinger equation	469
DONG WANG	
Global existence of smooth solutions to exponential wave maps in FLRW spacetimes	489
CHANG-HUA WEI and NING-AN LAI	