

# *Pacific Journal of Mathematics*

**CLUSTER TILTING MODULES  
AND NONCOMMUTATIVE PROJECTIVE SCHEMES**

KENTA UYAMA

Volume 289    No. 2

August 2017



# CLUSTER TILTING MODULES AND NONCOMMUTATIVE PROJECTIVE SCHEMES

KENTA UHEYAMA

We study the relationship between equivalences of noncommutative projective schemes and cluster tilting modules. In particular, we prove the following result. Let  $A$  be an AS-Gorenstein algebra of dimension  $d \geq 2$  and tails  $A$  the noncommutative projective scheme associated to  $A$ . If  $\text{gldim}(\text{tails } A) < \infty$  and  $A$  has a  $(d-1)$ -cluster tilting module  $X$  with the property that its graded endomorphism algebra is  $\mathbb{N}$ -graded, then the graded endomorphism algebra  $B$  of a basic  $(d-1)$ -cluster tilting submodule of  $X$  is a two-sided noetherian  $\mathbb{N}$ -graded AS-regular algebra over  $B_0$  of global dimension  $d$  such that tails  $B$  is equivalent to tails  $A$ .

## 1. Introduction

Artin and Zhang [1994] introduced the notion of a noncommutative projective scheme, and established a fundamental and comprehensive theory of noncommutative projective schemes. Since the study of the categories of coherent sheaves on commutative projective schemes (or their derived categories) is of increasing importance in algebraic geometry, the study of noncommutative projective schemes has been a major project in noncommutative projective geometry.

Let  $A, A'$  be right noetherian graded algebras, and tails  $A$ , tails  $A'$  the noncommutative projective schemes associated to  $A$  and  $A'$  respectively. Clearly, if  $A \cong A'$  as graded algebras, then tails  $A \cong$  tails  $A'$ . It is well known that the converse does not hold, so the following question is a natural one to ask.

**Question 1.1.** Given a right noetherian graded algebra  $A$ , can we find a better homogeneous coordinate ring of tails  $A$ ? That is, can we find a better graded algebra  $B$  (e.g.,  $\text{gldim } B < \infty$ ) such that tails  $B \cong$  tails  $A$ ?

For example, take the commutative graded algebra  $A = k[x, y, z, w]/(xw - yz)$ . Then tails  $A$  ( $\cong \text{coh } \mathbb{P}^1 \times \mathbb{P}^1$ ) is not equivalent to tails  $k[x_1, \dots, x_n]$  ( $\cong \text{coh } \mathbb{P}^{n-1}$ ), but we can find the noncommutative graded algebra  $B = k\langle x, y \rangle / (x^2y - yx^2, y^2x - xy^2)$

---

The author was supported by JSPS Grant-in-Aid for Young Scientists (B) 15K17503.

MSC2010: primary 16S38; secondary 14A22, 16G50, 16W50.

Keywords: noncommutative projective scheme, cluster tilting module, AS-Gorenstein algebra, AS-regular algebra, ASF-regular algebra.

of global dimension 3 such that  $\text{tails } A \cong \text{tails } B$ , so the above question has a significant meaning in noncommutative projective geometry.

The purpose of this paper is to give an answer to Question 1.1 by investigating cluster tilting modules. Cluster tilting modules are crucial in the study of higher-dimensional analogues of Auslander–Reiten theory, and also attract attention from the viewpoint of Van den Bergh’s noncommutative crepant resolutions. In particular, cluster tilting modules have been extensively studied for a certain class of algebras, called orders, including commutative Cohen–Macaulay rings and finite-dimensional algebras (see [Iyama 2008]). One of the goals of this paper is to develop cluster tilting theory for nonorders in terms of noncommutative projective geometry.

The main result of this paper is as follows. Let  $A$  be a two-sided noetherian connected graded algebra satisfying  $\chi$  and let  $X$  be a finitely generated graded right  $A$ -module. If  $A$  is an AS-Gorenstein algebra of dimension  $d \geq 2$  and  $X$  is a  $(d-1)$ -cluster tilting module satisfying some additional conditions, then the graded endomorphism algebra  $B = \underline{\text{End}}_A(X)$  is a two-sided noetherian ASF-regular algebra of global dimension  $d$  such that the functors

$$\text{tails } B \rightarrow \text{tails } A \text{ induced by } - \otimes_B X$$

and

$$\text{tails } B^{\text{op}} \rightarrow \text{tails } A^{\text{op}} \text{ induced by } \underline{\text{Hom}}_A(X, A) \otimes_B -$$

are equivalences (Theorem 3.10 (1)). Moreover, a certain converse statement also holds (Theorem 3.10 (2)). As a corollary of this result, we can answer Question 1.1.

**Theorem 1.2** (Corollary 3.12). *Let  $A$  be an AS-Gorenstein algebra of dimension  $d \geq 2$ . If  $\text{gldim}(\text{tails } A) < \infty$  and  $A$  has a  $(d-1)$ -cluster tilting module  $X$  with the property that its graded endomorphism algebra is  $\mathbb{N}$ -graded, then the graded endomorphism algebra  $B$  of a basic  $(d-1)$ -cluster tilting submodule of  $X$  is a two-sided noetherian  $\mathbb{N}$ -graded AS-regular algebra over  $B_0$  of global dimension  $d$  such that  $\text{tails } B \cong \text{tails } A$ .*

We note that the notions of ASF-regular and AS-regular over  $R$  were recently introduced by Minamoto and Mori [2011], and these are natural generalizations of AS-regular algebras for  $\mathbb{N}$ -graded (not necessarily connected graded) algebras. A comparison theorem for these algebras is described in Theorem 2.10 (see also Corollary 2.11).

## 2. Preliminaries

Throughout, let  $k$  be a field. A graded  $k$ -vector space  $V = \bigoplus_{i \in \mathbb{Z}} V_i$  is called locally finite if  $\dim_k V_i < \infty$  for all  $i \in \mathbb{Z}$ , and it is called left (resp. right) bounded if  $V_i = 0$  for all  $i \ll 0$  (resp.  $i \gg 0$ ). We denote by  $DV = \underline{\text{Hom}}_k(V, k)$  the graded vector space dual of a locally finite graded  $k$ -vector space  $V$ .

In this paper, a graded algebra means a  $\mathbb{Z}$ -graded algebra over  $k$  unless otherwise stated. For a graded algebra  $A$ , we denote by  $\text{GrMod } A$  the category of graded right  $A$ -modules with  $A$ -module homomorphisms of degree 0, and by  $\text{grmod } A$  the full subcategory consisting of finitely generated graded  $A$ -modules. Note that if  $A$  is right noetherian, then  $\text{grmod } A$  is an abelian category. We denote by  $A^{\text{op}}$  the opposite algebra of  $A$ , and by  $A^e = A^{\text{op}} \otimes_k A$  the enveloping algebra. The category of graded left  $A$ -modules is identified with  $\text{GrMod } A^{\text{op}}$ , and the category of graded  $A$ - $A$  bimodules is identified with  $\text{GrMod } A^e$ .

For a graded module  $M \in \text{GrMod } A$  and an integer  $n \in \mathbb{Z}$ , we define the shift  $M(n) \in \text{GrMod } A$  by  $M(n)_i := M_{n+i}$  for  $i \in \mathbb{Z}$ . Note that the rule  $M \mapsto M(n)$  is a  $k$ -linear autoequivalence for  $\text{GrMod } A$  and  $\text{grmod } A$ , called the shift functor. For  $M, N \in \text{GrMod } A$ , we write  $\text{Ext}_{\text{GrMod } A}^i(M, N)$  for the extension group in  $\text{GrMod } A$ , and define

$$\underline{\text{Ext}}_A^i(M, N) := \bigoplus_{i \in \mathbb{Z}} \text{Ext}_{\text{GrMod } A}^i(M, N(i)).$$

Let  $A$  be a right noetherian locally finite  $\mathbb{N}$ -graded algebra. For  $M \in \text{GrMod } A$  and an integer  $n \in \mathbb{Z}$ , we define the truncated submodule  $M_{\geq n} \in \text{GrMod } A$  by  $M_{\geq n} := \bigoplus_{i \geq n} M_i$ . We say that an element  $x$  of a graded module  $M \in \text{GrMod } A$  is torsion if there exists a positive integer  $n$  such that  $x A_{\geq n} = 0$ . We denote by  $t(M)$  the submodule of  $M$  consisting of all torsion elements. A graded module  $M \in \text{GrMod } A$  is called torsion if  $M = t(M)$ , and torsion-free if  $t(M) = 0$ . We denote by  $\text{Tors } A$  (resp.  $\text{tors } A$ ) the full subcategory of  $\text{GrMod } A$  (resp.  $\text{grmod } A$ ) consisting of torsion modules. One can define the Serre quotient categories

$$\text{Tails } A = \text{GrMod } A / \text{Tors } A \quad \text{and} \quad \text{tails } A = \text{grmod } A / \text{tors } A.$$

Note that  $\text{tails } A$  is the full subcategory of noetherian objects of  $\text{Tails } A$ . The quotient functor is denoted by  $\pi : \text{GrMod } A \rightarrow \text{Tails } A$ . We often denote by  $\mathcal{M} = \pi M \in \text{Tails } A$  the image of  $M \in \text{GrMod } A$ . Note that the shift functor preserves torsion modules, so it induces a  $k$ -linear autoequivalence  $\mathcal{M} \mapsto \mathcal{M}(n)$  for  $\text{Tails } A$  and  $\text{tails } A$ , again called the shift functor. For  $\mathcal{M}, \mathcal{N} \in \text{Tails } A$ , we write  $\text{Ext}_{\text{Tails } A}^i(\mathcal{M}, \mathcal{N})$  for the extension group in  $\text{Tails } A$ , and define

$$\underline{\text{Ext}}_{\mathcal{A}}^i(\mathcal{M}, \mathcal{N}) := \bigoplus_{i \in \mathbb{Z}} \text{Ext}_{\text{Tails } A}^i(\mathcal{M}, \mathcal{N}(i)).$$

See [Artin and Zhang 1994, Section 7] for details on Ext groups in  $\text{tails } A$ . Following Serre's theorem and the Gabriel–Rosenberg reconstruction theorem,  $\text{tails } A$  is called the noncommutative projective scheme associated to  $A$  (see [Artin and Zhang 1994] for details). We define the global dimension of  $\text{tails } A$  by

$$\text{gldim}(\text{tails } A) = \sup\{i \mid \text{Ext}_{\text{tails } A}^i(\mathcal{M}, \mathcal{N}) \neq 0 \text{ for some } \mathcal{M}, \mathcal{N} \in \text{tails } A\}.$$

If  $\text{gldim } A < \infty$ , it is clear that  $\text{gldim}(\text{tails } A) < \infty$ . The condition  $\text{gldim}(\text{tails } A) < \infty$  is considered as a noncommutative graded isolated singularity property (see [Ueyama 2013; 2015; Mori and Ueyama 2016a; 2016b]).

Recall that we say that  $\chi_i$  holds for  $A$  if  $\underline{\text{Ext}}_A^j(A/A_{\geq 1}, M)$  is finite-dimensional over  $k$  for every  $M \in \text{grmod } A$  and every  $j \leq i$ , and we say that  $\chi$  holds for  $A$  if  $\chi_i$  holds for every  $i$ . The condition  $\chi_i$  plays an essential role in the study of the noncommutative projective scheme  $\text{tails } A$  (see [Artin and Zhang 1994; Yekutieli and Zhang 1997] for details).

We call  $(C, \mathcal{O}, s)$  an algebraic triple if it consists of a  $k$ -linear abelian category  $C$ , an object  $\mathcal{O} \in C$ , and a  $k$ -linear autoequivalence  $s \in \text{Aut}_k C$ .

**Definition 2.1** [Artin and Zhang 1994]. Let  $(C, \mathcal{O}, s)$  be an algebraic triple. We say that the pair  $(\mathcal{O}, s)$  is ample for  $C$  if

- (Am1) for every object  $\mathcal{M} \in C$ , there are positive integers  $r_1, \dots, r_p \in \mathbb{N}^+$  and an epimorphism  $\bigoplus_{i=1}^p s^{-r_i} \mathcal{O} \rightarrow \mathcal{M}$  in  $C$ , and
- (Am2) for every epimorphism  $\mathcal{M} \rightarrow \mathcal{N}$  in  $C$ , there is an integer  $n_0$  such that the induced map  $\text{Hom}_C(s^{-n} \mathcal{O}, \mathcal{M}) \rightarrow \text{Hom}_C(s^{-n} \mathcal{O}, \mathcal{N})$  is surjective for every  $n \geq n_0$ .

We define the graded algebra associated to an algebraic triple  $(C, \mathcal{O}, s)$  by  $B(C, \mathcal{O}, s) := \bigoplus_{i \in \mathbb{Z}} \text{Hom}_C(\mathcal{O}, s^i \mathcal{O})$ . Moreover, for any object  $\mathcal{M} \in C$ , it is known that  $\bigoplus_{i \in \mathbb{Z}} \text{Hom}_C(\mathcal{O}, s^i \mathcal{M})$  has a natural graded right  $B(C, \mathcal{O}, s)$ -module structure.

**Theorem 2.2** [Artin and Zhang 1994, Corollary 4.6 (1)]. *Let  $(C, \mathcal{O}, s)$  be an algebraic triple. If  $\mathcal{O} \in C$  is a noetherian object,  $\dim_k \text{Hom}_C(\mathcal{O}, \mathcal{M}) < \infty$  for all  $\mathcal{M} \in C$ , and  $(\mathcal{O}, s)$  is ample for  $C$ , then  $B = B(C, \mathcal{O}, s)_{\geq 0}$  is a right noetherian locally finite  $\mathbb{N}$ -graded algebra satisfying  $\chi_1$ , and the functor*

$$C \rightarrow \text{tails } B, \quad \mathcal{F} \mapsto \pi \left( \bigoplus_{i \in \mathbb{N}} \text{Hom}_C(\mathcal{O}, s^i \mathcal{F}) \right)$$

*induces an equivalence of algebraic triples  $(C, \mathcal{O}, s) \rightarrow (\text{tails } B, \mathcal{B}, (1))$ .*

Let  $A$  be an  $\mathbb{N}$ -graded algebra. Then the augmentation ideal  $A_{\geq 1}$  is denoted by  $\mathfrak{m}$ . We define the functor  $\underline{\Gamma}_{\mathfrak{m}} : \text{GrMod } A \rightarrow \text{GrMod } A$  by

$$\underline{\Gamma}_{\mathfrak{m}}(-) = \lim_{n \rightarrow \infty} \underline{\text{Hom}}_A(A/A_{\geq n}, -).$$

The derived functor of  $\underline{\Gamma}_{\mathfrak{m}}$  is denoted by  $R\underline{\Gamma}_{\mathfrak{m}}(-)$ , and its cohomologies are denoted by  $\underline{H}_{\mathfrak{m}}^i(-) = h^i(R\underline{\Gamma}_{\mathfrak{m}}(-))$ . For a graded module  $M \in \text{GrMod } A$ , we define

$$\text{depth } M = \inf\{i \mid \underline{H}_{\mathfrak{m}}^i(M) \neq 0\} \quad \text{and} \quad \text{ldim } M = \sup\{i \mid \underline{H}_{\mathfrak{m}}^i(M) \neq 0\}.$$

See [Yekutieli 1992; Van den Bergh 1997] for basic properties of  $R\underline{\Gamma}_{\mathfrak{m}}(-)$ .

If  $A$  is an  $\mathbb{N}$ -graded algebra with  $A_0 = k$ , then  $A$  is called connected graded. Note that a right noetherian connected graded algebra is locally finite.

**Definition 2.3.** A two-sided noetherian connected graded algebra  $A$  is called AS-Gorenstein (resp. AS-regular) of dimension  $d$  and of Gorenstein parameter  $\ell$  if

- $\text{injdim}_A A = \text{injdim}_{A^{\text{op}}} A = d < \infty$  (resp.  $\text{gldim } A = d < \infty$ ), and
- $\underline{\text{Ext}}_A^i(k, A) \cong \underline{\text{Ext}}_{A^{\text{op}}}^i(k, A) \cong \begin{cases} k(\ell) & \text{if } i = d, \\ 0 & \text{if } i \neq d. \end{cases}$

For  $M \in \text{GrMod } A$  and a graded algebra automorphism  $\sigma \in \text{GrAut } A$ , we define the twist  $M_\sigma \in \text{GrMod } A$  by  $M_\sigma = M$  as a graded  $k$ -vector space with the new right action  $m * a = m\sigma(a)$ . Let  $A$  be an AS-Gorenstein algebra. Then it is well known that  $A$  has a balanced dualizing complex  $DR\Gamma_{\text{m}}(A) \cong DR\Gamma_{\text{m}^{\text{op}}}(A) \cong A_\nu(-\ell)[d]$  in  $D(\text{GrMod } A^e)$  with some graded algebra automorphism  $\nu \in \text{GrAut } A$ . This graded algebra automorphism  $\nu \in \text{GrAut } A$  is called the generalized Nakayama automorphism. We define  $\omega_A := A_\nu(-\ell) \in \text{GrMod } A^e$ .

**Definition 2.4.** Let  $A$  be an AS-Gorenstein algebra of dimension  $d$  and  $M \in \text{grmod } A$ . Then  $M$  is called a graded maximal Cohen–Macaulay module if  $\text{depth } M = \text{ldim } M = d$ .

Let  $A$  be an AS-Gorenstein algebra. Then  $A$  is a graded maximal Cohen–Macaulay  $A$ -module. It is well known that  $M \in \text{grmod } A$  is graded maximal Cohen–Macaulay if and only if  $\underline{\text{Ext}}_A^i(M, A) = 0$  for all  $i \neq 0$ . See [Mori 2003] for basic properties of maximal Cohen–Macaulay modules.

We write  $\text{CM}^{\text{gr}}(A)$  for the full subcategory of  $\text{grmod } A$  consisting of graded maximal Cohen–Macaulay modules. If  $M \in \text{grmod } A$ , then we define  $M^\dagger = \underline{\text{Hom}}_A(M, A) \in \text{grmod } A^{\text{op}}$ . Similarly, if  $N \in \text{grmod } A^{\text{op}}$ , then we define  $N^\dagger = \underline{\text{Hom}}_{A^{\text{op}}}(N, A) \in \text{grmod } A$ . It is well known that the contravariant functors

$$(2-1) \quad \text{CM}^{\text{gr}}(A) \xrightleftharpoons[\text{(-)}^\dagger]{\text{(-)}^\dagger} \text{CM}^{\text{gr}}(A^{\text{op}})$$

define a duality. For  $M \in \text{CM}^{\text{gr}}(A)$ , we put  $B = \underline{\text{End}}_A(M)$ ,  $C = \underline{\text{End}}_{A^{\text{op}}}(M^\dagger)$ . Then

$$(2-2) \quad C \cong \underline{\text{End}}_{A^{\text{op}}}(M^\dagger) \cong \underline{\text{End}}_A(M)^{\text{op}} \cong B^{\text{op}}$$

as graded algebras by the above duality.

The following theorem, called the maximal Cohen–Macaulay approximation theorem, plays a key role in this paper.

**Theorem 2.5.** *Let  $A$  be an AS-Gorenstein algebra. For any  $M \in \text{grmod } A$ , there exists a short exact sequence*

$$0 \rightarrow L \rightarrow Z \rightarrow M \rightarrow 0$$

*in  $\text{grmod } A$  such that  $Z \in \text{CM}^{\text{gr}}(A)$  and  $\text{injdim}_A L < \infty$ . Moreover,  $\underline{\text{Ext}}_A^i(X, L) = 0$  holds for any  $X \in \text{CM}^{\text{gr}}(A)$  and any  $i \geq 1$ .*

*Proof.* This follows from [Mori 2003, Proposition 5.3] and [Ueyama 2015, Lemma 3.5].  $\square$

Recently, Minamoto and Mori [2011] introduced the two notions of an  $\mathbb{N}$ -graded (not necessarily connected graded) AS-regular algebra.

**Definition 2.6** [Minamoto and Mori 2011, Definition 3.1]. A locally finite  $\mathbb{N}$ -graded algebra  $B$  is called AS-regular over  $R = B_0$  of dimension  $d$  and of Gorenstein parameter  $\ell$  if

- $\text{gldim } B = \text{gldim } B^{\text{op}} = d < \infty$ , and
- $\text{RHom}_B(B_0, B) \cong \text{RHom}_{B^{\text{op}}}(B_0, B) \cong (DB_0)(\ell)[-d]$  in  $\text{D}(\text{GrMod } B_0)$  and in  $\text{D}(\text{GrMod } B_0^{\text{op}})$ .

**Remark 2.7.** For the purpose of this paper, we do not require  $\text{gldim } B_0 < \infty$ .

**Definition 2.8** [Minamoto and Mori 2011, Definition 3.9]. A locally finite  $\mathbb{N}$ -graded algebra  $B$  is called ASF-regular of dimension  $d$  and of Gorenstein parameter  $\ell$  if

- $\text{gldim } B = \text{gldim } B^{\text{op}} = d < \infty$ , and
- $\text{R}\Gamma_{\text{m}}(B) \cong \text{R}\Gamma_{\text{m}^{\text{op}}}(B) \cong (DB)(\ell)[-d]$  in  $\text{D}(\text{GrMod } B)$  and in  $\text{D}(\text{GrMod } B^{\text{op}})$ .

By expanding the notion of an AS-regular algebra to  $\mathbb{N}$ -graded algebras, Minamoto and Mori [2011] gave a nice correspondence between  $\mathbb{N}$ -graded AS-regular algebras over  $R$  of dimension  $d$  with  $\text{gldim } R < \infty$  and quasi-Fano algebras of global dimension  $d - 1$ . This result provides a strong connection between noncommutative projective geometry and representation theory of finite-dimensional algebras. See [Herschend et al. 2014; Minamoto and Mori 2011; Mori 2015] for details.

At the end of this section, we give a comparison theorem for the two notions of AS-regular algebras.

**Lemma 2.9.** *Let  $B$  be a two-sided noetherian ASF-regular algebra, and  $C$  a two-sided noetherian locally finite  $\mathbb{N}$ -graded algebra. Then for any  $M \in \text{GrMod } C^{\text{op}} \otimes B$ ,*

$$D \text{R}\Gamma_{\text{m}}(M) \cong \text{RHom}_B(M, D \text{R}\Gamma_{\text{m}}(B))$$

*in  $\text{D}(\text{GrMod } B^{\text{op}} \otimes C)$ .*

*Proof.* Van den Bergh [1997] gave a theory on local duality for connected graded algebras. One can check that the results in [Van den Bergh 1997, Sections 3–6] hold with no essential change for a noetherian locally finite  $\mathbb{N}$ -graded algebra (see also [Reyes et al. 2014, Remark 3.6; 2017, Lemma 3.2 (1)]). This follows from the locally finite  $\mathbb{N}$ -graded version of [Van den Bergh 1997, Theorem 5.1].  $\square$

**Theorem 2.10.** *If  $B$  is a two-sided noetherian ASF-regular algebra of dimension  $d$  and of Gorenstein parameter  $\ell$ , then  $B$  is an AS-regular algebra over  $B_0$  of dimension  $d$  and of Gorenstein parameter  $\ell$ .*



*Proof.* There exists an algebra automorphism  $\mu \in \text{GrAut } B$  such that  $D \text{R}\Gamma_{\text{m}}(B) \cong B_{\mu}(-\ell)[d]$  in  $D(\text{GrMod } B^e)$ , so we have

$$\begin{aligned} \text{R}\underline{\text{Hom}}_B(B_0, B) &\cong \text{R}\underline{\text{Hom}}_B(B_0, B_{\mu}(-\ell)[d])_{\mu^{-1}(\ell)[-d]} \\ &\cong \text{R}\underline{\text{Hom}}_B(B_0, D \text{R}\Gamma_{\text{m}}(B))_{\mu^{-1}(\ell)[-d]} \\ &\cong D \text{R}\Gamma_{\text{m}}(B_0)_{\mu^{-1}(\ell)[-d]} \\ &\cong D(B_0)_{\mu^{-1}(\ell)[-d]} \end{aligned}$$

in  $D(\text{GrMod } B^{\text{op}} \otimes B_0)$ , and so  $\text{R}\underline{\text{Hom}}_B(B_0, B) \cong D(B_0)(\ell)[-d]$  in  $D(\text{GrMod } B_0)$  and in  $D(\text{GrMod } B_0^{\text{op}})$ . Hence the result follows.  $\square$

**Corollary 2.11.** *Let  $B$  be a two-sided noetherian locally finite  $\mathbb{N}$ -graded algebra with  $\text{gldim } B_0 < \infty$ . Then  $B$  is ASF-regular if and only if it is AS-regular over  $B_0$ .*

*Proof.* This is a combination of Theorem 2.10 and [Minamoto and Mori 2011, Theorem 3.12].  $\square$

### 3. Main result

In this section, we prove the main result (Theorem 3.10) and give an example of its use. First we introduce a condition which we require for the main result.

**Definition 3.1.** Let  $A$  be an AS-Gorenstein algebra with the generalized Nakayama automorphism  $\nu \in \text{GrAut } A$ . Then  $X \in \text{GrMod } A$  is called  $\nu$ -stable if  $X_{\nu} \cong X$  as graded right  $A$ -modules.

Since  $A_{\nu} \cong A$  in  $\text{GrMod } A$ ,  $A$  is always  $\nu$ -stable. Clearly, if  $A$  is symmetric, that is, the generalized Nakayama automorphism of  $A$  is the identity, then every  $M \in \text{GrMod } A$  is  $\nu$ -stable.

**Example 3.2.** Let  $S$  be an AS-regular algebra and  $G$  a finite subgroup of  $\text{GrAut } S$  such that  $\text{char } k$  does not divide  $|G|$ . If  $S^G$  is AS-Gorenstein, then  $S \in \text{GrMod } S^G$  is  $\nu$ -stable by [Ueyama 2013, Lemma 5.8].

**Lemma 3.3.** *Let  $A$  be an AS-Gorenstein algebra and let  $X \in \text{CM}^{\text{gr}}(A)$ . If  $X$  is  $\nu$ -stable, then  $X \cong X_{\nu^{-1}}$  in  $\text{GrMod } A$  and  $X^{\dagger}$  is  $\nu$ -stable in  $\text{GrMod } A^{\text{op}}$ .*

*Proof.* It is easy to check that  $X \cong X_{\nu\nu^{-1}} \cong X_{\nu^{-1}}$  in  $\text{GrMod } A$ , and

$$\nu(X^{\dagger}) \cong \underline{\text{Hom}}_A(X, {}_{\nu}A) \cong \underline{\text{Hom}}_A(X, A_{\nu^{-1}}) \cong \underline{\text{Hom}}_A(X_{\nu}, A) \cong \underline{\text{Hom}}_A(X, A) \cong X^{\dagger}$$

in  $\text{GrMod } A^{\text{op}}$ .  $\square$

The notion of an  $n$ -cluster tilting module plays an important role in representation theory of orders, especially higher-dimensional analogues of Auslander–Reiten theory. It can be regarded as a natural generalization of the classical notion of Cohen–Macaulay representation-finiteness.

**Definition 3.4.** Let  $A$  be an AS-Gorenstein algebra. For a positive integer  $n \in \mathbb{N}^+$ , a graded maximal Cohen–Macaulay module  $X \in \text{CM}^{\text{gr}}(A)$  is called  $n$ -cluster tilting if

$$\begin{aligned} \text{add}\{X(i) \mid i \in \mathbb{Z}\} &= \{M \in \text{CM}^{\text{gr}}(A) \mid \underline{\text{Ext}}_A^i(X, M) = 0 \text{ for } 0 < i < n\} \\ &= \{M \in \text{CM}^{\text{gr}}(A) \mid \underline{\text{Ext}}_A^i(M, X) = 0 \text{ for } 0 < i < n\}, \end{aligned}$$

where  $\text{add}\{X(i) \mid i \in \mathbb{Z}\}$  is the full subcategory of  $\text{grmod } A$  consisting of direct summands of finite direct sums of shifts of  $X$ .

Let  $A$  be a noetherian locally finite  $\mathbb{N}$ -graded algebra. Then it is known that  $\text{grmod } A$  has the Krull–Schmidt property, i.e., each finitely generated graded module is a direct sum of a uniquely determined set of indecomposable graded modules. Recall that  $M \in \text{grmod } A$  is called basic if each indecomposable direct summand occurs exactly once (up to isomorphism and degree shift of grading) in a direct sum decomposition. The following proposition says that if  $A$  has a  $(d-1)$ -cluster tilting module and  $\text{gldim}(\text{tails } A) < \infty$ , then it has a  $\nu$ -stable  $(d-1)$ -cluster tilting module.

**Proposition 3.5.** *Let  $A$  be an AS-Gorenstein algebra of dimension  $d \geq 2$ , Gorenstein parameter  $\ell$ , and  $\text{gldim}(\text{tails } A) < \infty$ . If  $X \in \text{CM}^{\text{gr}}(A)$  is a basic  $(d-1)$ -cluster tilting module, then  $X$  is  $\nu$ -stable.*

*Proof.* By [Ueyama 2013, Corollary 4.5], we see that the stable category  $\underline{\text{CM}}^{\text{gr}}(A)$  has the Serre functor  $-\otimes_A A_\nu(-\ell)[d-1]$ . Since

$$\underline{\text{Ext}}_A^i(X, X) \cong \bigoplus_{s \in \mathbb{Z}} \text{Hom}_{\underline{\text{CM}}^{\text{gr}}(A)}(X, X(s)[i]) = 0$$

for any  $0 < i < d-1$ , we have  $\underline{\text{Ext}}_A^i(X, X_\nu) = 0$  for any  $0 < i < d-1$  by using the Serre functor of  $\underline{\text{CM}}^{\text{gr}}(A)$ , so  $X_\nu \in \text{add}\{X(i) \mid i \in \mathbb{Z}\}$ . Since  $X$  is basic,  $X_\nu$  is also basic, so it follows that  $X_\nu$  is a direct summand of  $X$ . Similarly, we can show that  $X_{\nu^{-1}}$  is a direct summand of  $X$ , so  $X$  is a direct summand of  $X_\nu$ . Hence the result follows.  $\square$

Next we prepare some lemmas which we need to prove the main result.

**Lemma 3.6.** *Let  $A$  be a graded algebra and  $X$  a graded right  $A$ -module containing  $A$  as a direct summand. Then  $X$  is a finitely generated graded left projective module over  $\underline{\text{End}}_A(X)$ . Moreover, for any  $M \in \text{GrMod } A$ , there is a natural isomorphism*

$$M \cong \underline{\text{Hom}}_A(X, M) \otimes_{\underline{\text{End}}_A(X)} X$$

in  $\text{GrMod } A$ .

*Proof.* Put  $B := \underline{\text{End}}_A(X)$ . Since  $X \cong A \oplus Y$  for some  $Y \in \text{GrMod } A$ , we have

$$B = \underline{\text{Hom}}_A(X, X) \cong \underline{\text{Hom}}_A(A, X) \oplus \underline{\text{Hom}}_A(Y, X) \cong X \oplus \underline{\text{Hom}}_A(Y, X)$$

in  $\text{GrMod } B^{\text{op}}$ , so  $X$  is finitely generated graded left projective over  $B$ . Moreover, one can verify that  $\underline{\text{End}}_{B^{\text{op}}}(X) \cong A$  as graded algebras. Thus, for any  $M \in \text{GrMod } A$ , we have

$$\underline{\text{Hom}}_A(X, M) \otimes_B X \cong \underline{\text{Hom}}_A(\underline{\text{End}}_{B^{\text{op}}}(X), M) \cong \underline{\text{Hom}}_A(A, M) \cong M$$

as graded right  $A$ -modules.  $\square$

**Lemma 3.7.** *Let  $A$  be an AS-Gorenstein algebra, and  $X \in \text{CM}^{\text{gr}}(A)$  such that  $X$  contains  $A$  as a direct summand. If  $\underline{\text{End}}_A(X)$  is right noetherian and  $M \in \text{grmod } A$ , then  $\underline{\text{Hom}}_A(X, M)$  is a finitely generated graded right  $\underline{\text{End}}_A(X)$ -module.*

*Proof.* Put  $B := \underline{\text{End}}_A(X)$ . By Theorem 2.5, we have an exact sequence

$$0 \rightarrow \underline{\text{Hom}}_A(X, L) \rightarrow \underline{\text{Hom}}_A(X, Z) \rightarrow \underline{\text{Hom}}_A(X, M) \rightarrow 0$$

in  $\text{GrMod } B$ , where  $Z \in \text{CM}^{\text{gr}}(A)$  and  $\text{injdim}_A L < \infty$ , so it is enough to show that  $\underline{\text{Hom}}_A(X, Z)$  is finitely generated. Since  $A$  is AS-Gorenstein and  $Z \in \text{CM}^{\text{gr}}(A)$ , we have a graded monomorphism  $Z \rightarrow F$  in  $\text{GrMod } A$ , where  $F$  is a finitely generated free  $A$ -module. Now  $A$  is a direct summand of  $X$ , so there exist graded monomorphisms  $Z \rightarrow \hat{X}$  in  $\text{GrMod } A$  and  $\underline{\text{Hom}}_A(X, Z) \rightarrow \underline{\text{Hom}}_A(X, \hat{X})$  in  $\text{GrMod } B$ , where  $\hat{X}$  is a finite direct sum of shifts of  $X$ . Since  $\underline{\text{Hom}}_A(X, \hat{X})$  is finitely generated and  $B$  is right noetherian,  $\underline{\text{Hom}}_A(X, Z)$  is also finitely generated.  $\square$

**Lemma 3.8.** *Let  $B$  be a two-sided noetherian locally finite  $\mathbb{N}$ -graded algebra. For any  $M \in \text{GrMod } B^e$ ,*

$$\text{R}\Gamma_{\text{m}}(\text{R}\Gamma_{\text{m}^{\text{op}}}(M)) \cong \text{R}\Gamma_{\text{m}^{\text{op}}}(\text{R}\Gamma_{\text{m}}(M))$$

*in  $\text{D}(\text{GrMod } B^e)$ .*

*Proof.* One can show the locally finite  $\mathbb{N}$ -graded version of [Van den Bergh 1997, Lemma 4.5], so the assertion holds.  $\square$

**Lemma 3.9.** *Let  $B$  be a two-sided noetherian ASF-regular algebra with an idempotent  $e$ . If  $M \in \text{grmod } B$  is finite-dimensional over  $k$  such that  $Me = 0$ , then  $\underline{\text{Ext}}_B^i(M, eB) = 0$  for any  $i$ .*

*Proof.* Since  $B$  is ASF-regular, we have

$$\begin{aligned} \text{R}\underline{\text{Hom}}_B(M, eB) &\cong e \text{R}\underline{\text{Hom}}_B(M, D \text{R}\Gamma_{\text{m}}(B)(\ell)[-d]) \\ &\cong e \text{R}\underline{\text{Hom}}_B(M, D \text{R}\Gamma_{\text{m}}(B))(\ell)[-d]. \end{aligned}$$

Moreover, by Lemma 2.9,

$$e \text{R}\underline{\text{Hom}}_B(M, D \text{R}\Gamma_{\text{m}}(B))(\ell)[-d] \cong e D \text{R}\Gamma_{\text{m}}(M)(\ell)[-d].$$

Since  $M \in \text{grmod } B$  is finite-dimensional over  $k$ ,  $R\Gamma_{\mathfrak{m}}(M) \cong M$ , so we have  $\underline{\text{Ext}}_B^i(M, eB) \cong eD \underline{H}_{\mathfrak{m}}^{d-i}(M)(\ell) = 0$  for all  $i \neq d$ , and  $\underline{\text{Ext}}_B^d(M, eB) \cong eD(M)(\ell) \cong D(Me)(\ell) = 0$  because  $Me = 0$ .  $\square$

Now we are ready to prove the main result of this paper.

**Theorem 3.10.** *Let  $A$  be a two-sided noetherian connected graded algebra satisfying  $\chi$  on both sides, and let  $X \in \text{grmod } A$ . We consider the following conditions:*

(A)  *$A$  and  $X$  satisfy*

(A1)  *$A$  is an AS-Gorenstein algebra of dimension  $d \geq 2$ ,*

(A2)  *$X$  is a  $(d-1)$ -cluster tilting module,*

(A3)  *$\underline{\text{End}}_A(X)_{<0} = 0$ , and*

(A4)  *$\underline{\text{Ext}}_A^1(X, M)$  and  $\underline{\text{Ext}}_A^1(M, X)$  are finite-dimensional over  $k$  for any  $M \in \text{CM}^{\text{gr}}(A)$ .*

(B)  *$B := \underline{\text{End}}_A(X)$  satisfies*

(B1)  *$B$  is a two-sided noetherian ASF-regular algebra of dimension  $d \geq 2$ ,*

(B2)  *$-\otimes_B X$  induces an equivalence functor  $\text{tails } B \xrightarrow{\sim} \text{tails } A$ , and*

(B3)  *$X^\dagger \otimes_B -$  induces an equivalence functor  $\text{tails } B^{\text{op}} \xrightarrow{\sim} \text{tails } A^{\text{op}}$ .*

*Then:*

(1) *If (A) is fulfilled and  $X$  is either  $\nu$ -stable or basic, then (B) holds.*

(2) *If (B) is fulfilled and  $X$  contains  $A$  as a direct summand, then (A) holds.*

*Proof of (1) in Theorem 3.10.* Suppose that  $A$  and  $X$  satisfy (A), and  $X$  is either  $\nu$ -stable or basic. Since  $A$  is indecomposable, (A2) implies that  $X$  is a graded maximal Cohen–Macaulay  $A$ -module containing a shift of  $A$  as a direct summand. By properties of degree shifts, we may assume that  $X$  contains  $A$  as a direct summand without loss of generality.

The proof is divided into several steps:

(a)  $(\mathcal{X}, (1))$  is ample for tails  $A$ ,

(b)  $B$  is right noetherian, and (B2) holds,

(c)  $\text{gldim } B = d$ ,

(d)  $\underline{H}_{\mathfrak{m}}^i(B) = 0$  for any  $i \neq d$ ,

(e)  $(\mathcal{X}^\dagger, (1))$  is ample for tails  $A^{\text{op}}$ ,

(f)  $B$  is left noetherian, and (B3) holds,

(g)  $\text{gldim } B^{\text{op}} = d$ ,

(h)  $\underline{H}_{\mathfrak{m}^{\text{op}}}^i(B) = 0$  for any  $i \neq d$ , and

(i)  $\underline{H}_{\mathfrak{m}}^d(B) \cong \underline{H}_{\mathfrak{m}^{\text{op}}}^d(B) \cong (DB)(\ell)$  in  $\text{GrMod } B$  and in  $\text{GrMod } B^{\text{op}}$ .

*Proof of (a).* We show that  $(\mathcal{X}, (1))$  is ample for tails  $A$ . Since  $A$  is a direct summand of  $X$ , it is easy to check that the condition (Am1) is satisfied. Let  $f : \mathcal{M} \rightarrow \mathcal{N}$  be an epimorphism in tails  $A$ . It gives a short exact sequence  $0 \rightarrow \mathcal{K} \rightarrow \mathcal{M} \rightarrow \mathcal{N} \rightarrow 0$  in tails  $A$ . We take a finitely generated graded module  $K$  such that  $\pi K = \mathcal{K}$ . By Theorem 2.5, there exists an exact sequence

$$0 \rightarrow L \rightarrow Z \rightarrow K \rightarrow 0$$

in  $\text{grmod } A$  such that  $Z \in \text{CM}^{\text{gr}}(A)$  and  $\text{injdim}_A L < \infty$ . Furthermore, Theorem 2.5 also implies that  $\underline{\text{Ext}}_A^1(X, K) \cong \underline{\text{Ext}}_A^1(X, Z)$ , so  $\underline{\text{Ext}}_A^1(X, K)$  is finite-dimensional over  $k$  by (A4). By [Artin and Zhang 1994, Corollary 7.3 (2)], it follows that  $\underline{\text{Ext}}_A^1(\mathcal{X}, \mathcal{K})$  is right bounded; thus we have  $\text{Ext}_A^1(\mathcal{X}(-n), \mathcal{K}) = \underline{\text{Ext}}_A^1(\mathcal{X}, \mathcal{K})_n = 0$  for all  $n \gg 0$ . Since

$$\text{Hom}_A(\mathcal{X}(-n), \mathcal{M}) \rightarrow \text{Hom}_A(\mathcal{X}(-n), \mathcal{N}) \rightarrow \text{Ext}_A^1(\mathcal{X}(-n), \mathcal{K})$$

is exact,  $\text{Hom}_A(\mathcal{X}(-n), \mathcal{M}) \rightarrow \text{Hom}_A(\mathcal{X}(-n), \mathcal{N})$  is surjective for all  $n \gg 0$ , so the condition (Am2) is also satisfied; hence  $(\mathcal{X}, (1))$  is ample for tails  $A$ .

*Proof of (b).* We show that  $B$  is right noetherian and (B2) is satisfied. The basic idea of the proof comes from [Mori and Ueyama 2016a, Section 2]. Since  $\text{depth}_A X = d \geq 2$ , we have

$$B = \underline{\text{End}}_A(X) = B(\text{grmod } A, X, (1)) \cong B(\text{tails } A, \mathcal{X}, (1))$$

by [Mori 2013, Lemma 3.3]. By using (A3) and Theorem 2.2, it follows that  $B = B_{\geq 0} \cong B(\text{tails } A, \mathcal{X}, (1))_{\geq 0}$  is right noetherian locally finite. Moreover, the functor

$$F := \pi \circ \underline{\text{Hom}}_A(\mathcal{X}, -) : \text{tails } A \rightarrow \text{tails } B$$

is an equivalence. For  $M \in \text{grmod } A$ , there exists  $n \in \mathbb{Z}$  such that

$$\underline{\text{Hom}}_A(\mathcal{X}, \mathcal{M})_{\geq n} \cong \underline{\text{Hom}}_A(X, M)_{\geq n}$$

in  $\text{grmod } B$  by [Artin and Zhang 1994, Corollary 7.3 (2)], so the functor  $F$  is induced by the functor  $\underline{\text{Hom}}_A(X, -) : \text{grmod } A \rightarrow \text{grmod } B$ . By using Lemma 3.6, we see that the functor  $\text{tails } B \rightarrow \text{tails } A$  induced by  $- \otimes_B X : \text{grmod } B \rightarrow \text{grmod } A$  is an equivalence functor quasi-inverse to  $F$ .

*Proof of (c).* Here we show that  $\text{gldim } B = d$ . First let us explain that we can construct a graded right  $\text{add}\{X(i) \mid i \in \mathbb{Z}\}$ -approximation of  $M \in \text{grmod } A$ . Since  $\underline{\text{Hom}}_A(X, M)$  is a finitely generated graded right  $B$ -module by Lemma 3.7, we can take  $f_i \in \text{Hom}_{\text{GrMod } A}(X(s_i), M)$  such that  $f_1, \dots, f_n$  generate  $\underline{\text{Hom}}_A(X, M)$ . Thus for any  $f \in \text{Hom}_{\text{GrMod } A}(X(t), M)$ , there exist graded homomorphisms  $g_i \in$

$\text{Hom}_{\text{GrMod } A}(X(t), X(s_i))$  such that

$$\begin{array}{ccc} X(s_1) \oplus \cdots \oplus X(s_n) & \xrightarrow{(f_1 \ \cdots \ f_n)} & M \\ & \nwarrow \quad \nearrow f & \\ & X(t) & \end{array}$$

$\begin{pmatrix} g_1 \\ \vdots \\ g_n \end{pmatrix}$

commutes. We can check that  $\phi := (f_1 \ \cdots \ f_n) : \bigoplus_{i=1}^n X(s_i) \rightarrow M$  is surjective, and

$$\underline{\text{Hom}}_A\left(X, \bigoplus_{i=1}^n X(s_i)\right) \xrightarrow{\underline{\text{Hom}}(X, \phi)} \underline{\text{Hom}}_A(X, M)$$

is also surjective.

By the above arguments, the proof of  $\text{gldim } B \leq d$  is along the same lines as that of [Dao and Huneke 2013, Theorem 3.6] (see also [Ueyama 2013, Theorem 5.10]). Let  $N \in \text{grmod } B$  and take a projective presentation  $P_1 \rightarrow P_0 \rightarrow N \rightarrow 0$ . Since we can write  $P_i \cong \underline{\text{Hom}}_A(X, X_i)$ , where  $X_i \in \text{add}\{X(i) \mid i \in \mathbb{Z}\}$  for each  $i$ , we have an exact sequence

$$0 \rightarrow \underline{\text{Hom}}_A(X, M_1) \rightarrow \underline{\text{Hom}}_A(X, X_1) \rightarrow \underline{\text{Hom}}_A(X, X_0) \rightarrow N \rightarrow 0$$

in  $\text{grmod } B$  such that  $0 \rightarrow M_1 \rightarrow X_1 \rightarrow X_0$  is exact in  $\text{grmod } A$ . By using a graded right  $\text{add}\{X(i) \mid i \in \mathbb{Z}\}$ -approximation, we have a module  $X_2 \in \text{add}\{X(i) \mid i \in \mathbb{Z}\}$  and a surjection  $X_2 \rightarrow M_1$  such that  $\underline{\text{Hom}}_A(X, X_2) \rightarrow \underline{\text{Hom}}_A(X, M_1)$  is also surjective. Let  $M_2$  be the kernel of  $X_2 \rightarrow M_1$ . Then it is easy to see that  $\underline{\text{Ext}}_A^1(X, M_2) = 0$ . Continuing in this way inductively, we can make exact sequences

$$0 \rightarrow M_{d-1} \rightarrow X_{d-1} \rightarrow \cdots \rightarrow X_2 \rightarrow X_1 \xrightarrow{\xi} X_0 \rightarrow \text{Coker } \xi \rightarrow 0$$

in  $\text{grmod } A$ , and

$$0 \rightarrow \underline{\text{Hom}}_A(X, M_{d-1}) \rightarrow P_{d-1} \rightarrow \cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow N \rightarrow 0$$

in  $\text{grmod } B$ , where  $P_i = \underline{\text{Hom}}_A(X, X_i)$ . Furthermore we see  $\underline{\text{Ext}}_A^j(X, M_i) = 0$  for any  $2 \leq i \leq d-1$  and any  $0 < j < i$ . If  $M_{d-1} = 0$ , then clearly  $\text{projdim}_B N \leq d-1$ . We now consider the case  $M_{d-1} \neq 0$ . Since  $X_i \in \text{CM}^{\text{gr}}(A)$ , it follows from the depth lemma (see [Bruns and Herzog 1998, Proposition 1.2.9]) that  $M_{d-1}$  is graded maximal Cohen–Macaulay. Moreover, since  $\underline{\text{Ext}}_A^j(X, M_{d-1}) = 0$  for  $0 < j < d-1$ , it follows that  $M_{d-1} \in \text{add}\{X(i) \mid i \in \mathbb{Z}\}$  by (A2). Thus we obtain  $\text{projdim}_B N \leq d$ .

To see that  $\text{gldim } B = d$ , consider a graded projective resolution of  $\underline{\text{Hom}}_A(X, k)$  in  $\text{grmod } B$ , namely,

$$\cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow \underline{\text{Hom}}_A(X, k) \rightarrow 0.$$

Applying  $-\otimes_B X$  to the above exact sequence, we have an exact sequence

$$\cdots \rightarrow P_2 \otimes_B X \rightarrow P_1 \otimes_B X \rightarrow P_0 \otimes_B X \rightarrow \underline{\mathrm{Hom}}_A(X, k) \otimes_B X \rightarrow 0$$

in  $\mathrm{grmod} A$ . By Lemma 3.6,  $\underline{\mathrm{Hom}}_A(X, k) \otimes_B X \cong k$ . Since each  $P_i \otimes_B X$  is a direct summand of finite direct sums of shifts of  $X$ , it is graded maximal Cohen–Macaulay, so  $\mathrm{projdim}_B \underline{\mathrm{Hom}}_A(X, k) \leq d-1$  gives a contradiction to the fact that  $\mathrm{depth}_A k = 0$ . Thus  $\mathrm{gldim} B = d$ .

*Proof of (d).* We next show that  $\underline{H}_m^i(B) = 0$  for  $i \neq d$ . By the arguments in the proof of (a) and (b), we see that

$$\begin{array}{ccc} (\mathrm{grmod} A, X, (1)) & \xrightarrow{\pi} & (\mathrm{tails} A, \mathcal{X}, (1)) \\ \uparrow -\otimes_B X & & \cong \downarrow F \\ (\mathrm{grmod} B, B, (1)) & \xrightarrow{\pi} & (\mathrm{tails} B, \mathcal{B}, (1)) \end{array}$$

commutes, and the graded algebra homomorphism

$$\underline{\mathrm{End}}_B(B) = B(\mathrm{grmod} B, B, (1)) \rightarrow B(\mathrm{tails} B, \mathcal{B}, (1)) = \underline{\mathrm{End}}_{\mathcal{B}}(\mathcal{B})$$

induced by the natural functor  $\pi$  is an isomorphism. This says that the natural map  $\varphi$  appearing in the exact sequence

$$0 \rightarrow \underline{H}_m^0(B) \rightarrow \underline{\mathrm{End}}_B(B) \xrightarrow{\varphi} \underline{\mathrm{End}}_{\mathcal{B}}(\mathcal{B}) \rightarrow \underline{H}_m^1(B) \rightarrow 0$$

in [Artin and Zhang 1994, Proposition 7.2 (2)] is an isomorphism. Thus  $\underline{H}_m^0(B) = \underline{H}_m^1(B) = 0$ .

Since we already have the equivalence functor in (B2), it follows that

$$(3-1) \quad \underline{\mathrm{Ext}}_{\mathcal{B}}^i(\mathcal{B}, \mathcal{B}) \cong \underline{\mathrm{Ext}}_{\mathcal{A}}^i(\mathcal{X}, \mathcal{X})$$

for any  $i$ . Using  $\mathrm{depth}_A X = d$  and (A2), we have  $\underline{\mathrm{Ext}}_{\mathcal{A}}^i(\mathcal{X}, \mathcal{X}) \cong \underline{\mathrm{Ext}}_{\mathcal{A}}^i(X, X) = 0$  for any  $1 \leq i \leq d-2$ , so  $\underline{\mathrm{Ext}}_{\mathcal{B}}^i(\mathcal{B}, \mathcal{B}) = 0$  for any  $1 \leq i \leq d-2$ . Thus  $\underline{H}_m^i(B) = 0$  for  $2 \leq i \leq d-1$  by [Artin and Zhang 1994, Theorem 7.2 (2)]. Furthermore  $B$  has global dimension  $d$  by (c), so  $\underline{H}_m^i(B) = 0$  for  $i \geq d+1$ .

*Proof of (e).* By the duality (2-1), we see that (A) is equivalent to the following:

(A<sup>op</sup>)  $A^{\mathrm{op}}$  and  $X^\dagger$  satisfy

(A1<sup>op</sup>)  $A^{\mathrm{op}}$  is an AS-Gorenstein algebra of dimension  $d \geq 2$ ,

(A2<sup>op</sup>)  $X^\dagger$  is a  $(d-1)$ -cluster tilting module,

(A3<sup>op</sup>)  $\underline{\mathrm{End}}_{A^{\mathrm{op}}}(X^\dagger)_{<0} = 0$ , and

(A4<sup>op</sup>)  $\underline{\mathrm{Ext}}_{A^{\mathrm{op}}}^1(N, X^\dagger)$  and  $\underline{\mathrm{Ext}}_{A^{\mathrm{op}}}^1(X^\dagger, N)$  are finite-dimensional over  $k$  for any  $N \in \mathrm{CM}^{\mathrm{gr}}(A^{\mathrm{op}})$ .

Hence the same argument as that in the proof of (a) implies that  $(X^\dagger, (1))$  is ample for tails  $A^{\text{op}}$ .

*Proof of (f).* Since  $(X^\dagger, (1))$  is ample for tails  $A^{\text{op}}$ , it follows from the same argument as that in the proof of (b) that  $C := \underline{\text{End}}_{A^{\text{op}}}(X^\dagger)$  is right noetherian and  $-\otimes_C X^\dagger$  induces an equivalence functor  $\text{tails } C \xrightarrow{\sim} \text{tails } A^{\text{op}}$ . However,  $B^{\text{op}} \cong C$  as graded algebras by (2-2), so we obtain that  $B$  is left noetherian and  $X^\dagger \otimes_B -$  induces an equivalence functor  $\text{tails } B^{\text{op}} \xrightarrow{\sim} \text{tails } A^{\text{op}}$ .

*Proof of (g) and (h).* By the arguments in the proofs of (e) and (f), the proof of (g) (resp. (h)) is obtained in the same way as that of (c) (resp. (d)).

*Proof of (i).* By (b) and (c),  $\text{tails } A \cong \text{tails } B$  and  $\text{gldim } B < \infty$ , so it follows that tails  $A$  has finite global dimension. Thus the derived category  $D^b(\text{tails } A)$  has the Serre functor  $-\otimes_A \omega_A[d-1] \cong -\otimes_A \mathcal{A}_v(-\ell)[d-1]$  by [de Naeghel and Van den Bergh 2004, Theorem A.4]. Moreover, if  $X$  is basic, then it is  $\nu$ -stable by Proposition 3.5. Using the equivalence in (B2) and the fact that  $X$  is  $\nu$ -stable, for any  $\mathcal{M} = \pi M \in \text{tails } B$ , we have natural isomorphisms

$$\begin{aligned}
 \underline{\text{Hom}}_B(\mathcal{M}, \mathcal{B}(-\ell)) &\cong \underline{\text{Hom}}_A(\pi(M \otimes_B X), \pi X(-\ell)) \\
 &\cong \underline{\text{Hom}}_A(\pi(M \otimes_B X_{v^{-1}}), \pi X_{v^{-1}}(-\ell)) \\
 &\cong \underline{\text{Hom}}_A(\pi(M \otimes_B X_{v^{-1}}), \pi X(-\ell)) \quad (\text{by Lemma 3.3}) \\
 &\cong D \underline{\text{Ext}}_A^{d-1}(\pi X(-\ell), \pi(M \otimes_B X_{v^{-1}} \otimes_A \mathcal{A}_v(-\ell))) \\
 &\cong D \underline{\text{Ext}}_A^{d-1}(\pi X(-\ell), \pi(M \otimes_B X)(-\ell)) \\
 &\cong D \underline{\text{Ext}}_B^{d-1}(\mathcal{B}, \mathcal{M})
 \end{aligned}$$

as graded  $k$ -vector spaces. It follows that  $\mathcal{B}(-\ell)$  is a dualizing sheaf of tails  $B$  in the sense of [Yekutieli and Zhang 1997]. By [Artin and Zhang 1994, Proposition 7.10 (3)], it is easy to check that  $\text{cd}(\text{tails } B) = d-1$ . Moreover, by Theorem 2.2, we see that  $B$  is a right noetherian locally finite graded algebra satisfying  $\chi_1$ , so it follows that

$$\text{Hom}_B(\mathcal{B}, \mathcal{B}(-\ell)) \cong D \underline{\text{Ext}}_B^{d-1}(\mathcal{B}, \mathcal{B})$$

in  $\text{GrMod } B$  by the proof of [Yekutieli and Zhang 1997, Theorem 2.3 (2)]. Hence we obtain

$$(3-2) \quad B(-\ell) \cong \text{Hom}_B(\mathcal{B}, \mathcal{B})(-\ell) \cong D \underline{\text{Ext}}_B^{d-1}(\mathcal{B}, \mathcal{B}) \cong D \underline{H}_m^d(B)$$

in  $\text{GrMod } B$  by (d). Dually, (f), (g), (h), and Lemma 3.3 imply

$$(3-3) \quad B(-\ell) \cong D \underline{H}_{m^{\text{op}}}^d(B)$$



in  $\text{GrMod } B^{\text{op}}$ . In addition, we see that  $\underline{H}_m^d(B)$  and  $\underline{H}_{m^{\text{op}}}^d(B)$  are right bounded graded  $B^e$ -modules, so it follows from Lemma 3.8 that

$$\begin{aligned}\underline{H}_{m^{\text{op}}}^d(B) &\cong \text{R}\underline{\Gamma}_m(\underline{H}_{m^{\text{op}}}^d(B)) \cong \text{R}\underline{\Gamma}_m(\underline{H}_{m^{\text{op}}}^d(B)[-d])[d] \\ &\cong \text{R}\underline{\Gamma}_m(\text{R}\underline{\Gamma}_{m^{\text{op}}}(B))[d] \cong \text{R}\underline{\Gamma}_{m^{\text{op}}}(\text{R}\underline{\Gamma}_m(B))[d] \\ &\cong \text{R}\underline{\Gamma}_{m^{\text{op}}}(\underline{H}_m^d(B)[-d])[d] \cong \text{R}\underline{\Gamma}_{m^{\text{op}}}(\underline{H}_m^d(B)) \cong \underline{H}_m^d(B)\end{aligned}$$

in  $\text{D}(\text{GrMod } B^e)$ . Therefore our assertion follows from (3-2) and (3-3).

Hence the proof of Theorem 3.10 (1) is now complete.  $\square$

*Proof of (2) in Theorem 3.10.* Suppose that  $X$  satisfies (B), and contains  $A$  as a direct summand. Clearly (A3) is satisfied by (B1).

First we show that (A1) holds. Since  $A$  is a direct summand of  $X$ , there exists an idempotent  $e \in B$  such that  $eBe \cong \underline{\text{End}}_A(A) \cong A$  as graded algebras and  $Be \cong \underline{\text{Hom}}_A(A, X) \cong X$  as graded  $B$ - $A$  bimodules. Then (B2) says that the functor  $\text{tails } B \rightarrow \text{tails } eBe$  induced by  $-\otimes_B Be$  is an equivalence, so it follows that  $B/(e)$  is finite-dimensional over  $k$  by [Mori and Ueyama 2016b, Lemma 3.17]. Let  $\rho$  be the composition map

$$Be \otimes_{eBe}^L eB \xrightarrow{\text{nat.}} Be \otimes_{eBe} eB \xrightarrow{\text{mult.}} B.$$

We have the triangle

$$Be \otimes_{eBe}^L eB \xrightarrow{\rho} B \rightarrow C \rightarrow Be \otimes_{eBe}^L eB[-1]$$

in  $\text{D}(\text{grmod } B)$ . Applying  $-\otimes_B^L Be$  implies  $C \otimes_B^L Be = 0$ . It follows that  $h^i(C)e = 0$  for all  $i$ , and thus  $\underline{\text{Ext}}_B^j(h^i(C), eB) = 0$  for all  $i, j$  by Lemma 3.9. By using a hypercohomology spectral sequence [Weibel 1994, 5.7.9], we obtain for all  $p$  that  $h^p(\text{R}\underline{\text{Hom}}_B(C, eB)) = 0$ , so  $\text{R}\underline{\text{Hom}}_B(C, eB) = 0$ . Applying  $\text{R}\underline{\text{Hom}}_B(-, eB)$  to the above triangle induces

$$\text{R}\underline{\text{Hom}}_B(Be \otimes_{eBe}^L eB, eB) \cong \text{R}\underline{\text{Hom}}_B(B, eB) \cong eB.$$

On the other hand, we have

$$\begin{aligned}\text{R}\underline{\text{Hom}}_B(Be \otimes_{eBe}^L eB, eB) &\cong \text{R}\underline{\text{Hom}}_{eBe}(Be, \text{R}\underline{\text{Hom}}_B(eB, eB)) \\ &\cong \text{R}\underline{\text{Hom}}_{eBe}(Be, eBe),\end{aligned}$$

so it follows that

$$(3-4) \quad \underline{\text{Ext}}_A^i(X, A) \cong \underline{\text{Ext}}_{eBe}^i(Be, eBe) = 0$$

for all  $i > 0$ .

Let  $M \in \text{grmod } A$ . Since  $\text{gldim } B = d$ , we can take a projective resolution

$$0 \rightarrow P_m \rightarrow \cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow M \otimes_{eBe} eB \rightarrow 0$$

in  $\text{grmod } B$  with  $m \leq d$ . Applying  $-\otimes_B Be$  to the above exact sequence, we have an exact sequence

$$0 \rightarrow P_m \otimes_B Be \rightarrow \cdots \rightarrow P_2 \otimes_B Be \rightarrow P_1 \otimes_B Be \rightarrow P_0 \otimes_B Be \rightarrow M \rightarrow 0.$$

Since each  $P_j \otimes_B Be \in \text{add}\{Be(i) \mid i \in \mathbb{Z}\} \cong \text{add}\{X(i) \mid i \in \mathbb{Z}\}$ , by (3-4), we see that  $\underline{\text{Ext}}_A^{m+1}(M, A) = 0$ . Thus  $\text{injdim}_A A \leq d$ . By (B1), (B2), and (B3),  $\text{gldim}(\text{tails } A) < \infty$  and  $\text{gldim}(\text{tails } A^{\text{op}}) < \infty$ . Moreover,  $A$  satisfies  $\chi$  on both sides, so it has a dualizing complex by [Van den Bergh 1997, Theorem 6.3]. It follows from [Dong and Wu 2009, Theorem 3.6] that  $A$  is AS-Gorenstein. If  $\text{injdim}_A A \leq d-1$ , then  $\text{gldim}(\text{tails } B) = \text{gldim}(\text{tails } A) \leq d-2$  by the Serre duality [de Naeghel and Van den Bergh 2004, Theorem A.4]. This is a contradiction to the fact that  $\underline{\text{Ext}}_B^{d-1}(B, B) \cong \underline{H}_m^d(B) \neq 0$ . Hence  $A$  is AS-Gorenstein of dimension  $d$ .

To show that (A2) holds, it is enough to show that

- (a)  $X \in \text{CM}^{\text{gr}}(A)$  and  $\underline{\text{Ext}}_A^i(X, X) = 0$  for any  $0 < i < d-1$ ,
- (b)  $M \in \text{CM}^{\text{gr}}(A)$  satisfying  $\underline{\text{Ext}}_A^i(X, M) = 0$  for any  $0 < i < d-1$  belongs to  $\text{add}\{X(i) \mid i \in \mathbb{Z}\}$ , and
- (c)  $M \in \text{CM}^{\text{gr}}(A)$  satisfying  $\underline{\text{Ext}}_A^i(M, X) = 0$  for any  $0 < i < d-1$  belongs to  $\text{add}\{X(i) \mid i \in \mathbb{Z}\}$ .

By (3-4), we see that  $X \in \text{CM}^{\text{gr}}(A)$ . By (B2), the functor  $\text{tails } B \rightarrow \text{tails } A$  induced by  $-\otimes_A X$  is an equivalence, so  $\underline{\text{Ext}}_A^i(\mathcal{X}, \mathcal{X}) \cong \underline{\text{Ext}}_B^i(B, B)$  for any  $i$ . Using  $\text{depth}_A X = d$  and [Artin and Zhang 1994, Theorem 7.2 (2)], it follows that

$$\underline{\text{Ext}}_A^i(X, X) \cong \underline{\text{Ext}}_A^i(\mathcal{X}, \mathcal{X}) \cong \underline{\text{Ext}}_B^i(B, B) \cong \underline{H}_m^{i+1}(B)$$

for any  $0 < i < d-1$ . Hence (a) holds by (B1).

We now give the proof of (b). Let  $M \in \text{CM}^{\text{gr}}(A)$  be such that  $\underline{\text{Ext}}_A^i(X, M) = 0$  for any  $0 < i < d-1$ . Since we know that  $A$  is AS-Gorenstein, taking a free resolution of  $M^\dagger$  in  $\text{grmod } A^{\text{op}}$  and applying  $(-)^\dagger$ , we have an exact sequence

$$0 \rightarrow M \rightarrow F_0 \rightarrow \cdots \rightarrow F_{d-3} \rightarrow Y \rightarrow 0$$

in  $\text{grmod } A$ , where each  $F_i$  is a graded free  $A$ -module and  $Y \in \text{CM}^{\text{gr}}(A)$ . Then we can make an exact sequence

$$0 \rightarrow \underline{\text{Hom}}_A(X, M) \rightarrow \underline{\text{Hom}}_A(X, F_0) \rightarrow \cdots \rightarrow \underline{\text{Hom}}_A(X, F_{d-3}) \rightarrow \underline{\text{Hom}}_A(X, Y) \rightarrow 0$$

in  $\text{GrMod } B$  because  $\underline{\text{Ext}}_A^i(X, M) = 0$  for  $0 < i < d-1$ . Moreover, taking a free presentation of  $Y^\dagger \in \text{CM}^{\text{gr}}(A^{\text{op}})$  and applying  $\text{Hom}_A(X, (-)^\dagger)$ , we have an exact sequence

$$0 \rightarrow \text{Hom}_A(X, Y) \rightarrow \text{Hom}_A(X, F_{d-2}) \rightarrow \text{Hom}_A(X, F_{d-1})$$

in  $\text{GrMod } B$ , where  $F_{d-2}$  and  $F_{d-1}$  are graded free  $A$ -modules. By the assumption that  $A$  is a direct summand of  $X$ , each  $\underline{\text{Hom}}_A(X, F_i)$  is a graded right projective  $B$ -module. Since any graded right  $B$ -module has projective dimension at most  $d$  by (B1), it holds that  $\text{projdim}_B \underline{\text{Hom}}_A(X, M) = 0$ . This means that  $\underline{\text{Hom}}_A(X, M)$  is finitely generated graded right projective over  $B$  by Lemma 3.7, so  $M \in \text{add}\{X(i) \mid i \in \mathbb{Z}\}$  by Lemma 3.6. Hence (b) is proved.

We next give the proof of (c). Let  $M \in \text{CM}^{\text{gr}}(A)$  be such that  $\underline{\text{Ext}}_A^i(M, X) = 0$  for any  $0 < i < d - 1$ . Then  $\underline{\text{Ext}}_{A^{\text{op}}}^i(X^\dagger, M^\dagger) = 0$  for any  $0 < i < d - 1$ . Since  $B^{\text{op}} \cong \underline{\text{End}}_{A^{\text{op}}}(X^\dagger)$  is also ASF-regular, the same method as that in the proof of (b) yields  $M^\dagger \in \text{add}\{X^\dagger(i) \mid i \in \mathbb{Z}\}$ . Thus we see  $M \in \text{add}\{X(i) \mid i \in \mathbb{Z}\}$  and we obtain (c).

In addition, since we have  $\text{gldim } B < \infty$  and  $\text{tails } B \cong \text{tails } A$  by (B1) and (B2), it follows that  $\text{gldim}(\text{tails } A) < \infty$ , so (A4) is satisfied by [Ueyama 2013, Lemma 5.7].

Hence the proof of Theorem 3.10(2) is finished.  $\square$

**Remark 3.11.** By observing the proof of Theorem 3.10, we notice that the condition (A4) in Theorem 3.10 can be replaced by the condition

$$(A4') \quad \text{gldim}(\text{tails } A) < \infty.$$

In this sense, it can be said that (A4) corresponds to the graded isolated singularity property.

Hence we obtain the following result.

**Corollary 3.12.** *Let  $A$  be an AS-Gorenstein algebra of dimension  $d \geq 2$  and of Gorenstein parameter  $\ell$ . Assume that  $\text{gldim}(\text{tails } A) < \infty$  and  $A$  has a  $(d-1)$ -cluster tilting module  $X \in \text{CM}^{\text{gr}}(A)$  satisfying  $\underline{\text{End}}_A(X)_{<0} = 0$ . Then a basic  $(d-1)$ -cluster tilting module  $Y$  can be extracted from  $X$ , and in addition  $B = \underline{\text{End}}_A(Y)$  is a two-sided noetherian AS-regular algebra over  $B_0$  of dimension  $d$  and of Gorenstein parameter  $\ell$  such that  $\text{tails } B \cong \text{tails } A$ .*

*Proof.* By the proof of Theorem 3.10 (1), we see that the Gorenstein parameter of  $B$  is given by the Gorenstein parameter of  $A$ , so the statement follows from Theorem 3.10, Remark 3.11 and Theorem 2.10.  $\square$

It is clear that (A2), (A3), (A4) and the  $\nu$ -stable assumption are conditions for a “right”  $A$ -module  $X$ . On the other hand, we see that (B1), (B2) and (B3) are “two-sided” conditions for  $B = \underline{\text{End}}_A(X)$ . Thus Theorem 3.10 (1) asserts that one-sided conditions for  $X$  imply two-sided conditions for  $B = \underline{\text{End}}_A(X)$ .

We now consider the case that  $A$  is a noncommutative quotient singularity.

**Example 3.13.** Let  $S$  be an AS-regular algebra of dimension  $d \geq 2$ . Let  $G$  be a finite subgroup of  $\text{GrAut } S$  such that  $\text{hdet } g = 1$  for each  $g \in G$  (in the sense of Jørgensen and Zhang [2000]) and  $\text{char } k$  does not divide  $|G|$ . Assume that  $S * G / (e)$  is finite-dimensional over  $k$ , where  $e = \frac{1}{|G|} \sum_{g \in G} 1 * g \in S * G$ . Then

- $S^G$  is AS-Gorenstein of dimension  $d \geq 2$  by [Jørgensen and Zhang 2000, Theorem 3.3],
- $S \in \text{CM}^{\text{gr}}(S^G)$  is a  $(d-1)$ -cluster tilting module by [Mori and Ueyama 2016a, Theorems 3.10 and 3.15],
- $\text{End}_{S^G}(S)_{<0} = (S * G)_{<0} = 0$  by [Mori and Ueyama 2016a, Theorem 3.10],
- $\text{gldim}(\text{tails } S^G) < \infty$  by [Mori and Ueyama 2016a, Theorem 3.10, Lemma 2.12],
- $S$  is a  $\nu$ -stable  $S^G$ -module by Example 3.2,

so the assumption of Theorem 3.10 (1) is satisfied. If fact, it follows from [Mori and Ueyama 2016a, Corollary 3.6, Theorem 3.10] that  $B = \text{End}_{S^G}(S)$  is a two-sided noetherian AS-regular algebra over  $B_0 \cong kG$  of dimension  $d$  such that  $\text{tails } \text{End}_{S^G}(S) \cong \text{tails } S^G$ . Hence Theorem 3.10 is regarded as a detailed version of this phenomenon.

For the rest of this paper, we construct a noncommutative quadric hypersurface which gives a concrete example of Theorem 3.10. See [Smith and Van den Bergh 2013, Section 5] and [Ueyama 2015, Section 4] for detailed information.

**Example 3.14.** In this example, we assume that  $k$  is algebraically closed of characteristic 0. Let

$$S = k\langle x, y, z \rangle / (xy + yx - z^2, xz + zx, yz + zy), \quad \deg x = \deg y = \deg z = 1.$$

Then  $S$  is a Koszul AS-regular algebra of dimension  $d_S = 3$  and Gorenstein parameter  $\ell_S = 3$  with a central regular element  $x^2 + y^2 \in S_2$ . Let

$$A = S/(x^2 + y^2).$$

Then  $A$  is a Koszul AS-Gorenstein algebra of dimension  $d_A = 2$  and Gorenstein parameter  $\ell_A = 1$ . Take  $w := x^2 \in A_2^1$  so that  $S^1 \cong A^1/(w)$ . We define a finite-dimensional algebra  $C(A)$  by  $C(A) := A^1[w^{-1}]_0$  (see [Smith and Van den Bergh 2013, Lemma 5.1]). It is easy to check that  $C(A) \cong k[t]/(t^4 - 1) \cong k^4$  as algebras. By [Ueyama 2015, Proposition 4.1],  $A$  is of finite Cohen–Macaulay representation type, so it has a 1-cluster tilting module. It follows from [Ueyama 2015, Theorem 3.4] that  $\text{gldim}(\text{tails } A) < \infty$ . By [Ueyama 2015, Proposition 4.4] and [Mori 2006, Theorem 3.8], indecomposable nonprojective maximal Cohen–Macaulay  $A$ -modules (up to isomorphism and degree shift of grading) are parametrized by points of  $\text{Proj } A^1 = \mathcal{V}(xy + z^2, x^2 - y^2) \subset \mathbb{P}^2$ . Using this, one can check that the graded  $A$ -module

$$X := A \oplus X_1 \oplus X_2 \oplus X_3 \oplus X_4$$

is a representation generator of  $\text{CM}^{\text{gr}}(A)$ , that is, a 1-cluster tilting module, where

$$\begin{aligned} X_1 &:= A/(x - y + z)A, & X_2 &:= A/(x - y - z)A, \\ X_3 &:= A/(x + y + iz)A, & X_4 &:= A/(x + y - iz)A \end{aligned}$$

and  $i$  is a primitive fourth root of unity. Clearly  $X$  is basic, so it is  $\nu$ -stable. Since generators of  $X$  are concentrated in degree 0, every graded  $A$ -module homomorphism  $X \rightarrow X(-s)$  has to be zero for any positive integer  $s \in \mathbb{N}^+$ , so we have  $\underline{\text{End}}(X)_{<0} = 0$ . Hence we obtain that  $B = \underline{\text{End}}_A(X)$  is a two-sided noetherian AS-regular algebra over  $B_0$  of dimension 2 and of Gorenstein parameter 1 satisfying  $\text{tails } B \cong \text{tails } A$  by Corollary 3.12. We can calculate that the Hilbert series  $H_B(t)$  is  $9(1+t)/(1-t)^2$ . Furthermore  $B_0$  is isomorphic to the path algebra  $kQ$ , where

$$Q = \begin{array}{ccc} & \bullet & \\ & \searrow & \swarrow \\ \bullet & & \bullet \\ & \nearrow & \nwarrow \\ & \bullet & \end{array},$$

so  $\text{gldim } B_0 < \infty$ . Hence we obtain

$$\text{D}^b(\text{tails } A) \cong \text{D}^b(\text{tails } B) \cong \text{D}^b(\text{mod } B_0) \cong \text{D}^b(\text{mod } kQ)$$

by [Minamoto and Mori 2011, Theorem 4.14]. As a remark, for any 2-dimensional commutative Cohen–Macaulay algebra  $C$  of finite Cohen–Macaulay representation type generated in degree 1, it follows that  $\text{tails } C \cong \text{tails } k[x, y] \cong \text{coh } \mathbb{P}^1$  by [Eisenbud and Herzog 1988, theorem on page 347], so  $\text{D}^b(\text{tails } C) \cong \text{D}^b(\text{coh } \mathbb{P}^1) \cong \text{D}^b(\text{mod } kQ')$ , where  $Q' = \bullet \rightrightarrows \bullet$ . Thus we see that  $\text{D}^b(\text{tails } A) \cong \text{D}^b(\text{tails } B) \not\cong \text{D}^b(\text{tails } C)$ .

### Acknowledgment

The author thanks the referee for a careful reading of the manuscript and helpful suggestions.

### References

- [Artin and Zhang 1994] M. Artin and J. J. Zhang, “Noncommutative projective schemes”, *Adv. Math.* **109**:2 (1994), 228–287. MR Zbl
- [Bruns and Herzog 1998] W. Bruns and J. Herzog, *Cohen–Macaulay rings*, 2nd ed., Cambridge Studies in Advanced Mathematics **39**, Cambridge Univ. Press, 1998. Zbl
- [Dao and Huneke 2013] H. Dao and C. Huneke, “Vanishing of Ext, cluster tilting modules and finite global dimension of endomorphism rings”, *Amer. J. Math.* **135**:2 (2013), 561–578. MR Zbl
- [Dong and Wu 2009] Z.-C. Dong and Q.-S. Wu, “Non-commutative Castelnuovo–Mumford regularity and AS-regular algebras”, *J. Algebra* **322**:1 (2009), 122–136. MR Zbl
- [Eisenbud and Herzog 1988] D. Eisenbud and J. Herzog, “The classification of homogeneous Cohen–Macaulay rings of finite representation type”, *Math. Ann.* **280**:2 (1988), 347–352. MR Zbl
- [Herschend et al. 2014] M. Herschend, O. Iyama, and S. Oppermann, “ $n$ -representation infinite algebras”, *Adv. Math.* **252** (2014), 292–342. MR Zbl
- [Iyama 2008] O. Iyama, “Auslander–Reiten theory revisited”, pp. 349–397 in *Trends in representation theory of algebras and related topics*, edited by A. Skowroński, Eur. Math. Soc., Zürich, 2008. MR Zbl
- [Jørgensen and Zhang 2000] P. Jørgensen and J. J. Zhang, “Gourmet’s guide to Gorensteinness”, *Adv. Math.* **151**:2 (2000), 313–345. MR Zbl

- [Minamoto and Mori 2011] H. Minamoto and I. Mori, “The structure of AS-Gorenstein algebras”, *Adv. Math.* **226**:5 (2011), 4061–4095. MR Zbl
- [Mori 2003] I. Mori, “Homological properties of balanced Cohen–Macaulay algebras”, *Trans. Amer. Math. Soc.* **355**:3 (2003), 1025–1042. MR Zbl
- [Mori 2006] I. Mori, “Co-point modules over Koszul algebras”, *J. London Math. Soc.* (2) **74**:3 (2006), 639–656. MR Zbl
- [Mori 2013] I. Mori, “McKay-type correspondence for AS-regular algebras”, *J. Lond. Math. Soc.* (2) **88**:1 (2013), 97–117. MR Zbl
- [Mori 2015] I. Mori, “Regular modules over 2-dimensional quantum Beilinson algebras of type  $S''$ ”, *Math. Z.* **279**:3–4 (2015), 1143–1174. MR Zbl
- [Mori and Ueyama 2016a] I. Mori and K. Ueyama, “Ample group action on AS-regular algebras and noncommutative graded isolated singularities”, *Trans. Amer. Math. Soc.* **368**:10 (2016), 7359–7383. MR Zbl
- [Mori and Ueyama 2016b] I. Mori and K. Ueyama, “Stable categories of graded maximal Cohen–Macaulay modules over noncommutative quotient singularities”, *Adv. Math.* **297** (2016), 54–92. MR Zbl
- [de Naeghel and Van den Bergh 2004] K. de Naeghel and M. Van den Bergh, “Ideal classes of three-dimensional Sklyanin algebras”, *J. Algebra* **276**:2 (2004), 515–551. MR Zbl
- [Reyes et al. 2014] M. Reyes, D. Rogalski, and J. J. Zhang, “Skew Calabi–Yau algebras and homological identities”, *Adv. Math.* **264** (2014), 308–354. MR Zbl
- [Reyes et al. 2017] M. Reyes, D. Rogalski, and J. J. Zhang, “Skew Calabi–Yau triangulated categories and Frobenius Ext-algebras”, *Trans. Amer. Math. Soc.* **369**:1 (2017), 309–340. MR Zbl
- [Smith and Van den Bergh 2013] S. P. Smith and M. Van den Bergh, “Noncommutative quadric surfaces”, *J. Noncommut. Geom.* **7**:3 (2013), 817–856. MR Zbl
- [Ueyama 2013] K. Ueyama, “Graded maximal Cohen–Macaulay modules over noncommutative graded Gorenstein isolated singularities”, *J. Algebra* **383** (2013), 85–103. MR Zbl
- [Ueyama 2015] K. Ueyama, “Noncommutative graded algebras of finite Cohen–Macaulay representation type”, *Proc. Amer. Math. Soc.* **143**:9 (2015), 3703–3715. MR Zbl
- [Van den Bergh 1997] M. Van den Bergh, “Existence theorems for dualizing complexes over noncommutative graded and filtered rings”, *J. Algebra* **195**:2 (1997), 662–679. MR Zbl
- [Weibel 1994] C. A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics **38**, Cambridge Univ. Press, 1994. MR Zbl
- [Yekutieli 1992] A. Yekutieli, “Dualizing complexes over noncommutative graded algebras”, *J. Algebra* **153**:1 (1992), 41–84. MR Zbl
- [Yekutieli and Zhang 1997] A. Yekutieli and J. J. Zhang, “Serre duality for noncommutative projective schemes”, *Proc. Amer. Math. Soc.* **125**:3 (1997), 697–707. MR Zbl

Received April 13, 2016. Revised February 6, 2017.

KENTA UHEYAMA  
 DEPARTMENT OF MATHEMATICS  
 FACULTY OF EDUCATION  
 HIROSAKI UNIVERSITY  
 1 BUNKYOCHO, HIROSAKI  
 AOMORI 036-8560  
 JAPAN  
 k-ueyama@hirosaki-u.ac.jp

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Igor Pak  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pak.pjm@gmail.com](mailto:pak.pjm@gmail.com)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 289    No. 2    August 2017

---

Regular representations of completely bounded maps	257
B. V. RAJARAMA BHAT, NIRUPAMA MALLICK and K. SUMESH	
Ball convex bodies in Minkowski spaces	287
THOMAS JAHN, HORST MARTINI and CHRISTIAN RICHTER	
Local constancy of dimension of slope subspaces of automorphic forms	317
JOACHIM MAHNKOPF	
Weakening idempotency in $K$ -theory	381
VLADIMIR MANUILOV	
On Langlands quotients of the generalized principal series isomorphic to their Aubert duals	395
IVAN MATIĆ	
Exact Lagrangian fillings of Legendrian $(2, n)$ torus links	417
YU PAN	
Elementary calculation of the cohomology rings of real Grassmann manifolds	443
RUSTAM SADYKOV	
Cluster tilting modules and noncommutative projective schemes	449
KENTA UYAMA	
Concentration for a biharmonic Schrödinger equation	469
DONG WANG	
Global existence of smooth solutions to exponential wave maps in FLRW spacetimes	489
CHANG-HUA WEI and NING-AN LAI	



0030-8730(201708)289:2;1-O