

*Pacific
Journal of
Mathematics*

**GLOBAL EXISTENCE OF SMOOTH SOLUTIONS
TO EXPONENTIAL WAVE MAPS IN FLRW SPACETIMES**

CHANG-HUA WEI AND NING-AN LAI

Volume 289 No. 2

August 2017

GLOBAL EXISTENCE OF SMOOTH SOLUTIONS TO EXPONENTIAL WAVE MAPS IN FLRW SPACETIMES

CHANG-HUA WEI AND NING-AN LAI

We consider the Cauchy problem of exponential wave maps in Friedmann–Lemaître–Robertson–Walker (FLRW) spacetimes. Using a weighted energy estimate, we show that the smooth solution of the Cauchy problem for exponential wave maps in FLRW spacetimes exists globally for small initial data.

1. Introduction

Wave maps are the Lorentzian counterparts of harmonic maps. Due to their significance in geometry and physics, wave maps have attracted much attention, and many achievements have been made in recent decades [Choquet-Bruhat 2000; Gu 1980; Klainerman and Machedon 1993; 1995; Shatah and Struwe 1998; 2002; Tao 2001a; 2001b; 2008]. In this paper, we investigate exponential wave maps which were introduced by Eells and Lemaire [1992]. The relationship between wave maps and exponential wave maps has been studied by Chiang and Yang [2007] from the geometric point of view. In this work we study the PDE aspect of the exponential wave maps on a curved Lorentzian manifold.

In this paper, we consider exponential wave maps on a special class of FLRW spacetimes, whose metric takes the form $g = -dt^2 + a^2(t) d\sigma^2$ for suitable $a(t)$. The FLRW metric is an exact solution of Einstein's field equations; it describes a homogeneous and isotropic universe. For more details on this metric, we refer to the book [Hawking and Ellis 1973]. In particular, in local coordinates (t, x_1, \dots, x_m) , with $m \geq 1$, we consider the following metric

$$(1-1) \quad g = -dt^2 + t^l \sum_{i=1}^m dx_i^2,$$

where $l > 2$ and $t > 0$. Here we remark that for the case $a(t) = e^t$, the metric g turns to be the de Sitter metric and much progress has been made on the study of wave type equations on de Sitter spacetimes; see [Kong and Wei 2013; Yagdjian 2012; 2009; Yagdjian and Galstian 2008; 2009].

MSC2010: 35L05.

Keywords: exponential wave map, FLRW, weighted energy estimate.

Before our main result, we give a brief introduction to exponential wave maps.

Basic equation and the main result.

Definition 1.1. An exponential wave map $u : M^{1+m} \rightarrow N$ from a $(1+m)$ -dimensional Lorentzian manifold $(M^{1+m}, g_{\mu\nu})$ to an n -dimensional manifold $(N, h_{\alpha\beta})$ is a critical point of the exponential energy functional

$$(1-2) \quad E(u) := \int_M \exp(|du|^2) d\text{Vol}_M = \int_M \exp(h_{\alpha\beta} \partial_\mu u^\alpha \partial_\nu u^\beta g^{\mu\nu}) d\text{Vol}_M,$$

where $d\text{Vol}_M$ is the volume element of M .

We consider the exponential wave map taking M^{1+m} as the background manifold and \mathbb{R} as the target manifold N in Definition 1.1. Therefore, the exponential energy functional takes the explicit form

$$(1-3) \quad E(u) = \int_M \exp\left(-(\partial_t u)^2 + t^{-l} \sum_{i=1}^m (\partial_i u)^2\right) t^{lm/2} dt dx_1 \cdots dx_m,$$

where $\partial_t u = \frac{\partial u}{\partial t}$ and $\partial_i u = \frac{\partial u}{\partial x_i}$; the corresponding Euler–Lagrange equation reads

$$(1-4) \quad \frac{\partial}{\partial t} \left[-2\partial_t u \exp\left(-(\partial_t u)^2 + t^{-l} \sum_{i=1}^m (\partial_i u)^2\right) t^{lm/2} \right] \\ + \sum_{i=1}^m \frac{\partial}{\partial x_i} \left[2t^{-l} \partial_i u \exp\left(-(\partial_t u)^2 + t^{-l} \sum_{i=1}^m (\partial_i u)^2\right) t^{lm/2} \right] = 0,$$

i.e., with $\Delta := \sum_{i=1}^m \partial_i^2$,

$$(1-5) \quad -\partial_t^2 u + t^{-l} \Delta u - \frac{lm}{2t} \partial_t u = \partial_t u \partial_t \left[-(\partial_t u)^2 + t^{-l} \sum_{i=1}^m (\partial_i u)^2 \right] \\ - \sum_{i=1}^m t^{-l} \partial_i u \partial_i \left[-(\partial_t u)^2 + t^{-l} \sum_{i=1}^m (\partial_i u)^2 \right].$$

Remark 1.2. When $l = 0$, (1-5) is exactly the exponential wave map on Minkowski spacetime. Huh [2013] obtained a global existence result for sufficiently small initial data.

Define the wave operator as

$$(1-6) \quad \square_g = \frac{1}{\sqrt{|\det g|}} \partial_\mu (\sqrt{|\det g|} g^{\mu\nu} \partial_\nu) = -\partial_t^2 - \frac{lm}{2t} \partial_t + t^{-l} \sum_{i=1}^m \partial_i^2,$$

where the Greek index μ ranges from 0 to m and $\partial_0 := \partial_t$. In (1-6) and throughout the paper, we use the Einstein summation convention, which says that one always takes the summation over the repeated upper and lower indices.

Let

$$(1-7) \quad Q(\varphi, \psi) = -(\partial_t \varphi)(\partial_t \psi) + t^{-l} \sum_{i=1}^m (\partial_i \varphi)(\partial_i \psi).$$

Then (1-5) can be rewritten as

$$(1-8) \quad \square_g u = -Q(u, Q(u, u)).$$

From now on, we consider (1-8) with the following initial data

$$(1-9) \quad t = 1 : \quad u(1, x) = \epsilon f(x), \quad u_t(1, x) = \epsilon q(x),$$

where ϵ is a small parameter, f and q are smooth functions with compact support.

The main result of this paper is as follows:

Theorem 1.3. *There exists a positive constant ϵ_0 , such that the Cauchy problem (1-8)-(1-9) has a unique smooth solution on $[1, +\infty) \times \mathbb{R}^m$ for any $\epsilon \in [0, \epsilon_0]$.*

The structure of the paper. The paper is organized as follows. In Section 2, we attain pointwise decay estimates of the smooth solution to the linear wave equation by weighted energy estimates. The main theorem is proved in Section 3 by the continuity method and a careful analysis of the nonlinearities. Several further discussions are presented in Section 4.

2. Decay estimates for the linear wave equation

In this section, we investigate the pointwise decay estimates of smooth solution to the following linear wave equation on FLRW spacetimes

$$(2-1) \quad \square_g \varphi = 0,$$

where the metric g is given by (1-1). It will play an important role in the study of nonlinear (1-8).

For multi-index $I = (I_1, \dots, I_m)$ with $|I| = \sum_{j=1}^m |I_j|$, denote

$$D = \{\partial_1, \dots, \partial_m\} \quad \text{and} \quad D^I = \partial_1^{I_1} \dots \partial_m^{I_m}.$$

Furthermore, $\tilde{I} \leq I$ means $\tilde{I}_i \leq I_i$ for $i = 1, \dots, m$.

For any function $\varphi(x) = \varphi(x_1, \dots, x_m)$, we define

$$\|\varphi(x)\|_{L^2} = \left(\int_{\mathbb{R}^m} |\varphi(x)|^2 dx \right)^{1/2}, \quad \|\varphi(x)\|_{L^\infty} := \text{ess sup}_{x \in \mathbb{R}^m} |\varphi(x)|,$$

and

$$\|\varphi(x)\|_{H^s} = \left(\sum_{i=0}^s (\|D^i \varphi(x)\|_{L^2})^2 \right)^{1/2},$$

where s is an integer and $dx = dx^1 \dots dx^m$.

From here on, we also use $A \lesssim B$ to denote $A \leq CB$ for some positive constant C .

Corresponding to the (2-1) and the specific metric g given by (1-1), we define the energy momentum tensor as

$$(2-2) \quad T_{\mu\nu}(\varphi) = (\partial_\mu\varphi)(\partial_\nu\varphi) - \frac{1}{2}g_{\mu\nu}|\nabla\varphi|^2,$$

where

$$|\nabla\varphi|^2 = g^{\kappa\lambda}\partial_\kappa\varphi\partial_\lambda\varphi = -(\partial_t\varphi)^2 + \sum_{i=1}^m t^{-l}(\partial_i\varphi)^2.$$

Given a vector field $V = V^\mu\partial_\mu$, define the compatible currents

$$(2-3) \quad J_\mu^V(\varphi) = T_{\mu\nu}(\varphi)V^\nu$$

and

$$(2-4) \quad K^V(\varphi) = \Pi_{\mu\nu}^V T^{\mu\nu}(\varphi),$$

where $\Pi_{\mu\nu}^V$ denotes the deformation tensor defined by

$$(2-5) \quad \Pi_{\mu\nu}^V = \frac{1}{2}(\nabla_\mu V_\nu + \nabla_\nu V_\mu),$$

and

$$\nabla_\mu V_\nu = g(\nabla_\mu V, \partial_\nu).$$

In a constant t -slice, the induced volume form is

$$(2-6) \quad d\text{Vol}_t = t^{lm/2} dx_1 \cdots dx_m.$$

With above notations, we have for $i, j = 1, \dots, m$ and $i \neq j$

$$(2-7) \quad T_{00}(\varphi) = \frac{1}{2} \left((\partial_t\varphi)^2 + \sum_{i=1}^m t^{-l} (\partial_i\varphi)^2 \right),$$

$$(2-8) \quad T_{0i} = \partial_t\varphi\partial_i\varphi, \quad T_{ij} = \partial_i\varphi\partial_j\varphi,$$

and

$$(2-9) \quad T_{ii} = \frac{1}{2} \left(t^l (\partial_t\varphi)^2 + (\partial_i\varphi)^2 - \sum_{j \neq i} (\partial_j\varphi)^2 \right).$$

We recall the following energy identity without proof; see [Alinhac 2010].

Lemma 2.1. *For a solution φ of the equation $\square_g\varphi = f$, we have*

$$(2-10) \quad \nabla^\mu T_{\mu\nu} = (\square_g\varphi)(\partial_\nu\varphi) \quad \text{and} \quad \nabla^\mu J_\mu^V(\varphi) = K^V(\varphi) + \square_g\varphi \cdot V(\varphi).$$

The energy density $e(V, \nu)$ of the function φ at time t with respect to a past pointed timelike vector field V is the nonnegative number given by

$$(2-11) \quad e(V, \nu) = J_\alpha^V \nu^\alpha = T_{\alpha\beta}(\varphi) V^\beta \nu^\alpha,$$

where ν^α is the α -th component of the past oriented unit normal vector $\nu = -\partial_t$.

Given a past pointed vector field V , by Lemma 2.1 and the divergence theorem, we easily get the following lemma:

Lemma 2.2. *Integrating (2-10) over the spacetime domain $D = \{[1, t] \times \mathbb{R}^m\}$,*

$$(2-12) \quad \int_{\Sigma_t} J_\alpha^V \nu^\alpha d\text{Vol}_t - \int_{\Sigma_1} J_\alpha^V \nu^\alpha d\text{Vol}_1 = \int_1^t \int_{\Sigma_\tau} (K^V(\varphi) + \square_g \varphi V(\varphi)) d\text{Vol}_\tau d\tau,$$

where Σ_t denotes the spacelike hypersurface with $t = \text{constant}$.

From now on, we take $V = -\partial_t$ so that

$$(2-13) \quad \Pi_{\mu\nu}^V = -\frac{1}{2}(g_{\nu\kappa} \Gamma_{\mu t}^\kappa + g_{\mu\kappa} \Gamma_{\nu t}^\kappa).$$

Then for $i = 1, \dots, m$,

$$(2-14) \quad \Pi_{ii}^V = -\frac{l}{2} t^{l-1},$$

and for $i \neq j$,

$$(2-15) \quad \Pi_{ij}^V = 0, \quad \Pi_{0i}^V = 0, \quad \text{and} \quad \Pi_{00}^V = 0,$$

where we have assumed that $x_0 = t$. By (2-4), we have

$$(2-16) \quad K^{-\partial_t}(\varphi) = \frac{l}{4} \left[(m-2) \sum_{i=1}^m \frac{1}{t^{l+1}} (\partial_i \varphi)^2 - \frac{m}{t} (\partial_t \varphi)^2 \right].$$

By (2-11), we have

$$(2-17) \quad e(V, \nu) = \frac{1}{2} \left((\partial_t \varphi)^2 + \sum_{i=1}^m t^{-l} (\partial_i \varphi)^2 \right).$$

For a constant t -slice, we define

$$(2-18) \quad \begin{cases} E_0^{l, I_0}(t) = \frac{1}{2} \int_{\Sigma_t} (\partial_t \partial_t^{I_0} D^I \varphi)^2 d\text{Vol}_t, \\ E_1^{l, I_0}(t) = \frac{1}{2} \int_{\Sigma_t} \left(\sum_{i=1}^m t^{-l} (\partial_i \partial_t^{I_0} D^I \varphi)^2 \right) d\text{Vol}_t \end{cases}$$

and

$$(2-19) \quad E^{l, I_0}(t) = E_0^{l, I_0}(t) + E_1^{l, I_0}(t),$$

where I_0 and $|I|$ are nonnegative integers, $|I|$ is the magnitude of any multi-index I .

Then, by above calculations and Lemma 2.2, the following zeroth-order energy identity holds:

Lemma 2.3. *The energy identity (2-12) can be rewritten as*

$$(2-20) \quad E^{0,0}(t) - E^{0,0}(1) = \int_1^t \left[-\frac{l}{\tau} \left(\frac{m}{2} \right) E_0^{0,0}(\tau) + \frac{l}{\tau} \left(\frac{m-2}{2} \right) E_1^{0,0}(\tau) \right] d\tau.$$

Corollary 2.4. *We have*

$$(2-21) \quad \frac{d}{dt} E^{0,0}(t) = -\frac{lm}{2t} E_0^{0,0}(t) + \frac{l(m-2)}{2t} E_1^{0,0}(t).$$

According to the explicit expression of FLRW metric (1-1), it is easy to see that the operator D is a Killing vector field, which means

$$\Pi_{\mu\nu}^D = 0,$$

where D denotes the spacial derivatives. Thus, the structure of the equation (2-1) will not change if we take D^J as a commutator, i.e.,

$$(2-22) \quad \square_g(D^J \varphi) = 0.$$

By (2-22) and Corollary 2.4, for $I_0 = 0$, we have the following corollary.

Corollary 2.5. *For arbitrary nonnegative multi-index J , we have*

$$(2-23) \quad \frac{d}{dt} E^{J,0}(t) = -\frac{lm}{2t} E_0^{J,0}(t) + \frac{(m-2)l}{2t} E_1^{J,0}(t).$$

Define

$$(2-24) \quad \begin{cases} f^{I,I_0}(t) = E^{I,I_0}(t)t^{(2-m)l/2}, \\ f_0^{I,I_0}(t) = E_0^{I,I_0}(t)t^{(2-m)l/2}, \\ f_1^{I,I_0}(t) = E_1^{I,I_0}(t)t^{(2-m)l/2}. \end{cases}$$

Then we obtain the lemma:

Lemma 2.6. *The function $f^{I,0}(t)$ is uniformly bounded, provided that $E^{I,0}(1)$ is bounded for arbitrary multi-index I .*

Proof. By (2-24), since $m \geq 1$,

$$(2-25) \quad \begin{aligned} \frac{d}{dt} f^{I,0}(t) &= \left(\frac{d}{dt} E^{I,0}(t) \right) t^{(2-m)l/2} + \frac{(2-m)l}{2} E^{I,0}(t) t^{(2-m)l/2} t^{-1} \\ &= t^{(2-m)l/2} \left[-\frac{lm}{2t} E_0^{I,0}(t) + \frac{(m-2)l}{2t} E_1^{I,0}(t) + \frac{(2-m)l}{2t} E^{I,0}(t) \right] \\ &= -\frac{l(m-1)}{t} t^{(2-m)l/2} E_0^{I,0}(t) \leq 0. \end{aligned}$$

Thus, $f^{I,0}(t)$ is monotonically decreasing and we have

$$f^{I,0}(t) \leq f^{I,0}(1) = E^{I,0}(1). \quad \square$$

Next, we consider the equation with higher-order derivatives on t .

Lemma 2.7. *For $I_0 > 0$, differentiating (2-22) by $\partial_t^{I_0}$, we have*

$$(2-26) \quad \square_g(\partial_t^{I_0} D^J \varphi) = \sum_{I'_0=1}^{I_0} C_{I'_0,l,m} t^{-(I_0-I'_0+2)} \partial_t^{I'_0} D^J \varphi + \sum_{i=1}^m \sum_{I''_0=0}^{I_0-1} C_{I''_0,l,m} t^{-l-(I_0-I''_0)} \partial_i^2 \partial_t^{I''_0} D^J \varphi,$$

where $C_{I'_0,l,m}$ and $C_{I''_0,l,m}$ are constants depending on I'_0, I''_0, l , and m .

Proof. With the notation $D^J \varphi = v$, it suffices to prove

$$(2-27) \quad \square_g(\partial_t^{I_0} v) = \sum_{I'_0=1}^{I_0} C_{I'_0,l,m} t^{-(I_0-I'_0+2)} \partial_t^{I'_0} v + \sum_{i=1}^m \sum_{I''_0=0}^{I_0-1} C_{I''_0,l,m} t^{-l-(I_0-I''_0)} \partial_i^2 \partial_t^{I''_0} v.$$

By (2-22), we know that

$$(2-28) \quad \square_g v = 0.$$

Since ∂_t is not a Killing vector field, it does not commute with the operator \square_g , and a direct calculation gives

$$(2-29) \quad [\square_g, \partial_t] = -\frac{lm}{2} t^{-2} \partial_t + lt^{-l-1} \sum_{i=1}^m \partial_i^2.$$

Now, we can prove the lemma by induction on I_0 .

When $I_0 = 1$,

$$(2-30) \quad \square_g(\partial_t v) = [\square_g, \partial_t]v + \partial_t(\square_g v) = -\frac{lm}{2} t^{-2} \partial_t v + lt^{-l-1} \sum_{i=1}^m \partial_i^2 v.$$

Thus, the lemma holds for $I_0 = 1$.

Suppose the lemma holds for $I_0 - 1$. Then, for I_0 ,

$$(2-31) \quad \begin{aligned} \square_g(\partial_t^{I_0} v) &= [\square_g, \partial_t](\partial_t^{I_0-1} v) + \partial_t(\square_g \partial_t^{I_0-1} v) \\ &= \left(-\frac{lm}{2} t^{-2} \partial_t + lt^{-l-1} \sum_{i=1}^m \partial_i^2 \right) \partial_t^{I_0-1} v \\ &\quad + \partial_t \left(\sum_{I'_0=1}^{I_0-1} C_{I'_0,l,m} t^{-(I_0-1-I'_0+2)} \partial_t^{I'_0} v \right. \\ &\quad \left. + \sum_{i=1}^m \sum_{I''_0=0}^{I_0-2} C_{I''_0,l,m} t^{-l-(I_0-1-I''_0)} \partial_i^2 \partial_t^{I''_0} v \right), \end{aligned}$$

which we rewrite as

$$\begin{aligned} \square_g(\partial_t^{I_0} v) &= -\frac{lm}{2} t^{-2} \partial_t^{I_0} v + l t^{-l-1} \sum_{i=1}^m \partial_i^2 \partial_t^{I_0-1} v \\ &\quad + \sum_{I'_0=1}^{I_0} C_{I'_0, l, m} t^{-(I_0-I'_0+2)} \partial_t^{I'_0} v + \sum_{i=1}^m \sum_{I''_0=0}^{I_0-1} C_{I''_0, l, m} t^{-l-(I_0-I''_0)} \partial_i^2 \partial_t^{I''_0} v. \end{aligned}$$

By grouping together the last two terms on the right-hand-side, we see that (2-27) holds for arbitrary I_0 . □

We define the total energy of M -th order as

$$(2-32) \quad F^M(t) = \sum_{|I|+I_0 \leq M} f^{I, I_0}(t).$$

Lemma 2.8. *We have*

$$(2-33) \quad F^M(t) \leq C_M F^M(1),$$

where C_M is a positive constant depending only on M , and $F^M(1)$ is determined by the initial data.

Proof. For arbitrary $|I| + I_0 \leq M$ and $I_0 \geq 1$, by Lemmas 2.2, 2.6, 2.7, and Corollary 2.5,

$$\begin{aligned} (2-34) \quad \frac{d}{dt} E^{I, I_0}(t) &= -\frac{lm}{2t} E_0^{I, I_0}(t) + \frac{l(m-2)}{2t} E_1^{I, I_0}(t) \\ &\quad - \int_{\Sigma_t} \left(\sum_{I'_0=1}^{I_0} C_{I'_0, l, m} t^{-(I_0-I'_0+2)} \partial_t^{I'_0} D^J \varphi \right. \\ &\quad \left. + \sum_{i=1}^m \sum_{I''_0=0}^{I_0-1} C_{I''_0, l, m} t^{-l-(I_0-I''_0)} \partial_i^2 \partial_t^{I''_0} \varphi \right) \partial_t^{I_0+1} D^I \varphi \, d \text{Vol}_t. \end{aligned}$$

Then, by Lemma 2.6

$$\begin{aligned} (2-35) \quad \frac{d}{dt} f^{I, I_0}(t) &= \left(\frac{d}{dt} E^{I, I_0}(t) \right) t^{(2-m)l/2} + \frac{(2-m)l}{2} E^{I, I_0}(t) t^{(2-m)l/2-1} \\ &\leq \left| t^{(2-m)l/2} \int_{\Sigma_t} \left(\sum_{I'_0=1}^{I_0} C_{I'_0, l, m} t^{-(I_0-I'_0+2)} \partial_t^{I'_0} D^J \varphi \right. \right. \\ &\quad \left. \left. + \sum_{i=1}^m \sum_{I''_0=0}^{I_0-1} C_{I''_0, l, m} t^{-l-(I_0-I''_0)} \partial_i^2 \partial_t^{I''_0} \varphi \right) \partial_t^{I_0+1} D^I \varphi \, d \text{Vol}_t \right|. \end{aligned}$$

By Hölder’s inequality, for arbitrary $|I| + I_0 \leq M$,

$$(2-36) \quad \frac{d}{dt} f^{I, I_0}(t) \leq C_M (t^{-(I_0-I'_0+2)} F^M(t) + t^{-l/2-(I_0-I''_0)} F^M(t)),$$

where $1 \leq I'_0 \leq I_0$, $0 \leq I''_0 \leq I_0 - 1$, which is given by Lemma 2.7.

Summing over all $|I| + I_0 \leq M$ and using (2-36),

$$(2-37) \quad \frac{d}{dt} F^M(t) \leq C_M (t^{-(I_0 - I'_0 + 2)} F^M(t) + t^{-l/2 - (I_0 - I''_0)} F^M(t)).$$

By virtue of (2-37), one gets

$$(2-38) \quad F^M(t) \leq F^M(1) \exp\left(\int_1^t C_M (t^{-(I_0 - I'_0 + 2)} + t^{-l/2 - (I_0 - I''_0)}) dt\right) \leq C_M F^M(1),$$

where for the second inequality we use the facts that $I_0 - I'_0 + 2 \geq 2$, $I_0 - I''_0 \geq 1$, and $l > 0$. □

Remark 2.9. The constant C_M may vary from line to line in the above proof.

We define

$$(2-39) \quad e_0^{I, I_0}(t) = \frac{1}{2} \|\partial_t(\partial_t^{I_0} D^I \varphi)\|_{L^2}^2 = \frac{1}{2} \int_{\mathbb{R}^m} (\partial_t \partial_t^{I_0} D^I \varphi)^2 dx,$$

$$(2-40) \quad e_1^{I, I_0}(t) = \frac{1}{2} \left\| \sum_{i=1}^m \partial_i(\partial_t^{I_0} D^I \varphi) \right\|_{L^2}^2 = \frac{1}{2} \int_{\mathbb{R}^m} \sum_{i=1}^m (\partial_i \partial_t^{I_0} D^I \varphi)^2 dx$$

and

$$(2-41) \quad e^{I, I_0}(t) = e_0^{I, I_0}(t) + e_1^{I, I_0}(t).$$

By (2-18), (2-39), and (2-40),

$$(2-42) \quad E_0^{I, I_0}(t) = 1/2 \int_{\mathbb{R}^m} (\partial_t \partial_t^{I_0} D^I \varphi)^2 t^{lm/2} dx = t^{lm/2} e_0^{I, I_0}(t)$$

and

$$(2-43) \quad E_1^{I, I_0}(t) = \frac{1}{2} \int_{\mathbb{R}^m} \sum_{i=1}^m (\partial_i \partial_t^{I_0} D^I \varphi)^2 t^{-l} t^{lm/2} dx = t^{(m-2)l/2} e_1^{I, I_0}(t).$$

We obtain the following lemma directly from Lemma 2.8 and (2-42)–(2-43):

Lemma 2.10. *The following decay estimates hold:*

$$(2-44) \quad e_0^{I, I_0}(t) \leq t^{-l} f^{I, I_0}(1) \quad \text{and} \quad e_1^{I, I_0}(t) \leq f^{I, I_0}(1).$$

By Lemma 2.10 and the Sobolev embedding, one easily obtain the following L^∞ estimates:

Lemma 2.11. *For any $|I| \geq |J| + \lceil \frac{m}{2} + 1 \rceil$ and $i = 1, \dots, m$,*

$$(2-45) \quad \|\partial_t(\partial_t^{I_0} D^J \varphi)(t)\|_{L^\infty} \leq C_I t^{-l/2} \left(\sum_{K=0}^I f^{K, I_0}(t) \right)^{1/2} \quad \text{for } I_0 \geq 0$$

and

$$(2-46) \quad \|\partial_i(D^J \varphi)(t)\|_{L^\infty} \leq C_I \left(\sum_{K=0}^I f^{K,0}(t) \right)^{1/2},$$

provided that $f^{K,I_0}(t)$ ($K = 0, \dots, I$) is bounded.

3. The proof of Theorem 1.3

In this section, we will prove the global existence of smooth solutions to exponential wave maps (1-8) for small initial data (1-9). Since (1-8) can be reduced to a symmetric hyperbolic system, via standard results [Majda 1984; Alinhac 2010], there exists a unique local solution in the function space H^s , for $s > 1 + \frac{m}{2}$. Thus, for the global existence result, it suffices to prove a priori estimate; see Lemma 3.7.

To obtain the global existence of smooth solution to Eq. (1-8), one needs to derive the equation satisfied by the higher-order derivatives of the solution first.

Lemma 3.1. *Differentiating (1-8) by the spacial derivatives D^I ,*

$$(3-1) \quad \square_g(D^I u) = \sum_{I_1+I_2+I_3=I} -Q(D^{I_1} u, Q(D^{I_2} u, D^{I_3} u)).$$

Proof. Since D is a Killing vector field, it suffices to prove

$$(3-2) \quad D^I Q(u, v) = \sum_{I_1+I_2=I} C_{I_1, I_2} Q(D^{I_1} u, D^{I_2} v).$$

By Leibniz's rule,

$$(3-3) \quad \begin{aligned} D^I Q(u, v) &= D^I \left(-\partial_t u \partial_t v + t^{-l} \sum_{i=1}^m (\partial_i u \partial_i v) \right) \\ &= \sum_{I_1+I_2=I} C_{I_1, I_2} \left(\partial_t D^{I_1} u \partial_t D^{I_2} v + t^{-l} \sum_{i=1}^m (\partial_i D^{I_1} u) (\partial_i D^{I_2} v) \right) \\ &= \sum_{I_1+I_2=I} C_{I_1, I_2} Q(D^{I_1} u, D^{I_2} v). \end{aligned}$$

Let

$$(3-4) \quad v = Q(u, u).$$

By (3-2), we have

$$(3-5) \quad \begin{aligned} D^I Q(u, Q(u, u)) &= \sum_{I_1+I_2=I} C_{I_1, I_2} Q(D^{I_1} u, D^{I_2} Q(u, u)) \\ &= \sum_{I_1+I_{21}+I_{22}=I+I_2} C_{I_1, I_{21}, I_{22}} Q(D^{I_1} u, Q(D^{I_{21}} u, D^{I_{22}} u)). \end{aligned}$$

Thus,

$$(3-6) \quad \square_g(D^I u) = [\square_g, D^I]u + D^I(\square_g u) = -D^I Q(u, Q(u, u)).$$

From (3-5) and (3-6), we complete the proof of the lemma. □

For higher-order derivatives with respect to t , we have the following lemmas:

Lemma 3.2. *For any smooth functions $v(t, x)$ and $w(t, x)$,*

$$(3-7) \quad \begin{aligned} \partial_t^{I_0} Q(v, w) &= \sum_{I_{01}+I_{02}=I_0} Q(\partial_t^{I_{01}} v, \partial_t^{I_{02}} w) + \sum_{i=1}^m \sum_{I_{01}+I_{02}<I_0} t^{-l-(I_0-I_{01}-I_{02})} \partial_i \partial_t^{I_{01}} v \partial_i \partial_t^{I_{02}} w. \end{aligned}$$

Proof. We prove it by induction. For $I_0 = 1$, we have

$$\begin{aligned} \partial_t Q(v, w) &= \partial_t \left[-\partial_t v \partial_t w + \sum_{i=1}^m t^{-l} \partial_i v \partial_i w \right] \\ &= -\partial_t^2 v \partial_t w - \partial_t v \partial_t^2 w + \sum_{i=1}^m t^{-l} \partial_{ii}^2 v \partial_i w + \sum_{i=1}^m t^{-l} \partial_i v \partial_{ii}^2 w - \sum_{i=1}^m l t^{-l-1} \partial_i v \partial_i w \\ &= Q(\partial_t v, w) + Q(v, \partial_t w) - \sum_{i=1}^m t^{-l-(1-0)} \partial_i v \partial_i w. \end{aligned}$$

Therefore, the lemma holds for $I_0 = 1$.

Assume that the lemma holds for $I_0 - 1$ for arbitrary $I_0 > 1$; then for I_0 ,

$$\begin{aligned} \partial_t^{I_0} Q(v, w) &= \partial_t (\partial_t^{I_0-1} Q(v, w)) \\ &= \partial_t \left(\sum_{I_{01}+I_{02}=I_0-1} Q(\partial_t^{I_{01}} v, \partial_t^{I_{02}} w) + \sum_{i=1}^m \sum_{I_{01}+I_{02}<I_0-1} t^{-l-(I_0-I_{01}-I_{02})} \partial_i^{I_{01}} \partial_i v \partial_i^{I_{02}} \partial_i w \right) \\ &= \sum_{I_{01}+I_{02}=I_0-1} (Q(\partial_t^{I_{01}+1} v, \partial_t^{I_{02}} w) + Q(\partial_t^{I_{01}} v, \partial_t^{I_{02}+1} w)) \\ &\quad + \sum_{i=1}^m \sum_{I_{01}+I_{02}<I_0-1} t^{-1-(I_0+1-I_{01}-I_{02})} \partial_i^{I_{01}} \partial_i v \partial_i^{I_{02}} \partial_i w \\ &\quad + \sum_{i=1}^m \sum_{I_{01}+I_{02}<I_0-1} t^{-1-(I_0+1-I_{01}-I_{02})} (\partial_t^{I_{01}+1} \partial_i v \partial_t^{I_{02}} \partial_i w + \partial_t^{I_{01}} \partial_i v \partial_t^{I_{02}+1} \partial_i w) \\ &= \sum_{I_{01}+I_{02}=I_0} Q(\partial_t^{I_{01}} v, \partial_t^{I_{02}} w) + \sum_{i=1}^m \sum_{I_{01}+I_{02}<I_0} t^{-l-(I_0-I_{01}-I_{02})} \partial_i^{I_{01}} \partial_i v \partial_i^{I_{02}} \partial_i w. \end{aligned}$$

Thus, the lemma holds for arbitrary I_0 . □

We define

$$S(I_0) = \{(I_0, I_{01}, I_{02}, I_{021}, I_{022}) \mid I_{01} + I_{02} < I_0, I_{021} + I_{022} < I_{02}\}.$$

Lemma 3.3. *For any smooth functions $u(t, x)$, $v(t, x)$, and $w(t, x)$, and any non-negative integer I_0 ,*

$$\begin{aligned} (3-8) \quad \partial_t^{I_0} Q(u, Q(v, w)) &= \sum_{I_{01}+I_{02}+I_{03}=I_0} Q(\partial_t^{I_{01}} u, Q(\partial_t^{I_{02}} v, \partial_t^{I_{03}} w)) \\ &+ \sum_{i=1}^m \sum_{S(I_0)} t^{-l-(I_{02}-I_{021}-I_{022})} Q(\partial_t^{I_{01}} u, \partial_t^{I_{021}} \partial_i v \partial_t^{I_{022}} \partial_i w) \\ &+ \sum_{i=1}^m \sum_{S(I_0)} t^{-l-1-(I_{02}-I_{021}-I_{022})} \partial_t^{I_{01}} \partial_i u \partial_t^{I_{021}} \partial_i v \partial_t^{I_{022}} \partial_i w \\ &+ \sum_{i=1}^m \sum_{S(I_0)} t^{-l-(I_0-I_{01}-I_{02})} \partial_t^{I_{01}} \partial_i u \partial_i Q(\partial_t^{I_{021}} v, \partial_t^{I_{022}} w) \\ &+ \sum_{i=1}^m \sum_{S(I_0)} t^{2l+(I_0-I_{01}-I_{02})+(I_{02}-I_{021}-I_{022})} \partial_t^{I_{01}} \partial_i u \partial_i (\partial_t^{I_{021}} \partial_i v \partial_t^{I_{022}} \partial_i w). \end{aligned}$$

Proof. By Lemma 3.2, let $p = Q(v, w)$, so

$$\begin{aligned} (3-9) \quad \partial_t^{I_0} Q(u, Q(v, w)) &= \partial_t^{I_0} Q(u, p) \\ &= \sum_{I_{01}+I_{02}=I_0} Q(\partial_t^{I_{01}} u, \partial_t^{I_{02}} p) + \sum_{I_{01}+I_{02}<I_0} t^{-l-(I_0-I_{01}-I_{02})} \partial_t^{I_{01}} \partial_i u \partial_t^{I_{02}} \partial_i p, \end{aligned}$$

since

$$(3-10) \quad \partial_t^{I_{02}} p = \sum_{I_{021}+I_{022}=I_{02}} Q(\partial_t^{I_{021}} v, \partial_t^{I_{022}} w) + \sum_{I_{021}+I_{022}<I_{02}} t^{-l-(I_{02}-I_{021}-I_{022})} \partial_t^{I_{021}} \partial_i v \partial_t^{I_{022}} \partial_i w.$$

Inserting (3-10) into (3-9), we obtain the lemma. □

Combining Lemmas 2.1, 3.1, 3.2 and 3.3, we obtain an important lemma:

Lemma 3.4. *Differentiating (1-8) by $\partial_t^{I_0} D^I$ for any nonnegative integer I_0 and multi-index I , we have*

$$\begin{aligned}
 (3-11) \quad & \square_g(\partial_t^{I_0} D^I u) \\
 &= \sum_{I'_0=1}^{I_0} t^{-(I_0-I'_0+2)} \partial_t^{I'_0} D^I u + \sum_{i=1}^m \sum_{I''_0=0}^{I_0-1} t^{-l-(I_0-I''_0)} \partial_t^2 \partial_t^{I''_0} D^I u \\
 &\quad + \sum_{I_{01}+I_{02}+I_{03}=I_0} Q(\partial_t^{I_{01}} D^{I_1} u, Q(\partial_t^{I_{02}} D^{I_2} u, \partial_t^{I_{03}} D^{I_3} u)) \\
 &\quad + \sum_{S(I_0)} t^{-l-(I_{02}-I_{021}-I_{022})} Q(\partial_t^{I_{01}} D^{I_1} u, \partial_t^{I_{021}} \partial_t(D^{I_2} u) \partial_t^{I_{022}} \partial_t(D^{I_3} u)) \\
 &\quad + \sum_{S(I_0)} t^{-l-1-(I_{02}-I_{021}-I_{022})} \partial_t^{I_{01}}(D^{I_1} \partial_t u) \partial_t^{I_{021}}(D^{I_2} \partial_t u) \partial_t^{I_{022}}(D^{I_3} \partial_t u) \\
 &\quad + \sum_{S(I_0)} t^{-l-(I_0-I_{01}-I_{02})} \partial_t^{I_{01}}(D^{I_1} \partial_t u) \partial_t Q(\partial_t^{I_{021}} D^{I_2} u, \partial_t^{I_{022}}(D^{I_3} u)) \\
 &\quad + \sum_{S(I_0)} t^{-E} \partial_t^{I_{01}} \partial_t(D^{I_1} u) \partial_t(\partial_t^{I_{021}}(D^{I_2} \partial_t u) \partial_t^{I_{022}}(D^{I_3} \partial_t u)) \\
 &:= \sum_{I'_0=1}^{I_0} t^{-(I_0-I'_0+2)} \partial_t^{I'_0} D^I u + \sum_{i=1}^m \sum_{I''_0=0}^{I_0-1} t^{-l-(I_0-I''_0)} \partial_t^2 \partial_t^{I''_0} D^I u \\
 &\quad + Q(u, Q(u, \partial_t^{I_0} D^I u)) + R,
 \end{aligned}$$

where $E = 2l + (I_0 - I_{01} - I_{02}) + (I_{02} - I_{021} - I_{022})$, $I_1 + I_2 + I_3 = I$, and R denotes the remaining terms.

Remark 3.5. For convenience, in Lemmas 3.1–3.4, we have omitted the numerical constants in front of each term at the right hand of (3-1), (3-7), (3-8), and (3-11). It does not affect the estimates in the sequel since they are all universal constants.

Remark 3.6. In the following estimates, we distinguish the $Q(u, Q(u, \partial_t^{I_0} D^I u))$ term from others, since it contains the highest-order derivatives of $\partial_t^{I_0} D^I u$.

Based on Lemma 3.4, we have the following energy estimate. It plays a key role in the proof of Theorem 1.3.

Lemma 3.7. *The following generalized energy inequality holds in the maximal development of the smooth solution of Cauchy problem (1-8), (1-9):*

$$\begin{aligned}
 (3-12) \quad & F^M(t) \\
 & \lesssim F^M(1) + \int_1^t (\max\{\tau^{-(I_0-I'_0+2)}, \tau^{-l/2-(I_0-I''_0)}\} + \tau^{-l/2} F^M(\tau)) F^M(\tau) \, d\tau.
 \end{aligned}$$

provided $F^M(t) \ll 1$, where $F^M(t)$ is defined by (2-32) with $M \geq m + 2$.

Proof. We will prove the lemma in three steps.

Energy estimates based on (3-11). Taking $V = -\partial_t$ in Lemma 2.2, one gets

$$(3-13) \quad E^{I,J}(t) - E^{I,J}(1) = \int_1^t \int_{\Sigma_\tau} (K^V(\partial_t^J D^I u) + \square_g(\partial_t^J D^I u)V(\partial_t^J D^I u)) d\text{Vol}_\tau d\tau,$$

which is equivalent to

$$(3-14) \quad \frac{d}{dt} E^{I,J}(t) = \int_{\Sigma_t} (K^V(\partial_t^J D^I u(t)) + \square_g(\partial_t^J D^I u)V(\partial_t^J D^I u)) d\text{Vol}_t.$$

By (3-14) and (2-24),

$$(3-15) \quad \begin{aligned} & \frac{d}{dt} f^{I,J}(t) \\ &= t^{(2-m)l/2} \left(\frac{d}{dt} E^{I,J}(t) \right) + \frac{(2-m)l}{2} t^{(2-m)l/2} t^{-1} E^{I,J}(t) \\ &= t^{(2-m)l/2} \int_{\Sigma_t} (K^V(\partial_t^J D^I u(t)) + \square_g(\partial_t^J D^I u)V(\partial_t^J D^I u)) d\text{Vol}_t \\ & \quad + \frac{(2-m)l}{2} t^{-1} t^{(2-m)l/2} E^{I,J}(t) \\ &\leq t^{(2-m)l/2} \int_{\Sigma_t} \left(\sum_{J'=1}^J t^{-(J-J'+2)} \partial_t^{J'} D^I u + \sum_{i=1}^m \sum_{J''=0}^{J-1} t^{-l-(J-J'')} \partial_i^2 \partial_t^{J''} D^I u \right) \\ & \quad \times V(\partial_t^J D^I u) d\text{Vol}_t \\ & \quad + t^{(2-m)l/2} \int_{\Sigma_t} (Q(u, Q(u, \partial_t^J D^I u)) + R)V(\partial_t^J D^I u) d\text{Vol}_t. \end{aligned}$$

We first estimate the term that contains the second-order derivatives of $\partial_t^J D^I u$, i.e.,

$$(3-16) \quad \int_{\mathbb{R}^m} Q(u, Q(u, \partial_t^J D^I u)) V(\partial_t^J D^I u) t^{(2-m)l/2} t^{ml/2} dx.$$

Energy estimates based on (3-16). Let

$$(3-17) \quad v = \partial_t^J D^I u.$$

Expanding the term $Q(u, Q(u, v))$, one obtains

$$\begin{aligned} & Q(u, Q(u, v)) \\ &= -\partial_t u \partial_t \left(-\partial_t u \partial_t v + t^{-l} \sum_{i=1}^m \partial_i u \partial_i v \right) + t^{-l} \sum_{j=1}^m \partial_j u \partial_j \left(-\partial_t u \partial_t v + t^{-l} \sum_{i=1}^m \partial_i u \partial_i v \right) \end{aligned}$$

which yields

$$\begin{aligned}
 & Q(u, Q(u, v)) \\
 &= (\partial_t u)^2 \partial_t^2 v - 2t^{-l} \sum_{i=1}^m \partial_t u \partial_i u \partial_{ii}^2 v + \sum_{i,j=1}^m t^{-2l} \partial_i u \partial_j u \partial_{ij}^2 v + \partial_t u \partial_t^2 u \partial_t v \\
 &\quad - t^{-l} \sum_{i=1}^m \partial_t u \partial_{ii}^2 u \partial_t v + lt^{-l-1} \sum_{i=1}^m \partial_t u \partial_i u \partial_i v \\
 &\quad - t^{-l} \sum_{j=1}^m \partial_j u \partial_{jt}^2 u \partial_t v + t^{-2l} \sum_{i,j=1}^m \partial_j u \partial_{ij}^2 u \partial_i v \\
 &= A + B + C + L,
 \end{aligned}$$

where

$$(3-18) \quad A = (\partial_t u)^2 \partial_t^2 v, \quad B = -2t^{-l} \sum_{i=1}^m \partial_t u \partial_i u \partial_{ii}^2 v, \quad C = \sum_{i,j=1}^m t^{-2l} \partial_i u \partial_j u \partial_{ij}^2 v,$$

and

$$\begin{aligned}
 (3-19) \quad L &= \partial_t u \partial_t^2 u \partial_t v - t^{-l} \sum_{i=1}^m \partial_t u \partial_{ii}^2 u \partial_t v + lt^{-l-1} \sum_{i=1}^m \partial_t u \partial_i u \partial_i v \\
 &\quad - t^{-l} \sum_{j=1}^m \partial_j u \partial_{jt}^2 u \partial_t v + t^{-2l} \sum_{i,j=1}^m \partial_j u \partial_{ij}^2 u \partial_i v.
 \end{aligned}$$

In view of (3-16) and (3-18), and integrating by parts, we have

$$\begin{aligned}
 (3-20) \quad & \int_{\mathbb{R}^m} A \partial_t v t^{(2-m)l/2} t^{ml/2} dx \\
 &= \int_{\mathbb{R}^m} (\partial_t u)^2 \partial_t^2 v \partial_t v t^l dx \\
 &= \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^m} (\partial_t u)^2 (\partial_t v)^2 t^l dx \\
 &\quad - \int_{\mathbb{R}^m} \partial_t u \partial_t^2 u (\partial_t v)^2 t^l dx - \frac{l}{2} \int_{\mathbb{R}^m} (\partial_t u)^2 (\partial_t v)^2 t^{-1} t^l dx \\
 &= \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^m} (\partial_t u)^2 (\partial_t v)^2 t^l dx + A_1 + A_2.
 \end{aligned}$$

and

$$\begin{aligned}
 (3-21) \quad & \int_{\mathbb{R}^m} B \partial_t v t^l dx \\
 &= \int_{\mathbb{R}^m} \left(2t^{-l} \sum_{i=1}^m \partial_t u \partial_i u \partial_{it}^2 v \right) \partial_t v t^l dx \\
 &= \sum_{i=1}^m \partial_i \left(\int_{\mathbb{R}^m} \partial_t u \partial_i u (\partial_t v)^2 dx \right) \\
 &\quad - \sum_{i=1}^m \int_{\mathbb{R}^m} t^{-l} \partial_{ii}^2 u \partial_i u (\partial_t v)^2 t^l dx - \sum_{i=1}^m \int_{\mathbb{R}^m} t^{-l} \partial_t u \partial_i^2 u (\partial_t v)^2 t^l dx \\
 &:= \sum_{i=1}^m \partial_i \left(\int_{\mathbb{R}^m} \partial_t u \partial_i u (\partial_t v)^2 dx \right) + B_1 + B_2.
 \end{aligned}$$

Also,

$$\begin{aligned}
 (3-22) \quad & \int_{\mathbb{R}^m} C \partial_t v t^l dx \\
 &= - \sum_{i,j=1}^m \int_{\mathbb{R}^m} t^{-2l} \partial_i u \partial_j u \partial_{ij}^2 v \partial_t v t^l dx \\
 &= \sum_{i,j=1}^m \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^m} t^{-2l} \partial_i u \partial_j u \partial_i v \partial_j v t^l dx \\
 &\quad + \sum_{i,j=1}^m \int_{\mathbb{R}^m} t^{-2l} \partial_{ij}^2 u \partial_j u \partial_i v \partial_t v t^l dx + \int_{\mathbb{R}^m} t^{-2l} \partial_i u \partial_j^2 u \partial_i v \partial_t v t^l dx \\
 &\quad - \sum_{i,j=1}^m \int_{\mathbb{R}^m} t^{-2l} \partial_{ii}^2 u \partial_j u \partial_i v \partial_j v t^l dx - \sum_{i,j=1}^m \int_{\mathbb{R}^m} \frac{l}{2} t^{-2l-1} \partial_i u \partial_j u \partial_i v \partial_j v t^l dx \\
 &= \sum_{i,j=1}^m \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^m} t^{-2l} \partial_i u \partial_j u \partial_i v \partial_j v t^l dx + C_1 + C_2 + C_3 + C_4.
 \end{aligned}$$

Next, we estimate these terms via $F^M(t)$. By Lemma 2.11 and the assumption $M \geq m + 2$, we obtain easily

$$(3-23) \quad |A_i| \lesssim t^{-l} [F^M(t)]^2, \quad i = 1, 2,$$

$$(3-24) \quad |B_i| \lesssim t^{-l} [F^M(t)]^2, \quad i = 1, 2,$$

$$(3-25) \quad |C_i| \lesssim t^{-3l/2} [F^M(t)]^2, \quad i = 1, 2, 3,$$

and

$$(3-26) \quad |C_4| \lesssim t^{-l-1} [F^M(t)]^2.$$

For the remaining terms L , we have

$$(3-27) \quad \left| \int_{\mathbb{R}^m} L \partial_t v t^{(2-m)l/2} t^{ml/2} dx \right| \lesssim (t^{-l} + t^{-2l} + t^{-l-1}) [F^M(t)]^2.$$

At last, we estimate the lower-order terms

$$(3-28) \quad \int_{\mathbb{R}^m} R \partial_t v t^{(2-m)l/2} t^{ml/2} dx,$$

where R is defined by (3-11).

Energy estimates based on (3-28). We have

$$(3-29) \quad \int_{\mathbb{R}^m} R \partial_t v t^l dx = \int_{\mathbb{R}^m} \left(\sum_{i=1}^5 R_i \right) \partial_t v t^l dx,$$

where R_i ($i = 1, \dots, 5$) is defined by (3-11).

For R_1 , we have

$$(3-30) \quad \begin{aligned} & Q(\partial_t^{I_{01}} D^{I_1} u, Q(\partial_t^{I_{02}} D^{I_2} u, \partial_t^{I_{03}} D^{I_3} u)) \\ &= -\partial_t(\partial_t^{I_{01}} D^{I_1} u) \partial_t \left[-\partial_t(\partial_t^{I_{02}} D^{I_2} u) \partial_t(\partial_t^{I_{03}} D^{I_3} u) \right. \\ & \qquad \qquad \qquad \left. + t^{-l} \sum_{i=1}^m \partial_i(\partial_t^{I_{02}} D^{I_2} u) \partial_i(\partial_t^{I_{03}} D^{I_3} u) \right] \\ & \qquad \qquad \qquad + t^{-l} \sum_{j=1}^m \partial_j(\partial_t^{I_{01}} D^{I_1} u) \partial_j \left[-\partial_t(\partial_t^{I_{02}} D^{I_2} u) \partial_t(\partial_t^{I_{03}} D^{I_3} u) \right. \\ & \qquad \qquad \qquad \left. + t^{-l} \sum_{i=1}^m \partial_i(\partial_t^{I_{02}} D^{I_2} u) \partial_i(\partial_t^{I_{03}} D^{I_3} u) \right], \end{aligned}$$

so

$$(3-31) \quad \left| \int_{\mathbb{R}^m} R_1 \partial_t v t^{(2-m)l/2} t^{ml/2} dx \right| \lesssim (t^{-l} + t^{-l/2}) [F^M(t)]^2.$$

For R_2 , by direct calculations,

$$(3-32) \quad \begin{aligned} & Q(\partial_t^{I_{01}} D^{I_1} u, \partial_t^{I_{021}} \partial_i(D^{I_2} u) \partial_t^{I_{022}} \partial_i(D^{I_3} u)) \\ &= -\partial_t(\partial_t^{I_{01}} D^{I_1} u) \partial_t(\partial_t^{I_{021}} \partial_i(D^{I_2} u) \partial_t^{I_{022}} \partial_i(D^{I_3} u)) \\ & \qquad \qquad \qquad + t^{-l} \sum_{i=1}^m \partial_i(\partial_t^{I_{01}} D^{I_1} u) \partial_i(\partial_t^{I_{021}} \partial_i(D^{I_2} u) \partial_t^{I_{022}} \partial_i(D^{I_3} u)), \end{aligned}$$

so

$$(3-33) \quad \left| \int_{\mathbb{R}^m} R_2 \partial_t v t^{-l-(I_0-I_{021}-I_{022})} t^{(2-m)l/2} t^{ml/2} dx \right| \\ \lesssim (t^{-3l/2-(I_0-I_{021}-I_{022})} + t^{-l/2-(I_0-I_{021}-I_{022})}) [F^M(t)]^2.$$

For R_3 , we have

$$(3-34) \quad \left| \int_{\mathbb{R}^m} R_3 \partial_t v t^{-2l-(I_0-I_{01}-I_{02})-(I_{02}-I_{021}-I_{022})} t^{(2-m)l/2} t^{ml/2} dx \right| \\ \lesssim t^{-l/2-(I_0-I_{01}-I_{02})-(I_{02}-I_{021}-I_{022})} [F^M(t)]^2.$$

For R_4 ,

$$(3-35) \quad Q(\partial_t^{I_{02}} D^{I_2} u, \partial_t^{I_{03}} D^{I_3} u) \\ = -\partial_t(\partial_t^{I_{02}} D^{I_2} u) \partial_t(\partial_t^{I_{03}} D^{I_3} u) + t^{-l} \sum_{i=1}^m \partial_i(\partial_t^{I_{02}} D^{I_2} u) \partial_i(\partial_t^{I_{03}} D^{I_3} u),$$

so

$$(3-36) \quad \left| \int_{\mathbb{R}^m} R_4 \partial_t v t^{-l-(I_0-I_{01}-I_{02})} t^{(2-m)l/2} t^{ml/2} dx \right| \lesssim t^{-l/2-(I_0-I_{01}-I_{02})} [F^M(t)]^2.$$

For R_5 , we obtain

$$(3-37) \quad \left| \int_{\mathbb{R}^m} R_5 \partial_t v t^{-2l-(I_0-I_{01}-I_{02})-(I_{02}-I_{021}-I_{022})} t^{(2-m)l/2} t^{ml/2} dx \right| \\ \lesssim t^{-l/2-(I_0-I_{01}-I_{02})-(I_{02}-I_{021}-I_{022})} [F^M(t)]^2.$$

Combining these estimates above, (2-37) and (3-15)–(3-37), and summing up $|I| + J \leq M$, we have

$$(3-38) \quad \frac{d}{dt} \tilde{F}^M(t) \lesssim \max\{t^{-I_0-I'_0+2}, t^{-l/2-(I_0-I''_0)}\} F^M(t) \\ + \max\{t^{-l}, t^{-l-1}, t^{-l/2}, t^{-2l}, t^{-3l/2-(I_0-I_{021}-I_{022})}, \\ t^{-l/2-(I_0-I_{01}-I_{02})-(I_{02}-I_{021}-I_{022})}\} [F^M(t)]^2.$$

Here, we define $\tilde{F}^M(t)$ to be

$$\tilde{F}^M(t) = \sum_{|I|+J \leq M} f^{I,J}(t) + \frac{1}{2} \int_{\mathbb{R}^m} ((\partial_t u)^2 (\partial_t v)^2 t^l + \partial_t u \partial_j u \partial_i v \partial_j v t^{-2l}) dx.$$

It is easy to see that, if $F(t) \ll 1$, then there exists a positive constant C such that

$$C^{-1} F^M(t) \leq \tilde{F}^M(t) \leq C F^M(t).$$

Integrating (3-38) over $[1, t]$,

$$(3-39) \quad F^M(t) \lesssim F^M(1) + \int_1^t (\max\{\tau^{-(I_0-I'_0+2)}, \tau^{-l/2-(I_0-I''_0)}\} + \tau^{-l/2} F^M(\tau)) F^M(\tau) d\tau. \quad \square$$

Based on Lemma 3.7, we can prove Theorem 1.3 by the bootstrap method.

Proof of Theorem 1.3. Set

$$(3-40) \quad E = \{t \in [1, T) : F^M(s) \leq A\epsilon^2 \text{ for } 1 \leq s \leq t\}.$$

For the proof of the main result, it suffices to show that for any $t \in E$, the assumption $F^M(t) \leq A\epsilon^2 \ll 1$ will imply $F^M(t) \leq (A/2)\epsilon^2$, provided that A is sufficiently large and ϵ is sufficiently small.

By (3-12) and Gronwall's lemma, there exists a positive constant C such that

$$(3-41) \quad F^M(t) \leq C F^M(1) \exp \int_1^t (\max\{\tau^{-(I_0-I'_0+2)}, \tau^{-l/2-(I_0-I''_0)}\} + \tau^{-l/2} A\epsilon^2) d\tau.$$

Choosing

$$\epsilon_0 = \sqrt{\frac{\ln(2)}{100A}} \quad \text{and} \quad A \geq 16CD^M(1),$$

where $D^M(1)\epsilon^2 = F^M(1)$, then for any $\epsilon \in [0, \epsilon_0]$ and $l > 2$, we have

$$(3-42) \quad \begin{aligned} F^M(t) &\leq C F^M(1) \exp \int_1^\infty (\max\{\tau^{-(I_0-I'_0+2)}, \tau^{-l/2-(I_0-I''_0)}\} + \tau^{-l/2} A\epsilon) d\tau \\ &\leq \frac{A}{2} \epsilon^2. \end{aligned}$$

Then we can argument by contradiction and get the global existence result. □

4. Some discussions

In this paper, we have proved the global existence of smooth solutions to exponential wave maps on some special FLRW spacetimes, which are important and interesting in geometry and physics. Along with the development of Lorentzian geometry, the study of classical field theory and evolution equations is generalized to curved spacetimes. We believe that this field will attract better attention in the near future. Equations on a curved background, especially the solutions to the Einstein field equation, take a much more important role in physics. Confined by our knowledge, we only consider the easy case here, but there still exist a lot of problems that are worth focusing on.

For large initial data or large initial energy, whether the smooth solution of exponential wave maps exists globally or not is an interesting problem in the field

of PDEs. Since wave maps are a class of equations with good structure, the study of the large data problem is under consideration now; one can refer to [Wang and Yu 2013; Yang 2015] on Minkowski spacetime.

Acknowledgements

Wei was supported by the Scientific Research Foundation of Zhejiang Sci-Tech University (Grant No. 16062021-Y). Lai was supported by NSFC (Grant No. 11501273) and high level talent project of Lishui City (Grant No. 2016RC25).

References

- [Alinhac 2010] S. Alinhac, *Geometric analysis of hyperbolic differential equations: an introduction*, London Mathematical Society Lecture Note Series **374**, Cambridge University Press, 2010. MR Zbl
- [Chiang and Yang 2007] Y.-J. Chiang and Y.-H. Yang, “Exponential wave maps”, *J. Geom. Phys.* **57**:12 (2007), 2521–2532. MR Zbl
- [Choquet-Bruhat 2000] Y. Choquet-Bruhat, “Global wave maps on Robertson–Walker spacetimes”, *Nonlinear Dynam.* **22**:1 (2000), 39–47. MR Zbl
- [Eells and Lemaire 1992] J. Eells and L. Lemaire, “Some properties of exponentially harmonic maps”, pp. 129–136 in *Partial differential equations* (Warsaw, 1990), vol. 1, edited by B. Bojarski et al., Banach Center Publ. **27-1**, Polish Acad. Sci., Warsaw, 1992. MR Zbl
- [Gu 1980] C. H. Gu, “On the Cauchy problem for harmonic maps defined on two-dimensional Minkowski space”, *Comm. Pure Appl. Math.* **33**:6 (1980), 727–737. MR Zbl
- [Hawking and Ellis 1973] S. W. Hawking and G. F. R. Ellis, *The large scale structure of space-time*, Cambridge Monographs on Mathematical Physics **1**, Cambridge Univ. Press, London, 1973. MR Zbl
- [Huh 2013] H. Huh, “Global solutions of the exponential wave equation with small initial data”, *Bull. Korean Math. Soc.* **50**:3 (2013), 811–821. MR Zbl
- [Klainerman and Machedon 1993] S. Klainerman and M. Machedon, “Space-time estimates for null forms and the local existence theorem”, *Comm. Pure Appl. Math.* **46**:9 (1993), 1221–1268. MR Zbl
- [Klainerman and Machedon 1995] S. Klainerman and M. Machedon, “Smoothing estimates for null forms and applications”, *Duke Math. J.* **81**:1 (1995), 99–133. MR Zbl
- [Kong and Wei 2013] D.-X. Kong and C.-H. Wei, “Lifespan of smooth solutions for timelike extremal surface equation in de Sitter spacetime”, preprint, 2013. To appear in *J. Math. Phys.* arXiv
- [Majda 1984] A. Majda, *Compressible fluid flow and systems of conservation laws in several space variables*, Applied Mathematical Sciences **53**, Springer, New York, 1984. MR Zbl
- [Shatah and Struwe 1998] J. Shatah and M. Struwe, *Geometric wave equations*, Courant Lecture Notes in Mathematics **2**, Courant Institute of Mathematical Sciences, New York, 1998. MR Zbl
- [Shatah and Struwe 2002] J. Shatah and M. Struwe, “The Cauchy problem for wave maps”, *Int. Math. Res. Not.* **2002**:11 (2002), 555–571. MR Zbl
- [Tao 2001a] T. Tao, “Global regularity of wave maps, I: Small critical Sobolev norm in high dimension”, *Internat. Math. Res. Notices* **2001**:6 (2001), 299–328. MR Zbl
- [Tao 2001b] T. Tao, “Global regularity of wave maps, II: Small energy in two dimensions”, *Comm. Math. Phys.* **224**:2 (2001), 443–544. MR Zbl

- [Tao 2008] T. Tao, “Global regularity of wave maps, III: Large energy from \mathbb{R}^{1+2} to hyperbolic spaces”, preprint, 2008. arXiv
- [Wang and Yu 2013] J. Wang and P. Yu, “Long time solutions for wave maps with large data”, *J. Hyperbolic Differ. Equ.* **10**:2 (2013), 371–414. MR Zbl
- [Yagdjian 2009] K. Yagdjian, “The semilinear Klein–Gordon equation in de Sitter spacetime”, *Discrete Contin. Dyn. Syst. Ser. S* **2**:3 (2009), 679–696. MR Zbl
- [Yagdjian 2012] K. Yagdjian, “Global existence of the scalar field in de Sitter spacetime”, *J. Math. Anal. Appl.* **396**:1 (2012), 323–344. MR Zbl
- [Yagdjian and Galstian 2008] K. Yagdjian and A. Galstian, “Fundamental solutions of the wave equation in Robertson–Walker spaces”, *J. Math. Anal. Appl.* **346**:2 (2008), 501–520. MR Zbl
- [Yagdjian and Galstian 2009] K. Yagdjian and A. Galstian, “Fundamental solutions for the Klein–Gordon equation in de Sitter spacetime”, *Comm. Math. Phys.* **285**:1 (2009), 293–344. MR Zbl
- [Yang 2015] S. Yang, “Global solutions of nonlinear wave equations with large data”, *Selecta Math. (N.S.)* **21**:4 (2015), 1405–1427. MR Zbl

Received February 23, 2015. Revised January 27, 2016.

CHANG-HUA WEI
DEPARTMENT OF MATHEMATICS
ZHEJIANG SCI-TECH UNIVERSITY
HANGZHOU, 310018
CHINA
changhuawei1986@gmail.com

NING-AN LAI
DEPARTMENT OF MATHEMATICS, SCIENCES COLLEGE
LISHUI UNIVERSITY
LISHUI, 323000
CHINA
hyayue@gmail.com

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Igor Pak
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pak.pjm@gmail.com

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 289 No. 2 August 2017

Regular representations of completely bounded maps	257
B. V. RAJARAMA BHAT, NIRUPAMA MALLICK and K. SUMESH	
Ball convex bodies in Minkowski spaces	287
THOMAS JAHN, HORST MARTINI and CHRISTIAN RICHTER	
Local constancy of dimension of slope subspaces of automorphic forms	317
JOACHIM MAHNKOPF	
Weakening idempotency in K -theory	381
VLADIMIR MANUILOV	
On Langlands quotients of the generalized principal series isomorphic to their Aubert duals	395
IVAN MATIĆ	
Exact Lagrangian fillings of Legendrian $(2, n)$ torus links	417
YU PAN	
Elementary calculation of the cohomology rings of real Grassmann manifolds	443
RUSTAM SADYKOV	
Cluster tilting modules and noncommutative projective schemes	449
KENTA UEYAMA	
Concentration for a biharmonic Schrödinger equation	469
DONG WANG	
Global existence of smooth solutions to exponential wave maps in FLRW spacetimes	489
CHANG-HUA WEI and NING-AN LAI	



0030-8730(201708)289:2;1-O