

*Pacific  
Journal of  
Mathematics*

**HESSIAN EQUATIONS ON CLOSED HERMITIAN MANIFOLDS**

DEKAI ZHANG



## HESSIAN EQUATIONS ON CLOSED HERMITIAN MANIFOLDS

DEKAI ZHANG

We solve the complex Hessian equation on closed Hermitian manifolds, which generalizes the Kähler case proven by Hou, Ma and Wu and Dinev and Kołodziej. Solving the equation can be reduced to the derivation of a priori second-order estimates. We introduce a new method to prove the  $C^0$  estimate. The  $C^2$  estimate can be derived if we use the auxiliary function which is mainly due to Hou, Ma and Wu and Tosatti and Weinkove.

### 1. Introduction

Let  $(M, \omega)$  be a closed Hermitian manifold of complex dimension  $n \geq 2$ . In this paper, we study the Hessian equation

$$(1-1) \quad \begin{cases} \binom{n}{k} \omega_u^k \wedge \omega^{n-k} = e^f \omega^n, \\ \sup_M u = 0, \\ \omega_u = \omega + \sqrt{-1} \partial \bar{\partial} u \in \Gamma_k(M), \end{cases}$$

where  $\binom{n}{k} = n!/(k!(n-k)!)$ ,  $\Gamma_k(M)$  is a convex cone (see (2-2) in Section 2) and  $1 \leq k \leq n$ .

The complex Hessian equation is an important class of fully nonlinear elliptic equations. It arises naturally from many significant geometric problems. When  $k = 1$ , it is the classical Laplacian equation. For  $k = n$ , equation (1-1) is the complex Monge–Ampère equation

$$(1-2) \quad \omega_u^n = e^f \omega^n, \quad \sup_M u = 0.$$

Yau [1978] solved equation (1-2) on compact Kähler manifolds, and his solution is now known as Calabi–Yau theorem. For general Hermitian manifolds, (1-2) has been solved by Cherrier [1987] for dimension 2. Guan and Li [2010] and Zhang [2010] obtained  $C^1$  and  $C^2$  estimates for dimension  $n \geq 2$ . Finally, Tosatti and Weinkove [2010] derived the  $C^0$  estimate and thus solved (1-2) for arbitrary dimension.

While  $1 < k < n$ , equation (1-1) has more complicated structure and also is closely related to many important geometric problems. For example, for  $k = 2$ , it

---

*MSC2010:* 35J60, 53C55.

*Keywords:* Hessian equations, closed Hermitian manifolds, a priori estimates.

relates to Fu and Yau's [2008] generalization of the Strominger system which comes from superstring theory. Several significant results about the Fu–Yau equation have been obtained by Phong, Picard and Zhang [Phong et al. 2016a; 2016b; 2017]. When  $k = n - 1$ , it has similar features to the Monge–Ampère type equation in the study of Gauduchon conjecture by Tosatti and Weinkove [2015; 2017] and Tosatti, Weinkove and Székelyhidi [Székelyhidi et al. 2015].

We now come back to the complex Hessian equation. To solve it, it is crucial to derive the *a priori* estimates up to second-order. If  $(M, \omega)$  is a Kähler manifold, Hou, Ma and Wu [Hou et al. 2010] proved

$$(1-3) \quad \max |\partial\bar{\partial}u|_g \leq C(1 + \max |\nabla u|_g^2),$$

where  $C$  does not depend on the gradient bound of the solution.

They also pointed out that (1-3) may be adapted to the blow up analysis to get the gradient estimate. Later on, combining (1-3) with a blow up argument, Dinew and Kołodziej [2017] obtained the gradient estimate. Then equation (1-1) can be solved on Kähler manifolds.

In this paper, we solve the complex Hessian equation on closed Hermitian manifolds. More precisely,

**Theorem 1.1.** *Let  $(M, g)$  be a closed Hermitian manifold of complex dimension  $n \geq 2$  and  $f$  be a smooth real function on  $M$ . Then there exist a unique real number  $b$  and a unique smooth real function  $u$  on  $M$  solving*

$$(1-4) \quad \binom{n}{k} \omega_u^k \wedge \omega^{n-k} = e^{f+b} \omega^n, \quad \omega_u \in \Gamma_k(M), \quad \sup_M u = 0.$$

We use the continuity method to solve problem (1-4). The openness follows from implicit function theory. The closeness argument can be reduced to *a priori* estimates up to the second order by the standard Evans–Krylov theory. Actually, we can derive the zero-order estimate and the second-order estimate of solutions of equation (1-1) and thus use the blow up method to obtain the gradient estimate.

For the complex Monge–Ampère equation on closed Hermitian manifolds, Tosatti and Weinkove [2010] derived  $C^0$  estimate by proving a Cherrier-type inequality which was originally proved in [Cherrier 1987]. For the Hessian equation (1-1), we can prove a similar Cherrier-type inequality by a new method which combines an inductive argument with key inequalities for  $k$ -th elementary symmetric functions in [Chou and Wang 2001]. For the  $C^2$  estimate, the main difficulty is that there are new terms of the form  $T * D^3u$ , where  $T$  is the torsion of  $\omega$ . To control these terms, we use the auxiliary function due to Tosatti and Weinkove [2013]. The auxiliary function originally comes from Hou, Ma and Wu [Hou et al. 2010]. For the Hessian equation, the main difference is that for equation (1-1) we need to apply some

lemmas for the  $k$ -th elementary symmetric functions which were proved by Hou, Ma and Wu [Hou et al. 2010].

The rest of the paper is organized as follows. In Section 2, we give some preliminaries. In Section 3, a Cherrier-type inequality is derived, and then we obtain the  $C^0$  estimate. In Section 4, we prove the  $C^2$  estimate by a similar auxiliary function used in [Tosatti and Weinkove 2013].

Székelyhidi [2015] has also obtained similar results, but our methods are different.

## 2. Preliminaries

Let  $(M, g)$  be a closed Hermitian manifold and let  $\nabla$  denote the Chern connection of  $g$ . In this section we give some preliminaries about the  $k$ -th elementary symmetric function and the commutation formula of covariant derivatives.

**Elementary symmetric functions.** The  $k$ -th elementary symmetric function is defined by

$$\sigma_k(\lambda) = \sum_{1 \leq i_1 < \dots < i_k \leq n} \lambda_{i_1} \cdots \lambda_{i_k},$$

where  $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n$ . Let  $\lambda(a_{i\bar{j}})$  denote the eigenvalues of the Hermitian matrix  $\{a_{i\bar{j}}\}$ ; we define

$$\sigma_k(a_{i\bar{j}}) = \sigma_k(\lambda(a_{i\bar{j}})).$$

The definition of  $\sigma_k$  can be naturally extended to a Hermitian manifold. Indeed, let  $A^{1,1}(M, \mathbb{R})$  be the space of smooth real  $(1, 1)$ -forms on  $M$ ; for  $\chi \in A^{1,1}(M, \mathbb{R})$  we define

$$\sigma_k(\chi) = \binom{n}{k} \frac{\chi^k \wedge \omega^{n-k}}{\omega^n}.$$

### Definition 2.1.

$$(2-1) \quad \Gamma_k := \{\lambda \in \mathbb{R}^n : \sigma_j(\lambda) > 0, j = 1, \dots, k\}.$$

Similarly, we define  $\Gamma_k$  on  $M$  as follows

$$(2-2) \quad \Gamma_k(M) := \{\chi \in A^{1,1}(M, \mathbb{R}) : \sigma_j(\chi) > 0, j = 1, \dots, k\}.$$

Furthermore,  $\sigma_r(\lambda|i_1 \cdots i_l)$ , with  $i_1, \dots, i_l$  being distinct, denotes the  $r$ -th symmetric function with  $\lambda_{i_1} = \dots = \lambda_{i_l} = 0$ . For more details about elementary symmetric functions, one can see the lecture notes [Wang 2009].

To prove the  $C^0$  estimate, we need the following lemma for elementary symmetric functions:

**Lemma 2.2.** Suppose that  $\lambda \in \Gamma_k$ ,  $3 \leq k \leq n$  and  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ . Then there exists a positive constant  $C$  depending only on  $k$  and  $n$ , such that for  $1 \leq i \leq k-2$ ,

$$(2-3) \quad |\lambda_{j_1} \lambda_{j_2} \cdots \lambda_{j_i}| \leq (n-3)^{k-2} \sigma_i(\lambda|j), \quad 1 \leq j_1 < j_2 < \dots < j_i \leq n, \\ j_l \neq j, \quad 1 \leq l \leq i, \quad 1 \leq j \leq n.$$

*Proof.* Since

$$\sum_{p=k}^n \lambda_p = \sigma_1(\lambda|12 \cdots k-1) > 0, \quad \text{and} \quad \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n,$$

then

$$(2-4) \quad |\lambda_p| \leq (n-k)\lambda_k, \quad k+1 \leq p \leq n.$$

We first prove the lemma for  $k=3$ . In this case, one needs to prove

$$|\lambda_l| \leq C\sigma_1(\lambda|j) \quad \text{for } 1 \leq j, l \leq n \text{ and } l \neq j.$$

Since  $\sigma_1(\lambda|j) = \lambda_l + \sigma_1(\lambda|jl)$ ,  $\lambda_l \leq \sigma_1(\lambda|j)$ . Now, if  $\lambda_l < 0$ , then  $l \geq 4$ . By (2-4),

$$|\lambda_l| \leq (n-3)\lambda_3 \leq \sigma_1(\lambda|j), \quad 4 \leq l \leq n.$$

Then the lemma follows for  $k=3$ .

Next we prove the lemma for the general  $k$ ,  $3 \leq k \leq n$ .

If  $j > i$ , since  $i \leq k-2$ ,  $\lambda|j \in \Gamma_{i+1}$ , applying [Lin and Trudinger 1994, p. 322, (19)] yields  $\sigma_i(\lambda|j) \geq \lambda_1 \cdots \lambda_i$ . Since  $1 \leq l \leq i \leq k-2$ , by (2-4) we have

$$|\lambda_{jl}| \leq \max\{\lambda_l, (n-k)\lambda_k\} \leq (n-k)\lambda_l.$$

Then

$$(2-5) \quad |\lambda_{j_1} \lambda_{j_2} \cdots \lambda_{j_i}| \leq (n-k)^i \lambda_1 \cdots \lambda_i \leq (n-k)^{k-2} \sigma_i(\lambda|j).$$

If  $j \leq i$ , applying [Lin and Trudinger 1994, p. 322, (19)] yields

$$\sigma_i(\lambda|j) \geq \lambda_1 \cdots \lambda_{j-1} \lambda_{j+1} \cdots \lambda_{i+1}.$$

Note  $j_l \neq j$ , so

$$|\lambda_{jl}| \leq \begin{cases} (n-3)\lambda_l, & j_l < j, \\ (n-3)\lambda_{l+1}, & j_l > j. \end{cases}$$

Therefore, we have

$$(2-6) \quad |\lambda_{j_1} \lambda_{j_2} \cdots \lambda_{j_i}| \leq (n-k)^i \lambda_1 \cdots \lambda_{j-1} \lambda_{j+1} \cdots \lambda_{i+1} \leq (n-k)^{k-2} \sigma_i(\lambda|j).$$

Combining (2-5) and (2-6), we obtain

$$|\lambda_{j_1} \lambda_{j_2} \cdots \lambda_{j_i}| \leq (n-3)^{k-2} \sigma_i(\lambda|j), \quad 1 \leq j_1 < j_2 < \cdots < j_i \leq n,$$

$$j_l \neq j, \quad 1 \leq l \leq i, \quad 1 \leq j \leq n. \quad \square$$

By Lemma 2.2, we immediately obtain the following lemma which is a key ingredient in proving Lemma 3.2:

**Lemma 2.3.** *There exists a positive constant  $C$  depending only on  $(M, \omega)$  and  $n$  such that*

$$(2-7) \quad \left| \frac{\sqrt{-1}\partial u \wedge \bar{\partial} u \wedge \omega_u^{i-1} \wedge T_i}{\omega^n} \right| \leq C \frac{\sqrt{-1}\partial u \wedge \bar{\partial} u \wedge \omega_u^i \wedge \omega^{n-i-1}}{\omega^n},$$

where  $T_i$  is defined as the combinations of  $\omega, \partial\omega, \bar{\partial}\omega$ ; more precisely,

$$T_i = \sum_{0 \leq 3p+2q \leq n-i} \omega^{n-i-3p-2q} \wedge (\sqrt{-1})^p (\partial\omega)^p \wedge (\bar{\partial}\omega)^q \wedge (\sqrt{-1})^q (\bar{\partial}\bar{\partial}\omega)^q$$

for  $1 \leq i \leq k-1$ .

*Proof.* For  $x \in M$ , we choose the coordinates such that

$$\omega(x) = \sqrt{-1} \sum_{j=1}^n dz^j \wedge d\bar{z}^j, \quad \omega_u(x) = \sqrt{-1} \sum_{j=1}^n \lambda_j dz^j \wedge d\bar{z}^j,$$

and  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ . Write  $T_i$  as follows:

$$T_i = (\sqrt{-1})^{n-i} (T_i)_{l_1 \dots l_{n-i}, \bar{m}_1, \dots \bar{m}_{n-i}} dz^{l_1} \wedge \dots \wedge dz^{l_{n-i}} \wedge d\bar{z}^{m_1} \wedge \dots \wedge d\bar{z}^{m_{n-i}}.$$

Then

$$(2-8) \quad \begin{aligned} \left| \frac{\sqrt{-1}\partial u \wedge \bar{\partial} u \wedge \omega_u^i \wedge T_i}{\omega^n} \right| &\leq C \sum_{j,l=1}^n \sum_{1 \leq j_1 < \dots < j_i \leq n, j \neq j_l} |u_j| |u_{\bar{j}}| |\lambda_{j_1} \lambda_{j_2} \dots \lambda_{j_i}| \\ &\leq C \sum_{j=1}^n \sum_{\substack{1 \leq j_1 < \dots < j_i \leq n \\ j_i \neq j, 1 \leq l \leq i}} |u_j|^2 |\lambda_{j_1} \lambda_{j_2} \dots \lambda_{j_i}| \\ &\leq C \sum_{j=1}^n \sigma_i(\lambda|j) |u_j|^2 \\ &= C \frac{\sqrt{-1}\partial u \wedge \bar{\partial} u \wedge \omega_u^i \wedge \omega^{n-i-1}}{\omega^n}, \end{aligned}$$

where we have used Lemma 2.2 in the last inequality and  $C$  depends on the bound of the torsion and the curvature of  $(M, \omega)$ .  $\square$

**Commutation formula of covariant derivatives.** We have, in local complex coordinates  $z_1, \dots, z_n$ ,

$$(2-9) \quad g_{i\bar{j}} = g\left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^j}\right), \quad \{g^{i\bar{j}}\} = \{g_{i\bar{j}}\}^{-1}$$

For the Chern connection  $\nabla$ , we denote the covariant derivatives

$$(2-10) \quad u_i = \nabla_{\partial/\partial z^i} u, \quad u_{i\bar{j}} = \nabla_{\partial/\partial \bar{z}^j} \nabla_{\partial/\partial z^i} u, \quad u_{i\bar{j}k} = \nabla_{\partial/\partial z^k} \nabla_{\partial/\partial \bar{z}^j} \nabla_{\partial/\partial z^i} u.$$

We use commutation formulas for covariant derivatives on Hermitian manifolds which can be found in [Tosatti and Weinkove 2013]:

$$(2-11) \quad u_{i\bar{j}l} = u_{l\bar{j}i} - T_{li}^p u_{p\bar{j}}, \quad u_{pi\bar{j}} = u_{p\bar{j}i} + u_q R_{i\bar{j}p}^q, \quad u_{i\bar{p}\bar{j}} = u_{i\bar{j}\bar{p}} - \overline{T_{jp}^q} u_{i\bar{q}}.$$

$$(2-12) \quad u_{i\bar{j}l\bar{m}} = u_{l\bar{m}i\bar{j}} + u_{p\bar{j}} R_{l\bar{m}i}^p - u_{p\bar{m}} R_{i\bar{j}l}^p - T_{li}^p u_{p\bar{m}i} - \overline{T_{mj}^p} u_{l\bar{p}i} - T_{li}^p \overline{T_{mj}^q} u_{p\bar{q}}.$$

### 3. Zero-order estimate

In this section we derive the zero-order estimate by proving a Cherrier-type inequality and the lemmas in [Tosatti and Weinkove 2010]. Since the constant  $b$  in Theorem 1.1 satisfies

$$|b| \leq \sup_M |f| + C,$$

where  $C$  is a positive constant depending only on  $(M, \omega)$ . Thus, we will assume  $b = 0$  for convenience.

**Theorem 3.1.** *Let  $u$  be a solution of Theorem 1.1. Then there exists a constant  $C$  depending only on  $(M, \omega)$ ,  $n$ ,  $k$  and  $\sup_M |f|$  such that*

$$\sup_M |u| \leq C.$$

Due to Tosatti and Weinkove's results, finding the zero-order estimate can be reduced to deriving a Cherrier-type inequality which was first proved by Cherrier [1987]. For the Hessian equation, we use a new method which combines an inductive argument with the key Lemma 2.3. Even for the Monge–Ampère equation, our proof is different from that in [Tosatti and Weinkove 2010].

**Lemma 3.2.** *There exist constants  $p_0$  and  $C$  depending only on  $(M, \omega)$ ,  $n$ ,  $k$  and  $\sup_M |f|$  such that for any  $p \geq p_0$*

$$\int_M |\partial e^{-(p/2)u}|_g^2 \omega^n \leq Cp \int_M e^{-pu} \omega^n.$$

**Remark 3.3.** Recently, applying our key Lemma 2.2, Sun [2017] also proved the lemma above.

*Proof.* By the equation, we have

$$\omega_u^k \wedge \omega^{n-k} - \omega^n = \left( \frac{e^f}{\binom{n}{k}} - 1 \right) \omega^n \leq C_0 \omega^n,$$

where  $C_0$  is a constant depending only on  $\sup_M f$ ,  $n$  and  $k$ . On the other hand,

$$(3-1) \quad \omega_u^k \wedge \omega^{n-k} - \omega^n = (\omega_u^k - \omega^k) \wedge \omega^{n-k} = \sqrt{-1} \partial \bar{\partial} u \wedge \alpha,$$

where  $\alpha = \sum_{i=1}^k \omega_u^{i-1} \wedge \omega^{n-i}$ .

Now multiply both sides in (3-1) by  $e^{-pu}$  and integrate by parts:

$$\begin{aligned}
 (3-2) \quad C_0 \int_M e^{-pu} \omega^n &\geq \int_M e^{-pu} \sqrt{-1} \partial \bar{\partial} u \wedge \alpha \\
 &= - \int_M \partial e^{-pu} \sqrt{-1} \bar{\partial} u \wedge \alpha + \int_M e^{-pu} \sqrt{-1} \bar{\partial} u \wedge \partial \alpha \\
 &= p \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \alpha - \frac{1}{p} \int_M \sqrt{-1} \bar{\partial} e^{-pu} \wedge \partial \alpha \\
 &= p \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \alpha + \frac{1}{p} \int_M e^{-pu} \sqrt{-1} \bar{\partial} \partial \alpha \\
 &:= A + B,
 \end{aligned}$$

where we denote

$$A = p \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \left( \sum_{i=1}^k \omega_u^{i-1} \wedge \omega^{n-i} \right), \quad B = \frac{1}{p} \int_M e^{-pu} \sqrt{-1} \bar{\partial} \partial \alpha.$$

Our goal is to use term  $A$  to control term  $B$ . Direct calculation gives

$$\partial \alpha = n \sum_{i=1}^{k-1} \omega_u^{i-1} \wedge \omega^{n-i-1} \wedge \partial \omega + (n-k) \omega_u^{k-1} \wedge \omega^{n-k-1} \wedge \partial \omega,$$

and

$$\begin{aligned}
 \bar{\partial} \partial \alpha &= (n-k)(n-k-1) \omega_u^{k-1} \wedge \omega^{n-k-2} \wedge \bar{\partial} \omega \wedge \partial \omega + (n-k) \omega_u^{k-1} \wedge \omega^{n-k-1} \wedge \bar{\partial} \partial \omega \\
 &\quad + (n-k)(n+k-1) \omega_u^{k-2} \wedge \omega^{n-k-1} \wedge \bar{\partial} \omega \wedge \partial \omega \\
 &\quad + n(n-1) \sum_{i=0}^{k-3} \omega_u^i \wedge \omega^{n-i-3} \wedge \bar{\partial} \omega \wedge \partial \omega + n \sum_{i=0}^{k-2} \omega_u^i \wedge \omega^{n-i-2} \wedge \bar{\partial} \partial \omega.
 \end{aligned}$$

Then we have

$$\begin{aligned}
 B &= \frac{(n-k)(n-k-1)}{p} \int_M e^{-pu} \omega_u^{k-1} \wedge \omega^{n-k-2} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial \omega \\
 &\quad + \frac{(n-k)}{p} \int_M e^{-pu} \omega_u^{k-1} \wedge \omega^{n-k-1} \wedge \sqrt{-1} \bar{\partial} \partial \omega \\
 &\quad + \frac{(n+k-1)(n-k)}{p} \int_M e^{-pu} \omega_u^{k-2} \wedge \omega^{n-k-1} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial \omega \\
 &\quad + \frac{n(n-1)}{p} \sum_{i=0}^{k-3} \int_M e^{-pu} \omega_u^i \wedge \omega^{n-i-3} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial \omega \\
 &\quad + \frac{n}{p} \sum_{i=0}^{k-2} \int_M e^{-pu} \omega_u^i \wedge \omega^{n-i-2} \wedge \sqrt{-1} \bar{\partial} \partial \omega.
 \end{aligned}$$

When  $k = 2$ , term  $B$  just becomes

$$\begin{aligned}
 B &= \frac{(n-2)(n-3)}{p} \int_M e^{-pu} \omega_u \wedge \omega^{n-4} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial \omega \\
 &\quad + \frac{(n-2)}{p} \int_M e^{-pu} \omega_u \wedge \omega^{n-3} \wedge \sqrt{-1} \bar{\partial} \partial \omega \\
 &\quad \quad + \frac{(n+1)(n-2)}{p} \int_M e^{-pu} \omega^{n-3} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial \omega \\
 &= \frac{(n-2)(n-3)}{p} \int_M e^{-pu} \sqrt{-1} \partial \bar{\partial} u \wedge \omega^{n-4} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial \omega \\
 &\quad + \frac{(n-2)}{p} \int_M e^{-pu} \sqrt{-1} \partial \bar{\partial} u \wedge \omega^{n-3} \wedge \sqrt{-1} \bar{\partial} \partial \omega \\
 &\quad \quad + \frac{2(n-1)(n-2)}{p} \int_M e^{-pu} \omega^{n-3} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial \omega \\
 &\quad \quad + \frac{(n-2)}{p} \int_M e^{-pu} \omega^{n-2} \wedge \sqrt{-1} \bar{\partial} \partial \omega \\
 (3-3) \quad &\geq \frac{(n-2)(n-3)}{p} \int_M e^{-pu} \sqrt{-1} \partial \bar{\partial} u \wedge \omega^{n-4} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial \omega \\
 &\quad + \frac{(n-2)}{p} \int_M e^{-pu} \sqrt{-1} \partial \bar{\partial} u \wedge \omega^{n-3} \wedge \sqrt{-1} \bar{\partial} \partial \omega - \frac{C_1}{p} \int_M e^{-pu} \omega^n.
 \end{aligned}$$

We next use integration by parts to deal with the first term and second term on the right-hand side of the above equality. Indeed,

$$\begin{aligned}
 (3-4) \quad &\int_M e^{-pu} \sqrt{-1} \partial \bar{\partial} u \wedge \omega^{n-4} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial \omega \\
 &= p \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega^{n-4} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial \omega \\
 &\quad + \int_M e^{-pu} \sqrt{-1} \bar{\partial} u \wedge \sqrt{-1} \partial (\omega^{n-4} \wedge \bar{\partial} \omega \wedge \partial \omega) \\
 &= p \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega^{n-4} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial \omega \\
 &\quad + \frac{1}{p} \int_M e^{-pu} \sqrt{-1} \bar{\partial} \partial (\omega^{n-4} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial \omega) \\
 &\geq -p C_1 \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega^{n-1} - \frac{C_1}{p} \int_M e^{-pu} \omega^n \\
 &\geq -C_1 A - \frac{C_1}{p} \int_M e^{-pu} \omega^n.
 \end{aligned}$$

A similar calculation gives

$$(3-5) \quad \int_M e^{-pu} \sqrt{-1} \partial \bar{\partial} u \wedge \omega^{n-3} \wedge \sqrt{-1} \bar{\partial} \partial \omega \geq -C_1 A - \frac{C_1}{p} \int_M e^{-pu} \omega^n.$$

Inserting (3-4) and (3-5) into (3-3), we have

$$B \geq -\frac{C_1}{p} A - \frac{C_1}{p} \int_M e^{-pu} \omega^n.$$

By (3-2) and choosing  $p_0 = 2C_1 + 1$ , we obtain for  $p \geq p_0$

$$\frac{A}{2} \leq \left(1 - \frac{C_1}{p}\right) A \leq \left(\frac{C_1}{p} + C_0\right) \int_M e^{-pu} \omega^n \leq (C_0 + 1) \int_M e^{-pu} \omega^n.$$

By (3-7) below, we thus prove the lemma.

For the general  $k$ ,  $3 \leq k \leq n$ , we claim that there exist constants  $C_{1i}$  depending only on  $n, k$  and  $(M, \omega)$  such that the following holds for  $0 \leq i \leq k-1$ :

$$(3-6) \quad \begin{aligned} & \int_M e^{-pu} \omega_u^i \wedge T_i \\ & \geq -p C_{1i} \sum_{j=0}^{k-2} \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega_u^j \wedge \omega^{n-j-1} - C_{1i} \int_M e^{-pu} \omega^n, \end{aligned}$$

where  $T_i$  is defined as the combinations of  $\omega, \partial\omega, \bar{\partial}\omega$ ; more precisely

$$T_i = \sum_{0 \leq 3p+2q \leq n-i} \omega^{n-i-3p-2q} \wedge (\sqrt{-1})^p (\partial\omega)^p \wedge (\bar{\partial}\omega)^q \wedge (\sqrt{-1})^q (\bar{\partial}\bar{\partial}\omega)^q.$$

We use the claim (3-6) to prove the lemma:

$$\begin{aligned} B & \geq -C_1 \sum_{i=2}^k \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega_u^{k-i} \wedge \omega^{n+i-k-1} - \frac{C_1}{p} \int_M e^{-pu} \omega^n \\ & \geq -\frac{C_1}{p} A - \frac{C_1}{p} \int_M e^{-pu} \omega^n. \end{aligned}$$

Thus we have

$$\left(1 - \frac{C_1}{p}\right) A \leq \left(\frac{C_1}{p} + C_0\right) \int_M e^{-pu} \omega^n.$$

Now we choose  $p_0 = 2C_1 + 1$ , then for any  $p \geq p_0$ ,

$$p^2 \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega^{n-1} \leq 2p(C_0 + 1) \int_M e^{-pu} \omega^n.$$

Therefore we have

$$\begin{aligned}
 (3-7) \quad \int_M |\partial e^{-(p/2)u}|_g^2 \omega^n &= \frac{np^2}{4} \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega^{n-1} \\
 &\leq \frac{np(C_0 + 1)}{2} \int_M e^{-pu} \omega^n \\
 &= pC \int_M e^{-pu} \omega^n.
 \end{aligned}$$

Now, we prove the claim (3-6) by an inductive argument. When  $i = 1$ ,

$$\begin{aligned}
 \int_M e^{-pu} \omega_u \wedge T_1 &= \int_M e^{-pu} \omega \wedge T_1 + \int_M e^{-pu} \sqrt{-1} \partial \bar{\partial} u \wedge T_1 \\
 &= \int_M e^{-pu} \omega \wedge T_1 - \int_M \partial e^{-pu} \wedge \sqrt{-1} \bar{\partial} u \wedge T_1 + \int_M e^{-pu} \sqrt{-1} \bar{\partial} u \wedge \partial T_1 \\
 &= \int_M e^{-pu} \omega \wedge T_1 + p \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge T_1 - \frac{1}{p} \int_M \sqrt{-1} \bar{\partial} e^{-pu} \wedge \partial T_1 \\
 &= p \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge T_1 + \int_M e^{-pu} \omega \wedge T_1 - \frac{1}{p} \int_M e^{-pu} \wedge \sqrt{-1} \partial \bar{\partial} T_1 \\
 &\geq -C_1 p \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega^{n-1} - C_1 \int_M e^{-pu} \omega^n.
 \end{aligned}$$

Suppose that the claim is true for  $l \leq i-1$ ; we will prove that the claim is also true for  $l = i$ . Indeed,

$$\begin{aligned}
 \int_M e^{-pu} \omega_u^i \wedge T_i &= \int_M e^{-pu} \omega_u^{i-1} \wedge \omega \wedge T_i + \int_M e^{-pu} \sqrt{-1} \partial \bar{\partial} u \wedge \omega_u^{i-1} \wedge T_i \\
 &= \int_M e^{-pu} \omega_u^{i-1} \wedge \omega \wedge T_i + p \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega_u^{i-1} \wedge T_i \\
 &\quad + \int_M e^{-pu} \bar{\partial} u \wedge \sqrt{-1} \partial (\omega_u^{i-1} \wedge T_i) \\
 &:= A_{i,1} + A_{i,2} + A_{i,3}.
 \end{aligned}$$

By the induction,

$$\begin{aligned}
 A_{i,1} &= \int_M e^{-pu} \omega_u^{i-1} \wedge \omega \wedge T_i \\
 &\geq -pC_{1i}(n, k, \omega) \sum_{j=0}^{k-2} \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega_u^j \wedge \omega^{n-j-1} \\
 &\quad - C_{1i}(n, k, \omega) \int_M e^{-pu} \omega^n.
 \end{aligned}$$

By the inequality (2-7) in Lemma 2.3, we have

$$(3-8) \quad \begin{aligned} A_{i,2} &= p \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega_u^{i-1} \wedge T_i \\ &\geq -p C_{2i} \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega_u^{i-1} \wedge \omega^{n-i}. \end{aligned}$$

Now we deal with the term  $A_{i,3}$ :

$$\begin{aligned} A_{i,3} &= \int_M e^{-pu} \bar{\partial} u \wedge \sqrt{-1} \partial (\omega_u^{i-1} \wedge T_i) \\ &= \frac{1}{p} \int_M e^{-pu} \sqrt{-1} \bar{\partial} \partial (\omega_u^{i-1} \wedge T_i) \\ &= \frac{(i-1)(i-2)}{p} \int_M e^{-pu} \sqrt{-1} \omega_u^{i-3} \wedge \bar{\partial} \omega \wedge \partial \omega \wedge T_i \\ &\quad + \frac{i-1}{p} \int_M e^{-pu} \omega_u^{i-2} \wedge \sqrt{-1} \bar{\partial} (\partial \omega \wedge T_i) + \frac{i-1}{p} \int_M e^{-pu} \omega_u^{i-2} \wedge \sqrt{-1} \bar{\partial} \omega \wedge \partial T_i \\ &\quad - \frac{1}{p} \int_M e^{-pu} \omega_u^{i-1} \wedge \sqrt{-1} \partial \bar{\partial} T_i \\ &= \frac{(i-1)(i-2)}{p} \int_M e^{-pu} \sqrt{-1} \omega_u^{i-3} \wedge \bar{\partial} \omega \wedge \partial \omega \wedge T_i \\ &\quad + \frac{i-1}{p} \int_M e^{-pu} \omega_u^{i-2} \wedge [\sqrt{-1} \bar{\partial} (\partial \omega \wedge T_i) + \sqrt{-1} \bar{\partial} \omega \wedge \partial T_i] \\ &\quad - \frac{1}{p} \int_M e^{-pu} \omega_u^{i-1} \wedge \sqrt{-1} \partial \bar{\partial} T_i \\ &\geq -p C_{3i} \sum_{j=0}^{k-2} \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega_u^j \wedge \omega^{n-j-1} - C_{3i}(n, k, \omega) \int_M e^{-pu} \omega^n. \end{aligned}$$

For the last inequality, we have used the induction.  $\square$

#### 4. Second-order estimate

In this section we use the auxiliary function in [Tosatti and Weinkove 2013] which is modified by the auxiliary function in [Hou et al. 2010] to derive the second-order estimate of the form (1-3). The difficulty arises from the third-order derivatives. Locally the equation is

$$(4-1) \quad \sigma_k(\omega_u) = e^f.$$

**Theorem 4.1.** *There exists a uniform constant  $C$  depending only on  $(M, \omega)$ ,  $n$ ,  $k$  and  $f$  such that*

$$(4-2) \quad \max |\partial \bar{\partial} u|_g \leq C(1 + \max |\nabla u|_g^2).$$

*Proof.* Denote  $w_{i\bar{j}} = g_{i\bar{j}} + u_{i\bar{j}}$  and let  $\xi \in T^{1,0}M$ ,  $|\xi|_g^2 = 1$ .

We use the auxiliary function similar to that in [Tosatti and Weinkove 2013]:

$$H(x, \xi) = \log(w_{k\bar{l}}\xi^k\bar{\xi}^l) + c_0 \log(g^{k\bar{l}}w_{p\bar{l}}w_{k\bar{q}}\xi^p\bar{\xi}^q) + \varphi(|\nabla u|_g^2) + \psi(u),$$

where  $\varphi, \psi$  are given by

$$\begin{aligned}\varphi(s) &= -\frac{1}{2} \log\left(1 - \frac{s}{2K}\right), & 0 \leq s \leq K-1, \\ \psi(t) &= -A \log\left(1 + \frac{t}{2L}\right), & -L+1 \leq t \leq 0,\end{aligned}$$

for

$$K := \sup_M |\nabla u|_g^2 + 1, \quad L = \sup_M |u| + 1, \quad A := 2L(C_0 + 1),$$

where  $A_0$  is a constant to be determined later and  $c_0$  is a small positive constant depending only on  $n$  and will be determined later. By [Hou et al. 2010], we have

$$(4-3) \quad \frac{1}{2K} \geq \varphi' \geq \frac{1}{4K} > 0, \quad \varphi'' = 2(\varphi')^2 > 0.$$

$$(4-4) \quad \frac{A}{L} \geq -\psi' \geq \frac{A}{2L} = C_0 + 1, \quad \psi'' \geq \frac{2\varepsilon_0}{1-\varepsilon_0}(\psi')^2, \quad \text{for } \varepsilon_0 \leq \frac{1}{2A+1}.$$

These inequalities will be used below.

Suppose  $H(x, \xi)$  attains its maximum at the point  $x_0$  in the direction  $\xi_0$ . Then we choose local coordinates  $\{\partial/\partial z^1, \dots, \partial/\partial z^n\}$  near  $x_0$  such that

$$g_{i\bar{j}}(x_0) = \delta_{ij}, \quad u_{i\bar{j}} = u_{i\bar{i}}(x_0)\delta_{ij}, \quad \lambda_i = w_{i\bar{i}}(x_0) = 1 + u_{i\bar{i}}(x_0) \quad \text{with } \lambda_1 \geq \dots \geq \lambda_n.$$

We want to prove that

$$H(x_0, \xi) \leq H\left(x_0, \frac{\partial}{\partial z^1}\right) \quad \text{for all } \xi \in T^{1,0}M, \quad |\xi|_g^2 = 1, \quad \sum_{i,j} w_{i\bar{j}}(x_0)\xi^i\bar{\xi}^j > 0$$

by choosing  $c_0$  small enough. In fact, at  $x_0$  we have

$$\log(w_{k\bar{l}}\xi^k\bar{\xi}^l) + c_0 \log(g^{k\bar{l}}w_{p\bar{l}}w_{k\bar{q}}\xi^p\bar{\xi}^q) = \log\left(\sum_{k=1}^n w_{kk}|\xi^k|^2\right) + c_0 \log\left(\sum_{k=1}^n |w_{kk}|^2|\xi^k|^2\right).$$

If  $w_{n\bar{n}} \geq -w_{1\bar{1}}$ , which is always satisfied when  $n \leq 3$ , then  $w_{i\bar{i}}^2 \leq w_{1\bar{1}}$ . Thus we have  $H(x_0, \xi) \leq H(x_0, \partial/\partial z^1)$ .

Now we suppose that  $w_{n\bar{n}} < -w_{1\bar{1}}$ . Thus we have  $n \geq 4$ . Let  $i_0$  be the smallest integer satisfying  $w_{i\bar{i}} < -w_{1\bar{1}}$ . Then  $i_0 \geq k+1$ . By  $|w_{i\bar{i}}| < (n-2)w_{1\bar{1}}$ , so

$$\begin{aligned} & \log \sum_{i=1}^n w_{i\bar{i}} |\xi^i|^2 + c_0 \log \sum_{i=1}^n |w_{i\bar{i}}|^2 |\xi^i|^2 \\ & \leq \log w_{1\bar{1}} \left( \sum_{i=1}^{i_0-1} |\xi^i|^2 - \sum_{i=i_0}^n |\xi^i|^2 \right) + c_0 \log \left( w_{1\bar{1}}^2 \sum_{i=1}^{i_0-1} |\xi^i|^2 + (n-2)^2 w_{1\bar{1}}^2 \sum_{i=1}^{i_0-1} |\xi^i|^2 \right) \\ & = \log w_{1\bar{1}} (1-2t) + c_0 \log w_{1\bar{1}}^2 (1-t+(n-2)^2 t) := h(t), \end{aligned}$$

where  $t = \sum_{i=i_0}^n |\xi^i|^2 \in (0, \frac{1}{2})$ .

By choosing  $c_0 = 2/((n-2)^2 - 1)$ , we have  $h'(t) \leq 0$ . Then

$$h(t) \leq h(0) = \log(w_{1\bar{1}}) + c_0 \log w_{1\bar{1}}^2.$$

Consequently, we obtain

$$H(x_0, \xi) \leq H\left(x_0, \frac{\partial}{\partial z^1}\right) \quad \text{for all } \xi \in T^{1,0}M, |\xi|_g^2 = 1, \sum_{i,j} \eta_{ij}(x_0) \xi^i \xi^j > 0,$$

by choosing  $c_0 = 2/((n-2)^2 - 1)$  when  $n \geq 4$  and  $c_0 = 1$  when  $n \leq 3$ .

We extend  $\xi_0$  near  $x_0$  by  $\xi_0 = (g_{1\bar{1}})^{-1/2} (\partial/\partial z^1)$ . Consider the function

$$Q(x) = H(x, \xi_0) = \log(g_{1\bar{1}}^{-1} w_{1\bar{1}}) + c_0 \log(g_{1\bar{1}}^{-1} g^{k\bar{l}} w_{1\bar{l}} w_{k\bar{l}}) + \varphi(|\nabla u|_g^2) + \psi(u).$$

We will calculate  $F^{i\bar{j}} Q_{i\bar{j}}$  at  $x_0$  to get the estimate; all the calculations are taken at  $x_0$ . For simplicity, we denote  $\xi = \xi_0$  in the following. By  $\langle \xi, \bar{\xi} \rangle_g = |\xi|_g^2 = 1$ , differentiating both sides, we obtain at  $x_0$

$$\begin{aligned} 0 &= \frac{\partial}{\partial z^i} \langle \xi, \bar{\xi} \rangle_g = \langle \nabla_{\partial/\partial z^i} \xi, \bar{\xi} \rangle_g + \langle \xi, \nabla_{\partial/\partial z^i} \bar{\xi} \rangle_g \\ &= \left\langle \xi^k, i \frac{\partial}{\partial z^k}, \bar{\xi}^l \frac{\partial}{\partial z^l} \right\rangle_g + \left\langle \xi^k \frac{\partial}{\partial z^k}, \bar{\xi}^l, i \frac{\partial}{\partial z^l} \right\rangle_g \\ &= g_{k\bar{l}} \xi^k, i \bar{\xi}^l + g_{k\bar{l}} \xi^k \bar{\xi}^l, i \\ (4-5) \quad &= \xi^1, i + \bar{\xi}^1, i. \end{aligned}$$

We also have the basic formula for  $\xi \in T^{1,0}M$ :

$$\begin{aligned} \bar{\xi}^k, i &= \frac{\partial \bar{\xi}^k}{\partial z^i} = \frac{\partial \overline{\xi^k}}{\partial \bar{z}^i} = \overline{\xi^k, i}, \quad \xi^k, \bar{i} = \frac{\partial \xi^k}{\partial \bar{z}^i} = \frac{\partial \overline{\xi^k}}{\partial z^i} = \overline{\xi^k, i} \\ (4-6) \quad \bar{\xi}^k, i &= \frac{\partial \bar{\xi}^k}{\partial z^i} = \frac{\partial \overline{\xi^k}}{\partial z^i} = \overline{\xi^k, i}, \quad \xi^k, \bar{i} = \frac{\partial \xi^k}{\partial z^i} = \frac{\partial \overline{\xi^k}}{\partial \bar{z}^i} = \overline{\xi^k, i} \end{aligned}$$

Direct calculations give

$$\begin{aligned} Q_i &= \frac{(w_{k\bar{l}}\xi^k\bar{\xi}^l)_i}{w_{k\bar{l}}\xi^k\bar{\xi}^l} + c_0 \frac{(g^{p\bar{q}} w_{k\bar{q}} w_{p\bar{l}} \xi^k \bar{\xi}^l)_i}{g^{p\bar{q}} w_{k\bar{q}} w_{p\bar{l}} \xi^k \bar{\xi}^l} + \varphi_i + \psi_i, \\ Q_{i\bar{i}} &= \frac{(w_{k\bar{l}}\xi^k\bar{\xi}^l)_{i\bar{i}}}{w_{k\bar{l}}\xi^k\bar{\xi}^l} - \frac{(w_{k\bar{l}}\xi^k\bar{\xi}^l)_i (w_{k\bar{l}}\xi^k\bar{\xi}^l)_{\bar{i}}}{(w_{k\bar{l}}\xi^k\bar{\xi}^l)^2} + c_0 \frac{(g^{p\bar{q}} w_{k\bar{q}} w_{p\bar{l}} \xi^k \bar{\xi}^l)_{i\bar{i}}}{g^{p\bar{q}} w_{k\bar{q}} w_{p\bar{l}} \xi^k \bar{\xi}^l} \\ &\quad - c_0 \frac{(g^{p\bar{q}} w_{k\bar{q}} w_{p\bar{l}} \xi^k \bar{\xi}^l)_i (g^{p\bar{q}} w_{k\bar{q}} w_{p\bar{l}} \xi^k \bar{\xi}^l)_{\bar{i}}}{(g^{p\bar{q}} w_{k\bar{q}} w_{p\bar{l}} \xi^k \bar{\xi}^l)^2} + \varphi_{i\bar{i}} + \psi_{i\bar{i}}. \end{aligned}$$

Next, we want to simplify  $Q_i$  and  $Q_{i\bar{i}}$ . By (4-5), we have

$$(w_{k\bar{l}}\xi^k\bar{\xi}^l)_i = w_{k\bar{l},i}\xi^k\bar{\xi}^l + w_{k\bar{l}}\xi^k,_i\bar{\xi}^l + w_{k\bar{l}}\xi^k\bar{\xi}^l,_i = w_{1\bar{l},i} + w_{1\bar{l}}(\xi^1,_i + \bar{\xi}^1,_i) = w_{1\bar{l}i},$$

Thus we have

$$\begin{aligned} (g^{p\bar{q}} w_{k\bar{q}} w_{p\bar{l}} \xi^k \bar{\xi}^l)_i &= g^{p\bar{q}} w_{k\bar{q}i} w_{p\bar{l}} \xi^k \bar{\xi}^l + g^{p\bar{q}} w_{k\bar{q}} w_{p\bar{l}i} \xi^k \bar{\xi}^l + g^{p\bar{q}} w_{k\bar{q}} w_{p\bar{l}} \xi^k,_i \bar{\xi}^l + g^{p\bar{q}} w_{k\bar{q}} w_{p\bar{l}} \xi^k \bar{\xi}^l,_i \\ &= w_{1\bar{l}}(w_{1\bar{l}i} + w_{1\bar{l}i}) + w_{1\bar{l}}^2(\xi^1,_i + \bar{\xi}^1,_i) \\ &= 2w_{1\bar{l}}w_{1\bar{l}i}. \end{aligned}$$

Therefore, we obtain the simplified formula for  $Q_i$  at  $x_0$ :

$$(4.7) \quad Q_i = \frac{w_{1\bar{l}i}}{w_{1\bar{l}}} + c_0 \frac{2w_{1\bar{l}i}}{w_{1\bar{l}}} + \varphi_i + \psi_i = (1 + 2c_0) \frac{w_{1\bar{l}i}}{w_{1\bar{l}}} + \varphi_i + \psi_i = 0$$

Similar calculations give

$$\begin{aligned} (w_{k\bar{l}}\xi^k\bar{\xi}^l)_{i\bar{i}} &= [w_{k\bar{l}i}\xi^k\bar{\xi}^l + w_{k\bar{l}}(\xi^k,_i\bar{\xi}^l + \xi^k\bar{\xi}^l,_i)]_{\bar{i}} \\ &= w_{k\bar{l}i\bar{i}}\xi^k\bar{\xi}^l + w_{k\bar{l}i}(\xi^k,_i\bar{\xi}^l + \xi^k\bar{\xi}^l,_i) + w_{k\bar{l}\bar{i}}(\xi^k,_i\bar{\xi}^l + \xi^k\bar{\xi}^l,_i) \\ &\quad + w_{k\bar{l}}(\xi^k,_i\bar{\xi}^l + \xi^k,_i\bar{\xi}^l + \xi^k,_i\bar{\xi}^l + \xi^k\bar{\xi}^l,_i) \\ &= w_{1\bar{l}i\bar{i}} + w_{k\bar{l}i}\xi^k,_i + w_{1\bar{l}i}\bar{\xi}^l,_i + w_{k\bar{l}\bar{i}}\xi^k,_i + w_{1\bar{l}\bar{i}}\bar{\xi}^l,_i \\ &\quad + w_{1\bar{l}}(\xi^1,_i\bar{i} + \bar{\xi}^1,_i) + w_{k\bar{k}}(\xi^k,_i\bar{\xi}^k,_i + \xi^k,_i\bar{\xi}^k,_i) \\ &= w_{1\bar{l}i\bar{i}} + 2 \sum_{k \neq 1} \operatorname{Re}(w_{k\bar{l}i}\xi^k,_i + w_{1\bar{k}i}\bar{\xi}^k,_i) + w_{1\bar{l}}(\xi^1,_i\bar{i} + \bar{\xi}^1,_i) \\ &\quad + w_{k\bar{k}}(|\xi^k,_i|^2 + |\xi^k,_i|^2). \end{aligned}$$

The last equality holds because we use (4-2) and (4-5) and the fact

$$w_{k\bar{l}i}\xi^k,_i + w_{1\bar{l}i}\bar{\xi}^l,_i = 2 \operatorname{Re}(w_{k\bar{l}i}\xi^k,_i), \quad w_{1\bar{l}i}\bar{\xi}^l,_i + w_{k\bar{l}i}\xi^k,_i = 2 \operatorname{Re}(w_{1\bar{k}i}\bar{\xi}^k,_i).$$

We can also calculate

$$\begin{aligned}
& (g^{p\bar{q}} w_{k\bar{q}} w_{p\bar{l}} \xi^k \bar{\xi}^l)_{i\bar{i}} \\
&= g^{p\bar{q}} (w_{k\bar{q}i} w_{p\bar{l}} \xi^k \bar{\xi}^l + w_{k\bar{q}} w_{p\bar{l}i} \xi^k \bar{\xi}^l + w_{k\bar{q}} w_{p\bar{l}} \xi^k_i \bar{\xi}^l + w_{k\bar{q}} w_{p\bar{l}} \xi^k \bar{\xi}^l_i)_{\bar{i}} \\
&= g^{p\bar{q}} (w_{k\bar{q}i\bar{i}} w_{p\bar{l}} \xi^k \bar{\xi}^l + w_{k\bar{q}i} w_{p\bar{l}\bar{i}} \xi^k \bar{\xi}^l + w_{k\bar{q}i} w_{p\bar{l}} \xi^k_i \bar{\xi}^l + w_{k\bar{q}i} w_{p\bar{l}} \xi^k \bar{\xi}^l_i) \\
&\quad + g^{p\bar{q}} (w_{k\bar{q}i} w_{p\bar{l}i} \xi^k \bar{\xi}^l + w_{k\bar{q}} w_{p\bar{l}i} \xi^k \bar{\xi}^l + w_{k\bar{q}} w_{p\bar{l}i} \xi^k_i \bar{\xi}^l + w_{k\bar{q}} w_{p\bar{l}i} \xi^k \bar{\xi}^l_i) \\
&\quad + g^{p\bar{q}} (w_{k\bar{q}i} w_{p\bar{l}} \xi^k_i \bar{\xi}^l + w_{k\bar{q}} w_{p\bar{l}i} \xi^k_i \bar{\xi}^l + w_{k\bar{q}} w_{p\bar{l}} \xi^k_{i\bar{i}} \bar{\xi}^l + w_{k\bar{q}} w_{p\bar{l}} \xi^k_i \bar{\xi}^l_{\bar{i}}) \\
&\quad + g^{p\bar{q}} (w_{k\bar{q}i} w_{p\bar{l}} \xi^k \bar{\xi}^l_i + w_{k\bar{q}} w_{p\bar{l}i} \xi^k \bar{\xi}^l_i + w_{k\bar{q}} w_{p\bar{l}} \xi^k_i \bar{\xi}^l_i + w_{k\bar{q}} w_{p\bar{l}} \xi^k \bar{\xi}^l_{i\bar{i}}) \\
&= w_{1\bar{1}i\bar{i}} w_{1\bar{1}} + w_{1\bar{p}i} w_{p\bar{1}\bar{i}} + w_{k\bar{1}i} w_{1\bar{1}} \xi^k_i + w_{1\bar{p}i} w_{p\bar{p}} \bar{\xi}^p_{\bar{i}} \\
&\quad + w_{1\bar{p}i} w_{p\bar{1}i} + w_{1\bar{1}} w_{1\bar{1}i\bar{i}} + w_{p\bar{p}} w_{p\bar{1}i} \xi^p_{\bar{i}} + w_{1\bar{1}} w_{1\bar{1}i} \bar{\xi}^l_{\bar{i}} \\
&\quad + w_{k\bar{1}i} w_{1\bar{1}} \xi^k_i + w_{p\bar{p}} w_{p\bar{1}i} \xi^p_{\bar{i}} + w_{1\bar{1}}^2 \xi^1_{i\bar{i}} + w_{p\bar{p}}^2 \xi^p_{\bar{i}} \bar{\xi}^p_{\bar{i}} \\
&\quad + w_{1\bar{p}i} w_{p\bar{p}} \bar{\xi}^p_{\bar{i}} + w_{1\bar{1}} w_{1\bar{1}i} \bar{\xi}^l_{\bar{i}} + w_{p\bar{p}}^2 \xi^p_{\bar{i}} \bar{\xi}^p_{\bar{i}} + w_{1\bar{1}}^2 \bar{\xi}^1_{i\bar{i}} \\
&= 2w_{1\bar{1}} w_{1\bar{1}i\bar{i}} + |w_{1\bar{p}i}|^2 + |w_{1\bar{p}i}|^2 + 2w_{1\bar{1}} \operatorname{Re}(w_{p\bar{1}i} \xi^p_{\bar{i}} + w_{p\bar{1}i} \xi^p_{\bar{i}}) \\
&\quad + 2w_{p\bar{p}} \operatorname{Re}(w_{1\bar{p}i} \bar{\xi}^p_{\bar{i}} + w_{p\bar{1}i} \xi^p_{\bar{i}}) + w_{p\bar{p}}^2 (|\xi^p_{\bar{i}}|^2 + |\xi^p_{\bar{i}}|^2) + w_{1\bar{1}}^2 (\xi^1_{i\bar{i}} + \bar{\xi}^1_{i\bar{i}})
\end{aligned}$$

Therefore we simplify  $Q_{i\bar{i}}$  at  $x_0$  as follows

$$\begin{aligned}
Q_{i\bar{i}} &= (1 + 2c_0) \frac{w_{1\bar{1}i\bar{i}}}{w_{1\bar{1}}} + \frac{c_0}{w_{1\bar{1}}^2} \sum_{p \neq 1} (|w_{1\bar{p}i}|^2 + |w_{1\bar{p}i}|^2) \\
&\quad - (1 + 2c_0) \frac{|w_{1\bar{1}i}|^2}{w_{1\bar{1}}^2} + (**)_{i\bar{i}} + \varphi_{i\bar{i}} + \psi_{i\bar{i}},
\end{aligned}$$

where  $(**)_{i\bar{i}}$  is given by

$$\begin{aligned}
(**)_{i\bar{i}} &= \frac{2}{w_{1\bar{1}}} \sum_{k \neq 1} \operatorname{Re}(w_{k\bar{1}i} \xi^k_{\bar{i}} + w_{1\bar{k}i} \bar{\xi}^k_{\bar{i}}) + \xi^1_{i\bar{i}} + \bar{\xi}^1_{i\bar{i}} + \frac{w_{k\bar{k}}}{w_{1\bar{1}}} (|\xi^k_{\bar{i}}|^2 + |\xi^k_{\bar{i}}|^2) \\
&\quad + \frac{2c_0}{w_{1\bar{1}}} \sum_{p \neq 1} \operatorname{Re}(w_{p\bar{1}i} \xi^p_{\bar{i}} + w_{p\bar{1}i} \bar{\xi}^p_{\bar{i}}) + \sum_{p \neq 1} \frac{2c_0 w_{p\bar{p}}}{w_{1\bar{1}}^2} \operatorname{Re}(w_{1\bar{p}i} \bar{\xi}^p_{\bar{i}} + w_{p\bar{1}i} \xi^p_{\bar{i}}) \\
&\quad + \frac{2c_0 w_{p\bar{p}}^2}{w_{1\bar{1}}^2} (|\xi^p_{\bar{i}}|^2 + |\xi^p_{\bar{i}}|^2) + c_0 (\xi^1_{i\bar{i}} + \bar{\xi}^1_{i\bar{i}}).
\end{aligned}$$

For this term  $(**)_{i\bar{i}}$ , we have the estimate

$$(**)_{i\bar{i}} \geq -\frac{c_0}{2w_{1\bar{1}}^2} \sum_{p \neq 1} (|w_{1\bar{p}i}|^2 + |w_{1\bar{p}i}|^2) - C,$$

where  $C$  is a positive constant depending only on  $(M, \omega)$ .

Let  $F(\omega_u) = (\sigma_k(\omega_u))^{1/k}$ . We denote by

$$F^{i\bar{j}} = \frac{\partial F}{\partial w_{i\bar{j}}}, \quad F^{i\bar{j}, p\bar{q}} = \frac{\partial^2 F}{\partial w_{i\bar{j}} \partial w_{p\bar{q}}},$$

where  $(w_u)_{i\bar{j}} = g_{i\bar{j}} + u_{i\bar{j}}$ . Then, the positive definite matrix  $(F^{i\bar{j}}(\omega_u))$  is diagonalized at the point  $x_0$ . More precisely,

$$(4-8) \quad F^{i\bar{j}}(\omega_u) = \delta_{ij} F^{i\bar{i}}(\omega_u) = \frac{1}{k} [\sigma_k(\lambda)]^{1/k-1} \sigma_{k-1}(\lambda|i) \delta_{ij}.$$

Furthermore, at  $x_0$ ,

$$(4-9) \quad F^{i\bar{j}, p\bar{q}}(\omega_u) = \begin{cases} F^{i\bar{i}, p\bar{p}}, & \text{if } i = j, p = q; \\ F^{i\bar{p}, p\bar{i}}, & \text{if } i = q, p = j, i \neq p; \\ 0, & \text{otherwise,} \end{cases}$$

in which

$$\begin{aligned} F^{i\bar{i}, p\bar{p}} &= \frac{1}{k} [\sigma_k(\lambda)]^{1/k-1} (1 - \delta_{ip}) \sigma_{k-2}(\lambda|ip) \\ &\quad + \frac{1}{k} \left( \frac{1}{k} - 1 \right) [\sigma_k(\lambda)]^{1/k-2} \sigma_{k-1}(\lambda|i) \sigma_{k-1}(\lambda|p), \\ F^{i\bar{p}, p\bar{i}} &= -\frac{1}{k} [\sigma_k(\lambda)]^{1/k-1} \sigma_{k-2}(\lambda|ip). \end{aligned}$$

We have, in addition, at  $x_0$

$$(4-10) \quad \sum_{i=1}^n F^{i\bar{i}} w_{i\bar{i}} = \sum_{i=1}^n F^{i\bar{i}} \lambda_i = \sigma_k^{1/k} = e^{f/k}.$$

By the maximum principal, we have

$$\begin{aligned} (4-11) \quad 0 &\geq F^{i\bar{j}} Q_{i\bar{j}} = F^{i\bar{i}} Q_{i\bar{i}} \\ &\geq (1 + 2c_0) \sum_{i=1}^n \frac{F^{i\bar{i}} u_{1\bar{i}i\bar{i}}}{w_{1\bar{i}}} + \frac{c_0}{2} \sum_{i=1}^n \sum_{p \neq 1} \frac{F^{i\bar{i}} |u_{1\bar{p}i}|^2}{w_{1\bar{i}}^2} \\ &\quad - (1 + 2c_0) \sum_{i=1}^n \frac{F^{i\bar{i}} |u_{1\bar{i}i}|^2}{w_{1\bar{i}}^2} + \psi' \sum_{i=1}^n F^{i\bar{i}} u_{i\bar{i}} + \psi'' \sum_{i=1}^n F^{i\bar{i}} |u_i|^2 \\ &\quad + \varphi'' \sum_{i=1}^n F^{i\bar{i}} |\nabla u|_i^2 |\nabla u|_{\bar{i}}^2 + \varphi' \sum_{i,p=1}^n F^{i\bar{i}} (|u_{p\bar{i}}|^2 + |u_{pi}|^2) \\ &\quad + \varphi' \sum_{i,p=1}^n F^{i\bar{i}} (u_{p\bar{i}i} u_{\bar{p}} + u_{\bar{p}ii} u_p) - C_1 \sum_{i=1}^n F^{i\bar{i}} \\ &:= I_1 + I_2 + I_3 + I_4 + I_5 + I_6 + I_7 + I_8 - C_1 \sum_{i=1}^n F^{i\bar{i}} \end{aligned}$$

The equation can be written as  $F(\omega_u) = e^{f/k} := h$ . Differentiating this, we get

$$\sum_{i,j=1}^n F^{i\bar{j}} u_{i\bar{j}l} = \nabla_l F = h_l, \quad \sum_{i,j=1}^n F^{i\bar{j}} u_{i\bar{j}l\bar{m}} + \sum_{i,j,p,q=1}^n F^{i\bar{j}, p\bar{q}} u_{i\bar{j}l} u_{p\bar{q}\bar{m}} = h_{l\bar{m}}.$$

and

$$\sum_{i=1}^n F^{i\bar{i}} u_{i\bar{i}1\bar{1}} = h_{1\bar{1}} - \sum_{i,j,p,q=1}^n F^{i\bar{j}, p\bar{q}} u_{i\bar{j}1} u_{p\bar{q}\bar{1}}.$$

By commuting the covariant derivatives formula (2-12), we have

$$(4-12) \quad \begin{aligned} \sum_{i=1}^n F^{i\bar{i}} u_{1\bar{l}i\bar{i}} &= \sum_{i=1}^n F^{i\bar{i}} u_{i\bar{i}1\bar{1}} + \sum_{i=1}^n F^{i\bar{i}} (u_{1\bar{1}} - \sum_{i=1}^n u_{i\bar{i}}) R_{i\bar{i}1\bar{1}} \\ &\quad + \sum_{i=1}^n F^{i\bar{i}} \left( \sum_{p=1}^n T_{1i}^p u_{p\bar{l}\bar{i}} + \sum_{q=1}^n \bar{T}_{1i}^q u_{1\bar{q}i} - \sum_{p=1}^n |T_{1i}^p|^2 u_{p\bar{p}} \right). \end{aligned}$$

Inserting (4-12) into the term  $I_1$ , we have

$$(4-13) \quad \begin{aligned} I_1 &= (1+2c_0) \sum_{i=1}^n \frac{F^{i\bar{i}} u_{1\bar{l}i\bar{i}}}{w_{1\bar{l}}} \\ &= (1+2c_0) \sum_{i=1}^n \frac{F^{i\bar{i}} u_{i\bar{i}1\bar{1}}}{w_{1\bar{l}}} + (1+2c_0) \sum_{i=1}^n \frac{F^{i\bar{i}} (u_{1\bar{1}} - u_{i\bar{i}}) R_{i\bar{i}1\bar{1}}}{w_{1\bar{l}}} \\ &\quad + 2(1+2c_0) \sum_{i,p=1}^n F^{i\bar{i}} \operatorname{Re} \left( \frac{T_{1i}^p u_{p\bar{l}\bar{i}}}{w_{1\bar{l}}} \right) - (1+2c_0) \sum_{i,p=1}^n F^{i\bar{i}} \frac{|T_{1i}^p|^2 u_{p\bar{p}}}{w_{1\bar{l}}} \\ &= (1+2c_0) \frac{h_{1\bar{1}}}{w_{1\bar{l}}} - (1+2c_0) \sum_{i,j,p,q=1}^n \frac{F^{i\bar{j}, p\bar{q}} u_{i\bar{j}1} u_{p\bar{q}\bar{1}}}{w_{1\bar{l}}} \\ &\quad + (1+2c_0) \sum_{i=1}^n \frac{F^{i\bar{i}} (u_{1\bar{1}} - u_{i\bar{i}}) R_{i\bar{i}1\bar{1}}}{w_{1\bar{l}}} + 2(1+2c_0) \sum_i^n F^{i\bar{i}} \operatorname{Re} \left( \frac{T_{1i}^1 u_{1\bar{l}\bar{i}}}{w_{1\bar{l}}} \right) \\ &\quad + 2(1+2c_0) \sum_{i=1}^n F^{i\bar{i}} \operatorname{Re} \left( \sum_{p \neq 1}^n \frac{T_{1i}^p u_{p\bar{l}\bar{i}}}{w_{1\bar{l}}} \right) - (1+2c_0) \sum_{i,p=1}^n F^{i\bar{i}} \frac{|T_{1i}^p|^2 u_{p\bar{p}}}{w_{1\bar{l}}} \\ &:= I_{11} + I_{12} + I_{13} + I_{14} + I_{15} + I_{16}. \end{aligned}$$

We estimate each term in this sum. First we have

$$I_{11} + I_{13} + I_{16} \geq -C_1 - 3(nC_2 + C_3) \sum_{i=1}^n F^{i\bar{i}},$$

where we have supposed that  $\sup_M |T|^2_g \leq C_2$ ,  $\sup_M |R| \leq C_3$ .

We claim  $I_{15} + I_2 \geq -18n^2C_2 \sum_{i=1}^n F^{i\bar{i}}$ . Indeed, since  $\frac{1}{n^2} \leq c_0 \leq 1$ , we have

$$\begin{aligned} I_{15} + I_2 &= \frac{c_0}{2} \sum_{i=1}^n \sum_{p \neq 1} \frac{F^{i\bar{i}} |u_{1\bar{p}i}|^2}{w_{1\bar{1}}^2} + 2(1+2c_0) \sum_{i=1}^n F^{i\bar{i}} \operatorname{Re} \left( \sum_{p \neq 1} \frac{T_{1i}^p u_{p\bar{i}}}{w_{1\bar{1}}} \right) \\ &= \frac{c_0}{2} \sum_{i=1}^n F^{i\bar{i}} \sum_{p \neq 1} \left| \frac{u_{1\bar{p}i}}{w_{1\bar{1}}} + \frac{2(1+2c_0)}{c_0} T_{1i}^p \right|^2 - \frac{2(1+2c_0)^2}{c_0} \sum_{i=1}^n \sum_{p \neq 1} F^{i\bar{i}} |T_{1i}^p|^2 \\ &\geq -\frac{2(1+2c_0)^2}{c_0} \sum_{i=1}^n \sum_{p \neq 1} F^{i\bar{i}} |T_{1i}^p|^2 \\ &\geq -18n^2 C_2 \sum_{i=1}^n F^{i\bar{i}}. \end{aligned}$$

Then we obtain

$$\begin{aligned} (4-14) \quad I_1 + I_2 &\geq -(1+2c_0) \sum_{i,j,p,q=1}^n \frac{F^{i\bar{j}, p\bar{q}} u_{ij} u_{pq\bar{1}}}{w_{1\bar{1}}} + 2(1+2c_0) \sum_{i=1}^n F^{i\bar{i}} \operatorname{Re} \left( \frac{T_{1i}^1 u_{1\bar{i}}}{w_{1\bar{1}}} \right) \\ &\quad - (21n^2 C_2 + 3C_3) \sum_{i=1}^n F^{i\bar{i}} - C_1. \end{aligned}$$

For terms  $I_7 + I_8$ , we claim

$$(4-15) \quad I_7 + I_8 \geq \frac{1}{2} \varphi' \sum_{i=1}^n F^{i\bar{i}} |u_{ii}|_1^2 - (C_2 + C_3) \sum_{i=1}^n F^{i\bar{i}} - C_1.$$

Indeed, by the commutation formula for covariant derivatives (2-11) in Section 2,

$$u_{pi\bar{i}} = u_{i\bar{i}p} + T_{pi}^i u_{i\bar{i}} + u_q R_{i\bar{i}p\bar{q}}, \quad u_{\bar{p}i\bar{i}} = u_{i\bar{p}\bar{i}} = u_{i\bar{i}\bar{p}} - \overline{T_{ip}^i} u_{i\bar{i}}.$$

Then

$$\begin{aligned} \sum_{i=1}^n F^{i\bar{i}} u_{pi\bar{i}} &= \sum_{i=1}^n F^{i\bar{i}} u_{i\bar{i}p} + \sum_{i=1}^n F^{i\bar{i}} (T_{pi}^i u_{i\bar{i}} + u_q R_{i\bar{i}p\bar{q}}) \\ &= h_p + \sum_{i=1}^n F^{i\bar{i}} (T_{pi}^i u_{i\bar{i}} + u_q R_{i\bar{i}p\bar{q}}) \\ \sum_{i=1}^n F^{i\bar{i}} u_{\bar{p}i\bar{i}} &= \sum_{i=1}^n F^{i\bar{i}} u_{i\bar{i}\bar{p}} + \sum_{i=1}^n F^{i\bar{i}} \overline{T_{ip}^i} u_{i\bar{i}} = h_{\bar{p}} + \sum_{i=1}^n F^{i\bar{i}} \overline{T_{ip}^i} u_{i\bar{i}} \end{aligned}$$

Inserting the above formula into  $I_8$ , we obtain

$$\begin{aligned}
 (4-16) \quad I_8 &= \varphi' \sum_{i,p=1}^n F^{i\bar{i}} (u_{p\bar{i}i} u_{\bar{p}} + u_{\bar{p}i\bar{i}} u_p) \\
 &= \varphi' \sum_{p=1}^n u_{\bar{p}} \left[ h_p + \sum_{i=1}^n F^{i\bar{i}} (T_{pi}^i u_{i\bar{i}} + u_q R_{i\bar{i}p\bar{q}}) \right] + \varphi' \sum_{p=1}^n u_p \left[ h_{\bar{p}} - \sum_{i=1}^n F^{i\bar{i}} \overline{T_{ip}^i} u_{i\bar{i}} \right] \\
 &= 2\varphi' \sum_{i,p=1}^n F^{i\bar{i}} u_{i\bar{i}} \operatorname{Re}(u_{\bar{p}} T_{pi}^i) + \varphi' \sum_{p=1}^n \left[ 2 \operatorname{Re}(u_{\bar{p}} h_p) + \sum_{i,q=1}^n u_{\bar{p}} u_q F^{i\bar{i}} R_{i\bar{i}p\bar{q}} \right] \\
 &= I_{81} + I_{82}.
 \end{aligned}$$

For the term  $I_{82}$ , we have

$$I_{82} \geq -C_3 \sum_{i=1}^n F^{i\bar{i}} - C_1.$$

For the term  $I_{81}$ , we obtain

$$\begin{aligned}
 I_{81} + I_7 &= 2\varphi' \sum_{i,p=1}^n F^{i\bar{i}} u_{i\bar{i}} \operatorname{Re}(u_{\bar{p}} T_{pi}^i) + \varphi' \sum_{i,p=1}^n F^{i\bar{i}} (|u_{p\bar{i}}|^2 + |u_{pi}|^2) \\
 &\geq \varphi' \sum_{i=1}^n F^{i\bar{i}} \left[ |u_{i\bar{i}}|^2 + 2u_{i\bar{i}} \operatorname{Re} \left( \sum_{p=1}^n u_{\bar{p}} T_{pi}^i \right) \right] \\
 &= \varphi' \sum_{i=1}^n F^{i\bar{i}} \left| \frac{u_{i\bar{i}}}{2} + 2 \sum_{p=1}^n u_p \overline{T_{pi}^i} \right|^2 + \frac{3}{4} \varphi' \sum_{i=1}^n F^{i\bar{i}} |u_{i\bar{i}}|^2 - 4\varphi' \sum_{i=1}^n F^{i\bar{i}} \left| \sum_{p=1}^n u_p \overline{T_{pi}^i} \right|^2 \\
 &\geq \frac{1}{2} \varphi' \sum_{i=1}^n F^{i\bar{i}} |u_{i\bar{i}}|^2 - C_2 \sum_{i=1}^n F^{i\bar{i}}.
 \end{aligned}$$

Thus we have proved the above claim (4-15). Moreover, applying (4-10) yields

$$\begin{aligned}
 \psi' \sum_{i=1}^n F^{i\bar{i}} u_{i\bar{i}} &= \psi' \sum_{i=1}^n F^{i\bar{i}} (\lambda_i - 1) \\
 &= \psi' h - \psi' \sum_{i=1}^n F^{i\bar{i}} \geq -2(C_0 + 1) \sup_M e^{f/k} - \psi' \sum_{i=1}^n F^{i\bar{i}}.
 \end{aligned}$$

Similarly,

$$\begin{aligned}
\frac{1}{2}\varphi' \sum_{i=1}^n F^{i\bar{i}} |u_{i\bar{i}}|^2 &= \frac{1}{2}\varphi' \sum_{i=1}^n F^{i\bar{i}} (\lambda_i - 1)^2 \\
&= \frac{1}{2}\varphi' \sum_{i=1}^n F^{i\bar{i}} \lambda_i^2 - \varphi' \sum_{i=1}^n F^{i\bar{i}} \lambda_i + \frac{1}{2}\varphi' \sum_{i=1}^n F^{i\bar{i}} \\
&= \frac{1}{2}\varphi' \sum_{i=1}^n F^{i\bar{i}} \lambda_i^2 - \varphi' h + \frac{1}{2}\varphi' \sum_{i=1}^n F^{i\bar{i}} \\
&\geq \frac{1}{2}\varphi' \sum_{i=1}^n F^{i\bar{i}} \lambda_i^2 - \frac{1}{2} \sup_M e^{f/k} + \frac{1}{2}\varphi' \sum_{i=1}^n F^{i\bar{i}}.
\end{aligned}$$

Inserting these terms into (4-11), we obtain

$$\begin{aligned}
(4-17) \quad 0 &\geq F^{i\bar{i}} Q_{i\bar{i}} \\
&\geq -(1+2c_0) \sum_{i,j,p,q=1}^n \frac{F^{ij\bar{j}, p\bar{q}} u_{ij\bar{1}} u_{p\bar{q}\bar{1}}}{w_{1\bar{1}}} + 2(1+2c_0) \sum_{i=1}^n F^{i\bar{i}} \operatorname{Re} \left( \frac{T_{1i}^1 u_{1\bar{1}\bar{i}}}{w_{1\bar{1}}} \right) \\
&\quad - (1+2c_0) \sum_{i=1}^n \frac{F^{i\bar{i}} |u_{1\bar{1}\bar{i}}|^2}{w_{1\bar{1}}^2} \\
&\quad + \varphi'' \sum_{i=1}^n F^{i\bar{i}} |\nabla u|_i^2 |\nabla u|_{\bar{i}}^2 + \psi'' \sum_{i=1}^n F^{i\bar{i}} |u_i|^2 + \frac{1}{2}\varphi' \sum_{i=1}^n F^{i\bar{i}} \lambda_i^2 \\
&\quad + \left( -\psi' + \frac{1}{2}\varphi' - 22n^2 C_2 - 4C_3 \right) \sum_{i=1}^n F^{i\bar{i}} - C_1 \\
&= A_1 + A_2 + A_3 + A_4 + A_5 + A_6 \\
&\quad + \left( -\psi' + \frac{1}{2}\varphi' - 22n^2 C_2 - 4C_3 \right) \sum_{i=1}^n F^{i\bar{i}} - C_1,
\end{aligned}$$

where  $C_1$  is a positive constant depending only on  $C_0$ ,  $\sup e^{f/k}$ ,  $\sup |\nabla(e^{f/k})|^2$  and  $\sup |\partial\bar{\partial}(e^{f/k})|$ .

Let  $\varepsilon = \frac{1}{4}\delta \leq \frac{1}{16}$  and  $\delta = 1/(2A + 1)$ , where  $A = 2L(C_0 + 1)$  and  $C_0 = 31n^2 C_2 + 4C_3$ . We divide into two cases to derive the estimate, which is similar to [Hou et al. 2010].

**Case 1:**  $\lambda_n < -\varepsilon\lambda_1$ .

By condition (4-7), for  $1 \leq i \leq n$ , we have

$$\begin{aligned}
-(1+2c_0)^2 \left| \frac{u_{1\bar{1}i}}{w_{1\bar{1}}} \right|^2 &= -|\varphi'| |\nabla u|_i^2 + \psi' |u_i|^2 \geq -2(\varphi')^2 |\nabla u|_i^2 |\nabla u|_{\bar{i}}^2 - 2(\psi')^2 |u_i|^2 \\
&= -\varphi'' |\nabla u|_i^2 |\nabla u|_{\bar{i}}^2 - 2(\psi')^2 |u_i|^2.
\end{aligned}$$

This gives

$$\begin{aligned} A_2 &= 2(1+2c_0) \sum_{i \neq 1} F^{ii} \operatorname{Re} \left( \frac{T_{1i}^1 u_{1\bar{i}}}{w_{1\bar{i}}} \right) \\ &\geq -c_0 \sum_{i \neq 1} F^{ii} \left| \frac{u_{1\bar{i}}}{w_{1\bar{i}}} \right|^2 - \frac{(1+2c_0)^2}{c_0} \sum_{i \neq 1} F^{ii} |T_{1i}^1|^2 \\ &\geq -c_0 \sum_{i \neq 1} F^{ii} \left| \frac{u_{1\bar{i}}}{w_{1\bar{i}}} \right|^2 - 9n^2 C_2 \sum_{i \neq 1} F^{ii} |T_{1i}^1|^2 \end{aligned}$$

Thus

$$\begin{aligned} A_2 + A_3 &\geq -(1+3c_0) \sum_{i=1}^n \frac{F^{ii} |u_{1\bar{i}}|^2}{w_{1\bar{i}}^2} - 9n^2 C_2 \sum_{i \neq 1} F^{ii} |T_{1i}^1|^2 \\ &\geq -(1+2c_0)^2 \sum_{i=1}^n \frac{F^{ii} |u_{1\bar{i}}|^2}{w_{1\bar{i}}^2} - 9n^2 C_2 \sum_{i=1}^n F^{ii} \\ &= -A_4 - 2(\psi')^2 \sum_{i=1}^n F^{ii} |u_i|^2 - 9n^2 C_2 \sum_{i=1}^n F^{ii}. \end{aligned}$$

We therefore obtain

$$(4-18) \quad A_2 + A_3 + A_4 \geq -2(\psi')^2 \sum_{i=1}^n F^{ii} |u_i|^2 - 9n^2 C_2 \sum_{i=1}^n F^{ii}.$$

Using the inequality

$$\sum_{i=1}^n F^{ii} \lambda_i^2 \geq F^{n\bar{n}} \lambda_n^2 > \varepsilon^2 F^{n\bar{n}} \lambda_1^2 \geq \frac{\varepsilon^2}{n} \sum_{i=1}^n F^{ii} \lambda_1^2,$$

we have

$$(4-19) \quad A_6 = \frac{1}{2} \varphi' \sum_{i=1}^n F^{ii} \lambda_i^2 \geq \frac{\varepsilon^2}{2n} \varphi' \sum_{i=1}^n F^{ii} \lambda_1^2.$$

Combining (4-17) and (4-18) (4-19), we obtain

$$\begin{aligned} 0 &\geq \sum_{i=1}^n F^{ii} Q_{i\bar{i}} \geq \frac{\varepsilon^2}{2n} \varphi' \sum_{i=1}^n F^{ii} \lambda_1^2 - 2(\psi')^2 \sum_{i=1}^n F^{ii} |u_i|^2 \\ &\quad + \left( -\psi' + \frac{1}{2} \varphi' - 31n^2 C_2 - 4C_3 \right) \sum_{i=1}^n F^{ii} - C_1 \\ &\geq \left( \frac{\varepsilon^2}{8nK} \lambda_1^2 - 8K(C_0 + 1)^2 \right) \sum_{i=1}^n F^{ii} - C_1 \\ &\geq \frac{\varepsilon^2}{8nK} \lambda_1^2 - 8K(C_0 + 1)^2 - C_1, \end{aligned}$$

where we use the fact that  $\sum_{i=1}^n F^{ii} \geq 1$ , which follows from the definition of  $F^{ii}$  and the Newton–Maclaurin inequality.

Hence, we obtain

$$\lambda_1 \leq 8\sqrt{2}(2A+1)\sqrt{nK(8K(C_0+1)^2+C_1)} \leq CK.$$

**Case 2:**  $\lambda_n > -\varepsilon\lambda_1$ .

Let  $I = \{i \in \{1, \dots, n\} | \sigma_{k-1}(\lambda|i) \geq \varepsilon^{-1}\sigma_{k-1}(\lambda|1)\}$ . Obviously,  $1 \notin I$  and  $i \in I$  if and only if  $F^{ii} > \varepsilon^{-1}F^{1\bar{1}}$ . We first treat those indices which are not in  $I$ . By (4-7), we have

$$\begin{aligned} & -(1+2c_0) \sum_{i \notin I} \frac{F^{i\bar{i}}|u_{1\bar{i}}|^2}{w_{1\bar{i}}^2} + 2(1+2c_0) \sum_{i \notin I} F^{i\bar{i}} \operatorname{Re} \frac{T_{1i}^1 u_{1\bar{i}}}{w_{1\bar{i}}} \\ & \geq -(1+2c_0)^2 \sum_{i \notin I} \frac{F^{i\bar{i}}|u_{1\bar{i}}|^2}{w_{1\bar{i}}^2} - \frac{(1+2c_0)^2}{c_0} \sum_{i \notin I} F^{i\bar{i}} |T_{1i}^1|^2 \\ & = -\varphi'' \sum_{i \notin I} F^{i\bar{i}} |\nabla u|_i^2 |\nabla u|_{\bar{i}}^2 - 2(\psi')^2 \sum_{i \notin I} F^{i\bar{i}} |u_i|^2 - 9n^2 C_2 \sum_{i \notin I} F^{i\bar{i}} |T_{1i}^1|^2 \\ & \geq -\varphi'' \sum_{i \notin I} F^{i\bar{i}} |\nabla u|_i^2 |\nabla u|_{\bar{i}}^2 - 2\varepsilon^{-1} K(\psi')^2 F^{1\bar{1}} - 9n^2 C_2 \sum_{i=1}^n F^{i\bar{i}}. \end{aligned}$$

Substituting the above inequality into (4-17) yields

$$\begin{aligned} (4-20) \quad 0 & \geq F^{i\bar{i}} Q_{i\bar{i}} \\ & \geq -(1+2c_0) \sum_{i,j,p,q=1}^n \frac{F^{i\bar{j}, p\bar{q}} u_{i\bar{j}} u_{p\bar{q}}}{w_{1\bar{i}}} + 2(1+2c_0) \sum_{i \in I} F^{i\bar{i}} \operatorname{Re} \left( \frac{T_{1i}^1 u_{1\bar{i}}}{w_{1\bar{i}}} \right) \\ & \quad - (1+2c_0) \sum_{i \in I} \frac{F^{i\bar{i}} |u_{1\bar{i}}|^2}{w_{1\bar{i}}^2} + \varphi'' \sum_{i \in I} F^{i\bar{i}} |\nabla u|_i^2 |\nabla u|_{\bar{i}}^2 + \psi'' \sum_{i=1}^n F^{i\bar{i}} |u_i|^2 \\ & \quad + \frac{1}{2} \varphi' \sum_{i=1}^n F^{i\bar{i}} \lambda_i^2 - 2\varepsilon^{-1} K(\psi')^2 F^{1\bar{1}} \\ & \quad + \left( -\psi' + \frac{1}{2} \varphi' - 31n^2 C_2 - 4C_3 \right) \sum_{i=1}^n F^{i\bar{i}} - C_1 \\ & = B_1 + B_2 + B_3 + B_4 + B_5 + B_6 + B_7 + B_8. \end{aligned}$$

Firstly, we have

$$B_6 + B_7 = \frac{1}{2} \varphi' \sum_{i=1}^n F^{i\bar{i}} \lambda_i^2 - 2\varepsilon^{-1} K(\psi')^2 F^{1\bar{1}} \geq \frac{1}{4} \varphi' \sum_{i=1}^n F^{i\bar{i}} \lambda_i^2,$$

where we assume that  $\frac{1}{4}\varphi' F^{1\bar{1}} \lambda_1^2 \geq 2\varepsilon^{-1} K(\psi')^2 F^{1\bar{1}}$  (for otherwise  $\frac{1}{4}\varphi' F^{1\bar{1}} \lambda_1^2 \leq 2\varepsilon^{-1} K(\psi')^2 F^{1\bar{1}}$ , i.e.,  $\lambda_1 \leq CK$ ).

We next use  $B_1$  to cancel the other terms containing the third derivatives of  $u$ .

As the proof in [Hou et al. 2010, p. 559], we have

$$\lambda_1 \sigma_{k-2}(\lambda|1i) \geq (1 - 2\varepsilon) \sigma_{k-1}(\lambda|i) \quad \text{for } i \in I.$$

Then

$$-\lambda_1 F^{i\bar{1}, 1\bar{i}} = \frac{F^{1-k}}{k} \lambda_1 \sigma_{k-2}(\lambda|1i) \geq \frac{F^{1-k}}{k} (1 - 2\varepsilon) \sigma_{k-1}(\lambda|i) = (1 - 2\varepsilon) F^{i\bar{i}}.$$

Since  $u_{i\bar{1}1} = u_{1\bar{1}i} - T_{1i}^1(\lambda_1 - 1)$ , we get

$$\begin{aligned} B_1 &= -\frac{1+2c_0}{\lambda_1} \sum_{i,j,p,q=1}^n F^{ij, p\bar{q}} u_{ij1} u_{p\bar{q}\bar{1}} \\ &\geq -\frac{1+2c_0}{\lambda_1^2} \sum_{i \in I} \lambda_1 F^{i\bar{1}, 1\bar{i}} u_{i\bar{1}1} u_{1\bar{i}\bar{1}} \\ &\geq \frac{1+2c_0}{\lambda_1^2} (1 - 2\varepsilon) \sum_{i \in I} F^{i\bar{i}} |u_{1\bar{1}i} - T_{1i}^1(\lambda_1 - 1)|^2, \end{aligned}$$

and

$$B_2 = \frac{2(1+2c_0)}{\lambda_1} \sum_{i \in I} F^{i\bar{i}} \operatorname{Re}(T_{1i}^1 u_{1\bar{1}i}).$$

From (4-7), we have

$$\begin{aligned} B_4 &= \varphi'' \sum_{i \in I} F^{i\bar{i}} |\nabla u|_i^2 |\nabla u|_{\bar{i}}^2 \\ &= 2 \sum_{i \in I} F^{i\bar{i}} \left| (1 + 2c_0) \frac{u_{1\bar{1}i}}{w_{1\bar{1}}} + \psi' u_i \right|^2 \\ &\geq 2(1 + 2c_0)^2 \delta \sum_{i \in I} F^{i\bar{i}} \frac{|u_{1\bar{1}i}|^2}{w_{1\bar{1}}^2} - \frac{2\delta}{1 - \delta} (\psi')^2 \sum_{i \in I} F^{i\bar{i}} |u_i|^2 \\ &\geq 2(1 + 2c_0)^2 \delta \sum_{i \in I} F^{i\bar{i}} \frac{|u_{1\bar{1}i}|^2}{w_{1\bar{1}}^2} - B_5, \end{aligned}$$

where we use  $(2\delta/(1 - \delta))(\psi')^2 = \psi''$  by choosing  $\delta = 1/(2A + 1)$ .

So we get

$$B_3 + B_4 + B_5 \geq -(1 + 2c_0) \frac{[1 - 2(1 + 2c_0)\delta]}{\lambda_1^2} \sum_{i \in I} F^{i\bar{i}} |u_{1\bar{1}i}|^2.$$

Then we conclude

$$\begin{aligned}
& B_1 + B_2 + B_3 + B_4 + B_5 \\
& \geq \frac{1+2c_0}{\lambda_1^2} (1-2\varepsilon) \sum_{i \in I} F^{i\bar{i}} |u_{1\bar{i}} - T_{1i}^1(\lambda_1 - 1)|^2 \\
& \quad - (1+2c_0) \frac{[1-2(1+2c_0)\delta]}{\lambda_1^2} \sum_{i \in I} F^{i\bar{i}} |u_{1\bar{i}}|^2 \\
& \quad + \frac{2(1+2c_0)}{\lambda_1^2} \sum_{i \in I} F^{i\bar{i}} \operatorname{Re}(\lambda_1 T_{1i}^1 u_{1\bar{i}}) \\
& = \frac{1+2c_0}{\lambda_1^2} \sum_{i \in I} F^{i\bar{i}} \{(1-2\varepsilon) |u_{1\bar{i}} - T_{1i}^1(\lambda_1 - 1)|^2 \\
& \quad - (1-2(1+2c_0)\delta) |u_{1\bar{i}}|^2 + 2 \operatorname{Re}(\lambda_1 T_{1i}^1 u_{1\bar{i}})\} \\
& = \frac{1+2c_0}{\lambda_1^2} \sum_{i \in I} F^{i\bar{i}} \{(2(1+2c_0)\delta - 2\varepsilon) |u_{1\bar{i}}|^2 \\
& \quad + 2[2\varepsilon(\lambda_1 - 1) + 1] \operatorname{Re}(T_{1i}^1 u_{1\bar{i}}) + (1-2\varepsilon)(\lambda_1 - 1)^2 |T_{1i}^1|^2\} \\
& \geq 0,
\end{aligned}$$

where the last inequality holds if we choose  $\varepsilon = \frac{1}{4}\delta \leq \frac{1}{16}$ . In fact,

$$\begin{aligned}
\Delta = B^2 - 4AC &= 4[2\varepsilon(\lambda_1 - 1) + 1]^2 - 4(1-2\varepsilon)(\lambda_1 - 1)^2(2(1+2c_0)\delta - 2\varepsilon) \\
&\leq 36\varepsilon^2(\lambda_1 - 1)^2 - 4(1-2\varepsilon)(\lambda_1 - 1)^2(2(1+2c_0)\delta - 2\varepsilon) \\
&\leq 4(\lambda_1 - 1)^2(9\varepsilon^2 - 2(1-2\varepsilon)((1+2c_0)\delta) + 2\varepsilon(1-2\varepsilon)) \\
&\leq 4(\lambda_1 - 1)^2(5\varepsilon^2 + 2\varepsilon - \delta) \\
&\leq 4(\lambda_1 - 1)^2(4\varepsilon - \delta) \\
&= 0.
\end{aligned}$$

Then we finally obtain

$$\begin{aligned}
0 &\geq \frac{1}{4}\varphi' \sum_{i=1}^n F^{i\bar{i}} |u_{i\bar{i}}|^2 + \left(-\psi' + \frac{1}{2}\varphi' - C_2 - C_3\right) \sum_{i=1}^n F^{i\bar{i}} - C_1 \\
&= \left(-\psi' + \frac{1}{2}\varphi' - C_2 - C_3\right) \sum_{i=1}^n F^{i\bar{i}} + \frac{1}{4}\varphi' \sum_{i=1}^n F^{i\bar{i}} |u_{i\bar{i}}|^2 - C_1 \\
&\geq \sum_{i=1}^n F^{i\bar{i}} + \frac{1}{16K} \sum_{i=1}^n F^{i\bar{i}} \lambda_i^2 - C_1,
\end{aligned}$$

where we use  $-\psi' \geq C_0 + 1$  and  $C_0 = 31n^2C_2 + 4C_3$ .

In particular,  $\sum_{i=1}^n F^{ii} \leq C$ . By Lemma 2.2 in [Hou et al. 2010], we have  $F^{1\bar{1}} \geq c(n, k)/C_1^{k-1}$ , where  $c(n, k)$  is a positive constant depending only on  $n$  and  $k$ . Then we get the desired estimate

$$\lambda_1 \leq \frac{4C_1^{k/2}}{c(n, k)^{1/2}} \sqrt{K},$$

where  $C_1$  is given in (4-17).  $\square$

### Acknowledgment

We would like to thank Professor Xinan Ma for suggesting the problem and his great encouragement. We thank Professor Shengli Kong and Professor Chuanqiang Chen for careful reading and suggestions. We are also grateful to the referees for their careful reading and many helpful comments.

### References

- [Cherrier 1987] P. Cherrier, “Équations de Monge–Ampère sur les variétés hermitiennes compactes”, *Bull. Sci. Math.* (2) **111**:4 (1987), 343–385. MR Zbl
- [Chou and Wang 2001] K.-S. Chou and X.-J. Wang, “A variational theory of the Hessian equation”, *Comm. Pure Appl. Math.* **54**:9 (2001), 1029–1064. MR Zbl
- [Dinew and Kołodziej 2017] S. Dinew and S. Kołodziej, “Liouville and Calabi–Yau type theorems for complex Hessian equations”, *Amer. J. Math.* **139**:2 (2017), 403–415. MR Zbl
- [Fu and Yau 2008] J.-X. Fu and S.-T. Yau, “The theory of superstring with flux on non-Kähler manifolds and the complex Monge–Ampère equation”, *J. Differential Geom.* **78**:3 (2008), 369–428. MR Zbl
- [Guan and Li 2010] B. Guan and Q. Li, “Complex Monge–Ampère equations and totally real submanifolds”, *Adv. Math.* **225**:3 (2010), 1185–1223. MR Zbl
- [Hou et al. 2010] Z. Hou, X.-N. Ma, and D. Wu, “A second order estimate for complex Hessian equations on a compact Kähler manifold”, *Math. Res. Lett.* **17**:3 (2010), 547–561. MR Zbl
- [Lin and Trudinger 1994] M. Lin and N. S. Trudinger, “On some inequalities for elementary symmetric functions”, *Bull. Austral. Math. Soc.* **50**:2 (1994), 317–326. MR Zbl
- [Phong et al. 2016a] D. H. Phong, S. Picard, and X. Zhang, “The Fu–Yau equation with negative slope parameter”, *Invent. Math.* (online publication December 2016).
- [Phong et al. 2016b] D. H. Phong, S. Picard, and X. Zhang, “On estimates for the Fu–Yau generalization of a Strominger system”, *J. Reine Angew. Math.* (online publication October 2016).
- [Phong et al. 2017] D. H. Phong, S. Picard, and X. Zhang, “Geometric flows and Strominger systems”, *Math. Z.* (online publication March 2017).
- [Sun 2017] W. Sun, “On uniform estimate of complex elliptic equations on closed Hermitian manifolds”, *Commun. Pure Appl. Anal.* **16**:5 (2017), 1553–1570. MR Zbl
- [Székelyhidi 2015] G. Székelyhidi, “Fully non-linear elliptic equations on compact Hermitian manifolds”, preprint, 2015. To appear in *J. Differential Geom.* arXiv
- [Székelyhidi et al. 2015] G. Székelyhidi, V. Tosatti, and B. Weinkove, “Gauduchon metrics with prescribed volume form”, preprint, 2015. arXiv

- [Tosatti and Weinkove 2010] V. Tosatti and B. Weinkove, “The complex Monge–Ampère equation on compact Hermitian manifolds”, *J. Amer. Math. Soc.* **23**:4 (2010), 1187–1195. MR Zbl
- [Tosatti and Weinkove 2013] V. Tosatti and B. Weinkove, “Hermitian metrics,  $(n - 1, n - 1)$  forms and Monge–Ampère equations”, preprint, 2013. arXiv
- [Tosatti and Weinkove 2015] V. Tosatti and B. Weinkove, “On the evolution of a Hermitian metric by its Chern–Ricci form”, *J. Differential Geom.* **99**:1 (2015), 125–163. MR Zbl
- [Tosatti and Weinkove 2017] V. Tosatti and B. Weinkove, “The Monge–Ampère equation for  $(n - 1)$ -plurisubharmonic functions on a compact Kähler manifold”, *J. Amer. Math. Soc.* **30**:2 (2017), 311–346. MR Zbl
- [Wang 2009] X.-J. Wang, “The  $k$ -Hessian equation”, pp. 177–252 in *Geometric analysis and PDEs*, edited by A. Ambrosetti et al., Lecture Notes in Math. **1977**, Springer, Dordrecht, 2009. MR Zbl
- [Yau 1978] S. T. Yau, “On the Ricci curvature of a compact Kähler manifold and the complex Monge–Ampère equation, I”, *Comm. Pure Appl. Math.* **31**:3 (1978), 339–411. MR Zbl
- [Zhang 2010] X. Zhang, “A priori estimates for complex Monge–Ampère equation on Hermitian manifolds”, *Int. Math. Res. Not.* **2010**:19 (2010), 3814–3836. MR Zbl

Received October 30, 2015. Revised January 18, 2017.

DEKAI ZHANG  
SCHOOL OF MATHEMATICAL SCIENCE  
UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA  
HEFEI, ANHUI  
CHINA  
dekzhang@mail.ustc.edu.cn

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
blasius@math.ucla.edu

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
balmer@math.ucla.edu

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
liu@math.ucla.edu

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
qing@cats.ucsc.edu

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
chari@math.ucr.edu

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
jhlu@maths.hku.hk

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
cooper@math.ucsb.edu

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
popa@math.ucla.edu

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
yang@math.princeton.edu

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2017 is US \$450/year for the electronic version, and \$625/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 291    No. 2    December 2017

---

Torsion pairs in silting theory	257
LIDIA ANGELERI HÜGEL, FREDERIK MARKS and JORGE VITÓRIA	
Transfinite diameter on complex algebraic varieties	279
DAVID A. COX and SIONE MA’U	
A universal construction of universal deformation formulas, Drinfeld twists and their positivity	319
CHIARA ESPOSITO, JONAS SCHNITZER and STEFAN WALDMANN	
Uniform stable radius, Lê numbers and topological triviality for line singularities	359
CHRISTOPHE EYRAL	
Rost invariant of the center, revisited	369
SKIP GARIBALDI and ALEXANDER S. MERKURJEV	
Moduli spaces of rank 2 instanton sheaves on the projective space	399
MARCOS JARDIM, MARIO MAICAN and ALEXANDER S. TIKHOMIROV	
A symmetric 2-tensor canonically associated to $Q$ -curvature and its applications	425
YUEH-JU LIN and WEI YUAN	
Gauge invariants from the powers of antipodes	439
CRIS NEGRON and SIU-HUNG NG	
Branching laws for the metaplectic cover of $\mathrm{GL}_2$	461
SHIV PRAKASH PATEL	
Hessian equations on closed Hermitian manifolds	485
DEKAI ZHANG	



0030-8730(201712)291:2;1-W