

*Pacific
Journal of
Mathematics*

**GLOBAL EXISTENCE AND BLOWUP OF
SMOOTH SOLUTIONS OF 3-D POTENTIAL EQUATIONS
WITH TIME-DEPENDENT DAMPING**

FEI HOU, INGO WITT AND HUICHENG YIN

GLOBAL EXISTENCE AND BLOWUP OF SMOOTH SOLUTIONS OF 3-D POTENTIAL EQUATIONS WITH TIME-DEPENDENT DAMPING

FEI HOU, INGO WITT AND HUICHENG YIN

In this paper, we are concerned with the global existence and blowup of smooth solutions of the 3-D irrotational compressible Euler equation with time-dependent damping

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u + pI_3) = -\alpha(t)\rho u, \\ \rho(0, x) = \bar{\rho} + \varepsilon\rho_0(x), \quad u(0, x) = \varepsilon u_0(x), \end{cases}$$

where $x \in \mathbb{R}^3$, the frictional coefficient $\alpha(t) = \mu/(1+t)^\lambda$ with $\mu > 0$ and $\lambda \geq 0$, $\bar{\rho} > 0$ is a constant, $\rho_0, u_0 \in C_0^\infty(\mathbb{R}^3)$, $(\rho_0, u_0) \neq 0$, $\rho(0, x) > 0$, $\operatorname{curl} u_0 \equiv 0$, and $\varepsilon > 0$ is sufficiently small. For $0 \leq \lambda \leq 1$, we show that there exists a global $C^\infty([0, \infty) \times \mathbb{R}^3)$ -smooth solution (ρ, u) by introducing and establishing some uniform time-weighted energy estimates of (ρ, u) , while for $\lambda > 1$, in general, the smooth solution (ρ, u) blows up in finite time. Therefore, $\lambda = 1$ appears to be the critical value for the global existence of small amplitude smooth solution (ρ, u) .

1. Introduction

In this paper, we are concerned with the global existence and blowup of smooth solutions of the three-dimensional irrotational compressible Euler equations with time-dependent damping

$$(1-1) \quad \begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u + pI_3) = -\alpha(t)\rho u, \\ \rho(0, x) = \bar{\rho} + \varepsilon\rho_0(x), \quad u(0, x) = \varepsilon u_0(x), \end{cases}$$

Hou and Yin were supported by the NSFC (No. 11571177) and the Priority Academic Program Development of Jiangsu Higher Education Institutions. Ingo Witt was partly supported by the DFG via the Sino-German project Analysis of Partial Differential Equations and Applications.

MSC2010: primary 35L70; secondary 35L65, 35L67, 76N15.

Keywords: compressible Euler equations, damping, time-weighted energy inequality,

Klainerman–Sobolev inequality, blowup, hypergeometric function.

where $x = (x_1, x_2, x_3)$, ρ , $u = (u_1, u_2, u_3)$, and p stand for the density, velocity, and pressure, respectively, I_3 is the 3×3 identity matrix, the frictional coefficient $\alpha(t) = \mu/(1+t)^\lambda$ with $\mu > 0$ and $\lambda \geq 0$, and $u_0 = (u_{1,0}, u_{2,0}, u_{3,0})$,

$$\text{curl } u_0 = (\partial_2 u_{3,0} - \partial_3 u_{2,0}, \partial_3 u_{1,0} - \partial_1 u_{3,0}, \partial_1 u_{2,0} - \partial_2 u_{1,0}) \equiv 0.$$

The equation of state of the gases is assumed to be $p(\rho) = A\rho^\gamma$, where $A > 0$ and $\gamma > 1$ are constants. Furthermore, $\bar{\rho} > 0$ is a constant, $\rho_0, u_0 \in C_0^\infty(\mathbb{R}^3)$, $(\rho_0, u_0) \not\equiv 0$, $\rho(0, x) > 0$, and $\varepsilon > 0$ is sufficiently small. With respect to the physical background of (1-1), it can be found in [Dafermos 1995].

For $\mu = 0$ in $\alpha(t)$, (1-1) is the standard compressible Euler equation. It is well known that C^∞ -smooth solution (ρ, u) of (1-1) will in general blow up in finite time. For the extensive literature on blowup results and the blowup mechanism for (ρ, u) , see [Alinhac 1999a; 1999b; 1993; Christodoulou 2007; Christodoulou and Miao 2014; Christodoulou and Lisibach 2016; Ding et al. 2016; Hörmander 1997; Sideris 1997; 1985; Speck 2016; Yin and Qiu 1999; Yin 2004] and so on.

For $\lambda = 0$ in $\alpha(t)$, it has been shown that (1-1) admits a global C^∞ -smooth solution (ρ, u) and the large time behavior of (ρ, u) is governed by a parabolic equation derived by using Darcy’s law; see [Dafermos 1995; Hsiao and Serre 1996; Hsiao and Liu 1992; Kawashima and Yong 2004; Nishihara 1997; Pan and Zhao 2009; Sideris et al. 2003; Tan and Guochun 2012; Wang and Yang 2001].

For $\mu > 0$ and $\lambda > 0$ in $\alpha(t)$, an interesting problem arises: does the C^∞ -smooth solution (ρ, u) of (1-1) blow up in finite time or does it exist globally? In this paper, we will systematically study this problem under the assumption of $\text{curl } u_0 \equiv 0$. In this case it is not hard to see that $\text{curl } u(t, \cdot) \equiv 0$ for all $t \geq 0$ as long as the smooth solution (ρ, u) of (1-1) exists. Then one can introduce a potential function $\varphi = \varphi(t, x)$ such that $u = \nabla\varphi$ (here and below, $\nabla = \nabla_x$), where the C^∞ scalar function φ has a compact support in x (as $u(t, \cdot)$ has a compact support for any fixed $t \geq 0$ in view of $u_0 \in C_0^\infty(\mathbb{R}^3)$ and admits a finite propagation speed which holds for hyperbolic systems). Substituting $u = \nabla\varphi$ into the second equation of (1-1), we obtain

$$(1-2) \quad \partial_t \varphi + \frac{1}{2} |\nabla\varphi|^2 + h(\rho) + \frac{\mu}{(1+t)^\lambda} \varphi = 0,$$

where $h'(\rho) = c^2(\rho)/\rho$ with $c(\rho) = \sqrt{p'(\rho)}$ and $h(\bar{\rho}) = 0$.

From $h'(\rho) > 0$ for $\rho > 0$ we have that

$$(1-3) \quad \rho = h^{-1} \left(- \left(\partial_t \varphi + \frac{1}{2} |\nabla\varphi|^2 + \frac{\mu}{(1+t)^\lambda} \varphi \right) \right),$$

where $\bar{\rho} = h^{-1}(0)$ and h^{-1} is the inverse function of $h = h(\rho)$.

Substituting (1-3) into the first equation of (1-1) yields

$$(1-4) \quad \partial_t^2 \varphi - c^2(\rho) \Delta \varphi + 2 \sum_{k=1}^3 (\partial_k \varphi) \partial_{t_k}^2 \varphi + \sum_{i,k=1}^3 (\partial_i \varphi) (\partial_k \varphi) \partial_{i_k}^2 \varphi + \frac{\mu}{(1+t)^\lambda} |\nabla \varphi|^2 + \partial_t \left(\frac{\mu}{(1+t)^\lambda} \varphi \right) = 0.$$

As for the initial data $\varphi(0, x)$ and $\partial_t \varphi(0, x)$ for (1-4): Obviously, $\varphi(0, x) = \varepsilon \varphi_0(x)$, where

$$\varphi_0(x) = \int_{-\infty}^{x_1} u_{1,0}(s, x_2, x_3) ds.$$

Note that $\varphi_0 \in C_0^\infty(\mathbb{R}^3)$ in view of $\text{curl } u_0 \equiv 0$ and $u_0 \in C_0^\infty(\mathbb{R}^3)$. Furthermore, from (1-2) we infer that $\partial_t \varphi(0, x) = \varepsilon \varphi_1(x) + \varepsilon^2 g(x, \varepsilon)$, where

$$\varphi_1 = - \left(\mu \varphi_0 + \frac{c^2(\bar{\rho})}{\bar{\rho}} \rho_0 \right)$$

and

$$g(x, \varepsilon) = -\rho_0^2(x) \int_0^1 \left(\frac{c^2(\rho)}{\rho} \right)' \Big|_{\rho=\bar{\rho}+\theta \varepsilon \rho_0(x)} d\theta - \frac{1}{2} \sum_{i=1}^3 u_{i,0}^2(x).$$

Notice that $g(x, \varepsilon)$ is smooth in (x, ε) and has compact support in x . Consequently, studying problem (1-1) under the assumption $\text{curl } u_0 \equiv 0$ is equivalent to investigating the problem

$$(1-5) \quad \left\{ \begin{array}{l} \partial_t^2 \varphi - c^2(\rho) \Delta \varphi + 2 \sum_{k=1}^3 (\partial_k \varphi) \partial_{t_k}^2 \varphi + \sum_{i,k=1}^3 (\partial_i \varphi) (\partial_k \varphi) \partial_{i_k}^2 \varphi + \frac{\mu}{(1+t)^\lambda} |\nabla \varphi|^2 + \partial_t \left(\frac{\mu}{(1+t)^\lambda} \varphi \right) = 0, \\ \varphi(0, x) = \varepsilon \varphi_0(x), \quad \partial_t \varphi(0, x) = \varepsilon \varphi_1(x) + \varepsilon^2 g(x, \varepsilon). \end{array} \right.$$

Here we mention that

$$c^2(\rho) = c^2(\bar{\rho}) - (\gamma - 1) \left(\partial_t \varphi + \frac{1}{2} |\nabla \varphi|^2 + \frac{\mu}{(1+t)^\lambda} \varphi \right)$$

which follows by direct computation.

We now state the first main result of this paper.

Theorem 1.1 (global existence for $0 \leq \lambda \leq 1$). *Suppose that $\text{curl } u_0 \equiv 0$. If $\mu > 0$ and $0 \leq \lambda \leq 1$, then, for $\varepsilon > 0$ small enough, (1-5) admits a global C^∞ -smooth solution φ . As a consequence, (1-1) has a global C^∞ -smooth solution (ρ, u) which fulfills $\rho > 0$ and which is uniformly bounded for $t \geq 0$ together with all its derivatives.*

Remark. The principal part of the linearization of the equation in (1-5) about $(\rho, \varphi) = (\bar{\rho}, 0)$ is

$$(1-6) \quad \mathcal{L}(\dot{\varphi}) \equiv \partial_t^2 \dot{\varphi} - c^2(\bar{\rho}) \Delta \dot{\varphi} + \frac{\mu}{(1+t)^\lambda} \partial_t \dot{\varphi} - \frac{\mu \lambda}{(1+t)^{\lambda+1}} \dot{\varphi}.$$

For the linear operator \mathcal{L}_0 with

$$\mathcal{L}_0(\dot{\varphi}) \equiv \partial_t^2 \dot{\varphi} - c^2(\bar{\rho}) \Delta \dot{\varphi} + \frac{\mu}{(1+t)^\lambda} \partial_t \dot{\varphi},$$

which appears as part of (1-6), it is shown in [Wirth 2006; 2007] that the large-term behavior of solutions $\dot{\varphi}$ of $\mathcal{L}_0(\dot{\varphi}) = 0$ depends on the value of λ . For $0 \leq \lambda < 1$ it is the same as the large-term behavior of solutions of the linear heat equation $\partial_t \dot{\varphi} - c^2(\bar{\rho}) \Delta \dot{\varphi} = 0$, while for $\lambda > 1$ it is the same as the large-term behavior of solutions of the linear wave equation $\partial_t^2 \dot{\varphi} - c^2(\bar{\rho}) \Delta \dot{\varphi} = 0$. In addition, precise microlocal large-term decay properties of solutions $\dot{\varphi}$ of $\mathcal{L}(\dot{\varphi}) = 0$ have been established in [do Nascimento and Wirth 2015] for a special range of values of λ and μ . It seems to be difficult, however, to apply these microlocal estimates to attack the quasilinear problem (1-5). (In general, those microlocal estimates are useful when treating semilinear damped wave equations; see [D'Abbico and Reissig 2014; D'Abbico et al. 2015].)

Remark. For the 1-D Burgers equation with time-dependent damping term

$$(1-7) \quad \begin{cases} \partial_t w + w \partial_x w = -\frac{\mu}{(1+t)^\lambda} w, & (t, x) \in \mathbb{R}_+ \times \mathbb{R}, \\ w(0, x) = \varepsilon w_0(x), \end{cases}$$

where $\mu > 0$ and $\lambda \geq 0$ are constants, $w_0 \in C_0^\infty(\mathbb{R})$, $w_0 \not\equiv 0$, and $\varepsilon > 0$ is sufficiently small, one concludes by the method of characteristics that

$$\begin{cases} T_\varepsilon = \infty & \text{if } 0 \leq \lambda < 1 \text{ or } \lambda = 1, \mu > 1, \\ T_\varepsilon < \infty & \text{if } \lambda > 1 \text{ or } \lambda = 1, 0 \leq \mu \leq 1, \end{cases}$$

where T_ε is the lifespan of the C^∞ -smooth solution w of (1-7). Therefore, $\lambda = 1$ again appears to be the critical value for the global existence of smooth solutions w of (1-7) in the presence of the damping term

$$\frac{\mu}{(1+t)^\lambda} w.$$

Remark. The smallness of $\varepsilon > 0$ in Theorem 1.1 is necessary in order to guarantee the global existence of smooth solution (ρ, u) . Indeed, as in [Sideris et al. 2003], large amplitude smooth solution of (1-1) may blow up in finite time even for $0 \leq \lambda \leq 1$. See also Theorem 4.1.

Next we concentrate on the case of $\lambda > 1$. As in [Sideris 1985], introduce the two functions

$$q_0(l) = \int_{|x|>l} \frac{(|x| - l)^2}{|x|} (\rho(0, x) - \bar{\rho}) dx,$$

$$q_1(l) = \int_{|x|>l} \frac{|x|^2 - l^2}{|x|^3} x \cdot (\rho u)(0, x) dx.$$

Before stating our blowup result for problem (1-1) with $\lambda > 1$, we require to introduce a special hypergeometric function $\Psi(a, b, c; z)$, where the constants a and b satisfy $a + b = 1$ and $ab = \frac{1}{2}\mu\lambda$, $c \in \mathbb{R}^+$, the variable $z \in \mathbb{R}$, and

$$\Psi(a, b, c; z) = \sum_{n=0}^{+\infty} \frac{(a)_n (b)_n}{n! (c)_n} z^n$$

with $(a)_n = a(a + 1) \cdots (a + n - 1)$ and $(a)_0 = 1$. It is known from [Erdélyi et al. 1953] that $\Psi(a, b, c; z)$ is an analytic function of z in $(-1, 1)$ and $\Psi(a, b, c; 0) = \Psi(a + 1, b + 1, c; 0) = 1$. Therefore, there exists a small constant $\delta_0 > 0$ depending on a and b (i.e., μ and λ) such that for $-\frac{1}{2}\delta_0 \leq z \leq 0$,

$$(1-8) \quad \frac{1}{2} \leq \Psi(a, b, 1; z), \Psi(a + 1, b + 1, 2; z) \leq \frac{3}{2}.$$

Theorem 1.2 (blowup for $\lambda > 1$). *Suppose $\text{supp } \rho_0, \text{supp } u_0 \subseteq \{x : |x| \leq M\}$ and let*

$$(1-9) \quad q_0(l) > 0,$$

$$(1-10) \quad q_1(l) \geq 0$$

hold for all $l \in (\tilde{M}, M)$, where \tilde{M} is some fixed constant satisfying $0 \leq \tilde{M} < M$. Moreover, we assume that there exist two constants $M_0 \geq \tilde{M}$ and $\Lambda \geq \frac{3}{2}\mu\lambda$ such that

$$(1-11) \quad q_1(l) \geq \Lambda q_0(l),$$

holds for all $l \in (M_0, M)$, where $M - M_0 < \delta_0$ and δ_0 is given in (1-8). If $\mu > 0$ and $\lambda > 1$, then there exists an $\varepsilon_0 > 0$ such that, for $0 < \varepsilon \leq \varepsilon_0$, the lifespan T_ε of C^∞ -smooth solution (ρ, u) of (1-1) is finite.

Remark. It is not hard to find a large number of initial data $(\rho, u)(0, x)$ such that (1-9)–(1-11) are satisfied. For instance, choosing $\rho_0(x) > 0$ and $u_0(x) = x\rho_0(x)\Lambda/\bar{\rho}$, then we get (1-9)–(1-11).

Remark. Sideris [1985] showed the formation of singularities in three-dimensional compressible equations under the assumptions of (1-9)–(1-10). However, in order to prove the blowup result of smooth solution (ρ, u) to problem (1.1) and overcome the difficulty arisen by the time-dependent frictional coefficient $\mu/(1 + t)^\lambda$ with $\mu > 0$

and $\lambda > 1$, we pose an extra assumption (1-11) except (1-9)–(1-10), which leads to the nonnegativity of $P(t, l)$ in (3-7) so that an ordinary typed blowup inequalities (3-23)–(3-24) can be established. One can see more details in Section 3.

Let us indicate the proofs of Theorems 1.1 and 1.2. To prove Theorem 1.1, we first introduce the function $\psi = \varphi/(1+t)^\lambda$ which fulfills the second-order quasilinear wave equation

$$\partial_t^2 \psi - \Delta \psi + \frac{\mu}{(1+t)^\lambda} \partial_t \psi + \frac{2\lambda}{1+t} \partial_t \psi - \frac{\lambda(1-\lambda)}{(1+t)^2} \psi = Q(\psi, \partial \psi, \partial^2 \psi),$$

where $Q(\psi, \partial \psi, \partial^2 \psi)$ stands for an error term which is of the second order in $(\psi, \partial \psi, \partial^2 \psi)$; $\partial = (\partial_t, \nabla)$. Then, in order to establish the global existence of ψ , we introduce the time-weighted energy

$$E_N(\psi)(t) = \sum_{0 \leq |a| \leq N} \int_{\mathbb{R}^3} ((1+t)^{2\lambda} |\partial \Gamma^a \psi|^2 + |\Gamma^a \psi|^2) dx,$$

where $N \geq 8$ is a fixed number, $\Gamma = (\Gamma_0, \Gamma_1, \dots, \Gamma_7) = (\partial, \Omega, S)$ with $\Omega = (\Omega_1, \Omega_2, \Omega_3) = x \wedge \nabla$, $S = t\partial_t + \sum_{k=1}^3 x_k \partial_k$, and $\Gamma^a = \Gamma_0^{a_0} \Gamma_1^{a_1} \dots \Gamma_7^{a_7}$. Note that the vector fields Γ which appear in the definition of the energy $E_N(\psi)(t)$ only comprise part of the standard Klainerman vector fields $\{\partial, \Omega, S, H\}$, where $H = (H_1, H_2, H_3) = (x_1 \partial_t + t \partial_1, x_2 \partial_t + t \partial_2, x_3 \partial_t + t \partial_3)$. This is due to the fact that the equation in (1-5) is not invariant under the Lorentz transformations H in view of the presence of the time-dependent damping. By a rather technical and involved analysis of the resulting equation for ψ , we eventually show that $E_N(\psi)(t) \leq \frac{1}{2} K^2 \varepsilon^2$ when $E_N(\psi)(t) \leq K^2 \varepsilon^2$ is assumed for some suitably large constant $K > 0$ and small $\varepsilon > 0$. Here we emphasize that the condition of $0 \leq \lambda \leq 1$ plays an essential role in the process of deriving the uniform boundedness of $E_N(\psi)(t)$ (see Lemmas 2.3–2.5). This, together with the continuous induction argument, yields the global existence of ψ and further completes the proof of Theorem 1.1 for $0 \leq \lambda \leq 1$. To prove the blowup result of Theorem 1.2 for $\lambda > 1$, as in [Sideris 1985], we derive a related second-order ordinary differential inequality. From this and assumptions (1-9)–(1-11), an upper bound of the lifespan T_ε is derived by making essential use of $\lambda > 1$. In this way the proof of Theorem 1.2 is completed. In Theorem 4.1, we show that for large data smooth solution (ρ, u) of (1-1), even in case $0 \leq \lambda \leq 1$, (ρ, u) will in general blow up in finite time. In addition, the proof on the nonnegativity of $P(t, l)$, which is introduced in (3-1), is given in the Appendix.

Throughout, we shall use the following notation and conventions:

- ∇ stands for ∇_x ;
- $r = |x| = \sqrt{x_1^2 + x_2^2 + x_3^2}$;
- $\langle r - t \rangle = (1 + (r - t)^2)^{1/2}$;

- $\|u(t, x)\| = \left(\int_{\mathbb{R}^3} |u(t, x)|^2 dx\right)^{1/2}$ and $\|u(t, x)\|_{L^\infty} = \sup_{x \in \mathbb{R}^3} |u(t, x)|$;
- Γ denotes one of the vector fields $\{\partial, S, \Omega\}$ on $\mathbb{R}_+ \times \mathbb{R}^3$, where $\partial = (\partial_t, \nabla)$, $S = t\partial_t + \sum_{k=1}^3 x_k \partial_k$, $\Omega = (\Omega_1, \Omega_2, \Omega_3) = x \wedge \nabla$;
- β is the solution of $\beta'(t) = \frac{\mu}{(1+t)^\lambda} \beta(t)$ for $t \geq 0$, $\beta(0) = 1$, i.e.,

$$(1-12) \quad \beta(t) \equiv \begin{cases} e^{\frac{\mu}{1-\lambda}[(1+t)^{1-\lambda}-1]}, & \lambda \geq 0, \lambda \neq 1, \\ (1+t)^\mu, & \lambda = 1; \end{cases}$$

- $c(\bar{\rho}) = 1$ will be assumed throughout (introduce $X = x/c(\bar{\rho})$ as a new space coordinate if necessary).

2. Global existence for small amplitude in case $0 \leq \lambda \leq 1$

Throughout this section, $C > 0$ stands for a generic constant which is independent of K, ε , and t .

We start by recalling a Sobolev-type inequality (see [Klainerman 1987]).

Lemma 2.1. *Let $u = u(t, x)$ be a smooth function of $(t, x) \in [0, \infty) \times \mathbb{R}^3$. Then*

$$(2-1) \quad |u(t, x)| \leq C(1+r)^{-1} \sum_{|a| \leq 2} \|\Gamma^a u(t, x)\|.$$

Moreover, we shall make use of the following inequalities (see [Klainerman and Sideris 1996, Lemma 3.1 and Theorem 5.1]).

Lemma 2.2. *For $u \in C^2([0, \infty) \times \mathbb{R}^3)$,*

$$(2-2) \quad \|(r-t)\nabla\partial u(t, x)\| \leq C \left(\sum_{|b| \leq 1} \|\partial\Gamma^b u(t, x)\| + t\|\square u(t, x)\| \right),$$

$$(2-3) \quad (1+r)\langle r-t \rangle |\nabla\partial u(t, x)| \leq C \left(\sum_{|b| \leq 3} \|\partial\Gamma^b u(t, x)\| + t\|\square u(t, x)\| \right),$$

where $\square = \partial_t^2 - \Delta = \partial_t^2 - \sum_{k=1}^3 \partial_k^2$.

We now reformulate problem (1-5). Let $\psi = \varphi/(1+t)^\lambda$. From (1-5) and $c(\bar{\rho}) = 1$ we then have

$$(2-4) \quad \square\psi + \frac{\mu}{(1+t)^\lambda} \partial_t \psi + \frac{2\lambda}{1+t} \partial_t \psi - \frac{\lambda(1-\lambda)}{(1+t)^2} \psi = Q(\psi, \partial\psi, \partial^2\psi),$$

where

$$Q(\psi, \partial\psi, \partial^2\psi) = (c^2(\rho) - 1)\Delta\psi - 2(1+t)^\lambda \partial_t \nabla\psi \cdot \nabla\psi - 2\lambda(1+t)^{\lambda-1} |\nabla\psi|^2 - \mu |\nabla\psi|^2 - (1+t)^{2\lambda} \sum_{1 \leq i, j \leq 3} (\partial_i \psi)(\partial_j \psi) \partial_{ij}^2 \psi.$$

We define a time-weighted energy for (2-4),

$$E_N(\psi(t)) = \sum_{0 \leq |a| \leq N} \int_{\mathbb{R}^3} ((1+t)^{2\lambda} |\partial \Gamma^a \psi|^2 + |\Gamma^a \psi|^2) dx,$$

where $N \geq 8$ is a fixed number. Moreover, we assume that for any $t \geq 0$,

$$(2-5) \quad E_N(\psi(t)) \leq K^2 \varepsilon^2,$$

where $K > 0$ is a suitably large constant. It follows from (2-1) and (2-5) that, for all $|a| \leq N - 2$,

$$(2-6) \quad \begin{aligned} |\partial \Gamma^a \psi| &\leq C(1+r)^{-1} \sum_{|b| \leq 2} \|\Gamma^b \partial \Gamma^a \psi(t, x)\| \\ &\leq C(1+r)^{-1} \sum_{|b| \leq N} \|\partial \Gamma^b \psi(t, x)\| \\ &\leq C(1+r)^{-1} (1+t)^{-\lambda} \sqrt{E_N(\psi(t))} \\ &\leq CK\varepsilon(1+r)^{-1} (1+t)^{-\lambda} \end{aligned}$$

and

$$(2-7) \quad |\Gamma^a \psi| \leq C(1+r)^{-1} \sum_{|b| \leq N} \|\Gamma^b \psi(t, x)\| \leq CK\varepsilon(1+r)^{-1}.$$

In view of Lemma 2.2 and (2-5), we have

Lemma 2.3. *Let ψ be a solution of (2-4). Then, for all $|a| \leq N - 3$ and $0 \leq \lambda \leq 1$, we have the pointwise estimate*

$$(2-8) \quad \|\nabla \partial \Gamma^a \psi\|_{L^\infty} \leq CK\varepsilon(1+t)^{-2\lambda}.$$

Moreover, for $0 \leq l \leq N - 1$, the weighted L^2 estimate

$$(2-9) \quad \begin{aligned} \sum_{|b| \leq l} \|(r-t) \nabla \partial \Gamma^b \psi(t, x)\| \\ \leq C \sum_{|c| \leq l+1} \|\partial \Gamma^c \psi(t, x)\| + C(1+t)^{1-\lambda} \sum_{|c| \leq l} \|\nabla \Gamma^c \psi(t, x)\| \\ + C(1-\lambda)(1+t)^{-1} \sum_{|c| \leq l} \|\Gamma^c \psi(t, x)\| \end{aligned}$$

holds.

Proof. It follows from (2-3)–(2-4) and (2-6)–(2-7) that

$$\begin{aligned}
 & (1+t) \sum_{|a| \leq N-3} |\nabla \partial \Gamma^a \psi| \\
 & \leq C \sum_{|a| \leq N-3} (1+r) \langle r-t \rangle |\nabla \partial \Gamma^a \psi| \\
 & \leq C \sum_{|c| \leq N} \|\partial \Gamma^c \psi\| + Ct \sum_{|a| \leq N-3} \|\square \Gamma^a \psi\| \\
 & \leq CK\varepsilon(1+t)^{-\lambda} + C(1+t)^{1-\lambda} \sum_{|a| \leq N-3} \|\partial_t \Gamma^a \psi\| + C(1+t)^{-1} \sum_{|a| \leq N-3} \|\Gamma^a \psi\| \\
 & \quad + C(1+t) \sum_{|b|+|c| \leq N-3} \|\nabla \partial \Gamma^b \psi \Gamma^c \psi\| + C(1+t)^{1+\lambda} \sum_{|a| \leq N-3} \|\Gamma^a (\partial_t \nabla \psi \cdot \nabla \psi)\| \\
 & \leq CK\varepsilon(1+t)^{1-2\lambda} + CK\varepsilon(1+t) \sum_{|a| \leq N-3} \|\nabla \partial \Gamma^a \psi\|_{L^\infty},
 \end{aligned}$$

which derives (2-7) in view of the smallness of $\varepsilon > 0$.

By (2-2), (2-6)–(2-8) and (2-4), we have that, for $l \leq N-1$,

$$\begin{aligned}
 & (2-10) \\
 & \sum_{|b| \leq l} \|\langle r-t \rangle \nabla \partial \Gamma^b \psi\| \\
 & \leq C \sum_{|c| \leq l+1} \|\partial \Gamma^c \psi\| + Ct \sum_{|b| \leq l} \|\Gamma^b \square \psi\| \\
 & \leq C \sum_{|c| \leq l+1} \|\partial \Gamma^c \psi\| + C(1+t)^{1-\lambda} \sum_{|c| \leq l} \|\nabla \Gamma^c \psi\| + C(1-\lambda)(1+t)^{-1} \sum_{|c| \leq l} \|\Gamma^c \psi\| \\
 & \quad + C(1+t)^{1+\lambda} \sum_{|b| \leq l} \|\Gamma^b (\partial_t \nabla \psi \cdot \nabla \psi)\| \\
 & \quad + C(1+t) \sum_{\substack{|c| \leq N-3, \\ |b| \leq l-|c|}} \|\langle r-t \rangle^{-1} \Gamma^c \psi\|_{L^\infty} \|\langle r-t \rangle \nabla \partial \Gamma^b \psi\| \\
 & \quad + C(1+t) \sum_{\substack{2-N \leq |c| \leq l, \\ |b| \leq l+2-N}} \|(1+r) \nabla \partial \Gamma^b \psi\|_{L^\infty} \|(1+r)^{-1} \Gamma^c \psi\| \\
 & \leq C \sum_{|c| \leq l+1} \|\partial \Gamma^c \psi\| + C(1+t)^{1-\lambda} \sum_{|c| \leq l} \|\nabla \Gamma^c \psi\| + C(1-\lambda)(1+t)^{-1} \sum_{|c| \leq l} \|\Gamma^c \psi\| \\
 & \quad + CK\varepsilon \sum_{|b| \leq l} \|\langle r-t \rangle \nabla \partial \Gamma^b \psi\| + CK\varepsilon(1+t)^{1-\lambda} \sum_{2-N \leq |c| \leq l} \|(1+r)^{-1} \Gamma^c \psi\|.
 \end{aligned}$$

Note that $\Gamma^c \psi(t, x)$ is supported in $\{x : |x| \leq t + M\}$. Then it follows from Hardy inequality that

$$(2-11) \quad \|(1+r)^{-1} \Gamma^c \psi\| \leq C \|\nabla \Gamma^c \psi\|.$$

Substituting (2-11) into (2-10) and applying the smallness of ε , we derive (2-9). \square

Next we derive the time-weighted energy estimate for the solution ψ of (2-4).

Lemma 2.4. *Let $\mu > 0$ and $\lambda \in (0, 1]$. Under assumption (2-5), for all $t > 0$ and $N \geq 8$, it holds that*

$$(2-12) \quad \sum_{0 \leq |a| \leq N} \int_{\mathbb{R}^3} ((1+t)^{2\lambda} |\partial \partial^a \psi|^2 + \psi^2) dx + C \sum_{0 \leq |a| \leq N} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \partial^a \psi|^2 dx d\tau \\ \leq C\varepsilon^2 + C(1+K\varepsilon) \int_0^t A(\tau) \sum_{0 \leq |a| \leq N} \int_{\mathbb{R}^3} ((1+\tau)^{2\lambda} |\partial \partial^a \psi|^2 + \psi^2) dx d\tau,$$

where $A(\cdot)$ stands for a generic nonnegative function such that $A \in L^1((0, \infty))$, and $\|A\|_{L^1}$ is independent of K but dependent on μ and λ .

Proof. First we show (2-12) in case $|a| = 0$. Multiplying (2-4) by $m(1+t)^{2\lambda} \partial_t \psi + (1+t)^{2\lambda-1} \psi$ yields by a direct computation

$$(2-13) \quad \frac{1}{2} \partial_t [m(1+t)^{2\lambda} |\partial \psi|^2 + 2(1+t)^{2\lambda-1} \psi \partial_t \psi + (\mu(1+t)^{\lambda-1} + 2\lambda(1+t)^{2\lambda-2}) \psi^2] \\ + \operatorname{div}(\dots) + (\mu m(1+t)^\lambda + (\lambda m - 1)(1+t)^{2\lambda-1}) (\partial_t \psi)^2 \\ + (1 - \lambda m)(1+t)^{2\lambda-1} |\nabla \psi|^2 + \frac{\mu}{2} (1-\lambda)(1+t)^{\lambda-2} \psi^2 \\ + C_1(\lambda - 1)(1+t)^{2\lambda-2} \psi \partial_t \psi + C_2(\lambda - 1)(1+t)^{2\lambda-3} \psi^2 \\ = (m(1+t)^{2\lambda} \partial_t \psi + (1+t)^{2\lambda-1} \psi) Q(\psi, \partial \psi, \partial^2 \psi),$$

where the constant $m > 0$ will be determined later and C_i ($i = 1, 2$) are suitable constants. Note that in the square bracket of the first line in (2-13),

$$(2-14) \quad m(1+t)^{2\lambda} |\partial \psi|^2 + 2(1+t)^{2\lambda-1} \psi \partial_t \psi + (\mu(1+t)^{\lambda-1} + 2\lambda(1+t)^{2\lambda-2}) \psi^2 \\ = m(1+t)^{2\lambda} \left(\frac{1}{3} |\partial_t \psi|^2 + |\nabla \psi|^2 \right) + \left(\mu(1+t)^{\lambda-1} + \left(2\lambda - \frac{3}{2m} \right) (1+t)^{2\lambda-2} \right) \psi^2 \\ + \left(\sqrt{\frac{2m}{3}} (1+t)^\lambda \partial_t \psi + \sqrt{\frac{3}{2m}} (1+t)^{\lambda-1} \psi \right)^2.$$

We choose $m > 0$ to fulfill

$$\lambda < \frac{1}{m} < \min\{\mu + \lambda, 2\lambda\};$$

together with $\lambda \leq 1$ (i.e., $2\lambda - 2 \leq \lambda - 1 \leq 0$), this yields that (2-14) is equivalent to

$$(1+t)^{2\lambda} |\partial \psi|^2 + (1+t)^{\lambda-1} \psi^2.$$

On the other hand, the coefficients

$$\mu m(1+t)^\lambda + (\lambda m - 1)(1+t)^{2\lambda-1}$$

and

$$(1 - \lambda m)(1+t)^{2\lambda-1}$$

of $(\partial_t \psi)^2$ and $|\nabla \psi|^2$ in the left-hand side of (2-13) are both positive.

Then integrating (2-13) over $[0, t] \times \mathbb{R}^3$ yields

$$\begin{aligned} (2-15) \quad & \int_{\mathbb{R}^3} ((1+t)^{2\lambda} |\partial \psi|^2 + (1+t)^{\lambda-1} \psi^2) dx \\ & + C \int_0^t \int_{\mathbb{R}^3} ((1+\tau)^\lambda (\partial_t \psi)^2 + (1+\tau)^{2\lambda-1} |\nabla \psi|^2 + (1+\tau)^{\lambda-2} \psi^2) dx d\tau \\ & \leq C\varepsilon^2 + \int_0^t A(\tau) \int_{\mathbb{R}^3} (1+\tau)^{\lambda-1} \psi^2 dx d\tau \\ & + C \left| \int_0^t \int_{\mathbb{R}^3} (m(1+\tau)^{2\lambda} \partial_t \psi + (1+\tau)^{2\lambda-1} \psi) Q(\psi, \partial \psi, \partial^2 \psi) dx d\tau \right|. \end{aligned}$$

Next we improve the time-weighted estimate of ψ in the left-hand side of (2-15).

Multiplying both sides of (2-4) by $(1+t)^\lambda \psi$ yields by direct computation

$$\begin{aligned} & \partial_t \left((1+t)^\lambda \psi \partial_t \psi + \frac{\mu}{2} \psi^2 \right) + \operatorname{div}(\dots) - (1+t)^\lambda (\partial_t \psi)^2 - \lambda(1+t)^{\lambda-1} \psi \partial_t \psi \\ & + (1+t)^\lambda |\nabla \psi|^2 + 2\lambda(1+t)^{\lambda-1} \psi \partial_t \psi + \lambda(\lambda-1)(1+t)^{\lambda-2} \psi^2 \\ & = (1+t)^\lambda \psi Q(\psi, \partial \psi, \partial^2 \psi). \end{aligned}$$

From this and (2-15), we can choose the multiplier

$$m(1+t)^{2\lambda} \partial_t \psi + (1+t)^{2\lambda-1} \psi + \kappa(1+t)^\lambda \psi$$

for (2-4) with a small $\kappa > 0$ and then obtain

$$\begin{aligned} (2-16) \quad & \int_{\mathbb{R}^3} ((1+t)^{2\lambda} |\partial \psi|^2 + \psi^2) dx + C \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \psi|^2 dx d\tau \\ & \leq C\varepsilon^2 + \int_0^t A(\tau) \int_{\mathbb{R}^3} \psi^2 dx d\tau \\ & + C \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \psi) Q(\psi, \partial \psi, \partial^2 \psi) dx d\tau \right| \\ & + C \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda \psi Q(\psi, \partial \psi, \partial^2 \psi) dx d\tau \right|. \end{aligned}$$

Next we derive the time-weighted estimates of $\partial^a \psi$ with $1 \leq |a| \leq N$. Taking ∂^a on both sides of (2-4) yields

$$\begin{aligned} \square \partial^a \psi + \frac{\mu}{(1+t)^\lambda} \partial_t \partial^a \psi + \frac{2\lambda}{1+t} \partial_t \partial^a \psi \\ = \partial^a Q(\psi, \partial \psi, \partial^2 \psi) + \sum_{1 \leq |b| \leq |a|} \frac{1}{(1+t)^\lambda} (1 + \mathcal{O}((1+t)^{\lambda-1})) \partial^b \psi \\ - \lambda(\lambda-1) \partial^a \left(\frac{1}{(1+t)^2} \right) \psi. \end{aligned}$$

Exactly as for (2-16), multiplying this by

$$m(1+t)^{2\lambda} \partial_t \partial^a \psi + (1+t)^{2\lambda-1} \partial^a \psi + \kappa(1+t)^\lambda \partial^a \psi,$$

we obtain

$$\begin{aligned} (2-17) \quad \sum_{0 \leq |a| \leq N} \int_{\mathbb{R}^3} ((1+t)^{2\lambda} |\partial \partial^a \psi|^2 + \psi^2) dx + C \sum_{0 \leq |a| \leq N} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \partial^a \psi|^2 dx d\tau \\ \leq C\varepsilon^2 + \int_0^t A(\tau) \int_{\mathbb{R}^3} \psi^2 dx d\tau \\ + C \sum_{0 \leq |a| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \partial^a \psi) \partial^a Q(\psi, \partial \psi, \partial^2 \psi) dx d\tau \right| \\ + C \sum_{0 \leq |a| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda (\partial^a \psi) \partial^a Q(\psi, \partial \psi, \partial^2 \psi) dx d\tau \right|. \end{aligned}$$

We now deal with the last two terms in the right-hand side of (2-17). We first analyze the integrand $(1+t)^{2\lambda} (\partial_t \partial^a \psi) \partial^a Q(\psi, \partial \psi, \partial^2 \psi)$ of the penultimate term. Direct computation yields

$$\begin{aligned} \partial^a Q(\psi, \partial \psi, \partial^2 \psi) \\ = (c^2(\rho) - 1) \Delta \partial^a \psi - 2(1+t)^\lambda \nabla \partial_t \partial^a \psi \cdot \nabla \psi - (1+t)^{2\lambda} (\partial_i \psi) (\partial_j \psi) \partial_{ij}^2 \partial^a \psi + \text{l.o.t.} \end{aligned}$$

and

$$\begin{aligned} (2-18) \quad (1+t)^{2\lambda} (\partial_t \partial^a \psi) \partial^a Q(\psi, \partial \psi, \partial^2 \psi) \\ = \operatorname{div}((1+t)^{2\lambda} (c^2(\rho) - 1) (\partial_t \partial^a \psi) \nabla \partial^a \psi) - \operatorname{div}((1+t)^{3\lambda} |\partial_t \partial^a \psi|^2 \nabla \psi) \\ - \frac{1}{2} \partial_t ((1+t)^{2\lambda} (c^2(\rho) - 1) |\nabla \partial^a \psi|^2) \\ + (1+t)^{3\lambda} |\partial_t \partial^a \psi|^2 \Delta \psi + \lambda(1+t)^{2\lambda-1} (c^2(\rho) - 1) |\nabla \partial^a \psi|^2 \\ + \frac{1}{2} (1+t)^{2\lambda} (c^2(\rho))' \partial_t \rho |\nabla \partial^a \psi|^2 \\ - (1+t)^{4\lambda} (\partial_i \psi) (\partial_j \psi) (\partial_{ij}^2 \partial^a \psi) \partial_t \partial^a \psi + \text{l.o.t.}, \end{aligned}$$

where here and below l.o.t. designates lower-order terms which are of the form

$$(\partial^{b_1} \psi)(\partial^{b_2} \psi) \dots (\partial^{b_l} \psi)$$

(multiplied by $\partial \partial^a \psi$ or $\partial^a \psi$) with $l \geq 2$ and $1 \leq |b_1| + \dots + |b_l| \leq |a| + 1$. Here we are concerned with the top-order derivatives only. Note that the term $(1+t)^{4\lambda}(\partial_i \psi)(\partial_j \psi)(\partial_{ij}^2 \partial^a \psi) \partial_t \partial^a \psi$ in (2-18) can be expressed as

$$(2-19) \quad (1+t)^{4\lambda}(\partial_i \psi)(\partial_j \psi)(\partial_{ij}^2 \partial^a \psi) \partial_t \partial^a \psi \\ = \frac{1}{2} \{ \partial_i ((1+t)^{4\lambda}(\partial_i \psi)(\partial_j \psi)(\partial_j \partial^a \psi) \partial_t \partial^a \psi) \\ + \partial_j ((1+t)^{4\lambda}(\partial_i \psi)(\partial_j \psi)(\partial_i \partial^a \psi) \partial_t \partial^a \psi) \\ - \partial_t ((1+t)^{4\lambda}(\partial_i \psi)(\partial_j \psi)(\partial_i \partial^a \psi) \partial_j \partial^a \psi) \\ + \partial_t ((1+t)^{4\lambda}(\partial_i \psi) \partial_j \psi) (\partial_i \partial^a \psi) \partial_j \partial^a \psi + \text{l.o.t.} \}.$$

Similarly, for the integrand of

$$\left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda (\partial^a \psi) \partial^a Q(\psi, \partial \psi, \partial^2 \psi) dx d\tau \right|,$$

one has

$$(2-20) \quad (1+t)^\lambda \partial^a \psi \partial^a Q(\psi, \partial \psi, \partial^2 \psi) \\ = \operatorname{div}((1+t)^\lambda (c^2(\rho) - 1) \nabla(\partial^a \psi) \partial^a \psi) - \frac{1}{2} \partial_i ((1+t)^{3\lambda} (\partial_i \psi) \partial^a (|\nabla \psi|^2) \partial^a \psi) \\ - \partial_t ((1+t)^\lambda \partial^a (|\nabla \psi|^2) \partial^a \psi) - (1+t)^\lambda (c^2(\rho) - 1) |\nabla \partial^a \psi|^2 \\ - (1+t)^\lambda (c^2(\rho))' \nabla \rho \cdot \nabla(\partial^a \psi) \partial^a \psi + \lambda (1+t)^{\lambda-1} \partial^a (|\nabla \psi|^2) \partial^a \psi \\ + (1+t)^\lambda \partial^a (|\nabla \psi|^2) \partial_t \partial^a \psi + \frac{1}{2} (1+t)^{3\lambda} (\Delta \psi) \partial^a (|\nabla \psi|^2) \partial^a \psi \\ + \frac{1}{2} (1+t)^{3\lambda} \nabla \psi \cdot \nabla(\partial^a \psi) \partial^a (|\nabla \psi|^2) + \text{l.o.t.}$$

From the expression $(\partial^{b_1} \psi)(\partial^{b_2} \psi) \dots (\partial^{b_l} \psi)$ ($l \geq 2$, $1 \leq |b_1| + \dots + |b_l| \leq N+1$) of the lower-order terms one readily obtains that there exists at most one b_j ($1 \leq j \leq l$) such that

$$\left[\frac{N+3}{2} \right] < |b_j| \leq N+1.$$

Moreover, $\left[\frac{N+3}{2} \right] \leq N-2$ by $N \geq 8$. Thus, applying (2-5)–(2-7) and subsequently substituting (2-18)–(2-20) into (2-17) completes the proof of Lemma 2.4. \square

Next we focus on the general time-weighted energy estimate of $\partial \Gamma^a \psi$ with $0 \leq |a| \leq N$ and $N \geq 8$.

Lemma 2.5 (time-weighted energy estimate of $\partial\Gamma^a\psi$ for $|a| \leq N$). *Let $\mu > 0$ and $\lambda \in (0, 1]$. Under assumption (2-5), we have that, for $t > 0$,*

$$(2-21) \quad \begin{aligned} & \sum_{0 \leq |a| \leq N} \int_{\mathbb{R}^3} ((1+t)^{2\lambda} |\partial\Gamma^a\psi|^2 + |\Gamma^a\psi|^2) dx \\ & + C \sum_{0 \leq |a| \leq N} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial\Gamma^a\psi|^2 dx d\tau \\ & \leq C\varepsilon^2 + C(1+K\varepsilon) \int_0^t A(\tau) \sum_{0 \leq |a| \leq N} \int_{\mathbb{R}^3} ((1+\tau)^{2\lambda} |\partial\Gamma^a\psi|^2 + \psi^2) dx d\tau, \end{aligned}$$

where the function A has been defined in Lemma 2.4.

Proof. Writing $\Gamma^a = \tilde{\Gamma}^b \partial^c$ with $\tilde{\Gamma} \in \{\Omega, S\}$, we will use induction on $|b|$ to prove (2-21). In view of Lemma 2.4, it is enough to assume that $|c| = 0$.

Suppose that (2-21) holds for $|b| \leq l-1$, where $1 \leq l \leq N$. We then intend to establish (2-21) for $|b| = l$.

Acting with $\tilde{\Gamma}^a$ (where $a = b$ and $|b| = l$) on both sides of (2-4) yields

$$(2-22) \quad \begin{aligned} & \square \tilde{\Gamma}^a \psi + \frac{\mu}{(1+t)^\lambda} \partial_t \tilde{\Gamma}^a \psi + \frac{2\lambda}{1+t} \partial_t \tilde{\Gamma}^a \psi \\ & = \sum_{|b_1| < |b|} \tilde{\Gamma}^{b_1} \partial^c \square \psi + \tilde{\Gamma}^a Q(\psi, \partial\psi, \partial^2\psi) \\ & \quad - \left[\tilde{\Gamma}^a, \frac{\mu}{(1+t)^\lambda} \partial_t \right] \psi - \left[\tilde{\Gamma}^a, \frac{2\lambda}{1+t} \partial_t \right] \psi + \tilde{\Gamma}^a ((\lambda-1)(1+t)^{-2} \psi). \end{aligned}$$

Starting from (2-22), as in the proof of Lemma 2.4, we can choose the multiplier

$$m(1+t)^{2\lambda} \partial_t \tilde{\Gamma}^a \psi + (1+t)^{2\lambda-1} \tilde{\Gamma}^a \psi + \kappa(1+t)^\lambda \tilde{\Gamma}^a \psi$$

to derive (2-21). For the commutators, we see from (2-4) that

$$(2-23) \quad \begin{aligned} & \left| \int_0^t \int_{\mathbb{R}^3} \left[\tilde{\Gamma}^a, \frac{\mu}{(1+t)^\lambda} \partial_t \right] \psi (1+t)^\lambda \tilde{\Gamma}^a \psi dx d\tau \right| \\ & \leq C \sum_{|a_1| < |a|} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda \square \tilde{\Gamma}^{a_1} \psi \tilde{\Gamma}^a \psi dx d\tau \right| \\ & \quad + C \sum_{|a_1| < |a|} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda \tilde{\Gamma}^{a_1} Q(\psi, \partial\psi, \partial^2\psi) \tilde{\Gamma}^a \psi dx d\tau \right| \\ & \quad + C \sum_{|a_1| < |a|} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{\lambda-1} \tilde{\Gamma}^a \psi (\partial_t \tilde{\Gamma}^{a_1} \psi + (1-\lambda)(1+\tau)^{-1} \tilde{\Gamma}^{a_1} \psi) dx d\tau \right| \end{aligned}$$

$$\begin{aligned}
 &\leq C\varepsilon^2 + C \sum_{|a_1| < |a|} \left| \int_{\mathbb{R}^3} (1+t)^\lambda \partial_t \tilde{\Gamma}^{a_1} \psi \tilde{\Gamma}^a \psi \, dx \right| \\
 &\quad + C \sum_{|a_1| < |a|} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda \tilde{\Gamma}^{a_1} \mathcal{Q}(\psi, \partial\psi, \partial^2\psi) \tilde{\Gamma}^a \psi \, dx \, d\tau \right| \\
 &\quad + C \sum_{|a_1| < |a|} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{\lambda-1} \tilde{\Gamma}^a \psi (\partial_t \tilde{\Gamma}^{a_1} \psi + (1-\lambda)(1+\tau)^{-1} \tilde{\Gamma}^{a_1} \psi) \, dx \, d\tau \right| \\
 &\quad + C \sum_{|a_1| < |a|} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda \partial \tilde{\Gamma}^{a_1} \psi \partial \tilde{\Gamma}^a \psi \, dx \, d\tau \right|.
 \end{aligned}$$

By the finite propagation speed, we have for $a > 0$

$$(2-24) \quad |\tilde{\Gamma}^a \psi| \leq C(1+t) \sum_{|a_1| < |a|} |\partial \tilde{\Gamma}^{a_1} \psi|.$$

It follows from (2-23)–(2-24) and a direct computation that

$$\begin{aligned}
 (2-25) \quad &\sum_{\substack{|b|=l, \\ |c| \leq N-l}} \int_{\mathbb{R}^3} ((1+t)^{2\lambda} |\partial \tilde{\Gamma}^b \partial^c \psi|^2 + |\tilde{\Gamma}^b \partial^c \psi|^2) \, dx \\
 &\quad + C \sum_{\substack{|b|=l, \\ |c| \leq N-l}} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \tilde{\Gamma}^b \partial^c \psi|^2 \, dx \, d\tau \\
 &\leq C\varepsilon^2 + CE_{l-1}(\psi(t)) + C \sum_{\substack{|b_1| < l, \\ |c_1| \leq N-|b_1|}} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \tilde{\Gamma}^{b_1} \partial^{c_1} \psi|^2 \, dx \, d\tau \\
 &\quad + C(1+K\varepsilon) \int_0^t A(\tau) \sum_{\substack{|b_1| \leq l, \\ |c_1| \leq N-|b_1|}} \int_{\mathbb{R}^3} ((1+\tau)^{2\lambda} |\partial \tilde{\Gamma}^{b_1} \partial^{c_1} \psi|^2 + |\tilde{\Gamma}^{b_1} \partial^{c_1} \psi|^2) \, dx \, d\tau \\
 &\quad + C \sum_{\substack{|b_1| \leq l, \\ |c_1| \leq N-|b_1|}} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) \tilde{\Gamma}^{b_1} \partial^{c_1} \mathcal{Q}(\psi, \partial\psi, \partial^2\psi) \, dx \, d\tau \right| \\
 &\quad + C \sum_{\substack{|b_1| \leq l, \\ |c_1| \leq N-|b_1|}} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda (\tilde{\Gamma}^a \psi) \tilde{\Gamma}^{b_1} \partial^{c_1} \mathcal{Q}(\psi, \partial\psi, \partial^2\psi) \, dx \, d\tau \right|.
 \end{aligned}$$

Next we deal with the last two terms in the right-hand side of (2-25). Note that

$$c^2(\rho) - 1 = -G(\psi, \partial\psi) \int_0^1 (c^2)′(-sG(\psi, \partial\psi)) \, ds,$$

where $G(\psi, \partial\psi) = (1+t)^\lambda \partial_t \psi + (1+t)^{\lambda-1} \psi + (1+t)^{2\lambda} |\nabla \psi|^2 / 2 + \mu \psi$. From this, it is readily seen that the typical terms in $Q(\psi, \partial\psi, \partial^2\psi)$ are of the form $\psi \Delta \psi$, $(1+t)^\lambda \partial_t \nabla \psi \cdot \nabla \psi$, and $(1+t)^{2\lambda} (\partial_i \psi) (\partial_j \psi) \partial_{ij} \psi$. We analyze them separately. Without loss of generality, we assume $|c_1| = 0$ in the last two terms of (2-25); the treatment of the other cases is easier.

Part A: *Estimates of*

$$\sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) \tilde{\Gamma}^{b_1} (\psi \Delta \psi) dx d\tau \right|$$

and

$$\sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda (\tilde{\Gamma}^a \psi) \tilde{\Gamma}^{b_1} (\psi \Delta \psi) dx d\tau \right|.$$

Note that

$$\tilde{\Gamma}^{b_1} (\psi \Delta \psi) = I_1 + I_2 + I_3,$$

where

$$\begin{aligned} I_1 &= \psi \Delta \tilde{\Gamma}^{b_1} \psi, \\ I_2 &= \sum_{\substack{|b_1|=|b_2|+|b_3|, \\ 1 \leq |b_2| \leq N-2}} (\tilde{\Gamma}^{b_2} \psi) \Delta \tilde{\Gamma}^{b_3} \psi, \\ I_3 &= \sum_{\substack{|b_1|=|b_2|+|b_3|, \\ N-1 \leq |b_2| \leq t}} (\tilde{\Gamma}^{b_2} \psi) \Delta \tilde{\Gamma}^{b_3} \psi. \end{aligned}$$

In view of $b_1 = a$ and

$$\begin{aligned} &(1+t)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) \psi \Delta \tilde{\Gamma}^a \psi \\ &= \operatorname{div}((1+t)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) \psi \nabla \tilde{\Gamma}^a \psi) + \frac{1}{2} \partial_t ((1+t)^{2\lambda} |\nabla \tilde{\Gamma}^a \psi|^2 \psi) \\ &\quad - (1+t)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) \nabla \psi \cdot \nabla \tilde{\Gamma}^a \psi - \lambda (1+t)^{\lambda-1} |\nabla \tilde{\Gamma}^a \psi|^2 \psi - \frac{1}{2} (1+t)^{2\lambda} |\nabla \tilde{\Gamma}^a \psi|^2 \partial_t \psi, \end{aligned}$$

we have by an integration by parts and (2-6)–(2-7)

$$\begin{aligned} (2-26) \quad &\left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) I_1 dx d\tau \right| \\ &\leq C\varepsilon^2 + CK\varepsilon \sum_{0 \leq |a| \leq N} \int_{\mathbb{R}^3} (1+t)^{2\lambda} |\partial \tilde{\Gamma}^a \psi|^2 dx \\ &\quad + CK\varepsilon \sum_{0 \leq |a| \leq N} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \tilde{\Gamma}^a \psi|^2 dx d\tau. \end{aligned}$$

Moreover, it follows from (2-7) and (2-9) that

$$\begin{aligned}
 (2-27) \quad & \int_{\mathbb{R}^3} |(1+t)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) I_2| dx \\
 & \leq (1+t)^{2\lambda} \|\langle r-t \rangle^{-1} \tilde{\Gamma}^{b_2} \psi\|_{L^\infty} \cdot \|\partial_t \tilde{\Gamma}^a \psi\| \cdot \|\langle r-t \rangle \Delta \tilde{\Gamma}^{b_3} \psi\| \\
 & \leq CK\varepsilon(1+t)^\lambda \|\partial_t \tilde{\Gamma}^a \psi\| \sum_{|b_4| \leq |b_3|+1} (\|\nabla \tilde{\Gamma}^{b_4} \psi\| + (1-\lambda)(1+t)^{-1} \|\tilde{\Gamma}^{b_4} \psi\|) \\
 & \leq CK\varepsilon(1+t)^\lambda \|\partial_t \tilde{\Gamma}^a \psi\| \sum_{|b_4| \leq |b_3|+1} \|\nabla \tilde{\Gamma}^{b_4} \psi\| + CK\varepsilon(1+t)^\lambda \|\partial_t \tilde{\Gamma}^a \psi\|^2 \\
 & \quad + CK\varepsilon(1-\lambda)(1+t)^{\lambda-2} \sum_{|b_4| \leq |b_3|+1} \|\tilde{\Gamma}^{b_4} \psi\|^2.
 \end{aligned}$$

On the other hand, we have that by (2-6) and Hardy's inequality

$$\begin{aligned}
 (2-28) \quad & \int_{\mathbb{R}^3} |(1+t)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) I_3| dx \\
 & \leq (1+t)^{2\lambda} \|(1+r) \Delta \tilde{\Gamma}^{b_3} \psi\|_{L^\infty} \cdot \|\partial_t \tilde{\Gamma}^a \psi\| \|(1+r)^{-1} \tilde{\Gamma}^{b_2} \psi\| \\
 & \leq CK\varepsilon(1+t)^\lambda \|\partial_t \tilde{\Gamma}^a \psi\| \sum_{|b_4| \leq |b_2|} \|\nabla \tilde{\Gamma}^{b_4} \psi\|.
 \end{aligned}$$

Combining (2-26)–(2-28) together with $0 \leq \lambda \leq 1$ (this means that the coefficient $CK\varepsilon(1-\lambda)(1+t)^{\lambda-2}$ of $\sum_{|b_4| \leq |b_3|+1} \|\tilde{\Gamma}^{b_4} \psi\|^2$ in the last line of (2-27) is nonnegative and in $L^1(0, \infty)$) yields

$$\begin{aligned}
 (2-29) \quad & \sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \Gamma^a \psi) \Gamma^{b_1} (\psi \Delta \psi) dx d\tau \right| \\
 & \leq C\varepsilon^2 + CK\varepsilon \sum_{0 \leq |a| \leq N} \int_{\mathbb{R}^3} (1+t)^{2\lambda} |\partial \tilde{\Gamma}^a \psi|^2 dx \\
 & \quad + CK\varepsilon \sum_{0 \leq |a| \leq N} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \tilde{\Gamma}^a \psi|^2 dx d\tau \\
 & \quad + CK\varepsilon \sum_{|b_4| \leq N} \int_0^t A(\tau) \int_{\mathbb{R}^3} |\tilde{\Gamma}^{b_4} \psi|^2 dx d\tau.
 \end{aligned}$$

Note that

$$\begin{aligned}
 (1+t)^\lambda (\tilde{\Gamma}^a \psi) \tilde{\Gamma}^{b_1} (\psi \Delta \psi) &= \sum_{|b_2|+|b_3|=|b_1|} (1+t)^\lambda (\tilde{\Gamma}^a \psi) (\tilde{\Gamma}^{b_2} \psi) \Delta \tilde{\Gamma}^{b_3} \psi \\
 &= \operatorname{div} \left(\sum_{|b_2|+|b_3|=|b_1|} (1+t)^\lambda (\tilde{\Gamma}^a \psi) (\tilde{\Gamma}^{b_2} \psi) \nabla \tilde{\Gamma}^{b_3} \psi \right) + \sum_{i=4}^5 I_i,
 \end{aligned}$$

where

$$I_4 = - \sum_{\substack{|b_2| \leq N-2, \\ |b_2|+|b_3|=|b_1|}} (1+t)^\lambda (\tilde{\Gamma}^{b_2} \psi) (\nabla \tilde{\Gamma}^a \psi) \cdot (\nabla \tilde{\Gamma}^{b_3} \psi),$$

$$I_5 = - \sum_{\substack{N-1 \leq |b_2| \leq l-1, \\ |b_2|+|b_3|=|b_1|}} (1+t)^\lambda (\tilde{\Gamma}^{b_2} \psi) (\nabla \tilde{\Gamma}^a \psi) \cdot (\nabla \tilde{\Gamma}^{b_3} \psi) \\ - \sum_{|b_2|+|b_3|=|b_1|} (1+t)^\lambda (\tilde{\Gamma}^a \psi) (\nabla \tilde{\Gamma}^{b_2} \psi) \cdot (\nabla \tilde{\Gamma}^{b_3} \psi).$$

Therefore, by (2-7) and Hardy's inequality, we have

$$\int_{\mathbb{R}^3} |I_4| dx \leq CK\varepsilon (1+t)^\lambda \|\nabla \tilde{\Gamma}^a \psi\| \sum_{|b_1|+3-N \leq |b_3| \leq N} \|\nabla \tilde{\Gamma}^{b_3} \psi\|$$

and

$$\int_{\mathbb{R}^3} |I_5| dx \leq CK\varepsilon \|(1+r)^{-1} \tilde{\Gamma}^{b_2} \psi \nabla \tilde{\Gamma}^a \psi\|_{L^1} \leq CK\varepsilon \|\nabla \tilde{\Gamma}^{b_2} \psi\| \|\nabla \tilde{\Gamma}^a \psi\|.$$

This yields

$$(2-30) \quad \sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda (\tilde{\Gamma}^a \psi) \tilde{\Gamma}^{b_1} (\psi \Delta \psi) dx d\tau \right| \\ \leq CK\varepsilon \sum_{0 \leq |a| \leq N} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \tilde{\Gamma}^a \psi|^2 dx d\tau.$$

Part B: *Estimates of*

$$\sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) \tilde{\Gamma}^{b_1} ((1+\tau)^\lambda \partial_t \nabla \psi \cdot \nabla \psi) dx d\tau \right|$$

and

$$\sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda (\tilde{\Gamma}^a \psi) \tilde{\Gamma}^{b_1} ((1+\tau)^\lambda \partial_t \nabla \psi \cdot \nabla \psi) dx d\tau \right|.$$

One has

$$\tilde{\Gamma}^{b_1} ((1+t)^\lambda \partial_t \nabla \psi \cdot \nabla \psi) \\ = (1+t)^\lambda \partial_t \nabla \tilde{\Gamma}^{b_1} \psi \cdot \nabla \psi + \sum_{N-2 \leq |b_2| \leq l-1} (1+t)^\lambda (\partial_t \nabla \tilde{\Gamma}^{b_2} \psi) \nabla \tilde{\Gamma}^{b_3} \psi \\ + \sum_{|b_2| \leq N-3} (1+t)^\lambda (\partial_t \nabla \tilde{\Gamma}^{b_2} \psi) \nabla \tilde{\Gamma}^{b_3} \psi \\ = \Pi_1 + \Pi_2 + \Pi_3.$$

By (2-8), we have

$$(2-31) \quad \sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) \Pi_1 dx d\tau \right| \\ \leq CK\varepsilon \sum_{0 \leq |a| \leq N} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \tilde{\Gamma}^a \psi|^2 dx d\tau.$$

In addition, it follows from (2-6), (2-9) and a direct computation that

$$(2-32) \quad (1+t)^{2\lambda} \|(\partial_t \Gamma^a \psi) \Pi_2\|_{L^1} \\ \leq (1+t)^{3\lambda} \sum_{|b_2| \leq N-4} \|\langle r-t \rangle^{-1} \nabla \Gamma^{b_2} \psi\|_{L^\infty} \cdot \|\partial_t \Gamma^a \psi\| \cdot \|\langle r-t \rangle \partial_t \nabla \Gamma^{b_2} \psi\| \\ \leq CK\varepsilon (1+t)^\lambda \|\partial_t \Gamma^a \psi\| \sum_{|c| \leq |b_2|+1} (\|\nabla \Gamma^c \psi\| + (1-\lambda)(1+t)^{-1} \|\Gamma^c \psi\|) \\ \leq CK\varepsilon (1+t)^\lambda \|\partial_t \tilde{\Gamma}^a \psi\| \sum_{|b_4| \leq |b_3|+1} \|\nabla \tilde{\Gamma}^{b_4} \psi\| + CK\varepsilon (1+t)^\lambda \|\partial_t \tilde{\Gamma}^a \psi\|^2 \\ + CK\varepsilon (1-\lambda)(1+t)^{\lambda-2} \sum_{|b_4| \leq |b_3|+1} \|\tilde{\Gamma}^{b_4} \psi\|^2.$$

Treating Π_3 , we obtain by (2-8)

$$(2-33) \quad \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) \Pi_3 dx d\tau \right| \leq CK\varepsilon \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \tilde{\Gamma}^a \psi|^2 dx d\tau.$$

Collecting (2-31)–(2-33) together with $0 \leq \lambda \leq 1$ (this means that the coefficient $CK\varepsilon(1-\lambda)(1+t)^{\lambda-2}$ of $\sum_{|b_4| \leq |b_3|+1} \|\tilde{\Gamma}^{b_4} \psi\|^2$ in the last line of (2-32) is nonnegative and in $L^1(0, \infty)$) yields

$$(2-34) \quad \sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \Gamma^a \psi) \Gamma^{b_1} ((1+t)^\lambda \partial_t \nabla \psi \cdot \nabla \psi) dx d\tau \right| \\ \leq CK\varepsilon \sum_{0 \leq |a| \leq N} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \tilde{\Gamma}^a \psi|^2 dx d\tau \\ + CK\varepsilon \sum_{|b_4| \leq N} \int_0^t A(\tau) \int_{\mathbb{R}^3} |\tilde{\Gamma}^{b_4} \psi|^2 dx d\tau.$$

In addition, one notes that

$$2(1+t)^{2\lambda} (\tilde{\Gamma}^a \psi) \tilde{\Gamma}^a (\partial_t \nabla \psi \cdot \nabla \psi) \\ = \sum_{|c| \leq |a|} \partial_t ((1+t)^{2\lambda} \tilde{\Gamma}^a \psi \Gamma^c (|\nabla \psi|^2)) \\ - 2\lambda(1+t)^{2\lambda-1} (\tilde{\Gamma}^a \psi) \tilde{\Gamma}^c (|\nabla \psi|^2) - (1+t)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) \tilde{\Gamma}^c (|\nabla \psi|^2).$$

From this, (2-6) and Hardy's inequality, we have

$$(2-35) \quad \sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda (\tilde{\Gamma}^a \psi) \tilde{\Gamma}^{b_1} ((1+\tau)^\lambda \partial_t \nabla \psi \cdot \nabla \psi) dx d\tau \right| \\ \leq C\varepsilon^2 + CK\varepsilon \sum_{0 \leq |a| \leq N} \int_{\mathbb{R}^3} (1+t)^{2\lambda} |\partial \tilde{\Gamma}^a \psi|^2 dx \\ + CK\varepsilon \sum_{0 \leq |a| \leq N} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \tilde{\Gamma}^a \psi|^2 dx d\tau.$$

Part C: *Estimates of*

$$\sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) \tilde{\Gamma}^{b_1} ((1+\tau)^{2\lambda} (\partial_i \psi)(\partial_j \psi) \partial_{ij} \psi) dx d\tau \right|$$

and

$$\sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda (\tilde{\Gamma}^a \psi) \tilde{\Gamma}^{b_1} ((1+\tau)^{2\lambda} (\partial_i \psi)(\partial_j \psi) \partial_{ij} \psi) dx d\tau \right|.$$

A direct computation yields

$$\begin{aligned} & \tilde{\Gamma}^{b_1} ((\partial_i \psi)(\partial_j \psi) \partial_{ij} \psi) \\ &= \partial_i \psi \partial_j \psi \partial_{ij} \tilde{\Gamma}^{b_1} \psi + \sum_{N-2 \leq |b_2| \leq |b_1|-1} (\nabla^2 \tilde{\Gamma}^{b_2} \psi)(\nabla \tilde{\Gamma}^{b_3} \psi) \nabla \tilde{\Gamma}^{b_4} \psi \\ & \quad + \sum_{|b_2| \leq N-3} (\nabla^2 \tilde{\Gamma}^{b_2} \psi)(\nabla \tilde{\Gamma}^{b_3} \psi) \nabla \tilde{\Gamma}^{b_4} \psi \\ &= \text{III}_1 + \text{III}_2 + \text{III}_3. \end{aligned}$$

As in the treatment of II_1 in Part B, we have

$$(2-36) \quad \sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) \text{III}_1 dx d\tau \right| \\ \leq C\varepsilon^2 + CK\varepsilon \sum_{0 \leq |a| \leq N} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \tilde{\Gamma}^a \psi|^2 dx d\tau.$$

By (2-6) and (2-9), for the term III_2 , we have

$$(2-37) \quad (1+t)^{4\lambda} \|(\partial_t \tilde{\Gamma}^a \psi)(\nabla^2 \tilde{\Gamma}^{b_2} \psi)(\nabla \tilde{\Gamma}^{b_3} \psi) \nabla \tilde{\Gamma}^{b_4} \psi\|_{L^1} \\ \leq (1+t)^{4\lambda} \langle r-t \rangle^{-1} \|(\nabla \tilde{\Gamma}^{b_3} \psi) \nabla \tilde{\Gamma}^{b_4} \psi\|_{L^\infty} \cdot \|\partial_t \tilde{\Gamma}^a \psi\| \cdot \|(r-t) \nabla^2 \tilde{\Gamma}^{b_2} \psi\| \\ \leq CK\varepsilon (1+t)^\lambda \|\partial_t \tilde{\Gamma}^a \psi\| \sum_{|b_4| \leq |b_3|+1} \|\nabla \tilde{\Gamma}^{b_4} \psi\| + CK\varepsilon (1+t)^\lambda \|\partial_t \tilde{\Gamma}^a \psi\|^2 \\ + CK\varepsilon (1-\lambda) (1+t)^{\lambda-2} \sum_{|b_4| \leq |b_3|+1} \|\tilde{\Gamma}^{b_4} \psi\|^2.$$

By (2-6) and (2-8), for the term III_3 , one has

$$(2-38) \quad (1+t)^{4\lambda} \|(\partial_t \tilde{\Gamma}^a \psi)(\nabla^2 \tilde{\Gamma}^{b_2} \psi)(\nabla \tilde{\Gamma}^{b_3} \psi) \nabla \tilde{\Gamma}^{b_4} \psi\|_{L^1} \\ \leq CK\varepsilon(1+t)^\lambda \|\partial_t \tilde{\Gamma}^a \psi\| \sum_{|c| \leq |b_1|} \|\nabla \tilde{\Gamma}^c \psi\|.$$

Collecting (2-36)–(2-38) together with $0 \leq \lambda \leq 1$ (this means that the coefficient $CK\varepsilon(1-\lambda)(1+t)^{\lambda-2}$ of $\sum_{|b_4| \leq |b_3|+1} \|\tilde{\Gamma}^{b_4} \psi\|^2$ in the last line of (2-37) is nonnegative and in $L^1(0, \infty)$) yields

$$(2-39) \quad \sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^{2\lambda} (\partial_t \tilde{\Gamma}^a \psi) \tilde{\Gamma}^{b_1} ((1+\tau)^{2\lambda} (\partial_i \psi)(\partial_j \psi) \partial_{ij} \psi) dx d\tau \right| \\ \leq CK\varepsilon \sum_{0 \leq |a| \leq N} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \tilde{\Gamma}^a \psi|^2 dx d\tau \\ + CK\varepsilon \sum_{|b_4| \leq N} \int_0^t A(\tau) \int_{\mathbb{R}^3} |\tilde{\Gamma}^{b_4} \psi|^2 dx d\tau.$$

In addition,

$$2(1+t)^{3\lambda} (\Gamma^a \psi) \Gamma^{b_1} ((\partial_i \psi)(\partial_j \psi) \partial_{ij} \psi) \\ = \text{div}((1+t)^{3\lambda} (\Gamma^a \psi)(\nabla \psi) \Gamma^{b_1} (|\nabla \psi|^2)) - (1+t)^{3\lambda} (\nabla \Gamma^a \psi)(\nabla \psi) \Gamma^{b_1} (|\nabla \psi|^2) \\ - (1+t)^{3\lambda} (\Gamma^a \psi)(\Delta \psi) \Gamma^{b_1} (|\nabla \psi|^2) \\ + \sum_{|b_2| \leq |b_1|-1} (1+t)^{3\lambda} (\Gamma^a \psi)(\nabla^2 \Gamma^{b_2} \psi)(\nabla \Gamma^{b_3} \psi) \nabla \Gamma^{b_4} (|\psi|^2).$$

Together with (2-6) and Hardy's inequality this yields

$$(2-40) \quad \sum_{|b_1| \leq N} \left| \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda (\Gamma^a \psi) \Gamma^{b_1} ((1+\tau)^{2\lambda} (\partial_i \psi)(\partial_j \psi) \partial_{ij} \psi) dx d\tau \right| \\ \leq CK\varepsilon \sum_{|a| \leq N} \int_0^t \int_{\mathbb{R}^3} (1+\tau)^\lambda |\partial \Gamma^a \psi|^2 dx d\tau.$$

Therefore, substituting (2-29)–(2-30), (2-34)–(2-35), and (2-39)–(2-40) into (2-25) and utilizing the smallness of $\varepsilon > 0$ gives (2-21). \square

Based on Lemmas 2.4 and 2.5, we now prove Theorem 1.1.

Proof of Theorem 1.1. By Lemmas 2.4 and 2.5, one has that, for fixed $N \geq 8$,

$$E_N(t) \leq C\varepsilon^2 + C(1+K\varepsilon) \int_0^t A(t') E_N(t') dt'.$$

Choosing the constants $K > 0$ large and $\varepsilon > 0$ small, by Gronwall’s inequality one gets that, for any $t \geq 0$,

$$E_N(t) \leq e^{C(1+K\varepsilon)\|A(t)\|_{L^1}} E_N(0) \leq \frac{1}{2} K^2 \varepsilon^2.$$

Thus, [Theorem 1.1](#) is proved by the assumption that $E_N(t) \leq K^2 \varepsilon^2$ and a continuous induction argument. □

3. Blowup for small data in case $\lambda > 1$

In this section, we shall prove the blowup result of [Theorem 1.2](#) which is valid in case $\lambda > 1$.

Proof of Theorem 1.2. We divide the proof into two cases.

Case I: $\gamma = 2$. Let (ρ, u) be a smooth solution of (1-1). For $l > 0$, we define

$$(3-1) \quad P(t, l) = \int_{|x|>l} \eta(x, l)(\rho(t, x) - \bar{\rho}) dx,$$

where

$$\eta(x, l) = |x|^{-1}(|x| - l)^2.$$

Employing the first equation in (1-1) and an integration by parts, we see that

$$\begin{aligned} \partial_t P(t, l) &= \int_{|x|>l} \eta(x, l) \partial_t (\rho(t, x) - \bar{\rho}) dx = - \int_{|x|>l} \eta(x, l) \operatorname{div}(\rho u)(t, x) dx \\ &= \int_{|x|>l} (\nabla_x \eta)(x, l) \cdot (\rho u)(t, x) dx, \end{aligned}$$

where we have used the fact that $\eta(x, l) = 0$ on $|x| = l$ and that $u(t, x) = 0$ for $|x| \geq t + M$.

By differentiating $\partial_t P(t, l)$ again and using the second equation in (1-1), we find that

$$\begin{aligned} (3-2) \quad \partial_t^2 P(t, l) &= \int_{|x|>l} (\nabla_x \eta)(x, l) \cdot \partial_t (\rho u)(t, x) dx \\ &= - \sum_{i,j} \int_{|x|>l} (\partial_{x_i} \eta) \partial_{x_j} (\rho u_i u_j) dx - \int_{|x|>l} (\nabla_x \eta)(x, l) \cdot \nabla(p - \bar{p}) dx \\ &\quad - \frac{\mu}{(1+t)^\lambda} \int_{|x|>l} (\nabla_x \eta)(x, l) \cdot (\rho u)(t, x) dx, \end{aligned}$$

where $\nabla_x \eta(x, l) = |x|^{-3}(|x|^2 - l^2)x$ vanishes on $|x| = l$ and $\bar{p} = p(\bar{\rho})$. Integration by parts implies that

$$\begin{aligned}
 (3-3) \quad \partial_t^2 P(t, l) + \frac{\mu}{(1+t)^\lambda} \partial_t P(t, l) &= \sum_{i,j} \int_{|x|>l} (\partial_{x_i x_j}^2 \eta) \rho u_i u_j \, dx + \int_{|x|>l} (\Delta \eta)(p - \bar{p}) \, dx \\
 &\equiv J_1(t, l) + J_2(t, l),
 \end{aligned}$$

where we have used that $p - \bar{p}$ vanishes for $|x| \geq t + M$. A direct computation of $\partial_{x_i x_j}^2 \eta$ shows that

$$\begin{aligned}
 (3-4) \quad J_1(t, l) &= \int_{|x|>l} \frac{2l^2}{|x|^3} \rho \left(\frac{x}{|x|} \cdot u \right)^2 \, dx \\
 &\quad - \int_{|x|>l} \frac{|x|^2 - l^2}{|x|^3} \rho \left(\frac{x}{|x|} \cdot u \right)^2 \, dx + \int_{|x|>l} \frac{|x|^2 - l^2}{|x|^3} \rho |u|^2 \, dx \geq 0.
 \end{aligned}$$

On the other hand, notice that

$$(3-5) \quad \partial_t^2 \eta(x, l) = 2|x|^{-1} = \Delta_x \eta(x, l).$$

Then

$$(3-6) \quad J_2(t, l) = \int_{|x|>l} \partial_t^2 \eta(x, l)(p(t, x) - \bar{p}) \, dx = \partial_t^2 \int_{|x|>l} \eta(x, l)(p(t, x) - \bar{p}) \, dx,$$

where we have used the fact that η and $\partial_t \eta$ vanish on $|x| = l$. Combining (3-3)–(3-6), we arrive at

$$(3-7) \quad \partial_t^2 P(t, l) - \partial_t^2 P(t, l) + \frac{\mu}{(1+t)^\lambda} \partial_t P(t, l) = f(t, l) \equiv J_1(t, l) + G(t, l) \geq G(t, l),$$

where

$$(3-8) \quad G(t, l) = \partial_t^2 \int_{|x|>l} \eta(x, l)(p - \bar{p} - (\rho - \bar{\rho})) \, dx = \int_{|x|>l} 2|x|^{-1}(p - \bar{p} - (\rho - \bar{\rho})) \, dx.$$

Thanks to $\gamma = 2$ and the sound speed $\bar{c} = \sqrt{2A\bar{\rho}} = 1$, we have

$$(3-9) \quad p - \bar{p} - (\rho - \bar{\rho}) = A(\rho^2 - \bar{\rho}^2 - 2\bar{\rho}(\rho - \bar{\rho})) = A(\rho - \bar{\rho})^2.$$

Substituting (3-9) into (3-8) gives

$$G(t, l) \geq 0.$$

For M_0 satisfying the condition (1-11), let $\Sigma \equiv \{(t, l) : t \geq 0, t + M_0 \leq l \leq t + M\}$ be the strip domain. By applying Riemann’s representation (see [Courant and Hilbert

1962, §5.5]) with the assumptions (1-9)–(1-11), we see that the solution $P(t, l)$ to (3-7) is nonnegative in Σ . We put its proof in the [Appendix](#). Rewrite (3-7) as

$$\partial_t^2 P(t, l) - \partial_l^2 P(t, l) + \frac{\mu}{(1+t)^\lambda} (\partial_t P(t, l) - \partial_l P(t, l)) = f(t, l) - \frac{\mu}{(1+t)^\lambda} \partial_l P(t, l).$$

By the method of characteristics we have

$$\begin{aligned} P(t, l) &= \frac{1}{2} P(0, l+t) + \frac{1}{2\beta(t)} P(0, l-t) + \frac{1}{2} \int_0^t \frac{1}{\beta(\tau)} \frac{\mu}{(1+\tau)^\lambda} P(0, l+t-2\tau) d\tau \\ &\quad + \int_0^t \frac{1}{\beta(\tau)} \partial_t P(0, l+t-2\tau) d\tau + \frac{1}{2} \int_0^t \int_{l-t+\tau}^{l+t-\tau} \frac{\beta(\tau)}{\beta\left(\frac{l+t+\tau-y}{2}\right)} f(\tau, y) dy d\tau \\ &\quad + \frac{1}{2} \int_0^t \frac{\beta(\tau)}{\beta(t)} \frac{\mu}{(1+\tau)^\lambda} P(\tau, l-t+\tau) d\tau \\ &\quad + \frac{1}{2} \int_0^t \int_\tau^t \frac{\beta(\tau)}{\beta(s)} \frac{\mu^2}{(1+\tau)^\lambda (1+s)^\lambda} P(\tau, l+t-2s+\tau) ds d\tau \\ &\quad - \frac{1}{2} \int_0^t \frac{\mu}{(1+\tau)^\lambda} P(\tau, l+t-\tau) d\tau; \end{aligned}$$

see (1-12). Together with assumptions (1-9)–(1-10) and $P(t, l) \geq 0$ in Σ this yields, for $l \geq t + M_0$,

$$\begin{aligned} (3-10) \quad P(t, l) &\geq \frac{1}{2\beta(t)} q_0(l-t) + \frac{1}{2} \int_0^t \int_{l-t+\tau}^{l+t-\tau} \frac{\beta(\tau)}{\beta\left(\frac{l+t+\tau-y}{2}\right)} G(\tau, y) dy d\tau \\ &\quad - \frac{1}{2} \int_0^t \frac{\mu}{(1+\tau)^\lambda} P(\tau, l+t-\tau) d\tau. \end{aligned}$$

Define the function

$$(3-11) \quad F(t) \equiv \int_0^t (t-\tau) \int_{\tau+M_0}^{\tau+M} P(\tau, l) \frac{dl}{l} d\tau.$$

Then, by (3-10), we have that

$$\begin{aligned} (3-12) \quad F''(t) &= \int_{t+M_0}^{t+M} P(t, l) \frac{dl}{l} \\ &\geq \frac{1}{2\beta(t)} \int_{t+M_0}^{t+M} q_0(l-t) \frac{dl}{l} + \frac{1}{2} \int_{t+M_0}^{t+M} \int_0^t \int_{l-t+\tau}^{l+t-\tau} \frac{\beta(\tau)}{\beta\left(\frac{l+t+\tau-y}{2}\right)} G(\tau, y) dy d\tau \frac{dl}{l} \\ &\quad - \frac{1}{2} \int_{t+M_0}^{t+M} \int_0^t \frac{\mu}{(1+\tau)^\lambda} P(\tau, l+t-\tau) d\tau \frac{dl}{l} \\ &\equiv J_3 + J_4 - J_5. \end{aligned}$$

From $\lambda > 1$ and assumption (1-9), we see that

$$(3-13) \quad J_3 \geq \frac{c_1}{t+M} \int_{t+M_0}^{t+M} q_0(l-t) dl = \frac{c_1}{t+M} \int_{M_0}^M q_0(l) dl = \frac{c_2 \varepsilon}{t+M},$$

where $c_1, c_2 > 0$ are constants independent of ε . Note that $P(\tau, y)$ is supported in $\{y : y \leq \tau + M\}$ and nonnegative in Σ . Hence, there exists a constant $C_1 > 0$ such that

$$(3-14) \quad J_5 \leq \frac{C_1}{(1+t)^\lambda} \int_0^t \int_{\tau+M_0}^{\tau+M} P(\tau, y) \frac{dy}{y} d\tau = \frac{C_1}{(1+t)^\lambda} F'(t).$$

Substituting (3-14) into (3-12) gives

$$(3-15) \quad F''(t) + \frac{C_1}{(1+t)^\lambda} F'(t) \geq J_3 + J_4.$$

To bound J_4 from below, we write

$$(3-16) \quad \begin{aligned} J_4 &= \frac{1}{2} \int_0^{t-M_1} \int_{\tau+M_0}^{\tau+M} G(\tau, y) \int_{t+M_0}^{y+t-\tau} \frac{\beta(\tau)}{\beta\left(\frac{l+t+\tau-y}{2}\right)} \frac{dl}{l} dy d\tau \\ &\quad + \frac{1}{2} \int_{t-M_1}^t \int_{\tau+M_0}^{2t-\tau+M_0} G(\tau, y) \int_{t+M_0}^{y+t-\tau} \frac{\beta(\tau)}{\beta\left(\frac{l+t+\tau-y}{2}\right)} \frac{dl}{l} dy d\tau \\ &\quad + \frac{1}{2} \int_{t-M_1}^t \int_{2t-\tau+M_0}^{\tau+M} G(\tau, y) \int_{y-t+\tau}^{y+t-\tau} \frac{\beta(\tau)}{\beta\left(\frac{l+t+\tau-y}{2}\right)} \frac{dl}{l} dy d\tau \\ &\equiv J_{4,1} + J_{4,2} + J_{4,3}, \end{aligned}$$

where $M_1 = (M - M_0)/2$. For $t < M_1$, $t - M_1$ in the limits of integration is replaced by 0. By $\lambda > 1$, for the integrand in $J_{4,1}$ we have that

$$(3-17) \quad \int_{t+M_0}^{y+t-\tau} \frac{\beta(\tau)}{\beta\left(\frac{l+t+\tau-y}{2}\right)} \frac{dl}{l} \geq c \frac{y - \tau - M_0}{t + M} \geq c \frac{(t - \tau)(y - \tau - M_0)^2}{(t + M)^2}.$$

Analogously, for the integrands in $J_{4,2}$ and $J_{4,3}$ we have that

$$(3-18) \quad \int_{t+M_0}^{y+t-\tau} \frac{\beta(\tau)}{\beta\left(\frac{l+t+\tau-y}{2}\right)} \frac{dl}{l} \geq c \frac{(t - \tau)(y - \tau - M_0)^2}{(t + M)^2}$$

and

$$(3-19) \quad \int_{y-t+\tau}^{y+t-\tau} \frac{\beta(\tau)}{\beta\left(\frac{l+t+\tau-y}{2}\right)} \frac{dl}{l} \geq c \frac{t - \tau}{t + M} \geq c \frac{(t - \tau)(y - \tau - M_0)^2}{(t + M)^2},$$

where $c > 0$ is a constant. Substituting (3-17)–(3-19) into (3-16) yields

$$J_4 \geq \frac{c}{(t + M)^2} \int_0^t (t - \tau) \int_{\tau+M_0}^{\tau+M} (y - \tau - M_0)^2 \partial_y^2 \tilde{G}(\tau, y) dy d\tau,$$

where $\tilde{G}(t, l) = \int_{|x|>l} \eta(x, l)(p - \bar{p} - (\rho - \bar{\rho})) dx$. Note that $\tilde{G}(\tau, y) = \partial_y \tilde{G}(\tau, y) = 0$ for $y = \tau + M$. Thus, it follows from the integration by parts together with (3-8)–(3-9) that

$$\begin{aligned}
 (3-20) \quad J_4 &\geq \frac{c}{(t+M)^2} \int_0^t (t-\tau) \int_{\tau+M_0}^{\tau+M} \tilde{G}(\tau, y) dy d\tau \\
 &\geq \frac{c}{(t+M)^2} \int_0^t (t-\tau) \int_{\tau+M_0}^{\tau+M} \int_{|x|>y} \eta(x, y)(\rho(\tau, x) - \bar{\rho})^2 dx dy d\tau \\
 &\equiv \frac{c}{(t+M)^2} J_6.
 \end{aligned}$$

By applying the Cauchy–Schwartz inequality to $F(t)$ defined by (3-11), we arrive at

$$(3-21) \quad F^2(t) \leq J_6 \int_0^t (t-\tau) \int_{\tau+M_0}^{\tau+M} \int_{y<|x|<\tau+M} \eta(x, y) dx \frac{dy}{y^2} d\tau \equiv J_6 J_7.$$

We estimate J_7 as

$$\begin{aligned}
 (3-22) \quad J_7 &= \int_0^t (t-\tau) \int_{\tau+M_0}^{\tau+M} \int_{y<|x|<\tau+M} \frac{(|x|-y)^2}{|x|} dx \frac{dy}{y^2} d\tau \\
 &= \int_0^t (t-\tau) \int_{\tau+M_0}^{\tau+M} \int_y^{\tau+M} 4\pi l(l-y)^2 dl \frac{dy}{y^2} d\tau \\
 &\leq C \int_0^t (t-\tau) \int_{\tau+M_0}^{\tau+M} (\tau+M)(\tau+M-y)^3 \frac{dy}{y^2} d\tau \\
 &\leq C \int_0^t (t-\tau)(\tau+M) \int_{\tau+M_0}^{\tau+M} \frac{dy}{y^2} d\tau \\
 &\leq C \int_0^t \frac{t-\tau}{\tau+M} d\tau \leq C(t+M) \log(t/M+1).
 \end{aligned}$$

Combining (3-13), (3-15) and (3-20)–(3-22) gives the ordinary differential inequalities

$$(3-23) \quad F''(t) + \frac{C_1}{(1+t)^\lambda} F'(t) \geq \frac{c_2 \varepsilon}{t+M}, \quad t \geq 0,$$

$$(3-24) \quad F''(t) + \frac{C_1}{(1+t)^\lambda} F'(t) \geq C[(t+M)^3 \log(t/M+1)]^{-1} F^2(t), \quad t \geq 0.$$

Next, we apply (3-23)–(3-24) to prove that the lifespan T_ε of smooth solution $F(t)$ is finite for all $0 < \varepsilon \leq \varepsilon_0$. The fact that $F(0) = F'(0) = 0$, together with (3-23), yields

$$(3-25) \quad F'(t) \geq C\varepsilon \log(t/M+1), \quad t \geq 0,$$

$$(3-26) \quad F(t) \geq C\varepsilon(t+M) \log(t/M+1), \quad t \geq t_1 \equiv Me^2,$$

where the constant $C > 0$ is independent of ε . Substituting (3-26) into (3-24) derives

$$F''(t) + \frac{C_1}{(1+t)^\lambda} F'(t) \geq C\varepsilon^2(t+M)^{-1} \log(t/M+1), \quad t \geq t_1,$$

which leads to the improvement

$$(3-27) \quad F(t) \geq C\varepsilon^2(t+M) \log^2(t/M+1), \quad t \geq t_2 \equiv Me^3 > t_1.$$

Substituting this into (3-24) derives

$$(3-28) \quad F''(t) + \frac{C_1}{(1+t)^\lambda} F'(t) \geq C\varepsilon^2(t+M)^{-2} \log(t/M+1)F(t), \quad t \geq t_2.$$

It follows from (3-25) that $F'(t) \geq 0$ for $t \geq 0$. Then multiplying (3-28) by $F'(t)$ and integrating from t_3 (which will be chosen later) to t yield

$$F'(t)^2 \geq C_2 F'(t_3)^2 + C_3 \varepsilon^2 \int_{t_3}^t (s+M)^{-2} \log(s/M+1) [F(s)^2]' ds.$$

Integrating by parts yields

$$(3-29) \quad \begin{aligned} F'(t)^2 &\geq C_2 F'(t_3)^2 \\ &+ C_3 \varepsilon^2 \left((t+M)^{-2} \log(t/M+1) F(t)^2 - (t_3+M)^{-2} \log(t_3/M+1) F(t_3)^2 \right) \\ &- \int_{t_3}^t \left(\frac{\log(s/M+1)}{(s+M)^2} \right)' F(s)^2 ds, \quad t \geq t_3, \end{aligned}$$

where

$$\left(\frac{\log(s/M+1)}{(s+M)^2} \right)' \leq 0$$

for $t \geq t_3 \geq t_2$. On the other hand, (3-23) implies

$$\left(e^{-\frac{C_1}{\lambda-1}[(1+t)^{1-\lambda}-1]} F'(t) \right)' \geq 0, \quad t \geq 0,$$

which yields for $0 \leq t \leq \tau$

$$(3-30) \quad F'(t) \leq e^{\frac{C_1}{\lambda-1}[(1+t)^{1-\lambda}-(1+\tau)^{1-\lambda}]} F'(\tau).$$

Together with $F(0) = 0$, this yields

$$(3-31) \quad F(t) = \int_0^t F'(s) ds \leq C_4 t F'(t), \quad t > 0.$$

Choose

$$(3-32) \quad t_3 = M \left(e^{\frac{C_2}{2C_3 C_4 \varepsilon^2}} - 1 \right)$$

which satisfies $2C_3C_4 \log(t_3/M + 1)\varepsilon^2 = C_2$. Together with (3-29) and (3-31), this yields

$$(3-33) \quad F'(t) \geq \sqrt{C_3}\varepsilon(t + M)^{-1} \log^{\frac{1}{2}}(t/M + 1)F(t), \quad t \geq t_3.$$

By integrating (3-33) from t_3 to t , we arrive at

$$\log \frac{F(t)}{F(t_3)} \geq C\varepsilon \log^{\frac{3}{2}}\left(\frac{t + M}{t_3 + M}\right), \quad t \geq t_3.$$

If $t \geq t_4 \equiv Ct_3^2$, we then have

$$\log \frac{F(t)}{F(t_3)} \geq 8 \log(t/M + 1).$$

Together with (3-27) for $F(t_3)$, this yields

$$(3-34) \quad F(t) \geq C\varepsilon^2(t + M)^8, \quad t \geq t_4.$$

Substituting this into (3-24) derives

$$F''(t) + \frac{C_1}{(1 + t)^\lambda} F'(t) \geq C\varepsilon F(t)^{\frac{3}{2}}, \quad t \geq t_4.$$

Multiplying this differential inequality by $F'(t)$ and integrating from t_4 to t yields

$$F'(t)^2 \geq C\varepsilon(F(t)^{\frac{5}{2}} - F(t_4)^{\frac{5}{2}}).$$

On the other hand, (3-30) and (3-31) imply that, for $t \geq t_4$,

$$F(t) = F'(\xi)(t - t_4) + F(t_4) \geq CF'(t_4)(t - t_4) \geq CF(t_4) \frac{t - t_4}{t_4},$$

where $t_4 \leq \xi \leq t$. If $t \geq t_5 \equiv Ct_4$, then we have

$$F(t)^{\frac{5}{2}} - F(t_4)^{\frac{5}{2}} \geq \frac{1}{2}F(t)^{\frac{5}{2}}.$$

Thus

$$(3-35) \quad F'(t) \geq C\sqrt{\varepsilon}F(t)^{\frac{5}{4}}, \quad t \geq t_5.$$

If $T_\varepsilon > 2t_5$, then integrating (3-35) from t_5 to T_ε derives

$$F(t_5)^{-\frac{1}{4}} - F(T_\varepsilon)^{-\frac{1}{4}} \geq C\sqrt{\varepsilon}T_\varepsilon.$$

We see from (3-34) and $t_5 = Ct_3^2$ that

$$F(t_5) \geq C\varepsilon^2 e^{C/\varepsilon^2},$$

which together with $F(T_\varepsilon) > 0$ is a contradiction. Thus, $T_\varepsilon \leq 2t_5 = Ct_3^2$. From the choice of t_3 in (3-32), we see that $T_\varepsilon \leq e^{C/\varepsilon^2}$.

Case 2: $\gamma > 1$ and $\gamma \neq 2$. Recall that the sound speed is $\bar{c} = \sqrt{\gamma A \bar{\rho}^{\gamma-1}} = 1$. Instead of (3-9) we have

$$p - \bar{p} - (\rho - \bar{\rho}) = A(\rho^\gamma - \bar{\rho}^\gamma - \gamma \bar{\rho}^{\gamma-1}(\rho - \bar{\rho})) \equiv A\psi(\rho, \bar{\rho}).$$

The convexity of ρ^γ for $\gamma > 1$ implies that $\psi(\rho, \bar{\rho})$ is positive for $\rho \neq \bar{\rho}$. Applying Taylor's theorem, we have

$$\psi(\rho, \bar{\rho}) \geq C(\gamma, \bar{\rho}) \Phi_\gamma(\rho, \bar{\rho}),$$

where $C(\gamma, \bar{\rho})$ is a positive constant and Φ_γ is given by

$$\Phi_\gamma(\rho, \bar{\rho}) = \begin{cases} (\bar{\rho} - \rho)^\gamma, & \rho < \frac{1}{2}\bar{\rho}, \\ (\rho - \bar{\rho})^2, & \frac{1}{2}\bar{\rho} \leq \rho \leq 2\bar{\rho}, \\ (\rho - \bar{\rho})^\gamma, & \rho > 2\bar{\rho}. \end{cases}$$

For $\gamma > 2$, we have that $(\bar{\rho} - \rho)^\gamma = (\bar{\rho} - \rho)^2(\bar{\rho} - \rho)^{\gamma-2} \geq C(\gamma, \bar{\rho})(\rho - \bar{\rho})^2$ for $2\rho < \bar{\rho}$ and $(\rho - \bar{\rho})^\gamma = (\rho - \bar{\rho})^2(\rho - \bar{\rho})^{\gamma-2} \geq C(\gamma, \bar{\rho})(\rho - \bar{\rho})^2$ for $\rho > 2\bar{\rho}$. Thus, $\psi(\rho, \bar{\rho}) \geq C(\gamma, \bar{\rho})(\rho - \bar{\rho})^2$. In this case, Theorem 1.2 can be shown completely analogously to Case 1.

Next we treat the case $1 < \gamma < 2$. We define $F(t)$ as in (3-11),

$$F(t) = \int_0^t \int_{\tau+M_0}^{\tau+M} \frac{1}{l} \int_{|x|>l} \frac{(|x|-l)^2}{|x|} (\rho(\tau, x) - \bar{\rho}) dx dl d\tau.$$

Similarly to the case of $\gamma = 2$, we have

$$(3-36) \quad F''(t) \geq J_3 + J_4 - J_5,$$

where

$$\begin{aligned} J_3 &\geq \frac{C\varepsilon}{t+M}, \\ J_4 &\geq C(t+M)^{-2} \tilde{J}_6, \\ J_5 &\leq \frac{C_1}{(1+t)^\lambda} F'(t), \end{aligned}$$

and

$$\tilde{J}_6 = \int_0^t (t-\tau) \int_{\tau+M_0}^{\tau+M} \int_{|x|>y} \frac{(|x|-y)^2}{|x|} \Phi_\gamma(\rho(\tau, x) - \bar{\rho}) dx dy d\tau.$$

Denote $\Omega_1 = \{(\tau, x) : \bar{\rho} \leq \rho(\tau, x) \leq 2\bar{\rho}\}$, $\Omega_2 = \{(\tau, x) : \rho(\tau, x) > 2\bar{\rho}\}$, and $\Omega_3 = \{(\tau, x) : \rho(\tau, x) < \bar{\rho}\}$. Divide $F(t)$ into a sum of the three integrals over the domains Ω_i ($1 \leq i \leq 3$)

$$F(t) = F_1(t) + F_2(t) + F_3(t) \equiv \int_{\Omega_1} \dots + \int_{\Omega_2} \dots + \int_{\Omega_3} \dots$$

Corresponding to the three parts of $F(t)$, we define $\tilde{J}_6 \equiv \tilde{J}_{6,1} + \tilde{J}_{6,2} + \tilde{J}_{6,3}$. In view of $F(t) \geq 0$ and $F_3(t) \leq 0$, we have

$$F(t) \leq F_1(t) + F_2(t).$$

Applying Hölder's inequality for the domains Ω_1 and Ω_2 , we obtain that

$$\begin{aligned} F(t) &\leq \tilde{J}_{6,1}^{\frac{1}{2}} \left(\int_0^t (t-\tau) \int_{\tau+M_0}^{\tau+M} \frac{1}{y^2} \int_{y<|x|\leq\tau+M} \frac{(|x|-y)^2}{|x|} dx dy d\tau \right)^{\frac{1}{2}} \\ &\quad + \tilde{J}_{6,2}^{\frac{1}{\gamma}} \left(\int_0^t (t-\tau) \int_{\tau+M_0}^{\tau+M} \frac{1}{y^{\frac{\gamma}{\gamma-1}}} \int_{y<|x|\leq\tau+M} \frac{(|x|-y)^2}{|x|} dx dy d\tau \right)^{\frac{\gamma-1}{\gamma}} \\ &\leq \tilde{J}_6^{\frac{1}{2}} (t+M)^{\frac{1}{2}} \log^{\frac{1}{2}}(t/M+1) + \tilde{J}_6^{\frac{1}{\gamma}} (t+M)^{\frac{\gamma-1}{\gamma}} \\ &= (\tilde{J}_6(t+M)^{-1})^{\frac{1}{2}} (t+M) \log^{\frac{1}{2}}(t/M+1) + (\tilde{J}_6(t+M)^{-1})^{\frac{1}{\gamma}} (t+M). \end{aligned}$$

In view of $1 < \gamma < 2$, we have $\frac{1}{2\gamma} < \frac{1}{2} < \frac{1}{\gamma}$. Applying Young's inequality yields

$$F(t) \leq \left((\tilde{J}_6(t+M)^{-1})^{\frac{1}{2\gamma}} + (\tilde{J}_6(t+M)^{-1})^{\frac{1}{\gamma}} \right) (t+M) \log^{\frac{1}{2}}(t/M+1), \quad t \geq \tilde{t}_1 \equiv Me.$$

Together with the fact that $F(t) \geq C\varepsilon(t+M) \log(t/M+1)$, this yields

$$\tilde{J}_6 \geq CF(t)^\gamma (t+M)^{1-\gamma} \log^{-\frac{\gamma}{2}}(t/M+1), \quad t \geq \tilde{t}_1.$$

Substituting this into (3-36) yields

$$(3-37) \quad F''(t) + \frac{C_1}{(1+t)^\lambda} F'(t) \geq \frac{C\varepsilon}{t+M}, \quad t \geq 0,$$

$$(3-38) \quad F''(t) + \frac{C_1}{(1+t)^\lambda} F'(t) \geq CF(t)^\gamma (t+M)^{-1-\gamma} \log^{-\frac{\gamma}{2}}(t/M+1), \quad t \geq \tilde{t}_1.$$

Substituting $F(t) \geq C\varepsilon(t+M) \log(t/M+1)$ into (3-38) yields

$$F''(t) + \frac{C_1}{(1+t)^\lambda} F'(t) \geq C\varepsilon^\gamma (t+M)^{-1} \log^{\frac{\gamma}{2}}(t/M+1).$$

Integrating this yields

$$F(t) \geq C\varepsilon^\gamma (t+M) \log^{\frac{\gamma+2}{2}}(t/M+1).$$

Substituting this into (3-38) again gives

$$\begin{aligned} F''(t) + \frac{C_1}{(1+t)^\lambda} F'(t) \\ \geq C\varepsilon^{\gamma^2} (t+M)^{-1} \log^{\frac{\gamma(\gamma+1)}{2}}(t/M+1) = C\varepsilon^{\gamma^2} (t+M)^{-1} \log^{\frac{\gamma(\gamma^2-1)}{2(\gamma-1)}}(t/M+1). \end{aligned}$$

Repeating this process n times, we see that

$$(3-39) \quad F''(t) + \frac{C_1}{(1+t)^\lambda} F'(t) \geq C\varepsilon\gamma^n (t+M)^{-1} \log^{\frac{\gamma(\gamma^n-1)}{2(\gamma-1)}} (t/M+1),$$

where $n = [\log_\gamma 2]$. Solving (3-39) yields

$$F(t) \geq C\varepsilon\gamma^n (t+M) \log^{\frac{\gamma(\gamma^n-1)}{2(\gamma-1)}+1} (t/M+1), \quad t \geq \tilde{t}_2,$$

where $\tilde{t}_2 > 0$ is a constant only depending on γ . Substituting this into (3-38) derives

$$(3-40) \quad F''(t) + \frac{C_1}{(1+t)^\lambda} F'(t) \geq CF(t)\varepsilon\gamma^{n(\gamma-1)}(t+M)^{-2} \log^{\frac{\gamma^{n+1}-2}{2}} (t/M+1), \quad t \geq \tilde{t}_2,$$

where $\frac{1}{2}(\gamma^{n+1} - 2) > 0$ by the choice of $n = [\log_\gamma 2]$. Since (3-40) is analogous to (3-28), as in Case 1, we can choose

$$\tilde{t}_3 = O\left(e^{C\varepsilon \frac{2\gamma^n(\gamma-1)}{\gamma^{n+1}-2}}\right)$$

such that

$$F'(t) \geq C\varepsilon \frac{\gamma^n(\gamma-1)}{2} (t+M)^{-1} \log^{\frac{\gamma^{n+1}-2}{4}} (t/M+1)F(t), \quad t \geq \tilde{t}_3,$$

which is similar to (3-33) and yields

$$(3-41) \quad F(t) \geq C\varepsilon^{C_\gamma} (t+M)^{\frac{2(\gamma+2)}{\gamma-1}}, \quad t \geq \tilde{t}_4 \equiv C\tilde{t}_3^2,$$

where $C_\gamma > 0$ is a constant depending on γ . Substituting (3-41) into (3-38) yields

$$(3-42) \quad F''(t) + \frac{C_1}{(1+t)^\lambda} F'(t) \geq C\varepsilon^{C_\gamma} F(t)^{\frac{\gamma+1}{2}}, \quad t \geq \tilde{t}_4.$$

Multiplying (3-42) by $F'(t)$ and integrating over the variable t as in Case 1, we have

$$F'(t) \geq C\varepsilon^{C_\gamma} F(t)^{\frac{\gamma+3}{4}}, \quad t \geq \tilde{t}_5 \equiv C\tilde{t}_4.$$

Together with $\gamma > 1$ and the choice of \tilde{t}_3 , this yields $T_\varepsilon < \infty$.

Both Case 1 and Case 2 complete the proof of **Theorem 1.2**. □

4. Blowup for large data

In this section, we establish a blowup result for large amplitude smooth solutions of (1-1) which is valid for all $\lambda \geq 0$. More precisely, instead of (1-1) we consider

the Cauchy problem

$$(4-1) \quad \begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u + pI_3) = -\frac{\mu}{(1+t)^\lambda} \rho u, \\ \rho(0, x) = \bar{\rho} + \tilde{\rho}_0(x), \quad u(0, x) = \tilde{u}_0(x), \end{cases}$$

where $\tilde{\rho}_0, \tilde{u}_0 \in C_0^\infty(\mathbb{R}^3)$, $\operatorname{supp} \tilde{\rho}_0, \operatorname{supp} \tilde{u}_0 \subseteq B(0, M) \equiv \{x : |x| \leq M\}$, and $\rho(0, \cdot) > 0$. Motivated by the treatment of the special case of $\lambda = 0$ in [Sideris et al. 2003], we introduce the functions

$$\begin{aligned} H(t) &\equiv \int_{\mathbb{R}^3} x \cdot (\rho u)(t, x) \, dx, & L(t) &\equiv \int_{\mathbb{R}^3} (\rho(t, x) - \bar{\rho}) \, dx, \\ \gamma(t) &\equiv (t + M)^2 \left(L(0) + \frac{4\pi^2 \bar{\rho}}{3} (t + M)^3 \right), \end{aligned}$$

and also remind the reader of the definition of the function β in (1-12).

Then we have the following result:

Theorem 4.1. *Suppose that $L(0) \geq 0$ and*

$$(4-2) \quad H(0) \int_0^{T^*} \frac{d\tau}{\gamma(\tau)\beta(\tau)} > 1.$$

for some $T^* > 0$. Then $T < T^*$ holds for any solution $(\rho, u) \in C^1([0, T] \times \mathbb{R}^3)$ of (4-1).

Proof. From the first equation of (4-1), we see that

$$L'(t) = - \int_{\mathbb{R}^3} \operatorname{div}(\rho u) \, dx = 0,$$

which implies $L(t) = L(0)$. Applying the second equation of (4-1), we find that

$$H'(t) = \int_{\mathbb{R}^3} x \cdot \partial_t(\rho u)(t, x) \, dx = \int_{\mathbb{R}^3} x \cdot \left[-\operatorname{div}(\rho u \otimes u) - \nabla p - \frac{\mu}{(1+t)^\lambda} \rho u \right] \, dx.$$

An integration by parts gives

$$(4-3) \quad H'(t) + \frac{\mu}{(1+t)^\lambda} H(t) = \int_{\mathbb{R}^3} (\rho|u|^2 + 3(p(\rho) - p(\bar{\rho}))) \, dx.$$

Note that the convexity of $p = A\rho^\gamma$ for $\gamma > 1$ and $c(\bar{\rho}) = 1$ imply that

$$(4-4) \quad \int_{\mathbb{R}^3} (p(\rho) - p(\bar{\rho})) \, dx \geq \int_{\mathbb{R}^3} A\gamma \bar{\rho}^{\gamma-1} (\rho - \bar{\rho}) \, dx = L(0).$$

Furthermore, by applying the Cauchy–Schwartz inequality to $H(t)$ and taking into account $\text{supp } u(t, \cdot) \subseteq B(0, M + t)$ for any fixed $t \geq 0$, we have

$$\begin{aligned}
 (4-5) \quad H(t)^2 &\leq \left(\int_{\mathbb{R}^3} \rho |u|^2 dx \right) \left(\int_{|x| \leq t+M} \rho |x|^2 dx \right) \\
 &\leq (t + M)^2 \left(L(0) + \frac{4\pi^2 \bar{\rho}}{3} (t + M)^3 \right) \int_{\mathbb{R}^3} \rho |u|^2 dx \\
 &= \gamma(t) \int_{\mathbb{R}^3} \rho |u|^2 dx.
 \end{aligned}$$

Substituting (4-4)–(4-5) into (4-3) yields

$$(4-6) \quad H'(t) + \frac{\mu}{(1+t)^\lambda} H(t) \geq \frac{H(t)^2}{\gamma(t)} + 3L(0).$$

Together with $L(0) \geq 0$ and $H(0) > 0$ due to (4-2), this shows that $H(t) > 0$ for all $t \in [0, T]$. Denoting $G(t) \equiv \beta(t)H(t)$, from (1-12) and (4-6) we then get that

$$(4-7) \quad G'(t) \geq \frac{G^2(t)}{\gamma(t)\beta(t)}.$$

Now suppose that $T \geq T^*$. Then integrating (4-7) from 0 to T yields

$$\frac{1}{H(0)} - \frac{1}{G(T)} \geq \int_0^T \frac{d\tau}{\gamma(\tau)\beta(\tau)} \geq \int_0^{T^*} \frac{d\tau}{\gamma(\tau)\beta(\tau)},$$

which is a contradiction in view of $G(T) > 0$ and (4-2). □

Appendix: Proof of the nonnegativity of $P(t, l)$ in $\Sigma \equiv \{(t, l) : t \geq 0, t + M_0 \leq l \leq t + M\}$

We fixed a point $A = (t_A, l_A) \in \Sigma$. In the characteristic coordinates $\xi = 1 + t - l$ and $\zeta = 1 + t + l$, (3-7) can be written as

$$(A-1) \quad \mathcal{L}\bar{P} \equiv \partial_{\xi\zeta}^2 \bar{P} + \frac{2^{\lambda-2}\mu}{(\xi + \zeta)^\lambda} (\partial_\xi \bar{P} + \partial_\zeta \bar{P}) = \frac{\bar{f}}{4},$$

where $\bar{P}(\xi, \zeta) \equiv P\left(\frac{\xi+\zeta}{2} - 1, \frac{\zeta-\xi}{2}\right)$. The adjoint operator \mathcal{L}^* of \mathcal{L} has the form

$$(A-2) \quad \mathcal{L}^*\mathcal{R} \equiv \partial_{\xi\zeta}^2 \mathcal{R} - \frac{2^{\lambda-2}\mu}{(\xi + \zeta)^\lambda} (\partial_\xi \mathcal{R} + \partial_\zeta \mathcal{R}) + \frac{2^{\lambda-1}\mu\lambda}{(\xi + \zeta)^{\lambda+1}} \mathcal{R}.$$

For the point $A = (\xi_A, \zeta_A)$ with $\xi_A + \zeta_A = 2(1 + t_A) \geq 2$, write $B = (2 - \zeta_A, \zeta_A)$ and $C = (\xi_A, 2 - \xi_A)$, and let \mathcal{D} the domain surrounded by the triangle ABC (see Figure 1 below).




Figure 1. (ξ, ζ) -plane.

Let the numbers a and b satisfy $a + b = 1$ and $ab = \frac{1}{2}\mu\lambda$. We define

$$(A-3) \quad z \equiv -\frac{(\xi_A - \xi)(\zeta_A - \zeta)}{(\xi_A + \zeta_A)(\xi + \zeta)}$$

and

$$(A-4) \quad \mathcal{R}(\xi, \zeta; \xi_A, \zeta_A) \equiv \left[\frac{\beta(\xi + \zeta - 1)}{\beta(\xi_A + \zeta_A - 1)} \right]^{2\lambda-2} \Psi(a, b, 1; z);$$

here the definition of function β is given in (1-12) and Ψ is the hypergeometric function. From this and direct calculation, we infer

$$(A-5) \quad \mathcal{L}^*\mathcal{R} = \left[\frac{2^{\lambda-2}\mu\lambda}{(\xi + \zeta)^{\lambda+1}} - \frac{\mu\lambda}{2(\xi + \zeta)^2} - \frac{4^{\lambda-2}\mu^2}{(\xi + \zeta)^{2\lambda}} \right] \mathcal{R}.$$

On the other hand, from (A-1)–(A-2) we arrive at

$$\mathcal{R}\mathcal{L}\bar{P} - \bar{P}\mathcal{L}^*\mathcal{R} = \partial_\zeta \left(\mathcal{R}\partial_\xi \bar{P} + \frac{2^{\lambda-2}\mu}{(\xi + \zeta)^\lambda} \mathcal{R}\bar{P} \right) - \partial_\xi \left(\bar{P}\partial_\zeta \mathcal{R} - \frac{2^{\lambda-2}\mu}{(\xi + \zeta)^\lambda} \mathcal{R}\bar{P} \right).$$

Integrating this over \mathcal{D} yields

$$(A-6) \quad \begin{aligned} \bar{P}(A) &= \frac{1}{2}\mathcal{R}(C; A)\bar{P}(C) + \frac{1}{2}\mathcal{R}(B; A)\bar{P}(B) \\ &+ \iint_{\mathcal{D}} (\mathcal{R}\mathcal{L}\bar{P} - \bar{P}\mathcal{L}^*\mathcal{R}) d\xi d\zeta + \int_{BC} \left(\frac{1}{2}\mathcal{R}\partial_\xi \bar{P} - \frac{1}{2}\bar{P}\partial_\xi \mathcal{R} + \frac{\mu}{4}\mathcal{R}\bar{P} \right) d\xi \\ &+ \left(\frac{1}{2}\bar{P}\partial_\zeta \mathcal{R} - \frac{1}{2}\mathcal{R}\partial_\zeta \bar{P} - \frac{\mu}{4}\mathcal{R}\bar{P} \right) d\zeta. \end{aligned}$$




Figure 2. (t, l) -plane.

Returning to the variable (t, l) (see [Figure 2](#)), we find in the second line of [\(A-6\)](#) that

$$\begin{aligned}
 \text{(A-7)} \quad \int_{BC} \dots &= \int_B^C \left[\frac{1}{4} \mathcal{R}(\partial_t - \partial_l) P - \frac{1}{4} P(\partial_t - \partial_l) \mathcal{R} + \frac{\mu}{4} \mathcal{R} P \right] (-dl) \\
 &\quad + \left[\frac{1}{4} P(\partial_t + \partial_l) \mathcal{R} - \frac{1}{4} \mathcal{R}(\partial_t + \partial_l) P - \frac{\mu}{4} \mathcal{R} P \right] dl \\
 &= \int_{l_A - t_A}^{l_A + t_A} \left[\frac{\mu}{2} \mathcal{R} P + \frac{1}{2} \mathcal{R} \partial_t P - \frac{1}{2} P \partial_t \mathcal{R} \right] \Big|_{t=0} dl \\
 &= \int_{l_A - t_A}^{l_A + t_A} \beta(t_A)^{-\frac{1}{2}} \left[\Psi(a, b, 1; z|_{t=0}) \left(\frac{\mu}{4} q_0(l) + \frac{1}{2} q_1(l) \right) \right. \\
 &\quad \left. - \frac{\mu \lambda}{4} \Psi(a + 1, b + 1, 2; z|_{t=0}) q_0(l) z_t|_{t=0} \right] dl,
 \end{aligned}$$

where we have used the formula $\Psi'(a, b, c; z) = \frac{ab}{c} \Psi(a + 1, b + 1, c + 1; z)$ (see [\[Erdélyi et al. 1953, page 58\]](#)). From the definition [\(A-3\)](#), we arrive at

$$z = - \frac{(t_A - l_A - t + l)(t_A + l_A - t - l)}{4(1 + t_A)(1 + t)}$$

and

$$\text{(A-8)} \quad z_t|_{t=0} = \frac{t_A}{2(1 + t_A)} - z|_{t=0}.$$

If $(t, l) \in \Sigma \cap \bar{\mathcal{D}}$, we infer

$$\text{(A-9)} \quad 0 \geq z \geq -\frac{1}{2}(M - M_0) \geq -\frac{1}{2}\delta_0,$$

which implies that [\(1-8\)](#) holds. This, together with [\(A-7\)](#)–[\(A-9\)](#) and the assumption [\(1-11\)](#) of $\Lambda \geq \frac{3}{2}\mu\lambda$, yields that the integral in the second line of [\(A-6\)](#) is nonnegative.

Next we prove that $P(t, l) \geq 0$ for all $(t, l) \in \Sigma$. Define

$$\bar{t} \equiv \inf\{t : \exists l \in (t + M_0, t + M) \text{ such that } P(t, l) < 0\}.$$

From assumption (1-9), we get $\bar{t} > 0$. If $\bar{t} < +\infty$, we see that there exists $\bar{l} \in (\bar{t} + M_0, \bar{t} + M)$ such that $P(\bar{t}, \bar{l}) = 0$. Moreover, we have $P(t, l) \geq 0$ for $t < \bar{t}$. Choose $A = (t_A, l_A) = (\bar{t}, \bar{l})$ in (A-6). From (A-4)–(A-5) and (1-8) we infer $\mathcal{L}^*\mathcal{R} \leq 0$ for $\lambda > 1$ and $(t, l) \in \Sigma \cap \mathcal{D}$. It follows from $f(t, l) \geq 0$ in (3-7), (1-8), (1-9), and (A-6) that

$$P(\bar{t}, \bar{l}) \geq \frac{1}{2}\mathcal{R}(C; A)P(0, \bar{l} - \bar{t}) + \iint_{\Sigma \cap \mathcal{D}} (\mathcal{R}\mathcal{L}\bar{P} - \bar{P}\mathcal{L}^*\mathcal{R}) d\xi d\zeta \geq \frac{1}{4}q_0(\bar{l} - \bar{t}) > 0,$$

which is a contradiction with $P(\bar{t}, \bar{l}) = 0$. Consequently, we conclude that $\bar{t} = +\infty$ and $P(t, l) \geq 0$ for all $(t, l) \in \Sigma$.

Acknowledgment

Yin Huicheng wishes to express his gratitude to Professor Michael Reissig, Technical University Bergakademie Freiberg, Germany, for his interest in this problem and some fruitful discussions in the past.

References

- [Alinhac 1993] S. Alinhac, “Temps de vie des solutions régulières des équations d’Euler compressibles axisymétriques en dimension deux”, *Invent. Math.* **111**:3 (1993), 627–670. [MR](#) [Zbl](#)
- [Alinhac 1999a] S. Alinhac, “Blowup of small data solutions for a quasilinear wave equation in two space dimensions”, *Ann. of Math.* (2) **149**:1 (1999), 97–127. [MR](#) [Zbl](#)
- [Alinhac 1999b] S. Alinhac, “Blowup of small data solutions for a class of quasilinear wave equations in two space dimensions, II”, *Acta Math.* **182**:1 (1999), 1–23. [MR](#) [Zbl](#)
- [Christodoulou 2007] D. Christodoulou, *The formation of shocks in 3-dimensional fluids*, European Mathematical Society, Zürich, 2007. [MR](#) [Zbl](#)
- [Christodoulou and Lisibach 2016] D. Christodoulou and A. Lisibach, “Shock development in spherical symmetry”, *Ann. PDE* **2**:1 (2016), art. id. 3. [MR](#)
- [Christodoulou and Miao 2014] D. Christodoulou and S. Miao, *Compressible flow and Euler’s equations*, Surveys of Modern Mathematics **9**, International Press, Somerville, MA, 2014. [MR](#) [Zbl](#)
- [Courant and Hilbert 1962] R. Courant and D. Hilbert, *Methods of mathematical physics, II: Partial differential equations*, Interscience, New York, 1962. [MR](#) [Zbl](#)
- [D’Abbico and Reissig 2014] M. D’Abbico and M. Reissig, “Semilinear structural damped waves”, *Math. Methods Appl. Sci.* **37**:11 (2014), 1570–1592. [MR](#) [Zbl](#)
- [D’Abbico et al. 2015] M. D’Abbico, S. Lucente, and M. Reissig, “A shift in the Strauss exponent for semilinear wave equations with a not effective damping”, *J. Differential Equations* **259**:10 (2015), 5040–5073. [MR](#) [Zbl](#)
- [Dafermos 1995] C. M. Dafermos, “A system of hyperbolic conservation laws with frictional damping”, *Z. Angew. Math. Phys.* **46**:special issue (1995), 294–307. [MR](#) [Zbl](#)

- [Ding et al. 2016] B. Ding, I. Witt, and H. Yin, “The small data solutions of general 3-D quasilinear wave equations, II”, *J. Differential Equations* **261**:2 (2016), 1429–1471. [MR](#) [Zbl](#)
- [Erdélyi et al. 1953] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher transcendental functions, I*, McGraw-Hill, New York, 1953. [MR](#) [Zbl](#)
- [Hörmander 1997] L. Hörmander, *Lectures on nonlinear hyperbolic differential equations*, Math. Appl. **26**, Springer, Berlin, 1997. [MR](#) [Zbl](#)
- [Hsiao and Liu 1992] L. Hsiao and T.-P. Liu, “Convergence to nonlinear diffusion waves for solutions of a system of hyperbolic conservation laws with damping”, *Comm. Math. Phys.* **143**:3 (1992), 599–605. [MR](#) [Zbl](#)
- [Hsiao and Serre 1996] L. Hsiao and D. Serre, “Global existence of solutions for the system of compressible adiabatic flow through porous media”, *SIAM J. Math. Anal.* **27**:1 (1996), 70–77. [MR](#) [Zbl](#)
- [Kawashima and Yong 2004] S. Kawashima and W.-A. Yong, “Dissipative structure and entropy for hyperbolic systems of balance laws”, *Arch. Ration. Mech. Anal.* **174**:3 (2004), 345–364. [MR](#) [Zbl](#)
- [Klainerman 1987] S. Klainerman, “Remarks on the global Sobolev inequalities in the Minkowski space \mathbb{R}^{n+1} ”, *Comm. Pure Appl. Math.* **40**:1 (1987), 111–117. [MR](#) [Zbl](#)
- [Klainerman and Sideris 1996] S. Klainerman and T. C. Sideris, “On almost global existence for nonrelativistic wave equations in 3D”, *Comm. Pure Appl. Math.* **49**:3 (1996), 307–321. [MR](#) [Zbl](#)
- [do Nascimento and Wirth 2015] W. N. do Nascimento and J. Wirth, “Wave equations with mass and dissipation”, *Adv. Differential Equations* **20**:7-8 (2015), 661–696. [MR](#) [Zbl](#)
- [Nishihara 1997] K. Nishihara, “Asymptotic behavior of solutions of quasilinear hyperbolic equations with linear damping”, *J. Differential Equations* **137**:2 (1997), 384–395. [MR](#) [Zbl](#)
- [Pan and Zhao 2009] R. Pan and K. Zhao, “The 3D compressible Euler equations with damping in a bounded domain”, *J. Differential Equations* **246**:2 (2009), 581–596. [MR](#) [Zbl](#)
- [Sideris 1985] T. C. Sideris, “Formation of singularities in three-dimensional compressible fluids”, *Comm. Math. Phys.* **101**:4 (1985), 475–485. [MR](#) [Zbl](#)
- [Sideris 1997] T. C. Sideris, “Delayed singularity formation in 2D compressible flow”, *Amer. J. Math.* **119**:2 (1997), 371–422. [MR](#) [Zbl](#)
- [Sideris et al. 2003] T. C. Sideris, B. Thomases, and D. Wang, “Long time behavior of solutions to the 3D compressible Euler equations with damping”, *Comm. Partial Differential Equations* **28**:3-4 (2003), 795–816. [MR](#) [Zbl](#)
- [Speck 2016] J. Speck, *Shock formation in small-data solutions to 3D quasilinear wave equations*, Mathematical Surveys and Monographs **214**, American Mathematical Society, Providence, RI, 2016. [MR](#) [Zbl](#)
- [Tan and Guochun 2012] Z. Tan and W. Guochun, “Large time behavior of solutions for compressible Euler equations with damping in \mathbb{R}^3 ”, *J. Differential Equations* **252**:2 (2012), 1546–1561. [MR](#) [Zbl](#)
- [Wang and Yang 2001] W. Wang and T. Yang, “The pointwise estimates of solutions for Euler equations with damping in multi-dimensions”, *J. Differential Equations* **173**:2 (2001), 410–450. [MR](#) [Zbl](#)
- [Wirth 2006] J. Wirth, “Wave equations with time-dependent dissipation, I: Non-effective dissipation”, *J. Differential Equations* **222**:2 (2006), 487–514. [MR](#) [Zbl](#)
- [Wirth 2007] J. Wirth, “Wave equations with time-dependent dissipation, II: Effective dissipation”, *J. Differential Equations* **232**:1 (2007), 74–103. [MR](#) [Zbl](#)
- [Yin 2004] H. Yin, “Formation and construction of a shock wave for 3-D compressible Euler equations with the spherical initial data”, *Nagoya Math. J.* **175** (2004), 125–164. [MR](#) [Zbl](#)

[Yin and Qiu 1999] H. Yin and Q. Qiu, “The blowup of solutions for 3-D axisymmetric compressible Euler equations”, *Nagoya Math. J.* **154** (1999), 157–169. [MR](#) [Zbl](#)

Received December 31, 2016.

FEI HOU
DEPARTMENT OF MATHEMATICS AND IMS
NANJING UNIVERSITY
NANJING
CHINA
houfeimath@gmail.com

INGO WITT
MATHEMATICAL INSTITUTE
UNIVERSITY OF GÖTTINGEN
GÖTTINGEN
GERMANY
iwitt@uni-math.gwdg.de

HUICHENG YIN
SCHOOL OF MATHEMATICAL SCIENCES
JIANGSU PROVINCIAL KEY LABORATORY FOR NUMERICAL SIMULATION OF LARGE SCALE
COMPLEX SYSTEMS
NANJING NORMAL UNIVERSITY
NANJING
CHINA
huicheng@nju.edu.cn
05407@njnu.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 292 No. 2 February 2018

Locally helical surfaces have bounded twisting	257
DAVID BACHMAN, RYAN DERBY-TALBOT and ERIC SEDGWICK	
Superconvergence to freely infinitely divisible distributions	273
HARI BERCOVICI, JIUN-CHAU WANG and PING ZHONG	
Norm constants in cases of the Caffarelli–Kohn–Nirenberg inequality	293
AKSHAY L. CHANILLO, SAGUN CHANILLO and ALI MAALAOU	
Noncommutative geometry of homogenized quantum $\mathfrak{sl}(2, \mathbb{C})$	305
ALEX CHIRVASITU, S. PAUL SMITH and LIANG ZE WONG	
A generalization of “Existence and behavior of the radial limits of a bounded capillary surface at a corner”	355
JULIE N. CRENSHAW, ALEXANDRA K. ECHART and KIRK E. LANCASTER	
Norms in central simple algebras	373
DANIEL GOLDSTEIN and MURRAY SCHACHER	
Global existence and blowup of smooth solutions of 3-D potential equations with time-dependent damping	389
FEI HOU, INGO WITT and HUICHENG YIN	
Formal confluence of quantum differential operators	427
BERNARD LE STUM and ADOLFO QUIRÓS	
Rigidity of Hawking mass for surfaces in three manifolds	479
JIACHENG SUN	
Addendum to “A strong multiplicity one theorem for SL_2 ”	505
QING ZHANG	