

Pacific Journal of Mathematics

**CONSTRUCTION OF A RAPOPORT-ZINK SPACE FOR $GU(1, 1)$
IN THE RAMIFIED 2-ADIC CASE**

DANIEL KIRCH

CONSTRUCTION OF A RAPOPORT–ZINK SPACE FOR $\mathrm{GU}(1, 1)$ IN THE RAMIFIED 2-ADIC CASE

DANIEL KIRCH

Let $F|\mathbb{Q}_2$ be a finite extension. In this paper, we construct an RZ-space \mathcal{N}_E for split $\mathrm{GU}(1, 1)$ over a ramified quadratic extension $E|F$. For this, we first introduce the naive moduli problem $\mathcal{N}_E^{\mathrm{naive}}$ and then define $\mathcal{N}_E \subseteq \mathcal{N}_E^{\mathrm{naive}}$ as a canonical closed formal subscheme, using the so-called straightening condition. We establish an isomorphism between \mathcal{N}_E and the Drinfeld moduli problem, proving the 2-adic analogue of a theorem of Kudla and Rapoport. The formulation of the straightening condition uses the existence of certain polarizations on the points of the moduli space $\mathcal{N}_E^{\mathrm{naive}}$. We show the existence of these polarizations in a more general setting over any quadratic extension $E|F$, where $F|\mathbb{Q}_p$ is a finite extension for any prime p .

1. Introduction	341
2. Preliminaries on quaternion algebras and hermitian forms	347
3. The moduli problem in the case R-P	353
4. The moduli problem in the case R-U	368
5. A theorem on the existence of polarizations	382
Acknowledgements	388
References	388

1. Introduction

Rapoport–Zink spaces (RZ-spaces for short) are moduli spaces of p -divisible groups endowed with additional structure. Rapoport and Zink [1996] studied two major classes of RZ-spaces, called EL type and PEL type. The abbreviations EL and PEL indicate, in analogy to the case of Shimura varieties, whether the extra structure comes in the form of Endomorphisms and Level structure or in the form of Polarizations, Endomorphisms and Level structure. Rapoport and Zink [1996] developed a theory of these spaces, including important theorems about the existence of local models and nonarchimedean uniformization of Shimura varieties, for the EL type and for the PEL type whenever $p \neq 2$.

MSC2010: 11G18, 14G35.

Keywords: Rapoport–Zink spaces, Shimura varieties, bad reduction, unitary group, exceptional isomorphism.

The blanket assumption $p \neq 2$ made by Rapoport and Zink in the PEL case is by no means of a cosmetic nature, but originates with various serious difficulties that arise for $p = 2$. However, we recall that one can still use their definition in that case to obtain “naïve” moduli spaces that still satisfy basic properties like being representable by a formal scheme.

In this paper, we construct the 2-adic Rapoport–Zink space \mathcal{N}_E corresponding to the group of unitary similitudes of size 2 relative to any (wildly) ramified quadratic extension $E|F$, where $F|\mathbb{Q}_2$ is a finite extension. It is given as the closed formal subscheme of the corresponding naïve RZ-space $\mathcal{N}_E^{\text{naïve}}$ described by the so-called “straightening condition”, which is defined below. The main result of this paper is a natural isomorphism $\eta : \mathcal{M}_{Dr} \xrightarrow{\sim} \mathcal{N}_E$, where \mathcal{M}_{Dr} is Deligne’s formal model of the Drinfeld upper half-plane (cf. [Boutot and Carayol 1991]). This result is in analogy with Kudla and Rapoport’s construction [2014] of a corresponding isomorphism for $p \neq 2$ and also for $p = 2$ when $E|F$ is an unramified extension. The formal scheme \mathcal{M}_{Dr} solves a certain moduli problem of p -divisible groups and, in this way, it carries the structure of an RZ-space of EL type. In particular, \mathcal{M}_{Dr} is defined even for $p = 2$.

As in [Kudla and Rapoport 2014], there are natural group actions by $\mathrm{SL}_2(F)$ and the split $\mathrm{SU}_2(F)$ on the spaces \mathcal{M}_{Dr} and \mathcal{N}_E , respectively. The isomorphism η is hence a geometric realization of the exceptional isomorphism of these groups. As a consequence, one cannot expect a similar result in higher dimensions. Of course, the existence of “good” RZ-spaces is still expected, but a general definition will probably need a different approach.

The study of residue characteristic 2 is interesting and important for the following reasons: First of all, from the general philosophy of RZ-spaces and, more generally, of local Shimura varieties [Rapoport and Viehmann 2014], it follows that there should be a uniform approach for all primes p . In this sense, the present paper is in the same spirit as the recent constructions of RZ-spaces of Hodge type of W. Kim [2013], Howard and Pappas [2017] and Bültel and Pappas [2017]. Second, Rapoport–Zink spaces have been used to determine the arithmetic intersection numbers of special cycles on Shimura varieties [Kudla et al. 2006]; in this kind of problem, it is necessary to deal with all places, even those of residue characteristic 2. Finally, studying the cases of residue characteristic 2 also throws light on the cases previously known. In the specific case at hand, the methods we develop also give a simplification of the proof for $p \neq 2$ of Kudla and Rapoport [2014]; see Remark 5.3 (2).

We will now explain the results of this paper in greater detail. Let F be a finite extension of \mathbb{Q}_2 and $E|F$ a ramified quadratic extension. Following [Jacobowitz 1962], we consider the following dichotomy for this extension (see Section 2):

(R-P) There is a uniformizer $\pi_0 \in F$ such that $E = F[\Pi]$ with $\Pi^2 + \pi_0 = 0$. Then the rings of integers O_F of F and O_E of E satisfy $O_E = O_F[\Pi]$.

(R-U) $E|F$ is given by an Eisenstein equation of the form $\Pi^2 - t\Pi + \pi_0 = 0$. Here, π_0 is again a uniformizer in F and $t \in \mathcal{O}_F$ satisfies $\pi_0|t|2$. We still have $\mathcal{O}_E = \mathcal{O}_F[\Pi]$. Note that in this case $E|F$ is generated by a square root of the unit $1 - 4\pi_0/t^2$ in F .

An example of an extension of type R-P is $\mathbb{Q}_2(\sqrt{-2})|\mathbb{Q}_2$, whereas $\mathbb{Q}_2(\sqrt{-1})|\mathbb{Q}_2$ is of type R-U. Note that for $p > 2$, any ramified quadratic extension over \mathbb{Q}_p is of the form R-P.

Our results in the cases R-P and R-U are similar, but different. We first describe the results in the case R-P. Let $E|F$ be of type R-P.

We first define a naive moduli problem $\mathcal{N}_E^{\text{naive}}$, which merely copies the definition from $p \neq 2$ (cf. [Kudla and Rapoport 2014]). Let \check{F} be the completion of the maximal unramified extension of F and $\check{\mathcal{O}}_F$ its ring of integers. Then $\mathcal{N}_E^{\text{naive}}$ is a set-valued functor on $\mathrm{Nilp}_{\check{\mathcal{O}}_F}$, the category of $\check{\mathcal{O}}_F$ -schemes where π_0 is locally nilpotent. For $S \in \mathrm{Nilp}_{\check{\mathcal{O}}_F}$, the set $\mathcal{N}_E^{\text{naive}}(S)$ is the set of equivalence classes of tuples $(X, \iota, \lambda, \varrho)$. Here, X/S is a formal \mathcal{O}_F -module of height 4 and dimension 2, equipped with an action $\iota : \mathcal{O}_E \rightarrow \mathrm{End}(X)$. This action satisfies the Kottwitz condition of signature $(1, 1)$, i.e., for any $\alpha \in \mathcal{O}_E$, the characteristic polynomial of $\iota(\alpha)$ on $\mathrm{Lie} X$ is given by

$$\mathrm{char}(\mathrm{Lie} X, T \mid \iota(\alpha)) = (T - \alpha)(T - \bar{\alpha}).$$

Here, $\alpha \mapsto \bar{\alpha}$ denotes the Galois conjugation of $E|F$. The right side of this equation is a polynomial with coefficients in \mathcal{O}_S via the structure map $\mathcal{O}_F \hookrightarrow \check{\mathcal{O}}_F \rightarrow \mathcal{O}_S$. The third entry λ is a principal polarization $\lambda : X \rightarrow X^\vee$ such that the induced Rosati involution satisfies $\iota(\alpha)^* = \iota(\bar{\alpha})$ for all $\alpha \in \mathcal{O}_E$. (Here, X^\vee is the dual of X as a formal \mathcal{O}_F -module.) Finally, ϱ is a quasi-isogeny of height 0 (and compatible with all previous data) to a fixed framing object $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ over $\bar{k} = \check{\mathcal{O}}_F/\pi_0$. This framing object is unique up to isogeny under the condition that

$$\{\varphi \in \mathrm{End}^0(\mathbb{X}, \iota_{\mathbb{X}}) \mid \varphi^*(\lambda_{\mathbb{X}}) = \lambda_{\mathbb{X}}\} \simeq \mathrm{U}(C, h),$$

for a split $E|F$ -hermitian vector space (C, h) of dimension 2; see Proposition 3.2.

Recall that this is exactly the definition used in [Kudla and Rapoport 2014] for the ramified case with $p > 2$. There, $\mathcal{N}_E = \mathcal{N}_E^{\text{naive}}$ and we have natural isomorphism

$$\eta : \mathcal{M}_{Dr} \xrightarrow{\sim} \mathcal{N}_E,$$

where \mathcal{M}_{Dr} is the Drinfeld moduli problem mentioned above.

However, for $p = 2$, it turns out that the definition of $\mathcal{N}_E^{\text{naive}}$ is not the “correct” one in the sense that it is not isomorphic to the Drinfeld moduli problem. Hence this naive definition of the moduli space is not in line with the results from [Kudla and Rapoport 2014] and the general philosophy of (conjectural) local Shimura varieties (see [Rapoport and Viehmann 2014]). In order to remedy this, we will describe a

new condition on $\mathcal{N}_E^{\text{naive}}$, which we call the *straightening condition*, and show that this cuts out a closed formal subscheme $\mathcal{N}_E \subseteq \mathcal{N}_E^{\text{naive}}$ that is naturally isomorphic to \mathcal{M}_{Dr} . Interestingly, the straightening condition is not trivial on the rigid-analytic generic fiber of $\mathcal{N}_E^{\text{naive}}$ (as originally assumed by the author), but it cuts out an (admissible) open and closed subspace; see [Remark 3.13](#).

We would like to explicate the defect of the naive moduli space. For this, let us recall the definition of \mathcal{M}_{Dr} . It is a functor on $\text{Nilp}_{\check{O}_F}$, mapping a scheme S to the set $\mathcal{M}_{Dr}(S)$ of equivalence classes of tuples (X, ι_B, ϱ) . Again, X/S is a formal O_F -module of height 4 and dimension 2. Let B be the quaternion division algebra over F and O_B its ring of integers. Then ι_B is an action of O_B on X , satisfying the *special* condition of Drinfeld (see [\[Boutot and Carayol 1991\]](#) or [Section 3C](#) below). The last entry ϱ is an O_B -linear quasi-isogeny of height 0 to a fixed framing object $(\mathbb{X}, \iota_{\mathbb{X}, B})$ over \bar{k} . This framing object is unique up to isogeny (cf. [\[Boutot and Carayol 1991, II. Proposition 5.2\]](#)).

Fix an embedding $O_E \hookrightarrow O_B$ and consider the involution $b \mapsto b^* = \Pi b' \Pi^{-1}$ on B , where $b \mapsto b'$ is the standard involution. By work of Drinfeld (see [Proposition 3.14](#) below), there exists a principal polarization $\lambda_{\mathbb{X}}$ on the framing object $(\mathbb{X}, \iota_{\mathbb{X}, B})$ of \mathcal{M}_{Dr} such that the induced Rosati involution satisfies $\iota_{\mathbb{X}, B}(b)^* = \iota_{\mathbb{X}, B}(b^*)$ for all $b \in O_B$. This polarization is unique up to a scalar in O_F^\times . Furthermore, for any $(X, \iota_B, \varrho) \in \mathcal{M}_{Dr}(S)$, the pullback $\lambda = \varrho^*(\lambda_{\mathbb{X}})$ is a principal polarization on X .

We now set

$$\eta(X, \iota_B, \varrho) = (X, \iota_B|_{O_E}, \lambda, \varrho).$$

By [Lemma 3.15](#), this defines a closed embedding $\eta : \mathcal{M}_{Dr} \hookrightarrow \mathcal{N}_E^{\text{naive}}$. But η is far from being an isomorphism, as the following proposition shows:

Proposition 1.1. *The induced map $\eta(\bar{k}) : \mathcal{M}_{Dr}(\bar{k}) \rightarrow \mathcal{N}_E^{\text{naive}}(\bar{k})$ is not surjective.*

Let us sketch the proof here. Using Dieudonné theory, we can write $\mathcal{N}_E^{\text{naive}}(\bar{k})$ naturally as a union

$$\mathcal{N}_E^{\text{naive}}(\bar{k}) = \bigcup_{\Lambda \subseteq C} \mathbb{P}(\Lambda / \Pi \Lambda)(\bar{k}),$$

where the union runs over all O_E -lattices Λ in the hermitian vector space (C, h) that are Π^{-1} -modular, i.e., the dual Λ^\sharp of Λ with respect to h is given by $\Lambda = \Pi^{-1} \Lambda^\sharp$ (see [Lemma 3.7](#)). By [\[Jacobowitz 1962\]](#), there exist different types (i.e., $U(C, h)$ -orbits) of such lattices $\Lambda \subseteq C$ that are parametrized by their norm ideal $\text{Nm}(\Lambda) = \langle \{h(x, x) | x \in \Lambda\} \rangle \subseteq F$. In the case at hand, $\text{Nm}(\Lambda)$ can be any ideal with $2O_F \subseteq \text{Nm}(\Lambda) \subseteq O_F$. It is easily checked (see [Section 2](#)) that the norm ideal of Λ is minimal, that is $\text{Nm}(\Lambda) = 2O_F$, if and only if Λ admits a basis consisting of isotropic vectors, and hence we call these lattices *hyperbolic*. Now, the image under η of $\mathcal{M}_{Dr}(\bar{k})$ is the union of all lines $\mathbb{P}(\Lambda / \Pi \Lambda)(\bar{k})$ where $\Lambda \subseteq C$ is hyperbolic. This is a consequence of [Remark 3.12](#) and [Theorem 3.16](#) below.

On the framing object $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ of $\mathcal{N}_E^{\mathrm{naive}}$, there exists a principal polarization $\tilde{\lambda}_{\mathbb{X}}$ such that the induced Rosati involution is the identity on \mathcal{O}_E . This polarization is unique up to a scalar in \mathcal{O}_E^{\times} (see [Theorem 5.2 \(1\)](#)). On C , the polarization $\tilde{\lambda}_{\mathbb{X}}$ induces an E -linear alternating form b , such that $\det b$ and $\det h$ differ only by a unit (for a fixed basis of C). After possibly rescaling b by a unit in \mathcal{O}_E^{\times} , a Π^{-1} -modular lattice $\Lambda \subseteq C$ is hyperbolic if and only if $b(x, y) + h(x, y) \in 2\mathcal{O}_F$ for all $x, y \in \Lambda$. This enables us to describe the “hyperbolic” points of $\mathcal{N}_E^{\mathrm{naive}}$ (i.e., those that lie on a projective line corresponding to a hyperbolic lattice $\Lambda \subseteq C$) in terms of polarizations.

We now formulate the closed condition that characterizes \mathcal{N}_E as a closed formal subscheme of $\mathcal{N}_E^{\mathrm{naive}}$. For a suitable choice of $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ and $\tilde{\lambda}_{\mathbb{X}}$, we may assume that $\frac{1}{2}(\lambda_{\mathbb{X}} + \tilde{\lambda}_{\mathbb{X}})$ is a polarization on \mathbb{X} . The following definition is a reformulation of [Definition 3.11](#).

Definition 1.2. Let $S \in \mathrm{Nilp}_{\check{\mathcal{O}}_F}$. An object $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\mathrm{naive}}(S)$ satisfies the *straightening* condition if $\lambda_1 = \frac{1}{2}(\lambda + \tilde{\lambda})$ is a polarization on X . Here, $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$.

We remark that $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$ is a polarization on X . This is a consequence of [Theorem 5.2](#), which states the existence of certain polarizations on points of a larger moduli space \mathcal{M}_E containing $\mathcal{N}_E^{\mathrm{naive}}$; see below.

For $S \in \mathrm{Nilp}_{\check{\mathcal{O}}_F}$, let $\mathcal{N}_E(S) \subseteq \mathcal{N}_E^{\mathrm{naive}}(S)$ be the subset of all tuples $(X, \iota, \lambda, \varrho)$ that satisfy the straightening condition. By [\[Rapoport and Zink 1996, Proposition 2.9\]](#), this defines a closed formal subscheme $\mathcal{N}_E \subseteq \mathcal{N}_E^{\mathrm{naive}}$. An application of Drinfeld’s proposition ([Proposition 3.14](#); see also [\[Boutot and Carayol 1991\]](#)) shows that the image of \mathcal{M}_{Dr} under η lies in \mathcal{N}_E . The main theorem in the R-P case can now be stated as follows; see [Theorem 3.16](#).

Theorem 1.3. $\eta : \mathcal{M}_{Dr} \rightarrow \mathcal{N}_E$ is an isomorphism of formal schemes.

This concludes our discussion of the R-P case. From now on, we assume that $E|F$ is of type R-U.

In the case R-U, we have to make some adaptations for $\mathcal{N}_E^{\mathrm{naive}}$. For $S \in \mathrm{Nilp}_{\check{\mathcal{O}}_F}$, let $\mathcal{N}_E^{\mathrm{naive}}(S)$ be the set of equivalence classes of tuples $(X, \iota, \lambda, \varrho)$ with (X, ι) as in the R-P case. But now, the polarization $\lambda : X \rightarrow X^{\vee}$ is supposed to have kernel $\ker \lambda = X[\Pi]$ (in contrast to the R-P case, where λ is a principal polarization). As before, the Rosati involution of λ induces the conjugation on \mathcal{O}_E . There exists a framing object $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ over $\mathrm{Spec} \bar{k}$ for $\mathcal{N}_E^{\mathrm{naive}}$, which is unique up to isogeny under the condition that

$$\{\varphi \in \mathrm{End}^0(\mathbb{X}, \iota_{\mathbb{X}}) \mid \varphi^*(\lambda_{\mathbb{X}}) = \lambda_{\mathbb{X}}\} \simeq \mathrm{U}(C, h),$$

where (C, h) is a split $E|F$ -hermitian vector space of dimension 2 (see [Proposition 4.1](#)). Finally, ϱ is a quasi-isogeny of height 0 from X to \mathbb{X} , respecting all structure.

Fix an embedding $E \hookrightarrow B$. Using some subtle choices of elements in B (these are described in [Lemma 2.3 \(2\)](#)) and by Drinfeld's proposition, we can construct a polarization λ as above for any $(X, \iota_B, \varrho) \in \mathcal{M}_{Dr}(S)$. This induces a closed embedding

$$\eta : \mathcal{M}_{Dr} \rightarrow \mathcal{N}_E^{\text{naive}}, \quad (X, \iota_B, \varrho) \mapsto (X, \iota_B|_{O_E}, \lambda, \varrho).$$

We can write $\mathcal{N}_E^{\text{naive}}(\bar{k})$ as a union of projective lines,

$$\mathcal{N}_E^{\text{naive}}(\bar{k}) = \bigcup_{\Lambda \subseteq C} \mathbb{P}(\Lambda / \Pi \Lambda)(\bar{k}),$$

where the union now runs over all self-dual O_E -lattices $\Lambda \subseteq (C, h)$ with $\text{Nm}(\Lambda) \subseteq \pi_0 O_F$. As in the R-P case, these lattices $\Lambda \subseteq C$ are classified up to isomorphism by their norm ideal $\text{Nm}(\Lambda)$. Since Λ is self-dual with respect to h , the norm ideal can be any ideal satisfying $t O_F \subseteq \text{Nm}(\Lambda) \subseteq O_F$. We call Λ *hyperbolic* when the norm ideal is minimal, i.e., $\text{Nm}(\Lambda) = t O_F$. Equivalently, the lattice Λ has a basis consisting of isotropic vectors. Recall that here t is the element showing up in the Eisenstein equation for the R-U extension $E|F$ and that $\pi_0|t|2$. Hence there exists at least one type of self-dual lattices $\Lambda \subseteq C$ with $\text{Nm}(\Lambda) \subseteq \pi_0 O_F$. In the case R-U, it may happen that $|t| = |\pi_0|$, in which case all lattices Λ in the description of $\mathcal{N}_E^{\text{naive}}(\bar{k})$ are hyperbolic.

The image of $\mathcal{M}_{Dr}(\bar{k})$ under η in $\mathcal{N}_E^{\text{naive}}(\bar{k})$ is the union of all projective lines corresponding to hyperbolic lattices. Unless $|t| = |\pi_0|$, it follows that $\eta(\bar{k})$ is not surjective and thus η cannot be an isomorphism. For the case $|t| = |\pi_0|$, we will show that η is an isomorphism on reduced loci $(\mathcal{M}_{Dr})_{\text{red}} \xrightarrow{\sim} (\mathcal{N}_E^{\text{naive}})_{\text{red}}$ (see [Remark 4.11](#)), but η is not an isomorphism of formal schemes. This follows from the nonflatness of the deformation ring for certain points of $\mathcal{N}_E^{\text{naive}}$; see [Section 4D](#).

On the framing object $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ of $\mathcal{N}_E^{\text{naive}}$, there exists a polarization $\tilde{\lambda}_{\mathbb{X}}$ such that $\ker \tilde{\lambda}_{\mathbb{X}} = \mathbb{X}[\Pi]$ and such that the Rosati involution induces the identity on O_E . After a suitable choice of $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ and $\tilde{\lambda}_{\mathbb{X}}$, we may assume that $\frac{1}{t}(\lambda_{\mathbb{X}} + \tilde{\lambda}_{\mathbb{X}})$ is a polarization on \mathbb{X} . The straightening condition for the R-U case is given as follows (see [Definition 4.10](#)).

Definition 1.4. Let $S \in \text{Nilp}_{\check{O}_F}$. An object $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\text{naive}}(S)$ satisfies the *straightening* condition if $\lambda_1 = \frac{1}{t}(\lambda + \tilde{\lambda})$ is a polarization on X . Here, $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$.

Note that $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$ is a polarization on X by [Theorem 5.2](#).

The straightening condition defines a closed formal subscheme $\mathcal{N}_E \subseteq \mathcal{N}_E^{\text{naive}}$ that contains the image of \mathcal{M}_{Dr} under η . The main theorem in the R-U case can now be stated as follows; compare [Theorem 4.14](#).

Theorem 1.5. $\eta : \mathcal{M}_{Dr} \rightarrow \mathcal{N}_E$ is an isomorphism of formal schemes.

When formulating the straightening condition in the R-U and the R-P case, we mentioned that $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$ is a polarization for any $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\mathrm{naive}}(S)$. This fact is a corollary of [Theorem 5.2](#), which states the existence of this polarization in the following more general setting.

Let $F|\mathbb{Q}_p$ be a finite extension for any prime p and $E|F$ an arbitrary quadratic extension. We consider the following moduli space \mathcal{M}_E of EL type. For $S \in \mathrm{Nilp}_{\check{O}_F}$, the set $\mathcal{M}_E(S)$ consists of equivalence classes of tuples (X, ι_E, ϱ) , where X is a formal O_F -module of height 4 and dimension 2 and ι_E is an O_E -action on X satisfying the Kottwitz condition of signature $(1, 1)$ as above. The entry ϱ is an O_E -linear quasi-isogeny of height 0 to a supersingular framing object $(\mathbb{X}, \iota_{\mathbb{X}, E})$.

The points of \mathcal{M}_E are equipped with polarizations in the following natural way; see [Theorem 5.2](#).

- Theorem 1.6.** (1) *There exists a principal polarization $\tilde{\lambda}_{\mathbb{X}}$ on $(\mathbb{X}, \iota_{\mathbb{X}, E})$ such that the Rosati involution induces the identity on O_E , i.e., $\iota(\alpha)^* = \iota(\alpha)$ for all $\alpha \in O_E$. This polarization is unique up to a scalar in O_E^\times .*
- (2) *Fix $\tilde{\lambda}_{\mathbb{X}}$ as in part (1). For any $S \in \mathrm{Nilp}_{\check{O}_F}$ and $(X, \iota_E, \varrho) \in \mathcal{M}_E(S)$, there exists a unique principal polarization $\tilde{\lambda}$ on X such that the Rosati involution induces the identity on O_E and such that $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$.*

If $p = 2$ and $E|F$ is ramified of R-P or R-U type, then there is a canonical closed embedding $\mathcal{N}_E \hookrightarrow \mathcal{M}_E$ that forgets about the polarization λ . In this way, it follows that $\tilde{\lambda}$ is a polarization for any $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\mathrm{naive}}(S)$.

The statement of [Theorem 1.6](#) can also be expressed in terms of an isomorphism of moduli spaces $\mathcal{M}_{E, \mathrm{pol}} \xrightarrow{\sim} \mathcal{M}_E$. Here $\mathcal{M}_{E, \mathrm{pol}}$ is a moduli space of PEL type, defined by mapping $S \in \mathrm{Nilp}_{\check{O}_F}$ to the set of tuples $(X, \iota, \tilde{\lambda}, \varrho)$ where $(X, \iota, \varrho) \in \mathcal{M}_E(S)$ and $\tilde{\lambda}$ is a polarization as in the theorem.

We now briefly describe the contents of the subsequent sections of this paper. In [Section 2](#), we recall some facts about the quadratic extensions of F , the quaternion algebra $B|F$ and hermitian forms. In [Sections 3](#) and [4](#), we define the moduli spaces $\mathcal{N}_E^{\mathrm{naive}}$, introduce the straightening condition describing $\mathcal{N}_E \subseteq \mathcal{N}_E^{\mathrm{naive}}$ and prove our main theorem in both the cases R-P and R-U. Although the techniques are quite similar in both cases, we decided to treat these cases separately, since the results in both cases differ in important details. Finally, in [Section 5](#) we prove [Theorem 1.6](#) on the existence of the polarizations $\tilde{\lambda}$.

2. Preliminaries on quaternion algebras and hermitian forms

Let $F|\mathbb{Q}_2$ be a finite extension. In this section we will recall some facts about the quadratic extensions of F , the quaternion division algebra $B|F$ and certain hermitian forms. For more information on quaternion algebras, see for example the

book by Vignéras [1980]. A systematic classification of hermitian forms over local fields has been done by Jacobowitz [1962].

Let $E|F$ be a quadratic field extension and denote by O_F , resp. O_E , the rings of integers. There are three mutually exclusive possibilities for $E|F$:

- $E|F$ is unramified. Then $E = F[\delta]$ for δ a square root of a unit in F . We can choose δ such that $\delta^2 = 1 + 4u$ for some $u \in O_F^\times$. In this case, $O_E = O_F[(1+\delta)/2]$. The element $\gamma = (1+\delta)/2$ satisfies the Eisenstein equation $\gamma^2 - \gamma - u = 0$. In the following we will write $F^{(2)}$ instead of E and $O_F^{(2)}$ instead of O_E when talking about the unramified extension of F .
- $E|F$ is ramified and E is generated by the square root of a uniformizer in F . That is, $E = F[\Pi]$ and Π is given by the Eisenstein equation $\Pi^2 + \pi_0 = 0$ for a uniformizing element $\pi_0 \in O_F$. We also have $O_E = O_F[\Pi]$. Following Jacobowitz, we will say $E|F$ is of type R-P (which stands for “ramified-prime”).
- Finally, $E|F$ can be given by an Eisenstein equation of the form $\Pi^2 - t\Pi + \pi_0 = 0$ for a uniformizer π_0 and $t \in O_F$ such that $\pi_0|t|2$. Then $E|F$ is ramified and $O_E = O_F[\Pi]$. Here, E is generated by the square root of a unit in F . Indeed, for $\vartheta = 1 - 2\Pi/t$ we have $\vartheta^2 = 1 - 4\pi_0/t^2 \in O_F^\times$. Thus $E|F$ is said to be of type R-U (for “ramified-unit”).

We will use this notation throughout the paper.

Remark 2.1. The isomorphism classes of quadratic extensions of F correspond to nontrivial equivalence classes of $F^\times/(F^\times)^2$. We have $F^\times/(F^\times)^2 \simeq H^1(G_F, \mathbb{Z}/2\mathbb{Z})$ for the absolute Galois group G_F of F and $\dim H^1(G_F, \mathbb{Z}/2\mathbb{Z}) = 2 + d$, where $d = [F : \mathbb{Q}_2]$ is the degree of F over \mathbb{Q}_2 (see, for example, [Neukirch et al. 2008, Corollary 7.3.9]).

A representative of an equivalence class in $F^\times/F^{\times 2}$ can be chosen to be either a prime or a unit, and exactly half of the classes are represented by prime elements, the others being represented by units. It follows that there are, up to isomorphism, 2^{1+d} different extensions $E|F$ of type R-P and $2^{1+d} - 2$ extensions of type R-U. (We have to exclude the trivial element $1 \in F^\times/F^{\times 2}$ and one unit element corresponding to the unramified extension.)

Lemma 2.2. *The inverse different of $E|F$ is given by $\mathfrak{D}_{E|F}^{-1} = \frac{1}{2\Pi} O_E$ in the case R-P and by $\mathfrak{D}_{E|F}^{-1} = \frac{1}{t} O_E$ in the case R-U.*

Proof. The inverse different is defined as

$$\mathfrak{D}_{E|F}^{-1} = \{\alpha \in E \mid \text{Tr}_{E|F}(\alpha O_E) \subseteq O_F\}.$$

It is enough to check the condition on the trace for the elements 1 and $\Pi \in O_E$. If we write $\alpha = \alpha_1 + \Pi\alpha_2$ with $\alpha_1, \alpha_2 \in F$, we get

$$\mathrm{Tr}_{E|F}(\alpha \cdot 1) = \alpha + \bar{\alpha} = 2\alpha_1 + \alpha_2(\Pi + \bar{\Pi}),$$

$$\mathrm{Tr}_{E|F}(\alpha \cdot \Pi) = \alpha\Pi + \bar{\alpha}\bar{\Pi} = \alpha_1(\Pi + \bar{\Pi}) + \alpha_2(\Pi^2 + \bar{\Pi}^2).$$

In the case R-P we have $\Pi + \bar{\Pi} = 0$ and $\Pi^2 + \bar{\Pi}^2 = 2\pi_0$, while in the case R-U, $\Pi + \bar{\Pi} = t$ and $\Pi^2 + \bar{\Pi}^2 = t^2 - 2\pi_0$. It is now easy to deduce that the inverse different is of the claimed form. \square

Over F , there exists up to isomorphism exactly one quaternion division algebra B , with unique maximal order O_B . For every quadratic extension $E|F$, there exists an embedding $E \hookrightarrow B$ and this induces an embedding $O_E \hookrightarrow O_B$. If $E|F$ is ramified, a basis for O_E as O_F -module is given by $(1, \Pi)$. We would like to extend this to an O_F -basis of O_B .

Lemma 2.3. (1) *If $E|F$ is of type R-P, there exists an embedding $F^{(2)} \hookrightarrow B$ such that $\delta\Pi = -\Pi\delta$. An O_F -basis of O_B is then given by $(1, \gamma, \Pi, \gamma \cdot \Pi)$, where $\gamma = (1 + \delta)/2$.*

(2) *If $E|F$ is of type R-U, there exists an embedding $E_1 \hookrightarrow B$, where $E_1|F$ is of type R-P with uniformizer Π_1 such that $\vartheta\Pi_1 = -\Pi_1\vartheta$. The tuple $(1, \vartheta, \Pi_1, \vartheta\Pi_1)$ is an F -basis of B .*

Furthermore, there is also an embedding $\tilde{E} \hookrightarrow B$ with $\tilde{E}|F$ of type R-U with elements $\tilde{\Pi}$ and $\tilde{\vartheta}$ as above, such that $\vartheta\tilde{\vartheta} = -\tilde{\vartheta}\vartheta$ and $\tilde{\vartheta}^2 = 1 + (t^2/\pi_0) \cdot u$ for some unit $u \in F$. In terms of this embedding, an O_F -basis of O_B is given by $(1, \Pi, \tilde{\Pi}, \Pi \cdot \tilde{\Pi}/\pi_0)$. Also,

$$(2-1) \quad \frac{\Pi \cdot \tilde{\Pi}}{\pi_0} = \gamma$$

for some embedding $F^{(2)} \hookrightarrow B$ of the unramified extension and $\gamma^2 - \gamma - u = 0$. Hence, $O_B = O_F[\Pi, \gamma]$ as O_F -algebra.

Proof. (1) This is [Vignéras 1980, II. Corollary 1.7].

(2) By [Vignéras 1980, I. Corollary 2.4], it suffices to find a uniformizer $\Pi_1^2 \in F^\times \setminus \mathrm{Nm}_{E|F}(E^\times)$ in order to prove the first part. But $\mathrm{Nm}_{E|F}(E^\times) \subseteq F^\times$ is a subgroup of order 2 and $F^{\times 2} \subseteq \mathrm{Nm}_{E|F}(E^\times)$. On the other hand, the residue classes of uniformizing elements in $F^\times/F^{\times 2}$ generate the whole group. Thus they cannot all be contained in $\mathrm{Nm}_{E|F}(E^\times)$.

For the second part, choose a unit $\delta \in F^{(2)}$ with $\delta^2 = 1 + 4u \in F^\times \setminus F^{\times 2}$ for some $u \in O_F^\times$ and set $\gamma = (1 + \delta)/2$. Let $\tilde{E}|F$ be of type R-U, generated by $\tilde{\vartheta}$ with $\tilde{\vartheta}^2 = 1 + (t^2/\pi_0) \cdot u$. We have to show that $\tilde{\vartheta}^2$ is not contained in $\mathrm{Nm}_{E|F}(E^\times)$.

Assume it is a norm, so $\tilde{\vartheta}^2 = \text{Nm}_{E|F}(b)$ for a unit $b \in E^\times$. Then b is of the form $b = 1 + x \cdot (t/\Pi)$ for some $x \in O_E$. Indeed, let ℓ be the Π -adic valuation of $b - 1$, i.e., $b = 1 + x \cdot \Pi^\ell$ and $x \in O_E^\times$. We have

$$(2-2) \quad 1 + (t^2/\pi_0) \cdot u = \text{Nm}_{E|F}(b) = 1 + \text{Tr}_{E|F}(x\Pi^\ell) + \text{Nm}_{E|F}(x\Pi^\ell).$$

Let v be the π_0 -adic valuation on F ; then $v(\text{Nm}_{E|F}(x\Pi^\ell)) = \ell$ and $v(\text{Tr}_{E|F}(x\Pi^\ell)) \geq v(t) + \lfloor \frac{\ell}{2} \rfloor$, by [Lemma 2.2](#). On the left-hand side, we have $v((t^2/\pi_0) \cdot u) = 2v(t) - 1$. Comparing the valuations on both sides of (2-2), the assumption $\ell < 2v(t) - 1$ now quickly leads to a contradiction.

Hence $\ell \geq 2v(t) - 1$ and $b = 1 + x \cdot (t/\Pi)$ for some $x \in O_E$. Again,

$$1 + (t^2/\pi_0) \cdot u = \text{Nm}_{E|F}(b) = 1 + \text{Tr}_{E|F}(xt/\Pi) + \text{Nm}_{E|F}(xt/\Pi).$$

An easy calculation shows that the residue $\bar{x} \in k = O_E/\Pi = O_F/\pi_0$ of x satisfies $u = x + x^2$. But this equation has no solution in k , since a solution of $\gamma^2 - \gamma - u = 0$ generates the unramified quadratic extension of F . It follows that $\tilde{\vartheta}^2$ cannot be a norm.

Using again [\[Vignéras 1980, I. Corollary 2.4\]](#), we find an embedding $\tilde{E} \hookrightarrow B$ such that $\vartheta \tilde{\vartheta} = -\tilde{\vartheta} \vartheta$.

We have $\Pi = t(1 + \vartheta)/2$ and $\tilde{\Pi} = \pi_0(1 + \tilde{\vartheta})/t$; thus

$$\frac{\Pi \cdot \tilde{\Pi}}{\pi_0} = \frac{(1 + \vartheta) \cdot (1 + \tilde{\vartheta})}{2} = \frac{1 + \vartheta + \tilde{\vartheta} + \vartheta \cdot \tilde{\vartheta}}{2},$$

and

$$\begin{aligned} (\vartheta + \tilde{\vartheta} + \vartheta \cdot \tilde{\vartheta})^2 &= \vartheta^2 + \tilde{\vartheta}^2 - \vartheta^2 \cdot \tilde{\vartheta}^2 \\ &= (1 - 4\pi_0/t^2) + (1 + t^2u/\pi_0) - (1 - 4\pi_0/t^2)(1 + t^2u/\pi_0) \\ &= 1 + 4u. \end{aligned}$$

Hence $\gamma \mapsto \Pi \cdot \tilde{\Pi}/\pi_0$ induces an embedding $F^{(2)} \hookrightarrow B$.

It remains to prove that the tuple $u = (1, \Pi, \tilde{\Pi}, \Pi \cdot \tilde{\Pi}/\pi_0)$ is a basis of O_B as O_F -module. By [\[Vignéras 1980, I. Corollary 4.8\]](#), it suffices to check that the discriminant

$$\text{disc}(u) = \det(\text{Trd}(u_i u_j)) \cdot O_F$$

is equal to $\text{disc}(O_B)$. An easy calculation shows $\det(\text{Trd}(u_i u_j)) \cdot O_F = \pi_0 O_F$ and then the assertion follows from [\[Vignéras 1980, V, II. Corollary 1.7\]](#). \square

For the remainder of this section, we will consider lattices Λ in a 2-dimensional E -vector space C with a split $E|F$ -hermitian¹ form h . Recall from [\[Jacobowitz 1962\]](#) that, up to isomorphism, there are two different $E|F$ -hermitian vector

¹Here and in the following, sesquilinear forms will be linear from the left and semilinear from the right.

spaces (C, h) of fixed dimension n , parametrized by the discriminant $\mathrm{disc}(C, h) \in F^\times / \mathrm{Nm}_{E|F}(E^\times)$. A hermitian space (C, h) is called *split* whenever $\mathrm{disc}(C, h) = 1$. In our case, where (C, h) is split of dimension 2, we can find a basis (e_1, e_2) of C with $h(e_i, e_i) = 0$ and $h(e_1, e_2) = 1$.

Denote by Λ^\sharp the dual of a lattice $\Lambda \subseteq C$ with respect to h . The lattice Λ is called Π^i -*modular* if $\Lambda = \Pi^i \Lambda^\sharp$ (resp. *unimodular* or *self-dual* when $i = 0$). In contrast to the p -adic case with $p > 2$, there exist Π^i -modular lattices of more than one type in our case (cf. [Jacobowitz 1962]):

Proposition 2.4. *Define the norm ideal $\mathrm{Nm}(\Lambda)$ of Λ by*

$$(2-3) \quad \mathrm{Nm}(\Lambda) = \langle \{h(x, x) | x \in \Lambda\} \rangle \subseteq F.$$

Any Π^i -modular lattice $\Lambda \subseteq C$ is determined up to the action of $\mathrm{U}(C, h)$ by the ideal $\mathrm{Nm}(\Lambda) = \pi_0^\ell \mathcal{O}_F \subseteq F$. For $i = 0$ or 1 , the exponent ℓ can be any integer such that

$$\begin{aligned} |2| &\leq |\pi_0|^\ell \leq |1| && \text{for } E|F \text{ R-P, unimodular } \Lambda, \\ |2\pi_0| &\leq |\pi_0|^\ell \leq |\pi_0| && \text{for } E|F \text{ R-P, } \Pi\text{-modular } \Lambda, \\ |t| &\leq |\pi_0|^\ell \leq |1| && \text{for } E|F \text{ R-U, unimodular } \Lambda, \\ |t| &\leq |\pi_0|^\ell \leq |\pi_0| && \text{for } E|F \text{ R-U, } \Pi\text{-modular } \Lambda, \end{aligned}$$

where $|\cdot|$ is the (normalized) absolute value on F . Two Π^i -modular lattices Λ and Λ' are isomorphic if and only if $\mathrm{Nm}(\Lambda) = \mathrm{Nm}(\Lambda')$. \square

For any other i , the possible values of ℓ for a given Π^i -modular lattice Λ are easily obtained by shifting. In fact, we can choose an integer j such that $\Pi^j \Lambda$ is either unimodular or Π -modular. Then $\mathrm{Nm}(\Lambda) = \pi_0^{-j} \mathrm{Nm}(\Pi^j \Lambda)$ and we can apply the proposition above.

Since (C, h) is split, any Π^i -modular lattice Λ contains an *isotropic* vector v (i.e., with $h(v, v) = 0$). After rescaling with a suitable power of Π , we can extend v to a basis of Λ . Hence there always exists a basis (e_1, e_2) of Λ such that h is represented by a matrix of the form

$$(2-4) \quad H_\Lambda = \begin{pmatrix} x & \bar{\Pi}^i \\ \Pi^i & \end{pmatrix}, \quad x \in F.$$

If $x = 0$ in this representation, then $\mathrm{Nm}(\Lambda) = \pi_0^\ell \mathcal{O}_F$ is as small as possible, or in other words, the absolute value of $|\pi_0|^\ell$ is minimal. On the other hand, whenever $|\pi_0|^\ell$ takes the minimal absolute value for a given Π^i -modular lattice Λ , there exists a basis (e_1, e_2) of Λ such that h is represented by H_Λ with $x = 0$. Indeed, this follows because the ideal $\mathrm{Nm}(\Lambda)$ already determines Λ up to isomorphism. In this case (when $x = 0$), we call Λ a *hyperbolic* lattice. By the arguments above, a

Π^i -modular lattice is thus hyperbolic if and only if its norm is minimal. In all other cases, where Λ is Π^i -modular but not hyperbolic, we have $\text{Nm}(\Lambda) = x O_F$.

For further reference, we explicitly write down the norm of a hyperbolic lattice for the cases that we need later. For other values of i , the norm can easily be deduced from this by shifting (see also [Jacobowitz 1962, Table 9.1]).

Lemma 2.5. *A Π^i -modular lattice Λ is hyperbolic if and only if*

$$\text{Nm}(\Lambda) = 2O_F \quad \text{for } E|F \text{ R-P, } i = 0 \text{ or } -1,$$

$$\text{Nm}(\Lambda) = tO_F \quad \text{for } E|F \text{ R-U, } i = 0 \text{ or } 1.$$

The norm ideal of Λ is minimal among all norm ideals for Π^i -modular lattices in C . □

In the following, we will only consider the cases $i = 0$ or -1 for $E|F$ R-P and the cases $i = 0$ or 1 for $E|F$ R-U, since these are the cases we will need later. We want to study the following question:

Question 2.6. Assume $E|F$ is R-P. Fix a Π^{-1} -modular lattice $\Lambda_{-1} \subseteq C$ (not necessarily hyperbolic). How many unimodular lattices $\Lambda_0 \subseteq \Lambda_{-1}$ are there and what norms $\text{Nm}(\Lambda_0)$ can appear? Dually, for a fixed unimodular lattice $\Lambda_0 \subseteq C$, how many Π^{-1} -modular lattices Λ_{-1} with $\Lambda_0 \subseteq \Lambda_{-1}$ exist and what are their norms?

We can ask the same question for $E|F$ R-U and unimodular, resp. Π -modular, lattices.

Of course, such an inclusion is always of index 1. The inclusions $\Lambda_0 \subseteq \Lambda_{-1}$ of index 1 correspond to lines in $\Lambda_{-1}/\Pi\Lambda_{-1}$. Denote by q the number of elements in the common residue field of O_F and O_E . Then there exist at most $q + 1$ such Π -modular lattices Λ_0 for a given Λ_{-1} . The same bound holds in the dual case, i.e., there are at most $q + 1$ Π^{-1} -modular lattices containing a given unimodular lattice Λ_0 . Propositions 2.7 and 2.8 below provide an exhaustive answer to Question 2.6. Since the proofs consist of a lengthy but simple case-by-case analysis, we will leave them to the interested reader.

Proposition 2.7. *Let $E|F$ be of type R-P.*

- (1) *Let $\Lambda_{-1} \subseteq C$ be a Π^{-1} -modular hyperbolic lattice. There are $q + 1$ hyperbolic unimodular lattices contained in Λ_{-1} .*
- (2) *Let $\Lambda_{-1} \subseteq C$ be a Π^{-1} -modular nonhyperbolic lattice. Let $\text{Nm}(\Lambda_{-1}) = \pi_0^\ell O_F$. Then Λ_{-1} contains one unimodular lattice Λ_0 with $\text{Nm}(\Lambda_0) = \pi_0^{\ell+1} O_F$ and q unimodular lattices of norm $\pi_0^\ell O_F$.*
- (3) *Let $\Lambda_0 \subseteq C$ be a unimodular hyperbolic lattice. There are two hyperbolic Π^{-1} -modular lattices $\Lambda_{-1} \supseteq \Lambda_0$ and $q - 1$ nonhyperbolic Π^{-1} -modular lattices $\Lambda_{-1} \supseteq \Lambda_0$ with $\text{Nm}(\Lambda_{-1}) = 2/\pi_0 O_F$.*

- (4) Let $\Lambda_0 \subseteq C$ be unimodular nonhyperbolic. Let $\mathrm{Nm}(\Lambda_0) = \pi_0^\ell O_F$. There exists one Π^{-1} -modular lattice $\Lambda_{-1} \supseteq \Lambda_0$ with $\mathrm{Nm}(\Lambda_{-1}) = \pi_0^\ell O_F$ and, unless $\ell = 0$, there are q nonhyperbolic Π^{-1} -modular lattices $\Lambda_{-1} \supseteq \Lambda_0$ with $\mathrm{Nm}(\Lambda_{-1}) = \pi_0^{\ell-1} O_F$.

Note that the total number of unimodular, resp. Π^{-1} -modular, lattices found for $\Lambda = \Lambda_{-1}$, resp. Λ_0 , is $q + 1$ except in the case of [Proposition 2.7 \(4\)](#) when $\ell = 0$. In that particular case, there is just one Π^{-1} -modular lattice contained in Λ_0 . The same phenomenon also appears in the case R-U; see part (2) of the following proposition.

Proposition 2.8. *Let $E|F$ be of type R-U.*

- (1) *Let $\Lambda_0 \subseteq C$ be a unimodular hyperbolic lattice. There are $q + 1$ hyperbolic Π -modular lattices $\Lambda_1 \subseteq \Lambda_0$.*
- (2) *Let $\Lambda_0 \subseteq C$ be unimodular nonhyperbolic with $\mathrm{Nm}(\Lambda_0) = \pi_0^\ell O_F$. There is one Π -modular lattice $\Lambda_1 \subseteq \Lambda_0$ with norm ideal $\mathrm{Nm}(\Lambda_1) = \pi_0^{\ell+1} O_F$ and if $\ell \neq 0$, there are also q nonhyperbolic Π -modular lattices $\Lambda_1 \subseteq \Lambda_0$ with $\mathrm{Nm}(\Lambda_1) = \pi_0^\ell O_F$.*
- (3) *Let $\Lambda_1 \subseteq C$ be a Π -modular hyperbolic lattice. There are two unimodular hyperbolic lattices containing Λ_1 and $q - 1$ unimodular lattices Λ_0 with $\Lambda_1 \subseteq \Lambda_0$ and $\mathrm{Nm}(\Lambda_0) = t/\pi_0 O_F$.*
- (4) *Let $\Lambda_1 \subseteq C$ be a Π -modular nonhyperbolic lattice and let $\mathrm{Nm}(\Lambda_1) = \pi_0^\ell O_F$. The lattice Λ_1 is contained in q unimodular lattices of norm $\pi_0^{\ell-1} O_F$ and in one unimodular lattice Λ_0 with $\mathrm{Nm}(\Lambda_0) = \pi_0^\ell O_F$.*

If $E|F$ is a quadratic extension of type R-U such that $|t| = |\pi_0|$, there exist only hyperbolic Π -modular lattices in C and hence case (4) of [Proposition 2.8](#) does not appear.

3. The moduli problem in the case R-P

Throughout this section, $E|F$ is a quadratic extension of type R-P, i.e., there exist uniformizing elements $\pi_0 \in F$ and $\Pi \in E$ such that $\Pi^2 + \pi_0 = 0$. Then $O_E = O_F[\Pi]$ for the rings of integers O_F and O_E of F and E , respectively. Let k be the common residue field with q elements, \bar{k} an algebraic closure, and \check{F} the completion of the maximal unramified extension of F , with ring of integers $\check{O}_F = W_{O_F}(\bar{k})$. Let σ be the lift of the Frobenius in $\mathrm{Gal}(\bar{k}|k)$ to $\mathrm{Gal}(\check{O}_F|O_F)$.

3A. The definition of the naive moduli problem $\mathcal{N}_E^{\mathrm{naive}}$. We first construct a functor $\mathcal{N}_E^{\mathrm{naive}}$ on $\mathrm{Nilp}_{\check{O}_F}$, the category of \check{O}_F -schemes S such that $\pi_0 \mathcal{O}_S$ is locally nilpotent. We consider tuples (X, ι, λ) , where

- X is a formal O_F -module over S of dimension 2 and height 4.

- $\iota : O_E \rightarrow \text{End}(X)$ is an action of O_E satisfying the *Kottwitz condition*: The characteristic polynomial of $\iota(\alpha)$ on $\text{Lie } X$ for any $\alpha \in O_E$ is

$$\text{char}(\text{Lie } X, T \mid \iota(\alpha)) = (T - \alpha)(T - \bar{\alpha}).$$

Here $\alpha \mapsto \bar{\alpha}$ is the nontrivial Galois automorphism and the right-hand side is a polynomial with coefficients in \mathcal{O}_S via the composition $O_F[T] \hookrightarrow \check{O}_F[T] \rightarrow \mathcal{O}_S[T]$.

- $\lambda : X \rightarrow X^\vee$ is a principal polarization on X such that the Rosati involution satisfies $\iota(\alpha)^* = \iota(\bar{\alpha})$ for $\alpha \in O_E$.

Definition 3.1. A *quasi-isogeny* (resp. an *isomorphism*) $\varphi : (X, \iota, \lambda) \rightarrow (X', \iota', \lambda')$ of two such tuples (X, ι, λ) and (X', ι', λ') over S is an O_E -linear quasi-isogeny of height 0 (resp. an O_E -linear isomorphism) $\varphi : X \rightarrow X'$ such that $\lambda = \varphi^*(\lambda')$.

Denote the group of quasi-isogenies $\varphi : (X, \iota, \lambda) \rightarrow (X, \iota, \lambda)$ by $\text{QIsog}(X, \iota, \lambda)$.

For $S = \text{Spec } \bar{k}$ we have the following proposition:

Proposition 3.2. *Up to isogeny, there exists precisely one tuple $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ over $\text{Spec } \bar{k}$ such that the group $\text{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ contains $\text{SU}(C, h)$ as a closed subgroup. Here $\text{SU}(C, h)$ is the special unitary group for a 2-dimensional E -vector space C with split $E|F$ -hermitian form h .*

Remark 3.3. If $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ is as in the proposition, we always have $\text{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}}) \cong \text{U}(C, h)$. This follows directly from the proof and gives a more natural way to describe the framing object. However, we will need the slightly stronger statement of the proposition later, in [Lemma 3.15](#).

Proof of Proposition 3.2. We first show uniqueness. Let $(X, \iota, \lambda)/\text{Spec } \bar{k}$ be such a tuple. Its (relative) rational Dieudonné module N_X is a 4-dimensional vector space over \check{F} with an action of E and an alternating form $\langle \cdot, \cdot \rangle$ such that for all $x, y \in N_X$,

$$(3-1) \quad \langle x, \Pi y \rangle = -\langle \Pi x, y \rangle.$$

The space N_X has the structure of a 2-dimensional vector space over $\check{E} = E \otimes_F \check{F}$ and we can define an $\check{E}|\check{F}$ -hermitian form on it via

$$(3-2) \quad h(x, y) = \langle \Pi x, y \rangle + \Pi \langle x, y \rangle.$$

The alternating form can be recovered from h by

$$(3-3) \quad \langle x, y \rangle = \text{Tr}_{\check{E}|\check{F}} \left(\frac{1}{2\Pi} \cdot h(x, y) \right).$$

Furthermore we have on N_X a σ -linear operator F , the Frobenius, and a σ^{-1} -linear operator V , the Verschiebung, that satisfy $VF = FV = \pi_0$. Recall that σ is the lift

of the Frobenius on \check{O}_F . Since $\langle \cdot, \cdot \rangle$ comes from a polarization, we have

$$\langle \mathbf{F}x, y \rangle = \langle x, \mathbf{V}y \rangle^\sigma$$

and

$$h(\mathbf{F}x, y) = h(x, \mathbf{V}y)^\sigma$$

for all $x, y \in N_X$. Let us consider the σ -linear operator $\tau = \Pi \mathbf{V}^{-1}$. Its slopes are all zero, since N_X is isotypical of slope $\frac{1}{2}$. (This follows from the condition on $\mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$.) We set $C = N_X^\tau$. This is a 2-dimensional vector space over E and $N_X = C \otimes_E \check{E}$. Now h induces an $E|F$ -hermitian form on C since

$$h(\tau x, \tau y) = h(-\mathbf{F}\Pi^{-1}x, \Pi\mathbf{V}^{-1}y) = -h(\Pi^{-1}x, \Pi y)^\sigma = h(x, y)^\sigma.$$

A priori, there are up to isomorphism two possibilities for (C, h) , either h is split on C or nonsplit. But automorphisms of (C, h) correspond to elements of $\mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$. The unitary groups of (C, h) for h split and h nonsplit are not isomorphic and they cannot contain each other as a closed subgroup. Hence the condition on $\mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ implies that h is split.

Assume now we have two different objects (X, ι, λ) and (X', ι', λ') as in the proposition. These give us isomorphic vector spaces (C, h) and (C', h') and an isomorphism between these extends to an isomorphism between N_X and N'_X (respecting all rational structure) which corresponds to a quasi-isogeny between (X, ι, λ) and (X', ι', λ') .

The existence of $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ now follows from the fact that a 2-dimensional E -vector space (C, h) with split $E|F$ -hermitian form contains a unimodular lattice Λ . Indeed, this gives us a lattice $M = \Lambda \otimes_{O_E} \check{O}_E \subseteq C \otimes_E \check{E}$. We extend h to $N = C \otimes_E \check{E}$ and define the \check{F} -linear alternating form $\langle \cdot, \cdot \rangle$ as in (3-3). Now M is unimodular with respect to $\langle \cdot, \cdot \rangle$, because $\frac{1}{2\Pi} \check{O}_E$ is the inverse different of $\check{E}|\check{F}$ (see Lemma 2.2). We choose the operators \mathbf{F} and \mathbf{V} on M such that $\mathbf{F}\mathbf{V} = \mathbf{V}\mathbf{F} = \pi_0$ and $\Lambda = M^\tau$ for $\tau = \Pi\mathbf{V}^{-1}$. This makes M a (relative) Dieudonné module and we define $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ as the corresponding formal O_F -module. \square

We fix such a framing object $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ over $\mathrm{Spec} \bar{k}$.

Definition 3.4. For arbitrary $S \in \mathrm{Nilp}_{\check{O}_F}$, let $\bar{S} = S \times_{\mathrm{Spf} \check{O}_F} \mathrm{Spec} \bar{k}$. Define $\mathcal{N}_E^{\mathrm{naive}}(S)$ as the set of equivalence classes of tuples $(X, \iota, \lambda, \varrho)$ over S , where (X, ι, λ) as above and

$$\varrho : X \times_S \bar{S} \rightarrow \mathbb{X} \times_{\mathrm{Spec} \bar{k}} \bar{S}$$

is a quasi-isogeny between the tuple (X, ι, λ) and the framing object $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ (after base change to \bar{S}). Two objects $(X, \iota, \lambda, \varrho)$ and $(X', \iota', \lambda', \varrho')$ are equivalent if and only if there exists an isomorphism $\varphi : (X, \iota, \lambda) \rightarrow (X', \iota', \lambda')$ such that $\varrho = \varrho' \circ (\varphi \times_S \bar{S})$.

Remark 3.5. (1) The morphism ϱ is a quasi-isogeny in the sense of [Definition 3.1](#), i.e., we have $\lambda = \varrho^*(\lambda_{\mathbb{X}})$. Similarly, we have $\lambda = \varphi^*(\lambda')$ for the isomorphism φ . We obtain an equivalent definition of $\mathcal{N}_E^{\text{naive}}$ if we replace strict equality by the condition that, locally on S , λ and $\varrho^*(\lambda_{\mathbb{X}})$ (resp. $\varphi^*(\lambda')$) only differ by a scalar in O_F^\times . This variant is used in the definition of RZ-spaces of PEL type for $p > 2$ in [\[Rapoport and Zink 1996\]](#). In this paper we will use the version with strict equality, since it simplifies the formulation of the straightening condition; see [Definition 3.11](#) below.

(2) $\mathcal{N}_E^{\text{naive}}$ is pro-representable by a formal scheme, formally locally of finite type over $\text{Spf } \check{O}_F$. This follows from [\[Rapoport and Zink 1996, Theorem 3.25\]](#).

As a next step, we use Dieudonné theory in order to get a better understanding of the special fiber of $\mathcal{N}_E^{\text{naive}}$. Let $N = N_{\mathbb{X}}$ be the rational Dieudonné module of the base point $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ of $\mathcal{N}_E^{\text{naive}}$. This is a 4-dimensional vector space over \check{F} , equipped with an E -action, an alternating form $\langle \cdot, \cdot \rangle$ and two operators V and F . As in the proof of [Proposition 3.2](#), the form $\langle \cdot, \cdot \rangle$ satisfies condition (3-1):

$$(3-4) \quad \langle x, \Pi y \rangle = -\langle \Pi x, y \rangle.$$

A point $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\text{naive}}(\bar{k})$ corresponds to an \check{O}_F -lattice $M_X \subseteq N$. It is stable under the actions of the operators V and F and of the ring O_E . Furthermore M_X is unimodular under $\langle \cdot, \cdot \rangle$, i.e., $M_X = M_X^\vee$, where

$$M_X^\vee = \{x \in N \mid \langle x, y \rangle \in \check{O}_F \text{ for all } y \in M_X\}.$$

We can regard N as a 2-dimensional vector space over \check{E} with the $\check{E}|\check{F}$ -hermitian form h defined by

$$(3-5) \quad h(x, y) = \langle \Pi x, y \rangle + \Pi \langle x, y \rangle.$$

Let $\check{O}_E = O_E \otimes_{O_F} \check{O}_F$. Then $M_X \subseteq N$ is an \check{O}_E -lattice and we have

$$M_X = M_X^\vee = M_X^\sharp,$$

where M_X^\sharp is the dual lattice of M_X with respect to h . The latter equality follows from the formula

$$(3-6) \quad \langle x, y \rangle = \text{Tr}_{\check{E}|\check{F}} \left(\frac{1}{2\Pi} \cdot h(x, y) \right)$$

and the fact that the inverse different of $E|F$ is $\mathfrak{D}_{E|F}^{-1} = \frac{1}{2\Pi} O_E$ (see [Lemma 2.2](#)). We can thus write the set $\mathcal{N}_E^{\text{naive}}(\bar{k})$ as

$$(3-7) \quad \mathcal{N}_E^{\text{naive}}(\bar{k}) = \{ \check{O}_E\text{-lattices } M \subseteq N_{\mathbb{X}} \mid M^\sharp = M, \pi_0 M \subseteq VM \subseteq M \}.$$

Let $\tau = \Pi V^{-1}$. This is a σ -linear operator on N with all slopes zero. The elements invariant under τ form a 2-dimensional E -vector space $C = N^\tau$. The hermitian form

h is invariant under τ , hence it induces a split hermitian form on C which we denote again by h . With the same proof as in [Kudla and Rapoport 2014, Lemma 3.2], we have:

Lemma 3.6. *Let $M \in \mathcal{N}_E^{\mathrm{naive}}(\bar{k})$. Then:*

- (1) $M + \tau(M)$ is τ -stable.
- (2) Either M is τ -stable and $\Lambda_0 = M^\tau \subseteq C$ is unimodular ($\Lambda_0^\sharp = \Lambda_0$) or M is not τ -stable and then $\Lambda_{-1} = (M + \tau(M))^\tau \subseteq C$ is Π^{-1} -modular ($\Lambda_{-1}^\sharp = \Pi\Lambda_{-1}$).

Under the identification $N = C \otimes_E \check{E}$, we get $M = \Lambda_0 \otimes_{O_E} \check{O}_E$ for a τ -stable Dieudonné lattice M . If M is not τ -stable, we have $M + \tau M = \Lambda_{-1} \otimes_{O_E} \check{O}_E$ and $M \subseteq \Lambda_{-1} \otimes_{O_E} \check{O}_E$ is a sublattice of index 1. The next lemma is the analogue of [Kudla and Rapoport 2014, Lemma 3.3].

Lemma 3.7. (1) *Fix a Π^{-1} -modular lattice $\Lambda_{-1} \subseteq C$. There is an injective map*

$$i_{\Lambda_{-1}} : \mathbb{P}(\Lambda_{-1}/\Pi\Lambda_{-1})(\bar{k}) \hookrightarrow \mathcal{N}_E^{\mathrm{naive}}(\bar{k})$$

mapping a line $\ell \subseteq (\Lambda_{-1}/\Pi\Lambda_{-1}) \otimes \bar{k}$ to its preimage in $\Lambda_{-1} \otimes \check{O}_E$. Identify $\mathbb{P}(\Lambda_{-1}/\Pi\Lambda_{-1})(\bar{k})$ with its image in $\mathcal{N}_E^{\mathrm{naive}}(\bar{k})$. Then $\mathbb{P}(\Lambda_{-1}/\Pi\Lambda_{-1})(\bar{k}) \subseteq \mathbb{P}(\Lambda_{-1}/\Pi\Lambda_{-1})(\bar{k})$ is the set of τ -invariant Dieudonné lattices $M \subseteq \Lambda_{-1} \otimes \check{O}_E$.

- (2) *The set $\mathcal{N}_E^{\mathrm{naive}}(\bar{k})$ is a union*

$$(3-8) \quad \mathcal{N}_E^{\mathrm{naive}}(\bar{k}) = \bigcup_{\Lambda_{-1} \subseteq C} \mathbb{P}(\Lambda_{-1}/\Pi\Lambda_{-1})(\bar{k}),$$

ranging over all Π^{-1} -modular lattices $\Lambda_{-1} \subseteq C$. The projective lines corresponding to the lattices Λ_{-1} and Λ'_{-1} intersect in $\mathcal{N}_E^{\mathrm{naive}}(\bar{k})$ if and only if $\Lambda_0 = \Lambda_{-1} \cap \Lambda'_{-1}$ is unimodular. In this case, their intersection consists of the point $M = \Lambda_0 \otimes \check{O}_E \in \mathcal{N}_E^{\mathrm{naive}}(\bar{k})$.

Proof. We only have to prove that the map $i_{\Lambda_{-1}}$ is well-defined. Denote by M the preimage of $\ell \subseteq (\Lambda_{-1}/\Pi\Lambda_{-1}) \otimes \bar{k}$ in $\Lambda_{-1} \otimes \check{O}_E$. We need to show that M is an element in $\mathcal{N}_E^{\mathrm{naive}}(\bar{k})$ under the identification of (3-7). It is clearly a sublattice of index 1 in $\Lambda_{-1} \otimes \check{O}_E$, stable under the actions of F , V and O_E .

Let $e_1 \in \Lambda_{-1} \otimes \check{O}_E$ such that $e_1 \otimes \bar{k}$ generates ℓ . We can extend this to a basis (e_1, e_2) of Λ_{-1} and with respect to this basis, h is represented by a matrix of the form

$$\begin{pmatrix} x & -\Pi^{-1} \\ \Pi^{-1} & y \end{pmatrix},$$

with $x, y \in \Pi^{-1}\check{O}_E \cap \check{O}_F = \check{O}_F$. The lattice $M \subseteq \Lambda_{-1} \otimes \check{O}_E$ is generated by e_1 and Πe_2 . With respect to this new basis, h is now given by the matrix

$$\begin{pmatrix} x & 1 \\ 1 & \pi_0 y \end{pmatrix}.$$

Since all entries of the matrix are integral, we have $M \subseteq M^\sharp$. But this already implies $M^\sharp = M$, because they both have index 1 in $\Lambda_{-1} \otimes \check{O}_E$. Thus $M \in \mathcal{N}_E^{\text{naive}}(\bar{k})$ and $i_{\Lambda_{-1}}$ is well-defined. \square

Remark 3.8. (1) Recall from [Proposition 2.4](#) that the isomorphism type of a Π^i -modular lattice $\Lambda \subseteq C$ only depends on its norm ideal $\text{Nm}(\Lambda) = \langle \{h(x, x) | x \in \Lambda\} \rangle = \pi_0^\ell O_F \subseteq F$. In the case that $\Lambda = \Lambda_0$ or Λ_{-1} is unimodular or Π^{-1} -modular, ℓ can be any integer such that $|1| \geq |\pi_0|^\ell \geq |2|$. In particular, there are always at least two possible values for ℓ . Recall from [Lemma 2.5](#) that Λ is *hyperbolic* if and only if $\text{Nm}(\Lambda) = 2O_F$.

(2) The intersection behavior of the projective lines in $\mathcal{N}_E^{\text{naive}}(\bar{k})$ can be deduced from [Proposition 2.7](#). In particular, for a given unimodular lattice $\Lambda_0 \subseteq C$ with $\text{Nm}(\Lambda_0) \subseteq \pi_0 O_F$, there are $q + 1$ lines intersecting in $M = \Lambda_0 \otimes \check{O}_E$. If $\text{Nm}(\Lambda_0) = O_F$, the lattice $M = \Lambda_0 \otimes \check{O}_E$ is only contained in one projective line. On the other hand, a projective line $\mathbb{P}(\Lambda_{-1}/\Pi\Lambda_{-1})(\bar{k}) \subseteq \mathcal{N}_E^{\text{naive}}(\bar{k})$ contains $q + 1$ points corresponding to unimodular lattices in C . By [Lemma 3.7 \(1\)](#), these are exactly the k -rational points of $\mathbb{P}(\Lambda_{-1}/\Pi\Lambda_{-1})$.

(3) If we restrict the union at the right-hand side of (3-8) to hyperbolic Π^{-1} -modular lattices $\Lambda_{-1} \subseteq C$ (i.e., $\text{Nm}(\Lambda_{-1}) = 2O_F$; see [Lemma 2.5](#)), we obtain a canonical subset $\mathcal{N}_E(\bar{k}) \subseteq \mathcal{N}_E^{\text{naive}}(\bar{k})$ and there is a description of \mathcal{N}_E as a pro-representable functor on $\text{Nilp}_{\check{O}_F}$ (see below). We will see later ([Theorem 3.16](#)) that \mathcal{N}_E is isomorphic to the Drinfeld moduli space \mathcal{M}_{D_F} , described in [[Boutot and Carayol 1991](#), I.3]. In particular, the underlying topological space of \mathcal{N}_E is connected. (The induced topology on the projective lines is the Zariski topology; see [Proposition 3.9](#).) Moreover, each projective line in $\mathcal{N}_E(\bar{k})$ has $q + 1$ intersection points and there are two projective lines intersecting in each such point (see also [Proposition 2.7](#)).

We fix such an intersection point $P \in \mathcal{N}_E(\bar{k})$. Now going back to $\mathcal{N}_E^{\text{naive}}(\bar{k})$, there are $q - 1$ additional lines going through $P \in \mathcal{N}_E^{\text{naive}}(\bar{k})$ that correspond to nonhyperbolic lattices in C (see [Proposition 2.7](#)). Each of these additional lines contains P as its only “hyperbolic” intersection point, all other intersection points on this line and the line itself correspond to unimodular, resp. Π^{-1} -modular, lattices $\Lambda \subseteq C$ of norm $\text{Nm}(\Lambda) = (2/\pi_0)O_F$ (whereas all hyperbolic lattices occurring have the norm ideal $2O_F$; see [Lemma 2.5](#)). Assume $\mathbb{P}(\Lambda/\Pi\Lambda)(\bar{k}) \subseteq \mathcal{N}_E^{\text{naive}}(\bar{k})$ is such a line and let $P' \in \mathbb{P}(\Lambda/\Pi\Lambda)(\bar{k})$ be an intersection point, where $P \neq P'$. There are again q more lines going through P' (always $q + 1$ in total) that correspond to lattices with norm ideal $\text{Nm}(\Lambda) = (2/\pi_0^2)O_F$, and these lines again have more intersection points and so on. This goes on until we reach lines $\mathbb{P}(\Lambda'/\Pi\Lambda')(\bar{k})$ with $\text{Nm}(\Lambda') = O_F$. Each of these lines contains q points that correspond to unimodular lattices $\Lambda_0 \subseteq C$ with $\text{Nm}(\Lambda_0) = O_F$. Such a lattice is only contained




Figure 1. The reduced locus of $\mathcal{N}_E^{\text{naive}}$ for $E|F$ of type R-P where $F|\mathbb{Q}_2$ has ramification index e and inertia degree f . Solid lines are given by subschemes $\mathcal{N}_{E,\Lambda}$ for hyperbolic lattices Λ .

in one Π^{-1} -modular lattice (see part (4) of Proposition 2.7). Hence, these points are only contained in one projective line, namely $\mathbb{P}(\Lambda'/\Pi\Lambda')(\bar{k})$.

In other words, each intersection point $P \in \mathcal{N}_E(\bar{k})$ has a “tail”, consisting of finitely many projective lines, which is the connected component of P in $(\mathcal{N}_E^{\text{naive}}(\bar{k}) \setminus \mathcal{N}_E(\bar{k})) \cup \{P\}$. Figure 1 shows a drawing of $(\mathcal{N}_E^{\text{naive}})_{\text{red}}$ for the cases $F = \mathbb{Q}_2$ (on the left-hand side) and $F|\mathbb{Q}_2$ a ramified quadratic extension (on the right-hand side). The “tails” are indicated by dashed lines.

Fix a Π^{-1} -modular lattice $\Lambda = \Lambda_{-1} \subseteq C$. Let X_Λ^+ be the formal \mathcal{O}_F -module over $\mathrm{Spec} \bar{k}$ associated to the Dieudonné lattice $M = \Lambda \otimes_{\check{\mathcal{O}}_E} \check{\mathcal{O}}_E \subseteq N$. It comes with a canonical quasi-isogeny

$$\varrho_\Lambda^+ : \mathbb{X} \rightarrow X_\Lambda^+$$

of F -height 1. We define a subfunctor $\mathcal{N}_{E,\Lambda} \subseteq \mathcal{N}_E^{\text{naive}}$ by mapping $S \in \mathrm{Nilp}_{\check{\mathcal{O}}_F}$ to

$$(3-9) \quad \mathcal{N}_{E,\Lambda}(S) = \{(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\text{naive}}(S) \mid (\varrho_\Lambda^+ \times S) \circ \varrho \text{ is an isogeny}\}.$$

Note that the condition of (3-9) is closed; cf. [Rapoport and Zink 1996, Proposition 2.9]. Hence $\mathcal{N}_{E,\Lambda}$ is representable by a closed formal subscheme of $\mathcal{N}_E^{\text{naive}}$. On geometric points, we have a bijection

$$(3-10) \quad \mathcal{N}_{E,\Lambda}(\bar{k}) \xrightarrow{\sim} \mathbb{P}(\Lambda/\Pi\Lambda)(\bar{k}),$$

as a consequence of Lemma 3.7 (1).

Proposition 3.9. *The reduced locus of $\mathcal{N}_E^{\text{naive}}$ is given by*

$$(\mathcal{N}_E^{\text{naive}})_{\text{red}} = \bigcup_{\Lambda \subseteq C} \mathcal{N}_{E,\Lambda},$$

where Λ runs over all Π^{-1} -modular lattices in C . For each Λ , there is an isomorphism of reduced schemes

$$\mathcal{N}_{E,\Lambda} \xrightarrow{\sim} \mathbb{P}(\Lambda/\Pi\Lambda),$$

inducing the map (3-10) on \bar{k} -valued points.

Proof. The embedding

$$(3-11) \quad \bigcup_{\Lambda \subseteq C} (\mathcal{N}_{E,\Lambda})_{\text{red}} \hookrightarrow (\mathcal{N}_E^{\text{naive}})_{\text{red}}$$

is closed, because each embedding $\mathcal{N}_{E,\Lambda} \subseteq \mathcal{N}_E^{\text{naive}}$ is closed and, locally on $(\mathcal{N}_E^{\text{naive}})_{\text{red}}$, the left-hand side is always only a finite union of $(\mathcal{N}_{E,\Lambda})_{\text{red}}$. It follows already that (3-11) is an isomorphism, since it is a bijection on \bar{k} -valued points (see (3-8) and (3-10)) and $(\mathcal{N}_E^{\text{naive}})_{\text{red}}$ is reduced by definition and locally of finite type over $\text{Spec } \bar{k}$ by Remark 3.5 (2).

For the second part of the proposition, we follow the proof presented in [Kudla and Rapoport 2014, Lemma 4.2]. Fix a Π^{-1} -modular lattice $\Lambda \subseteq C$ and let $M = \Lambda \otimes \check{O}_E \subseteq N$, as above. Now X_{Λ}^{+} is the formal O_F -module associated to M , but we also get a formal O_F -module X_{Λ}^{-} associated to the dual $M^{\sharp} = \Pi M$ of M . This comes with a natural isogeny

$$\text{nat}_{\Lambda} : X_{\Lambda}^{-} \rightarrow X_{\Lambda}^{+}$$

and a quasi-isogeny $\varrho_{\Lambda}^{-} : X_{\Lambda}^{-} \rightarrow \mathbb{X}$ of F -height 1. For $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\text{naive}}(S)$ where $S \in \text{Nilp}_{\check{O}_F}$, we consider the composition

$$\varrho_{\Lambda,X}^{-} = \varrho^{-1} \circ (\varrho_{\Lambda}^{-} \times S) : (X_{\Lambda}^{-} \times S) \rightarrow X.$$

By [Kudla and Rapoport 2014, Lemma 4.2], this composition is an isogeny if and only if $(\varrho_{\Lambda}^{+} \times S) \circ \varrho$ is an isogeny, or, in other words, if and only if $(X, \iota, \lambda, \varrho) \in \mathcal{N}_{E,\Lambda}(S)$. Let $\mathbb{D}_{X_{\Lambda}^{-}}(S)$ be the (relative) Grothendieck–Messing crystal of X_{Λ}^{-} evaluated at S (cf. [Ahsendorf et al. 2016, Definition 3.24] or [Ahsendorf 2011, Section 5.2]). This is a locally free \mathcal{O}_S -module of rank 4, isomorphic to $\Lambda/\pi_0\Lambda \otimes_{O_F} \mathcal{O}_S$. The kernel of $\mathbb{D}(\text{nat}_{\Lambda})(S)$ is given by $(\Lambda/\Pi\Lambda) \otimes_{O_F} \mathcal{O}_S$, locally a direct summand of rank 2 of $\mathbb{D}_{X_{\Lambda}^{-}}(S)$. For any $(X, \iota, \lambda, \varrho) \in \mathcal{N}_{E,\Lambda}(S)$, the kernel of $\varrho_{\Lambda,X}^{-}$ is contained in $\ker(\text{nat}_{\Lambda})$. It follows from [Vollaard and Wedhorn 2011, Corollary 4.7] (see also [Kudla and Rapoport 2014, Proposition 4.6]) that $\ker \mathbb{D}(\varrho_{\Lambda,X}^{-})(S)$ is locally a direct summand of rank 1 of $(\Lambda/\Pi\Lambda) \otimes_{O_F} \mathcal{O}_S$. This induces a map

$$\mathcal{N}_{E,\Lambda}(S) \rightarrow \mathbb{P}(\Lambda/\Pi\Lambda)(S),$$

functorial in S , and the arguments of [Vollaard and Wedhorn 2011, Section 4.7] show that it is an isomorphism. (One easily checks that their results indeed carry over to the relative setting over O_F .) \square

3B. Construction of the closed formal subscheme $\mathcal{N}_E \subseteq \mathcal{N}_E^{\mathrm{naive}}$. We now use a result from [Section 5](#). By [Theorem 5.2](#) and [Remark 5.1 \(2\)](#), there exists a principal polarization $\tilde{\lambda}_{\mathbb{X}} : \mathbb{X} \rightarrow \mathbb{X}^{\vee}$ on $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$, unique up to a scalar in O_E^{\times} , such that the induced Rosati involution is the identity on O_E . Furthermore, for any $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\mathrm{naive}}(S)$, the pullback $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$ is a principal polarization on X .

The next proposition is crucial for the construction of \mathcal{N}_E . Recall the notion of a *hyperbolic* lattice from [Proposition 2.4](#) and the subsequent discussion.

Proposition 3.10. *It is possible to choose $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ and $\tilde{\lambda}_{\mathbb{X}}$ such that*

$$\lambda_{\mathbb{X},1} = \frac{1}{2}(\lambda_{\mathbb{X}} + \tilde{\lambda}_{\mathbb{X}}) \in \mathrm{Hom}(\mathbb{X}, \mathbb{X}^{\vee}).$$

Fix such a choice and let $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\mathrm{naive}}(\bar{k})$. Then, $\frac{1}{2}(\lambda + \tilde{\lambda}) \in \mathrm{Hom}(X, X^{\vee})$ if and only if $(X, \iota, \lambda, \varrho) \in \mathcal{N}_{E,\Lambda}(\bar{k})$ for some hyperbolic lattice $\Lambda \subseteq C$.

Proof. The polarization $\tilde{\lambda}_{\mathbb{X}}$ on \mathbb{X} induces an alternating form $(,)$ on the rational Dieudonné module $N = M_{\mathbb{X}} \otimes_{\check{O}_F} \check{F}$. For all $x, y \in N$, the form $(,)$ satisfies

$$(\mathbf{F}x, y) = (x, \mathbf{V}y)^{\sigma},$$

$$(\Pi x, y) = (x, \Pi y).$$

It induces an \check{E} -alternating form b on N via

$$b(x, y) = \delta((\Pi x, y) + \Pi(x, y)),$$

where $\delta \in \check{O}_F$ is a unit generating the unramified quadratic extension of F , chosen such that $\delta^{\sigma} = -\delta$ and $(1 + \delta)/2 \in \check{O}_F$; see page [348](#). On the other hand, we can describe $(,)$ in terms of b ,

$$(3-12) \quad (x, y) = \mathrm{Tr}_{\check{E}|\check{F}}\left(\frac{1}{2\Pi\delta} \cdot b(x, y)\right).$$

The form b is invariant under $\tau = \Pi \mathbf{V}^{-1}$, since

$$b(\tau x, \tau y) = b(-\mathbf{F}\Pi^{-1}x, \Pi \mathbf{V}^{-1}y) = b(\Pi^{-1}x, \Pi y)^{\sigma} = b(x, y)^{\sigma}.$$

Hence b defines an E -linear alternating form on $C = N^{\tau}$, which we again denote by b . Denote by \langle , \rangle the alternating form on $M_{\mathbb{X}}$ induced by the polarization $\lambda_{\mathbb{X}}$ and let h be the corresponding hermitian form; see [\(3-2\)](#). On $N_{\mathbb{X}}$, we define the alternating form \langle , \rangle_1 by

$$\langle x, y \rangle_1 = \frac{1}{2}(\langle x, y \rangle + (x, y)).$$

This form is integral on $M_{\mathbb{X}}$ if and only if $\lambda_{\mathbb{X},1} = \frac{1}{2}(\lambda_{\mathbb{X}} + \tilde{\lambda}_{\mathbb{X}})$ is a polarization on \mathbb{X} .

We choose $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ such that it corresponds to a unimodular hyperbolic lattice $\Lambda_0 \subseteq (C, h)$ under the identifications of [\(3-7\)](#) and [Lemma 3.6](#). There exists a basis

(e_1, e_2) of Λ_0 such that

$$(3-13) \quad h \hat{=} \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, \quad b \hat{=} \begin{pmatrix} & u \\ -u & \end{pmatrix}$$

for some $u \in E^\times$. Since $\tilde{\lambda}_{\mathbb{X}}$ is principal, the alternating form b is perfect on Λ_0 , thus $u \in O_E^\times$. After rescaling $\tilde{\lambda}_{\mathbb{X}}$, we may assume that $u = 1$. We now have

$$\frac{1}{2}(h(x, y) + b(x, y)) \in O_E,$$

for all $x, y \in \Lambda_0$. Thus $\frac{1}{2}(h + b)$ is integral on $M_{\mathbb{X}} = \Lambda_0 \otimes_{O_E} \check{O}_E$. This implies that

$$\begin{aligned} \langle x, y \rangle_1 &= \frac{1}{2}(\langle x, y \rangle + \langle x, y \rangle) = \frac{1}{2} \operatorname{Tr}_{\check{E}|\check{F}} \left(\frac{1}{2\Pi} \cdot h(x, y) + \frac{1}{2\Pi\delta} \cdot b(x, y) \right) \\ &= \operatorname{Tr}_{\check{E}|\check{F}} \left(\frac{1}{4\Pi} (h(x, y) + b(x, y)) \right) + \operatorname{Tr}_{\check{E}|\check{F}} \left(\frac{1-\delta}{4\Pi\delta} \cdot b(x, y) \right) \in \check{O}_F \end{aligned}$$

for all $x, y \in M_{\mathbb{X}}$. Indeed, in the definition of b , the unit δ has been chosen such that $(1 + \delta)/2 \in \check{O}_F$, so the second summand is in \check{O}_F . The first summand is integral, since $\frac{1}{2}(h + b)$ is integral. It follows that $\lambda_{\mathbb{X},1} = \frac{1}{2}(\lambda_{\mathbb{X}} + \tilde{\lambda}_{\mathbb{X}})$ is a polarization on \mathbb{X} .

Let $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\text{naive}}(\bar{k})$ and assume that $\lambda_1 = \frac{1}{2}(\lambda + \tilde{\lambda}) = \varrho^*(\lambda_{\mathbb{X},1})$ is a polarization on X . Then $\langle \cdot, \cdot \rangle_1$ is integral on the Dieudonné module $M \subseteq N$ of X . By the above calculation, this is equivalent to $\frac{1}{2}(h + b)$ being integral on M . In particular, this implies that

$$h(x, x) = h(x, x) + b(x, x) \in 2\check{O}_F,$$

for all $x \in M$. Let $\Lambda = (M + \tau(M))^\tau$. Then $h(x, x) \in 2O_F$ for all $x \in \Lambda$; hence $\operatorname{Nm}(\Lambda) \subseteq 2O_F$. By [Lemma 2.5](#) and the bound of norm ideals, we have $\operatorname{Nm}(\Lambda) = 2O_F$ and Λ is a hyperbolic lattice. It follows that $(X, \iota, \lambda, \varrho) \in \mathcal{N}_{E,\Lambda'}(\bar{k})$ for some hyperbolic Π^{-1} -modular lattice $\Lambda' \subseteq C$. Indeed, if $M^\tau \subsetneq \Lambda$ then Λ is Π^{-1} -modular and $\Lambda' = \Lambda$. If $M^\tau = \Lambda$ then it is contained in some Π^{-1} -modular hyperbolic lattice Λ' by [Proposition 2.7](#).

Conversely, assume that $(X, \iota, \lambda, \varrho) \in \mathcal{N}_{E,\Lambda}(\bar{k})$ for some hyperbolic lattice $\Lambda \subseteq C$. It suffices to show that $\frac{1}{2}(h + b)$ is integral on Λ . Indeed, it follows that $\frac{1}{2}(h + b)$ is integral on the Dieudonné module M . Thus $\langle \cdot, \cdot \rangle_1$ is integral on M and this is equivalent to $\lambda_1 = \frac{1}{2}(\lambda + \tilde{\lambda}) \in \operatorname{Hom}(X, X^\vee)$.

Let $\Lambda' \subseteq C$ be the Π^{-1} -modular lattice generated by e_1 and $\Pi^{-1}e_2$, where (e_1, e_2) is the basis of the lattice Λ_0 corresponding to the framing object $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$. By [\(3-13\)](#), h and b have the following form with respect to the basis $(e_1, \Pi^{-1}e_2)$,

$$h \hat{=} \begin{pmatrix} & -\Pi^{-1} \\ \Pi^{-1} & \end{pmatrix}, \quad b \hat{=} \begin{pmatrix} & \Pi^{-1} \\ -\Pi^{-1} & \end{pmatrix}.$$

In particular, Λ' is hyperbolic and $\frac{1}{2}(h + b)$ is integral on Λ' . By [Proposition 2.4](#), there exists an automorphism $g \in \operatorname{SU}(C, h)$ mapping Λ onto Λ' . Since $\det g = 1$,

the alternating form b is invariant under g . It follows that $\frac{1}{2}(h + b)$ is also integral on Λ . \square

From now on, we assume $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ and $\tilde{\lambda}_{\mathbb{X}}$ chosen in a way such that

$$\lambda_{\mathbb{X},1} = \frac{1}{2}(\lambda_{\mathbb{X}} + \tilde{\lambda}_{\mathbb{X}}) \in \mathrm{Hom}(\mathbb{X}, \mathbb{X}^{\vee}).$$

Note that this determines the polarization $\tilde{\lambda}_{\mathbb{X}}$ up to a scalar in $1 + 2O_E$. If we replace $\tilde{\lambda}_{\mathbb{X}}$ by $\tilde{\lambda}'_{\mathbb{X}} = \tilde{\lambda}_{\mathbb{X}} \circ \iota_{\mathbb{X}}(1 + 2u)$ for some $u \in O_E$, then $\lambda'_{\mathbb{X},1} = \lambda_{\mathbb{X},1} + \tilde{\lambda}_{\mathbb{X}} \circ \iota_{\mathbb{X}}(u)$.

We can now formulate the straightening condition.

Definition 3.11. Let $S \in \mathrm{Nilp}_{\check{O}_F}$. An object $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\mathrm{naive}}(S)$ satisfies the *straightening condition* if

$$(3-14) \quad \lambda_1 \in \mathrm{Hom}(X, X^{\vee}),$$

where $\lambda_1 = \frac{1}{2}(\lambda + \tilde{\lambda}) = \varrho^*(\lambda_{\mathbb{X},1})$.

This definition is clearly independent of the choice of the polarization $\tilde{\lambda}_{\mathbb{X}}$. We define \mathcal{N}_E as the functor that maps $S \in \mathrm{Nilp}_{\check{O}_F}$ to the set of all tuples $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\mathrm{naive}}(S)$ that satisfy the straightening condition. By [Rapoport and Zink 1996, Proposition 2.9], \mathcal{N}_E is representable by a closed formal subscheme of $\mathcal{N}_E^{\mathrm{naive}}$.

Remark 3.12. The reduced locus of \mathcal{N}_E can be written as

$$(\mathcal{N}_E)_{\mathrm{red}} = \bigcup_{\Lambda \subseteq C} \mathcal{N}_{E,\Lambda} \simeq \bigcup_{\Lambda \subseteq C} \mathbb{P}(\Lambda / \Pi\Lambda),$$

where we take the unions over all *hyperbolic* Π^{-1} -modular lattices $\Lambda \subseteq C$. By Proposition 2.7 and Lemma 3.7, each projective line contains $q + 1$ points corresponding to unimodular lattices and there are two lines intersecting in each such point. Recall from Remark 3.8 (1) that there exist nonhyperbolic Π^{-1} -modular lattices $\Lambda \subseteq C$; thus we have $\mathcal{N}_E(\bar{k}) \neq \mathcal{N}_E^{\mathrm{naive}}(\bar{k})$, and in particular $(\mathcal{N}_E)_{\mathrm{red}} \neq (\mathcal{N}_E^{\mathrm{naive}})_{\mathrm{red}}$.

Remark 3.13. As has been pointed out to the author by A. Genestier, the straightening condition is not trivial on the rigid-analytic generic fiber of $\mathcal{N}_E^{\mathrm{naive}}$. However, we can show that it is open and closed. Since a proper study of the generic fiber would go beyond the scope of this paper, we restrain ourselves to indications rather than complete proofs.

Let C be an algebraically closed extension of F and \mathcal{O}_C its ring of integers. Take a point $x = (X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\mathrm{naive}}(\mathcal{O}_C)$ and consider its 2-adic Tate module $T_2(x)$. It is a free O_E -module of rank 2 and λ endows $T_2(x)$ with a perfect (nonsplit) hermitian form h . If $x \in \mathcal{N}_E(\mathcal{O}_C)$, then the straightening condition implies that $(T_2(x), h)$ is a lattice with minimal norm² $\mathrm{Nm}(T_2(x))$ in the vector space $V_2(x) = T_2(x) \otimes_{O_E} E$ (see Proposition 2.4 and [Jacobowitz 1962]). But $V_2(x)$ also contains

²Calling this lattice “hyperbolic” doesn’t make much sense here since it is anisotropic.

self-dual lattices with nonminimal norm ideal. Let $\Lambda \subseteq V_2(x)$ be such a lattice with $\text{Nm}(\Lambda) \neq \text{Nm}(T_2(x))$. Let Λ' be the intersection of $T_2(x)$ and Λ in $V_2(x)$. The inclusions $\Lambda' \hookrightarrow \Lambda$ and $\Lambda' \hookrightarrow T_2(x)$ define canonically a formal O_F -module Y with $T_2(Y) = \Lambda'$ and a quasi-isogeny $\varphi : X \rightarrow Y$. By inheriting all data, Y becomes a point in $\mathcal{N}_E^{\text{naive}}(\mathcal{O}_C)$ that does not satisfy the straightening condition.

To see that the straightening condition is open and closed on the generic fiber, consider the universal formal O_F -module $\mathcal{X} = (\mathcal{X}, \iota_{\mathcal{X}}, \lambda_{\mathcal{X}})$ over $\mathcal{N}_E^{\text{naive}}$ and let $T_2(\mathcal{X})$ be its Tate module. Then $T_2(\mathcal{X})$ is a locally constant sheaf over $\mathcal{N}_E^{\text{naive,rig}}$ with respect to the étale topology. The polarization $\lambda_{\mathcal{X}}$ defines a hermitian form h on $T_2(\mathcal{X})$. Since $T_2(\mathcal{X})$ is a locally constant sheaf, the norm ideal $\text{Nm}(T_2(\mathcal{X}))$ with respect to h (see [Proposition 2.4](#)) is locally constant as well. Hence the locus where $\text{Nm}(T_2(\mathcal{X}))$ is minimal is open and closed in $\mathcal{N}_E^{\text{naive,rig}}$. But this is exactly $\mathcal{N}_E^{\text{rig}} \subseteq \mathcal{N}_E^{\text{naive,rig}}$.

3C. The isomorphism to the Drinfeld moduli problem. We now recall the Drinfeld moduli problem \mathcal{M}_{Dr} on $\text{Nilp}_{\check{O}_F}$. Let B be the quaternion division algebra over F and O_B its ring of integers. Let $S \in \text{Nilp}_{\check{O}_F}$. Then $\mathcal{M}_{Dr}(S)$ is the set of equivalence classes of objects (X, ι_B, ϱ) , where

- X is a formal O_F -module over S of dimension 2 and height 4;
- $\iota_B : O_B \rightarrow \text{End}(X)$ is an action of O_B on X satisfying the *special* condition, i.e., $\text{Lie } X$ is, locally on S , a free $(\mathcal{O}_S \otimes_{O_F} O_F^{(2)})$ -module of rank 1, where $O_F^{(2)} \subseteq O_B$ is any embedding of the unramified quadratic extension of O_F into O_B (cf. [\[Boutot and Carayol 1991\]](#));
- $\varrho : X \times_S \bar{S} \rightarrow \mathbb{X} \times_{\text{Spec } \bar{k}} \bar{S}$ is an O_B -linear quasi-isogeny of height 0 to a fixed framing object $(\mathbb{X}, \iota_{\mathbb{X}}) \in \mathcal{M}_{Dr}(\bar{k})$.

Such a framing object exists and is unique up to isogeny. By a proposition of Drinfeld, cf. [\[Boutot and Carayol 1991, p. 138\]](#), there always exist polarizations on these objects, as follows:

Proposition 3.14 [\[Drinfeld 1976\]](#). *Let $\Pi \in O_B$ a uniformizer with $\Pi^2 \in O_F$ and let $b \mapsto b'$ be the standard involution of B . Then $b \mapsto b^* = \Pi b' \Pi^{-1}$ is another involution on B .*

- (1) *There exists a principal polarization $\lambda_{\mathbb{X}} : \mathbb{X} \rightarrow \mathbb{X}^{\vee}$ on \mathbb{X} with associated Rosati involution $b \mapsto b^*$. It is unique up to a scalar in O_F^{\times} .*
- (2) *Let $\lambda_{\mathbb{X}}$ be as in (1). For $(X, \iota_B, \varrho) \in \mathcal{M}_{Dr}(S)$, there exists a unique principal polarization*

$$\lambda : X \rightarrow X^{\vee}$$

with Rosati involution $b \mapsto b^$ such that $\varrho^*(\lambda_{\mathbb{X}}) = \lambda$ on \bar{S} .*

We now relate \mathcal{M}_{Dr} and \mathcal{N}_E . For this, we fix an embedding $E \hookrightarrow B$. Any choice of a uniformizer $\Pi \in O_E$ with $\Pi^2 \in O_F$ induces the same involution $b \mapsto b^* = \Pi b' \Pi^{-1}$ on B .

For the framing object $(\mathbb{X}, \iota_{\mathbb{X}})$ of \mathcal{M}_{Dr} , let $\lambda_{\mathbb{X}}$ be a polarization associated to this involution by [Proposition 3.14 \(1\)](#). Denote by $\iota_{\mathbb{X}, E}$ the restriction of $\iota_{\mathbb{X}}$ to $O_E \subseteq O_B$. For any object $(X, \iota_B, \varrho) \in \mathcal{M}_{Dr}(S)$, let λ be the polarization with Rosati involution $b \mapsto b^*$ that satisfies $\varrho^*(\lambda_{\mathbb{X}}) = \lambda$; see [Proposition 3.14 \(2\)](#). Let ι_E be the restriction of ι_B to O_E .

Lemma 3.15. *$(\mathbb{X}, \iota_{\mathbb{X}, E}, \lambda_{\mathbb{X}})$ is a framing object for $\mathcal{N}_E^{\mathrm{naive}}$. Furthermore, the map*

$$(X, \iota_B, \varrho) \mapsto (X, \iota_E, \lambda, \varrho)$$

induces a closed immersion of formal schemes

$$\eta : \mathcal{M}_{Dr} \hookrightarrow \mathcal{N}_E^{\mathrm{naive}}.$$

Proof. There are two things to check: that $\mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ contains $\mathrm{SU}(C, h)$ as a closed subgroup and that ι_E satisfies the Kottwitz condition. Indeed, once these two assertions hold, we can take $(\mathbb{X}, \iota_{\mathbb{X}, E}, \lambda_{\mathbb{X}})$ as a framing object for $\mathcal{N}_E^{\mathrm{naive}}$ and the morphism η is well-defined. For any $S \in \mathrm{Nilp}_{\check{O}_F}$, the map $\eta(S)$ is injective, because (X, ι_B, ϱ) and $(X', \iota'_B, \varrho') \in \mathcal{M}_{Dr}(S)$ map to the same point in $\mathcal{N}_E^{\mathrm{naive}}(S)$ under η if and only if the quasi-isogeny $\varrho' \circ \varrho$ on \bar{S} lifts to an isomorphism on S , i.e., if and only if (X, ι_B, ϱ) and (X', ι'_B, ϱ') define the same point in $\mathcal{M}_{Dr}(S)$. The functor

$$F : S \mapsto \{(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\mathrm{naive}}(S) \mid \iota \text{ extends to an } O_B\text{-action}\}$$

is pro-representable by a closed formal subscheme of $\mathcal{N}_E^{\mathrm{naive}}$ by [\[Rapoport and Zink 1996, Proposition 2.9\]](#). Now, the formal subscheme $\eta(\mathcal{M}_{Dr}) \subseteq F$ is given by the special condition. But the special condition is open and closed (see [\[Rapoport and Zink 2017, p. 7\]](#)), thus η is a closed embedding.

It remains to show the two assertions from the beginning of this proof. We first check the condition on $\mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$. Let $G_{(\mathbb{X}, \iota_{\mathbb{X}})}$ be the group of O_B -linear quasi-isogenies $\varphi : (\mathbb{X}, \iota_{\mathbb{X}}) \rightarrow (\mathbb{X}, \iota_{\mathbb{X}})$ of height 0 such that the induced homomorphism of Dieudonné modules has determinant 1. Then we have (noncanonical) isomorphisms $G_{(\mathbb{X}, \iota_{\mathbb{X}})} \simeq \mathrm{SL}_{2, F}$ and $\mathrm{SL}_{2, F} \simeq \mathrm{SU}(C, h)$, since h is split. The uniqueness of the polarization $\lambda_{\mathbb{X}}$ (up to a scalar in O_F^\times) implies that $G_{(\mathbb{X}, \iota_{\mathbb{X}})} \subseteq \mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$. This is a closed embedding of linear algebraic groups over F , since a quasi-isogeny $\varphi \in \mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ lies in $G_{(\mathbb{X}, \iota_{\mathbb{X}})}$ if and only if it is O_B -linear and has determinant 1, and these are closed conditions on $\mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$.

Finally, the special condition implies the Kottwitz condition for any element $b \in O_B$ (see [\[Rapoport and Zink 2017, Proposition 5.8\]](#)), i.e., the characteristic

polynomial for the action of $\iota(b)$ on $\text{Lie } X$ is

$$\text{char}(\text{Lie } X, T \mid \iota(b)) = (T - b)(T - b'),$$

where the right-hand side is a polynomial in $\mathcal{O}_S[T]$ via the structure homomorphism $\mathcal{O}_F \hookrightarrow \check{\mathcal{O}}_F \rightarrow \mathcal{O}_S$. From this, the second assertion follows. \square

Let $\mathcal{O}_F^{(2)} \subseteq \mathcal{O}_B$ be an embedding such that conjugation with Π induces the nontrivial Galois action on $\mathcal{O}_F^{(2)}$, as in [Lemma 2.3 \(1\)](#). Fix a generator $\gamma = (1 + \delta)/2$ of $\mathcal{O}_F^{(2)}$ with $\delta^2 \in \mathcal{O}_F^\times$. On $(\mathbb{X}, \iota_{\mathbb{X}})$, the principal polarization $\tilde{\lambda}_{\mathbb{X}}$ given by

$$\tilde{\lambda}_{\mathbb{X}} = \lambda_{\mathbb{X}} \circ \iota_{\mathbb{X}}(\delta)$$

has a Rosati involution that induces the identity on \mathcal{O}_E . For any $(X, \iota_B, \varrho) \in \mathcal{M}_{Dr}(S)$, we set $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}}) = \lambda \circ \iota_B(\delta)$. The tuple $(X, \iota_E, \lambda, \varrho) = \eta(X, \iota_B, \varrho)$ satisfies the straightening condition [\(3-14\)](#), since

$$\lambda_1 = \frac{1}{2}(\lambda + \tilde{\lambda}) = \lambda \circ \iota_B(\gamma) \in \text{Hom}(X, X^\vee).$$

In particular, the tuple $(\mathbb{X}, \iota_{\mathbb{X}, E}, \lambda_{\mathbb{X}})$ is a framing object of \mathcal{N}_E and η induces a natural transformation

$$(3-15) \quad \eta : \mathcal{M}_{Dr} \hookrightarrow \mathcal{N}_E.$$

Note that this map does not depend on the above choices, as \mathcal{N}_E is a closed formal subscheme of $\mathcal{N}_E^{\text{naive}}$.

Theorem 3.16. *$\eta : \mathcal{M}_{Dr} \rightarrow \mathcal{N}_E$ is an isomorphism of formal schemes.*

We will first prove this on \bar{k} -valued points:

Lemma 3.17. *η induces a bijection $\eta(\bar{k}) : \mathcal{M}_{Dr}(\bar{k}) \rightarrow \mathcal{N}_E(\bar{k})$.*

Proof. We can identify the \bar{k} -valued points of \mathcal{M}_{Dr} with a subset $\mathcal{M}_{Dr}(\bar{k}) \subseteq \mathcal{N}_E^{\text{naive}}(\bar{k})$. The rational Dieudonné module N of \mathbb{X} is equipped with an action of B . Fix an embedding $F^{(2)} \hookrightarrow B$ as in [Lemma 2.3 \(1\)](#). This induces a $\mathbb{Z}/2$ -grading $N = N_0 \oplus N_1$ of N , where

$$N_0 = \{x \in N \mid \iota(a)x = ax \text{ for all } a \in F^{(2)}\},$$

$$N_1 = \{x \in N \mid \iota(a)x = \sigma(a)x \text{ for all } a \in F^{(2)}\}$$

for a fixed embedding $F^{(2)} \hookrightarrow \check{F}$. The operators V and F have degree 1 with respect to this decomposition. Recall that λ has Rosati involution $b \mapsto \Pi b' \Pi^{-1}$ on \mathcal{O}_B which restricts to the identity on $\mathcal{O}_F^{(2)}$. The subspaces N_0 and N_1 are therefore orthogonal with respect to $\langle \cdot, \cdot \rangle$.

Under the identification [\(3-7\)](#), a lattice $M \in \mathcal{M}_{Dr}(\bar{k})$ respects this decomposition, i.e., $M = M_0 \oplus M_1$ with $M_i = M \cap N_i$. Furthermore it satisfies the special condition

$$\dim M_0 / V M_1 = \dim M_1 / V M_0 = 1.$$

We already know that $\mathcal{M}_{Dr}(\bar{k}) \subseteq \mathcal{N}_E(\bar{k})$, so let us assume $M \in \mathcal{N}_E(\bar{k})$. We want to show that $M \in \mathcal{M}_{Dr}(\bar{k})$, i.e., that the lattice M is stable under the action of O_B on N and satisfies the special condition. It is stable under the O_B -action if and only if $M = M_0 \oplus M_1$ for $M_i = M \cap N_i$. Let $y \in M$ and $y = y_0 + y_1$ with $y_i \in N_i$. For any $x \in M$, we have

$$(3-16) \quad \langle x, y \rangle = \langle x, y_0 \rangle + \langle x, y_1 \rangle \in \check{O}_F.$$

We can assume that $\lambda_{\mathbb{X},1} = \lambda_{\mathbb{X}} \circ \iota_B(\gamma)$ with $\gamma \in O_F^{(2)}$ under our fixed embedding $F^{(2)} \hookrightarrow B$. Recall that $\gamma^\sigma = 1 - \gamma$ from page 348. Let $\langle \cdot, \cdot \rangle_1$ be the alternating form on M induced by $\lambda_{\mathbb{X},1}$. Then,

$$(3-17) \quad \langle x, y \rangle_1 = \gamma \cdot \langle x, y_0 \rangle + (1 - \gamma) \cdot \langle x, y_1 \rangle \in \check{O}_F.$$

From (3-16) and (3-17), it follows that $\langle x, y_0 \rangle$ and $\langle x, y_1 \rangle$ lie in \check{O}_F . Since $x \in M$ was arbitrary and $M = M^\vee$, this gives $y_0, y_1 \in M$. Hence M respects the decomposition of N and is stable under the action of O_B .

It remains to show that M satisfies the special condition: The alternating form $\langle \cdot, \cdot \rangle$ is perfect on M , thus the restrictions to M_0 and M_1 are perfect as well. If M is not special, we have $M_i = \mathbf{V} M_{i+1}$ for some $i \in \{0, 1\}$. But then, $\langle \cdot, \cdot \rangle$ cannot be perfect on M_i . In fact, for any $x, y \in M_{i+1}$,

$$\langle \mathbf{V}x, \mathbf{V}y \rangle^\sigma = \langle \mathbf{F}\mathbf{V}x, y \rangle = \pi_0 \cdot \langle x, y \rangle \in \pi_0 \check{O}_F.$$

Thus M is indeed special, i.e., $M \in \mathcal{M}_{Dr}(\bar{k})$, and this finishes the proof of the lemma. \square

Proof of Theorem 3.16. We already know that η is a closed embedding

$$\eta : \mathcal{M}_{Dr} \hookrightarrow \mathcal{N}_E.$$

Let $(\mathbb{X}, \iota_{\mathbb{X}})$ be the framing object of \mathcal{M}_{Dr} and choose an embedding $O_F^{(2)} \subseteq O_B$ and a generator γ of $O_F^{(2)}$ as in Lemma 2.3 (1). We take $(\mathbb{X}, \iota_{\mathbb{X},E}, \lambda_{\mathbb{X}})$ as a framing object for \mathcal{N}_E and set $\tilde{\lambda}_{\mathbb{X}} = \lambda_{\mathbb{X}} \circ \iota_{\mathbb{X}}(\delta)$.

Let $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E(S)$ and $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$. We have

$$\varrho^{-1} \circ \iota_{\mathbb{X}}(\gamma) \circ \varrho = \varrho^{-1} \circ \lambda_{\mathbb{X}}^{-1} \circ \lambda_{\mathbb{X},1} \circ \varrho = \lambda^{-1} \circ \lambda_1 \in \mathrm{End}(X),$$

where $\lambda_{\mathbb{X},1} = \frac{1}{2}(\lambda_{\mathbb{X}} + \tilde{\lambda}_{\mathbb{X}})$ and $\lambda_1 = \frac{1}{2}(\lambda + \tilde{\lambda})$. Since $O_B = O_F[\Pi, \gamma]$, this induces an O_B -action ι_B on X and makes ϱ an O_B -linear quasi-isogeny. We have to check that (X, ι_B, ϱ) satisfies the special condition.

Recall that the special condition is open and closed (see [Rapoport and Zink 2017, p. 7]), so η is an open and closed embedding. Furthermore, $\eta(\bar{k})$ is bijective and the reduced loci $(\mathcal{M}_{Dr})_{\mathrm{red}}$ and $(\mathcal{N}_E)_{\mathrm{red}}$ are locally of finite type over $\mathrm{Spec} \bar{k}$. Hence η induces an isomorphism on reduced subschemes. But any open and closed

embedding of formal schemes, that is, an isomorphism on the reduced subschemes, is already an isomorphism. \square

4. The moduli problem in the case R-U

Let $E|F$ be a quadratic extension of type R-U, generated by a uniformizer Π satisfying an Eisenstein equation of the form $\Pi^2 - t\Pi + \pi_0 = 0$ where $t \in O_F$ and $\pi_0|t|2$. Let O_F and O_E be the rings of integers of F and E . We have $O_E = O_F[\Pi]$. As in the case R-P, let k be the common residue field, \bar{k} an algebraic closure, \check{F} the completion of the maximal unramified extension with ring of integers $\check{O}_F = W_{O_F}(\bar{k})$ and σ the lift of the Frobenius in $\text{Gal}(\bar{k}|k)$ to $\text{Gal}(\check{O}_F|O_F)$.

4A. The naive moduli problem. Let $S \in \text{Nilp}_{\check{O}_F}$. Consider tuples (X, ι, λ) , where

- X is a formal O_F -module over S of dimension 2 and height 4.
- $\iota : O_E \rightarrow \text{End}(X)$ is an action of O_E on X satisfying the *Kottwitz condition*: The characteristic polynomial of $\iota(\alpha)$ for some $\alpha \in O_E$ is given by

$$\text{char}(\text{Lie } X, T \mid \iota(\alpha)) = (T - \alpha)(T - \bar{\alpha}).$$

Here $\alpha \mapsto \bar{\alpha}$ is the Galois conjugation of $E|F$ and the right-hand side is a polynomial in $\mathcal{O}_S[T]$ via the structure morphism $O_F \hookrightarrow \check{O}_F \rightarrow \mathcal{O}_S$.

- $\lambda : X \rightarrow X^\vee$ is a polarization on X with kernel $\ker \lambda = X[\Pi]$, where $X[\Pi]$ is the kernel of $\iota(\Pi)$. Further we demand that the Rosati involution of λ satisfies $\iota(\alpha)^* = \iota(\bar{\alpha})$ for all $\alpha \in O_E$.

We define quasi-isogenies $\varphi : (X, \iota, \lambda) \rightarrow (X', \iota', \lambda')$ and the group $\text{QIsog}(X, \iota, \lambda)$ as in [Definition 3.1](#).

Proposition 4.1. *Up to isogeny, there exists exactly one such tuple $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ over $S = \text{Spec } \bar{k}$ under the condition that the group $\text{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ contains a closed subgroup isomorphic to $\text{SU}(C, h)$ for a 2-dimensional E -vector space C with split $E|F$ -hermitian form h .*

Remark 4.2. As in the case R-P, we have $\text{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}}) \cong \text{U}(C, h)$ for $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ as in the proposition.

Proof of Proposition 4.1. We first show uniqueness of the object. Let $(X, \iota, \lambda)/\text{Spec } \bar{k}$ be a tuple as in the proposition and consider its rational Dieudonné module N_X . This is a 4-dimensional vector space over \check{F} equipped with an action of E and an alternating form \langle, \rangle such that

$$(4-1) \quad \langle x, \Pi y \rangle = \langle \bar{\Pi} x, y \rangle$$

for all $x, y \in N_X$. Let $\check{E} = \check{F} \otimes_F E$. We can see N_X as 2-dimensional vector space over \check{E} with a hermitian form h given by

$$(4-2) \quad h(x, y) = \langle \Pi x, y \rangle - \bar{\Pi} \langle x, y \rangle.$$

Let F and V be the σ -linear Frobenius and the σ^{-1} -linear Verschiebung on N_X . We have $FV = VF = \pi_0$ and, since $\langle \cdot, \cdot \rangle$ comes from a polarization,

$$\langle Fx, y \rangle = \langle x, Vy \rangle^\sigma.$$

Consider the σ -linear operator $\tau = \Pi V^{-1} = F\bar{\Pi}^{-1}$. The hermitian form h is invariant under τ :

$$h(\tau x, \tau y) = h(F\bar{\Pi}^{-1}x, \Pi V^{-1}y) = h(Fx, V^{-1}y) = h(x, y)^\sigma.$$

From the condition on $\mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ it follows that N_X is isotypical of slope $\frac{1}{2}$ and thus the slopes of τ are all zero. Let $C = N_X^\tau$. This is a 2-dimensional vector space over E with $N_X = C \otimes_E \check{E}$ and h induces an $E|F$ -hermitian form on C . A priori, there are two possibilities for (C, h) , either h is split or nonsplit. The group $\mathrm{U}(C, h)$ of automorphisms is isomorphic to $\mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$. But the unitary groups for h split and h nonsplit are not isomorphic and do not contain each other as a closed subgroup. Thus the condition on $\mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ implies that h is split.

Assume we are given two different objects (X, ι, λ) and (X', ι', λ') as in the proposition. Then there is an isomorphism between the spaces (C, h) and (C', h') extending to an isomorphism of N_X and $N_{X'}$ respecting all structure. This corresponds to a quasi-isogeny $\varphi : (X, \iota, \lambda) \rightarrow (X', \iota', \lambda')$.

Now we prove the existence of $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$. We start with a Π -modular lattice Λ in a 2-dimensional vector space (C, h) over E with split hermitian form. Then $M = \Lambda \otimes_{O_E} \check{O}_E$ is an \check{O}_E -lattice in $N = C \otimes_E \check{E}$. The σ -linear operator $\tau = 1 \otimes \sigma$ on N has slopes are all 0. We can extend h to N such that

$$h(\tau x, \tau y) = h(x, y)^\sigma,$$

for all $x, y \in N$. The operators F and V are given by $\tau = \Pi V^{-1} = F\bar{\Pi}^{-1}$. Finally, the alternating form $\langle \cdot, \cdot \rangle$ is defined via

$$\langle x, y \rangle = \mathrm{Tr}_{\check{E}|\check{F}} \left(\frac{1}{t\vartheta} \cdot h(x, y) \right),$$

for $x, y \in N$. The lattice $M \subseteq N$ is the Dieudonné module of $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$. We leave it to the reader to check that this is indeed an object as considered above. \square

We fix such an object $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ over $\mathrm{Spec} \bar{k}$ from the proposition. We define the functor $\mathcal{N}_E^{\mathrm{naive}}$ on $\mathrm{Nilp}_{\check{O}_F}$ as in [Definition 3.4](#).

Remark 4.3. $\mathcal{N}_E^{\text{naive}}$ is pro-representable by a formal scheme, formally locally of finite type over $\check{\mathcal{O}}_F$; cf. [Rapoport and Zink 1996, Theorem 3.25].

We now study the \bar{k} -valued points of the space $\mathcal{N}_E^{\text{naive}}$. Let $N = N_{\mathbb{X}}$ be the rational Dieudonné module of $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$. This is a 4-dimensional vector space over \check{F} , equipped with an action of E , with two operators F and V and an alternating form $\langle \cdot, \cdot \rangle$.

Let $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\text{naive}}(\bar{k})$. This corresponds to an $\check{\mathcal{O}}_F$ -lattice $M = M_X \subseteq N$ which is stable under the actions of F , V and O_E . The condition on the kernel of λ implies that $M = \Pi M^\vee$ for

$$M^\vee = \{x \in N \mid \langle x, y \rangle \in \check{\mathcal{O}}_F \text{ for all } y \in M\}.$$

The alternating form $\langle \cdot, \cdot \rangle$ induces an $\check{E}|\check{F}$ -hermitian form h on N , seen as a 2-dimensional vector space over \check{E} (see (4-2)):

$$h(x, y) = \langle \Pi x, y \rangle - \bar{\Pi} \langle x, y \rangle.$$

We can recover the form $\langle \cdot, \cdot \rangle$ from h via

$$(4-3) \quad \langle x, y \rangle = \text{Tr}_{\check{E}|\check{F}} \left(\frac{1}{t\vartheta} \cdot h(x, y) \right).$$

Since the inverse different of $E|F$ is $\mathfrak{D}_{E|F}^{-1} = \frac{1}{t} O_E$ (see Lemma 2.2), this implies that M is Π -modular with respect to h , as $\check{\mathcal{O}}_E$ -lattice in N . We denote the dual of M with respect to h by M^\sharp . There is a natural bijection

$$(4-4) \quad \mathcal{N}_E^{\text{naive}}(\bar{k}) = \{\check{\mathcal{O}}_E\text{-lattices } M \subseteq N \mid M = \Pi M^\sharp, \pi_0 M \subseteq VM \subseteq M\}.$$

Recall that $\tau = \Pi V^{-1}$ is a σ -linear operator on N with slopes all 0. Further $C = N^\tau$ is a 2-dimensional E -vector space with hermitian form h .

Lemma 4.4. *Let $M \in \mathcal{N}_E^{\text{naive}}(\bar{k})$. Then:*

- (1) $M + \tau(M)$ is τ -stable.
- (2) Either M is τ -stable and $\Lambda_1 = M^\tau \subseteq C$ is Π -modular with respect to h , or M is not τ -stable and then $\Lambda_0 = (M + \tau(M))^\tau \subseteq C$ is unimodular.

The proof is the same as that of [Kudla and Rapoport 2014, Lemma 3.2]. We identify N with $C \otimes_E \check{E}$. For any τ -stable lattice $M \in \mathcal{N}_E^{\text{naive}}(\bar{k})$, we have $M = \Lambda_1 \otimes_{O_E} \check{\mathcal{O}}_E$. If $M \in \mathcal{N}_E^{\text{naive}}(\bar{k})$ is not τ -stable, there is an inclusion $M \subseteq \Lambda_0 \otimes_{O_E} \check{\mathcal{O}}_E$ of index 1. Recall from Proposition 2.4 that the isomorphism class of a Π -modular or unimodular lattice $\Lambda \subseteq C$ is determined by the norm ideal

$$\text{Nm}(\Lambda) = \langle \{h(x, x) \mid x \in \Lambda\} \rangle.$$

There are always at least two types of unimodular lattices. However, not all of them appear in the description of $\mathcal{N}_E^{\text{naive}}(\bar{k})$.

Lemma 4.5. (1) *Let $\Lambda \subseteq C$ be a unimodular lattice with $\mathrm{Nm}(\Lambda) \subseteq \pi_0 O_F$. There is an injection*

$$i_\Lambda : \mathbb{P}(\Lambda / \Pi\Lambda)(\bar{k}) \hookrightarrow \mathcal{N}_E^{\mathrm{naive}}(\bar{k}),$$

that maps a line $\ell \subseteq \Lambda / \Pi\Lambda \otimes_k \bar{k}$ to its inverse image under the canonical projection

$$\Lambda \otimes_{O_E} \check{O}_E \rightarrow \Lambda / \Pi\Lambda \otimes_k \bar{k}.$$

The k -valued points $\mathbb{P}(\Lambda / \Pi\Lambda)(k) \subseteq \mathbb{P}(\Lambda / \Pi\Lambda)(\bar{k})$ are mapped to τ -invariant Dieudonné modules $M \subseteq \Lambda \otimes_{O_E} \check{O}_E$ under this embedding.

(2) *Identify $\mathbb{P}(\Lambda / \Pi\Lambda)(\bar{k})$ with its image under i_Λ . The set $\mathcal{N}_E^{\mathrm{naive}}(\bar{k})$ can be written as*

$$\mathcal{N}_E^{\mathrm{naive}}(\bar{k}) = \bigcup_{\Lambda \subseteq C} \mathbb{P}(\Lambda / \Pi\Lambda)(\bar{k}),$$

where the union is taken over all lattices $\Lambda \subseteq C$ with $\mathrm{Nm}(\Lambda) \subseteq \pi_0 O_F$.

Proof. Let $\Lambda \subseteq C$ be a unimodular lattice. For any line $\ell \in \mathbb{P}(\Lambda / \Pi\Lambda)(\bar{k})$, denote its preimage in $\Lambda \otimes \check{O}_E$ by M . The inclusion $M \subseteq \Lambda \otimes \check{O}_E$ has index 1 and M is an \check{O}_E -lattice with $\Pi(\Lambda \otimes \check{O}_E) \subseteq M$. Furthermore $\Lambda \otimes \check{O}_E$ is τ -invariant by construction, hence $\Pi(\Lambda \otimes \check{O}_E) = V(\Lambda \otimes \check{O}_E) = F(\Lambda \otimes \check{O}_E)$. It follows that M is stable under the actions of F and V . Thus $M \in \mathcal{N}_E^{\mathrm{naive}}(\bar{k})$ if and only if $M = \Pi M^\sharp$. The hermitian form h induces a symmetric form s on $\Lambda / \Pi\Lambda$. Now M is Π -modular if and only if it is the preimage of an isotropic line $\ell \subseteq \Lambda / \Pi\Lambda \otimes \bar{k}$. Note that s is also antisymmetric since we are in characteristic 2.

We first consider the case $\mathrm{Nm}(\Lambda) \subseteq \pi_0 O_F$. We can find a basis of Λ such that h has the form

$$H_\Lambda = \begin{pmatrix} x & 1 \\ 1 & \end{pmatrix}, \quad x \in \pi_0 O_F;$$

see (2-4). It follows that the induced form s is even alternating (because $x \equiv 0 \pmod{\pi_0}$). Hence any line in $\Lambda / \Pi\Lambda \otimes \bar{k}$ is isotropic. This implies that i_Λ is well-defined, proving part (1) of the lemma.

Now assume that $\mathrm{Nm}(\Lambda) = O_F$. There is a basis (e_1, e_2) of Λ such that h is represented by

$$H_\Lambda = \begin{pmatrix} 1 & 1 \\ 1 & \end{pmatrix}.$$

The induced form s is given by the same matrix and $\ell = \bar{k} \cdot e_2$ is the only isotropic line in $\Lambda / \Pi\Lambda$. Since ℓ is already defined over k , the corresponding lattice $M \in \mathcal{N}_E^{\mathrm{naive}}(\bar{k})$ is of the form $M = \Lambda_1 \otimes \check{O}_E$ for a Π -modular lattice $\Lambda_1 \subseteq \Lambda$. But, by Proposition 2.8, any Π -modular lattice in C is contained in a unimodular lattice Λ' with $\mathrm{Nm}(\Lambda') \subseteq \pi_0 O_F$.




Figure 2. The reduced locus of $\mathcal{N}_E^{\text{naive}}$ for an R-U extension $E|F$ where e and f are the ramification index and the inertia degree of $F|\mathbb{Q}_2$ and $v(t)$ is the π_0 -adic valuation of t . We always have $1 \leq v(t) \leq e$. The solid lines lie in $\mathcal{N}_E \subseteq \mathcal{N}_E^{\text{naive}}$.

It follows that we can write $\mathcal{N}_E^{\text{naive}}(\bar{k})$ as a union

$$\mathcal{N}_E^{\text{naive}}(\bar{k}) = \bigcup_{\Lambda \subseteq C} \mathbb{P}(\Lambda / \Pi \Lambda)(\bar{k}),$$

where the union is taken over all unimodular lattices $\Lambda \subseteq C$ with $\text{Nm}(\Lambda) \subseteq \pi_0 O_F$. This shows the second part of the lemma. □

Remark 4.6. We can use Proposition 2.8 to describe the intersection behavior of the projective lines in $\mathcal{N}_E^{\text{naive}}(\bar{k})$. A τ -invariant point $M \in \mathcal{N}_E^{\text{naive}}(\bar{k})$ corresponds to the Π -modular lattice $\Lambda_1 = M^\tau \subseteq C$. If $\text{Nm}(\Lambda_1) \subseteq \pi_0^2 O_F$, there are $q + 1$ lines going through M . If $\text{Nm}(\Lambda_1) = \pi_0 O_F$, the point M is contained in one or two lines, depending on whether Λ_1 is hyperbolic or not; see parts (3) and (4) of Proposition 2.8. The former case (i.e., Λ_1 is hyperbolic) appears if and only if $\pi_0 O_F = \text{Nm}(\Lambda_1) = t O_F$ (see Lemma 2.5). This happens only for a specific type of R-U extension $E|F$; see page 348. We refer to Remark 4.8, Remark 4.11 and Section 4D for a further discussion of this special case.

On the other hand, each projective line in $\mathcal{N}_E^{\text{naive}}(\bar{k})$ contains $q + 1$ τ -invariant points. Such a τ -invariant point M is an intersection point of two or more projective lines if and only if $|t| = |\pi_0|$ or $\Lambda_1 = M^\tau \subseteq C$ has a norm ideal satisfying $\text{Nm}(\Lambda_1) \subseteq \pi_0^2 O_F$.

Let $\Lambda \subseteq C$ as in Lemma 4.5. We denote by X_Λ^+ the formal O_F -module corresponding to the Dieudonné module $M = \Lambda \otimes \check{O}_E$. There is a canonical quasi-isogeny

$$\varrho_\Lambda^+ : \mathbb{X} \rightarrow X_\Lambda^+$$

of F -height 1. For $S \in \text{Nilp}_{\check{O}_F}$, we define

$$\mathcal{N}_{E,\Lambda}(S) = \{(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\text{naive}}(S) \mid (\varrho_\Lambda^+ \times S) \circ \varrho \text{ is an isogeny}\}.$$

By [Rapoport and Zink 1996, Proposition 2.9], the functor $\mathcal{N}_{E,\Lambda}$ is representable by a closed formal subscheme of $\mathcal{N}_E^{\mathrm{naive}}$. On geometric points, we have

$$(4-5) \quad \mathcal{N}_{E,\Lambda}(\bar{k}) \xrightarrow{\sim} \mathbb{P}(\Lambda/\Pi\Lambda)(\bar{k}),$$

as follows from Lemma 4.5 (1).

Proposition 4.7. *The reduced locus of $\mathcal{N}_E^{\mathrm{naive}}$ is a union*

$$(\mathcal{N}_E^{\mathrm{naive}})_{\mathrm{red}} = \bigcup_{\Lambda \subseteq C} \mathcal{N}_{E,\Lambda},$$

where Λ runs over all unimodular lattices in C with $\mathrm{Nm}(\Lambda) \subseteq \pi_0 O_F$. For each Λ , there exists an isomorphism

$$\mathcal{N}_{E,\Lambda} \xrightarrow{\sim} \mathbb{P}(\Lambda/\Pi\Lambda),$$

inducing the bijection (4-5) on \bar{k} -valued points.

The proof is analogous to that of Proposition 3.9.

Remark 4.8. Similar to Remark 3.8 (3), we let $(\mathcal{N}_E)_{\mathrm{red}} \subseteq (\mathcal{N}_E^{\mathrm{naive}})_{\mathrm{red}}$ be the union of all projective lines $\mathcal{N}_{E,\Lambda}$ corresponding to *hyperbolic* unimodular lattices $\Lambda \subseteq C$. Later, we will define \mathcal{N}_E as a functor on $\mathrm{Nilp}_{\check{O}_F}$ and show that $\mathcal{N}_E \simeq \mathcal{M}_{Dr}$, where \mathcal{M}_{Dr} is the Drinfeld moduli problem (see Theorem 4.14, a description of the formal scheme \mathcal{M}_{Dr} can be found in [Boutot and Carayol 1991, I.3]). In particular, $(\mathcal{N}_E)_{\mathrm{red}}$ is connected and each projective line in $(\mathcal{N}_E)_{\mathrm{red}}$ has $q + 1$ intersection points and there are two lines intersecting in each such point.

It might happen that $(\mathcal{N}_E)_{\mathrm{red}} = (\mathcal{N}_E^{\mathrm{naive}})_{\mathrm{red}}$ (see, for example, Figure 2(b)) if there are no nonhyperbolic unimodular lattices $\Lambda \subseteq C$ with $\mathrm{Nm}(\Lambda) \subseteq \pi_0 O_F$. In fact, this is the case if and only if $|t| = |\pi_0|$; see Proposition 2.4 and Lemma 2.5. (Note however that we still have $\mathcal{N}_E \neq \mathcal{N}_E^{\mathrm{naive}}$; see Remark 4.11 and Section 4D.)

Assume $|t| \neq |\pi_0|$ and let $P \in \mathcal{N}_E(\bar{k})$ be an intersection point. Then, as in the case where $E|F$ is of type R-P (compare Remark 3.8 (3)), the connected component of P in $((\mathcal{N}_E^{\mathrm{naive}})_{\mathrm{red}} \setminus (\mathcal{N}_E)_{\mathrm{red}}) \cup \{P\}$ consists of a finite union of projective lines (corresponding to nonhyperbolic lattices, by definition of $(\mathcal{N}_E)_{\mathrm{red}}$). In Figure 2(a), these components are indicated by dashed lines (they consist of just one projective line in that case).

4B. The straightening condition. As in the case R-P (see Section 3B) we use the results of Section 5 to define the straightening condition on $\mathcal{N}_E^{\mathrm{naive}}$. By Theorem 5.2 and Remark 5.1 (2), there exists a principal polarization $\tilde{\lambda}_{\mathbb{X}}^0$ on the framing object $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ such that the Rosati involution is the identity on O_E . We set $\tilde{\lambda}_{\mathbb{X}} = \tilde{\lambda}_{\mathbb{X}}^0 \circ \iota_{\mathbb{X}}(\Pi)$, which is again a polarization on \mathbb{X} with the Rosati involution inducing

the identity on O_E , but with kernel $\ker \tilde{\lambda}_{\mathbb{X}} = \mathbb{X}[\Pi]$. This polarization is unique up to a scalar in O_E^\times , i.e., any two polarizations $\tilde{\lambda}_{\mathbb{X}}$ and $\tilde{\lambda}'_{\mathbb{X}}$ with these properties satisfy

$$\tilde{\lambda}'_{\mathbb{X}} = \tilde{\lambda}_{\mathbb{X}} \circ \iota(\alpha),$$

for some $\alpha \in O_E^\times$. For any $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\text{naive}}(S)$,

$$\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}}) = \varrho^*(\tilde{\lambda}_{\mathbb{X}}^0) \circ \iota(\Pi)$$

is a polarization on X with kernel $\ker \tilde{\lambda} = X[\Pi]$; see [Theorem 5.2 \(2\)](#).

Recall that a unimodular or Π -modular lattice $\Lambda \subseteq C$ is called *hyperbolic* if there exists a basis (e_1, e_2) of Λ such that, with respect to this basis, h has the form

$$\begin{pmatrix} & \Pi^i \\ \bar{\Pi}^i & \end{pmatrix},$$

for $i = 0$ (resp. 1). By [Lemma 2.5](#), this is the case if and only if $\text{Nm}(\Lambda) = t O_F$.

Proposition 4.9. *For a suitable choice of $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ and $\tilde{\lambda}_{\mathbb{X}}$, the quasipolarization*

$$\lambda_{\mathbb{X},1} = \frac{1}{t}(\lambda_{\mathbb{X}} + \tilde{\lambda}_{\mathbb{X}})$$

is a polarization on \mathbb{X} . Let $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\text{naive}}(\bar{k})$ and $\tilde{\lambda} = \varrho^(\tilde{\lambda}_{\mathbb{X}})$. Then $\lambda_1 = \frac{1}{t}(\lambda + \tilde{\lambda})$ is a polarization if and only if $(X, \iota, \lambda, \varrho) \in \mathcal{N}_{E,\Lambda}(\bar{k})$ for a hyperbolic unimodular lattice $\Lambda \subseteq C$.*

Proof. On the rational Dieudonné module $N = M_{\mathbb{X}} \otimes_{\check{O}_F} \check{F}$, denote by $\langle \cdot, \cdot \rangle$, (\cdot, \cdot) and $\langle \cdot, \cdot \rangle_1$ the alternating forms induced by $\lambda_{\mathbb{X}}$, $\tilde{\lambda}_{\mathbb{X}}$ and $\lambda_{\mathbb{X},1}$, respectively. The form $\langle \cdot, \cdot \rangle_1$ is integral on $M_{\mathbb{X}}$ if and only if $\lambda_{\mathbb{X},1}$ is a polarization on \mathbb{X} . We have

$$(\mathbf{F}x, y) = (x, \mathbf{V}y)^\sigma,$$

$$(\Pi x, y) = (x, \bar{\Pi}y),$$

$$\langle x, y \rangle_1 = \frac{1}{t}(\langle x, y \rangle + (x, y))$$

for all $x, y \in N$. The form (\cdot, \cdot) induces an \check{E} -bilinear alternating form b on N by the formula

$$(4-6) \quad b(x, y) = c((\Pi x, y) - \bar{\Pi}(x, y)).$$

Here, c is a unit in \check{O}_E such that $c \cdot \sigma(c)^{-1} = \bar{\Pi}\Pi^{-1}$. Since

$$\frac{\bar{\Pi}}{\Pi} = \frac{t - \Pi}{\Pi} \in 1 + \frac{t}{\Pi} \check{O}_E,$$

we can even choose $c \in 1 + t\Pi^{-1}\check{O}_E$. The dual of M with respect to this form is again $M^\sharp = \Pi^{-1}M$, since

$$(x, y) = \text{Tr}_{\check{E}|\check{F}}\left(\frac{1}{t\vartheta c} \cdot b(x, y)\right),$$

and the inverse different of $E|F$ is given by $\mathfrak{D}_{E|F}^{-1} = t^{-1} O_E$; see [Lemma 2.2](#). Now b is invariant under the σ -linear operator $\tau = \Pi V^{-1} = F \bar{\Pi}^{-1}$, because

$$b(\tau x, \tau y) = b(F \bar{\Pi}^{-1} x, \Pi V^{-1} y) = \frac{c}{\sigma(c)} \cdot b(\bar{\Pi}^{-1} x, \Pi y)^\sigma = b(x, y)^\sigma.$$

Hence b defines an E -linear alternating form on C .

We choose the framing object $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ such that $M_{\mathbb{X}}$ is τ -invariant (see [Lemma 4.4](#)) and such that $\Lambda_1 = M_{\mathbb{X}}^\tau$ is hyperbolic. We can find a basis (e_1, e_2) of Λ_1 such that

$$h \cong \begin{pmatrix} & \Pi \\ \bar{\Pi} & \end{pmatrix}, \quad b \cong \begin{pmatrix} & u \\ -u & \end{pmatrix}$$

for some $u \in E^\times$. Since $\tilde{\lambda}_{\mathbb{X}}$ has the same kernel as $\lambda_{\mathbb{X}}$, we have $u = \bar{\Pi} u'$ for some unit $u' \in O_E^\times$. We can choose $\tilde{\lambda}_{\mathbb{X}}$ such that $u' = 1$ and $u = \bar{\Pi}$. Now $\frac{1}{t}(h(x, y) + b(x, y))$ is integral for all $x, y \in \Lambda_1$. Hence $\frac{1}{t}(h(x, y) + b(x, y))$ is also integral for all $x, y \in M_{\mathbb{X}}$. For all $x, y \in M_{\mathbb{X}}$, we have

$$\begin{aligned} \langle x, y \rangle_1 &= \frac{1}{t}(\langle x, y \rangle + (x, y)) = \frac{1}{t} \mathrm{Tr}_{\check{E}|\check{F}} \left(\frac{1}{t\vartheta} \cdot h(x, y) + \frac{1}{t\vartheta c} \cdot b(x, y) \right) \\ &= \mathrm{Tr}_{\check{E}|\check{F}} \left(\frac{1}{t^2\vartheta} \cdot (h(x, y) + b(x, y)) \right) + \mathrm{Tr}_{\check{E}|\check{F}} \left(\frac{1-c}{t^2\vartheta c} \cdot b(x, y) \right). \end{aligned}$$

The first summand is integral since $\frac{1}{t}(h(x, y) + b(x, y))$ is integral. The second summand is integral since $1 - c$ is divisible by $t\Pi^{-1}$ and $b(x, y)$ lies in $\Pi \check{O}_E$. It follows that the second summand above is integral as well. Hence \langle, \rangle_1 is integral on $M_{\mathbb{X}}$ and this implies that $\lambda_{\mathbb{X},1}$ is a polarization on \mathbb{X} .

Now let $(X, \iota, \lambda, \varrho) \in \mathcal{N}_{\check{E}}^{\mathrm{naive}}(\bar{k})$ and denote by $M \subseteq N$ its Dieudonné module. Assume that $\lambda_1 = t^{-1}(\lambda + \bar{\lambda})$ is a polarization on X . Then \langle, \rangle_1 is integral on M . But this is equivalent to $t^{-1}(h(x, y) + b(x, y))$ being integral for all $x, y \in M$. For $x = y$, we have

$$h(x, x) = h(x, x) + b(x, x) \in t\check{O}_F.$$

Let $\Lambda \subseteq C$ be the unimodular or Π -modular lattice given by $\Lambda = M^\tau$, resp. $\Lambda = (M + \tau(M))^\tau$; see [Lemma 4.4](#). Then $h(x, x) \in tO_F$ for all $x \in \Lambda$. Thus $\mathrm{Nm}(\Lambda) \subseteq tO_F$ and, by minimality, this implies that $\mathrm{Nm}(\Lambda) = tO_F$ and Λ is hyperbolic (see [Lemma 2.5](#)). Hence, in either case, the point corresponding to $(X, \iota, \lambda, \varrho)$ lies in $\mathcal{N}_{E,\Lambda'}$ for a hyperbolic lattice Λ' .

Conversely, assume that $(X, \iota, \lambda, \varrho) \in \mathcal{N}_{E,\Lambda}(\bar{k})$ for some hyperbolic lattice $\Lambda \subseteq C$. We want to show that λ_1 is a polarization on X . This follows if \langle, \rangle_1 is integral on M , or equivalently, if $t^{-1}(h(x, y) + b(x, y))$ is integral on M . For this, it is enough to show that $t^{-1}(h(x, y) + b(x, y))$ is integral on Λ . Let $\Lambda' \subseteq C$ be the unimodular lattice generated by $\bar{\Pi}^{-1}e_1$ and e_2 , where (e_1, e_2) is the basis of the Π -modular

lattice $\Lambda_1 = M_{\mathbb{X}}$. With respect to the basis $(\bar{\Pi}^{-1}e_1, e_2)$, we have

$$h \hat{=} \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, \quad b \hat{=} \begin{pmatrix} & 1 \\ -1 & \end{pmatrix}.$$

In particular, Λ' is a hyperbolic lattice and $t^{-1}(h + b)$ is integral on Λ' . By [Proposition 2.4](#), there exists an element $g \in \mathrm{SU}(C, h)$ with $g\Lambda = \Lambda'$. Since $\det g = 1$, the alternating form b is invariant under g . Thus $t^{-1}(h + b)$ is also integral on Λ . \square

From now on, we assume that $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ and $\tilde{\lambda}_{\mathbb{X}}$ are chosen in a way such that

$$\lambda_{\mathbb{X},1} = \frac{1}{t}(\lambda_{\mathbb{X}} + \tilde{\lambda}_{\mathbb{X}}) \in \mathrm{Hom}(\mathbb{X}, \mathbb{X}^{\vee}).$$

Definition 4.10. A tuple $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\mathrm{naive}}(S)$ satisfies the *straightening condition* if

$$(4-7) \quad \lambda_1 = \frac{1}{t}(\lambda + \tilde{\lambda}) \in \mathrm{Hom}(X, X^{\vee}).$$

This condition is independent of the choice of $\tilde{\lambda}_{\mathbb{X}}$. In fact, we can only change $\tilde{\lambda}_{\mathbb{X}}$ by a scalar of the form $1 + t\Pi^{-1}u$, $u \in O_E$. But if $\tilde{\lambda}'_{\mathbb{X}} = \tilde{\lambda}_{\mathbb{X}} \circ \iota(1 + t\Pi^{-1}u)$, then $\lambda'_{\mathbb{X},1} = \lambda_{\mathbb{X},1} + \tilde{\lambda}_{\mathbb{X}} \circ \iota(\Pi^{-1}u) = \lambda_{\mathbb{X},1} + \tilde{\lambda}_{\mathbb{X}}^0 \circ \iota(u)$ and $\lambda'_1 = \lambda_1 + \varrho^*(\tilde{\lambda}_{\mathbb{X}}^0) \circ \iota(u)$. Clearly, λ'_1 is a polarization if and only if λ_1 is one.

For $S \in \mathrm{Nilp}_{\check{O}_F}$, let $\mathcal{N}_E(S)$ be the set of all tuples $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\mathrm{naive}}(S)$ that satisfy the straightening condition. By [\[Rapoport and Zink 1996, Proposition 2.9\]](#), the functor \mathcal{N}_E is representable by a closed formal subscheme of $\mathcal{N}_E^{\mathrm{naive}}$.

Remark 4.11. The reduced locus of \mathcal{N}_E is given by

$$(\mathcal{N}_E)_{\mathrm{red}} = \bigcup_{\Lambda \subseteq C} \mathcal{N}_{E,\Lambda} \simeq \bigcup_{\Lambda \subseteq C} \mathbb{P}(\Lambda / \Pi\Lambda),$$

where the union goes over all *hyperbolic* unimodular lattices $\Lambda \subseteq C$. Note that, depending on the form of the R-U extension $E|F$, it may happen that all unimodular lattices are hyperbolic (when $|t| = |\pi_0|$) and in that case, we have $(\mathcal{N}_E)_{\mathrm{red}} = (\mathcal{N}_E^{\mathrm{naive}})_{\mathrm{red}}$. However, the equality does not extend to an isomorphism between \mathcal{N}_E and $\mathcal{N}_E^{\mathrm{naive}}$. This will be discussed in [Section 4D](#).

4C. The main theorem for the case R-U. As in the case R-P, we want to establish a connection to the Drinfeld moduli problem. Therefore, fix an embedding of E into the quaternion division algebra B . Let $(\mathbb{X}, \iota_{\mathbb{X}})$ be the framing object of the Drinfeld problem. We want to construct a polarization $\lambda_{\mathbb{X}}$ on \mathbb{X} with $\ker \lambda_{\mathbb{X}} = \mathbb{X}[\Pi]$ and Rosati involution given by $b \mapsto \vartheta b' \vartheta^{-1}$ on B . Here $b \mapsto b'$ denotes the standard involution on B .

By [Lemma 2.3 \(2\)](#), there exists an embedding $E_1 \hookrightarrow B$ of a ramified quadratic extension $E_1|F$ of type R-P, such that $\Pi_1\vartheta = -\vartheta\Pi_1$ for a prime element $\Pi_1 \in E_1$. From [Proposition 3.14 \(1\)](#) we get a principal polarization $\lambda_{\mathbb{X}}^0$ on \mathbb{X} with associated Rosati involution $b \mapsto \Pi_1 b' \Pi_1^{-1}$. If we assume fixed choices of E_1 and Π_1 , this is unique up to a scalar in O_F^\times . We define

$$\lambda_{\mathbb{X}} = \lambda_{\mathbb{X}}^0 \circ \iota_{\mathbb{X}}(\Pi_1 \vartheta).$$

Since $\lambda_{\mathbb{X}}^0$ is a principal polarization and $\Pi_1 \vartheta$ and Π have the same valuation in O_B , we have $\ker \lambda_{\mathbb{X}} = \mathbb{X}[\Pi]$. The Rosati involution of $\lambda_{\mathbb{X}}$ is $b \mapsto \vartheta b' \vartheta^{-1}$. On the other hand, any polarization on \mathbb{X} satisfying these two conditions can be constructed in this way (using the same choices for E_1 and Π_1). Hence:

- Lemma 4.12.** (1) *There exists a polarization $\lambda_{\mathbb{X}} : \mathbb{X} \rightarrow \mathbb{X}^\vee$, unique up to a scalar in O_F^\times , with $\ker \lambda_{\mathbb{X}} = \mathbb{X}[\Pi]$ and associated Rosati involution $b \mapsto \vartheta b' \vartheta^{-1}$.*
- (2) *Fix $\lambda_{\mathbb{X}}$ as in (1) and let $(X, \iota_B, \varrho) \in \mathcal{M}_{Dr}(S)$. There exists a unique polarization λ on X with $\ker \lambda = X[\Pi]$ and Rosati involution $b \mapsto \vartheta b' \vartheta^{-1}$ such that $\varrho^*(\lambda_{\mathbb{X}}) = \lambda$ on $\bar{S} = S \times_{\mathrm{spf} \check{O}_F} \bar{k}$.*

Note also that the involution $b \mapsto \vartheta b' \vartheta^{-1}$ does not depend on the choice of $\vartheta \in E$. We write $\iota_{\mathbb{X}, E}$ for the restriction of $\iota_{\mathbb{X}}$ to $E \subseteq B$ and, in the same manner, we write ι_E for the restriction of ι_B to E for any $(X, \iota_B, \varrho) \in \mathcal{M}_{Dr}(S)$. Fix a polarization $\lambda_{\mathbb{X}}$ of \mathbb{X} as in [Lemma 4.12 \(1\)](#). Accordingly for a tuple $(X, \iota_B, \varrho) \in \mathcal{M}_{Dr}(S)$, let λ be the polarization given by [Lemma 4.12 \(2\)](#).

Lemma 4.13. *The tuple $(\mathbb{X}, \iota_{\mathbb{X}, E}, \lambda_{\mathbb{X}})$ is a framing object of $\mathcal{N}_E^{\mathrm{naive}}$. Moreover, the map*

$$(X, \iota_B, \varrho) \mapsto (X, \iota_E, \lambda, \varrho)$$

induces a closed embedding of formal schemes

$$\eta : \mathcal{M}_{Dr} \hookrightarrow \mathcal{N}_E^{\mathrm{naive}}.$$

Proof. We follow the same argument as in the proof of [Lemma 3.15](#). Again it is enough to check that $\mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ contains $\mathrm{SU}(C, h)$ as a closed subgroup and that ι_E satisfies the Kottwitz condition.

By [[Rapoport and Zink 2017](#), Proposition 5.8], the special condition on ι_B implies the Kottwitz condition for ι_E . It remains to show that $\mathrm{SU}(C, h) \subseteq \mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$. But the group $G_{(\mathbb{X}, \iota_{\mathbb{X}})}$ of automorphisms of determinant 1 of $(\mathbb{X}, \iota_{\mathbb{X}})$ is isomorphic to $\mathrm{SL}_{2, F}$ and $G_{(\mathbb{X}, \iota_{\mathbb{X}})} \subseteq \mathrm{QIsog}(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ is a Zariski-closed subgroup by the same argument as in [Lemma 3.15](#). Hence the statement follows from the exceptional isomorphism $\mathrm{SL}_{2, F} \simeq \mathrm{SU}(C, h)$. \square

As a next step, we want to show that this already induces a closed embedding

$$(4-8) \quad \eta : \mathcal{M}_{Dr} \hookrightarrow \mathcal{N}_E.$$

Let $\tilde{E} \hookrightarrow B$ an embedding of a ramified quadratic extension $\tilde{E}|F$ of type R-U as in [Lemma 2.3 \(2\)](#). On the framing object $(\mathbb{X}, \iota_{\mathbb{X}})$ of \mathcal{M}_{Dr} , we define a polarization $\tilde{\lambda}_{\mathbb{X}}$ via

$$\tilde{\lambda}_{\mathbb{X}} = \lambda_{\mathbb{X}} \circ \iota_{\mathbb{X}}(\tilde{\vartheta}),$$

where $\tilde{\vartheta}$ is a unit in \tilde{E} of the form $\tilde{\vartheta}^2 = 1 + (t^2/\pi_0) \cdot u$; see [Lemma 2.3 \(2\)](#). The Rosati involution of $\tilde{\lambda}_{\mathbb{X}}$ induces the identity on O_E and we have

$$\begin{aligned} \lambda_{\mathbb{X},1} &= \frac{1}{t}(\lambda_{\mathbb{X}} + \tilde{\lambda}_{\mathbb{X}}) = \frac{1}{t} \cdot \lambda_{\mathbb{X}} \circ \iota_B(1 + \tilde{\vartheta}) = \lambda_{\mathbb{X}} \circ \iota_B(\tilde{\Pi}/\pi_0) \\ &= \lambda_{\mathbb{X}} \circ \iota_B(\Pi^{-1}\gamma) \in \text{Hom}(\mathbb{X}, \mathbb{X}^{\vee}), \end{aligned}$$

using the notation of [Lemma 2.3 \(2\)](#). For $(X, \iota_B, \varrho) \in \mathcal{M}_{Dr}(S)$, we set $\tilde{\lambda} = \lambda \circ \iota_B(\tilde{\vartheta})$. By the same calculation, we have $\lambda_1 = \frac{1}{t}(\lambda + \tilde{\lambda}) \in \text{Hom}(X, X^{\vee})$. Thus the tuple $(X, \iota_E, \lambda, \varrho) = \eta(X, \iota_B, \varrho)$ satisfies the straightening condition. Hence we get a closed embedding of formal schemes $\eta : \mathcal{M}_{Dr} \rightarrow \mathcal{N}_E$ which is independent of the choice of \tilde{E} .

Theorem 4.14. *$\eta : \mathcal{M}_{Dr} \rightarrow \mathcal{N}_E$ is an isomorphism of formal schemes.*

We first check this for \bar{k} -valued points:

Lemma 4.15. *η induces a bijection $\eta(\bar{k}) : \mathcal{M}_{Dr}(\bar{k}) \rightarrow \mathcal{N}_E(\bar{k})$.*

Proof. We only have to show surjectivity and we will use for this the Dieudonné theory description of $\mathcal{N}_E^{\text{naive}}(\bar{k})$; see [\(4-4\)](#). The rational Dieudonné module $N = N_{\mathbb{X}}$ of \mathbb{X} now carries additionally an action of B . The embedding $F^{(2)} \hookrightarrow B$ given by

$$(4-9) \quad \gamma \mapsto \frac{\Pi \cdot \tilde{\Pi}}{\pi_0},$$

(see [Lemma 2.3 \(2\)](#)) induces a $\mathbb{Z}/2$ -grading $N = N_0 \oplus N_1$. Here,

$$\begin{aligned} N_0 &= \{x \in N \mid \iota(a)x = ax \text{ for all } a \in F^{(2)}\}, \\ N_1 &= \{x \in N \mid \iota(a)x = \sigma(a)x \text{ for all } a \in F^{(2)}\} \end{aligned}$$

for a fixed embedding $F^{(2)} \hookrightarrow \check{F}$. The operators F and V have degree 1 with respect to this grading. The principal polarization

$$\lambda_{\mathbb{X},1} = \frac{1}{t}(\lambda_{\mathbb{X}} + \tilde{\lambda}_{\mathbb{X}}) = \lambda_{\mathbb{X}} \circ \iota_{\mathbb{X}}(\Pi^{-1}\gamma)$$

induces an alternating form $\langle \cdot, \cdot \rangle_1$ on N that satisfies

$$\langle x, y \rangle_1 = \langle x, \iota(\Pi^{-1}\gamma) \cdot y \rangle,$$

for all $x, y \in N$. Let $M \in \mathcal{N}_E(\bar{k}) \subseteq \mathcal{N}_E^{\mathrm{naive}}(\bar{k})$ be an \check{O}_F -lattice in N . We claim that $M \in \mathcal{M}_{Dr}(\bar{k})$. For this, it is necessary that M is stable under the action of $O_F^{(2)}$ (since $O_B = O_F[\Pi, \gamma] = O_F^{(2)}[\Pi]$; see [Lemma 2.3 \(2\)](#)) or equivalently, that M respects the grading of N , i.e., $M = M_0 \oplus M_1$ for $M_i = M \cap N_i$. Furthermore M has to satisfy the *special* condition:

$$\dim M_0 / VM_1 = \dim M_1 / VM_0 = 1.$$

We first show that $M = M_0 \oplus M_1$. Let $y = y_0 + y_1 \in M$ with $y_i \in N_i$. Since $M = \Pi M^\vee$, we have

$$\langle x, \iota(\Pi)^{-1}y \rangle = \langle x, \iota(\Pi)^{-1}y_0 \rangle + \langle x, \iota(\Pi)^{-1}y_1 \rangle \in \check{O}_F,$$

for all $x \in M$. Together with

$$\begin{aligned} \langle x, y \rangle_1 &= \langle x, y_0 \rangle_1 + \langle x, y_1 \rangle_1 = \langle x, \iota(\tilde{\Pi}/\pi_0)y_0 \rangle + \langle x, \iota(\tilde{\Pi}/\pi_0)y_1 \rangle \\ &= \gamma \cdot \langle x, \iota(\Pi^{-1})y_0 \rangle + (1 - \gamma) \cdot \langle x, \iota(\Pi^{-1})y_1 \rangle \in \check{O}_F, \end{aligned}$$

this implies that $\langle x, \iota(\Pi^{-1})y_0 \rangle$ and $\langle x, \iota(\Pi^{-1})y_1 \rangle$ lie in \check{O}_F for all $x \in M$. Hence, $y_0, y_1 \in M$ and this means that M respects the grading. It follows that M is stable under the action of O_B .

In order to show that M is special, note that

$$\langle Vx, Vy \rangle_1^\sigma = \langle FVx, y \rangle_1 = \pi_0 \cdot \langle x, y \rangle_1 \in \pi_0 \check{O}_F,$$

for all $x, y \in M$. The form $\langle \cdot, \cdot \rangle_1$ comes from a principal polarization, so it induces a perfect form on M . Now it is enough to show that also the restrictions of $\langle \cdot, \cdot \rangle_1$ to M_0 and M_1 are perfect. Indeed, if M was not special, we would have $M_i = VM_{i+1}$ for some i and this would contradict $\langle \cdot, \cdot \rangle_1$ being perfect on M_i . We prove that $\langle \cdot, \cdot \rangle_1$ is perfect on M_i by showing $\langle M_0, M_1 \rangle_1 \subseteq \pi_0 \check{O}_F$.

Let $x \in M_0$ and $y \in M_1$. Then,

$$\begin{aligned} \langle x, y \rangle_1 &= (1 - \gamma) \cdot \langle x, \iota(\Pi)^{-1}y \rangle, \\ \langle x, y \rangle_1 &= -\langle y, x \rangle_1 = -\gamma \cdot \langle y, \iota(\Pi)^{-1}x \rangle = \gamma \cdot \langle x, \iota(\bar{\Pi})^{-1}y \rangle. \end{aligned}$$

We take the difference of these two equations. From $\Pi \equiv \bar{\Pi} \pmod{\pi_0}$, it follows that $\langle x, \iota(\Pi)^{-1}y \rangle \equiv 0 \pmod{\pi_0}$ and thus also $\langle x, y \rangle_1 \equiv 0 \pmod{\pi_0}$. The form $\langle \cdot, \cdot \rangle_1$ is hence perfect on M_0 and M_1 and the special condition follows. This finishes the proof of [Lemma 4.15](#). \square

Proof of Theorem 4.14. Let $(\mathbb{X}, \iota_{\mathbb{X}})$ be a framing object for \mathcal{M}_{Dr} and let further

$$\eta(\mathbb{X}, \iota_{\mathbb{X}}) = (\mathbb{X}, \iota_{\mathbb{X}, E}, \lambda_{\mathbb{X}})$$

be the corresponding framing object for \mathcal{N}_E . We fix an embedding $F^{(2)} \hookrightarrow B$ as in [Lemma 2.3 \(2\)](#). For $S \in \text{Nilp}_{\check{O}_F}$, let $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E(S)$ and $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$. We have

$$\begin{aligned} \varrho^{-1} \circ \iota_{\mathbb{X}}(\gamma) \circ \varrho &= \varrho^{-1} \circ \iota_{\mathbb{X}}(\Pi) \circ \lambda_{\mathbb{X}}^{-1} \circ \lambda_{\mathbb{X},1} \circ \varrho \\ &= \iota(\Pi) \circ \lambda^{-1} \circ \lambda_1 \in \text{End}(X) \end{aligned}$$

for $\lambda_1 = t^{-1}(\lambda + \tilde{\lambda})$, since $\ker \lambda = X[\Pi]$. But $O_B = O_F[\Pi, \gamma]$ (see [Lemma 2.3 \(2\)](#)), so this already induces an O_B -action ι_B on X . It remains to show that (X, ι_B, ϱ) satisfies the *special* condition (see the discussion before [Proposition 3.14](#) for a definition).

The special condition is open and closed (see [\[Rapoport and Zink 2017, p. 7\]](#)) and η is bijective on \bar{k} -points. Hence η induces an isomorphism on reduced subschemes

$$(\eta)_{\text{red}} : (\mathcal{M}_{Dr})_{\text{red}} \xrightarrow{\sim} (\mathcal{N}_E)_{\text{red}},$$

because $(\mathcal{M}_{Dr})_{\text{red}}$ and $(\mathcal{N}_E)_{\text{red}}$ are locally of finite type over $\text{Spec } \bar{k}$. It follows that $\eta : \mathcal{M}_{Dr} \rightarrow \mathcal{N}_E$ is an isomorphism. \square

4D. Deformation theory of intersection points. In this section, we will study the deformation rings of certain geometric points in $\mathcal{N}_E^{\text{naive}}$ with the goal of proving that $\mathcal{N}_E \subseteq \mathcal{N}_E^{\text{naive}}$ is a strict inclusion even in the case $|t| = |\pi_0|$. In contrast to the non-2-adic case, we are not able to use the theory of local models (see [\[Pappas et al. 2013\]](#) for a survey) since there is in general no normal form for the lattices $\Lambda \subseteq C$; see [Proposition 2.4](#) and [\[Rapoport and Zink 1996, Theorem 3.16\]](#).³ Thus we will take the more direct approach of studying the deformations of a fixed point $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\text{naive}}(\bar{k})$ and using the theory of Grothendieck and Messing [\[Messing 1972\]](#).

Let $\Lambda \subseteq C$ be a Π -modular hyperbolic lattice. By [Lemma 4.5](#), there is a unique point $x = (X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\text{naive}}(\bar{k})$ with a τ -stable Dieudonné module $M \subseteq C \otimes_E \check{E}$ and $M^\tau = \Lambda$. Since Λ is hyperbolic, x satisfies the straightening condition, i.e., $x \in \mathcal{N}_E(\bar{k})$. (In [Figure 2](#), x would lie on the intersection of two solid lines.)

Let $\widehat{\mathcal{O}_{\mathcal{N}_E^{\text{naive}}, x}}$ be the formal completion of the local ring at x . It represents the following deformation functor Def_x . For an artinian \check{O}_F -algebra R with residue field \bar{k} , we have

$$\text{Def}_x(R) = \{(Y, \iota_Y, \lambda_Y)/R \mid Y_{\bar{k}} \cong X\},$$

where (Y, ι_Y, λ_Y) satisfies the usual conditions (see [Section 4A](#)) and the isomorphism $Y_{\bar{k}} \cong X$ is actually an isomorphism of tuples $(Y_{\bar{k}}, \iota_Y, \lambda_Y) \cong (X, \iota, \lambda)$ as in [Definition 3.1](#).

³It is possible to define a local model for the nonnaive spaces \mathcal{N}_E (also in the case R-P) and establish a local model diagram as in [\[Rapoport and Zink 1996, Definition 3.27\]](#). The local model is then isomorphic to the local model of the Drinfeld moduli problem. This will be part of a future paper of the author.

Now assume the quotient map $R \rightarrow \bar{k}$ is an O_F -pd-thickening (see [Ahsendorf 2011]). For example, this is the case when $\mathfrak{m}^2 = 0$ for the maximal ideal \mathfrak{m} of R . Then, by Grothendieck–Messing theory (see [Messing 1972] and [Ahsendorf 2011]), we get an explicit description of $\mathrm{Def}_X(R)$ in terms of liftings of the Hodge filtration:

The (relative) Dieudonné crystal $\mathbb{D}_X(R)$ of X evaluated at R is naturally isomorphic to the free R -module $\Lambda \otimes_{O_F} R$ and this isomorphism is equivariant under the action of O_E induced by ι and respects the perfect form $\Phi = \langle \cdot, \cdot \rangle \circ (1, \Pi^{-1})$ induced by $\lambda \circ \iota(\Pi^{-1})$. The Hodge filtration of X is given by $\mathcal{F}_X = V \cdot \mathbb{D}_X(\bar{k}) \cong \Pi \cdot (\Lambda \otimes_{O_F} \bar{k}) \subseteq \Lambda \otimes_{O_F} \bar{k}$.

A point $Y \in \mathrm{Def}_X(R)$ now corresponds, via Grothendieck–Messing, to a direct summand $\mathcal{F}_Y \subseteq \Lambda \otimes_{O_F} R$ of rank 2 lifting \mathcal{F}_X , stable under the O_E -action and totally isotropic with respect to Φ . Furthermore, it has to satisfy the Kottwitz condition (see Section 4A): For the action of $\alpha \in O_E$ on $\mathrm{Lie} Y = (\Lambda \otimes_{O_F} R)/\mathcal{F}_Y$, we have

$$\mathrm{char}(\mathrm{Lie} Y, T \mid \iota(\alpha)) = (T - \alpha)(T - \bar{\alpha}).$$

Let us now fix an O_E -basis (e_1, e_2) of Λ and let us write everything in terms of the O_F -basis $(e_1, e_2, \Pi e_1, \Pi e_2)$. Since Λ is hyperbolic, we can fix (e_1, e_2) such that h is represented by the matrix

$$h \hat{=} \begin{pmatrix} & \Pi \\ \bar{\Pi} & \end{pmatrix},$$

and then

$$\Phi = \mathrm{Tr}_{E|F} \frac{1}{t\vartheta} h(\cdot, \Pi^{-1}\cdot) \hat{=} \left(\begin{array}{c|c} t/\pi_0 & 1 \\ \hline -1 + t^2/\pi_0 & t \end{array} \right).$$

An R -basis (v_1, v_2) of \mathcal{F}_Y can now be chosen such that

$$(v_1 v_2) = \begin{pmatrix} y_{11} & y_{12} \\ y_{21} & y_{22} \\ 1 & \\ & 1 \end{pmatrix},$$

with $y_{ij} \in R$. As an easy calculation shows, the conditions on \mathcal{F}_Y above are now equivalent to the following conditions on the y_{ij} :

$$\begin{aligned} y_{11} + y_{22} &= t, \\ y_{11}y_{22} - y_{12}y_{21} &= \pi_0, \\ t \left(\frac{ty_{22}}{\pi_0} + 2 \right) &= y_{11} \left(\frac{ty_{22}}{\pi_0} + 2 \right) = y_{21} \left(\frac{ty_{22}}{\pi_0} + 2 \right) = y_{12} \left(\frac{ty_{22}}{\pi_0} + 2 \right) = 0. \end{aligned}$$

Let T be the closed subscheme of $\mathrm{Spec} O_F[y_{11}, y_{12}, y_{21}, y_{22}]$ given by these equations. Let T_y be the formal completion of the localization at the ideal generated by the y_{ij} and π_0 . Then we have $\mathrm{Def}_x(R) \cong T_y(R)$ for any O_F -pd-thickening $R \rightarrow \bar{k}$. In particular, the first infinitesimal neighborhoods of Def_x and T_y coincide. The first infinitesimal neighborhood of T_y is given by $\mathrm{Spec} O_F[y_{ij}]/((y_{ij})^2, y_{11} + y_{22} - t, \pi_0)$, hence T_y has Krull dimension 3 and so has Def_x . However, \mathcal{M}_{Dr} is regular of dimension 2; cf. [Boutot and Carayol 1991]. Thus:

Proposition 4.16. $\mathcal{N}_E^{\mathrm{naive}} \neq \mathcal{M}_{Dr}$, even when $|t| = |\pi_0|$.

Indeed, $\dim \widehat{\mathcal{O}}_{\mathcal{N}_E^{\mathrm{naive}}, x} = \dim \mathrm{Def}_x = 3 > 2 = \dim \widehat{\mathcal{O}}_{\mathcal{N}_E, x}$.

5. A theorem on the existence of polarizations

In this section, we will prove the existence of the polarization $\tilde{\lambda}$ for any $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E^{\mathrm{naive}}(S)$ as claimed in the Sections 3B and 4B in both the cases R-P and R-U. In fact, we will show more generally that $\tilde{\lambda}$ exists even for the points of a larger moduli space \mathcal{M}_E where we forget about the polarization λ .

We start with the definition of the moduli space \mathcal{M}_E . Let $F|\mathbb{Q}_p$ be a finite extension (not necessarily $p = 2$) and let $E|F$ be a quadratic extension (not necessarily ramified). We denote by O_F and O_E the rings of integers, by k the residue field of O_F and by \bar{k} the algebraic closure of k . Furthermore, \check{F} is the completion of the maximal unramified extension of F and \check{O}_F its ring of integers. Let B be the quaternion division algebra over F and O_B the ring of integers.

If $E|F$ is unramified, we fix a common uniformizer $\pi_0 \in O_F \subseteq O_E$. If $E|F$ is ramified and $p > 2$, we choose a uniformizer $\Pi \in O_E$ such that $\pi_0 = \Pi^2 \in O_F$. If $E|F$ is ramified and $p = 2$, we use the notation of Section 2 for the cases R-P and R-U.

For $S \in \mathrm{Nilp}_{\check{O}_F}$, let $\mathcal{M}_E(S)$ be the set of isomorphism classes of tuples (X, ι_E, ϱ) over S . Here, X is a formal O_F -module of dimension 2 and height 4 and ι_E is an action of O_E on X satisfying the Kottwitz condition for the signature $(1, 1)$, i.e., the characteristic polynomial for the action of $\iota_E(\alpha)$ on $\mathrm{Lie}(X)$ is

$$(5-1) \quad \mathrm{char}(\mathrm{Lie} X, T \mid \iota(\alpha)) = (T - \alpha)(T - \bar{\alpha}),$$

for any $\alpha \in O_E$, compare the definition of $\mathcal{N}_E^{\mathrm{naive}}$ in Sections 3 and 4. The last entry ϱ is an O_E -linear quasi-isogeny

$$\varrho : X \times_S \bar{S} \rightarrow \mathbb{X} \times_{\mathrm{Spec} \bar{k}} \bar{S},$$

of height 0 to the framing object $(\mathbb{X}, \iota_{\mathbb{X}, E})$ defined over $\mathrm{Spec} \bar{k}$. The framing object for \mathcal{M}_E is the Drinfeld framing object $(\mathbb{X}, \iota_{\mathbb{X}, B})$ where we restrict the O_B -action to O_E for an arbitrary embedding $O_E \hookrightarrow O_B$. The special condition on $(\mathbb{X}, \iota_{\mathbb{X}, B})$ implies the Kottwitz condition for any $\alpha \in O_E$ by [Rapoport and Zink 2017, Proposition 5.8].

Remark 5.1. (1) Up to isogeny, there is more than one pair (X, ι_E) over $\mathrm{Spec} \bar{k}$ satisfying the conditions above. Indeed, let N_X be the rational Dieudonné module of (X, ι_E) . This is a 4-dimensional \check{F} -vector space with an action of O_E . The Frobenius F on N_X commutes with the action of O_E . For a suitable choice of a basis of N_X , it may be of either of the following two forms,

$$F = \begin{pmatrix} & & 1 \\ & & \\ \pi_0 & & \\ & \pi_0 & \end{pmatrix} \sigma \quad \text{or} \quad F = \begin{pmatrix} \pi_0 & & \\ & \pi_0 & \\ & & 1 \\ & & & 1 \end{pmatrix} \sigma.$$

This follows from the classification of isocrystals; see, for example, [Rapoport and Zink 1996, p. 3]. In the left case, F is isoclinic of slope $1/2$ (the supersingular case), and in the right case, the slopes are 0 and 1 . Our choice of the framing object above assures that we are in the supersingular case, since the framing object for the Drinfeld moduli problem can be written as a product of two formal O_F -modules of dimension 1 and height 2 (cf. [Boutot and Carayol 1991, p. 136–137]).

(2) Let $p = 2$ and $E|F$ ramified of type R-P or R-U. We can identify the framing objects $(\mathbb{X}, \iota_{\mathbb{X}, E})$ for $\mathcal{N}_E^{\mathrm{naive}}$, \mathcal{M}_{Dr} and \mathcal{M}_E by Proposition 3.14 and Lemma 4.13. In this way, we obtain a forgetful morphism $\mathcal{N}_E^{\mathrm{naive}} \rightarrow \mathcal{M}_E$. This is a closed embedding, since the existence of a polarization λ for $(X, \iota_E, \varrho) \in \mathcal{M}_E(S)$ is a closed condition by [Rapoport and Zink 1996, Proposition 2.9].

By [Rapoport and Zink 1996, Theorem 3.25], \mathcal{M}_E is pro-representable by a formal scheme over $\mathrm{Spf} \check{O}_F$. We will prove the following theorem in this section.

Theorem 5.2. (1) *There exists a principal polarization $\tilde{\lambda}_{\mathbb{X}}$ on $(\mathbb{X}, \iota_{\mathbb{X}, E})$ such that the Rosati involution induces the identity on O_E , i.e., $\iota(\alpha)^* = \iota(\alpha)$ for all $\alpha \in O_E$. This polarization is unique up to a scalar in O_E^\times , that is, for any two polarizations $\tilde{\lambda}_{\mathbb{X}}$ and $\tilde{\lambda}'_{\mathbb{X}}$ of this form, there exists an element $\alpha \in O_E^\times$ such that $\tilde{\lambda}'_{\mathbb{X}} = \tilde{\lambda}_{\mathbb{X}} \circ \iota_{\mathbb{X}, E}(\alpha)$.*

(2) *Fix $\tilde{\lambda}_{\mathbb{X}}$ as in part (1). For any $S \in \mathrm{Nilp}_{\check{O}_F}^\circ$ and $(X, \iota_E, \varrho) \in \mathcal{M}_E(S)$, there exists a unique principal polarization $\tilde{\lambda}$ on X such that the Rosati involution induces the identity on O_E and such that $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$.*

Remark 5.3. (1) We will see later that this theorem describes a natural isomorphism between \mathcal{M}_E and another space $\mathcal{M}_{E, \mathrm{pol}}$ which solves the moduli problem for tuples $(X, \iota_E, \tilde{\lambda}, \varrho)$ where $\tilde{\lambda}$ is a principal polarization with Rosati involution the identity on O_E . This is an RZ-space for the symplectic group $\mathrm{GSp}_2(E)$ and thus the theorem gives us another geometric realization of an exceptional isomorphism of reductive groups, in this case $\mathrm{GSp}_2(E) \cong \mathrm{GL}_2(E)$.

Since there is no such isomorphism in higher dimensions, the theorem does not generalize to these cases and a different approach is needed to formulate the straightening condition.

(2) With [Theorem 5.2](#) established, one can give an easier proof of the isomorphism $\mathcal{N}_E \xrightarrow{\sim} \mathcal{M}_{Dr}$ for the cases where $E|F$ is unramified or $E|F$ is ramified and $p > 2$, which is the main theorem of [\[Kudla and Rapoport 2014\]](#). Indeed, the main part of the proof in that paper consists of Propositions 2.1 and 3.1, which claim the existence of a certain principal polarization λ_X^0 for any point $(X, \iota, \lambda, \varrho) \in \mathcal{N}_E(S)$. But there is a canonical closed embedding $\mathcal{N}_E \hookrightarrow \mathcal{M}_E$ and under this embedding, λ_X^0 is just the polarization $\tilde{\lambda}$ of [Theorem 5.2](#), for a suitable choice of $\tilde{\lambda}_{\mathbb{X}}$ on the framing object. More explicitly, using the notation on page 2 of [\[loc. cit.\]](#), we take $\tilde{\lambda}_{\mathbb{X}} = \lambda_{\mathbb{X}} \circ \iota_{\mathbb{X}}^{-1}(\Pi) = \lambda_{\mathbb{X}}^0 \circ \iota_{\mathbb{X}}(-\delta)$ in the unramified case and $\tilde{\lambda}_{\mathbb{X}} = \lambda_{\mathbb{X}} \circ \iota_{\mathbb{X}}(\zeta^{-1})$ in the ramified case.

We will split the proof of this theorem into several lemmata. As a first step, we use Dieudonné theory to prove the statement for all geometric points.

Lemma 5.4. *Part (1) of [Theorem 5.2](#) holds. Furthermore, for a fixed polarization $\tilde{\lambda}_{\mathbb{X}}$ on $(\mathbb{X}, \iota_{\mathbb{X}, E})$ and for any $(X, \iota_E, \varrho) \in \mathcal{M}_E(\bar{k})$, the pullback $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$ is a polarization on X .*

Proof. This follows almost immediately from the theory of affine Deligne–Lusztig varieties (see, for example, [\[Chen and Viehmann 2015\]](#)) since we are comparing the geometric points of RZ-spaces for the isomorphic groups $\mathrm{GL}_2(E)$ and $\mathrm{GSp}_2(E)$.

It is also possible to check this via a more direct computation using Dieudonné theory, as we will indicate briefly. Proceeding very similarly to [Proposition 3.2](#) or [Proposition 4.1](#) (cf. [\[Kudla and Rapoport 2014\]](#) in the unramified case), we can associate to \mathbb{X} a lattice Λ in the 2-dimensional E -vector space C (the Frobenius invariant points of the (rational) Dieudonné module). The choice of a principal polarization on \mathbb{X} with trivial Rosati involution corresponds now exactly to a choice of perfect alternating form on Λ . It immediately follows that such a polarization exists and that it is unique up to a scalar in O_E^\times .

For the second part, let $X \in \mathcal{M}_E(\bar{k})$ and $M \subseteq C \otimes_E \check{E}$ be its Dieudonné module. Since ϱ has height 0, we have

$$[M : M \cap (\Lambda \otimes_E \check{E})] = [(\Lambda \otimes_E \check{E}) : M \cap (\Lambda \otimes_E \check{E})],$$

and one easily checks that a perfect alternating form b on Λ is also perfect on M . \square

In the following, we fix a polarization $\tilde{\lambda}_{\mathbb{X}}$ on $(\mathbb{X}, \iota_{\mathbb{X}, E})$ as in [Theorem 5.2 \(1\)](#). Let $(X, \iota_E, \varrho) \in \mathcal{M}_E(S)$ for $S \in \mathrm{Nilp}_{\check{O}_F}$ and consider the pullback $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$. In general, this is only a quasipolarization. It suffices to show that $\tilde{\lambda}$ is a polarization on X . Indeed, since ϱ is O_E -linear and of height 0, this is then automatically a principal polarization on X such that the Rosati involution is the identity on O_E .

Define a subfunctor $\mathcal{M}_{E,\mathrm{pol}} \subseteq \mathcal{M}_E$ by

$$\mathcal{M}_{E,\mathrm{pol}}(S) = \{(X, \iota_E, \varrho) \in \mathcal{M}_E(S) \mid \tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}}) \text{ is a polarization on } X\}.$$

This is a closed formal subscheme by [Rapoport and Zink 1996, Proposition 2.9]. Moreover, Lemma 5.4 shows that $\mathcal{M}_{E,\mathrm{pol}}(\bar{k}) = \mathcal{M}_E(\bar{k})$.

Remark 5.5. Equivalently, we can describe $\mathcal{M}_{E,\mathrm{pol}}$ as follows. For $S \in \mathrm{Nilp}_{\check{O}_F}^{\circ}$, we define $\mathcal{M}_{E,\mathrm{pol}}(S)$ to be the set of equivalence classes of tuples $(X, \iota_E, \tilde{\lambda}, \varrho)$, where

- X is a formal O_F -module over S of height 4 and dimension 2,
- ι_E is an action of O_E on X that satisfies the Kottwitz condition in (5-1) and
- $\tilde{\lambda}$ is a principal polarization on X such that the Rosati involution induces the identity on O_E .
- Furthermore, we fix a framing object $(\mathbb{X}, \iota_{\mathbb{X},E}, \tilde{\lambda}_{\mathbb{X}})$ over $\mathrm{Spec} \bar{k}$, where $(\mathbb{X}, \iota_{\mathbb{X},E})$ is the framing object for \mathcal{M}_E and $\tilde{\lambda}_{\mathbb{X}}$ is a polarization as in Theorem 5.2 (1). Then ϱ is an O_E -linear quasi-isogeny

$$\varrho : X \times_S \bar{S} \rightarrow \mathbb{X} \times_{\mathrm{Spec} \bar{k}} \bar{S},$$

of height 0 such that, locally on \bar{S} , the (quasi-)polarizations $\varrho^*(\tilde{\lambda}_{\mathbb{X}})$ and $\tilde{\lambda}$ on X only differ by a scalar in O_E^{\times} , i.e., there exists an element $\alpha \in O_E^{\times}$ such that $\varrho^*(\tilde{\lambda}_{\mathbb{X}}) = \tilde{\lambda} \circ \iota_E(\alpha)$. Two tuples $(X, \iota_E, \tilde{\lambda}, \varrho)$ and $(X', \iota'_E, \tilde{\lambda}', \varrho')$ are equivalent if there exists an O_E -linear isomorphism $\varphi : X \xrightarrow{\sim} X'$ such that $\varphi^*(\tilde{\lambda}')$ and $\tilde{\lambda}$ only differ by a scalar in O_E^{\times} .

In this way, we give a definition for $\mathcal{M}_{E,\mathrm{pol}}$ by introducing extra data on points of the moduli space \mathcal{M}_E , instead of extra conditions. It is now clear that $\mathcal{M}_{E,\mathrm{pol}}$ describes a moduli problem for p -divisible groups of PEL type. It is easily checked that the two descriptions of $\mathcal{M}_{E,\mathrm{pol}}$ give rise to the same moduli space.

Theorem 5.2 now holds if and only if $\mathcal{M}_{E,\mathrm{pol}} = \mathcal{M}_E$. This equality is a consequence of the following statement.

Lemma 5.6. *For any point $x = (X, \iota_E, \varrho) \in \mathcal{M}_{E,\mathrm{pol}}(\bar{k})$, the embedding $\mathcal{M}_{E,\mathrm{pol}} \hookrightarrow \mathcal{M}_E$ induces an isomorphism of completed local rings $\widehat{\mathcal{O}}_{\mathcal{M}_{E,\mathrm{pol}},x} \cong \widehat{\mathcal{O}}_{\mathcal{M}_E,x}$.*

For the proof of this lemma, we use the theory of local models; cf. [Rapoport and Zink 1996, Chapter 3]. We postpone the proof of this lemma to the end of this section and we first introduce the local models $\mathbf{M}_E^{\mathrm{loc}}$ and $\mathbf{M}_{E,\mathrm{pol}}^{\mathrm{loc}}$ for \mathcal{M}_E and $\mathcal{M}_{E,\mathrm{pol}}$.

Let C be a 4-dimensional F -vector space with an action of E and let $\Lambda \subseteq C$ be an O_F -lattice that is stable under the action of O_E . Furthermore, let (\cdot, \cdot) be an F -bilinear alternating form on C with

$$(5-2) \quad (\alpha x, y) = (x, \alpha y)$$

for all $\alpha \in E$ and $x, y \in C$ and such that Λ is unimodular with respect to (\cdot, \cdot) . It is easily checked that (\cdot, \cdot) is unique up to an isomorphism of C that commutes with the E -action and that maps Λ to itself.

For an O_F -algebra R , let $M_E^{\text{loc}}(R)$ be the set of all direct summands $\mathcal{F} \subseteq \Lambda \otimes_{O_F} R$ of rank 2 that are O_E -linear and satisfy the *Kottwitz condition*. That means, for all $\alpha \in O_E$, the action of α on the quotient $(\Lambda \otimes_{O_F} R)/\mathcal{F}$ has the characteristic polynomial

$$\text{char}(\text{Lie } X, T \mid \alpha) = (T - \alpha)(T - \bar{\alpha}).$$

The subset $M_{E,\text{pol}}^{\text{loc}}(R) \subseteq M_E^{\text{loc}}(R)$ consists of all direct summands $\mathcal{F} \in M_E^{\text{loc}}(R)$ that are in addition totally isotropic with respect to (\cdot, \cdot) on $\Lambda \otimes_{O_F} R$.

The functor M_E^{loc} is representable by a closed subscheme of $\text{Gr}(2, \Lambda)_{O_F}$, the Grassmannian of rank 2 direct summands of Λ , and $M_{E,\text{pol}}^{\text{loc}}$ is representable by a closed subscheme of M_E^{loc} . In particular, both M_E^{loc} and $M_{E,\text{pol}}^{\text{loc}}$ are projective schemes over $\text{Spec } O_F$.

These local models have already been studied by Deligne and Pappas. In particular, we have:

Proposition 5.7 [Deligne and Pappas 1994]. $M_{E,\text{pol}}^{\text{loc}} = M_E^{\text{loc}}$. In other words, for an O_F -algebra R , any direct summand $\mathcal{F} \in M_E^{\text{loc}}(R)$ is totally isotropic with respect to (\cdot, \cdot) .

The moduli spaces \mathcal{M}_E and $\mathcal{M}_{E,\text{pol}}$ are related to the local models M_E^{loc} and $M_{E,\text{pol}}^{\text{loc}}$ via local model diagrams; cf. [Rapoport and Zink 1996, Chapter 3]. Let $\mathcal{M}_E^{\text{large}}$ be the functor that maps a scheme $S \in \text{Nilp}_{\check{O}_F}$ to the set of isomorphism classes of tuples $(X, \iota_E, \varrho; \gamma)$. Here,


$$(X, \iota_E, \varrho) \in \mathcal{M}_E(S),$$

and γ is an O_E -linear isomorphism

$$\gamma : \mathbb{D}_X(S) \xrightarrow{\sim} \Lambda \otimes_{O_F} \mathcal{O}_S.$$

On the left-hand side, $\mathbb{D}_X(S)$ denotes the (relative) Grothendieck–Messing crystal of X evaluated at S ; cf. [Ahsendorf 2011, Section 5.2].

Let $\widehat{M}_E^{\text{loc}}$ be the π_0 -adic completion of $M_E^{\text{loc}} \otimes_{O_F} \check{O}_F$. Then there is a local model diagram:



The morphism f on the left-hand side is the projection $(X, \iota_E, \varrho; \gamma) \mapsto (X, \iota_E, \varrho)$. The morphism g on the right-hand side maps $(X, \iota_E, \varrho; \gamma) \in \mathcal{M}_E^{\mathrm{large}}(S)$ to

$$\mathcal{F} = \ker(\Lambda \otimes_{\mathcal{O}_F} \mathcal{O}_S \xrightarrow{\gamma^{-1}} \mathbb{D}_X(S) \rightarrow \mathrm{Lie} X) \subseteq \Lambda \otimes_{\mathcal{O}_F} \mathcal{O}_S.$$

By [Rapoport and Zink 1996, Theorem 3.11], the morphism f is smooth and surjective. The morphism g is formally smooth by Grothendieck–Messing theory; see [Messing 1972, V.1.6], resp. [Ahsendorf 2011, Chapter 5.2] for the relative setting (i.e., when $\mathcal{O}_F \neq \mathbb{Z}_p$).

We also have a local model diagram for the space $\mathcal{M}_{E,\mathrm{pol}}$. We define $\mathcal{M}_{E,\mathrm{pol}}^{\mathrm{large}}$ as the fiber product $\mathcal{M}_{E,\mathrm{pol}}^{\mathrm{large}} = \mathcal{M}_{E,\mathrm{pol}} \times_{\mathcal{M}_E} \mathcal{M}_E^{\mathrm{large}}$. Then $\mathcal{M}_{E,\mathrm{pol}}^{\mathrm{large}}$ is closed formal subscheme of $\mathcal{M}_E^{\mathrm{large}}$ with the following moduli description. A point $(X, \iota_E, \varrho; \gamma) \in \mathcal{M}_E^{\mathrm{large}}(S)$ lies in $\mathcal{M}_{E,\mathrm{pol}}^{\mathrm{large}}(S)$ if $\tilde{\lambda} = \varrho^*(\tilde{\lambda}_{\mathbb{X}})$ is a principal polarization on X . In that case, $\tilde{\lambda}$ induces an alternating form $(,)^X$ on $\mathbb{D}_X(S)$ which, under the isomorphism γ , is equal to the form $(,)$ on $\Lambda \otimes_{\mathcal{O}_F} \mathcal{O}_S$, up to a unit in $\mathcal{O}_E \otimes_{\mathcal{O}_F} \mathcal{O}_S$.

The local model diagram for $\mathcal{M}_{E,\mathrm{pol}}$ now looks as follows.

$$(5-3) \quad \begin{array}{ccc} & \mathcal{M}_{E,\mathrm{pol}}^{\mathrm{large}} & \\ f_{\mathrm{pol}} \swarrow & & \searrow g_{\mathrm{pol}} \\ \mathcal{M}_{E,\mathrm{pol}} & & \widehat{\mathcal{M}}_{E,\mathrm{pol}}^{\mathrm{loc}} \end{array}$$

Here, $\widehat{\mathcal{M}}_{E,\mathrm{pol}}^{\mathrm{loc}}$ is the π_0 -adic completion of $\mathcal{M}_{E,\mathrm{pol}}^{\mathrm{loc}} \otimes_{\mathcal{O}_F} \check{\mathcal{O}}_F$ and f_{pol} and g_{pol} are the restrictions of the morphisms f and g above. Again, g_{pol} is formally smooth by Grothendieck–Messing theory and f_{pol} is smooth and surjective by construction.

We can now finish the proof of Lemma 5.6.

Proof of Lemma 5.6. We have the following commutative diagram.

$$(5-4) \quad \begin{array}{ccccc} \mathcal{M}_{E,\mathrm{pol}} & \xleftarrow{f_{\mathrm{pol}}} & \mathcal{M}_{E,\mathrm{pol}}^{\mathrm{large}} & \xrightarrow{g_{\mathrm{pol}}} & \widehat{\mathcal{M}}_{E,\mathrm{pol}}^{\mathrm{loc}} \\ \downarrow & & \downarrow & & \parallel \\ \mathcal{M}_E & \xleftarrow{f} & \mathcal{M}_E^{\mathrm{large}} & \xrightarrow{g} & \widehat{\mathcal{M}}_E^{\mathrm{loc}} \end{array}$$

The equality on the right-hand side follows from Proposition 5.7. The other vertical arrows are closed embeddings.

Let $x \in \mathcal{M}_{E,\mathrm{pol}}(\bar{k})$. By [Rapoport and Zink 1996, Proposition 3.33], there exists an étale neighborhood U of x in \mathcal{M}_E and section $s : U \rightarrow \mathcal{M}_E^{\mathrm{large}}$ such that $g \circ s$ is formally étale. Similarly, $U_{\mathrm{pol}} = U \times_{\mathcal{M}_E} \mathcal{M}_{E,\mathrm{pol}}$ and s_{pol} is the base change of s to U_{pol} . Then the composition $g_{\mathrm{pol}} \circ s_{\mathrm{pol}}$ is also formally étale. These formally étale

maps induce isomorphisms of local rings

$$\widehat{\mathcal{O}}_{\mathcal{M}_E, x} \xrightarrow{\sim} \widehat{\mathcal{O}}_{\widehat{\mathcal{M}}_E^{\text{loc}}, x'} \quad \text{and} \quad \widehat{\mathcal{O}}_{\mathcal{M}_{E, \text{pol}}, x} \xrightarrow{\sim} \widehat{\mathcal{O}}_{\widehat{\mathcal{M}}_{E, \text{pol}}^{\text{loc}}, x'}, \quad x' = s(g(x)).$$

By [Proposition 5.7](#), we have $\widehat{\mathcal{O}}_{\widehat{\mathcal{M}}_E^{\text{loc}}, x'} = \widehat{\mathcal{O}}_{\widehat{\mathcal{M}}_{E, \text{pol}}^{\text{loc}}, x'}$ and since this identification commutes with $g \circ s$ (resp. $g_{\text{pol}} \circ s_{\text{pol}}$), we get the desired isomorphism $\widehat{\mathcal{O}}_{\mathcal{M}_{E, \text{pol}}, x} \cong \widehat{\mathcal{O}}_{\mathcal{M}_E, x}$. \square

Acknowledgements

First of all, I am very grateful to my advisor, M. Rapoport, for suggesting this topic and for his constant support and helpful discussions. I thank the members of our Arbeitsgruppe in Bonn for numerous discussions, and I also would like to thank the audience of my two AG talks for many helpful questions and comments. Furthermore, I owe thanks to A. Genestier for many useful remarks and for pointing out a mistake in an earlier version of this paper. I would also like to thank the referee for helpful comments.

This work is the author's PhD thesis at the University of Bonn, which was supported by the SFB/TR45 “Periods, Moduli Spaces and Arithmetic of Algebraic Varieties” of the DFG (German Research Foundation). Parts of this paper were written during the fall semester program “New Geometric Methods in Number Theory and Automorphic Forms” at the MSRI in Berkeley.

References

- [Ahsendorf 2011] T. Ahsendorf, [“ \$\mathbb{C}\$ -displays and \$\pi\$ -divisible formal \$\mathbb{C}\$ -modules](#)”, Ph.D. thesis, Universität Bielefeld, 2011, available at <http://tinyurl.com/ahsendorf>.
- [Ahsendorf et al. 2016] T. Ahsendorf, C. Cheng, and T. Zink, [“ \$\mathbb{C}\$ -displays and \$\pi\$ -divisible formal \$\mathbb{C}\$ -modules](#)”, *J. Algebra* **457** (2016), 129–193. [MR](#) [Zbl](#)
- [Boutot and Carayol 1991] J.-F. Boutot and H. Carayol, “Uniformisation p -adique des courbes de Shimura: les théorèmes de Cherednik et de Drinfeld”, pp. 45–158 in *Courbes modulaires et courbes de Shimura* (Orsay, 1987/1988), Astérisque **196-197**, Société Mathématique de France, Paris, 1991. [MR](#) [Zbl](#)
- [Büttel and Pappas 2017] O. Büttel and G. Pappas, “ (G, μ) -displays and Rapoport–Zink spaces”, preprint, 2017. [arXiv](#)
- [Chen and Viehmann 2015] M. Chen and E. Viehmann, “Affine Deligne–Lusztig varieties and the action of J ”, 2015. To appear in *J. Alg. Geom.* [arXiv](#)
- [Deligne and Pappas 1994] P. Deligne and G. Pappas, [“Singularités des espaces de modules de Hilbert, en les caractéristiques divisant le discriminant”](#), *Compositio Math.* **90**:1 (1994), 59–79. [MR](#) [Zbl](#)
- [Drinfeld 1976] V. G. Drinfeld, “Coverings of p -adic symmetric domains”, *Funkcional. Anal. i Prilozhen.* **10**:2 (1976), 29–40. In Russian; translated in *Funct. Anal. Appl.* **10**:2 (1976), 107–115. [MR](#) [Zbl](#)
- [Howard and Pappas 2017] B. Howard and G. Pappas, [“Rapoport–Zink spaces for spinor groups”](#), *Compositio Math.* **153**:5 (2017), 1050–1118. [MR](#) [Zbl](#)

- [Jacobowitz 1962] R. Jacobowitz, “[Hermitian forms over local fields](#)”, *Amer. J. Math.* **84** (1962), 441–465. [MR](#) [Zbl](#)
- [Kim 2013] W. Kim, “Rapoport–Zink spaces of Hodge type”, preprint, 2013. [arXiv](#)
- [Kudla and Rapoport 2014] S. Kudla and M. Rapoport, “[An alternative description of the Drinfeld \$p\$ -adic half-plane](#)”, *Ann. Inst. Fourier (Grenoble)* **64**:3 (2014), 1203–1228. [MR](#) [Zbl](#)
- [Kudla et al. 2006] S. S. Kudla, M. Rapoport, and T. Yang, *[Modular forms and special cycles on Shimura curves](#)*, Annals of Mathematics Studies **161**, Princeton Univ. Press, 2006. [MR](#) [Zbl](#)
- [Messing 1972] W. Messing, *The crystals associated to Barsotti–Tate groups: with applications to abelian schemes*, Lecture Notes in Math. **264**, Springer, Berlin, 1972. [MR](#) [Zbl](#)
- [Neukirch et al. 2008] J. Neukirch, A. Schmidt, and K. Wingberg, *Cohomology of number fields*, 2nd ed., Grundlehren der Math. Wissenschaften **323**, Springer, Berlin, 2008. [MR](#) [Zbl](#)
- [Pappas et al. 2013] G. Pappas, M. Rapoport, and B. Smithling, “Local models of Shimura varieties, I: Geometry and combinatorics”, pp. 135–217 in *Handbook of moduli, III*, edited by G. Farkas and I. Morrison, Adv. Lect. Math. (ALM) **26**, Int. Press, Somerville, MA, 2013. [MR](#) [Zbl](#)
- [Rapoport and Viehmann 2014] M. Rapoport and E. Viehmann, “Towards a theory of local Shimura varieties”, *Münster J. Math.* **7**:1 (2014), 273–326. [MR](#) [Zbl](#)
- [Rapoport and Zink 1996] M. Rapoport and T. Zink, *Period spaces for p -divisible groups*, Annals of Mathematics Studies **141**, Princeton Univ. Press, 1996. [MR](#) [Zbl](#)
- [Rapoport and Zink 2017] M. Rapoport and T. Zink, “[On the Drinfeld moduli problem of \$p\$ -divisible groups](#)”, *Camb. J. Math.* **5**:2 (2017), 229–279. [MR](#) [Zbl](#)
- [Vignéras 1980] M.-F. Vignéras, *Arithmétique des algèbres de quaternions*, Lecture Notes in Math. **800**, Springer, Berlin, 1980. [MR](#) [Zbl](#)
- [Vollaard and Wedhorn 2011] I. Vollaard and T. Wedhorn, “[The supersingular locus of the Shimura variety of \$\mathrm{GU}\(1, n - 1\)\$, II](#)”, *Invent. Math.* **184**:3 (2011), 591–627. [MR](#) [Zbl](#)

Received May 24, 2017.

DANIEL KIRCH

INSTITUT DE MATHÉMATIQUES DE JUSSIEU - PARIS RIVE GAUCHE

IMJ-PRG

PARIS

FRANCE

dkirch@math.uni-bonn.de

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 293 No. 2 April 2018

Sums of CR functions from competing CR structures	257
DAVID E. BARRETT and DUSTY E. GRUNDMEIER	
On Tate duality and a projective scalar property for symmetric algebras	277
FLORIAN EISELE, MICHAEL GELINE, RADHA KESSAR and MARKUS LINCKELMANN	
Coaction functors, II	301
S. KALISZEWSKI, MAGNUS B. LANDSTAD and JOHN QUIGG	
Construction of a Rapoport–Zink space for $\mathrm{GU}(1, 1)$ in the ramified 2-adic case	341
DANIEL KIRCH	
Group and round quadratic forms	391
JAMES O’SHEA	
Dual operator algebras close to injective von Neumann algebras	407
JEAN ROYDOR	
Scalar curvature and singular metrics	427
YUGUANG SHI and LUEN-FAI TAM	
On the differentiability issue of the drift-diffusion equation with nonlocal Lévy-type diffusion	471
LIUTANG XUE and ZHUAN YE	