

*Pacific
Journal of
Mathematics*

GROUP AND ROUND QUADRATIC FORMS

JAMES O'SHEA

Volume 293 No. 2

April 2018

GROUP AND ROUND QUADRATIC FORMS

JAMES O'SHEA

We offer some elementary characterisations of group and round quadratic forms. These characterisations are applied to establish new (and recover existing) characterisations of Pfister forms. We establish “going-up” results for group and anisotropic round forms with respect to iterated Laurent series field extensions, which contrast with the established results with respect to rational function field extensions. For forms of two-power dimension, we determine when there exists a field extension over which the form becomes an anisotropic group form that is not round.

1. Introduction

A quadratic form is round if its value set coincides with the multiplicative group of similarity factors associated with the form. Thus, round forms constitute a prominent subclass of group forms, forms whose value sets are multiplicative groups. As roundness is one of the fundamental properties of Pfister forms, the class of forms that occupy a central role in quadratic form theory, it is unsurprising that this notion has had a number of important consequences. However, while the structure and behaviour of round forms has received extensive treatment in the literature, this class of forms is still not fully understood and, as suggested in [Lam 2005], merits further study. The broader class of group forms is, comparatively, little understood.

Our opening results, which are invoked throughout this article, record elementary characterisations of the classes of group and round forms (see Proposition 2.2 to Corollary 2.5). In Section 2, we apply these results to obtain new characterisations of Pfister forms (see Theorem 2.7), in addition to reproving established ones (see Corollary 2.8), and to extend Elman's classification of odd-dimensional round forms in accordance with our broader definition of roundness.

The group and round properties of a form are intrinsically linked to its base field of definition, and thus are sensitive to scalar extension. Alpers [1991] remarks that while general “going-down” results exist with respect to roundness, with round forms over odd-degree extensions being round over their base fields for example, no general results are known in the “going-up” direction. We establish such results

MSC2010: 11E04, 11E10, 11E81, 11E99, 12F20.

Keywords: quadratic forms, round forms, group forms, Pfister forms.

for group and anisotropic round forms with respect to iterated Laurent series fields (see Corollary 3.4), highlighting an interesting divergence in the behaviour of forms under extension to iterated Laurent series fields as opposed to rational function fields (see Remark 3.5).

Hsia and Johnson [1973a; 1973b] studied the problem of distinguishing between anisotropic group and round forms over local and global fields. In this spirit, we consider the following general question:

Question 1.1. For q an anisotropic form over F , does there exist an extension K/F such that q_K is an anisotropic group form that is not round?

Our characterisation of group forms allows for the construction of a generic field extension over which a form becomes an anisotropic group form. Thus, the adoption of Merkurjev's method of passing to iterated field extensions obtained by composing function fields of quadratic forms represents the natural approach to addressing the above question. While highlighting an obstruction to resolving Question 1.1 in general (see Proposition 3.6), we can employ this method to good effect in certain situations. In particular, Theorem 3.8 represents a complete answer to Question 1.1 with respect to forms of two-power dimension.

We let F denote a field of characteristic different from two, and recall that every nondegenerate quadratic form on a vector space over F can be diagonalised. We write $\langle a_1, \dots, a_n \rangle$ to denote the (n -dimensional) quadratic form on an n -dimensional F -vector space defined by $a_1, \dots, a_n \in F^\times$. We use the term "form" to refer to a nondegenerate quadratic form of positive dimension. If p and q are forms over F , we denote by $p \perp q$ their orthogonal sum and by $p \otimes q$ their tensor product. We use aq to denote $\langle a \rangle \otimes q$ for $a \in F^\times$. We write $p \simeq q$ to indicate that p and q are isometric, and say that p and q are *similar* if $p \simeq aq$ for some $a \in F^\times$. A form p is a *subform* of q if $q \simeq p \perp r$ for some form r , in which case we write $p \subseteq q$. For q a form over F and K/F a field extension, we will often employ the notation q_K when viewing q as a form over K via the canonical embedding. A form q represents $a \in F$ if there exists a vector v such that $q(v) = a$. We denote by $D_F(q)$ the set of values in F^\times represented by q . A form over F is *isotropic* if it represents zero nontrivially, and *anisotropic* otherwise. Every form q has a decomposition $q \simeq q_{\text{an}} \perp i(q) \times \langle 1, -1 \rangle$ where the anisotropic quadratic form q_{an} and the nonnegative integer $i(q)$ are uniquely determined. If a form q is isotropic over F , then $D_F(q) = F^\times$, as $((a+1)/2)^2 - ((a-1)/2)^2 = a$ for all $a \in F^\times$. A form q is *hyperbolic* if $i(q) = \frac{1}{2} \dim q$.

A form q is a *group form* over F if $D_F(q)$ is a subgroup of F^\times . The similarity factors of q constitute the group $G_F(q) = \{a \in F^\times \mid aq \simeq q\}$. Equivalently, $G_F(q) = \{a \in F^\times \mid \langle 1, -a \rangle \otimes q \text{ is hyperbolic}\}$. A group form q over F is said to be *round* if $D_F(q) = G_F(q)$. Equivalently, a form q is round over F if $D_F(q) \subseteq G_F(q)$, as if

$a \in D_F(q) \subseteq G_F(q)$, then $aq \simeq q$, whereby $1 \in D_F(q)$ and thus $G_F(q) \subseteq D_F(q)$. We use $H_F(q)$ to denote the set of products of two elements of $D_F(q)$. Per Lemma 2.1, we have that $H_F(q) = \{a \in F^\times \mid \langle 1, -a \rangle \otimes q \text{ is isotropic}\}$. For $n \in \mathbb{N}$, an n -fold Pfister form over F is a form isometric to $\langle 1, a_1 \rangle \otimes \cdots \otimes \langle 1, a_n \rangle$ for some $a_1, \dots, a_n \in F^\times$ (the form $\langle 1 \rangle$ is the 0-fold Pfister form). Isotropic Pfister forms are hyperbolic [Lam 2005, Theorem X.1.7]. Pfister forms are round (see [Lam 2005, Theorem X.1.8]). A form τ is a Pfister neighbour if τ is similar to a subform of a Pfister form π and $\dim \tau > \frac{1}{2} \dim \pi$.

We recall that every nonzero square class in $F((x))$, the Laurent series field in the variable x over F , can be represented by a or ax for some $a \in F^\times$, whereby every form over $F((x))$ can be written as $p \perp xq$ for p and q forms over F . We will often invoke the following folkloric result regarding isotropy over Laurent series fields.

Lemma 1.2. *Let p and q be forms over F . Considering $p \perp xq$ as a form over $F((x))$, we have that $i(p \perp xq) = i(p) + i(q)$.*

For a form q over F with $\dim q = n \geq 2$ and $q \not\simeq \langle 1, -1 \rangle$, the function field $F(q)$ of q is the quotient field of the integral domain $F[X_1, \dots, X_n]/(q(X_1, \dots, X_n))$ (this is the function field of the affine quadric $q(X) = 0$ over F). To avoid case distinctions, we set $F(q) = F$ if $\dim q = 1$ or $q \simeq \langle 1, -1 \rangle$. For q a form over F , we note the inclusion $F((x))(q) \subseteq F(q)((x))$, which we will apply in combination with Lemma 1.2. For all forms p over F and all extensions K/F such that q_K is isotropic, we have that $i(p_{F(q)}) \leq i(p_K)$ in accordance with Knebusch’s specialisation results [1976, Proposition 3.1 and Theorem 3.3]. In particular, letting $i_1(q)$ denote $i(q_{F(q)})$, we have that $i_1(q) \leq i(q_K)$ for all extensions K/F such that q_K is isotropic. Invoking the Cassels–Pfister Subform Theorem [Lam 2005, Theorem X.4.5] of Wadsworth [1975, Theorem 2] and Knebusch [1976, Lemma 4.5], for p and q anisotropic forms over F of dimension at least two such that $p_{F(q)}$ is hyperbolic, one has that $aq \subseteq bp$ for all $a \in D_F(q)$ and $b \in D_F(p)$. For q an anisotropic form over F of dimension at least two, it is known that $q_{F(q)}$ is hyperbolic if and only if q is similar to a Pfister form over F by [Lam 2005, Theorem X.4.14], a result of Wadsworth [1975, Theorem 5] and Knebusch [1976, Theorem 5.8]. We will regularly invoke Hoffmann’s separation theorem [1995, Theorem 1], which we recall below.

Theorem 1.3. *Let p and q be forms over F such that p is anisotropic. If*

$$\dim p \leq 2^n < \dim q$$

for some integer $n \geq 0$, then $p_{F(q)}$ is anisotropic.

In accordance with the above theorem and [Lam 2005, Exercise I.16], for q an anisotropic form over F , we note that $\dim q$ and $\dim q - i_1(q)$ belong to an interval of the form $[2^n, 2^{n+1}]$ for some $n \in \mathbb{N} \cup \{0\}$. We will also invoke the following isotropy criterion of Karpenko and Merkurjev [2003, Theorem 4.1].

Theorem 1.4. For p and q anisotropic forms over F such that $p_{F(q)}$ is isotropic,

- (i) $\dim p - i_1(p) \geq \dim q - i_1(q)$;
- (ii) $\dim p - i_1(p) = \dim q - i_1(q)$ if and only if $q_{F(p)}$ is isotropic.

We refer the reader to works by Vishik [2011] and Scully [2016a] for recent results in the spirit of Theorem 1.3 and Theorem 1.4.

2. Characterisations of group, round and Pfister forms

As above, the forms we consider are nondegenerate and of positive dimension over fields of characteristic different from two. In accordance with the associated definitions, we begin our study of the properties of a form q over F being group or round by considering the value set $D_F(q)$, the group of similarity factors $G_F(q)$, and the set of products of two elements of $D_F(q)$, usually denoted by $D_F(q)D_F(q)$.

Roussey [2005, Lemme 2.5.4] proved the following result in his thesis.

Lemma 2.1. For p and q forms over F , we have that

$$D_F(p)D_F(q) = \{a \in F^\times \mid p \perp -aq \text{ is isotropic}\}.$$

Proof. The statement clearly holds if either p or q is isotropic. Thus, assuming that p and q are anisotropic, we have that $p \perp -aq$ is isotropic if and only if there exist nonzero vectors v and w such that $p(v) - aq(w) = 0$. Thus, $p(v) = aq(w) \neq 0$, whereby $a = p(v)(q(w))^{-1}$. Hence,

$$a = p(v) \frac{1}{q(w)} = p(v)q\left(\frac{w}{q(w)}\right) \in D_F(p)D_F(q).$$

As $1 \in D_F(dq)$ for $d \in D_F(q)$, we have $p \perp -cdq$ is isotropic for $c \in D_F(p)$. \square

We let $H_F(q) = \{a \in F^\times \mid \langle 1, -a \rangle \otimes q \text{ is isotropic}\}$, giving $H_F(q) = D_F(q)D_F(q)$ in accordance with Lemma 2.1. As with $D_F(q)$ and $G_F(q)$, we may restrict our attention to the square classes contained in $H_F(q)$, since $H_F(q) = (F^\times)^2 H_F(q)$. Clearly we have that $(F^\times)^2 \subseteq G_F(q) \subseteq H_F(q)$ for all forms q over F . Moreover, if $1 \in D_F(q)$, then

$$(F^\times)^2 \subseteq G_F(q) \subseteq D_F(q) \subseteq H_F(q).$$

If q is isotropic over F , then q is a group form over F , with $D_F(q) = F^\times = H_F(q)$ in this case. Our opening result records that Lemma 2.1 may be applied to characterise group forms.

Proposition 2.2. A form q is a group form over F if and only if $H_F(q) \subseteq D_F(q)$.

Proof. Letting $a \in D_F(q)$, there exists a nonzero vector v such that $q(v) = a$, whereby $q(a^{-1}v) = (a^{-1})^2 q(v) = a^{-1} \in D_F(q)$. Thus, q is a group form over F if and only if $D_F(q)D_F(q) \subseteq D_F(q)$, so the result follows by invoking Lemma 2.1. \square

As group forms represent one, we thus obtain the following corollary.

Corollary 2.3. *A form q is a group form over F if and only if $H_F(q) = D_F(q)$.*

In accordance with our definition of roundness, if q is isotropic over F , then q is round over F if and only if q is hyperbolic or the nonzero form q_{an} is such that $D_F(q_{\text{an}}) = F^\times = G_F(q_{\text{an}})$. This observation follows from the fact that $G_F(q) = G_F(q_{\text{an}})$, in accordance with Witt cancellation and the fact that $D_F(q) = F^\times$ for q isotropic over F . Per [Lam 2005, Example X.1.15(5)], the form $q \simeq \langle 1, -1, 1, 1 \rangle$ over \mathbb{F}_3 is an example of an isotropic round form that is not hyperbolic.

Corollary 2.4. *A form q is round over F if and only if 1 is an element of $D_F(q)$ and $H_F(q) \subseteq G_F(q)$.*

Proof. If q is round over F , then $1 \in D_F(q) = G_F(q)$ and q is a group form over F . Invoking Proposition 2.2, it follows that $H_F(q) \subseteq D_F(q)$, whereby $H_F(q) \subseteq G_F(q)$.

Conversely, as $1 \in D_F(q)$, we recall that $G_F(q) \subseteq D_F(q) \subseteq H_F(q)$, whereby the equality $D_F(q) = G_F(q)$ follows from the assumption that $H_F(q) \subseteq G_F(q)$. \square

We note that, for q a round form over F , the inclusion $H_F(q) \subseteq G_F(q)$ can also be derived from [Wadsworth and Shapiro 1977, Proposition 1 and Theorem 1].

Addressing the question of distinguishing between the classes of group and round forms over a given field, it is reasonable to restrict one’s consideration to those forms that represent one, whereby the preceding characterisation may be simplified.

Corollary 2.5. *Let q be a form such that $1 \in D_F(q)$. The following are equivalent:*

- (i) q is round over F .
- (ii) $H_F(q) \subseteq G_F(q)$.
- (iii) $q \otimes \rho$ is anisotropic or hyperbolic for every 1-fold Pfister form ρ over F .
- (iv) $q \otimes \beta$ is anisotropic or hyperbolic for every 2-dimensional form β over F .
- (v) $q \otimes \pi$ is anisotropic or hyperbolic for every n -fold Pfister form π over F , for $n \in \mathbb{N}$.

Proof. Statements (i), (ii) and (iii) are equivalent by Corollary 2.4. Statements (iii) and (iv) are equivalent, as scaling does not affect isotropy. Statement (v) clearly implies (iii). Assuming (i), it follows that $q \otimes \pi$ is round for every Pfister form π over F , by Witt’s round form theorem [Lam 2005, Theorem X.1.14]. By repeatedly invoking statement (iii), we see that (v) follows. \square

Next, note that scalar multiples of Pfister forms representing one are Pfister forms.

Lemma 2.6. *A form q over F is a Pfister form if and only if q is similar to a Pfister form and $1 \in D_F(q)$.*

Proof. To establish the right-to-left implication, we let $q \simeq a\pi$ for $a \in F^\times$ and π a Pfister form over F . As $1 \in D_F(q)$, it follows that $a \in D_F(\pi)$, whereby $q \simeq \pi$ as π is round. \square

We can apply the above characterisations of round and group forms to obtain a new characterisation of Pfister forms.

Theorem 2.7. *Let q be an anisotropic form. The following are equivalent:*

- (i) q is a Pfister form over F .
- (ii) q is a round form over $K = F((x))(q \otimes \langle 1, -x \rangle)$.
- (iii) q is a group form over $K = F((x))(q \otimes \langle 1, -x \rangle)$.

Proof. As Pfister forms are round, and round forms are group, it suffices to prove that (iii) implies (i).

The field K is the function field of $q \otimes \langle 1, -x \rangle$ over $F((x))$, which is an anisotropic form in accordance with Lemma 1.2. We will first show that

$$D_K(q) \cap F^\times = D_F(q).$$

Let $a \in F^\times$ be such that $q \perp \langle -a \rangle$ is anisotropic over F and suppose, for the sake of contradiction, that $q \perp \langle -a \rangle$ is isotropic over K . Invoking Theorem 1.4 (i),

$$\dim(q \perp \langle -a \rangle) - i_1(q \perp \langle -a \rangle) \geq \dim(q \otimes \langle 1, -x \rangle) - i_1(q \otimes \langle 1, -x \rangle).$$

In accordance with Theorem 1.3 and [Lam 2005, Exercise I.16], there exists $n \in \mathbb{N}$ such that

$$\dim(q \perp \langle -a \rangle) - i_1(q \perp \langle -a \rangle) = \dim(q \otimes \langle 1, -x \rangle) - i_1(q \otimes \langle 1, -x \rangle) = 2^n,$$

whereby $\dim q = 2^n$. Hence, $q \otimes \langle 1, -x \rangle$ is isotropic over $F((x))(q \perp \langle -a \rangle)$, in accordance with Theorem 1.4 (ii). Invoking [Izhboldin 2000, Lemma 5.4 (3)], it thus follows that q is isotropic over $F(q \perp \langle -a \rangle)$. However, as $\dim q = 2^n$, this contradicts Theorem 1.3, thereby establishing the claim.

We have that $1 \in D_K(q)$ by assumption, whereby $1 \in D_F(q)$ by the statement proven above. As $x \in H_K(q)$ by construction, it follows that $x \in D_K(q)$ in accordance with Proposition 2.2, whereby the form $q \perp \langle -x \rangle$ becomes isotropic over $F((x))(q \otimes \langle 1, -x \rangle)$. Arguing as above, it follows that, for some $n \in \mathbb{N}$,

$$\dim(q \perp \langle -x \rangle) - i_1(q \perp \langle -x \rangle) = \dim(q \otimes \langle 1, -x \rangle) - i_1(q \otimes \langle 1, -x \rangle) = 2^n,$$

and $\dim q = 2^n$. Hence, $i_1(q \otimes \langle 1, -x \rangle) = \dim q$, in accordance with this equality, whereby $q \otimes \langle 1, -x \rangle$ becomes hyperbolic over $F((x))(q \otimes \langle 1, -x \rangle)$. Invoking [O'Shea 2016, Proposition 3.2], it follows that q is hyperbolic over $F(q)$. Thus, q is similar to a Pfister form over F by [Lam 2005, Theorem X.4.14]. As $1 \in D_F(q)$, the result follows by invoking Lemma 2.6. □

We can invoke the above result to reprove the following characterisations of Pfister forms due to Pfister (see [Pfister 1965, Satz 5 and Theorem 2], [Scharlau 1985, Theorem 4.4 p.153] or [Elman et al. 2008, Theorem 23.2]).

Corollary 2.8. *Let q be an anisotropic form over F . The following are equivalent:*

- (i) q is a Pfister form over F .
- (ii) q is round over K for every extension K/F .
- (iii) q is group over K for every extension K/F .

Similarly, we obtain the following corollary of Theorem 2.7.

Corollary 2.9. *For q an anisotropic form over F , let $K = F((x))(q \otimes \langle 1, -x \rangle)$.*

- (i) *Then q is group over every extension of F if and only if q is group over K , and*
- (ii) *q is round over every extension of F if and only if q is round over K .*

Remark 2.10. Per [Scharlau 1985, Theorem 4.4, p.153], Pfister established that, for q an anisotropic form of dimension n , the three statements in Corollary 2.8 are equivalent to each of the following statements:

- (i) $q(x_1, \dots, x_n) \in G_K(q)$ for $K = F(x_1, \dots, x_n)$.
- (ii) $q(x_1, \dots, x_n)q(x_{n+1}, \dots, x_{2n}) \in D_K(q)$ for $K = F(x_1, \dots, x_{2n})$.

Thus, for q an anisotropic form of dimension n , it follows that q is a Pfister form over F if and only if q is a round form over $F(x_1, \dots, x_n)$, and that q is a Pfister form over F if and only if q is a group form over $F(x_1, \dots, x_{2n})$.

In a similar spirit to the preceding results, we offer the following characterisation of scalar multiples of Pfister forms.

Proposition 2.11. *Letting q be an anisotropic form over F , the following are equivalent:*

- (i) q is similar to a Pfister form over F .
- (ii) q_K is round for all extensions K/F such that $1 \in D_K(q)$.
- (iii) q_K is round for all extensions K/F such that q_K is isotropic.

Proof. Assuming (i), Lemma 2.6 implies that q_K is a Pfister form for K/F such that $1 \in D_K(q)$, whereby (ii) follows. As (ii) clearly implies (iii), it suffices to show that (iii) implies (i).

Letting $K = F(q)((x))$, we have that

$$D_K(q) = K^\times = G_K(q).$$

As $x \in G_K(q)$, the form $q \otimes \langle 1, -x \rangle$ is hyperbolic over $F(q)((x))$. Invoking Lemma 1.2, it follows that q is hyperbolic over $F(q)$. Thus, q is similar to a Pfister form over F by [Lam 2005, Theorem X.4.14]. \square

We conclude this section by characterising the odd-dimensional round forms (see Elman's characterisation [1973] of odd-dimensional round forms in the situation where isotropic round forms are defined to be hyperbolic).

Proposition 2.12. *Let q be a form over F .*

- (i) *If $D_F(q) = (F^\times)^2$, then q is round over F .*
- (ii) *If $H_F(q) \neq (F^\times)^2$ and q is round over F , then q is even-dimensional.*

Proof. (i) If $D_F(q) = (F^\times)^2$, then $D_F(q) \subseteq G_F(q)$, whereby q is round over F .

(ii) Let $a \in H_F(q) \setminus (F^\times)^2$. As q is round over F , we have that $H_F(q) \subseteq G_F(q)$ by Corollary 2.4, whereby $q \perp -aq$ is hyperbolic over F . As a comparison of determinants yields the contradiction that $a \in (F^\times)^2$ for q odd-dimensional, the result follows. □

Adapting Elman's proof of [Elman 1973, Lemma], we obtain the following result as a corollary of Proposition 2.12.

Corollary 2.13. *Let q be an odd-dimensional form over F . If q is round, then*

$$q \simeq (2r + 1) \times \langle 1 \rangle$$

for some $r \in \mathbb{N} \cup \{0\}$. Moreover, the following are equivalent:

- (i) $(2k + 1) \times \langle 1 \rangle$ is round over F for some $k \in \mathbb{N}$,
- (ii) $(2n + 1) \times \langle 1 \rangle$ is round over F for every $n \in \mathbb{N} \cup \{0\}$,
- (iii) F is Pythagorean.

Corollary 2.14. *Let q be an odd-dimensional isotropic form over F . Then q is round over F if and only if F is quadratically closed.*

Proof. If q is round, then $D_F(q) = (F^\times)^2$ by Proposition 2.12. As q is isotropic, it follows that $D_F(q) = F^\times$, whereby F is quadratically closed.

If F is quadratically closed, then $q_{\text{an}} \simeq \langle 1 \rangle$, whereby q is round. □

Corollary 2.15. *If q is an odd-dimensional anisotropic round form over F , then $q \otimes \beta$ is anisotropic over F for every anisotropic 2-dimensional form β over F .*

Proof. Let $\beta \simeq b\langle 1, -a \rangle$ be anisotropic over F for $a, b \in F^\times$. Suppose, for the sake of contradiction, that $q \otimes \beta$ is isotropic over F . Hence, $q \otimes \langle 1, -a \rangle$ is hyperbolic over F by Corollary 2.4. By repeatedly invoking [Elman and Lam 1973, Proposition 2.2], it follows that there exist binary forms β_1, \dots, β_n over F such that $\beta_i \otimes \langle 1, -a \rangle$ is hyperbolic over F for $i = 1, \dots, n$ and such that

$$q \simeq \beta_1 \perp \dots \perp \beta_n,$$

whereby q is even-dimensional, a contradiction. □

Remark 2.16. The preceding result can also be derived from the fact that F is Pythagorean and real, whereby its Witt ring is torsion free (see [Elman et al. 2008, Theorem 23.2]).

3. Group and round forms over field extensions

Alpers [1991] considers roundness with respect to algebraic extensions, establishing “going-down” and “going-up” results in certain situations. In particular, he remarks that a general going-down result holds for odd-degree extensions by Springer’s theorem [Springer 1952] (see [Lam 2005, Theorem VII.2.7]). We generalise this remark below.

Proposition 3.1. *Let q be a form over F and let K be an extension of F .*

- (i) *Suppose that $D_K(q) \cap F^\times \subseteq D_F(q)$. If q_K is a group form, then q is a group form over F .*
- (ii) *Suppose that every anisotropic form over F of dimension at most $\dim q + 1$ is anisotropic over K . If q_K is a round form, then q is a round form over F .*

Proof. (i) As $D_K(q) \cap F^\times = D_F(q)$ follows from the assumption, if $D_K(q)$ is a group it readily follows that $D_F(q)$ is a group.

(ii) If q is anisotropic over F , the assumption on K implies that $D_K(q) \cap F^\times = D_F(q)$ and $G_K(q) \cap F^\times = G_F(q)$. Applying this argument to q_{an} in the case where q is isotropic over F , the result follows. \square

Thus, as a consequence of the above, group and round forms satisfy going-down results with respect to purely transcendental extensions. Per Remark 2.10, going-up results do not hold for group or round forms with respect to rational function fields. However, we do have the following result with respect to Laurent series fields:

Proposition 3.2. *Let q be a form over F and let $K = F((x))$.*

- (i) *q is a group form over F if and only if q is a group form over K .*
- (ii) *If q is anisotropic, then q is round over F if and only if q is round over K .*

Proof. We remark that anisotropic forms over F are anisotropic over K .

(i) As isotropic forms are trivially group, we may assume, without loss of generality, that q is anisotropic over F . We consider the set $H_K(q)$, recalling that every nonzero square class in K can be represented by a or ax for some $a \in F^\times$. Invoking Lemma 1.2, it is apparent that $q \perp -axq$ is anisotropic over K for $a \in F^\times$. For $a \in F^\times$ such that $q \perp -aq$ is isotropic over K , it follows that $q \perp -aq$ is isotropic over F , whereby $q \perp \langle -a \rangle$ is isotropic over F by Proposition 2.2. Thus, as $q \perp \langle -a \rangle$ is isotropic over K , it follows from Proposition 2.2 that q is a group form over K .

For the converse statement, we may invoke Proposition 3.1 (i).

(ii) As $1 \in D_F(q)$ if and only if $1 \in D_K(q)$, we may argue as in the preceding proof of (i), with respect to Corollary 2.4 as opposed to Proposition 2.2, to establish the “only if” statement. The “if” statement can be established by invoking Proposition 3.1 (ii). \square

Remark 3.3. We note the necessity of the restriction to anisotropic round forms in statement (ii) of the above result. If q is isotropic and round over $K = F((x))$, then it readily follows that q is isotropic and round over F , but the converse does not hold in general. In particular, the isotropic form $q \simeq \langle 1, -1, 1, 1 \rangle$ is round over \mathbb{F}_3 but it is not round over $\mathbb{F}_3((x))$, as $x \notin D_K((q_K)_{\text{an}})$ in accordance with Lemma 1.2.

Iterating the above, we obtain the following result.

Corollary 3.4. *Let q be a form over F and let $K = F((x_1)) \cdots ((x_n))$ for $n \in \mathbb{N}$.*

- (i) *q is a group form over F if and only if q is a group form over K .*
- (ii) *If q is anisotropic, then q is round over F if and only if q is round over K .*

Remark 3.5. We note that the above result demonstrates a divergence in the behaviour, with respect to the properties of being group or round, of forms over F extended to iterated Laurent series fields as opposed to rational function fields. Corollary 3.4 contrasts with Pfister's result, per Remark 2.10, that an anisotropic form of dimension n over F is a round form over $F(x_1, \dots, x_n)$ if and only if it is a group form over $F(x_1, \dots, x_{2n})$ if and only if it is a Pfister form over F .

Motivated by the problem of distinguishing between anisotropic group and round forms, as studied over particular fields in [Hsia and Johnson 1973a; Hsia and Johnson 1973b], the rest of this section is devoted to addressing Question 1.1.

In accordance with Corollary 2.5, if q is an anisotropic group form over F , one can resolve Question 1.1 by determining whether there exists a Pfister form π over F such that $q \otimes \pi$ is isotropic but not hyperbolic. If such a form q is odd-dimensional, Question 1.1 further reduces to the problem of determining whether $H_F(q) \setminus (F^\times)^2$ is empty, in accordance with Proposition 2.12.

The natural approach towards answering Question 1.1 in the case where q is not a group form over F is to consider its extension to the generic extension K/F such that q_K is a group form. However, one encounters the following obstruction:

Proposition 3.6. *Let q be an anisotropic form over F . If there exists $a \in H_F(q)$ such that $i_1(q \perp \langle -a \rangle) > 1$, then there does not exist an extension K/F such that q_K is an anisotropic group form.*

Proof. Let K/F be an extension such that q_K is a group form. Since

$$a \in H_F(q) \subseteq H_K(q),$$

it follows that $a \in D_K(q)$ by Corollary 2.3, whereby $q \perp \langle -a \rangle$ is isotropic over K . Since $i_1(q \perp \langle -a \rangle) > 1$, it follows that $i((q \perp \langle -a \rangle)_K) > 1$, whereby q_K is isotropic (see [Lam 2005, Exercise I.16]). □

The following example illustrates that, provided that $\dim q \neq 2^n$ for $n \in \mathbb{N} \cup \{0\}$, there exist fields F , forms q over F and scalars $a \in F^\times$ that satisfy the hypotheses of Proposition 3.6.

Example 3.7. Let L a field and $a \in L^\times$ be such that $q \perp \langle -a \rangle$ is an anisotropic Pfister neighbour, where $\dim q \neq 2^n$ for $n \in \mathbb{N} \cup \{0\}$. Letting $F = L(q \perp -aq)$, we have that $a \in H_F(q)$ and that $q \perp \langle -a \rangle$ is anisotropic over F , by Theorem 1.3. As $q \perp \langle -a \rangle$ is a Pfister neighbour of dimension $\neq 2^n + 1$ for $n \in \mathbb{N} \cup \{0\}$, it follows that $i_1(q \perp \langle -a \rangle) > 1$.

In contrast with the above example, letting q be an arbitrary anisotropic form over F of dimension 2^n for some $n \in \mathbb{N}$, the proof of the following theorem demonstrates that there does exist an extension K/F such that q_K is an anisotropic group form. Moreover, when possible, we can find an extension K/F such that q_K is an anisotropic group form that is not round.

Theorem 3.8. *Let q be an anisotropic form over F of dimension 2^n for $n \in \mathbb{N}$.*

- (i) *There exists an extension K/F such that q_K is an anisotropic group form.*
- (ii) *If q is similar to a Pfister form over F , then q_K is round for every extension K/F such that q_K is a group form.*
- (iii) *If q is not similar to a Pfister form over F , there exists an extension K/F such that q_K is an anisotropic group form that is not round.*

Proof. (i) In light of statement (iii), it suffices to prove this statement in the case where q is similar to a Pfister form over F . By Lemma 2.6, we have that q is a Pfister form if and only if it represents one. Thus, we may let $K = F$ in the case where $1 \in D_F(q)$. Otherwise, we may let $K = F(q \perp \langle -1 \rangle)$, as q_K is anisotropic by Theorem 1.3.

(ii) Let K/F be such that q_K is a group form. As $1 \in D_K(q)$, we have that q_K is a Pfister form by Lemma 2.6, whereby q_K is round.

(iii) If $1 \notin D_F(q)$, we may consider q as a form over $L = F(q \perp \langle -1 \rangle)$, whereby $1 \in D_L(q)$. In this case, q_L remains anisotropic by Theorem 1.3. As q is not similar to a Pfister form over F , it follows that q is not hyperbolic over $F(q)$ by [Lam 2005, Theorem X.4.14]. Since

$$i(q_{L(q)}) = i(q_{F(q)(q \perp \langle -1 \rangle)}),$$

we may invoke the Cassels–Pfister subform theorem [Lam 2005, Theorem X.4.5] to establish that q is not hyperbolic over $L(q)$, whereby it follows that q is not similar to a Pfister form over L by [Lam 2005, Theorem X.4.14]. Hence, we may assume, without loss of generality, that $1 \in D_F(q)$.

Let $K = F$ if q is a group form over F that is not round. Otherwise, let

$$L_0 = F((x))(q \otimes \langle 1, -x \rangle).$$

Since q is not similar to a Pfister form over F , it follows that q is not a group form over L_0 by Theorem 2.7. Hence, we have that $H_{L_0}(q) \setminus D_{L_0}(q)$ is a nonempty

set by Corollary 2.3 (in particular, the proof of Theorem 2.7 establishes that $x \in H_{L_0}(q) \setminus D_{L_0}(q)$). Consider the set

$$Q(L_0) = \{q \perp \langle -a \mid a \in H_{L_0}(q) \setminus D_{L_0}(q)\},$$

which is a nonempty set of anisotropic forms over L_0 . For $i \geq 0$, we inductively define L_{i+1} to be the compositum of all function fields of forms in $Q(L_i)$. For all L_i and $a \in H_{L_i}(q) \setminus D_{L_i}(q)$, we have that q is anisotropic over $L_i(q \perp \langle -a \rangle)$ by Theorem 1.3. Hence, letting $K = \bigcup_{i=0}^\infty L_i$, it follows that q_K is anisotropic. Moreover, as $H_K(q) = D_K(q)$ by construction, it follows that q_K is a group form by Corollary 2.3.

It remains to show that q_K is not round. By construction, we have that $q \perp -xq$ is isotropic over L_0 , whereby $x \in H_K(q)$. Suppose, for the sake of contradiction, that $q \perp -xq$ is hyperbolic over K . Hence, for some $i \in \mathbb{N} \cup \{0\}$, there exists an extension L'_i/L_i and $a \in (L'_i)^\times$ such that $((q \perp -xq)_{L'_i})_{\text{an}}$ is hyperbolic over $L'_i(q \perp \langle -a \rangle)$. As a consequence of Elman and Lam's representation theorem [1972, Theorem 1.4], there exists a form p over L'_i such that $\dim p < \dim q$ and

$$((q \perp -xq)_{L'_i})_{\text{an}} \simeq p \perp -xp$$

(see [Hoffmann 1996, Lemma 3.1]). Hence, invoking [Karpenko and Merkurjev 2003, Corollary 4.2], it follows that

$$\dim(p \perp -xp) - i((p \perp -xp)_{L'_i(q \perp \langle -a \rangle)}) \geq \dim(q \perp \langle -a \rangle) - i_1(q \perp \langle -a \rangle).$$

As $i_1(q \perp \langle -a \rangle) = 1$ by Theorem 1.3 and [Lam 2005, Exercise I.16], it follows that

$$\dim(p \perp -xp) - \frac{\dim(p \perp -xp)}{2} \geq \dim q,$$

in contradiction to the fact that $\dim p < \dim q$. Hence, having obtained our desired contradiction, we may conclude that $x \notin G_K(q)$, whereby q_K is not round by Corollary 2.5. □

Remark 3.9. Scully [2016b, Main Theorem] recently established that, for p and q anisotropic forms over F of dimension at least two with $2^i < \dim q \leq 2^{i+1}$, if $p_{F(q)}$ is hyperbolic, then $\dim p = 2^{i+1}k$ for some $k \in \mathbb{N}$. One may invoke this result to shorten the final component of the above proof.

Per Example 3.7, in order to answer Question 1.1 in the case where $\dim q \neq 2^n$ for $n \in \mathbb{N}$, we require some additional assumptions regarding the form q over F . Orderings are a useful tool in this regard, with their behaviour with respect to function-field extensions being governed by the following result due to Elman, Lam and Wadsworth [Elman et al. 1979, Theorem 3.5] and, independently, Knebusch [Gentile and Shapiro 1978, Lemma 10].

Theorem 3.10. *Let q be a form of dimension at least two over a real field F . An ordering P of F extends to $F(q)$ if and only if q is indefinite at P .*

Invoking Theorem 3.10, we can resolve Question 1.1 in the case where q is a positive-definite form over a real field.

Proposition 3.11. *Let F be a real field. Let q be a form over F that is positive definite with respect to some ordering of F . If q is not similar to a Pfister form over F , there exists an extension K/F such that q_K is an anisotropic group form that is not round.*

Proof. Let P be an ordering of F such that q is positive definite with respect to P . If $1 \notin D_F(q)$, we may consider q as a form over $L = F(q \perp \langle -1 \rangle)$, whereby P is an ordering of L by Theorem 3.10 and $1 \in D_L(q)$. Per the proof of Theorem 3.8 (iii), q is not hyperbolic over $L(q)$, and thus is not similar to a Pfister form over L . Hence, we may assume, without loss of generality, that $1 \in D_F(q)$.

Let $K = F$ if q is a group form over F that is not round. Otherwise, let

$$L_0 = F((x))(q \otimes \langle 1, -x \rangle).$$

Since q is not similar to a Pfister form over F , it follows that q is not a group form over L_0 by Theorem 2.7. Hence, we have that $H_{L_0}(q) \setminus D_{L_0}(q)$ is a nonempty set by Corollary 2.3. Moreover, by Theorem 3.10, there exist orderings of L_0 such that q is positive definite.

Let L/L_0 be an extension such that q is positive definite with respect to an ordering P of L and $H_L(q) \setminus D_L(q)$ is not empty. Let $a \in H_L(q) \setminus D_L(q)$, whereby $a \in P$ and the form $q \perp \langle -a \rangle$ has signature $\dim q - 1$ with respect to P . As P extends to $L(q \perp \langle -a \rangle)$, by Theorem 3.10, it thus follows that $i_1(q \perp \langle -a \rangle) = 1$. Hence, q is anisotropic over $L(q \perp \langle -a \rangle)$ by Theorem 1.4 (i).

Equipped with the above, we may now proceed with the argument in the proof of Theorem 3.8 (iii) to establish the existence of an extension K/L_0 such that q_K is an anisotropic group form with $x \in D_K(q) \setminus G_K(q)$. \square

As discussed in [O’Shea 2016], many properties of a form q over F are shared by its generic Pfister multiple $q \otimes \langle 1, x \rangle$ over $F((x))$. Invoking Proposition 2.2, we may show that this is also the case with respect to the group property.

Proposition 3.12. *A form q is a group form over F if and only if $q \otimes \langle 1, x \rangle$ is a group form over $K = F((x))$.*

Proof. Without loss of generality, we may assume that q is anisotropic over F .

Let q be a group form over F . As every nonzero square class in K can be represented by a or ax for some $a \in F^\times$, we first suppose that

$$q \otimes \langle 1, x \rangle \perp -ax(q \otimes \langle 1, x \rangle)$$

is isotropic over K for $a \in F^\times$. Then, because $x \in D_K(\langle 1, x \rangle)$, it follows that $q \otimes \langle 1, x \rangle \perp -a(q \otimes \langle 1, x \rangle)$ is isotropic over K in this case. Thus, supposing that $q \otimes \langle 1, x \rangle \perp -a(q \otimes \langle 1, x \rangle)$ is isotropic over K for $a \in F^\times$, it suffices to show that

$q \otimes \langle 1, x \rangle \perp \langle -a \rangle$ and $q \otimes \langle 1, x \rangle \perp \langle -ax \rangle$ are isotropic over K in accordance with Proposition 2.2. Invoking Lemma 1.2, it follows that $q \perp -aq$ is isotropic over F , whereby $q \perp \langle -a \rangle$ is isotropic over F by Proposition 2.2. Hence, it follows that $q \otimes \langle 1, x \rangle \perp \langle -a \rangle$ and $q \otimes \langle 1, x \rangle \perp \langle -ax \rangle$ are isotropic over K , as desired.

Conversely, let $q \otimes \langle 1, x \rangle$ be a group form over K . Letting $a \in F^\times$ be such that $q \perp -aq$ is isotropic over F , it follows that $q \otimes \langle 1, x \rangle \perp -a(q \otimes \langle 1, x \rangle)$ is isotropic over K , whereby $q \otimes \langle 1, x \rangle \perp -a$ is isotropic over K by Proposition 2.2. Hence, $q \perp \langle -a \rangle$ is isotropic over F by Lemma 1.2, whereby q is a group form over F by Proposition 2.2. \square

Combining Proposition 3.12 with Proposition 3.2 (ii) and [O'Shea 2016, Proposition 3.11], we obtain the following corollary.

Corollary 3.13. *Let q be an anisotropic group form over F that is not round. Then $q \otimes \langle 1, x \rangle$ is an anisotropic group form over $K = F((x))$ that is not round.*

Acknowledgements

I thank Sylvain Roussey for making his comprehensive PhD thesis available to me. I am grateful to Vincent Astier and Thomas Unger for helpful comments.

References

- [Alpers 1991] B. Alpers, "Round quadratic forms under algebraic extensions", *Pacific J. Math.* **147**:2 (1991), 213–229. MR Zbl
- [Elman 1973] R. Elman, "Rund forms over real algebraic function fields in one variable", *Proc. Amer. Math. Soc.* **41** (1973), 431–436. MR Zbl
- [Elman and Lam 1972] R. Elman and T. Y. Lam, "Pfister forms and K -theory of fields", *J. Algebra* **23** (1972), 181–213. MR Zbl
- [Elman and Lam 1973] R. Elman and T. Y. Lam, "Quadratic forms and the u -invariant, I", *Math. Z.* **131** (1973), 283–304. MR Zbl
- [Elman et al. 1979] R. Elman, T. Y. Lam, and A. R. Wadsworth, "Orderings under field extensions", *J. Reine Angew. Math.* **306** (1979), 7–27. MR Zbl
- [Elman et al. 2008] R. Elman, N. Karpenko, and A. Merkurjev, *The algebraic and geometric theory of quadratic forms*, American Mathematical Society Colloquium Publications **56**, American Mathematical Society, Providence, RI, 2008. MR Zbl
- [Gentile and Shapiro 1978] E. R. Gentile and D. B. Shapiro, "Conservative quadratic forms", *Math. Z.* **163**:1 (1978), 15–23. MR Zbl
- [Hoffmann 1995] D. W. Hoffmann, "Isotropy of quadratic forms over the function field of a quadric", *Math. Z.* **220**:3 (1995), 461–476. MR Zbl
- [Hoffmann 1996] D. W. Hoffmann, "Twisted Pfister forms", *Doc. Math.* **1** (1996), 67–102. MR Zbl
- [Hsia and Johnson 1973a] J. S. Hsia and R. P. Johnson, "Round and group quadratic forms over global fields", *J. Number Theory* **5** (1973), 356–366. MR Zbl
- [Hsia and Johnson 1973b] J. S. Hsia and R. P. Johnson, "Round and Pfister forms over $\mathbb{R}(t)$ ", *Pacific J. Math.* **49**:1 (1973), 101–108. MR Zbl

- [Izhboldin 2000] O. T. Izhboldin, “On the isotropy of low-dimensional forms over function fields of a quadric”, *Algebra i Analiz* **12**:5 (2000), 106–127. In Russian; translated in *St. Petersburg Math. J.* **12**:5 (2001), 791–806. MR Zbl
- [Karpenko and Merkurjev 2003] N. Karpenko and A. Merkurjev, “Essential dimension of quadrics”, *Invent. Math.* **153**:2 (2003), 361–372. MR Zbl
- [Knebusch 1976] M. Knebusch, “Generic splitting of quadratic forms, I”, *Proc. London Math. Soc.* (3) **33**:1 (1976), 65–93. MR Zbl
- [Lam 2005] T. Y. Lam, *Introduction to quadratic forms over fields*, Graduate Studies in Mathematics **67**, American Mathematical Society, Providence, RI, 2005. MR Zbl
- [O’Shea 2016] J. O’Shea, “Multiples of Pfister forms”, *J. Algebra* **449** (2016), 214–236. MR Zbl
- [Pfister 1965] A. Pfister, “Multiplikative quadratische Formen”, *Arch. Math. (Basel)* **16** (1965), 363–370. MR Zbl
- [Roussey 2005] S. Roussey, *Isotropie, corps de fonctions et équivalences birationnelles des formes quadratiques*, Ph.D. thesis, Université de Franche-Comté, 2005.
- [Scharlau 1985] W. Scharlau, *Quadratic and Hermitian forms*, Grundlehren der Math. Wissenschaften **270**, Springer, Berlin, 1985. MR Zbl
- [Scully 2016a] S. Scully, “A bound for the index of a quadratic form after scalar extension to the function field of a quadric”, preprint, 2016. arXiv
- [Scully 2016b] S. Scully, “Hyperbolicity and near hyperbolicity of quadratic forms over function fields of quadrics”, preprint, 2016. arXiv
- [Springer 1952] T. A. Springer, “Sur les formes quadratiques d’indice zéro”, *C. R. Acad. Sci. Paris* **234** (1952), 1517–1519. MR Zbl
- [Vishik 2011] A. Vishik, “Excellent connections in the motives of quadrics”, *Ann. Sci. Éc. Norm. Supér.* (4) **44**:1 (2011), 183–195. MR Zbl
- [Wadsworth 1975] A. R. Wadsworth, “Similarity of quadratic forms and isomorphism of their function fields”, *Trans. Amer. Math. Soc.* **208** (1975), 352–358. MR Zbl
- [Wadsworth and Shapiro 1977] A. R. Wadsworth and D. B. Shapiro, “On multiples of round and Pfister forms”, *Math. Z.* **157**:1 (1977), 53–62. MR Zbl

Received June 8, 2017. Revised July 23, 2017.

JAMES O’ SHEA
DEPARTMENT OF MATHEMATICS AND STATISTICS
NATIONAL UNIVERSITY OF IRELAND, MAYNOOTH
IRELAND
james.oshea@mu.ie

Previous affiliation:

SCHOOL OF MATHEMATICS, STATISTICS AND APPLIED MATHEMATICS
NATIONAL UNIVERSITY OF IRELAND, GALWAY
IRELAND

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 293 No. 2 April 2018

Sums of CR functions from competing CR structures	257
DAVID E. BARRETT and DUSTY E. GRUNDMEIER	
On Tate duality and a projective scalar property for symmetric algebras	277
FLORIAN EISELE, MICHAEL GELINE, RADHA KESSAR and MARKUS LINCKELMANN	
Coaction functors, II	301
S. KALISZEWSKI, MAGNUS B. LANDSTAD and JOHN QUIGG	
Construction of a Rapoport–Zink space for $GU(1, 1)$ in the ramified 2-adic case	341
DANIEL KIRCH	
Group and round quadratic forms	391
JAMES O’ SHEA	
Dual operator algebras close to injective von Neumann algebras	407
JEAN ROYDOR	
Scalar curvature and singular metrics	427
YUGUANG SHI and LUEN-FAI TAM	
On the differentiability issue of the drift-diffusion equation with nonlocal Lévy-type diffusion	471
LIUTANG XUE and ZHUAN YE	



0030-8730(201804)293:2;1-U