

*Pacific
Journal of
Mathematics*

**THREE-DIMENSIONAL SOL MANIFOLDS
AND COMPLEX KLEINIAN GROUPS**

WALDEMAR BARRERA, RENE GARCIA-LARA AND JUAN NAVARRETE

Volume 294 No. 1

May 2018

THREE-DIMENSIONAL SOL MANIFOLDS AND COMPLEX KLEINIAN GROUPS

WALDEMAR BARRERA, RENE GARCIA-LARA AND JUAN NAVARRETE

We give a topological description of the quotient space $\Omega(G)/G$, in the case when $G \subset \mathrm{PSL}(3, \mathbb{C})$ is a discrete subgroup acting on $\mathbb{P}_{\mathbb{C}}^2$ and the maximum number of complex projective lines in general position contained in the Kulkarni limit set $\Lambda(G) = \mathbb{P}_{\mathbb{C}}^2 \setminus \Omega(G)$ is equal to 4. Moreover, we give a topological description of the quotient space $\Omega(G)/G$ in the case when G is a lattice of the Heisenberg group.

1. Introduction

Complex Kleinian groups were introduced by José Seade and Alberto Verjovsky [2001]. A complex Kleinian group G is a subgroup of $\mathrm{PSL}(n+1, \mathbb{C})$ acting properly and discontinuously on a nonempty G -invariant open subset of $\mathbb{P}_{\mathbb{C}}^n$. We remark that complex Kleinian groups are discrete subgroups of $\mathrm{PSL}(n+1, \mathbb{C})$ but the converse is not necessarily true; for example, the group $\mathrm{PSL}(3, \mathbb{Z})$ is a discrete subgroup of $\mathrm{PSL}(3, \mathbb{C})$ which is not a complex Kleinian group. See [Barrera et al. 2014].

There is no standard definition of limit set for a complex Kleinian group, we use the following three notions: Kulkarni limit set, Myrberg limit set, or the complement of a maximal region of discontinuity which are discussed in detail in [Barrera et al. 2016]. However by some additional hypotheses on the action of G on the projective plane, all these concepts of limit set are equivalent; see [Barrera et al. 2011a]. In the classical theory of Kleinian groups, there is a theorem which states that discrete infinite subgroups contain one, two, or infinity points in its limit set. On the other hand, Angel Cano and José Seade show that every infinite discrete subgroup of $\mathrm{PSL}(3, \mathbb{C})$ has a complex projective line contained in its limit set (see [Cano and Seade 2014]), in consequence, the limit set of infinite subgroups of $\mathrm{PSL}(3, \mathbb{C})$ is an uncountable subset of $\mathbb{P}_{\mathbb{C}}^2$.

Thus, it is natural to say that $G \subset \mathrm{PSL}(3, \mathbb{C})$ is an *elementary complex Kleinian group* whenever its limit set contains a finite number of complex projective lines; see [Cano et al. 2013]. There is another kind of group whose limit set contains

MSC2010: primary 32F45, 37F30, 32Q45; secondary 22E40, 37F45.

Keywords: Kleinian groups, projective complex plane, discrete groups, limit set.

infinitely many complex projective lines but only finitely many in general position. We call these groups *elementary complex Kleinian groups of type II*.

In [Barrera et al. 2011b], the authors give an algebraic characterization of those complex Kleinian groups such that the maximum number of complex projective lines in general position contained in its Kulkarni limit set is equal to 4. In this article we describe the topology of the quotient space of these groups. In fact, we prove the following theorem:

Theorem 1.1. *Let $G \subset \mathrm{PSL}(3, \mathbb{C})$ be a torsion-free complex Kleinian group, such that the maximum number of complex projective lines in general position contained in its Kulkarni limit set is equal to 4, then:*

- (i) *The group G is isomorphic to a lattice of the group Sol (see Section 2F).*
- (ii) *If Ω_0 is a G -invariant connected component of the Kulkarni discontinuity region of G , then Ω_0/G is diffeomorphic to $(\mathrm{Sol}/G) \times \mathbb{R}$.*

Corollary 1.2. *There is a countable number of nonisomorphic complex Kleinian groups such that the maximum number of complex projective lines in general position contained in its Kulkarni limit set is equal to 4.*

Corollary 1.3. *Under the hypotheses of Theorem 1.1, Ω_0/G is a fiber bundle with base \mathbb{S}^1 and fiber $\mathbb{T}^2 \times \mathbb{R}$.*

Theorem 1.4. *If G is a lattice on the three-dimensional real Heisenberg group \mathcal{H} , then there exists a G -invariant open set $\Omega \subset \mathbb{P}_{\mathbb{C}}^2$ such that Ω/G is diffeomorphic to $(\mathcal{H}/G) \times \mathbb{R}$.*

This article is organized in the following way: In Section 2 we include some basic preliminaries about complex Kleinian groups, and a brief survey on complex Kleinian groups, such that the maximum number of complex lines contained in their Kulkarni limit set is equal to 4 [Barrera et al. 2011b]. In Section 3 we give an explicit smooth foliation of the bidisc $\mathbb{H} \times \mathbb{H}$ where the leaves are diffeomorphic copies of Sol . In Section 4, we study the geometry of the leaves and we show that the bidisc $\mathbb{H} \times \mathbb{H}$ is diffeomorphic to $\mathrm{Sol} \times \mathbb{R}$. Moreover this diffeomorphism is G -equivariant, where G denotes an hyperbolic toral group. In Section 5 we do some explicit computations to determine the Riemannian metrics of the leaves. For each leaf we obtain an isometric embedding of the group Sol to $\mathbb{H} \times \mathbb{H}$. Finally, we give a proof of Theorem 1.1.

Corollaries 1.2 and 1.3 are a consequence of Theorem 1.1 and [de la Harpe 2000, Proposition 30]. In Section 6 we give a proof of Theorem 1.4, the procedure is similar to the proof of Theorem 1.1, except that we have not a G -equivariant diffeomorphism between $\mathbb{C} \times \mathbb{H}$ and $\mathcal{H} \times \mathbb{R}$. However the proof can be done because the natural action of G on $\mathbb{C} \times \mathbb{H}$ translated to $\mathcal{H} \times \mathbb{R}$ is the classical action on the first factor of G on \mathcal{H} , and it is the trivial action on the second factor.

2. Preliminaries

The purpose of this section is to provide some definitions and results about complex Kleinian groups that will be helpful to the reader. For more details see [Cano et al. 2013; Barrera et al. 2011b; 2011a].

2A. Projective geometry. We recall that the complex projective plane $\mathbb{P}_{\mathbb{C}}^2$ is defined as the orbit space of the usual scalar multiplication action of the Lie group \mathbb{C}^* in $\mathbb{C}^3 \setminus \{\mathbf{0}\}$ and it is denoted by

$$\mathbb{P}_{\mathbb{C}}^2 := (\mathbb{C}^3 \setminus \{\mathbf{0}\})/\mathbb{C}^*.$$

This is a compact connected complex 2-dimensional manifold. Let $[\]: \mathbb{C}^3 \setminus \{\mathbf{0}\} \rightarrow \mathbb{P}_{\mathbb{C}}^2$ be the quotient map. If $\beta = \{e_1, e_2, e_3\}$ is the standard basis of \mathbb{C}^3 , we write $[e_j] = e_j$, for $j = 1, 2, 3$, and if $\mathbf{z} = (z_1, z_2, z_3) \in \mathbb{C}^3 \setminus \{\mathbf{0}\}$ then we write $[\mathbf{z}] = [z_1 : z_2 : z_3]$. Also, $\ell \subset \mathbb{P}_{\mathbb{C}}^2$ is said to be a complex line if $[\ell]^{-1} \cup \{\mathbf{0}\}$ is a complex linear subspace of dimension 2. Given two distinct points $[\mathbf{z}], [\mathbf{w}] \in \mathbb{P}_{\mathbb{C}}^2$, there is a unique complex projective line passing through $[\mathbf{z}]$ and $[\mathbf{w}]$. This kind of complex projective line is called a *line*, for short, and it is denoted by $\overleftrightarrow{[\mathbf{z}], [\mathbf{w}]}$. Consider the action of \mathbb{C}^* on $\text{GL}(3, \mathbb{C})$ given by the usual scalar multiplication, then

$$\text{PGL}(3, \mathbb{C}) = \text{GL}(3, \mathbb{C})/\mathbb{C}^*$$

is a Lie group whose elements are called projective transformations. Letting $[[\]]: \text{GL}(3, \mathbb{C}) \rightarrow \text{PGL}(3, \mathbb{C})$ be the quotient map, $g \in \text{PGL}(3, \mathbb{C})$ and $\mathbf{g} \in \text{GL}(3, \mathbb{C})$, we say that \mathbf{g} is a lift of g if $[[\mathbf{g}]] = g$. One can show that $\text{PGL}(3, \mathbb{C})$ is a Lie group which acts transitively, effectively and by biholomorphisms on $\mathbb{P}_{\mathbb{C}}^2$ by $[[\mathbf{g}]](\mathbf{w}) = [\mathbf{g}(\mathbf{w})]$, where $\mathbf{w} \in \mathbb{C}^3 \setminus \{\mathbf{0}\}$ and $\mathbf{g} \in \text{GL}(3, \mathbb{C})$.

We could have considered the action of the cube roots of unity $\{1, \omega, \omega^2\} \subset \mathbb{C}^*$ on $\text{SL}(3, \mathbb{C})$ given by the usual scalar multiplication, then

$$\text{PSL}(3, \mathbb{C}) = \text{SL}(3, \mathbb{C})/\{1, \omega, \omega^2\} \cong \text{PGL}(3, \mathbb{C}).$$

We denote by $M_{3 \times 3}(\mathbb{C})$ the space of all 3×3 matrices with entries in \mathbb{C} equipped with the standard topology. The quotient space

$$\text{End}(3, \mathbb{C}) := (M_{3 \times 3}(\mathbb{C}) \setminus \{\mathbf{0}\})/\mathbb{C}^*$$

is called the space of *pseudoprojective maps of $\mathbb{P}_{\mathbb{C}}^2$* and it is naturally identified with the projective space $\mathbb{P}_{\mathbb{C}}^8$. Since $\text{GL}(3, \mathbb{C})$ is an open, dense, \mathbb{C}^* -invariant set of $M_{3 \times 3}(\mathbb{C}) \setminus \{\mathbf{0}\}$, we get that the space of pseudoprojective maps of $\mathbb{P}_{\mathbb{C}}^2$ is a compactification of $\text{PGL}(3, \mathbb{C})$ (or $\text{PSL}(3, \mathbb{C})$). As in the case of projective maps, if s is an element in $M_{3 \times 3}(\mathbb{C}) \setminus \{\mathbf{0}\}$, then $[s]$ denotes the equivalence class of the matrix s in the space of pseudoprojective maps of $\mathbb{P}_{\mathbb{C}}^2$. Also, we say that $\mathbf{s} \in M_{3 \times 3}(\mathbb{C}) \setminus \{\mathbf{0}\}$ is a lift of the pseudoprojective map S whenever $[s] = S$.

Let S be an element in $(M_{3 \times 3}(\mathbb{C}) \setminus \{\mathbf{0}\})/\mathbb{C}^*$ and s a lift to $M_{3 \times 3}(\mathbb{C}) \setminus \{\mathbf{0}\}$ of S . The matrix s induces a nonzero linear transformation $s : \mathbb{C}^3 \rightarrow \mathbb{C}^3$, which is not necessarily invertible. Let $\text{Ker}(s) \subsetneq \mathbb{C}^3$ be its kernel and let $\text{Ker}(S)$ denote its projectivization to $\mathbb{P}_{\mathbb{C}}^2$, taking into account that $\text{Ker}(S) := \emptyset$ whenever $\text{Ker}(s) = \{(0, 0, 0)\}$.

2B. *Discontinuous actions on $\mathbb{P}_{\mathbb{C}}^2$.*

Definition 2.1. Let $G \subset \text{PSL}(3, \mathbb{C})$ be a group. We say that G is a *complex Kleinian group* if it acts properly and discontinuously on an open nonempty G -invariant set $U \subset \mathbb{P}_{\mathbb{C}}^2$, meaning that for each pair of compact subsets $C, D \subset U$, the set

$$\{g \in G : g(C) \cap D \neq \emptyset\}$$

is *finite*.

One of the main difficulties in deciding whether a group G is Kleinian complex is to find an open set verifying the definition above. In order to give an answer to this problem we study two mathematical concepts: the equicontinuity set of G and the Kulkarni discontinuity region of G . Now, we discuss each of these concepts.

2C. *The equicontinuity set.* The concept of equicontinuity has long been studied in mathematics. For convenience to the reader, we include the definition and notation that we use in this work.

Definition 2.2. The *equicontinuity set* for a family \mathcal{F} of endomorphisms of $\mathbb{P}_{\mathbb{C}}^2$, denoted $\text{Eq}(\mathcal{F})$, is defined as the set of points $z \in \mathbb{P}_{\mathbb{C}}^2$ for which there is an open neighborhood U of z such that $\{f|_U : f \in \mathcal{F}\}$ is a normal family.

This modern approach and ideas on this concept were studied by Angel Cano in his Ph.D thesis. However, thanks to a reference by Ravi Kulkarni to works of Myrberg, we found that some of these results had already been discovered, in an arcane mathematical language. However, it is fair to acknowledge Angel Cano for rediscovering these results and applying them successfully to the theory of complex Kleinian groups.

Definition 2.3. Let $G \subset \text{PSL}(3, \mathbb{C})$ be a discrete group. If

$$G' = \{S \text{ is a pseudoprojective map of } \mathbb{P}_{\mathbb{C}}^2 : S \text{ is a cluster point of } G\},$$

then the *Myrberg limit set* (see [Myrberg 1925]) is defined as the set

$$\Lambda_{\text{Myr}}(G) = \bigcup_{S \in G'} \text{Ker}(S).$$

Myrberg [1925] shows that G acts properly and discontinuously on $\mathbb{P}_{\mathbb{C}}^2 \setminus \Lambda_{\text{Myr}}(G)$.

Theorem 2.4 [Barrera et al. 2011a]. *If $G \subset \text{PSL}(3, \mathbb{C})$ is a discrete group, then:*

- (i) *The group G acts properly and discontinuously on $\text{Eq}(G)$.*

(ii) *The equicontinuity set of G satisfies:*

$$\text{Eq}(G) = \mathbb{P}_{\mathbb{C}}^2 \setminus \Lambda_{\text{Myr}}(G)$$

(iii) *If U is an open G -invariant subset such that $\mathbb{P}_{\mathbb{C}}^2 \setminus U$ contains at least three complex lines in general position, then $U \subset \text{Eq}(G)$.*

2D. Kulkarni discontinuity region. Kulkarni [1978] defined a limit set for groups of homeomorphisms acting on a locally compact Hausdorff space. For the reader's convenience, we explain this construction in the context of projective spaces.

Definition 2.5. Let $G \subset \text{PSL}(3, \mathbb{C})$ be a subgroup.

- The set $L_0(G)$ is defined as the closure of the set of points in $\mathbb{P}_{\mathbb{C}}^2$ with infinite isotropy group.
- The set $L_1(G)$ is defined as the closure of the set of cluster points of the orbit Gz , where z runs over $\mathbb{P}_{\mathbb{C}}^2 \setminus L_0(G)$.
- The set $L_2(G)$ is defined as the closure of the set of cluster points of the family of compact sets $\{g(K) : g \in G\}$, where K runs over all the compact subsets of

$$\mathbb{P}_{\mathbb{C}}^2 \setminus (L_0(G) \cup L_1(G)).$$

The *Kulkarni limit set* of G is defined as

$$\Lambda_{\text{Kul}}(G) = L_0(G) \cup L_1(G) \cup L_2(G).$$

The *Kulkarni discontinuity region* of G is defined as:

$$\Omega(G) = \mathbb{P}_{\mathbb{C}}^2 \setminus \Lambda_{\text{Kul}}(G).$$

Kulkarni [1978] proves that G acts properly and discontinuously on the set $\Omega(G)$. However, $\Omega(G)$ is not necessarily the maximal open subset of $\mathbb{P}_{\mathbb{C}}^2$ where G acts properly and discontinuously.

We notice that the Kulkarni limit set is a generalization of the classical limit set for a discrete subgroup of hyperbolic isometries acting on the sphere at infinity of hyperbolic space. In general, it is very hard to give an explicit computation of the Kulkarni limit set. In [Navarrete 2006; 2008], we can find these computations for the cyclic subgroups of $\text{PSL}(3, \mathbb{C})$ and for discrete subgroups of $\text{PU}(2, 1)$ acting on the complex projective plane $\mathbb{P}_{\mathbb{C}}^2$.

We could define the limit set as the complement of a maximal open set where the group acts properly and discontinuously, but in general there is no canonical way to build this G -invariant open set. On the other hand, when we ensure the existence of this maximal open set, this notion of limit set has good properties. See [Barrera et al. 2014].

2E. Four lines complex Kleinian groups. This section is devoted to complex Kleinian groups of $\mathrm{PSL}(3, \mathbb{C})$ such that the maximum number of complex projective lines in general position contained in its Kulkarni limit set is equal to 4. For simplicity we call groups of this kind *four lines complex Kleinian groups*. In [Barrera et al. 2011b], the authors give an algebraic characterization of four lines complex Kleinian groups. For the reader's convenience, we reproduce briefly the main ideas and the notation used there.

Letting $A \in \mathrm{SL}(2, \mathbb{Z})$, with $|\mathrm{tr}(A)| > 2$, we define the following discrete subgroup of $\mathrm{PSL}(3, \mathbb{C})$, called *hyperbolic toral group*.

$$G_A = \left\{ \begin{pmatrix} A^k & \mathbf{b} \\ \mathbf{0} & 1 \end{pmatrix} : \mathbf{b} \in M(2 \times 1, \mathbb{Z}), k \in \mathbb{Z} \right\},$$

The group G_A is a four lines complex Kleinian group and moreover if G is a four lines complex Kleinian group, then there exists a hyperbolic toral group G_A such that $[G : G_A] \leq 8$.

It is possible to conjugate the group G_A to a group, still denoted by G_A , where each element is of the form

$$\begin{pmatrix} \lambda^k & 0 & ny_0 + mx_0 \\ 0 & \lambda^{-k} & nx_0 + mz_0 \\ 0 & 0 & 1 \end{pmatrix}$$

where k, m and n run over \mathbb{Z} and λ is one of the eigenvalues of A . At this point it is not hard to see that the Kulkarni discontinuity region consists of four disjoint copies of $\mathbb{H}^\pm \times \mathbb{H}^\pm$, where \mathbb{H}^+ is the upper half plane and \mathbb{H}^- is the lower half plane.

2F. The Sol geometries. Sol is one of the eight geometries defined by William Thurston in his famous program of geometrization of compact three manifolds. The group Sol is defined as the space $\mathbb{R}^3 = \mathbb{R}^2 \times \mathbb{R}$ equipped with the group operation

$$\left(\begin{pmatrix} x_1 \\ y_1 \end{pmatrix}, t_1 \right) \cdot \left(\begin{pmatrix} x_2 \\ y_2 \end{pmatrix}, t_2 \right) = \left(\begin{pmatrix} x_1 + e^{t_1} x_2 \\ y_1 + e^{-t_1} y_2 \end{pmatrix}, t_1 + t_2 \right).$$

In fact, it is a Lie group and it is equipped with the left-invariant Riemannian metric: $ds^2 = e^{2t} dx^2 + e^{-2t} dy^2 + dt^2$. An interesting fact about the Sol geometries is given by the following theorem of [de la Harpe 2000], which we state for convenience:

Proposition 2.6. *Let A, B in $\mathrm{GL}(2, \mathbb{Z})$ be two matrices with traces of absolute value strictly larger than 2. The semidirect products $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$, and $\mathbb{Z}^2 \rtimes_B \mathbb{Z}$ considered as the matrix groups*

$$\left\{ \begin{pmatrix} A^k & \mathbf{b} \\ \mathbf{0} & 1 \end{pmatrix} : \mathbf{b} \in M(2 \times 1, \mathbb{Z}), k \in \mathbb{Z} \right\},$$

$$\left\{ \begin{pmatrix} B^k & \mathbf{b} \\ \mathbf{0} & 1 \end{pmatrix} : \mathbf{b} \in M(2 \times 1, \mathbb{Z}), k \in \mathbb{Z} \right\}$$

are isomorphic if and only if A is conjugate in $\mathrm{GL}(2, \mathbb{Z})$ to B or B^{-1} , and they are quasi-isometric in all cases.

The quotient spaces $\mathrm{Sol}/(\mathbb{Z}^2 \rtimes_A \mathbb{Z})$ are examples of compact three manifolds where the topological type is determined by the fundamental group. For more details about this subject see [Scott 1983; Thurston 1997; de la Harpe 2000].

3. Foliation of $\mathbb{H} \times \mathbb{H}$ by Sol

In [Barrera et al. 2011b], the authors introduce the concept of hyperbolic toral groups. These groups are matrix groups where the elements are given by

$$\begin{pmatrix} \lambda^k & 0 & ny_0 + mx_0 \\ 0 & \lambda^{-k} & nx_0 + mz_0 \\ 0 & 0 & 1 \end{pmatrix},$$

where λ is a fixed real number, $|\lambda| \neq 1$ and k, n, m run over \mathbb{Z} . It is not hard to check this group is isomorphic to $\mathbb{Z}^2 \rtimes_A \mathbb{Z}$. Moreover, a continuous version of this group is given in the following way. Let $A \in \mathrm{SL}(2, \mathbb{Z})$ be a hyperbolic automorphism with Jordan form

$$A = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}$$

and consider the set of matrices of the form

$$\begin{pmatrix} \lambda^t & 0 & x \\ 0 & \lambda^{-t} & y \\ 0 & 0 & 1 \end{pmatrix},$$

where t, x, y run over \mathbb{R} . It is not hard to check that this group is isomorphic to $\mathrm{Sol} = \mathbb{R}^2 \rtimes \mathbb{R}$. For convenience for our computations we use this representation of the Lie group Sol . In the sequel, we will use the product metric in $\mathbb{H} \times \mathbb{H}$, where we endow each copy of \mathbb{H} with a metric homothetic to the hyperbolic metric by a factor of $\frac{1}{2}$:

$$\frac{dx_1^2 + dy_1^2}{2y_1^2} + \frac{dx_2^2 + dy_2^2}{2y_2^2}.$$

Proposition 3.1. *Let $z_1, z_2 \in \mathbb{H}$. We define a natural action of Sol in $\mathbb{H} \times \mathbb{H}$ by*

$$\begin{pmatrix} \lambda^t & 0 & x \\ 0 & \lambda^{-t} & y \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \\ 1 \end{pmatrix} = \begin{pmatrix} \lambda^t z_1 + x \\ \lambda^{-t} z_2 + y \\ 1 \end{pmatrix}.$$

The natural action of the group Sol on $\mathbb{H} \times \mathbb{H}$ satisfies the following:

- (i) The action is free.
- (ii) For each $z = (z_1, z_2) \in \mathbb{H} \times \mathbb{H}$ the function $f_z : \text{Sol} \rightarrow \mathbb{H} \times \mathbb{H}$ defined by $f_z(g) = gz$ is a smooth embedding.
- (iii) If $\{e_1, e_2, e_3, e_4\}$ is the canonical basis of \mathbb{R}^4 , then

$$X = y_1 e_2 + y_2 e_4$$

is one of the two normal unitary vector fields to the embedding $f_z(\text{Sol})$ in $\mathbb{H} \times \mathbb{H}$, which therefore is a smooth vector field globally defined.

Proof. (i) Let $z = (z_1, z_2) \in \mathbb{H} \times \mathbb{H}$, and assume that $\gamma \in \text{Sol}$ is such that $\gamma \cdot z = z$. Let $z_k = x_k + iy_k$, where $k = 1, 2$. Taking imaginary parts in the action, we get

$$\lambda^t y_1 = y_1, \quad \lambda^{-t} y_2 = y_2,$$

so $\lambda^t = \lambda^{-t} = 1$, because imaginary parts can not be null. Taking real parts, we obtain $x_1 + x = x_1$ and $x_2 + y = x_2$, then $x = y = 0$.

(ii) From the definition of the action, it is clear that f_z is smooth in t, x, y , which parametrizes Sol . By straightforward computations, we have that df_z has the Jacobian matrix given by

$$[df_z] = \begin{pmatrix} \ln(\lambda) \lambda^t x_1 & 1 & 0 \\ \ln(\lambda) \lambda^t y_1 & 0 & 0 \\ -\ln(\lambda) \lambda^{-t} x_2 & 0 & 1 \\ -\ln(\lambda) \lambda^{-t} y_2 & 0 & 0 \end{pmatrix}.$$

Since $y_1 > 0$, the Jacobian matrix has rank 3. Therefore, f_z is an immersion.

If $z_k = x_k + iy_k$, and $z' = (x'_1, y'_1, x'_2, y'_2) \in \mathbb{H} \times \mathbb{H}$, define t such that $\lambda^t = y'_1/y_1$, and (x', y') such that

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} x'_1 - \frac{y'_1 x_1}{y_1} \\ x'_2 - \frac{y'_2 y_2}{y_1} \end{pmatrix}.$$

These values for (t, x', y') define a mapping F , from $\mathbb{H} \times \mathbb{H}$ to $\mathbb{R}^3 \cong \text{Sol}$, such that $F \circ f_z = \text{Id}$. Note that F is a left continuous inverse for f_z , and hence, f_z is an homeomorphism.

(iii) The given formula for the product metric implies that X is unitary. By the form of the Jacobian matrix, the tangent space to the leaf passing through $z = (z_1, z_2) \in \mathbb{H} \times \mathbb{H}$ is spanned by the vectors, $e_1, e_3, \lambda^t x_1 e_1 + \lambda^t y_1 e_2 - \lambda^{-t} x_2 e_3 - \lambda^{-t} y_2 e_4$.

A straightforward computation shows that $X = \lambda^t y_1 e_2 + \lambda^{-t} y_2 e_4$ is orthogonal to the spanning tangent vectors. Finally, the result is obtained by taking $t = 0$. \square

By this theorem, if we vary z , we obtain a foliation $f_z(\text{Sol})$ of $\mathbb{H} \times \mathbb{H}$ by copies of Sol . We proceed to show that this foliation is globally rectifiable, in the sense that it induces a diffeomorphism to $\mathbb{R}^3 \times \mathbb{R}$, such that the hyperplanes $\mathbb{R}^3 \times \{t\}$ correspond to the leaves, and are diffeomorphic to Sol . For details on the theory of foliations, the reader can consult [Candel and Conlon 2000].

4. Geometry of the leaves

In the previous section, we described how Sol induces a foliation in the space $\mathbb{H} \times \mathbb{H}$ and gave an explicit formula for a smooth vector field X , normal to any leaf of the foliation in a product metric, which is homothetic to the canonical metric in each hyperbolic factor. In this section, we study the dynamics of the integral curves for this normal field. Let

$$\psi(t) = (z_1(t), z_2(t))$$

be an integral curve of the field, where $z_k = x_k + iy_k$ is as before. From the definition of X , it follows that the integral curves satisfy the set of equations

$$\dot{x}_k = 0, \quad \dot{y}_k = y_k.$$

These equations can be readily solved to get constant solutions in the real part of each copy of the hyperbolic, and exponentials in the imaginary parts. The flow of the normal field defines a one-parameter family of diffeomorphisms in $\mathbb{H} \times \mathbb{H}$, denoted by $\psi_t(z_1, z_2)$, where

$$\psi_t(z_1, z_2) = (x_1, e^t y_1, x_2, e^t y_2), \quad t \in \mathbb{R},$$

and it satisfies the following:

Proposition 4.1. (i) *The flow ψ_t rules $\mathbb{H} \times \mathbb{H}$ by geodesics.*

(ii) *The action of f_z is **equivariant** with the action induced by the flow, that is,*

$$\psi_s \circ f_z = f_{\psi_s(z)}.$$

Proof. (i) Both curves

$$(x_1, e^t y_1) \quad \text{and} \quad (x_2, e^t y_2)$$

correspond to a parametrization of a vertical geodesic in \mathbb{H} with respect to the hyperbolic metric. Since the metric we consider is homothetic to the standard hyperbolic metric, with a constant factor, these parametrizations correspond to geodesics with respect to this metric as well. Since the metric in $\mathbb{H} \times \mathbb{H}$ is a product, the result follows (see [Gallot et al. 2004]).

(ii) $\psi_s \circ f_z(t, x, y) = (\lambda^t x_1 + x, \lambda^t e^s y_1, \lambda^{-t} x_2 + y, \lambda^{-t} e^s y_2),$

which is the same expression obtained calculating $f_{\psi_s(z)}$. □

Observe that if we reparametrize the presentation of Sol we have used so far by the change of coordinates

$$(t', x', y') \mapsto \left(\frac{t'}{\ln(\lambda)}, x', y' \right),$$

we recover the original description of the group as given in [Scott 1983], and that under this reparametrization, we can analyze the geometry of the foliation in simpler terms, i.e., we can and will assume that

$$\psi_s \circ f_z(t, x, y) = (e^t x_1 + x, e^{t+s} y_1, e^{-t} x_2 + y, e^{-t+s} y_2)$$

in order to analyze the metric properties of the foliation.

Proposition 4.2. *Let $z = (iy_1, iy_2)$. Consider the leaf $f_z : \text{Sol} \rightarrow \mathbb{H} \times \mathbb{H}$, then the pullback metric is*

$$dt^2 + \frac{e^{-2t}}{2y_1^2} dx^2 + \frac{e^{2t}}{2y_2^2} dy^2.$$

In particular, if $y_1 = y_2 = 1/\sqrt{2}$, f_z is an isometric embedding of Sol into $\mathbb{H} \times \mathbb{H}$.

Proof. We have

$$f_z(t, x, y) = (x, e^t y_1, y, e^{-t} y_2).$$

Therefore, the Jacobian matrix is

$$\begin{pmatrix} 0 & 1 & 0 \\ e^t y_1 & 0 & 0 \\ 0 & 0 & 1 \\ -e^{-t} y_2 & 0 & 0 \end{pmatrix}.$$

Applying the product metric to the basis vectors $e^t y_1 e_2 - e^{-t} y_2 e_4$, e_1 , e_3 we get the result. \square

In the sequel, unless otherwise established, z_0 will denote the special point $1/\sqrt{2}(i, i)$.

Corollary 4.3. *The leaves $f_{\psi_s(z_0)} : \text{Sol} \rightarrow \mathbb{H} \times \mathbb{H}$ can be identified with Sol, up to a homothety in the direction spanned by the x, y coordinates.*

Proof. We have

$$\psi_s \circ f_{z_0}(t, x, y) = \left(x, \frac{1}{\sqrt{2}} e^{t+s}, y, \frac{1}{\sqrt{2}} e^{-t+s} \right).$$

If we pullback the induced metric to Sol, we get

$$dt^2 + e^{-2(t+s)} dx^2 + e^{2(t-s)} dy^2.$$

Define $F_s : \text{Sol} \rightarrow \text{Sol}$ by $F_s(t, x, y) = (t, e^s x, e^s y)$. Another pullback with F_s turns the induced metric into the standard metric in Sol. \square

Proposition 4.4. *The foliation is globally rectifiable: there is a diffeomorphism $\Psi : \mathbb{R}^3 \times \mathbb{R} \rightarrow \mathbb{H} \times \mathbb{H}$, such that each hyperplane $\mathbb{R}^3 \times \{c\}$ is diffeomorphic to a leaf.*

Proof. Sol is diffeomorphic to \mathbb{R}^3 in a natural way. Any $\gamma \in \text{Sol}$ is uniquely determined by a triplet (t, x, y) . Define $\Psi : \mathbb{R}^3 \times \mathbb{R} \rightarrow \mathbb{H} \times \mathbb{H}$ by $\Psi(t, x, y, s) = \psi_s \circ f_{z_0}(\gamma)$. The function Ψ is injective because the action is free. Given $z' \in \mathbb{H} \times \mathbb{H}$, there is a leaf going through it, and since $\psi_s(z_0)$ traverses all the leaves, there exists a number s , such that $\psi_{-s}(z')$ is in the leaf passing through z_0 . Let $\gamma \in \text{Sol}$ be such that $\psi_{-s}(z') = f_{z_0}(\gamma)$. Therefore,

$$z' = \psi_s \circ f_{z_0}(\gamma),$$

which implies that Ψ is also surjective. Finally, the Jacobian of Ψ is

$$[d\Psi] = \begin{pmatrix} 0 & 1 & 0 & 0 \\ e^{t+s}/\sqrt{2} & 0 & 0 & e^{t+s}/\sqrt{2} \\ 0 & 0 & 1 & 0 \\ -e^{-t+s}/\sqrt{2} & 0 & 0 & e^{-t+s}/\sqrt{2} \end{pmatrix},$$

which is nondegenerated. By the inverse function theorem, Ψ is a diffeomorphism. The last claim follows from the fact that ψ_s maps leaves onto leaves. \square

Corollary 4.5. *The previous diffeomorphism can be modified, such that it not only maps the foliation to a Cartesian product globally, but also maps each leaf in the foliation isometrically to Sol.*

Proof. The pullback of the metric in $\mathbb{H} \times \mathbb{H}$ with the previous diffeomorphism is

$$dt^2 + e^{-2(t+s)}dx^2 + e^{2(t-s)}dy^2 + ds^2,$$

which is analogous to the expression in Corollary 4.3. Let

$$\tilde{\Psi}(t, x, y, s) = \Psi(t, e^s x, e^s y, s).$$

$\tilde{\Psi}$ is a leaf-preserving diffeomorphism such that, for fixed s , it isometrically maps Sol into the leaf $\mathbb{R}^3 \times \{s\}$. \square

5. Extrinsic geometry

Proposition 5.1. *Integral curves of the normal field X are geodesics.*

Proof. We previously found that the integral curves of the field are given by $\gamma(t) = (x_1, e^t y_1, x_2, e^t y_2)$. Let $\phi(t)$ be a smooth curve in \mathbb{H} with the homothetic metric. Then,

$$\|\dot{\phi}(t)\|^2 = \frac{\dot{x}^2 + \dot{y}^2}{2y^2},$$

which is half the standard hyperbolic square length [Gallot et al. 2004]. Therefore, a curve minimizes hyperbolic arc length if and only if it minimizes the homothetic metric arc length, i.e., geodesics in both cases are the same. It is a well known fact that the vertical curves $(x_k, e^t y_k)$ are geodesics in hyperbolic space. Finally, since γ can be projected in two geodesics and the metric is a product, γ is a geodesic in $\mathbb{H} \times \mathbb{H}$ (see 3.15 in [Gallot et al. 2004]). \square

Proposition 5.2. *There are isometries in $\mathbb{H} \times \mathbb{H}$ acting transitively and sending leaves onto leaves.*

Proof. We work in the $\mathbb{R}^3 \times \mathbb{R}$ picture with the $\tilde{\Psi}$ isometry. By straightforward calculations we have that the mappings

$$(t, x, y, s) \mapsto (t + t', e^{t'+s'}x + x', e^{-t'+s'}y + y', s + s')$$

are isometries. The first claim comes from the fact that given a pair of points (t_k, x_k, y_k, s_k) , there exists exactly one such isometry sending one onto another. That this isometry sends leaves onto leaves is obvious, since under this diffeomorphism, they correspond to hypersurfaces where s is constant. \square

We aim to calculate the distance between any pair of leaves. Recall that in any metric space, the distance from a point p to a set $S \neq \emptyset$ is given by the expression

$$d(p, S) = \inf\{d(p, x) : x \in S\}.$$

See [Munkres 2000] for details.

Proposition 5.3. *The separation between two leaves in $\mathbb{H} \times \mathbb{H}$ is constant. Moreover, if leaves are parametrized with the normal field affine parameter, then leaves' separation is given by the difference $|s - s'|$ between the parameters corresponding to any leaf.*

Proof. A point in a leaf can be parametrized as

$$\left(x, \frac{e^{s+t}}{\sqrt{2}}, y, \frac{e^{s-t}}{\sqrt{2}}\right),$$

where x, y, t are arbitrary, and s is the parameter corresponding to the leaf. Given a second point in another leaf, say,

$$(x', e^{s'+t'}/\sqrt{2}, y', e^{s'-t'}/\sqrt{2}),$$

and since the metric is a product, we can find a geodesic minimizing the arc length in $\mathbb{H} \times \mathbb{H}$, such that, in each factor \mathbb{H} , the distance is also minimized [Gallot et al. 2004]. On the other hand, the metric we use in each factor of $\mathbb{H} \times \mathbb{H}$ is half the hyperbolic

distance, for which a well-known formula gives us the distance [Anderson 2005]. Let ρ_k denote the distance in each factor with our metric, then,

$$\cosh(\sqrt{2}\rho_1) = 1 + \frac{2(x - x')^2 + (e^{s+t} - e^{s'+t'})^2}{2e^{s+t}e^{s'+t'}},$$

$$\cosh(\sqrt{2}\rho_2) = 1 + \frac{2(y - y')^2 + (e^{s-t} - e^{s'-t'})^2}{2e^{s-t}e^{s'-t'}},$$

where the $\sqrt{2}$ factor within the hyperbolic cosine is due to the factor relating the standard hyperbolic metric with ours. The previous expression shows that, in order to get the minimum distance, x' must be equal to x and y' to y . Simplifying the previous expressions for such values of x' and y' , we find

$$\cosh(\sqrt{2}\rho_1) = \cosh(s - s' + t - t'),$$

$$\cosh(\sqrt{2}\rho_2) = \cosh(s - s' + t' - t).$$

Therefore,

$$\rho_1 = \frac{|s - s' + t - t'|}{\sqrt{2}}, \quad \rho_2 = \frac{|s - s' + t' - t|}{\sqrt{2}},$$

and the distance in the product metric is given by $\sqrt{\rho_1^2 + \rho_2^2}$. In order for this distance to be a minimum, a short analysis shows that one must take $t' = t$, and the statement follows. \square

Proposition 5.4. *The principal curvatures of each leaf are -1 with multiplicity two, and 0 . The principal directions are determined by the integral curves of the vectors $\partial_x, \partial_y, \partial_t$ respectively.*

Proof. Recall the principal curvatures and directions for an orientable submanifold M are determined by the *shape operator*, S , which, in codimension 1, can be regarded as the mapping $TM \rightarrow TM$ given by $v_x \mapsto \nabla_{v_x} X$, where X is the normal field to the manifold, compatible with orientation (see [Spivak 1979, Chapter 1]). Here, the principal directions and curvatures are the shape operator eigenvectors, and eigenvalues. Consider a leaf embedded in $\mathbb{H} \times \mathbb{H}$,

$$\left(x, \frac{e^{-t-s}}{\sqrt{2}}, y, \frac{e^{t-s}}{\sqrt{2}} \right),$$

with normal field $X = x_2\partial_2 + x_4\partial_4$, where $x_2 = e^{-t-s}/\sqrt{2}$ and $x_4 = e^{t-s}/\sqrt{2}$. A calculation shows that

$$\nabla X = -dx_1 \otimes \partial_1 - dx_3 \otimes \partial_3,$$

i.e., the shape operator is diagonal, once expressed in the base for the tangent space to the leaf, spanned by the coordinate vectors ∂_1, ∂_3 , and the vector $-x_2\partial_2 + x_4\partial_4$, with eigenvalues $\{-1, -1, 0\}$ counted with multiplicity. \square

5A. Proof of Theorem 1.1. Let G be a complex Kleinian group with a maximum of four lines in general position contained in its limit set, then G acts properly and discontinuously in four disjoint copies of $\mathbb{H} \times \mathbb{H}$. Without loss of generality we can assume that $\mathbb{H} \times \mathbb{H}$ is G -invariant. By Proposition 3.1, if

$$\psi : \text{Sol} \times \mathbb{R} \rightarrow \mathbb{H} \times \mathbb{H}$$

is a G -equivariant diffeomorphism, then $(\mathbb{H} \times \mathbb{H})/G$ is diffeomorphic to $(\text{Sol}/G) \times \mathbb{R}$. We notice the topological type is perfectly determined by the group G . In fact, the group G is the fundamental group of the manifold $(\mathbb{H} \times \mathbb{H})/G$. We remember the Kulkarni discontinuity region is equal to four disjoint copies of $\mathbb{H} \times \mathbb{H}$, hence Ω/G is equal to four disjoint copies of $(\mathbb{H} \times \mathbb{H})/G$. We remark that if G represents a lattice of the Lie group Sol , then Sol/G is a compact 3 manifold. This last statement implies in some sense that Sol/G is the compact heart of $(\mathbb{H} \times \mathbb{H})/G$.

6. The Heisenberg group

Given a symplectic vector space, V , with symplectic form ω , recall the *Heisenberg* group, \mathcal{H} , is the space $V \times \mathbb{R}$, with the product operation given by

$$(v, t) * (w, s) = (v + w, t + s + \omega(v, w)).$$

An account of this group in the context of complex hyperbolic geometry can be found in [Cano et al. 2013]. If V is of dimension 2, and $\{\partial_p, \partial_q\}$ is a symplectic base for V , that is, $\omega(\partial_p, \partial_q) = 1$, a well-known fact from Lie group theory is that there is a faithful representation $\mathcal{H} \rightarrow \text{SL}(3, \mathbb{R})$ [Binz and Pods 2008], given by

$$(p\partial_p + q\partial_q t) \rightarrow \begin{pmatrix} 1 & p & t + \frac{1}{2}pq \\ 0 & 1 & q \\ 0 & 0 & 1 \end{pmatrix}.$$

We will use this representation and identify \mathcal{H} with a subgroup of $\text{SL}(3, \mathbb{R})$. Therefore, we will identify \mathcal{H} with \mathbb{R}^3 , with group structure,

$$(a, b, c) * (a', b', c') = (a + a', b + b', c + c' + ab'),$$

which corresponds to the matrix product

$$\begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & a' & c' \\ 0 & 1 & b' \\ 0 & 0 & 1 \end{pmatrix}.$$

With these identifications, there is a natural left action $\mathcal{H} \circlearrowleft \mathbb{C} \times \mathbb{H}$:

$$\begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} z \\ w \\ 1 \end{pmatrix} = \begin{pmatrix} z + aw + c \\ w + b \\ 1 \end{pmatrix},$$

which we will denote by $(a, b, c) * (z, w)$.

Proposition 6.1. *The action of \mathcal{H} in $\mathbb{C} \times \mathbb{H}$ is free.*

Proof. If $(a, b, c) * (z, w) = (z, w)$, then

$$\begin{aligned} z + aw + c &= z, \\ w + b &= w. \end{aligned}$$

From this linear system, one deduces that $a = b = c = 0$. □

Proposition 6.2. *For fixed $(z, w) \in \mathbb{C} \times \mathbb{H}$, the orbit $h \in \mathcal{H} \mapsto h * (z, w)$ defines a differentiable embedding $\mathcal{H} \hookrightarrow \mathbb{C} \times \mathbb{H}$.*

Proof. The map is injective, since the action is free. Let $w = p + qi$, the Jacobian matrix of the mapping in $(a, b, c) \in \mathcal{H}$ is given by

$$\begin{pmatrix} p & 0 & 1 \\ q & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Since the Jacobian has rank 3, the action defines a local diffeomorphism, and hence an embedding. □

Therefore, the action of \mathcal{H} defines a foliation of $\mathbb{C} \times \mathbb{H}$, in analogy with the foliation of $\mathbb{H} \times \mathbb{H}$ generated by Sol.

Proposition 6.3. *Consider $\mathbb{C} \times \mathbb{H}$ as a subset of \mathbb{R}^4 , but with the product metric of the euclidean metric in \mathbb{C} and the hyperbolic metric in \mathbb{H} . If e_1, \dots, e_4 denote the canonical coordinates in \mathbb{R}^4 , and (p, q) denotes the coordinates in \mathbb{H} , then the vector field $X = qe_4$ is unitary and orthogonal to any leaf of the foliation generated by \mathcal{H} .*

Proof. From Theorem 2.4, the vector fields $pe_1 + qe_2, e_3, e_1$ generate the tangent space to the orbit of $(z, w) \in \mathbb{C} \times \mathbb{H}$, where $w = p + qi$. Since the metric is a product, X is orthogonal to $pe_1 + qe_2$ and e_1 . Moreover, the metric in \mathbb{H} is conformal to the euclidean, and therefore X is orthogonal to e_3 . Finally, qe_4 is unitary in the hyperbolic metric. □

Corollary 6.4. *Let $(z, w) \in \mathbb{C} \times \mathbb{H}$, $z = x + yi$ and $w = p + qi$. The integral curves of X are geodesics.*

Proof. The integral curves of X are constant in the first factor, and vertical straight lines in the hyperbolic factor. □

Although in this case, the action induced by the normal field X is not equivariant, we can describe in a precise way the quotients $(\mathbb{C} \times \mathbb{H})/\Gamma$, where Γ is a discrete subgroup of \mathcal{H} . Moreover, if Γ acts properly discontinuously in $\mathbb{C} \times \mathbb{H}$, it has to act in the same way in Heisenberg, because the slices $\mathcal{H} \times \{qi\}$ are preserved. This is a general property of Lie groups that we prove in the following lemma.

Lemma 6.5. *Let X and Y be two locally compact spaces. If $\Gamma \circlearrowleft X \times Y$, and the action of $g \in \Gamma$ can be decomposed as $g \cdot (x, y) = (g \cdot x, y)$ then Γ acts properly discontinuously in X if and only if it acts properly discontinuously in $X \times Y$.*

Proof. Let $K \subset X$ be a compact set. Fix $y \in Y$. With the product topology, $K \times \{y\}$ is a compact set in $X \times Y$. One can easily verify the equality

$$\{g \in \Gamma : g \cdot K \cap K \neq \emptyset\} = \{g \in \Gamma : g \times 1 \cdot K \times \{y\} \cap K \times \{y\} \neq \emptyset\}.$$

If Γ acts properly discontinuously in $X \times Y$, the previous equality implies that it acts properly discontinuously in X . On the other hand, if $K \subset X \times Y$ is compact, the product topology together with the local compactness implies that we can find an open set $U \times V$, with $U \subset X$ and $V \subset Y$, such that \bar{U} is compact in X , \bar{V} is compact in Y , and $K \subset U \times V$. We have the contention

$$\{g \in \Gamma : gK \cap K \neq \emptyset\} \subset \{g \in \Gamma : g \cdot l(\bar{U} \times \bar{V}) \cap \bar{U} \times \bar{V} \neq \emptyset\}$$

Take $g \in \Gamma$ and $(x, y) \in \bar{U} \times \bar{V}$, such that $g \cdot (x, y) \in \bar{U} \times \bar{V}$. Since $g \cdot (x, y) = (g \cdot x, y)$, it follows that $g \cdot x \in \bar{U}$. Therefore, the second set in the previous contention is at the same time contained in

$$\{g \in \Gamma : g \cdot \bar{U} \cap \bar{U} \neq \emptyset\}.$$

If Γ acts properly discontinuously in X , this set has to be finite, and the same must be true for the set of intersections in $X \times Y$, that is, Γ acts properly discontinuously in $X \times Y$. \square

Proposition 6.6. *$\mathbb{C} \times \mathbb{H}$ is diffeomorphic to $\mathcal{H} \times \mathbb{R}$, where, up to diffeomorphism, \mathcal{H} acts on the first factor only.*

Proof. Let $\gamma = (a, b, c) \in \mathcal{H}$ and take $(0, qi) \in \mathbb{C} \times \mathbb{H}$. We can describe the orbits $\gamma \cdot (0, qi)$ explicitly:

$$\begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 0 \\ qi \\ 1 \end{pmatrix} = \begin{pmatrix} aqi + c \\ qi + b \\ 1 \end{pmatrix}.$$

Therefore, there is exactly one $(0, qi)$ in each orbit of the group action. Define $\Psi : \mathcal{H} \times \mathbb{R} \rightarrow \mathbb{C} \times \mathbb{H}$ as

$$\Psi(\gamma, q) = \gamma \cdot (0, qi).$$

It can be shown that Ψ is bijective. It is a diffeomorphism, since an explicit computation shows that $d\Psi$ maps the canonical vectors $T_{(\gamma, q)}\mathcal{H} \times \mathbb{R} \cong \mathbb{R}^4 \rightarrow T_{\gamma \cdot (0, qi)}\mathbb{C} \times \mathbb{H} \cong \mathbb{R}^4$:

$$\{\partial_1, \dots, \partial_4\} \mapsto \{q\partial_2, \partial_3, \partial_1, a\partial_2 + \partial_4\}.$$

The last assertion follows since the action is associative, i.e., $\gamma' \cdot (\gamma \cdot (0, qi)) = (\gamma' \cdot \gamma) \cdot (0, qi)$, and therefore, preserves the imaginary part on the second factor. \square

6A. Proof of Theorem 1.4. The proof is analogous to that of Theorem 1.1, the only difference is that we need the technical Lemma 6.5. A consequence of Theorem 1.4 is the following corollary:

Corollary 6.7. *If $\Gamma < \mathcal{H}$ is a discrete subgroup acting properly and discontinuously in $\mathbb{C} \times \mathbb{H}$, up to diffeomorphism, $(\mathbb{C} \times \mathbb{H})/\Gamma \cong (\mathcal{H}/\Gamma) \times \mathbb{R}$, and the quotient \mathcal{H}/Γ is a manifold whose fundamental group is $\pi_1(\mathcal{H}/\Gamma) \cong \Gamma$.*

Example 6.8. Let $\mathcal{H}_{\mathbb{Z}} < \mathcal{H}$ be the discrete subgroup of Heisenberg matrices with integer coefficients. It can be shown that the unit cube $K_{\mathbb{C}} = [0, 1]^3 \subset \mathcal{H}$ is a fundamental region for the action of $\mathcal{H}_{\mathbb{Z}}$ [Lukyanenko 2014]. The quotient $\mathcal{H}_{\mathbb{Z}} \backslash \mathcal{H}$ is an example of a nilmanifold, whose fundamental group is

$$\mathcal{H}_{\mathbb{Z}} \cong \langle m, n, k : [m, n] = k^4 \rangle;$$

see [Lukyanenko 2014]. In view of the previous results, $\mathcal{H}_{\mathbb{Z}}$ acts properly and discontinuously in $\mathbb{C} \times \mathbb{H}$, and the quotient $(\mathbb{C} \times \mathbb{H})/\mathcal{H}_{\mathbb{Z}}$ is a product of a nilmanifold times \mathbb{R} , whose fundamental group has the previous presentation.

References

- [Anderson 2005] J. W. Anderson, *Hyperbolic geometry*, 2nd ed., Springer, 2005. MR Zbl
- [Barrera et al. 2011a] W. Barrera, A. Cano, and J. P. Navarrete, “The limit set of discrete subgroups of $\mathrm{PSL}(3, \mathbb{C})$ ”, *Math. Proc. Cambridge Philos. Soc.* **150**:1 (2011), 129–146. MR Zbl
- [Barrera et al. 2011b] W. Barrera, A. Cano, and J. P. Navarrete, “Subgroups of $\mathrm{PSL}(3, \mathbb{C})$ with four lines in general position in its limit set”, *Conform. Geom. Dyn.* **15** (2011), 160–176. MR Zbl
- [Barrera et al. 2014] W. Barrera, A. Cano, and J. P. Navarrete, “Pappus’ theorem and a construction of complex Kleinian groups with rich dynamics”, *Bull. Braz. Math. Soc. (N.S.)* **45**:1 (2014), 25–52. MR Zbl
- [Barrera et al. 2016] W. Barrera, A. Cano, and J. Navarrete, “On the number of lines in the limit set for discrete subgroups of $\mathrm{PSL}(3, \mathbb{C})$ ”, *Pacific J. Math.* **281**:1 (2016), 17–49. MR Zbl
- [Binz and Pods 2008] E. Binz and S. Pods, *The geometry of Heisenberg groups*, Mathematical Surveys and Monographs **151**, Amer. Math. Soc., Providence, RI, 2008. MR Zbl
- [Candel and Conlon 2000] A. Candel and L. Conlon, *Foliations, I*, Graduate Studies in Mathematics **23**, Amer. Math. Soc., Providence, RI, 2000. MR Zbl
- [Cano and Seade 2014] A. Cano and J. Seade, “On discrete groups of automorphisms of $\mathbb{P}_{\mathbb{C}}^2$ ”, *Geom. Dedicata* **168** (2014), 9–60. MR Zbl
- [Cano et al. 2013] A. Cano, J. P. Navarrete, and J. Seade, *Complex Kleinian groups*, Progress in Mathematics **303**, Springer, 2013. MR Zbl
- [Gallot et al. 2004] S. Gallot, D. Hulin, and J. Lafontaine, *Riemannian geometry*, 3rd ed., Springer, 2004. MR Zbl
- [de la Harpe 2000] P. de la Harpe, *Topics in geometric group theory*, University of Chicago Press, 2000. MR Zbl
- [Kulkarni 1978] R. S. Kulkarni, “Groups with domains of discontinuity”, *Math. Ann.* **237**:3 (1978), 253–272. MR Zbl

- [Lukyanenko 2014] A. Lukyanenko, *Geometric mapping theory of the Heisenberg group, sub-Riemannian manifolds, and hyperbolic spaces*, Ph.D. thesis, University of Illinois at Urbana-Champaign, 2014, available at <https://search.proquest.com/docview/1652502708>. MR Zbl
- [Munkres 2000] J. R. Munkres, *Topology*, 2nd ed., Prentice Hall, Upper Saddle River, NJ, 2000. Zbl
- [Myrberg 1925] P. J. Myrberg, “Untersuchungen über die automorphen Funktionen beliebig vieler Variablen”, *Acta Math.* **46**:3-4 (1925), 215–336. MR Zbl
- [Navarrete 2006] J.-P. Navarrete, “On the limit set of discrete subgroups of $PU(2, 1)$ ”, *Geom. Dedicata* **122** (2006), 1–13. MR Zbl
- [Navarrete 2008] J.-P. Navarrete, “The trace function and complex Kleinian groups in $\mathbb{P}_{\mathbb{C}}^2$ ”, *Internat. J. Math.* **19**:7 (2008), 865–890. MR Zbl
- [Scott 1983] P. Scott, “The geometries of 3-manifolds”, *Bull. London Math. Soc.* **15**:5 (1983), 401–487. MR Zbl
- [Seade and Verjovsky 2001] J. Seade and A. Verjovsky, “Actions of discrete groups on complex projective spaces”, pp. 155–178 in *Laminations and foliations in dynamics, geometry and topology* (Stony Brook, NY, 1998), edited by M. Lyubich et al., *Contemp. Math.* **269**, Amer. Math. Soc., Providence, RI, 2001. MR Zbl
- [Spivak 1979] M. Spivak, *A comprehensive introduction to differential geometry*, vol. 3, 2nd ed., Publish or Perish, Wilmington, DE, 1979. MR Zbl
- [Thurston 1997] W. P. Thurston, *Three-dimensional geometry and topology, I*, edited by S. Levy, Princeton Mathematical Series **35**, Princeton University Press, 1997. MR Zbl

Received January 20, 2017. Revised June 5, 2017.

WALDEMAR BARRERA
 FACULTAD DE MATEMÁTICAS
 UNIVERSIDAD AUTÓNOMA DE YUCATÁN
 ANILLO PERIFÉRICO NORTE
 MÉRIDA
 MEXICO
 bvargas@correo.uady.mx

RENE GARCIA-LARA
 SCHOOL OF MATHEMATICS
 UNIVERSITY OF LEEDS
 LEEDS
 ENGLAND
 mmrig@leeds.ac.uk

JUAN NAVARRETE
 FACULTAD DE MATEMÁTICAS
 UNIVERSIDAD AUTÓNOMA DE YUCATÁN
 ANILLO PERIFÉRICO NORTE
 MÉRIDA
 MEXICO
 jp.navarrete@correo.uady.mx

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 294 No. 1 May 2018

Three-dimensional Sol manifolds and complex Kleinian groups	1
WALDEMAR BARRERA, RENE GARCIA-LARA and JUAN NAVARRETE	
On periodic points of symplectomorphisms on surfaces	19
MARTA BATORÉO	
Mixing properties for hom-shifts and the distance between walks on associated graphs	41
NISHANT CHANDGOTIA and BRIAN MARCUS	
Simultaneous construction of hyperbolic isometries	71
MATT CLAY and CAGLAR UYANIK	
A local weighted Axler–Zheng theorem in \mathbb{C}^n	89
ŽELJKO ČUČKOVIĆ, SÖNMEZ ŞAHUTOĞLU and YUNUS E. ZEYTUNCU	
Monotonicity and radial symmetry results for Schrödinger systems with fractional diffusion	107
JING LI	
Moduli spaces of stable pairs	123
YINBANG LIN	
Spark deficient Gabor frames	159
ROMANOS-DIOGENES MALIKIOSIS	
Ordered groups as a tensor category	181
DALE ROLFSEN	
Multiplication of distributions and a nonlinear model in elastodynamics	195
C. O. R. SARRICO	
Some Ambrose- and Galloway-type theorems via Bakry–Émery and modified Ricci curvatures	213
HOMARE TADANO	
Irreducible decomposition for local representations of quantum Teichmüller space	233
JÉRÉMY TOULISSE	



0030-8730(201805)294:1;1-T