

*Pacific  
Journal of  
Mathematics*

**A LOCAL WEIGHTED AXLER–ZHENG THEOREM IN  $\mathbb{C}^n$**

**ŽELJKO ČUČKOVIĆ, SÖNMEZ ŞAHUTOĞLU AND YUNUS E. ZEYTUNCU**

Volume 294 No. 1

May 2018



## A LOCAL WEIGHTED AXLER–ZHENG THEOREM IN $\mathbb{C}^n$

ŽELJKO ČUČKOVIĆ, SÖNMEZ ŞAHUTOĞLU AND YUNUS E. ZEYTUNCU

The well-known Axler–Zheng theorem characterizes compactness of finite sums of finite products of Toeplitz operators on the unit disk in terms of the Berezin transform of these operators. Subsequently this theorem was generalized to other domains and appeared in different forms, including domains in  $\mathbb{C}^n$  on which the  $\bar{\partial}$ -Neumann operator  $N$  is compact. In this work we remove the assumption on  $N$ , and we study weighted Bergman spaces on smooth bounded pseudoconvex domains. We prove a local version of the Axler–Zheng theorem characterizing compactness of Toeplitz operators in the algebra generated by symbols continuous up to the boundary in terms of the behavior of the Berezin transform at strongly pseudoconvex points. We employ a Forelli–Rudin type inflation method to handle the weights.

### 1. Introduction

**1.1. History.** In the theory of Bergman space operators on the open unit disk  $\mathbb{D}$ , the Axler–Zheng theorem [1998] provides an important characterization of compactness of a large class of operators in terms of their Berezin transforms. Specifically this theorem states that if  $S$  is a finite sum of finite products of Toeplitz operators on the Bergman space  $A^2(\mathbb{D})$  whose symbols are in  $L^\infty(\mathbb{D})$ , then  $S$  is compact if and only if  $BS(z)$ , the Berezin transform of  $S$ , tends to 0 as  $|z| \rightarrow 1$ . This theorem has been extended by Suárez [2007] to include all operators in the Toeplitz algebra in the unit ball in  $\mathbb{C}^n$ . Engliš [1999] extended the Axler–Zheng theorem to irreducible bounded symmetric domains and the unit polydisk. Mitkovski, Suárez and Wick [Mitkovski et al. 2013] proved a weighted version of Suárez’s result on the unit ball in  $\mathbb{C}^n$ . Using the techniques of several complex variables, Čučković and Şahutoğlu [2013] proved a version of the Axler–Zheng theorem on smooth bounded pseudoconvex domains on which the  $\bar{\partial}$ -Neumann operator is compact. The use of the  $\bar{\partial}$  techniques required that the operators in their theorem belong to the algebra  $\mathcal{T}(\bar{\Omega})$  which is the norm closed algebra generated by  $\{T_\phi : \phi \in C(\bar{\Omega})\}$ . Recently, Kreutzer [2014] generalized Čučković and Şahutoğlu’s result in a more abstract setting.

---

The work of the third author was partially supported by a grant from the Simons Foundation (#353525).  
*MSC2010:* primary 47B35; secondary 32W05.

*Keywords:* Axler–Zheng theorem, Toeplitz operators, pseudoconvex domains.

Our aim is to extend the previous result of Čučković and Şahutoğlu in two ways: Firstly, we want to remove the hypothesis of the compactness of the  $\bar{\partial}$ -Neumann operator on  $\Omega$ . We also want to consider weighted Bergman spaces. Our main theorem gives a local version of the Axler–Zheng theorem for a wide class of domains in  $\mathbb{C}^n$ . The novelty of our approach is to use the inflation of the domain argument pioneered by Forelli and Rudin [1974] and Ligočka [1989]. The second important ingredient is the B-regularity of the inflated domain which will give us the compactness of  $\bar{\partial}$ , thus replacing the assumption on the compactness of the  $\bar{\partial}$ -Neumann operator. As a corollary we obtain a weighted version of the Axler–Zheng theorem for strongly pseudoconvex domains, which itself is a new result.

**1.2. Preliminaries.** Let  $\Omega$  be a  $C^1$ -smooth bounded pseudoconvex domain in  $\mathbb{C}^n$  with a defining function  $\rho$ . We denote the boundary of  $\Omega$  by  $b\Omega$ . Let  $L^2(\Omega, (-\rho)^r)$  denote the square integrable functions on  $\Omega$  with respect to the measure  $(-\rho)^r dV$  where  $dV$  denotes the Lebesgue measure,  $r \geq 0$ , and

$$A^2(\Omega, (-\rho)^r) = \{f \in L^2(\Omega, (-\rho)^r) : f \text{ is holomorphic}\}.$$

Since  $A^2(\Omega, (-\rho)^r)$  is a closed subspace of  $L^2(\Omega, (-\rho)^r)$ , a bounded orthogonal projection,

$$P_r : L^2(\Omega, (-\rho)^r) \rightarrow A^2(\Omega, (-\rho)^r),$$

(called Bergman projection) exists.  $P_r$  is an integral operator of the form

$$P_r(f)(z) = \int_{\Omega} K^r(z, \xi) f(\xi) (-\rho)^r dV$$

for  $f \in L^2(\Omega, (-\rho)^r)$ . The integral kernel  $K^r(z, \xi)$  is called the Bergman kernel and the normalized Bergman kernel  $k_z^r(\xi)$  is defined as

$$k_z^r(\xi) = \frac{K^r(\xi, z)}{\sqrt{K^r(z, z)}}.$$

When  $r = 0$  we drop the superscript  $r$ ; that is,  $K = K_{\Omega}$  denotes the unweighted Bergman kernel and  $k_z$  denotes the unweighted normalized Bergman kernel. For a bounded operator  $T$  on  $A^2(\Omega, (-\rho)^r)$ , the Berezin transform  $B_r T$  of  $T$  is defined as

$$B_r T(z) = \langle T k_z^r, k_z^r \rangle_r,$$

where  $\langle \cdot, \cdot \rangle_r$  is the inner product on  $A^2(\Omega, (-\rho)^r)$ .

For  $\phi \in L^{\infty}(\Omega)$ , the weighted Toeplitz operator  $T_{\phi}^r$  and the weighted Hankel operator  $H_{\phi}^r$  are defined as

$$\begin{aligned} T_{\phi}^r &= P^r M_{\phi}, \\ H_{\phi}^r &= (I - P^r) M_{\phi}, \end{aligned}$$

where  $M_{\phi} : A^2(\Omega, (-\rho)^r) \rightarrow L^2(\Omega, (-\rho)^r)$  denotes the multiplication by  $\phi$ .

We use  $\mathcal{T}(\bar{\Omega}, (-\rho)^r)$  to denote the norm closed subalgebra of bounded linear operators on  $A^2(\Omega, (-\rho)^r)$  generated by the set of Toeplitz operators

$$\{T_\phi^r : \phi \in C(\bar{\Omega})\}.$$

For  $\phi \in L^\infty$  we define  $B_r\phi = B_r T_\phi$ .

In this paper we look at weighted Hankel and Toeplitz operators on various domains and various weighted spaces. Whenever we need to clarify where these operators are defined, we will use appropriate subscripts and superscripts. In particular, when we need to emphasize the underlying domain we will write  $P^\Omega$ ,  $K_\Omega(z, \xi)$ ,  $H_\phi^\Omega$ , and  $T_\phi^\Omega$ , where the Bergman spaces are unweighted. When we have weighted spaces and we need to indicate the domain and the weight we will write  $P^{\Omega,r}$ ,  $K_\Omega^r(z, \xi)$ ,  $H_\phi^{\Omega,r}$ , and  $T_\phi^{\Omega,r}$ .

**1.3. Main result.** We start with the following two definitions that capture the local structure of the main theorem. To motivate the following definition, if  $f_j \rightarrow f$  weakly in  $A^2(\Omega)$  then for any point  $p \in b\Omega$  and  $r > 0$  one can show that  $f_j \rightarrow f$  weakly in  $A^2(\Omega \cap B(p, r))$  where  $B(p, r)$  is the open ball centered at  $p$  with radius  $r$ .

**Definition 1.** Let  $r \geq 0$  and  $\Omega$  be a  $C^2$ -smooth bounded pseudoconvex domain in  $\mathbb{C}^n$  with a defining function  $\rho$ . Furthermore, let  $\{f_j\} \subset A^2(\Omega, (-\rho)^r)$  be a sequence and  $f \in A^2(\Omega, (-\rho)^r)$ . We say that  $\{f_j\}$  converges to  $f$  *weakly about strongly pseudoconvex points* if:

- (i)  $f_j \rightarrow f$  weakly in  $A^2(\Omega, (-\rho)^r)$  as  $j \rightarrow \infty$ .
- (ii) When  $\Gamma_\Omega$ , the set of the weakly pseudoconvex points in  $b\Omega$ , is nonempty, there exists an open neighborhood  $U$  of  $\Gamma_\Omega$  such that  $\|f_j - f\|_{L^2(U \cap \Omega, (-\rho)^r)} \rightarrow 0$  as  $j \rightarrow \infty$ .

We note that on strongly pseudoconvex domains, sequences converging weakly about strongly pseudoconvex points and weakly convergent sequences coincide.

**Definition 2.** Let  $r$ ,  $\Omega$ , and  $\rho$  be as above. Furthermore, let  $T : A^2(\Omega, (-\rho)^r) \rightarrow A^2(\Omega, (-\rho)^r)$  be a bounded linear operator. We say that  $T$  is *compact about strongly pseudoconvex points* if  $Tf_j \rightarrow Tf$  in  $A^2(\Omega, (-\rho)^r)$  whenever  $f_j \rightarrow f$  weakly about strongly pseudoconvex points.

**Remark 3.** As shown in Proposition 13 below, it is interesting that any Hankel operator with a symbol continuous on the closure of the domain is compact about strongly pseudoconvex points.

With the help of these two definitions, we state our main result as follows.

**Theorem 4.** *Let  $r$  be a nonnegative real number,  $\Omega$  be a  $C^2$ -smooth bounded pseudoconvex domain in  $\mathbb{C}^n$  with a defining function  $\rho$ , and  $T \in \mathcal{T}(\bar{\Omega}, (-\rho)^r)$ .*

Then  $T$  is compact about strongly pseudoconvex points on  $A^2(\Omega, (-\rho)^r)$  if and only if  $\lim_{z \rightarrow p} B_r T(z) = 0$  for any strongly pseudoconvex point  $p \in b\Omega$ .

If  $\Omega$  is a strongly pseudoconvex domain then we have the following corollary.

**Corollary 5.** *Let  $r$  be a nonnegative real number,  $\Omega$  be a  $C^2$ -smooth bounded strongly pseudoconvex domain in  $\mathbb{C}^n$  with a defining function  $\rho$ , and  $T$  be an element of  $\mathcal{T}(\bar{\Omega}, (-\rho)^r)$ . Then  $T$  is compact on  $A^2(\Omega, (-\rho)^r)$  if and only if  $\lim_{z \rightarrow p} B_r T(z) = 0$  for any  $p \in b\Omega$ .*

**Remark 6.** In the case of the unit ball  $\mathbb{B}^n$  in  $\mathbb{C}^n$  and  $\rho(z) = |z|^2 - 1$ , we partially recover [Mitkovski et al. 2013, Theorem 1.1]. Unlike the arguments on the unit ball, the proof of Corollary 5 does not require any explicit form for the weight or the weighted Bergman kernel.

## 2. Proof of Theorem 4

In this section, before we prove Theorem 4, we present some propositions and lemmas that encapsulate the technical details of the proof.

**Proposition 7.** *Let  $\Omega$  be a  $C^2$ -smooth bounded pseudoconvex domain in  $\mathbb{C}^n$  and  $\{T_j\}$  be a sequence of operators compact about strongly pseudoconvex points that converges to  $T$  in the operator norm. Then  $T$  is compact about strongly pseudoconvex points.*

*Proof.* Let  $\{f_j\}$  be a sequence in  $A^2(\Omega, (-\rho)^r)$  that converges to 0 weakly about strongly pseudoconvex points. Since  $f_j \rightarrow 0$  weakly there exists  $C > 0$  such that

$$\sup\{\|f_j\| : j = 1, 2, 3, \dots\} \leq C.$$

Then for any  $k$  we have

$$\|Tf_j\| \leq \|(T - T_k)f_j\| + \|T_k f_j\| \leq C\|T - T_k\| + \|T_k f_j\|.$$

Let  $\varepsilon > 0$  be given. Since  $T_j \rightarrow T$  in the operator norm, we choose  $k_\varepsilon$  such that  $\|T - T_{k_\varepsilon}\| \leq \varepsilon$ . Then

$$\limsup_{j \rightarrow \infty} \|Tf_j\| \leq C\varepsilon + \limsup_{j \rightarrow \infty} \|T_{k_\varepsilon} f_j\| \leq C\varepsilon.$$

Since  $\varepsilon > 0$  was arbitrary, we conclude that  $Tf_j \rightarrow 0$ . That is,  $T$  is compact about strongly pseudoconvex points.  $\square$

One of the key ideas in the proof is to use an inflated domain over  $\Omega$  to understand the weighted Bergman spaces. For this purpose, unless stated otherwise, for the rest of the paper,  $\Omega$  will be a bounded pseudoconvex domain in  $\mathbb{C}^n$  with  $C^2$ -smooth boundary,  $\rho$  will be a defining function for  $\Omega$ , and

$$(1) \quad \Omega_r^p = \{(z, w) \in \mathbb{C}^n \times \mathbb{C}^p : z \in \Omega \text{ and } \rho(z) + |w_1|^{2p/r} + \dots + |w_p|^{2p/r} < 0\},$$

where  $p$  is a positive integer and  $r$  is a real number such that  $0 < r \leq p$ . For a function  $f \in A^2(\Omega, (-\rho)^r)$ , we let  $F(z, w) = f(z)$  be the trivial extension of  $f$  to  $\Omega_r^p$ . It easily follows from an iterated integral argument that  $F \in A^2(\Omega_r^p)$ .

The following proposition is interesting in its own right as it gives a relationship between the Bergman kernels of the inflated domain and base.

**Proposition 8.** *Using the notation above,*

$$K_{\Omega}^r(z, \xi) = c_{p,r} K_{\Omega_r^p}(z, 0; \xi, 0),$$

where  $c_{p,r} = \int_{|\tilde{w}_1|^{2p/r} + \dots + |\tilde{w}_p|^{2p/r} < 1} dV(\tilde{w})$  and  $K_{\Omega_r^p}^r(z, \xi)$  is the weighted Bergman kernel of  $\Omega$  with weight  $(-\rho)^r$ .

*Proof.* We will follow a standard inflation argument (see, for instance, [Forelli and Rudin 1974; Ligočka 1989]). Since  $\Omega_r^p$  is a Hartogs domain with base  $\Omega$ , the Bergman kernel of  $\Omega_r^p$  can be written as

$$K_{\Omega_r^p}(z, w; \xi, \eta) = K_{\Omega_r^p}(z, 0; \xi, 0) + \sum_{|J| \geq 1} K_J(z, \xi) w^J \bar{\eta}^J,$$

where  $J$  is a multiindex with nonnegative entries. Then for  $f \in A^2(\Omega, (-\rho)^r)$  and  $z \in \Omega$  we have ( $F$  below is the trivial extension of  $f$ )

$$(2) \quad f(z) = \int_{\Omega_r^p} K_{\Omega_r^p}(z, 0; \xi, 0) F(\xi, \eta) dV(\xi, \eta) + \sum_{|J| \geq 1} \int_{\Omega_r^p} K_J(z, \xi) w^J \bar{\eta}^J F(\xi, \eta) dV(\xi, \eta).$$

However, the integrals under the sum on the right-hand side above all vanish.

Using the change of variables  $\tilde{w}_j = w_j / (-\rho(z))^{r/2p}$ , one can compute that

$$(3) \quad \int_{|w_1|^{2p/r} + \dots + |w_p|^{2p/r} < -\rho(z)} dV(w) = (-\rho(z))^r \int_{|\tilde{w}_1|^{2p/r} + \dots + |\tilde{w}_p|^{2p/r} < 1} dV(\tilde{w}).$$

We denote

$$(4) \quad c_{p,r} = \int_{|\tilde{w}_1|^{2p/r} + \dots + |\tilde{w}_p|^{2p/r} < 1} dV(\tilde{w}).$$

Then using (2), (3), and (4) we get

$$\begin{aligned} f(z) &= \int_{\Omega_r^p} K_{\Omega_r^p}(z, 0; \xi, 0) F(\xi, \eta) dV(\xi, \eta) \\ &= c_{p,r} \int_{\Omega} K_{\Omega_r^p}(z, 0; \xi, 0) f(\xi) (-\rho(\xi))^r dV(\xi). \end{aligned}$$

Therefore,  $c_{p,r} K_{\Omega_r^p}(z, 0; \xi, 0) = K_{\Omega}^r(z, \xi)$ . □

For a  $C^2$ -smooth function  $\rho$  around a point  $P \in \mathbb{C}^n$ ,  $X = (x_1, \dots, x_n) \in \mathbb{C}^n$ , and  $Y = (y_1, \dots, y_n) \in \mathbb{C}^n$ , we define the complex Hessian of  $\rho$  at  $P$  as

$$H_\rho(P; X, Y) = \sum_{j,k=1}^n \frac{\partial^2 \rho(P)}{\partial z_j \partial \bar{z}_k} x_j \bar{y}_k.$$

Furthermore, we use the notation  $H_\rho(P; X) = H_\rho(P; X, X)$ .

**Lemma 9.** *Let  $\Omega$  be a  $C^2$ -smooth bounded pseudoconvex domain in  $\mathbb{C}^n$ ,  $z_0 \in b\Omega$  be a strongly pseudoconvex point, and  $\Omega_r^p$  be defined as in (1). Then there exists  $s > 0$  such that  $(z, w) \in b\Omega_r^p$  is strongly pseudoconvex for  $|z - z_0| < s$  and  $w_k \neq 0$  for all  $1 \leq k \leq p$ .*

*Proof.* Let  $\tilde{\rho}(z, w) = \rho(z) + \lambda(w)$  where  $\lambda(w) = |w_1|^{2p/r} + \dots + |w_p|^{2p/r}$  and  $p \geq r$  an integer. Then  $\tilde{\rho}$  is a  $C^2$ -smooth function. Assume that  $Q = (z, w) \in b\Omega_r^p$  is near  $z_0$  and  $X$  is a complex tangential vector to  $b\Omega_r^p$  at  $Q$ . Then  $X$  can be written as  $X = X_n + X_p$  where  $X_n$  and  $X_p$  are the components of  $X$  in the  $z$  and  $w$  variables, respectively. Then

$$H_{\tilde{\rho}}(Q; X) = H_\rho(z; X_n) + H_{\tilde{\rho}}(Q; X_n, X_p) + H_{\tilde{\rho}}(Q; X_p, X_n) + H_\lambda(w; X_p).$$

However,  $H_{\tilde{\rho}}(Q; X_n, X_p) = H_{\tilde{\rho}}(Q; X_p, X_n) = 0$  as  $z$  and  $w$  are decoupled in  $\tilde{\rho}$ . Then

$$H_{\tilde{\rho}}(Q; X) = H_\rho(z; X_n) + H_\lambda(w; X_p).$$

Let  $\pi$  denote the projection from a neighborhood of  $b\Omega$  in  $\mathbb{C}^n$  onto  $b\Omega$ . Then  $X_n = X_t + X_v$  where  $X_t$  is a tangential vector to  $b\Omega$  at  $\pi z$  and  $X_v$  is a vector complex normal to  $b\Omega$  at  $\pi z$ . Then

$$H_\rho(z; X_n) = H_\rho(z; X_t) + H_\rho(z; X_t, X_v) + H_\rho(z; X_v, X_t) + H_\rho(z; X_v).$$

We note that the complex Hessian  $H_\rho$  changes continuously and  $w \rightarrow 0$  as  $z \rightarrow z_0$  (here we assume  $(z, w) \in b\Omega_r^p$ ). Furthermore,  $X_v \rightarrow 0$  as  $z \rightarrow z_0$  (as the complex normal to  $b\Omega$  at  $z_0$  is parallel to the complex normal to  $b\Omega_r^p$  at  $(z_0, 0)$ ). Then, using the fact that  $z_0$  is a strongly pseudoconvex point, we conclude that there exists  $s > 0$  so that

$$H_\rho(z; X_n) \geq \frac{H_\rho(\pi z; X_t)}{2} > 0$$

for  $|z - z_0| < s$  and  $X_t \neq 0$ . Also  $H_\lambda(w; X_p) > 0$  whenever  $X_p \neq 0$  and  $w_k \neq 0$  for all  $k$  as  $\lambda$  is strictly plurisubharmonic whenever  $w_k \neq 0$  for all  $k$ . Therefore,  $H_{\tilde{\rho}}(Q; X) > 0$  for  $Q = (z, w) \in b\Omega_r^p$  such that  $|z - z_0| < s$  and  $w_k \neq 0$  for all  $k$ .  $\square$

The following corollary follows from the previous lemma together with the fact that  $\Omega_r^p$  has  $C^2$ -smooth boundary for  $0 < r \leq p$ .



**Corollary 10.** *Let  $\Omega$  be a  $C^2$ -smooth bounded pseudoconvex domain in  $\mathbb{C}^n$ ,  $z_0 \in b\Omega$  be a strongly pseudoconvex point, and  $\Omega_r^p$  be defined as in (1). Then there exists  $\varepsilon > 0$  such that  $B((z_0, 0), \varepsilon) \cap \Omega_r^p$  is pseudoconvex.*

Next we will prove some statements about compactness of single Toeplitz and Hankel operators.

**Lemma 11.** *Let  $\phi \in L^\infty(\Omega)$ ,  $\{f_j\}$  be a bounded sequence in  $A^2(\Omega, (-\rho)^r)$  and  $F_j$  be the trivial extension of  $f_j$  to  $\Omega_r^p$  for each  $j$  where  $\Omega_r^p$  is defined as in (1). Assume that  $\{H_\phi^{\Omega_r^p} F_j\}$  is convergent in  $L^2(\Omega_r^p)$ . Then  $\{H_\phi^{\Omega, r} f_j\}$  is convergent in  $L^2(\Omega, (-\rho)^r)$ .*

*Proof.* We will abuse the notation and denote the trivial extension of  $\phi$  to  $\Omega_r^p$  by  $\phi$ . We assume that  $\{H_\phi^{\Omega_r^p} F_j\}$  is convergent (and hence Cauchy). Let

$$G_j(z, w) = (H_\phi^{\Omega_r^p} F_j)(z, w)$$

and  $g_j(z) = G_j(z, 0)$ . Then  $G_j$  is holomorphic in  $w$  because

$$\frac{\partial G_j}{\partial \bar{w}_k} = \frac{\partial}{\partial \bar{w}_k} (I - P^{\Omega_r^p})(F_j \phi) = \frac{\partial (F_j \phi)}{\partial \bar{w}_k} = 0$$

for all  $j$  and  $1 \leq k \leq p$ . We note that  $\partial(F_j \phi)/\partial \bar{w}_k = 0$  as  $F_j \phi$  is independent of  $w_k$ . Then  $|G_j(z, w) - G_k(z, w)|^2$  is subharmonic in  $w$  and using the mean value property for subharmonic functions together with (3) and (4) one can show that

$$\begin{aligned} & |g_j(z) - g_k(z)|^2 \\ & \leq \frac{1}{c_{p,r}(-\rho(z))^r} \int_{|w_1|^{2p/r} + \dots + |w_p|^{2p/r} < -\rho(z)} |G_j(z, w) - G_k(z, w)|^2 dV(w) \end{aligned}$$

for  $j = 1, 2, \dots$  and  $z \in \Omega$ . By integrating over  $\Omega$  we get

$$c_{p,r} \|g_j - g_k\|_{L^2_{(0,1)}(\Omega, (-\rho)^r)}^2 \leq \|G_j - G_k\|_{L^2_{(0,1)}(\Omega_r^p)}^2$$

for  $j, k = 1, 2, \dots$ . Then  $\{g_j\}$  is a Cauchy sequence in  $L^2_{(0,1)}(\Omega, (-\rho)^r)$  (and hence convergent) because  $\|G_j - G_k\|_{L^2_{(0,1)}(\Omega_r^p)} \rightarrow 0$  as  $j, k \rightarrow \infty$ .

Let  $h_j(z) = P^{\Omega_r^p}(\phi F_j)(z, 0)$ . Then

$$c_{r,p} \|h_j\|_{L^2(\Omega, (-\rho)^r)}^2 \leq \|P^{\Omega_r^p}(\phi F_j)\|_{L^2(\Omega_r^p)}^2 \leq \|\phi F_j\|_{L^2(\Omega_r^p)}^2 = c_{r,p} \|\phi f_j\|_{L^2(\Omega, (-\rho)^r)}^2 < \infty$$

for each  $j$ . Hence,  $h_j \in A^2(\Omega, (-\rho)^r)$  and  $(I - P^{\Omega, r})h_j = 0$  for all  $j$ . We get equality between the last terms above because  $F_j$  and  $\phi$  are independent of  $w$ . Now

$$\begin{aligned} (I - P^{\Omega, r})g_j &= (I - P^{\Omega, r})(\phi f_j - P^{\Omega_r^p}(\phi F_j)(\cdot, 0)) \\ &= (I - P^{\Omega, r})(\phi f_j) - (I - P^{\Omega, r})(h_j) = H_\phi^{\Omega, r} f_j. \end{aligned}$$

Therefore, the sequence  $\{H_\phi^{\Omega, r} f_j\}$  is convergent in  $L^2(\Omega, (-\rho)^r)$ .  $\square$

**Lemma 12.** *Let  $r$  be a nonnegative real number and  $\Omega$  be a  $C^2$ -smooth bounded pseudoconvex domain in  $\mathbb{C}^n$  with a defining function  $\rho$ . Assume that  $\phi \in C(\bar{\Omega})$  such that  $\phi(z) = 0$  if  $z$  is a strongly pseudoconvex point in  $b\Omega$ . Then  $T_\phi^r$  is compact about strongly pseudoconvex points on  $A^2(\Omega, (-\rho)^r)$ .*

*Proof.* Let  $\{f_j\}$  be a sequence in  $A^2(\Omega, (-\rho)^r)$  that (without loss of generality) converges to 0 weakly about strongly pseudoconvex points. Then  $f_j \rightarrow 0$  weakly as  $j \rightarrow \infty$  and there is a neighborhood  $U$  of weakly pseudoconvex points in  $b\Omega$  such that

$$\|f_j\|_{L^2(U \cap \Omega, (-\rho)^r)} \rightarrow 0 \text{ as } j \rightarrow \infty.$$

Using the uniform boundedness principle and the fact that  $f_j \rightarrow 0$  weakly we conclude that the sequence  $\{f_j\}$  is bounded in  $A^2(\Omega, (-\rho)^r)$ . Furthermore, Cauchy estimates together with Montel's theorem (and the fact that  $f_j \rightarrow 0$  weakly) imply that  $\{f_j\}$  converges to zero uniformly on compact subsets of  $\Omega$ . Using the fact that  $\phi = 0$  on strongly pseudoconvex points, one can show that  $\phi f_j \rightarrow 0$  in  $A^2(\Omega, (-\rho)^r)$ . Therefore,  $T_\phi^r f_j \rightarrow 0$  in  $A^2(\Omega, (-\rho)^r)$ . That is,  $T_\phi^r$  is compact about strongly pseudoconvex points on  $A^2(\Omega, (-\rho)^r)$ .  $\square$

Let  $\Omega$  be a domain in  $\mathbb{C}^n$ . Then  $z \in b\Omega$  is said to have a holomorphic (plurisubharmonic) peak function if there exists a holomorphic (plurisubharmonic)  $f$  that is continuous on  $\bar{\Omega}$  such that  $f(z) = 1$  and  $|f(w)| < 1$  (or  $f(w) < 1$  if  $f$  is plurisubharmonic) for  $w \in \bar{\Omega} \setminus \{z\}$ .

Next we show that any Hankel operator with a symbol continuous on the closure of the domain is compact about strongly pseudoconvex points.

**Proposition 13.** *Let  $r$  be a nonnegative real number,  $\Omega$  be a  $C^2$ -smooth bounded pseudoconvex domain in  $\mathbb{C}^n$  with a defining function  $\rho$ , and  $\phi \in C(\bar{\Omega})$ . Then*

$$H_\phi^r : A^2(\Omega, (-\rho)^r) \rightarrow L^2(\Omega, (-\rho)^r)$$

*is compact about strongly pseudoconvex points.*

*Proof.* We will prove more (see Corollary 14 below). First of all, for any  $\phi \in C(\bar{\Omega})$  there exists  $\{\phi_j\} \subset C^1(\bar{\Omega})$  such that  $\phi_j \rightarrow \phi$  uniformly on  $\bar{\Omega}$  as  $j \rightarrow \infty$ . Furthermore,  $\{H_{\phi_j}^r\}$  converges to  $H_\phi^r$  in the operator norm and, by Proposition 7, if  $H_{\phi_j}^r$  is compact about strongly pseudoconvex points for every  $j$  then so is  $H_\phi^r$ . Therefore, for the rest of the proof we will assume that  $\phi \in C^1(\bar{\Omega})$ . Secondly, the proof for  $r = 0$  does not require the inflation argument in the next paragraph and hence it is easier than the case  $r > 0$ . Since both proofs are similar, except for the inflation argument, in the rest of the proof, we will assume that  $r > 0$ .

Let  $z_0 \in b\Omega$  be a strongly pseudoconvex point. Then, by Corollary 10, the domain  $B((z_0, 0), \varepsilon) \cap \Omega_r^p$  is pseudoconvex for small  $\varepsilon$ . Let  $\varepsilon > 0$  be such that

$X_0 = b\Omega \cap \overline{B((z_0, 0), \varepsilon)} \subset \mathbb{C}^n$  consists of strongly pseudoconvex points. Let us define

$$Y = b\Omega_r^p \cap \overline{B((z_0, 0), \varepsilon)} \cap \{(z, w) \in \mathbb{C}^n \times \mathbb{C}^p : w_k = 0 \text{ for some } 1 \leq k \leq p\},$$

$$X_j = b\Omega_r^p \cap \overline{B((z_0, 0), \varepsilon)} \cap \{(z, w) \in \mathbb{C}^n \times \mathbb{C}^p : |w_k| \geq 1/j \text{ for all } 1 \leq k \leq p\},$$

for  $j = 1, 2, 3, \dots$ . Then  $X_0$  is B-regular as any point in  $X_0$  has a holomorphic (hence plurisubharmonic) peak function on  $\Omega \subset \mathbb{C}^n$ . The same function (by extending it trivially) is also a plurisubharmonic peak function on  $\Omega_r^p \subset \mathbb{C}^{n+p}$ . Hence,  $X_0 \times \{0\}$  is B-regular as a compact set in  $\mathbb{C}^{n+p}$ . Furthermore, Lemma 9 implies that we can shrink  $\varepsilon$ , if necessary, so that  $X_j$ 's are composed of strongly pseudoconvex points for  $j \geq 1$ . Hence,  $X_j$  is B-regular for every  $j = 0, 1, 2, \dots$ .

Next we will apply a similar idea to  $Y$  in lower dimensions. Let us define  $Y_1 = \bigcup_{m=1}^p Y_1^m$  where

$$Y_1^m = b\Omega_r^p \cap \overline{B((z_0, 0), \varepsilon)} \cap \{(z, w) \in \mathbb{C}^n \times \mathbb{C}^p : w_k = 0 \text{ for } k \neq m\}.$$

We can write  $Y_1^m$  as the union of  $X_0 \times \{0\}$  together with the compact sets

$$b\Omega_r^p \cap \overline{B((z_0, 0), \varepsilon)} \cap \{(z, w) \in \mathbb{C}^n \times \mathbb{C}^p : |w_m| \geq 1/j, w_k = 0 \text{ for } k \neq m\}$$

for  $j = 1, 2, 3, \dots$ . However, we can think of the sets above as subsets in  $\mathbb{C}^n \times \mathbb{C}$  and (by Lemma 9) they are composed of strongly pseudoconvex points. Hence, they are B-regular. Then [Sibony 1987, Proposition 1.9] implies that each  $Y_1^m$  is B-regular as it is a countable union of B-regular sets. Hence, applying Sibony's proposition again, we conclude that  $Y_1$  is B-regular. Similarly, we can define  $Y_2 \subset Y$  as a countable union of compact sets where all but at most two  $w_k$  are equal to 0. Using the same reasoning above adopted for  $Y_2$  we can conclude that  $Y_2$  is B-regular. In a similar fashion, we can define  $Y_l$  for  $1 \leq l \leq p-1$  and prove that all of them are B-regular. Hence  $Y = \left(\bigcup_{l=1}^p Y_l\right) \cup (X_0 \times \{0\})$  is B-regular. Then

$$b(\Omega_r^p \cap B((z_0, 0), \varepsilon)) \subset Y \cup \left(\bigcup_{j=1}^{\infty} X_j\right) \cup (X_0 \times \{0\}) \cup bB((z_0, 0), \varepsilon)$$

is B-regular (satisfies property (P) in Catlin's terminology) and, therefore, the  $\bar{\partial}$ -Neumann operator on  $\Omega_r^p \cap B((z_0, 0), \varepsilon)$  is compact [Straube 2010, Theorem 4.8; Catlin 1984]. Then  $H_\phi^{\Omega_r^p \cap B((z_0, 0), \varepsilon)}$  is compact (see [Straube 2010, Proposition 4.1]) and Lemma 11 implies that  $H_\phi^{\Omega \cap B((z_0, 0), \varepsilon), r}$  is compact.

Next we will use local compact solution operators to show that  $H_\phi^r$  is compact about strongly pseudoconvex points. Let  $\{f_j\} \subset A^2(\Omega, (-\rho)^r)$  be a sequence weakly convergent about strongly pseudoconvex points. Then there exists an open neighborhood  $U$  of the set of weakly pseudoconvex points in  $b\Omega$  such that

- (i)  $\{f_j\}$  is weakly convergent,
- (ii)  $\|f_j - f_k\|_{L^2(U \cap \Omega, (-\rho)^r)} \rightarrow 0$  as  $j, k \rightarrow \infty$ .

Let us choose  $\{p_k : k = 1, \dots, m\} \subset b\Omega \setminus U$  and  $\varepsilon_k > 0$  (for  $k = 1, \dots, m$ ) such that

- (i)  $b\Omega \setminus U \subset \bigcup_{k=1}^m B(p_k, \varepsilon_k)$ ,
- (ii)  $H_\phi^{k,r} = H_\phi^{B(p_k, \varepsilon_k) \cap \Omega, r}$  is compact on  $A^2(B(p_k, \varepsilon_k) \cap \Omega, (-\rho)^r)$  for  $k = 1, \dots, m$ .

Let us choose a strongly pseudoconvex domain  $\Omega_{-1} \Subset \Omega$  and smooth cut-off functions  $\chi_{-1} \in C_0^\infty(\Omega_{-1})$ ,  $\chi_0 \in C_0^\infty(U)$ , and  $\chi_k \in C_0^\infty(B(p_k, \varepsilon))$  for  $k = 1, \dots, m$  such that  $\sum_{k=-1}^m \chi_k \equiv 1$  on  $\bar{\Omega}$ .

Let  $H_\phi^{-1,r} = H_\phi^{\Omega_{-1}, r}$ ,  $H_\phi^{0,r} = H_\phi^{U \cap \Omega, r}$ , and  $g_j = \sum_{k=-1}^m \chi_k H_\phi^{k,r} f_j$ . We note that  $H_\phi^{-1,r}$  is compact as  $\Omega_{-1} \Subset \Omega$  is strongly pseudoconvex (and  $\rho < 0$  on the closure of  $\Omega_{-1}$ );  $\{H_\phi^{0,r} f_j\}$  is convergent as  $\{f_j\}$  is convergent in  $L^2(U \cap \Omega, (-\rho)^r)$ ; and by the previous part of this proof,  $H_\phi^{k,r}$  is compact for each  $k = 1, \dots, m$ . Therefore,  $\{g_j\}$  is convergent in  $L^2(\Omega, (-\rho)^r)$ . Furthermore,

$$\bar{\partial} g_j = f_j \bar{\partial} \phi + \sum_{k=-1}^m (\bar{\partial} \chi_k) H_\phi^{k,r} f_j.$$

Then  $\{\sum_{k=-1}^m (\bar{\partial} \chi_k) H_\phi^{k,r} f_j\}$  is a convergent sequence of  $\bar{\partial}$ -closed  $(0, 1)$ -forms as both  $\bar{\partial} g_j$  and  $f_j \bar{\partial} \phi$  are  $\bar{\partial}$ -closed. Let  $Z^r : L^2_{(0,1)}(\Omega, (-\rho)^r) \rightarrow L^2(\Omega, (-\rho)^r)$  be a bounded linear solution operator to  $\bar{\partial}$  (see [Hörmander 1965]). Let

$$h_j = g_j - Z^r \sum_{k=-1}^m (\bar{\partial} \chi_k) H_\phi^{k,r} f_j.$$

Then  $\{h_j\}$  is convergent and  $\bar{\partial} h_j = f_j \bar{\partial} \phi$ . So by taking projection on the orthogonal complement of  $A^2(\Omega, (-\rho)^r)$  we get  $(I - P^r)h_j = H_\phi^r f_j$ . Therefore,  $\{H_\phi^r f_j\}$  is convergent.  $\square$

Using the proof of the proposition above we get the following corollary.

**Corollary 14.** *Let  $r$  be a nonnegative real number and  $\Omega$  be a  $C^2$ -smooth bounded pseudoconvex domain in  $\mathbb{C}^n$  with a defining function  $\rho$ . Assume that  $\Omega$  satisfies property (P) of Catlin (or B-regularity of Sibony). Then*

- (i)  $\bar{\partial}$  has a compact solution operator on  $K^2_{(0,1)}(\Omega, (-\rho)^r)$ , the weighted  $\bar{\partial}$ -closed  $(0, 1)$ -forms,
- (ii)  $H_\phi^r : A^2(\Omega, (-\rho)^r) \rightarrow L^2(\Omega, (-\rho)^r)$  is compact for all  $\phi \in C(\bar{\Omega})$ .

*Proof.* Since (ii) follows from (i), we will only prove (i). By a theorem of Diederich and Fornæss [1977], there exists a  $C^2$ -smooth defining function  $\rho_1$  and  $0 < \eta \leq 1$  such that  $-(-\rho_1)^\eta$  is a strictly plurisubharmonic exhaustion function for  $\Omega$ . Since  $\rho_1$  and  $\rho$  are comparable on  $\bar{\Omega}$  it is enough to prove that  $\bar{\partial}$  has a compact solution operator on  $K^2_{(0,1)}(\Omega, (-\rho_1)^r)$ .

Let  $s = r/\eta \geq 0$  and  $q$  be an integer such that  $s \leq q$ . We define

$$\Omega_s^q = \{(z, w) \in \mathbb{C}^n \times \mathbb{C}^q : -(-\rho_1(z))^\eta + \lambda(w) < 0\},$$

where  $\lambda(w) = |w_1|^{2q/s} + \dots + |w_q|^{2q/s}$ . Then  $-(\rho_1)^\eta + \lambda$  is a bounded  $C^2$ -smooth plurisubharmonic function and  $\Omega_s^q$  is pseudoconvex. Furthermore, the first part of the proof of Proposition 13 shows that  $\Omega_s^q$  satisfies property (P).

Let  $\{f_j\}$  be a bounded sequence in  $K_{(0,1)}^2(\Omega, (-\rho_1)^r)$ . Then  $\{F_j\}$  is a bounded sequence in  $K_{(0,1)}^2(\Omega_s^q)$ . As shown in the first part of this proof,  $\Omega_s^q$  is a bounded (not necessarily  $C^2$ -smooth) pseudoconvex domain with property (P). Then  $\{\bar{\partial}^* N^{\Omega_s^q} F_j\}$  has a convergent subsequence in  $L^2(\Omega_s^q)$  where  $N^{\Omega_s^q}$  is the  $\bar{\partial}$ -Neumann operator on  $L_{(0,1)}^2(\Omega_s^q)$ . By the proof of Proposition 8 and the fact that  $\bar{\partial}^* N^{\Omega_s^q} F_j$  is holomorphic in  $w$ , we conclude that  $\bar{\partial}^* N^{\Omega_s^q} F_j(\cdot, 0) \in L^2(\Omega, (-\rho_1)^r)$ . Further,  $\bar{\partial} \bar{\partial}^* N^{\Omega_s^q} F_j(\cdot, 0) = f_j$  for all  $j$  and  $\{\bar{\partial}^* N^{\Omega_s^q} F_j(\cdot, 0)\}$  has a convergent subsequence in  $L^2(\Omega, (-\rho_1)^r)$ . Therefore,  $\bar{\partial}$  has a compact solution operator  $R \bar{\partial}^* N^{\Omega_s^q} E$  on  $K_{(0,1)}^2(\Omega, (-\rho_1)^r)$  where  $E$  is the trivial extension operator and  $R$  is the restriction from  $\Omega_s^q$  onto  $\Omega$ .  $\square$

The following lemma is essentially contained in the proof of [Arazy and Engliš 2001, Proposition 1.3]. We present it here for the convenience of the reader.

**Lemma 15.** *Let  $r$  be a nonnegative real number,  $\Omega$  be a bounded domain in  $\mathbb{C}^n$ , and  $\phi \in C(\bar{\Omega})$ . Assume that  $z_0 \in b\Omega$  has a holomorphic peak function. Then*

$$\lim_{z \rightarrow z_0} B_r T_\phi^r(z) = \phi(z_0).$$

*Proof.* First, we prove that for any neighborhood  $U$  of  $z_0$ ,

$$(5) \quad \int_{\Omega \setminus U} |k_z^r(w)|^2 (-\rho(w))^r dV(w) \rightarrow 0 \quad \text{as } z \rightarrow z_0.$$

Indeed, for given  $U$  and  $\varepsilon > 0$  first we choose a holomorphic peak function  $g$  such that  $|g(w)| \leq \varepsilon$  for all  $w \in \Omega \setminus U$ . This can be simply done by taking a high enough power of the holomorphic peak function  $g$ . Then we choose  $\delta > 0$  such that if  $|z - z_0| < \delta$  and  $z \in \Omega$  then  $|g(z)| > 1 - \varepsilon$ . In this case,

$$\begin{aligned} \int_U |k_z^r(w)|^2 (-\rho(w))^r dV(w) &\geq \int_U |g(w)| |k_z^r(w)|^2 (-\rho(w))^r dV(w) \\ &\geq \left| \int_\Omega g(w) |k_z^r(w)|^2 (-\rho(w))^r dV(w) \right| \\ &\quad - \left| \int_{\Omega \setminus U} g(w) |k_z^r(w)|^2 (-\rho(w))^r dV(w) \right| \\ &\geq |g(z)| - \int_{\Omega \setminus U} |g(w)| |k_z^r(w)|^2 (-\rho(w))^r dV(w) \\ &\geq 1 - \varepsilon - \varepsilon \int_{\Omega \setminus U} |k_z^r(w)|^2 (-\rho(w))^r dV(w) \\ &\geq 1 - 2\varepsilon, \end{aligned}$$

whenever  $|z - z_0| < \delta$ . This implies that for a given neighborhood  $U$  and  $\varepsilon > 0$ , there exists  $\delta > 0$  such that if  $|z - z_0| < \delta$  then

$$\int_{\Omega \setminus U} |k_z^r(w)|^2 (-\rho(w))^r dV(w) \leq \varepsilon.$$

This gives (5).

Now for  $\varepsilon > 0$ , we choose a neighborhood  $U$  of  $z$  such that  $|\phi(w) - \phi(z_0)| \leq \varepsilon$  for all  $w \in U$ . Then for this neighborhood  $U$  and the same  $\varepsilon$  we choose  $\delta > 0$  such that if  $|z - z_0| < \delta$  then  $\int_{\Omega \setminus U} |k_z^r(w)|^2 (-\rho(w))^r dV(w) \leq \varepsilon / (1 + 2\|\phi\|_{L^\infty})$ . In this case,

$$\begin{aligned} |B_r T_\phi^r(z) - \phi(z_0)| &\leq \int_{\Omega} |\phi(w) - \phi(z_0)| |k_z^r(w)|^2 (-\rho(w))^r dV(w) \\ &= \int_U |\phi(w) - \phi(z_0)| |k_z^r(w)|^2 (-\rho(w))^r dV(w) \\ &\quad + \int_{\Omega \setminus U} |\phi(w) - \phi(z_0)| |k_z^r(w)|^2 (-\rho(w))^r dV(w) \\ &\leq \varepsilon \int_U |k_z^r(w)|^2 (-\rho(w))^r dV(w) \\ &\quad + 2\|\phi\|_{L^\infty} \int_{\Omega \setminus U} |k_z^r(w)|^2 (-\rho(w))^r dV(w) \\ &\leq \varepsilon + \varepsilon = 2\varepsilon. \end{aligned}$$

This indeed concludes  $\lim_{z \rightarrow z_0} B_r T_\phi^r(z) = \phi(z_0)$ .  $\square$

We note that on any bounded domain, we have (see [Čučković and Şahutoğlu 2014, Lemma 1])

$$T_{\phi_2}^r T_{\phi_1}^r = T_{\phi_2 \phi_1}^r - H_{\phi_2}^{r*} H_{\phi_1}^r.$$

Using the fact above inductively one can prove the following lemma.

**Lemma 16.** *Let  $r$  be a nonnegative real number and  $\Omega$  be a  $C^1$ -smooth bounded domain in  $\mathbb{C}^n$  with a defining function  $\rho$ . Suppose  $\phi_1, \dots, \phi_m \in L^\infty(\Omega)$ . Then*

$$\begin{aligned} T_{\phi_m}^r T_{\phi_{m-1}}^r \cdots T_{\phi_2}^r T_{\phi_1}^r &= T_{\phi_m \phi_{m-1} \cdots \phi_2 \phi_1}^r - T_{\phi_m}^r T_{\phi_{m-1}}^r \cdots T_{\phi_3}^r H_{\phi_2}^{r*} H_{\phi_1}^r \\ &\quad - T_{\phi_m}^r T_{\phi_{m-1}}^r \cdots T_{\phi_4}^r H_{\phi_3}^{r*} H_{\phi_2 \phi_1}^r - \cdots - H_{\phi_m}^{r*} H_{\phi_{m-1} \cdots \phi_2 \phi_1}^r \\ &= T_{\phi_m \phi_{m-1} \cdots \phi_2 \phi_1}^r + S^r, \end{aligned}$$

where  $S^r$  is a finite sum of finite products of operators and each product starts with a Hankel operator.

Therefore, if the symbols  $\phi_1, \dots, \phi_m$  are continuous on  $\bar{\Omega}$  we can write

$$(6) \quad T_\phi^r \cdots T_{\phi_m}^r = T_{\phi_1 \cdots \phi_m}^r + S^r,$$

where  $S^r$  is a finite sum of finite products of operators such that each product starts with a Hankel operator with symbol continuous on  $\bar{\Omega}$ .

We state the lemma below for general weights  $\mu(z)$  (not only the ones of the form  $(-\rho)^k$ ) that are nonnegative (can vanish on the boundary) and continuous on  $\Omega$ . The weights of this form are called admissible weights (see [Pasternak-Winiarski 1990]) and the corresponding weighted Bergman projections and kernels are well defined. We say two weights  $\mu_1$  and  $\mu_2$  are comparable if there exists  $c > 0$  such that  $c^{-1}\mu_1 < \mu_2 < c\mu_1$  on  $\Omega$ .

**Lemma 17.** *Let  $\Omega$  be a domain in  $\mathbb{C}^n$  and  $\mu_1$  and  $\mu_2$  be comparable admissible weights. Let  $k_z^{\mu_j}$  be the normalized Bergman kernel corresponding to  $\mu_j$  for  $j = 1, 2$ , and  $z_0 \in b\Omega$ . Then  $k_z^{\mu_1} \rightarrow 0$  weakly as  $z \rightarrow z_0$  if and only if  $k_z^{\mu_2} \rightarrow 0$  weakly as  $z \rightarrow z_0$ .*

*Proof.* It is enough to show one direction. So we will show that if  $k_z^{\mu_1} \rightarrow 0$  weakly as  $z \rightarrow z_0$  then  $k_z^{\mu_2} \rightarrow 0$  weakly as  $z \rightarrow z_0$ . Since  $\mu_1$  and  $\mu_2$  are equivalent measures we have  $A^2(\Omega, d\mu_1) = A^2(\Omega, d\mu_2)$  and there exists  $C > 1$  such that

$$\frac{\|f\|_{\mu_1}}{C} \leq \|f\|_{\mu_2} \leq C\|f\|_{\mu_1}$$

for all  $f \in A^2(\Omega, d\mu_1)$ . We remind the reader that for  $z \in \Omega$  we have

$$K_{\mu_j}(z, z) = \sup\{|f(z)|^2 : \|f\|_{\mu_j} \leq 1\},$$

where  $K_{\mu_j}$  is the Bergman kernel corresponding to  $\mu_j$ . Then  $K_{\mu_1}$  and  $K_{\mu_2}$  are equivalent on the diagonal in the sense that there exists  $D = C^2 > 1$  such that

$$\frac{K_{\mu_1}(z, z)}{D} \leq K_{\mu_2}(z, z) \leq DK_{\mu_1}(z, z).$$

Now we assume that  $k_z^{\mu_1} \rightarrow 0$  weakly as  $z \rightarrow z_0$ . Let us fix  $f \in A^2(\Omega, d\mu_1)$ . Then we have

$$\frac{f(z)}{\sqrt{K_{\mu_1}(z, z)}} = \langle f, k_z^{\mu_1} \rangle_{\mu_1} \rightarrow 0 \quad \text{as } z \rightarrow z_0.$$

Then

$$\langle f, k_z^{\mu_2} \rangle_{\mu_2} = \frac{f(z)}{\sqrt{K_{\mu_2}(z, z)}} \rightarrow 0 \quad \text{as } z \rightarrow z_0.$$

Therefore, we showed that if  $k_z^{\mu_1} \rightarrow 0$  weakly as  $z \rightarrow z_0$  then  $k_z^{\mu_2} \rightarrow 0$  weakly as  $z \rightarrow z_0$ .  $\square$

Let  $\Omega$  be a pseudoconvex domain in  $\mathbb{C}^n$  and  $z_0 \in b\Omega$ . Then we call  $z_0$  a *bumping point* if for any  $\delta > 0$  there exists a pseudoconvex domain  $\Omega_1$  such that  $\{z_0\} \cup \Omega \subset \Omega_1 \subset \Omega \cup B(z_0, \delta)$ .

**Lemma 18.** *Let  $r$  be a nonnegative real number,  $\Omega$  be a bounded pseudoconvex domain in  $\mathbb{C}^n$  with Lipschitz boundary, and  $z_0 \in b\Omega$  be a bumping point. Then  $k_z^r \rightarrow 0$  weakly as  $z \rightarrow z_0$ .*

*Proof.* By Lemma 17, without loss of generality, we assume that  $\rho$  denotes the negative distance to the boundary of  $\Omega$ .

Let us fix  $f \in A^2(\Omega, (-\rho)^r)$  and choose  $r_1, r_2 > 0$  so that  $0 < r_1 < r_2$  and the outward unit vector  $\nu$  is transversal to  $B(z_0, 2r_2) \cap b\Omega$ . Since  $z_0$  is a bumping point we choose a bounded pseudoconvex domain  $\Omega_1$  such that

$$\{z_0\} \cup \Omega \subset \Omega_1 \subset \Omega \cup B(z_0, r_1).$$

So even though  $\Omega_1$  contains a small neighborhood of  $z_0$ , we have

$$\Omega \setminus B(z_0, r_1) = \Omega_1 \setminus B(z_0, r_1).$$

Let us choose  $\chi \in C_0^\infty(B(z_0, r_2))$  such that  $\chi \equiv 1$  on a neighborhood of  $\overline{B(z_0, r_1)}$ . For  $\varepsilon > 0$  small we define  $f_\varepsilon(z) = f(z - \varepsilon\nu)$  and  $g_\varepsilon = (1 - \chi)f + \chi f_\varepsilon$ . Then

- (i)  $f_\varepsilon \in A^2(\Omega \cap B(z_0, r_2), (-\rho)^r)$  and  $f_\varepsilon \rightarrow f$  in  $L^2(\Omega \cap B(z_0, r_2), (-\rho)^r)$ ,
- (ii)  $g_\varepsilon|_{\Omega \cap B(z_0, r_2)}$  is  $C^\infty$ -smooth and  $g_\varepsilon \rightarrow f$  in  $L^2(\Omega, (-\rho)^r)$  as  $\varepsilon \rightarrow 0$ .

Let  $\rho_1$  and  $\text{Supp}(\bar{\partial}\chi)$  denote the negative distance to the boundary of  $\Omega_1$  and the support of  $\bar{\partial}\chi$ , respectively. Then  $\text{Supp}(\bar{\partial}\chi) \cap \Omega = \text{Supp}(\bar{\partial}\chi) \cap \Omega_1$  and  $-\rho$  and  $-\rho_1$  are equivalent on the support of  $\bar{\partial}\chi$ . We note that  $g_\varepsilon$  is well defined on  $\Omega$  and not on  $\Omega_1$ . However,  $\bar{\partial}g_\varepsilon = 0$  on  $\Omega \cap B(z_0, r_1)$  as  $\bar{\partial}\chi = 0$  on  $B(z_0, r_1)$  for all small  $\varepsilon > 0$ . Hence  $\bar{\partial}g_\varepsilon$  can be extended trivially to be defined on  $\Omega_1$  as a  $\bar{\partial}$ -closed  $(0, 1)$ -form on  $\Omega_1$ . Then there exists  $C > 0$  such that

$$\|\bar{\partial}g_\varepsilon\|_{L^2(\Omega_1, (-\rho_1)^r)} \leq C \|f - f_\varepsilon\|_{L^2(\Omega \cap B(z_0, r_2), (-\rho)^r)} \|\bar{\partial}\chi\|_{L^\infty(B(z_0, r_2))} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

Next we will use Hörmander's theorem [1965] with the plurisubharmonic exponential weight  $-r \log(-\rho_1)$ . We note that  $-\log(-\rho_1)$  is plurisubharmonic because  $\Omega_1$  is pseudoconvex. Then using Hörmander's theorem we get a constant  $c_{\Omega_1} > 0$  (depending on  $\Omega_1$ ) and  $h_\varepsilon \in L^2(\Omega_1)$  such that  $\bar{\partial}h_\varepsilon = \bar{\partial}g_\varepsilon$  and

$$\|h_\varepsilon\|_{L^2(\Omega_1, (-\rho_1)^r)} \leq c_{\Omega_1} \|\bar{\partial}g_\varepsilon\|_{L^2(\Omega_1, (-\rho_1)^r)}.$$

Furthermore, since  $\bar{\partial}$  is elliptic on the interior and  $\bar{\partial}g_\varepsilon$  is  $C^\infty$ -smooth on  $\Omega_1$ , we have  $h_\varepsilon \in C^\infty(\Omega_1)$ .

We define  $\tilde{f}_n = g_{1/n} - h_{1/n}$  on  $\Omega$ . We note that while  $\bar{\partial}g_\varepsilon$  is even defined on all of  $\Omega_1$ ,  $g_\varepsilon$  and, hence,  $\tilde{f}_{1/n}$  in general, are not. Then we have

- (i)  $\tilde{f}_n \in A^2(\Omega, (-\rho)^r)$  and  $\tilde{f}_n \rightarrow f$  in  $A^2(\Omega, (-\rho)^r)$ ,
- (ii)  $\tilde{f}_n|_{\Omega \cap B(z_0, r_1)} \in C^\infty(\overline{\Omega \cap B(z_0, r_1)})$ .



So  $\{\tilde{f}_n\}$  is a sequence converging to  $f$  and each member of the sequence is smooth up to the boundary of  $\Omega$  on a neighborhood of  $z_0$ .

Finally, we will show weak convergence of  $k_z^r$  to 0 as  $z \rightarrow z_0$ .

$$\begin{aligned} |\langle f, k_z^r \rangle_{A^2(\Omega, (-\rho)^r)}| &\leq |\langle f - \tilde{f}_n, k_z^r \rangle_{A^2(\Omega, (-\rho)^r)}| + |\langle \tilde{f}_n, k_z^r \rangle_{A^2(\Omega, (-\rho)^r)}| \\ &\leq \|f - \tilde{f}_n\|_{L^2(\Omega, (-\rho)^r)} + \frac{|\tilde{f}_n(z)|}{\sqrt{K_r(z, z)}}. \end{aligned}$$

The first term on the right-hand side can be made arbitrarily small for large enough  $n$ , because  $\|f - \tilde{f}_n\|_{L^2(\Omega, (-\rho)^r)} \rightarrow 0$  as  $n \rightarrow \infty$ . So for  $\delta > 0$  given we choose  $n_\delta$  so that  $\|f - \tilde{f}_{n_\delta}\|_{L^2(\Omega, (-\rho)^r)} \leq \delta$ . Then since  $\tilde{f}_{n_\delta}$  is  $C^\infty$ -smooth on  $\overline{\Omega} \cap B(z_0, r_1)$  (and  $K_r(z, z) \rightarrow \infty$  as  $z \rightarrow z_0$ ), we conclude that  $|\tilde{f}_{n_\delta}(z)|/\sqrt{K_r(z, z)} \rightarrow 0$  as  $z \rightarrow z_0$ . Hence,  $\limsup_{z \rightarrow z_0} |\langle f, k_z^r \rangle| \leq \delta$  for arbitrary  $\delta > 0$ . Therefore,  $k_z^r \rightarrow 0$  weakly as  $z \rightarrow z_0$ .  $\square$

Now we are ready to prove Theorem 4.

*Proof of Theorem 4.* In the case  $r = 0$ , the proof of the theorem simplifies greatly as inflation and the related techniques are unnecessary. So we will prove the more difficult case,  $r > 0$ .

First we assume that  $T$  is compact about strongly pseudoconvex points. Let  $\Omega_r^p$  be defined as in (1) and  $z_0 \in b\Omega$  be a strongly pseudoconvex point. Since small  $C^2$ -perturbations of strongly pseudoconvex points stay pseudoconvex,  $z_0$  is a bumping point for  $\Omega$ . Then Lemma 18 implies that  $k_z^r \rightarrow 0$  weakly as  $z \rightarrow z_0$ . Furthermore, there exists an open neighborhood  $U$  of  $z_0$  such that weakly pseudoconvex points are contained in  $b\Omega \setminus \overline{U}$ ; and, as in the proof of (5), one can show that

$$\|k_z^r\|_{L^2(\Omega \setminus \overline{U}, (-\rho)^r)} \rightarrow 0 \text{ as } z \rightarrow z_0.$$

Therefore,  $\{k_z^r\}$  converges to 0 weakly about strongly pseudoconvex points as  $z \rightarrow z_0$ . Moreover, since  $T$  is compact about strongly pseudoconvex points (such operators map sequences of holomorphic functions weakly convergent about strongly pseudoconvex points to convergent sequences) we conclude that

$$B_r T(z) = \langle T k_z^r, k_z^r \rangle_{A^2(\Omega, (-\rho)^r)} \rightarrow 0$$

as  $z \rightarrow z_0$ .

Next we prove the other direction. As a first step we assume that  $T$  is a finite sum of finite products of Toeplitz operators on  $A^2(\Omega, (-\rho)^r)$  with symbols continuous on  $\overline{\Omega}$ . Furthermore, we assume that

$$\lim_{z \rightarrow z_0} B_r T(z) = 0$$

for any strongly pseudoconvex point  $z_0 \in b\Omega$ .

Lemma 16 implies that

$$(7) \quad T = T_\phi^r + S^r,$$

where  $\phi \in C(\overline{\Omega})$  and  $S^r$  is a sum of operators that start with a Hankel operator with symbol continuous on  $\overline{\Omega}$ .

Lemma 15 implies that

$$(8) \quad \lim_{z \rightarrow z_0} B_r T_\phi^r(z) = \phi(z_0)$$

as strongly pseudoconvex points have holomorphic peak functions (see [Range 1986, Theorem 1.13 in Ch VI]).

By Proposition 13, the operator  $H_\psi^r$  is compact about strongly pseudoconvex points for any  $\psi \in C(\overline{\Omega})$ . Then  $H_\psi^r k_z^r \rightarrow 0$  as  $z \rightarrow z_0$  for any  $\psi \in C(\overline{\Omega})$  because, as proven in the first part of this proof,  $k_z^r \rightarrow 0$  weakly about strongly pseudoconvex points as  $z \rightarrow z_0$ . Hence,  $B_r S^r(z) \rightarrow 0$  as  $z \rightarrow z_0$ . Combining this with (7) and (8) we can conclude that

$$\phi(z_0) = \lim_{z \rightarrow z_0} B_r T(z) = 0.$$

Since  $z_0$  was an arbitrary strongly pseudoconvex point, we have  $\phi = 0$  on all the strongly pseudoconvex boundary points. Then Lemma 12 and the fact that  $S^r$  is compact about strongly pseudoconvex points imply that  $T$  is compact about strongly pseudoconvex points.

Finally, we assume  $T \in \mathcal{T}(\overline{\Omega}, (-\rho)^r)$ . Then, using Lemma 16, for every  $\varepsilon > 0$  there exists  $\phi_\varepsilon \in C(\overline{\Omega})$  and an operator  $S_\varepsilon^r$ , compact about strongly pseudoconvex points, such that

$$\|T + T_{\phi_\varepsilon}^r + S_\varepsilon^r\| \leq \varepsilon.$$

Then for  $z \in \Omega$  we have

$$\begin{aligned} |B_r T(z) + B_r T_{\phi_\varepsilon}^r(z) + B_r S_\varepsilon^r(z)| &= |\langle T k_z^r + T_{\phi_\varepsilon}^r k_z^r + S_\varepsilon^r k_z^r, k_z^r \rangle_r| \\ &\leq \|T + T_{\phi_\varepsilon}^r + S_\varepsilon^r\| \\ &\leq \varepsilon. \end{aligned}$$

Since  $B_r S_\varepsilon^r(z) \rightarrow 0$  and  $B_r T_{\phi_\varepsilon}^r(z) \rightarrow \phi_\varepsilon(z_0)$  as  $z \rightarrow z_0$  (and we assume that  $B_r T(z) \rightarrow 0$  as  $z \rightarrow z_0$ ), we have  $|\phi_\varepsilon(z_0)| \leq \varepsilon$ . That is,  $|\phi_\varepsilon|$  is less than or equal to  $\varepsilon$  on strongly pseudoconvex points of  $\overline{\Omega}$ . We choose  $\psi_\varepsilon \in C(\overline{\Omega})$  such that  $\psi_\varepsilon = 0$  on strongly pseudoconvex boundary points of  $\Omega$  and

$$\sup\{|\psi_\varepsilon(z) - \phi_\varepsilon(z)| : z \in \overline{\Omega}\} \leq 2\varepsilon.$$

Then Lemma 12 implies that  $T_{\psi_\varepsilon}^r$  is compact about strongly pseudoconvex points and

$$\|T_{\phi_\varepsilon}^r - T_{\psi_\varepsilon}^r\| \leq 2\varepsilon.$$

Hence

$$\|T + T_{\psi_\varepsilon}^r + S_\varepsilon^r\| \leq \|T + T_{\phi_\varepsilon}^r + S_\varepsilon^r\| + \|T_{\psi_\varepsilon}^r - T_{\phi_\varepsilon}^r\| \leq 3\varepsilon.$$

Therefore,  $T$  is in the norm closure of the compact about strongly pseudoconvex points operators. Finally, Proposition 7 implies that  $T$  is compact about strongly pseudoconvex points.  $\square$

### Acknowledgements

Part of this work was done while Şahutoğlu was visiting Sabancı University. He thanks this institution for its hospitality and good working environment. We would like to thank the anonymous referee for pointing out and helping to fix some inaccuracies.

### References

- [Arazy and Engliš 2001] J. Arazy and M. Engliš, “Iterates and the boundary behavior of the Berezin transform”, *Ann. Inst. Fourier (Grenoble)* **51**:4 (2001), 1101–1133. MR Zbl
- [Axler and Zheng 1998] S. Axler and D. Zheng, “Compact operators via the Berezin transform”, *Indiana Univ. Math. J.* **47**:2 (1998), 387–400. MR Zbl
- [Catlin 1984] D. W. Catlin, “Global regularity of the  $\bar{\partial}$ -Neumann problem”, pp. 39–49 in *Complex analysis of several variables* (Madison, Wis., 1982), edited by Y.-T. Siu, Proc. Sympos. Pure Math. **41**, Amer. Math. Soc., Providence, RI, 1984. MR Zbl
- [Čučković and Şahutoğlu 2013] Ž. Čučković and S. Şahutoğlu, “Axler–Zheng type theorem on a class of domains in  $\mathbb{C}^n$ ”, *Integral Equations Operator Theory* **77**:3 (2013), 397–405. MR
- [Čučković and Şahutoğlu 2014] Ž. Čučković and S. Şahutoğlu, “Compactness of products of Hankel operators on convex Reinhardt domains in  $\mathbb{C}^2$ ”, *New York J. Math.* **20** (2014), 627–643. MR
- [Diederich and Fornæss 1977] K. Diederich and J. E. Fornæss, “Pseudoconvex domains: bounded strictly plurisubharmonic exhaustion functions”, *Invent. Math.* **39**:2 (1977), 129–141. MR Zbl
- [Engliš 1999] M. Engliš, “Compact Toeplitz operators via the Berezin transform on bounded symmetric domains”, *Integral Equations Operator Theory* **33**:4 (1999), 426–455. MR Zbl
- [Forelli and Rudin 1974] F. Forelli and W. Rudin, “Projections on spaces of holomorphic functions in balls”, *Indiana Univ. Math. J.* **24** (1974), 593–602. MR Zbl
- [Hörmander 1965] L. Hörmander, “ $L^2$  estimates and existence theorems for the  $\bar{\partial}$  operator”, *Acta Math.* **113** (1965), 89–152. MR
- [Kreutzer 2014] E. Kreutzer, *Toeplitz extensions and Berezin transforms*, master’s thesis, Universität des Saarlandes, 2014, Available at <https://www.math.uni-sb.de/ag/eschmeier/old/examen/>.
- [Ligocka 1989] E. Ligocka, “On the Forelli–Rudin construction and weighted Bergman projections”, *Studia Math.* **94**:3 (1989), 257–272. MR Zbl
- [Mitkovski et al. 2013] M. Mitkovski, D. Suárez, and B. D. Wick, “The essential norm of operators on  $A_\alpha^p(\mathbb{B}_n)$ ”, *Integral Equations Operator Theory* **75**:2 (2013), 197–233. MR Zbl
- [Pasternak-Winiarski 1990] Z. Pasternak-Winiarski, “On the dependence of the reproducing kernel on the weight of integration”, *J. Funct. Anal.* **94**:1 (1990), 110–134. MR Zbl
- [Range 1986] R. M. Range, *Holomorphic functions and integral representations in several complex variables*, Graduate Texts in Mathematics **108**, Springer, 1986. MR Zbl

[Sibony 1987] N. Sibony, “Une classe de domaines pseudoconvexes”, *Duke Math. J.* **55**:2 (1987), 299–319. MR Zbl

[Straube 2010] E. J. Straube, *Lectures on the  $\mathcal{L}^2$ -Sobolev theory of the  $\bar{\partial}$ -Neumann problem*, ESI Lectures in Mathematics and Physics **7**, European Mathematical Society (EMS), Zürich, 2010. MR Zbl

[Suárez 2007] D. Suárez, “The essential norm of operators in the Toeplitz algebra on  $A^p(\mathbb{B}_n)$ ”, *Indiana Univ. Math. J.* **56**:5 (2007), 2185–2232. MR Zbl

Received April 23, 2017. Revised September 25, 2017.

ŽELJKO ČUČKOVIĆ  
DEPARTMENT OF MATHEMATICS AND STATISTICS  
UNIVERSITY OF TOLEDO  
TOLEDO, OH  
UNITED STATES  
zeljko.cuckovic@utoledo.edu

SÖNMEZ ŞAHUTOĞLU  
DEPARTMENT OF MATHEMATICS AND STATISTICS  
UNIVERSITY OF TOLEDO  
TOLEDO, OH  
UNITED STATES

*Current address:*  
SABANCI UNIVERSITY  
TUZLA  
ISTANBUL  
TURKEY  
sonmez.sahutoglu@utoledo.edu

YUNUS E. ZEYTUNCU  
DEPARTMENT OF MATHEMATICS AND STATISTICS  
UNIVERSITY OF MICHIGAN-DEARBORN  
DEARBORN, MI  
UNITED STATES  
zeytuncu@umich.edu

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Wee Teck Gan  
Mathematics Department  
National University of Singapore  
Singapore 119076  
[matgwt@nus.edu.sg](mailto:matgwt@nus.edu.sg)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 294 No. 1 May 2018

---

Three-dimensional Sol manifolds and complex Kleinian groups	1
WALDEMAR BARRERA, RENE GARCIA-LARA and JUAN NAVARRETE	
On periodic points of symplectomorphisms on surfaces	19
MARTA BATORÉO	
Mixing properties for hom-shifts and the distance between walks on associated graphs	41
NISHANT CHANDGOTIA and BRIAN MARCUS	
Simultaneous construction of hyperbolic isometries	71
MATT CLAY and CAGLAR UYANIK	
A local weighted Axler–Zheng theorem in $\mathbb{C}^n$	89
ŽELJKO ČUČKOVIĆ, SÖNMEZ ŞAHUTOĞLU and YUNUS E. ZEYTUNCU	
Monotonicity and radial symmetry results for Schrödinger systems with fractional diffusion	107
JING LI	
Moduli spaces of stable pairs	123
YINBANG LIN	
Spark deficient Gabor frames	159
ROMANOS-DIOGENES MALIKIOSIS	
Ordered groups as a tensor category	181
DALE ROLFSEN	
Multiplication of distributions and a nonlinear model in elastodynamics	195
C. O. R. SARRICO	
Some Ambrose- and Galloway-type theorems via Bakry–Émery and modified Ricci curvatures	213
HOMARE TADANO	
Irreducible decomposition for local representations of quantum Teichmüller space	233
JÉRÉMY TOULISSE	



0030-8730(201805)294:1;1-T