

*Pacific
Journal of
Mathematics*

**MONOTONICITY AND RADIAL SYMMETRY RESULTS FOR
SCHRÖDINGER SYSTEMS WITH FRACTIONAL DIFFUSION**

JING LI

Volume 294 No. 1

May 2018

MONOTONICITY AND RADIAL SYMMETRY RESULTS FOR SCHRÖDINGER SYSTEMS WITH FRACTIONAL DIFFUSION

JING LI

We consider a nonlinear Schrödinger system with fractional diffusion

$$\begin{cases} (-\Delta)^{\alpha/2}u(x) + A(x)u(x) = v^p(x) & \text{in } \Omega, \\ (-\Delta)^{\beta/2}v(x) + B(x)v(x) = u^q(x) & \text{in } \Omega, \\ u(x) \geq 0, v(x) \geq 0 & \text{in } \Omega, \\ u(x) = v(x) = 0 & \text{on } \Omega^C, \end{cases}$$

where Ω is an unbounded parabolic domain. We first establish a narrow region principle. Using this principle and a direct method of moving planes, we obtain the monotonicity of nonnegative solutions and the Liouville-type result for the nonlinear Schrödinger system with fractional diffusion. We also obtain the radially symmetric result of positive solutions for the system in the unit ball when $A(x)$ and $B(x)$ are constants.

1. Introduction

We are interested in the following nonlinear Schrödinger system with fractional diffusion:

$$(1-1) \quad \begin{cases} (-\Delta)^{\alpha/2}u(x) + A(x)u(x) = v^p(x) & \text{in } \Omega, \\ (-\Delta)^{\beta/2}v(x) + B(x)v(x) = u^q(x) & \text{in } \Omega, \\ u(x) \geq 0, v(x) \geq 0 & \text{in } \Omega, \\ u(x) = v(x) = 0 & \text{on } \Omega^C, \end{cases}$$

where $\alpha, \beta \in (0, 2)$, $p, q > 1$, $A(x)$ and $B(x)$ are bounded from below and Ω is an unbounded parabolic domain in \mathbb{R}^n defined by

$$\Omega = \{x = (x', x_n) \in \mathbb{R}^n \mid x_n > |x'|^2, x' = (x_1, x_2, \dots, x_{n-1})\}.$$

The research is supported by the National Natural Science Foundation of China (No. 11271111 and No. 11601131).

MSC2010: 35J60.

Keywords: fractional Schrödinger system, narrow region principle, direct method of moving planes, monotonicity, radially symmetric.

Here, $(-\Delta)^{\alpha/2}$ and $(-\Delta)^{\beta/2}$ are nonlocal pseudodifferential operators defined by

$$(1-2) \quad (-\Delta)^{\alpha/2}u(x) = C_{n,\alpha} P.V. \int_{\mathbb{R}^n} \frac{u(x) - u(y)}{|x - y|^{n+\alpha}} dy,$$

$$(1-3) \quad (-\Delta)^{\beta/2}v(x) = C_{n,\beta} P.V. \int_{\mathbb{R}^n} \frac{v(x) - v(y)}{|x - y|^{n+\beta}} dy,$$

where $P.V.$ stands for the Cauchy principal value, and $C_{n,\alpha}$, $C_{n,\beta}$ are normalization positive constants. Let

$$F = L_\alpha \cap C_{loc}^{1,1}(\Omega), \quad G = L_\beta \cap C_{loc}^{1,1}(\Omega),$$

where

$$L_\alpha = \left\{ u \mid u \in L_{loc}^1, \int_{\mathbb{R}^n} \frac{|u(x)|}{1 + |x|^{n+\alpha}} dx < \infty \right\},$$

and

$$L_\beta = \left\{ v \mid v \in L_{loc}^1, \int_{\mathbb{R}^n} \frac{|v(x)|}{1 + |x|^{n+\beta}} dx < \infty \right\}.$$

For $u \in F$, $v \in G$, the integral on the left-hand side of the equations in (1-1) is well defined (see [Chen et al. 2017b]).

Linear and nonlinear equations and systems involving the fractional Laplacian have received growing attention in recent years. It can be used to model diverse physical phenomena, such as turbulence and water waves, molecular dynamics, and pseudorelativistic boson stars (see [Bouchaud and Georges 1990], [Caffarelli and Vasseur 2010], [Constantin 2006], [Tarasov and Zaslavsky 2006]). The operator $(-\Delta)^{\alpha/2}$ can also be used in mathematical finance (see [Applebaum 2009], [Bertoin 1996]). But they are still much less understood than nonfractional counterparts.

When $\alpha = \beta = 2$, $A(x) = B(x) = 0$, (1-1) becomes the classical Lane–Emden system:

$$(1-4) \quad \begin{cases} -\Delta u = v^p, \\ -\Delta v = u^q. \end{cases}$$

When $1/(p+1) + 1/(q+1) > (n-2)/n$, the system (1-4) has no positive radial solutions in all dimension (see [Mitidieri 1996]). D. G. de Figueiredo and P. L. Felmer [1994] studied a Liouville type theorem for (1-4) by introducing superharmonic functions when $n \geq 3$. The main tool they used is the method of moving planes. For $n = 3$, J. Serrin and H. Zou [1996] proved that the system (1-4) has no positive solutions when $1/(p+1) + 1/(q+1) > (n-2)/n$ under assumption that (u, v) has at most polynomial growth at infinity. After Serrin's work, there are some interesting works about Lane–Emden systems and related Schrödinger systems on whole space and half space; see [Montaru and Souplet 2014; Poláčik et al. 2007; Souplet 2009].

For classical semilinear elliptic system, the symmetry and monotonicity of positive solutions have been widely studied (see [Busca and Sirakov 2000; Chen and Li 2010; Liu and Ma 2012; 2013; Ma and Liu 2010]). A powerful tool to obtain these properties of such equations and systems is the method of moving planes, which was introduced by Alexandrov [1962]. Serrin [1971] and Gidas, Ni, Nirenberg [Gidas et al. 1979; 1981] adapted this method in partial differential equations and made great contributions to improving this method.

As we know, the fractional Laplacian is nonlocal; that is, it is not differentiable pointwise, but is globally integrable with respect to a singular kernel. The nonlocality causes the main difficulty in studying corresponding problems. To circumvent this difficulty, Caffarelli and Silvestre [2007] introduced the extension method that reduced this nonlocal problem in \mathbb{R}^n into a local one in \mathbb{R}_+^{n+1} through constructing a Dirichlet to Neumann operator of a degenerate elliptic equation. This extension method has been applied successfully to study equations involving the fractional Laplacian, and a series of fruitful results have been obtained; see [Brändle et al. 2013; Chen and Zhu 2016].

Due to technical restrictions, they have to assume $\alpha \geq 1$. It seems that this condition cannot be weakened if one wants to carry out the method of moving planes on the extended equation. Actually, the case $0 < \alpha < 1$ can be treated by considering the corresponding integral equation. Using the method of moving planes (or spheres) in integral forms [Chen and Li 2009; Chen et al. 2005a; 2005b; 2006; 2015; Fall and Weth 2016; Fang and Zhang 2013; Li and Ma 2008; Ma and Chen 2008; Ma and Zhao 2008, one can obtain the radial symmetry properties of the fractional Laplacian equation. For the fractional Laplacian system, here we mention the work by Zhuo, Chen, Cui and Yuan [Zhuo et al. 2016]. They considered the system

$$(1-5) \quad (-\Delta)^{\alpha/2} u_i(x) = f_i(u_1(x), u_2(x), \dots, u_m(x)), \quad x \in \mathbb{R}^n, \quad i = 1, 2, \dots, m.$$

By establishing the equivalence between (1-5) and its corresponding integral system, the authors obtained the symmetry result and the nonexistence of positive solutions.

Either by extension or by integral equations, one needs to impose extra conditions on the solutions. Can one carry out the method of moving planes directly on fractional equation? The answer was provided in [Jarohs and Weth 2016] by Jarohs and Weth. They introduced antisymmetric maximum principles and applied them to carry out the method of moving planes directly on nonlocal problems to show the symmetry of solutions. However, their maximum principles only apply to bounded regions.

Recently, Chen, Li and Li [Chen et al. 2017b] developed a direct method of moving planes to study the fractional Laplacian, which worked directly on the nonlocal operator. The key ingredients of this method are the antisymmetric properties. They used this property to develop some techniques needed in the direct method of moving planes in the whole space \mathbb{R}^n and the upper half space \mathbb{R}_+^n .

such as the narrow region principle, decay at infinity. The direct method of moving planes is very useful. This method has been applied to fully nonlinear fractional order-equations and systems in [Chen et al. 2017a]. In [Cheng et al. 2017], the authors considered the symmetry and monotonicity properties for positive solutions of fractional Laplacian equations by the direct method. Using the spirit of direct method of moving planes in [Chen et al. 2017b], Cai and Mei [2017] studied the fractional Lane–Emden system in \mathbb{R}^n and obtain the symmetry properties and Liouville-type result of positive solutions. Liu and Ma [2016] studied symmetry properties of the general fractional Laplacian system:

$$(1-6) \quad \begin{cases} (-\Delta)^{\alpha/2}u(x) = f(u, v) & \text{in } \mathbb{R}^n, \\ (-\Delta)^{\alpha/2}v(x) = g(u, v) & \text{in } \mathbb{R}^n, \\ u(x) \geq 0, v(x) \geq 0 & \text{in } \mathbb{R}^n \end{cases}$$

under a strong decay condition on the solutions at infinity.

The goal of this paper is to generalize the direct method of moving planes to the Schrödinger system. We first establish the narrow region principle for Schrödinger systems with fractional diffusion. We write $x = (x', x_n) \in \mathbb{R}^n$ with $x' = (x_1, x_2, \dots, x_{n-1})$. Assume $A(x)$ and $B(x)$ are independent of x_n , that is,

$$A(x) = A(x'), \quad B(x) = B(x').$$

Let

$$T_\lambda = \{x \in \mathbb{R}^n \mid x_n = \lambda, \lambda \in \mathbb{R}, \lambda > 0\}$$

be the moving plane and denote

$$H_\lambda = \{x \in \mathbb{R}^n \mid x_n < \lambda\}, \quad \Sigma_\lambda = \{x \in \Omega \mid 0 < x_n < \lambda\}.$$

For each point $x = (x', x_n) \in \Sigma_\lambda$, let $x^\lambda = (x', 2\lambda - x_n)$ be the reflection point about the plane T_λ . Denote

$$U_\lambda(x) = u(x^\lambda) - u(x) = u_\lambda(x) - u(x), \quad V_\lambda(x) = v(x^\lambda) - v(x) = v_\lambda(x) - v(x).$$

It follows that for $x \in \Sigma_\lambda$,

$$(1-7) \quad \begin{aligned} (-\Delta)^{\alpha/2}U_\lambda(x) &= (-\Delta)^{\alpha/2}u_\lambda(x) - (-\Delta)^{\alpha/2}u(x) \\ &= p\xi^{p-1}(x)V_\lambda(x) - A(x')U_\lambda(x), \end{aligned}$$

and

$$(1-8) \quad \begin{aligned} (-\Delta)^{\beta/2}V_\lambda(x) &= (-\Delta)^{\beta/2}v_\lambda(x) - (-\Delta)^{\beta/2}v(x) \\ &= q\eta^{q-1}(x)U_\lambda(x) - B(x')V_\lambda(x), \end{aligned}$$

where $\xi(x)$ is between $v_\lambda(x)$ and $v(x)$ and $\eta(x)$ is between $u_\lambda(x)$ and $u(x)$. It is obvious that $U_\lambda(x)$ and $V_\lambda(x)$ satisfy the antisymmetry property:

$$(1-9) \quad U_\lambda(x^\lambda) = -U_\lambda(x), \quad V_\lambda(x^\lambda) = -V_\lambda(x), \quad x \in H_\lambda.$$

Lemma 1.1 (narrow region principle). *Let $(u, v) \in F \times G$ be a nonnegative solution of system (1-1) with $0 < \alpha, \beta < 2$. Assume that $1 < p, q < \infty$, $A(x) = A(x')$ and $B(x) = B(x')$ are bounded from below in Ω , where $x = (x', x_n) \in \Omega$, $x' = (x_1, x_2, \dots, x_{n-1})$. Then for all systems (1-7) and (1-8) and for sufficiently small δ ,*

(i) *if there exists $x_1^* \in \Sigma_{\lambda, \delta} = \{x \in \Sigma_\lambda \mid \lambda - \delta < x_n < \lambda\}$ satisfying $U_\lambda(x_1^*) = \min_{x \in \bar{\Sigma}_\lambda} U_\lambda(x) < 0$, then*

$$V_\lambda(x_1^*) < U_\lambda(x_1^*) < 0;$$

(ii) *if there exists $x_2^* \in \Sigma_{\lambda, \delta} = \{x \in \Sigma_\lambda \mid \lambda - \delta < x_n < \lambda\}$ satisfying $V_\lambda(x_2^*) = \min_{x \in \bar{\Sigma}_\lambda} V_\lambda(x) < 0$, then*

$$U_\lambda(x_2^*) < V_\lambda(x_2^*) < 0.$$

Based on Lemma 1.1, we can obtain the following result.

Theorem 1.2. *Let $(u, v) \in F \times G$ be a nonnegative solution of system (1-1) with $0 < \alpha, \beta < 2$. If $1 < p, q < \infty$, $A(x) = A(x')$ and $B(x) = B(x')$ are bounded from below in Ω , where $x = (x', x_n) \in \Omega$, $x' = (x_1, x_2, \dots, x_{n-1})$. Then (u, v) is monotonically increasing in x_n .*

When $\alpha = \beta = 2$, this result is contained in the series papers of Berestycki, Caffarelli and Nirenberg [Berestycki and Nirenberg 1992; Berestycki et al. 1993; 1996; 1997] and they have used the classical method of moving planes. Hence our result by using the direct method of moving planes due to [Chen et al. 2017b] and [Jarohs and Weth 2016] can be considered as an extension of theirs to the nonlocal system.

As an immediate application, we obtain the following Liouville type result.

Corollary 1.3. *Let $(u, v) \in F \times G$ be a nonnegative solution of system (1-1) with $0 < \alpha, \beta < 2$. Assume that $1 < p, q < \infty$, $A(x) = A(x')$ and $B(x) = B(x')$ are bounded from below in Ω , where $x = (x', x_n) \in \Omega$, $x' = (x_1, x_2, \dots, x_{n-1})$. If*

$$(1-10) \quad \lim_{x \rightarrow \infty} u(x) = 0, \quad \lim_{x \rightarrow \infty} v(x) = 0,$$

Then $u \equiv 0, v \equiv 0$.

We consider the system (1-1) when $A(x) = A$ and $B(x) = B$, where A, B are two constants in the unit ball $\mathbb{B}_1(0)$ and obtain the radial symmetry and monotonicity of positive solutions.

Theorem 1.4. *Assume $(u, v) \in L_\alpha \cap C_{loc}^{1,1}(\mathbb{B}_1(0)) \times L_\beta \cap C_{loc}^{1,1}(\mathbb{B}_1(0))$ is a positive solution of the following system*

$$(1-11) \quad \begin{cases} (-\Delta)^{\alpha/2} u(x) + Au(x) = v^p(x) & \text{in } \mathbb{B}_1(0), \\ (-\Delta)^{\beta/2} v(x) + Bv(x) = u^q(x) & \text{in } \mathbb{B}_1(0), \\ u(x) \geq 0, v(x) \geq 0 & \text{in } \mathbb{B}_1(0), \\ u(x) = v(x) = 0 & \text{on } \mathbb{B}_1^C(0), \end{cases}$$

with $0 < \alpha, \beta < 2$ and $1 < p, q < \infty$. Then each positive solution $(u(x), v(x))$ must be radially symmetric and monotone decreasing about the origin.

The paper is organized as follows. Section 2 is devoted to proving Lemma 1.1, the narrow region principle for (1-1). In Section 3, we study the monotonicity of positive solutions of (1-1) in Ω and prove Theorem 1.2. Finally, the proof of Theorem 1.4 will be presented in Section 4. Note that in the following, C will be a positive constant which can be different from line to line.

2. Preliminaries

In this section, we will prove Lemma 1.1, which plays an important role in the proof of Theorem 1.2 and Theorem 1.4.

Proof. (i) Without loss of generality, let

$$x_1^* \in \Sigma_{\lambda, \delta} \quad \text{and} \quad U_\lambda(x_1^*) = \min_{x \in \Sigma_\lambda} U_\lambda(x) < 0.$$

It follows that

$$\begin{aligned} (-\Delta)^{\alpha/2} U_\lambda(x_1^*) &= C_{n, \alpha} P.V. \int_{\mathbb{R}^n} \frac{U_\lambda(x_1^*) - U_\lambda(y)}{|x_1^* - y|^{n+\alpha}} dy \\ &= C_{n, \alpha} P.V. \left(\int_{H_\lambda} \frac{U_\lambda(x_1^*) - U_\lambda(y)}{|x_1^* - y|^{n+\alpha}} dy + \int_{\mathbb{R}^n \setminus H_\lambda} \frac{U_\lambda(x_1^*) - U_\lambda(y)}{|x_1^* - y|^{n+\alpha}} dy \right) \\ &= C_{n, \alpha} P.V. \left(\int_{H_\lambda} \frac{U_\lambda(x_1^*) - U_\lambda(y)}{|x_1^* - y|^{n+\alpha}} dy + \int_{H_\lambda} \frac{U_\lambda(x_1^*) - U_\lambda(y^\lambda)}{|x_1^* - y^\lambda|^{n+\alpha}} dy \right). \end{aligned}$$

Note that $|x_1^* - y| \leq |x_1^* - y^\lambda|$ when $x_1^*, y \in \Sigma_\lambda$, and using the antisymmetry of $U_\lambda(x)$, we have

$$\begin{aligned} (-\Delta)^{\alpha/2} U_\lambda(x_1^*) &\leq C_{n, \alpha} P.V. \left(\int_{H_\lambda} \frac{U_\lambda(x_1^*) - U_\lambda(y)}{|x_1^* - y^\lambda|^{n+\alpha}} dy + \int_{H_\lambda} \frac{U_\lambda(x_1^*) + U_\lambda(y)}{|x_1^* - y^\lambda|^{n+\alpha}} dy \right) \\ &= C_{n, \alpha} P.V. \int_{H_\lambda} \frac{2U_\lambda(x_1^*)}{|x_1^* - y^\lambda|^{n+\alpha}} dy. \end{aligned}$$

Let $D = B_{2\delta}(x_1^*) \cap H_\lambda$. Then we obtain

$$(2-1) \quad \int_{H_\lambda} \frac{1}{|x_1^* - y^\lambda|^{n+\alpha}} dy \geq \int_D \frac{1}{|x_1^* - y^\lambda|^{n+\alpha}} dy \geq C \int_{B_{2\delta}(x_1^*)} \frac{1}{|x_1^* - y^\lambda|^{n+\alpha}} dy \geq \frac{C}{\delta^\alpha}.$$

Thus,

$$(-\Delta)^{\alpha/2} U_\lambda(x_1^*) \leq \frac{C U_\lambda(x_1^*)}{\delta^\alpha} < 0.$$

According to (1-7), we get

$$p\delta^{p-1}(x_1^*)V_\lambda(x_1^*) - A(x_1^*)U_\lambda(x_1^*) = (-\Delta)^{\alpha/2} U_\lambda(x_1^*) \leq \frac{C}{\delta^\alpha} U_\lambda(x_1^*) < 0.$$

Note that $A(x) = A(x')$ is bounded from below and $\xi(x)$ is between $v_\lambda(x)$ and $v(x)$, $v(x) \in G = L_\beta \cap C_{loc}^{1,1}(\Omega)$, hence, for δ sufficiently small, we have

$$(2-2) \quad V_\lambda(x_1^*) < \frac{A(x_1^*) + \frac{C}{\delta^\alpha}}{pv^{p-1}(x_1^*)} U_\lambda(x_1^*) < U_\lambda(x_1^*) < 0.$$

(ii) If there exists $x_2^* \in \Sigma_{\lambda,\delta} = \{x \in \Sigma_\lambda \mid \lambda - \delta < x_n < \lambda\}$ such that

$$V_\lambda(x_2^*) = \min_{x \in \bar{\Sigma}_\lambda} V_\lambda(x) < 0,$$

similarly to the proof of (i), we can obtain

$$(2-3) \quad (-\Delta)^{\beta/2} V_\lambda(x_2^*) \leq \frac{C V_\lambda(x_2^*)}{\delta^\beta} < 0.$$

Note that $B(x)$ is bounded from below and $u(x) \in F = L_\alpha \cap C_{loc}^{1,1}(\Omega)$, hence, for δ sufficiently small, according to (1-8), we have

$$(2-4) \quad U_\lambda(x_2^*) < \frac{B(x_2^*) + \frac{C}{\delta^\beta}}{qu^{q-1}(x_2^*)} V_\lambda(x_2^*) < V_\lambda(x_2^*) < 0.$$

Thus we have completed the proof of Lemma 1.1. \square

3. The proof of Theorem 1.2

In this section, we will carry out the direct method of moving planes on the solution $(u(x), v(x))$ along x_n direction to prove Theorem 1.2.

Proof. The proof of Theorem 1.2 is divided into two steps.

Step 1: We show that for $\lambda > 0$ sufficiently close to zero,

$$(3-1) \quad U_\lambda(x) \geq 0, \quad V_\lambda(x) \geq 0, \quad x \in \Sigma_\lambda.$$

If (3-1) does not hold, then there exists a point $x_1^* \in \Sigma_\lambda$ such that $U_\lambda(x_1^*) < 0$. Without loss of generality, we assume

$$U_\lambda(x_1^*) = \min_{x \in \bar{\Sigma}_\lambda} U_\lambda(x) < 0.$$

By Lemma 1.1, for $\lambda > 0$ sufficiently close to zero,

$$(3-2) \quad V_\lambda(x_1^*) < \frac{A(x_1^*) + \frac{C}{\lambda^\alpha}}{pv^{p-1}(x_1^*)} U_\lambda(x_1^*) < 0.$$

Note that $V_\lambda(x) = 0$ for $x \in T_\lambda$ and $V_\lambda(x) \geq 0$ for $x \in \partial \Sigma_\lambda$, hence, there exists $x_2^* \in \Sigma_\lambda$ such that

$$V_\lambda(x_2^*) = \min_{x \in \bar{\Sigma}_\lambda} V_\lambda(x) < 0.$$

Similarly to the proof of Lemma 1.1, for $\lambda > 0$ sufficiently close to zero, we have

$$(3-3) \quad U_\lambda(x_2^*) < \frac{B(x_2^*) + \frac{C}{\lambda^\beta}}{qu^{q-1}(x_2^*)} V_\lambda(x_2^*).$$

Therefore, together with (3-2) and (3-3), we get

$$(3-4) \quad \begin{aligned} V_\lambda(x_1^*) &< \frac{A(x_1^*) + \frac{C}{\lambda^\alpha}}{pv^{p-1}(x_1^*)} U_\lambda(x_1^*) \leq \frac{A(x_1^*) + \frac{C}{\lambda^\alpha}}{pv^{p-1}(x_1^*)} U_\lambda(x_2^*) \\ &< \frac{(A(x_1^*) + \frac{C}{\lambda^\alpha})}{pv^{p-1}(x_1^*)} \frac{(B(x_2^*) + \frac{C}{\lambda^\beta})}{qu^{q-1}(x_2^*)} V_\lambda(x_2^*) \\ &\leq \frac{(A(x_1^*) + \frac{C}{\lambda^\alpha})}{pv^{p-1}(x_1^*)} \frac{(B(x_2^*) + \frac{C}{\lambda^\beta})}{qu^{q-1}(x_2^*)} V_\lambda(x_1^*). \end{aligned}$$

Because $A(x) = A(x')$ and $B(x) = B(x')$ are bounded from below and $V_\lambda(x_1^*) < 0$, therefore, (3-4) is a contradiction for $\lambda > 0$ sufficiently close to zero and the proof of Step 1 is completed.

Step 1 provides a starting point. We start from such a small λ and move the plane T_λ up continuously in the direction of x_n -axis to its limiting position as long as (3-1) holds. Define

$$(3-5) \quad \lambda_0 = \sup\{\lambda > 0 \mid U_\mu(x) \geq 0, V_\mu(x) \geq 0, x \in \Sigma_\mu; \mu \leq \lambda\}.$$

Step 2: We prove

$$(3-6) \quad \lambda_0 = +\infty.$$

Before proceeding further, we investigate some properties of $U_{\lambda_0}(x)$ and $V_{\lambda_0}(x)$ for $x \in \Sigma_{\lambda_0}$.

Proposition 3.1. *If $U_{\lambda_0}(x) \equiv 0$, then $V_{\lambda_0}(x) \equiv 0$. If $V_{\lambda_0}(x) \equiv 0$, then $U_{\lambda_0}(x) \equiv 0$.*

Proof. If $U_{\lambda_0}(x) \equiv 0$, then (1-7) becomes

$$0 = (-\Delta)^{\alpha/2} U_{\lambda_0}(x) = p\xi^{p-1}(x) V_{\lambda_0}(x),$$

Obviously, we get $V_{\lambda_0}(x) \equiv 0$. □

Proposition 3.2. *If $U_{\lambda_0}(x) \not\equiv 0$ or $V_{\lambda_0}(x) \not\equiv 0$, then $U_{\lambda_0}(x) > 0$ and $V_{\lambda_0}(x) > 0$ for all $x \in \Sigma_{\lambda_0}$.*

Proof. Since we know that $U_{\lambda_0}(x) \geq 0$, $x \in \Sigma_{\lambda_0}$. If $U_{\lambda_0}(x) > 0$ does not hold, we assume that there exists some point $\tilde{x} \in \Sigma_{\lambda_0}$ such that $U_{\lambda_0}(\tilde{x}) = 0$.

$$\begin{aligned}
 (-\Delta)^{\alpha/2}U_{\lambda_0}(\tilde{x}) &= C_{n,\alpha}P.V. \int_{\mathbb{R}^n} \frac{U_{\lambda_0}(\tilde{x}) - U_{\lambda_0}(y)}{|\tilde{x} - y|^{n+\alpha}} dy \\
 &= C_{n,\alpha}P.V. \left(\int_{H_{\lambda_0}} \frac{-U_{\lambda_0}(y)}{|\tilde{x} - y|^{n+\alpha}} dy + \int_{\mathbb{R}^n \setminus H_{\lambda_0}} \frac{-U_{\lambda_0}(y)}{|\tilde{x} - y|^{n+\alpha}} dy \right) \\
 &= C_{n,\alpha}P.V. \left(\int_{H_{\lambda_0}} \frac{-U_{\lambda_0}(y)}{|\tilde{x} - y|^{n+\alpha}} dy + \int_{H_{\lambda_0}} \frac{-U_{\lambda_0}(y^\lambda)}{|\tilde{x} - y^\lambda|^{n+\alpha}} dy \right) \\
 (3-7) \quad &= C_{n,\alpha}P.V. \int_{H_{\lambda_0}} \left(\frac{1}{|\tilde{x} - y^\lambda|^{n+\alpha}} - \frac{1}{|\tilde{x} - y|^{n+\alpha}} \right) U_{\lambda_0}(y) dy.
 \end{aligned}$$

Note that $|\tilde{x} - y^\lambda| > |\tilde{x} - y|$, hence,

$$(3-8) \quad (-\Delta)^{\alpha/2}U_{\lambda_0}(\tilde{x}) < 0.$$

On the other hand, according to (1-7),

$$(3-9) \quad (-\Delta)^{\alpha/2}U_{\lambda_0}(\tilde{x}) = p\xi^{p-1}(\tilde{x})V_{\lambda_0}(\tilde{x}) - A(\tilde{x})U_{\lambda_0}(\tilde{x}) = p\xi^{p-1}(\tilde{x})V_{\lambda_0}(\tilde{x}) \geq 0.$$

Evidently, this is contradictory to (3-8). Consequently, we obtain $U_{\lambda_0}(x) > 0$ for all $x \in \Sigma_{\lambda_0}$. Then by Proposition 3.1, we get $V_{\lambda_0}(x) \not\equiv 0$. Similarly to above, we obtain $V_{\lambda_0}(x) > 0$ for all $x \in \Sigma_{\lambda_0}$, completing the proof. \square

Now, we start to prove (3-6). If $\lambda_0 < +\infty$, we will show

$$(3-10) \quad U_{\lambda_0}(x) \equiv 0, \quad x \in \Sigma_{\lambda_0}.$$

Then by Proposition 3.1, $V_{\lambda_0}(x) \equiv 0$, $x \in \Sigma_{\lambda_0}$. Thus, we obtain

$$(3-11) \quad u(x', 2\lambda_0) = u(x', 0),$$

$$(3-12) \quad v(x', 2\lambda_0) = v(x', 0).$$

But the left-hand side of (3-11) is positive, and the right-hand side of (3-11) is equal to zero. This is contradictory. The same holds for $v(x)$. Hence, (3-6) holds.

In the following, we will prove (3-10). If $U_{\lambda_0}(x) \not\equiv 0$, $x \in \Sigma_{\lambda_0}$, by Proposition 3.2, we have $U_{\lambda_0}(x) > 0$, $V_{\lambda_0}(x) > 0$, $x \in \Sigma_{\lambda_0}$. Hence, for small $\delta > 0$, there exists a positive constant c_0 such that

$$U_{\lambda_0}(x) \geq c_0 > 0, \quad V_{\lambda_0}(x) \geq c_0 > 0, \quad x \in \bar{\Sigma}_{\lambda_0-\delta}.$$

Since $U_\lambda(x)$ and $V_\lambda(x)$ depends on λ continuously, there exists $\varepsilon > 0$ and $\varepsilon < \delta$ such that for all $\lambda \in (\lambda_0, \lambda_0 + \varepsilon)$,

$$(3-13) \quad U_\lambda(x) \geq 0, \quad V_\lambda(x) \geq 0, \quad x \in \bar{\Sigma}_{\lambda_0-\delta}.$$

When $x \in \Sigma_\lambda \setminus \bar{\Sigma}_{\lambda_0-\delta}$, we also have $U_\lambda(x) \geq 0$, $V_\lambda(x) \geq 0$. If not, without loss of generality, we assume that there exists a point $\bar{x}_1 \in \Sigma_\lambda \setminus \bar{\Sigma}_{\lambda_0-\delta}$ such that

$$U_\lambda(\bar{x}_1) = \min_{x \in \Sigma_\lambda \setminus \bar{\Sigma}_{\lambda_0-\delta}} U_\lambda(x) < 0.$$

According to Lemma 1.1,

$$V_\lambda(\bar{x}_1) < \frac{A(\bar{x}_1) + \frac{C}{(\delta+\varepsilon)^\alpha}}{pv^{p-1}(\bar{x}_1)} U_\lambda(\bar{x}_1) < 0.$$

Hence, there exists $\bar{x}_2 \in \Sigma_\lambda \setminus \bar{\Sigma}_{\lambda_0-\delta}$ such that

$$V_\lambda(\bar{x}_2) = \min_{x \in \Sigma_\lambda \setminus \bar{\Sigma}_{\lambda_0-\delta}} V_\lambda(x) < 0.$$

By Lemma 1.1 again,

$$U_\lambda(\bar{x}_2) < \frac{B(\bar{x}_2) + \frac{C}{(\delta+\varepsilon)^\beta}}{qu^{q-1}(\bar{x}_2)} V_\lambda(\bar{x}_2) < 0.$$

Similarly to (3-4),

$$U_\lambda(\bar{x}_2) < \frac{\left(A(\bar{x}_1) + \frac{C}{(\delta+\varepsilon)^\alpha}\right) \left(B(\bar{x}_2) + \frac{C}{(\delta+\varepsilon)^\beta}\right)}{pv^{p-1}(\bar{x}_1) qu^{q-1}(\bar{x}_2)} U_\lambda(\bar{x}_2).$$

Since $A(x) = A(x')$ and $B(x) = B(x')$ are bounded from below and $U_\lambda(\bar{x}_2) < 0$, this is contradictory for δ and ε sufficiently small. Therefore, we obtain

$$(3-14) \quad U_\lambda(x) \geq 0, \quad V_\lambda(x) \geq 0, \quad x \in \Sigma_\lambda \setminus \bar{\Sigma}_{\lambda_0-\delta}.$$

Combining (3-13) and (3-14), we get that for $\lambda \in (\lambda_0, \lambda_0 + \varepsilon)$,

$$(3-15) \quad U_\lambda(x) \geq 0, \quad V_\lambda(x) \geq 0, \quad x \in \Sigma_\lambda.$$

(3-15) indicates that the plane T_{λ_0} can be moved up further. We have reached a contradiction with the definition of λ_0 . Hence, we must have $U_{\lambda_0}(x) \equiv 0$.

We have shown that $\lambda_0 = +\infty$ and $U_{\lambda_0}(x) \geq 0$, $V_{\lambda_0}(x) \geq 0$. It indicates that $u(x)$ and $v(x)$ are monotonically increasing in x_n , which completes the proof of Theorem 1.2. \square

Proof of Corollary 1.3. We have shown that $u(x)$ and $v(x)$ are monotonically increasing in x_n . In terms of $u(0) = v(0) = 0$ and the condition (1-10), we derive

$$u(x) \equiv 0, \quad v(x) \equiv 0 \quad \text{in } \Omega.$$

Thus we have completed the proof. \square

4. The proof of Theorem 1.4

In this section, we will apply Lemma 1.1 to prove Theorem 1.4. We consider the case $A(x) = A$ and $B(x) = B$ in system (1-1), where A, B are constants. In this case, (1-1) becomes (1-11).

Choose any direction to be the x_1 direction. We write $x = (x_1, x') \in \mathbb{R}^n$ with $x' = (x_2, x_3, \dots, x_n)$. Let

$$\hat{T}_\lambda = \{x \in \mathbb{R}^n \mid x_1 = \lambda, \lambda \in \mathbb{R}, \lambda > -1\}$$

be the moving planes and denote

$$\hat{H}_\lambda = \{x \in \mathbb{R}^n \mid x_1 < \lambda\}, \quad \hat{\Sigma}_\lambda = \{x \in \mathbb{B}_1(0) \mid -1 < x_1 < \lambda\}.$$

For each point $x = (x_1, x') \in \hat{\Sigma}_\lambda$, let $x^\lambda = (2\lambda - x_1, x')$ be the reflection point about the plane \hat{T}_λ and $U_\lambda(x), V_\lambda(x)$ defined as before. Then it follows that for $x \in \hat{\Sigma}_\lambda$,

$$(4-1) \quad (-\Delta)^{\alpha/2} U_\lambda(x) = p\xi^{p-1}(x)V_\lambda(x) - AU_\lambda(x),$$

and

$$(4-2) \quad (-\Delta)^{\beta/2} V_\lambda(x) = q\eta^{q-1}(x)U_\lambda(x) - BV_\lambda(x),$$

where $\xi(x)$ is between $v_\lambda(x)$ and $v(x)$ and $\eta(x)$ is between $u_\lambda(x)$ and $u(x)$.

Proof of Theorem 1.4. Step 1: We show that for $\lambda > -1$ sufficiently close to -1 ,

$$(4-3) \quad U_\lambda(x) \geq 0, \quad V_\lambda(x) \geq 0, \quad x \in \hat{\Sigma}_\lambda.$$

The proof is almost the same as Step 1 in the proof of Theorem 1.2.

Step 1 provides a starting point. We start from such a small λ and move the plane \hat{T}_λ continuously in the direction of x_1 -axis to its limiting position as long as (4-3) holds.

Step 2: Define

$$(4-4) \quad \lambda_0 = \sup\{\lambda > -1 \mid U_\mu(x) \geq 0, V_\mu(x) \geq 0, x \in \hat{\Sigma}_\mu; \mu \leq \lambda\}.$$

We will prove

$$(4-5) \quad \lambda_0 = 0.$$

If $\lambda_0 < 0$, we will show that the plane \hat{T}_λ can be moved further right. That is, there exists $\varepsilon > 0$ such that for all $\lambda \in (\lambda_0, \lambda_0 + \varepsilon)$,

$$(4-6) \quad U_\lambda(x) \geq 0, \quad V_\lambda(x) \geq 0, \quad x \in \hat{\Sigma}_\lambda.$$

This is a contradiction with the definition of λ_0 . Hence, we must have $\lambda_0 = 0$.

The proof of (4-6) is composed of two parts.

(a): We show that for $\varepsilon > 0$, $\delta > 0$ and $\varepsilon < \delta$, when $\lambda \in (\lambda_0, \lambda_0 + \varepsilon)$,

$$(4-7) \quad U_\lambda(x) \geq 0, \quad V_\lambda(x) \geq 0, \quad x \in \bar{\Sigma}_{\lambda_0 - \delta}.$$

We know that

$$U_{\lambda_0}(x) \geq 0, \quad V_{\lambda_0}(x) \geq 0, \quad x \in \hat{\Sigma}_{\lambda_0}.$$

In fact, if $\lambda_0 < 0$, we must have

$$U_{\lambda_0}(x) > 0, \quad V_{\lambda_0}(x) > 0, \quad x \in \hat{\Sigma}_{\lambda_0}.$$

If $U_{\lambda_0}(x) > 0$ does not hold, we assume that there exists some point $\tilde{x} \in \hat{\Sigma}_{\lambda_0}$ such that $U_{\lambda_0}(\tilde{x}) = 0$.

$$\begin{aligned} (-\Delta)^{\alpha/2} U_{\lambda_0}(\tilde{x}) &= C_{n,\alpha} P.V. \int_{\mathbb{R}^n} \frac{U_{\lambda_0}(\tilde{x}) - U_{\lambda_0}(y)}{|\tilde{x} - y|^{n+\alpha}} dy \\ &= C_{n,\alpha} P.V. \int_{\hat{H}_{\lambda_0}} \frac{-U_{\lambda_0}(y)}{|\tilde{x} - y|^{n+\alpha}} dy + \int_{\mathbb{R}^n \setminus \hat{H}_{\lambda_0}} \frac{-U_{\lambda_0}(y)}{|\tilde{x} - y|^{n+\alpha}} dy \\ &= C_{n,\alpha} P.V. \int_{\hat{H}_{\lambda_0}} \frac{-U_{\lambda_0}(y)}{|\tilde{x} - y|^{n+\alpha}} dy + \int_{\hat{H}_{\lambda_0}} \frac{-U_{\lambda_0}(y^\lambda)}{|\tilde{x} - y^\lambda|^{n+\alpha}} dy \\ &= C_{n,\alpha} P.V. \int_{\hat{H}_{\lambda_0}} \left(\frac{1}{|\tilde{x} - y^\lambda|^{n+\alpha}} - \frac{1}{|\tilde{x} - y|^{n+\alpha}} \right) U_{\lambda_0}(y) dy. \end{aligned}$$

Note that $|\tilde{x} - y^\lambda| > |\tilde{x} - y|$, hence,

$$(4-8) \quad (-\Delta)^{\alpha/2} U_{\lambda_0}(\tilde{x}) < 0.$$

On the other hand, according to (4-1),

$$(-\Delta)^{\alpha/2} U_{\lambda_0}(\tilde{x}) = p\xi^{p-1}(\tilde{x})V_\lambda(\tilde{x}) - AU_\lambda(\tilde{x}) = p\xi^{p-1}(\tilde{x})V_\lambda(\tilde{x}) \geq 0.$$

This is contradictory to (4-8). Consequently, we obtain $U_{\lambda_0}(x) > 0$ for all $x \in \hat{\Sigma}_{\lambda_0}$. Similarly, we can show $V_{\lambda_0}(x) > 0$ for all $x \in \hat{\Sigma}_{\lambda_0}$. Hence, for small $\delta > 0$, there exists a positive constant c_0 such that

$$U_{\lambda_0}(x) \geq c_0 > 0, \quad V_{\lambda_0}(x) \geq c_0 > 0, \quad x \in \bar{\Sigma}_{\lambda_0 - \delta}.$$

Since $U_\lambda(x)$ and $V_\lambda(x)$ depend on λ continuously, there exists $\varepsilon > 0$ with $\varepsilon < \delta$ such that for all $\lambda \in (\lambda_0, \lambda_0 + \varepsilon)$,

$$U_\lambda(x) \geq 0, \quad V_\lambda(x) \geq 0, \quad x \in \bar{\Sigma}_{\lambda_0 - \delta}.$$

(b): Using Lemma 1.1 and similarly to the proof of (3-14), we get

$$(4-9) \quad U_\lambda(x) \geq 0, \quad V_\lambda(x) \geq 0 \quad x \in \hat{\Sigma}_\lambda \setminus \bar{\Sigma}_{\lambda_0 - \delta}.$$

Together with **(a)** and **(b)**, we prove (4-6) is true for all $\lambda \in (\lambda_0, \lambda_0 + \varepsilon)$. Thus, we obtain $\lambda_0 = 0$ and $U_{\lambda_0}(x) \geq 0$, $V_{\lambda_0}(x) \geq 0$, $x \in \hat{\Sigma}_{\lambda_0}$.

Similarly, we move the plane T_λ from 1 to the left and show that

$$U_{\lambda_0}(x) \leq 0, V_{\lambda_0}(x) \leq 0, \quad x \in \hat{\Sigma}_{\lambda_0}.$$

Then we obtain that

$$\lambda_0 = 0 \quad \text{and} \quad U_{\lambda_0}(x) \equiv 0, V_{\lambda_0}(x) \equiv 0, \quad x \in \hat{\Sigma}_{\lambda_0}.$$

This indicates that $u(x)$ and $v(x)$ are symmetric about T_0 . Since the x_1 direction can be chosen arbitrarily, we have actually shown that $u(x)$ and $v(x)$ are radially symmetric about the origin. Thus, we have completed the proof of Theorem 1.4. \square

Acknowledgement

I express my deep thanks to my advisor Li Ma for guidance and I could not have finished this paper without his generous help. I thank the unknown referees very much for useful suggestions.

References

- [Alexandrov 1962] A. D. Alexandrov, “A characteristic property of spheres”, *Ann. Mat. Pura Appl.* (4) **58** (1962), 303–315. MR Zbl
- [Applebaum 2009] D. Applebaum, *Lévy processes and stochastic calculus*, 2nd ed., Cambridge Studies in Advanced Mathematics **116**, Cambridge Univ. Press, 2009. MR Zbl
- [Berestycki and Nirenberg 1992] H. Berestycki and L. Nirenberg, “Travelling fronts in cylinders”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **9**:5 (1992), 497–572. MR Zbl
- [Berestycki et al. 1993] H. Berestycki, L. Caffarelli, and L. Nirenberg, “Symmetry for elliptic equations in a half space”, pp. 27–42 in *Boundary value problems for partial differential equations and applications*, edited by J.-L. Lions and C. Baiocchi, RMA Res. Notes Appl. Math. **29**, Masson, Paris, 1993. MR Zbl
- [Berestycki et al. 1996] H. Berestycki, L. Caffarelli, and L. Nirenberg, “Inequalities for second-order elliptic equations with applications to unbounded domains, I”, *Duke Math. J.* **81**:2 (1996), 467–494. MR Zbl
- [Berestycki et al. 1997] H. Berestycki, L. Caffarelli, and L. Nirenberg, “Further qualitative properties for elliptic equations in unbounded domains”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **25**:1-2 (1997), 69–94. MR Zbl
- [Bertoin 1996] J. Bertoin, *Lévy processes*, Cambridge Tracts in Mathematics **121**, Cambridge Univ. Press, 1996. MR Zbl
- [Bouchaud and Georges 1990] J.-P. Bouchaud and A. Georges, “Anomalous diffusion in disordered media: statistical mechanisms, models and physical applications”, *Phys. Rep.* **195**:4-5 (1990), 127–293. MR
- [Brändle et al. 2013] C. Brändle, E. Colorado, A. de Pablo, and U. Sánchez, “A concave-convex elliptic problem involving the fractional Laplacian”, *Proc. Roy. Soc. Edinburgh Sect. A* **143**:1 (2013), 39–71. MR Zbl

- [Busca and Sirakov 2000] J. Busca and B. Sirakov, “Symmetry results for semilinear elliptic systems in the whole space”, *J. Differential Equations* **163**:1 (2000), 41–56. MR Zbl
- [Caffarelli and Silvestre 2007] L. Caffarelli and L. Silvestre, “An extension problem related to the fractional Laplacian”, *Comm. Partial Differential Equations* **32**:7-9 (2007), 1245–1260. MR Zbl
- [Caffarelli and Vasseur 2010] L. Caffarelli and A. Vasseur, “Drift diffusion equations with fractional diffusion and the quasi-geostrophic equation”, *Ann. of Math. (2)* **171**:3 (2010), 1903–1930. MR Zbl
- [Cai and Mei 2017] M. Cai and L. Mei, “Narrow region principle and Liouville type results for fractional Lane–Emden system”, *Complex Var. Elliptic Equ.* **62**:7 (2017), 1002–1014. MR Zbl
- [Chen and Li 2009] W. Chen and C. Li, “An integral system and the Lane–Emden conjecture”, *Discrete Contin. Dyn. Syst.* **24**:4 (2009), 1167–1184. MR Zbl
- [Chen and Li 2010] W. Chen and C. Li, *Methods on nonlinear elliptic equations*, AIMS Series on Differential Equations & Dynamical Systems **4**, American Inst. Math. Sci., Springfield, MO, 2010. MR Zbl
- [Chen and Zhu 2016] W. Chen and J. Zhu, “Indefinite fractional elliptic problem and Liouville theorems”, *J. Differential Equations* **260**:5 (2016), 4758–4785. MR Zbl
- [Chen et al. 2005a] W. Chen, C. Li, and B. Ou, “Classification of solutions for a system of integral equations”, *Comm. Partial Differential Equations* **30**:1-3 (2005), 59–65. MR Zbl
- [Chen et al. 2005b] W. Chen, C. Li, and B. Ou, “Qualitative properties of solutions for an integral equation”, *Discrete Contin. Dyn. Syst.* **12**:2 (2005), 347–354. MR Zbl
- [Chen et al. 2006] W. Chen, C. Li, and B. Ou, “Classification of solutions for an integral equation”, *Comm. Pure Appl. Math.* **59**:3 (2006), 330–343. MR Zbl
- [Chen et al. 2015] W. Chen, Y. Fang, and R. Yang, “Liouville theorems involving the fractional Laplacian on a half space”, *Adv. Math.* **274** (2015), 167–198. MR Zbl
- [Chen et al. 2017a] W. Chen, C. Li, and G. Li, “Maximum principles for a fully nonlinear fractional order equation and symmetry of solutions”, *Calc. Var. Partial Differential Equations* **56**:2 (2017), art. id. 29. MR Zbl
- [Chen et al. 2017b] W. Chen, C. Li, and Y. Li, “A direct method of moving planes for the fractional Laplacian”, *Adv. Math.* **308** (2017), 404–437. MR Zbl
- [Cheng et al. 2017] T. Cheng, G. Huang, and C. Li, “The maximum principles for fractional Laplacian equations and their applications”, *Commun. Contemp. Math.* **19**:6 (2017), art. id. 1750018. MR
- [Constantin 2006] P. Constantin, “Euler equations, Navier–Stokes equations and turbulence”, pp. 1–43 in *Mathematical foundation of turbulent viscous flows* (Martina Franca, Italy, 2003), edited by M. Cannone and T. Miyakawa, Lecture Notes in Math. **1871**, Springer, 2006. MR Zbl
- [Fall and Weth 2016] M. M. Fall and T. Weth, “Monotonicity and nonexistence results for some fractional elliptic problems in the half-space”, *Commun. Contemp. Math.* **18**:1 (2016), art. id. 1550012. MR Zbl
- [Fang and Zhang 2013] Y. Fang and J. Zhang, “Nonexistence of positive solution for an integral equation on a half-space \mathbb{R}_+^n ”, *Commun. Pure Appl. Anal.* **12**:2 (2013), 663–678. MR Zbl
- [de Figueiredo and Felmer 1994] D. G. de Figueiredo and P. L. Felmer, “A Liouville-type theorem for elliptic systems”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **21**:3 (1994), 387–397. MR Zbl
- [Gidas et al. 1979] B. Gidas, W.-M. Ni, and L. Nirenberg, “Symmetry and related properties via the maximum principle”, *Comm. Math. Phys.* **68**:3 (1979), 209–243. MR Zbl
- [Gidas et al. 1981] B. Gidas, W.-M. Ni, and L. Nirenberg, “Symmetry of positive solutions of nonlinear elliptic equations in \mathbb{R}^n ”, pp. 369–402 in *Mathematical analysis and applications, A*, edited by L. Nachbin, Adv. in Math. Suppl. Stud. **7a**, Academic Press, New York, 1981. MR Zbl

- [Jarohs and Weth 2016] S. Jarohs and T. Weth, “Symmetry via antisymmetric maximum principles in nonlocal problems of variable order”, *Ann. Mat. Pura Appl.* (4) **195**:1 (2016), 273–291. MR Zbl
- [Li and Ma 2008] C. Li and L. Ma, “Uniqueness of positive bound states to Schrödinger systems with critical exponents”, *SIAM J. Math. Anal.* **40**:3 (2008), 1049–1057. MR Zbl
- [Liu and Ma 2012] B. Liu and L. Ma, “Symmetry results for decay solutions of semilinear elliptic systems on half spaces”, *Nonlinear Anal.* **75**:6 (2012), 3167–3177. MR Zbl
- [Liu and Ma 2013] B. Liu and L. Ma, “Symmetry results for elliptic Schrödinger systems on half spaces”, *J. Math. Anal. Appl.* **401**:1 (2013), 259–268. MR Zbl
- [Liu and Ma 2016] B. Liu and L. Ma, “Radial symmetry results for fractional Laplacian systems”, *Nonlinear Anal.* **146** (2016), 120–135. MR Zbl
- [Ma and Chen 2008] L. Ma and D. Chen, “Radial symmetry and monotonicity for an integral equation”, *J. Math. Anal. Appl.* **342**:2 (2008), 943–949. MR Zbl
- [Ma and Liu 2010] L. Ma and B. Liu, “Symmetry results for decay solutions of elliptic systems in the whole space”, *Adv. Math.* **225**:6 (2010), 3052–3063. MR Zbl
- [Ma and Zhao 2008] L. Ma and L. Zhao, “Uniqueness of ground states of some coupled nonlinear Schrödinger systems and their application”, *J. Differential Equations* **245**:9 (2008), 2551–2565. MR Zbl
- [Mitidieri 1996] E. Mitidieri, “Nonexistence of positive solutions of semilinear elliptic systems in \mathbb{R}^N ”, *Differential Integral Equations* **9**:3 (1996), 465–479. MR Zbl
- [Montaru and Souplet 2014] A. Montaru and P. Souplet, “Symmetry of components and Liouville theorems for noncooperative elliptic systems on the half-space”, *C. R. Math. Acad. Sci. Paris* **352**:4 (2014), 321–325. MR Zbl
- [Poláčik et al. 2007] P. Poláčik, P. Quittner, and P. Souplet, “Singularity and decay estimates in superlinear problems via Liouville-type theorems, I: Elliptic equations and systems”, *Duke Math. J.* **139**:3 (2007), 555–579. MR Zbl
- [Serrin 1971] J. Serrin, “A symmetry problem in potential theory”, *Arch. Rational Mech. Anal.* **43** (1971), 304–318. MR Zbl
- [Serrin and Zou 1996] J. Serrin and H. Zou, “Non-existence of positive solutions of Lane–Emden systems”, *Differential Integral Equations* **9**:4 (1996), 635–653. MR Zbl
- [Souplet 2009] P. Souplet, “The proof of the Lane–Emden conjecture in four space dimensions”, *Adv. Math.* **221**:5 (2009), 1409–1427. MR Zbl
- [Tarasov and Zaslavsky 2006] V. E. Tarasov and G. M. Zaslavsky, “Fractional dynamics of systems with long-range interaction”, *Commun. Nonlinear Sci. Numer. Simul.* **11**:8 (2006), 885–898. MR Zbl
- [Zhuo et al. 2016] R. Zhuo, W. Chen, X. Cui, and Z. Yuan, “Symmetry and non-existence of solutions for a nonlinear system involving the fractional Laplacian”, *Discrete Contin. Dyn. Syst.* **36**:2 (2016), 1125–1141. MR Zbl

Received December 28, 2016. Revised June 1, 2017.

JING LI
 COLLEGE OF PHYSICS AND MATERIALS SCIENCE
 COLLEGE OF MATHEMATICS AND INFORMATION SCIENCE
 HENAN NORMAL UNIVERSITY
 XINXIANG
 CHINA
 jingli@htu.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 294 No. 1 May 2018

Three-dimensional Sol manifolds and complex Kleinian groups	1
WALDEMAR BARRERA, RENE GARCIA-LARA and JUAN NAVARRETE	
On periodic points of symplectomorphisms on surfaces	19
MARTA BATORÉO	
Mixing properties for hom-shifts and the distance between walks on associated graphs	41
NISHANT CHANDGOTIA and BRIAN MARCUS	
Simultaneous construction of hyperbolic isometries	71
MATT CLAY and CAGLAR UYANIK	
A local weighted Axler–Zheng theorem in \mathbb{C}^n	89
ŽELJKO ČUČKOVIĆ, SÖNMEZ ŞAHUTOĞLU and YUNUS E. ZEYTUNCU	
Monotonicity and radial symmetry results for Schrödinger systems with fractional diffusion	107
JING LI	
Moduli spaces of stable pairs	123
YINBANG LIN	
Spark deficient Gabor frames	159
ROMANOS-DIOGENES MALIKIOSIS	
Ordered groups as a tensor category	181
DALE ROLFSEN	
Multiplication of distributions and a nonlinear model in elastodynamics	195
C. O. R. SARRICO	
Some Ambrose- and Galloway-type theorems via Bakry–Émery and modified Ricci curvatures	213
HOMARE TADANO	
Irreducible decomposition for local representations of quantum Teichmüller space	233
JÉRÉMY TOULISSE	



0030-8730(201805)294:1;1-T