

*Pacific
Journal of
Mathematics*

ORDERED GROUPS AS A TENSOR CATEGORY

DALE ROLFSEN

ORDERED GROUPS AS A TENSOR CATEGORY

DALE ROLFSEN

It is a classical theorem that the free product of ordered groups is orderable. In this note we show that, using a method of G. Bergman, an ordering of the free product can be constructed in a functorial manner, in the category of ordered groups and order-preserving homomorphisms. With this functor interpreted as a tensor product this category becomes a tensor (or monoidal) category. Moreover, if $O(G)$ denotes the space of orderings of the group G with the natural topology, then for fixed groups F and G our construction can be considered a function $O(F) \times O(G) \rightarrow O(F * G)$. We show that this function is continuous and injective. Similar results hold for left-ordered groups.

1. Introduction

An ordered group $(G, <)$ is a group G together with a strict total ordering $<$ of its elements such that $x < y$ implies $xz < yz$ and $zx < zy$ for all $x, y, z \in G$. If such an ordering exists, G is said to be orderable. If $(F, <_F)$ and $(G, <_G)$ are ordered groups, a homomorphism $\phi : F \rightarrow G$ is said to be order-preserving (relative to $<_F, <_G$) if for all $x, y \in F$ we have $x <_F y \implies \phi(x) <_G \phi(y)$. In this case the reverse implication follows, and ϕ is necessarily injective.

A theorem of Vinogradov [1949] asserts that if F and G are orderable groups, then the free product $F * G$ (sometimes called the coproduct, as in [Bergman 1990]) is orderable. Other proofs of this can be found in [Johnson 1968; Passman 1977; Bergman 1990], and a generalization in [Chiswell 2012]. A proof given in [Botto Mura and Rhemtulla 1977] was unfortunately found to have a gap, as discussed in [Holland and Medvedev 1994; Chiswell 2014]. Yet another proof, in [Révész 1987], was also shown to have a gap [Medvedev 1991].

Here we show that a version of the construction in [Bergman 1990] is functorial in the following sense. Suppose $(F_i, <_{F_i})$, $i = 0, 1$, are ordered groups. We will

The author gratefully acknowledges the support of a grant from the Canadian Natural Sciences and Engineering Research Council. Thanks also to George Bergman, Adam Clay and Christian Kassel for very helpful comments on earlier versions of this paper, and to Victoria Lebed and Arnaud Mortier for providing an English translation of [Vinogradov 1949] (arxiv 1703.05781).

MSC2010: 18D10, 20F60.

Keywords: ordered group, free product, coproduct, tensor category, monoidal category.

construct an ordering \prec of $F_0 * F_1$, so that $(F_0 * F_1, \prec)$ is an ordered group, and write

$$\mathfrak{F}((F_0, \prec_{F_0}), (F_1, \prec_{F_1})) := (F_0 * F_1, \prec).$$

Theorem 1 shows that \mathfrak{F} is a (bi)functor in the category \mathfrak{C} of ordered groups and order-preserving homomorphisms. We will show in [Section 5](#) that this functor gives \mathfrak{C} the structure of a tensor, or monoidal, category.

Theorem 1. *Suppose that (F_i, \prec_{F_i}) , $i = 0, 1$, are ordered groups. Then the ordered group $(F_0 * F_1, \prec_F) = \mathfrak{F}((F_0, \prec_{F_0}), (F_1, \prec_{F_1}))$ has the following properties:*

- (1) \prec_F extends the given orderings of F_i as subgroups of $F_0 * F_1$.
- (2) If (G_i, \prec_{G_i}) , $i = 0, 1$, are ordered groups and

$$(G_0 * G_1, \prec_G) = \mathfrak{F}((G_0, \prec_{G_0}), (G_1, \prec_{G_1}))$$

and if $\phi_i : F_i \rightarrow G_i$, $i = 0, 1$, are homomorphisms which preserve the given orderings of F_i and G_i , then the homomorphism $\phi_0 * \phi_1 : F_0 * F_1 \rightarrow G_0 * G_1$ is order-preserving, relative to \prec_F, \prec_G .

In [Section 8](#), [Theorem 1](#) will be extended to free products of an arbitrary, possibly infinite, collection of ordered groups. We will typically use multiplicative notation for groups and use 1 to denote the identity element, though additive groups are also considered, with 0 as identity element. We may also use 1 to denote the unit of a ring (all rings we consider are assumed to have a unit), as well as the natural number.

Many of our results could have been proven using the original construction of Vinogradov. Like Bergman's, his proof involves embedding a free product of groups into a ring of matrices. Vinogradov's matrices are infinite dimensional upper triangular matrices, whereas Bergman's are 2×2 matrices with polynomial entries, a useful simplification.

2. Embedding free products in matrix rings

We use an observation of Bergman which generalizes the fact that the matrices $\begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix}$ freely generate a free subgroup of the multiplicative group of invertible 2×2 matrices with entries in the polynomial ring $\mathbb{Z}[t]$.

Consider a ring R without zero divisors and let F and G be multiplicative groups of nonzero elements of R . Let $M_2(R[t])$ be the ring of 2×2 matrices with entries in the polynomial ring $R[t]$. Then one can embed F in $M_2(R[t])$ by $f \mapsto \begin{pmatrix} f & 0 \\ 0 & 1 \end{pmatrix}$. But we can conjugate that by $\begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}$ to get a different embedding which has a highest degree in the upper right corner when $f \neq 1$:

$$\rho(f) = \begin{pmatrix} 1 & -t \\ 0 & 1 \end{pmatrix} \begin{pmatrix} f & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} f & (f-1)t \\ 0 & 1 \end{pmatrix}.$$

Similarly we embed G by

$$\rho(g) = \begin{pmatrix} 1 & 0 \\ (g-1)t & g \end{pmatrix}.$$

This then defines a multiplicative homomorphism $\rho : F * G \rightarrow M_2(R[t])$, which Bergman observes to be a faithful representation.

Proposition 2 [Bergman 1990, Corollary 12]. *With the assumptions stated in the preceding paragraph, $\rho : F * G \rightarrow M_2(R[t])$ is injective.*

Proof. Here is a sketch of a proof using a ping-pong argument. Let

$$f_k g_k f_{k-1} \cdots g_2 f_1 g_1 \neq 1$$

be a reduced word in $F * G$, with $f_i \in F, g_i \in G$ nonidentity elements (except possibly for $i \in \{1, k\}$). Assume that $g_1 \neq 1$, the other case with $g_1 = 1, f_1 \neq 1$ being similar. We need to show that the product of matrices $\rho(f_k)\rho(g_k) \cdots \rho(f_1)\rho(g_1)$ is not the identity matrix. Consider the set V of column vectors $\begin{pmatrix} A(t) \\ B(t) \end{pmatrix}$ with entries in $R[t]$ and partition that set into three parts $V = V_1 \sqcup V_2 \sqcup V_3$ according to their degrees as polynomials. Take V_1 to be the set of such pairs with $\deg A(t) > \deg B(t)$, V_2 the set with $\deg A(t) < \deg B(t)$ and V_3 the set with equal degree.

Apply $\rho(f_k)\rho(g_k) \cdots \rho(f_1)\rho(g_1)$ (on the left) to the vector $\begin{pmatrix} 1 \\ 1 \end{pmatrix} \in V_3$ and note that $\rho(g_1)$ sends $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ to $\begin{pmatrix} 1 \\ g_1+(g_1-1)t \end{pmatrix}$ which belongs to V_2 . Then $\rho(f_1)$ sends this result into V_1 , which is then sent to V_2 by $\rho(g_2)$, and so on. The end result, after multiplying all the matrices, will be in V_1 or V_2 , not V_3 , and so the product cannot be the identity matrix. \square

3. Constructing the ordering \prec

Suppose we are given two ordered groups, $(F_0, <_{F_0})$ and $(F_1, <_{F_1})$. To embed them in a ring, we take R to be the integral group ring of their direct product: $R = \mathbb{Z}(F_0 \times F_1)$. It is well known that integral group rings of orderable groups have no zero divisors (see, for example, [Botto Mura and Rhemtulla 1977] p. 155), so R has no zero divisors. Define a multiplicative homomorphism $\rho : F_0 * F_1 \rightarrow M_2(R[t])$ by

$$\rho(f_0) = \begin{pmatrix} f_0 & (f_0-1)t \\ 0 & 1 \end{pmatrix} \quad \rho(f_1) = \begin{pmatrix} 1 & 0 \\ (f_1-1)t & f_1 \end{pmatrix}, \quad f_i \in F_i.$$

By Proposition 2, ρ is faithful; it defines an isomorphism of $F_0 * F_1$ onto a multiplicative subgroup of $M_2(R[t])$.

We now turn to the task of defining the ordering, choosing a specific recipe among many described in [Bergman 1990]. First we order $F_0 \times F_1$ lexicographically, defining $(f_0, f_1) < (f'_0, f'_1)$ if $f_0 <_{F_0} f'_0$ or else $f_0 = f'_0$ and $f_1 <_{F_1} f'_1$. Then the

group ring $R = \mathbb{Z}(F_0 \times F_1)$ becomes an ordered ring¹ by declaring a nonzero element to be *positive* if the coefficient of the largest term (in the ordering $<$ of $F_0 \times F_1$) is a positive integer.

Note that as a ring element, $f_0 \in F_0$, which can be considered an abbreviation of $1(f_0, 1) \in R$, is considered positive even if $f_0 <_{F_0} 1$ and it would be called “negative” as a group element. In particular, the diagonal elements of the matrices displayed above are all positive.

Bergman then orders $M_2(R)$ as follows. Choose “an arbitrary order among the four ‘positions’ in a 2×2 matrix, and call a nonzero element of this module ‘positive’ if in the first position in which a nonzero coefficient occurs, the coefficient is in fact positive.” To be definite, we will choose the 1, 1 position to be first, the 2, 2 position to be second, and the off-diagonal positions ordered third and fourth in some fixed way. Now to order the matrix ring of polynomials, call an element M of $M_2(R[t])$ positive if it satisfies the following. Expand $M = M_0 + M_1t + \cdots + M_k t^k$, where each M_i belongs to $M_2(R)$. Let $n \geq 0$ be the least integer such that t^n has nonzero coefficient and say M is positive if and only if the first nonzero entry of M_n is positive in the ordered ring R .

Bergman points out that “the orderings of the positions can be the same for all n , but need not — there is a lot of freedom here.” But we will use the same ordering of the positions, as described, throughout.

Finally, define an ordering of $F_0 * F_1$ by declaring that $x < y$ if and only if $\rho(y) - \rho(x)$ is positive in $M_2(R[t])$.

4. Proof of [Theorem 1](#) and further properties of $<$

First we’ll argue that $(F_0 * F_1, <)$ is an ordered group. Clearly $<$ is a strict total ordering. To check invariance under multiplication, first note that every element of $\rho(F_0 * F_1)$ in $M_2(R[t])$, when expanded in powers of t , has constant term a diagonal matrix with positive entries. (See the proof of [Proposition 4](#) below to be more precise.) The product of such a matrix, on either side, with a positive matrix in $M_2(R[t])$ will again be positive. Thus, if $x, y, z \in F_0 * F_1$, one has $x < y \iff \rho(y) - \rho(x)$ is positive $\iff \rho(z)(\rho(y) - \rho(x)) = \rho(z y) - \rho(z x)$ is positive $\iff z x < z y$. Right invariance is proved similarly.

Next we will show that the ordering $<$ extends the given orderings $<_{F_0}$ and $<_{F_1}$. Suppose $f_0, f'_0 \in F_0$ and $f_0 <_{F_0} f'_0$. Then the difference between their images in $M_2(R[t])$ is the matrix $\begin{pmatrix} f'_0 - f_0 & * \\ 0 & 0 \end{pmatrix}$, and noting that $f'_0 - f_0$ is positive in R we conclude $f_0 < f'_0$. A similar argument shows that $<$ also extends $<_{F_1}$.

¹We understand an ordered ring $(R, <)$ to be an ordered group as an additive group, for which the positive cone $P = \{r \in R \mid 0 < r\}$ is also closed under multiplication.

This establishes the first part of [Theorem 1](#). To prove the second part, note that $\phi_0 \times \phi_1$ preserves the lexicographic orderings $<_F, <_G$ of $F_0 \times F_1$ and $G_0 \times G_1$, respectively. A homomorphism of groups naturally extends to a ring homomorphism of the integral group rings, and we see that if the group homomorphism preserves given orderings of the groups, then its extension takes “positive” elements of the group ring to positive elements. Then $\phi_0 \times \phi_1$ defines a ring homomorphism $R_F \rightarrow R_G$, where $R_F = \mathbb{Z}(F_0 \times F_1)$ and $R_G = \mathbb{Z}(G_0 \times G_1)$, which we will call $\phi_0 \times \phi_1$ again. This extends to a ring homomorphism $R_F[t] \rightarrow R_G[t]$, and further induces an additive homomorphism $M_2(R_F[t]) \rightarrow M_2(R_G[t])$, which we will again call $\phi_0 \times \phi_1$.

The diagram

$$\begin{array}{ccc} F_0 * F_1 & \xrightarrow{\rho} & M_2(R_F[t]) \\ \phi_0 * \phi_1 \downarrow & & \phi_0 \times \phi_1 \downarrow \\ G_0 * G_1 & \xrightarrow{\rho} & M_2(R_G[t]) \end{array}$$

is commutative (we have used the same symbol ρ for different maps, but defined analogously), and as already mentioned, $\phi_0 \times \phi_1$ takes positive matrix entries to positive matrix entries. We now argue that $\phi_0 * \phi_1$ is order-preserving, relative to $<_F, <_G$. Suppose $x, y \in F_0 * F_1$ and $x <_F y$. Then $\rho(y) - \rho(x)$ is positive, and therefore $\phi_0 \times \phi_1(\rho(y) - \rho(x))$ is positive in $M_2(R_G[t])$. But $\phi_0 \times \phi_1(\rho(y) - \rho(x)) = \phi_0 \times \phi_1(\rho(y)) - \phi_0 \times \phi_1(\rho(x)) = \rho(\phi_0 * \phi_1(y)) - \rho(\phi_0 * \phi_1(x))$, and since this is positive, we conclude that $\phi_0 * \phi_1(x) <_G \phi_0 * \phi_1(y)$. □

Corollary 3. *If $(F, <_F)$ and $(G, <_G)$ are ordered groups, then the ordered group $(F * G, <_*) := \mathfrak{F}((F, <_F), (G, <_G))$ has the properties that $<_*$ extends the orderings of F and G , and for any automorphisms $\phi : F \rightarrow F$ and $\psi : G \rightarrow G$ which preserve the given orderings, the automorphism $\phi * \psi : F * G \rightarrow F * G$ preserves the ordering $<_*$.*

Following the terminology used in [[Botto Mura and Rhemtulla 1977](#)], we will call a homomorphism $\phi : F \rightarrow G$ of ordered groups $(F, <_F)$ and $(G, <_G)$ an *order-homomorphism* (relative to the given orderings) if $x \leq_F y$ implies $\phi(x) \leq_G \phi(y)$ for all $x, y \in F$. Note that order-preserving homomorphisms are order-homomorphisms, and that order-homomorphisms need not be injective. Indeed, the order-preserving homomorphisms are exactly the order-homomorphisms which are injective. For example, using the lexicographic ordering of the direct product, the inclusions $F \rightarrow F \times G$ and $G \rightarrow F \times G$ are order-preserving, while the projection $F \times G \rightarrow F$ is an order-homomorphism. But the projection $F \times G \rightarrow G$ will not be an order-homomorphism, if the groups are nontrivial.

We’ll see that our construction of $<_*$ has similar properties. First note that [Theorem 1\(1\)](#) implies that the natural inclusion homomorphisms $F \rightarrow F * G$ and

$G \rightarrow F * G$ are order-preserving. There are also canonical maps $F * G \rightarrow F$, obtained by killing elements of G , and similarly $F * G \rightarrow G$. They combine to define a canonical homomorphism $\alpha : F * G \rightarrow F \times G$. Specifically, if $f_1 g_1 f_2 \cdots f_k g_k$ is an element of $F * G$, with $f_i \in F$ and $g_i \in G$, then $\alpha(f_1 g_1 f_2 \cdots f_k g_k) = (f_1 \cdots f_k, g_1 \cdots g_k)$.

Proposition 4. *Suppose that $(F, <_F)$ and $(G, <_G)$ are ordered groups. Then the canonical homomorphism $\alpha : F * G \rightarrow F \times G$ is an order-homomorphism, relative to the lexicographic ordering of $F \times G$ and the ordering $<$ for $F * G$.*

Proof. If $x \in F * G$ has image $\alpha(x) = (f, g) \in F \times G$, we observe that its image under the representation $\rho : F * G \rightarrow M_2(R[t])$ may be written

$$\rho(x) = \begin{pmatrix} f & 0 \\ 0 & g \end{pmatrix} + \text{terms of positive degree.}$$

The conclusion follows from our convention for ordering $M_2(R[t])$. \square

A subset $C \subset G$ of an ordered group $(G, <_G)$ is said to be *convex* if the inequalities $c <_G g <_G c'$, with $c, c' \in C$ imply that $g \in C$. For example, it is easy to see that if $(F, <_F)$ and $(G, <_G)$ are ordered groups and $\phi : F \rightarrow G$ is an order-homomorphism, then the kernel K of ϕ is a convex subgroup of F .

Corollary 5. *The kernel of the homomorphism $\alpha : F * G \rightarrow F \times G$ is convex, relative to the ordering $<$ of $F * G$.*

The kernel of $\alpha : F * G \rightarrow F \times G$ is known to be a free subgroup of $F * G$, freely generated by commutators of the form $f g f^{-1} g^{-1}$, where $1 \neq f \in F$ and $1 \neq g \in G$.

Corollary 6. *If $F * G$ is ordered by $<$, the canonical homomorphism $F * G \rightarrow F$ is an order-homomorphism, but $F * G \rightarrow G$ will not be an order-homomorphism, if the groups are nontrivial.*

Indeed, if $f <_F f'$ in F while $g' <_G g$ in G , we have, as elements of $F * G$, the inequality $f g < f' g'$. If the canonical map $F * G \rightarrow G$ were an order-homomorphism, we'd conclude $g <_G g'$, a contradiction. The asymmetry exposed by this corollary cannot be corrected, as the following observation shows. We note that, by the same proof, the proposition also applies to direct products.

Proposition 7. *If F and G are nontrivial ordered groups, then there is no ordering of $F * G$ for which both of the canonical homomorphisms $F * G \rightarrow F$ and $F * G \rightarrow G$ are order-homomorphisms.*

Proof. As above, choose $f, f' \in F$ and $g, g' \in G$ such that $f <_F f'$ and $g' <_G g$. Suppose $<$ is an ordering of $F * G$ for which the canonical homomorphisms $F * G \rightarrow F$ and $F * G \rightarrow G$ are order-homomorphisms, and compare $f g$ with $f' g'$. If $f g < f' g'$, then applying the map $F * G \rightarrow F$ implies that $f \leq_F f'$, a contradiction. Similarly, $f g > f' g'$ implies the contradiction $g \geq_G g'$. \square

5. Structure as a tensor category

Recall that \mathcal{C} denotes the category of ordered groups and order-preserving homomorphisms, and that $\mathfrak{F} : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ is a bifunctor. Let us rename \mathfrak{F} as follows, for ordered groups $(F_0, <_{F_0})$ and $(F_1, <_{F_1})$:

$$(F_0, <_{F_0}) \otimes (F_1, <_{F_1}) := \mathfrak{F}((F_0, <_{F_0}), (F_1, <_{F_1})) = (F_0 * F_1, <)$$

It is well known that the category of groups under free product is a tensor category, with unit the trivial group; see [Mac Lane 1998, p. 161], or the definition given in [Wikipedia 2017]. I am grateful to Christian Kassel for suggesting the following to me.

Theorem 8. *With the bifunctor \otimes the category \mathcal{C} is a tensor category, in other words, a monoidal category.*

For ordered groups $(F_0, <_{F_0}), (F_1, <_{F_1}), (F_2, <_{F_2})$, we have the isomorphism of groups

$$F_0 * (F_1 * F_2) \cong (F_0 * F_1) * F_2.$$

We need to check that the orderings constructed on both sides of this equivalence are the same under the isomorphism, in other words the isomorphism is order-preserving. But this follows from the observation that the lexicographic orderings on the direct products $F_0 \times (F_1 \times F_2)$ and $(F_0 \times F_1) \times F_2$, used in the respective orderings of $F_0 * (F_1 * F_2)$ and $(F_0 * F_1) * F_2$, both reduce to the lexicographic ordering of triples.

Similarly, the coherence relations involved in tensor categories follow from the observation that for ordered groups $(F_i, <_{F_i}), 0 \leq i \leq 3$, our orderings of the groups

$$(F_0 * F_1) * (F_2 * F_3), \quad (F_0 * (F_1 * F_2)) * F_3, \quad F_0 * ((F_1 * F_2) * F_3), \\ (F_0 * F_1) * (F_2 * F_3), \quad \text{and} \quad F_0 * (F_1 * (F_2 * F_3))$$

are identical (under their natural isomorphisms).

6. An application to braid groups

The original motivation for this study is the following application to the theory of braids. The braid group B_n acts by automorphisms on the free group \mathbb{F}_n , as observed by Artin [1925; 1947]. Free groups are orderable, and we may call a braid “order-preserving” if its image under the (faithful) Artin representation $B_n \rightarrow \text{Aut}(\mathbb{F}_n)$ preserves *some* ordering of \mathbb{F}_n ; see [Kin and Rolfsen 2016]. In that paper it is noted that a braid is order-preserving if and only if the complement of the link in S^3 consisting of the braid’s closure, plus the braid axis, has orderable fundamental group. It is used to show, for example, that of the two minimal volume orientable




Figure 1. $\alpha \in B_m$ (left), $\beta \in B_n$ (center), $\alpha \otimes \beta \in B_{m+n}$ (right).

hyperbolic 2-cusped 3-manifolds, one has an orderable fundamental group, while the group of the other is not orderable (although it is left-orderable).

Multiplication of braids is by concatenation, and the product of two order-preserving braids need not be order-preserving, as observed in [Kin and Rolfsen 2016]. There is also a tensor product operation $\otimes : B_m \times B_n \rightarrow B_{m+n}$ which forms an $m+n$ strand braid $\alpha \otimes \beta$ from an m -braid α and an n -braid β by placing them side by side with no crossing between the strands of α and those of β , as in Figure 1. See for example [Kassel and Turaev 2008, p. 69].

It is easy to see from the definition of Artin's representation that the automorphism of $\mathbb{F}_{m+n} \cong \mathbb{F}_m * \mathbb{F}_n$ corresponding to $\alpha \otimes \beta$ is just the free product of the automorphisms corresponding to α and β .

Corollary 9. *The tensor product $\alpha \otimes \beta$ of braids is order-preserving if and only if both α and β are order-preserving braids.*

Proof. One direction follows from Corollary 3. For if α and β preserve some orderings of \mathbb{F}_m and \mathbb{F}_n respectively, then $\alpha \otimes \beta$ preserves the corresponding ordering $<$ of $\mathbb{F}_m * \mathbb{F}_n \cong \mathbb{F}_{m+n}$. On the other hand, suppose $\alpha \otimes \beta$ preserves an ordering of $\mathbb{F}_{m+n} \cong \mathbb{F}_m * \mathbb{F}_n$. Considering \mathbb{F}_m and \mathbb{F}_n as the natural subgroups of $\mathbb{F}_m * \mathbb{F}_n$, we see that the action of $\alpha \otimes \beta$ leaves each of these subgroups invariant. Therefore the ordering of \mathbb{F}_{m+n} preserved by $\alpha \otimes \beta$ restricts to each of the subgroups making the action of the braids α and β order-preserving. \square

We note the multiple use of the tensor product symbol. Indeed, let us say that the ordered free group $(\mathbb{F}_n, <)$ represents the braid $\beta \in B_n$ if the automorphism of \mathbb{F}_n corresponding to β under the Artin representation preserves the ordering $<$. We have observed the following.

Proposition 10. *If $(\mathbb{F}_m, <)$ represents $\alpha \in B_m$ and $(\mathbb{F}_n, <')$ represents $\beta \in B_n$, then $(\mathbb{F}_m, <) \otimes (\mathbb{F}_n, <')$ represents $\alpha \otimes \beta \in B_{m+n}$.*

7. Continuity

The goal of this section is to establish that our construction is continuous in an appropriate sense. If $O(G)$ denotes the set of all (two-sided invariant) orderings of the group G , there is a natural topology on $O(G)$, defined below. Given orderable groups F and G , the construction defined in Section 3 can be considered a function

whose input is a pair of orderings $<_F$ and $<_G$ and the output is an ordering $<$ of $F * G$, in other words a function

$$O(F) \times O(G) \rightarrow O(F * G).$$

We'll see that it is both continuous and injective.

7.1. The space of orderings. The set of orderings $O(G)$ of the group G is endowed with a natural topology, as detailed by Sikora [2004]. See also [Dabkowska et al. 2007; Navas 2010]. Consider a specific ordering $<_G$ of G , and choose a *finite* number of inequalities among elements of G which are satisfied using $<_G$. Then a basic neighborhood of $<_G$ consists of all orderings of G for which all those inequalities remain true. Neighborhoods of this type form a basis for the topology we are considering. Equivalently, a neighborhood of $<_G$ is defined by choosing some finite set of elements of G which are positive (greater than the identity) using $<_G$. Then take the neighborhood to consist of all orderings of G under which that finite set remains positive.

It is known, and not difficult to show, that $O(G)$ is compact and totally disconnected. An isolated point of $O(G)$ is an ordering which is “finitely determined” in the sense that it is the only ordering of G for which some finite set of inequalities holds. Sikora [2004] showed that for $n \geq 2$, $O(\mathbb{Z}^n)$ has no isolated points, and is homeomorphic with the Cantor set. Whether $O(\mathbb{F}_n)$ has isolated points, for the free group \mathbb{F}_n , $n \geq 2$, is an open question at the time of writing.

7.2. Continuity of lexicographic ordering of direct products. As a warmup to our main result, we consider the lexicographic ordering of direct products $F \times G$ of ordered groups, as discussed in Section 3 (similar results would hold for the reverse lexicographic ordering). It may be considered a function

$$\mathfrak{L} : O(F) \times O(G) \rightarrow O(F \times G).$$

Proposition 11. \mathfrak{L} is continuous and injective.

Proof. We may assume both F and G are nontrivial groups; otherwise there is nothing to prove. For injectivity, suppose $<_F$ and $<'_F$ are orderings of F and that $<_G$ and $<'_G$ are orderings of G . Consider $< = \mathfrak{L}(<_F, <_G)$ and $<' = \mathfrak{L}(<'_F, <'_G)$. If $<_F$ and $<'_F$ are distinct, there must be an element $f \in F$ with $1 <_F f$ but $f <'_F 1$. Then we have, for any $g \in G$, that $1 < (f, g)$ and $(f, g) <' 1$. It follows that $<$ and $<'$ are distinct. Similarly, if $<_G$ and $<'_G$ are different, then one can find an element $(1, g) \in F * G$ with $(1, g)$ having different signs relative to the orderings $<$ and $<'$. This establishes injectivity.

To establish continuity, note that a basic neighborhood $\mathfrak{N}_<$ of $<$ in $O(F \times G)$

is defined by choosing some finite set of positive elements:

$$(f_1, g_1), \dots, (f_k, g_k), (1, g_{k+1}), \dots, (1, g_{k+l}).$$

Here we have

$$1 <_F f_1, \dots, 1 <_F f_k \quad \text{and} \quad 1 <_G g_{k+1}, \dots, 1 <_G g_{k+l},$$

whereas some of the list g_1, \dots, g_k may be negative in the ordering $<_G$. Possibly $k = 0$ or $l = 0$.

Continuity will be established if we can find neighborhoods $\mathfrak{N}_{<_F}$ of $<_F$ in $O(F)$ and $\mathfrak{N}_{<_G}$ of $<_G$ in $O(G)$ so that $\mathfrak{L}(\mathfrak{N}_{<_F} \times \mathfrak{N}_{<_G}) \subset \mathfrak{N}_{<}$. But this is straightforward: take $\mathfrak{N}_{<_F}$ to be the set of all orderings of F for which f_1, \dots, f_k are positive, and $\mathfrak{N}_{<_G}$ the set of all orderings of G under which g_{k+1}, \dots, g_{k+l} are positive. \square

7.3. Continuity of the ordering of free products. Recalling the construction in Section 3, we defined a function of ordered groups:

$$\mathfrak{F}((F, <_F), (G, <_G)) = (F * G, <).$$

By abuse of notation, if F and G are fixed, but orderings thereof are variable, we may write

$$\mathfrak{F}(<_F, <_G) = <.$$

Then we have a function of spaces of orderings:

$$\mathfrak{F} : O(F) \times O(G) \rightarrow O(F * G).$$

Theorem 12. \mathfrak{F} is continuous and injective.

Proof. One may prove injectivity as in Proposition 11; we leave the details to the reader. Note also that we proved continuity of the map \mathfrak{L} by showing that any finite set of inequalities in $F \times G$ would be implied (under \mathfrak{L}) by finitely many inequalities in F and in G .

We will argue similarly in this case; we'll try to avoid excessive notation and sketch the ideas. Suppose $<_F$ and $<_G$ are given orderings of F and G , respectively, and that $< = \mathfrak{F}(<_F, <_G)$ is the corresponding ordering of the free product $F * G$. A neighborhood $\mathfrak{N}_{<}$ of $<$ in the space $O(F * G)$ consists of all orderings of $F * G$ for which all members of some finite set x_1, \dots, x_k of elements of $F * G$ are positive, where $1 < x_i$ for $i = 1, \dots, k$. But note that $1 < x_i$ is equivalent to the matrix $\rho(x_i) - \rho(1)$ being positive in $M_2(\mathbb{Z}(F \times G)[t])$, and this is positive if the first nonzero entry of that matrix, expanded in powers of t , is positive. That entry, an element of $\mathbb{Z}(F \times G)$, is positive if the coefficient of its greatest group element, say (f_i, g_i) , is a positive integer. But the condition that (f_i, g_i) is the greatest group element appearing in that entry is equivalent to a finite number of inequalities in $F \times G$, using the lexicographic ordering. This in turn, as in Proposition 11, is implied by a finite number of inequalities in F and G which are in particular

satisfied using the orderings $<_F$ and $<_G$. Using the open neighborhoods $\mathfrak{N}_{<_F}$ of $<_F$ and $\mathfrak{N}_{<_G}$ of $<_G$ defined by those inequalities, we see $\mathfrak{F}(\mathfrak{N}_{<_F}, \mathfrak{N}_{<_G}) \subset \mathfrak{N}_{<}$, which establishes continuity of \mathfrak{F} . \square

Suppose, in the procedure for defining $<$ in Section 3, one used some ordering of $F \times G$ other than the lexicographic one, but otherwise defined $<$ in the same way. This then defines a function $O(F \times G) \rightarrow O(F * G)$, which we will call \mathfrak{M} , short for matrix construction. The proof of Theorem 12 actually shows that \mathfrak{M} is continuous. Our specific construction \mathfrak{F} may therefore be considered a composite

$$O(F) \times O(G) \xrightarrow{\mathfrak{L}} O(F \times G) \xrightarrow{\mathfrak{M}} O(F * G)$$

of two continuous functions, both injective.

8. Free product of arbitrarily many ordered groups

We now consider an arbitrary collection of ordered groups. For convenience, we assume the groups are indexed by an ordinal number γ and denote the collection by $\{(F_\alpha, <_{F_\alpha})\}_{\alpha < \gamma}$. So far we have been considering the case $\gamma = 2$.

Theorem 13. *Let $\gamma \geq 2$ be an ordinal. Suppose $\{(F_\alpha, <_{F_\alpha})\}_{\alpha < \gamma}$ is a collection of ordered groups and let $F := *_{\alpha < \gamma} F_\alpha$ denote the free product. Then there is an ordering $<_F$ of F , so that $(F, <_F)$ is an ordered group, denoted*

$$\mathfrak{F}(\{(F_\alpha, <_{F_\alpha})\}_{\alpha < \gamma}) := (F, <_F),$$

and such that the following hold:

- (1) For each $\alpha < \gamma$ the restriction of $<_F$ to the natural subgroup F_α of F equals $<_{F_\alpha}$.
- (2) If $\{(G_\alpha, <_{G_\alpha})\}_{\alpha < \gamma}$ is another collection of ordered groups with $G := *_{\alpha < \gamma} G_\alpha$ and

$$(G, <_G) = \mathfrak{F}(\{(G_\alpha, <_{G_\alpha})\}_{\alpha < \gamma}),$$

then for any collection $\phi_\alpha : F_\alpha \rightarrow G_\alpha$ of homomorphisms defined for all $\alpha < \gamma$ and which are order-preserving, relative to $<_{F_\alpha}$ and $<_{G_\alpha}$, the free product homomorphism $*_{\alpha < \gamma} \phi_\alpha : F \rightarrow G$ is order-preserving, relative to $<_F$ and $<_G$.

Proof. We will define the ordering of F by induction, possibly transfinite. For that reason, we'll call the ordering $<_\gamma$ and only later call it $<_F$ also. The base for the induction, for $\gamma = 2$, is Theorem 1, taking $<_2$ to be the ordering $<$ defined there. For induction we may assume that orderings $<_\beta$ have been defined for all the groups $*_{\alpha < \beta} F_\alpha$ for all $1 < \beta < \gamma$, and that they satisfy (1) and (2) with β replacing γ . Note that $*_{\alpha < \beta} F_\alpha$ is naturally a subgroup of $*_{\alpha < \gamma} F_\alpha$. To facilitate the induction, we'll prove that in addition to properties (1) and (2) of the theorem, $<_\gamma$ further satisfies:

(3) If $1 < \beta < \gamma$ the restriction of the ordering $<_\gamma$ to $*_{\alpha < \beta} F_\alpha$ coincides with $<_\beta$.

Again, by [Theorem 1](#) this is satisfied for the base case $\gamma = 2$. To construct $<_\gamma$ we consider two cases.

Case 1: γ is a successor ordinal: $\gamma = \beta + 1$. Since $<_\beta$ is by hypothesis already defined, and noting that F can be naturally identified with $(*_{\alpha < \beta} F_\alpha) * F_\beta$, we use the functor \mathfrak{F} defined in the proof of [Theorem 1](#) and take

$$(F, <_\gamma) \cong ((*_{\alpha < \beta} F_\alpha) * F_\beta, <_\gamma) := \mathfrak{F}((*_{\alpha < \beta} F_\alpha), <_\beta), (F_\beta, <_\beta)).$$

Case 2: γ is a limit ordinal. Then the group $*_{\alpha < \gamma} F_\alpha$ is the union of its subgroups $*_{\alpha < \beta} F_\alpha$ with $\beta < \gamma$. Thus to compare two group elements x, y in $*_{\alpha < \gamma} F_\alpha$, choose $\beta < \gamma$ for which $x, y \in *_{\alpha < \beta} F_\alpha$ and define $x <_\gamma y$ if and only if $x <_\beta y$. By property (3), which may be assumed for ordinals less than γ , this does not depend on choice of β .

In either case, it is routine to verify that the ordering $<_\gamma$ (also called $<_F$) satisfies the conditions (1), (2) and (3). □

9. Left-ordered groups

An ordering $<$ of the elements of a group G is a left-ordering if for all $f, g, h \in G$ one has

$$g < h \implies fg < fh;$$

in this case we call $(G, <)$ a left-ordered group. It is much easier than for the ordered case to see that the free product of left-ordered groups is left-orderable. For left-ordered groups $(F, <_F)$ and $(G, <_G)$ consider the short exact sequence

$$1 \rightarrow K \rightarrow F * G \rightarrow F \times G \rightarrow 1,$$

where $F * G \rightarrow F \times G$ is the canonical homomorphism. The kernel K is a free group, which is orderable, and one can left-order $F \times G$, lexicographically. Since left-orderability (unlike orderability) is always preserved under extensions, we conclude that $F * G$ is left-orderable.

On the other hand, our construction of the ordering $<$ for the free product of ordered groups may be revised in a straightforward way to the left-ordered (or right-ordered) situation. One must be a bit careful. For a left-ordered group $(G, <)$ the group ring $\mathbb{Z}(G)$ is not, strictly speaking, an ordered ring by our definition. For example if we have $g, g', h \in G$ with $g < g'$ but $gh > g'h$ then the ring elements $g' - g$ and h are positive, whereas their product $g'h - gh$ is not positive. However the product in the other order, $hg' - hg$, is necessarily positive, and more generally a positive element of $\mathbb{Z}(G)$ multiplied on the left by a monomial with positive coefficient remains positive. This is enough to establish left-invariance of $<$ in the proof of [Theorem 1](#).

Therefore, we conclude that all the results above remain true if “ordered” is replaced by “left-ordered” throughout. In particular, the category of left-ordered groups and order-preserving homomorphisms is also a tensor category using our functorial construction.

10. Concluding remarks

The ordering we construct is by no means canonical; for example other choices of ordering the direct product, or the entries of matrices, can lead to a different ordering of the free product which satisfies the conditions of [Theorem 1](#), and even defines a tensor category structure. Indeed, [Corollary 6](#) reveals the asymmetry of the construction. In a real sense, the first group in the free product of two groups is treated preferentially in our construction. It could as well have been the reverse.

The argument given here does not extend to the larger category of ordered groups and order-homomorphisms (which are not necessarily injective) as some positive matrix entries may be mapped to zero under such a map. Extending our results to this category seems to be an open question.

As noted in [\[Bergman 1990\]](#), much of this can be done in the more general setting of ordered semigroups; see also [\[Johnson 1968\]](#). We leave such generalization for the interested reader to contemplate.

References

- [Artin 1925] E. Artin, “[Theorie der Zöpfe](#)”, *Abh. Math. Sem. Univ. Hamburg* **4**:1 (1925), 47–72. [MR](#) [Zbl](#)
- [Artin 1947] E. Artin, “[Theory of braids](#)”, *Ann. of Math. (2)* **48** (1947), 101–126. [MR](#) [Zbl](#)
- [Bergman 1990] G. M. Bergman, “[Ordering coproducts of groups and semigroups](#)”, *J. Algebra* **133**:2 (1990), 313–339. [MR](#) [Zbl](#)
- [Botto Mura and Rhemtulla 1977] R. Botto Mura and A. Rhemtulla, *Orderable groups*, Lecture Notes in Pure and Applied Mathematics **27**, Dekker, New York, 1977. [MR](#) [Zbl](#)
- [Chiswell 2012] I. M. Chiswell, “[Ordering graph products of groups](#)”, *Internat. J. Algebra Comput.* **22**:4 (2012), art. id. 1250037. [MR](#) [Zbl](#)
- [Chiswell 2014] I. M. Chiswell, “[Ordering free products of groups](#)”, *Math. Slovaca* **64**:3 (2014), 707–726. [MR](#) [Zbl](#)
- [Dabkowska et al. 2007] M. A. Dabkowska, M. K. Dabkowski, V. S. Harizanov, J. H. Przytycki, and M. A. Veve, “[Compactness of the space of left orders](#)”, *J. Knot Theory Ramifications* **16**:3 (2007), 257–266. [MR](#) [Zbl](#)
- [Holland and Medvedev 1994] W. C. Holland and N. Y. Medvedev, “[A very large class of small varieties of lattice-ordered groups](#)”, *Comm. Algebra* **22**:2 (1994), 551–578. [MR](#) [Zbl](#)
- [Johnson 1968] R. E. Johnson, “[Free products of ordered semigroups](#)”, *Proc. Amer. Math. Soc.* **19** (1968), 697–700. [MR](#) [Zbl](#)
- [Kassel and Turaev 2008] C. Kassel and V. Turaev, *Braid groups*, Graduate Texts in Mathematics **247**, Springer, 2008. [MR](#) [Zbl](#)

- [Kin and Rolfsen 2016] E. Kin and D. Rolfsen, “Braids, orderings and minimal volume cusped hyperbolic 3-manifolds”, preprint, 2016. [arXiv](#)
- [Mac Lane 1998] S. Mac Lane, *Categories for the working mathematician*, 2nd ed., Graduate Texts in Mathematics **5**, Springer, 1998. [MR](#) [Zbl](#)
- [Medvedev 1991] N. Y. Medvedev, “A remark on a paper by G. Révész”, *Czechoslovak Math. J.* **41**:1 (1991), 51. In Russian; correction to [Révész 1987]. [MR](#) [Zbl](#)
- [Navas 2010] A. Navas, “On the dynamics of (left) orderable groups”, *Ann. Inst. Fourier (Grenoble)* **60**:5 (2010), 1685–1740. [MR](#) [Zbl](#)
- [Passman 1977] D. S. Passman, *The algebraic structure of group rings*, Wiley, New York, 1977. [MR](#) [Zbl](#)
- [Révész 1987] G. Révész, “A simple proof of Vinogradov’s theorem on the orderability of the free product of o -groups”, *Czechoslovak Math. J.* **37**:2 (1987), 310–312. [MR](#) [Zbl](#)
- [Sikora 2004] A. S. Sikora, “Topology on the spaces of orderings of groups”, *Bull. London Math. Soc.* **36**:4 (2004), 519–526. [MR](#) [Zbl](#)
- [Vinogradov 1949] A. A. Vinogradov, “On the free product of ordered groups”, *Mat. Sbornik N.S.* **25(67)**:1 (1949), 163–168. In Russian; a translation by V. Lebed and A. Mortier is available at <https://arxiv.org/abs/1703.05781>. [MR](#) [Zbl](#)
- [Wikipedia 2017] “Monoidal category”, web page, 2017, available at https://en.wikipedia.org/wiki/Monoidal_category.

Received April 9, 2017. Revised September 26, 2017.

DALE ROLFSEN
MATHEMATICS DEPARTMENT
UNIVERSITY OF BRITISH COLUMBIA
VANCOUVER, BC
CANADA
rolfsen@math.ubc.ca

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 294 No. 1 May 2018

Three-dimensional Sol manifolds and complex Kleinian groups	1
WALDEMAR BARRERA, RENE GARCIA-LARA and JUAN NAVARRETE	
On periodic points of symplectomorphisms on surfaces	19
MARTA BATORÉO	
Mixing properties for hom-shifts and the distance between walks on associated graphs	41
NISHANT CHANDGOTIA and BRIAN MARCUS	
Simultaneous construction of hyperbolic isometries	71
MATT CLAY and CAGLAR UYANIK	
A local weighted Axler–Zheng theorem in \mathbb{C}^n	89
ŽELJKO ČUČKOVIĆ, SÖNMEZ ŞAHUTOĞLU and YUNUS E. ZEYTUNCU	
Monotonicity and radial symmetry results for Schrödinger systems with fractional diffusion	107
JING LI	
Moduli spaces of stable pairs	123
YINBANG LIN	
Spark deficient Gabor frames	159
ROMANOS-DIOGENES MALIKIOSIS	
Ordered groups as a tensor category	181
DALE ROLFSEN	
Multiplication of distributions and a nonlinear model in elastodynamics	195
C. O. R. SARRICO	
Some Ambrose- and Galloway-type theorems via Bakry–Émery and modified Ricci curvatures	213
HOMARE TADANO	
Irreducible decomposition for local representations of quantum Teichmüller space	233
JÉRÉMY TOULISSE	