

*Pacific
Journal of
Mathematics*

**THE PROJECTIVE LINEAR SUPERGROUP AND THE
SUSY-PRESERVING AUTOMORPHISMS OF $\mathbb{P}^{1|1}$**

RITA FIORESI AND STEPHEN D. KWOK

THE PROJECTIVE LINEAR SUPERGROUP AND THE SUSY-PRESERVING AUTOMORPHISMS OF $\mathbb{P}^{1|1}$

RITA FIORESI AND STEPHEN D. KWOK

The purpose of this paper is to describe the projective linear supergroup, its relation with the automorphisms of the projective superspace and to determine the supergroup of SUSY-preserving automorphisms of $\mathbb{P}^{1|1}$.

1. Introduction

The works of Manin [1988; 1991] and more recently of Witten et al. [Witten 2012; Donagi and Witten 2015] have drawn attention to projective supergeometry and more specifically to SUSY curves and their moduli superspaces.

In this paper we study the automorphisms of the projective superspace $\mathbb{P}^{m|n}$ and its SUSY-preserving subsupergroup. We start by defining the projective linear supergroup $\mathrm{PGL}_{m|n}$, using the functor of points formalism, and then we show that this supergroup functor is indeed representable, that is, it is the functor of points of a superscheme. We achieve this by realizing $\mathrm{PGL}_{m|n}$ as a closed subsupergroup scheme of $\mathrm{GL}_{m^2+n^2|2mn}$, mimicking the ordinary procedure.

In relating this supergroup scheme to the automorphism supergroup of $\mathbb{P}^{m|n}$ we encounter a difficulty, not present in the ordinary setting, namely the fact that the Picard group of the projective superspace is not known in general and involves some difficulties. This is a consequence of the fact that the supergroup of automorphisms of the projective superspace is larger than $\mathrm{PGL}_{m|n}$ for $n > 1$. Nevertheless, going to the special case of $n = 1$, we are able to give the projective linear supergroup quite explicitly and to prove it coincides with the automorphisms of the projective superspace.

The question of singling out the SUSY-preserving automorphisms inside this supergroup was already settled over the complex field by Manin [1991] and Witten [2012]; we extend their considerations to an arbitrary algebraically closed field k , $\mathrm{char}(k) \neq 2$, and provide some extra details of their proofs.

The organization of this paper is as follows. In Section 2 we start by reviewing some generally known facts on the projective superspace and its functor of points to establish our notation. We then discuss line bundles and projective morphisms,

MSC2010: 14L30, 58A50.

Keywords: supergeometry, Lie theory, algebraic geometry.

proving, in [Proposition 2.3](#), that the Picard group of $\mathbb{P}^{m|1}$ is \mathbb{Z} . To our knowledge this result is new and gives insight into projective supergeometry. In [Section 3](#) we define the projective linear supergroup in terms of functor of points and we prove its representability by realizing it as a closed subsuperscheme of the general linear supergroup. Then, in [Section 4](#) we prove that the projective linear supergroup is the supergroup of automorphisms of the projective superspace in the case of one odd dimension. Though the approach in both [Sections 3](#) and [4](#) closely resembles the ordinary one, the results are novel in the supergeometric context. In [Section 5](#), we use the machinery developed previously to prove that the subsupergroup of $\text{Aut}(\mathbb{P}^{1|1})$ of SUSY-preserving automorphisms of $\mathbb{P}^{1|1}$ consists precisely of the irreducible component $(\text{SpO}_{2|1})^0$ of the $2|1$ -symplectic-orthogonal supergroup $\text{SpO}_{2|1}$ containing the identity. This section is a generalization of the claims made in [\[Manin 1991\]](#) regarding complex supergeometry and provides proofs for such claims for a generic algebraically closed field.

2. The projective superspace $\mathbb{P}^{m|n}$

In this section we want to recall different, but equivalent definitions of projective superspace and we describe the line bundles on it. For all of our notation and main definitions of supergeometry, we refer the reader to [\[Manin 1988; Deligne and Morgan 1999; Carmeli et al. 2011\]](#).

Let k be our ground ring.

We recall that, by definition, the functor of points of a superscheme $X = (|X|, \mathcal{O}_X)$ is the functor

$$X : (\text{sschemes})^o \rightarrow (\text{sets}), \quad X(S) = \text{Hom}_{(\text{sschemes})}(S, X), \quad X(\phi)(f) = f \circ \phi,$$

where (sschemes) denotes the category of superschemes (it is customary to use the same letter for X and its functor of points). Equivalently (see [\[Carmeli et al. 2011, Chapter 10\]](#)), we can view the functor of points of X as $X : (\text{salg}) \rightarrow (\text{sets})$:

$$X(R) = \text{Hom}_{(\text{sschemes})}(\underline{\text{Spec}} R, X), \quad X(\phi)(f) = f \circ \underline{\text{Spec}}(\phi),$$

where (salg) denotes the category of superalgebras (over k), (we shall use the same letter for this functor also). In fact the functor of points of a superscheme is determined by its behavior on the affine superscheme subcategory, which in turn is equivalent to the category of superalgebras; see [\[Carmeli et al. 2011, Chapter 10, Theorem 10.2.5\]](#). If $X = \underline{\text{Spec}} \mathcal{O}(X)$, that is, X is affine, we have that

$$X(R) = \text{Hom}_{(\text{sschemes})}(\underline{\text{Spec}} R, X) = \text{Hom}_{(\text{salg})}(\mathcal{O}(X), R),$$

where $\mathcal{O}(X)$ denotes the superalgebra of global sections of the sheaf of superalgebras \mathcal{O}_X . We say that the $X(R)$ are the R -points of the superscheme X .

The algebraic superscheme $\mathbb{P}^{m|n}$ is defined as the patching of the $m + 1$ affine superspaces $U_i = \underline{\text{Spec}} \mathcal{O}(U_i)$, with $\mathcal{O}(U_i) = \underline{\text{Spec}} k[x_0^i, \dots, \hat{x}_i^i, \dots, x_m^i, \xi_1^i, \dots, \xi_n^i]$ through the change of charts:

$$(1) \quad \begin{aligned} \phi_{ij} : \mathcal{O}(U_j)[(x_i^j)^{-1}] &\mapsto \mathcal{O}(U_i)[(x_j^i)^{-1}] \\ x_k^j &\mapsto x_k^i/x_j^i \\ x_j^j &\mapsto 1/x_j^i \\ \xi_k^j &\mapsto \xi_k^i/x_j^i, \end{aligned}$$

(where as usual \hat{x}_i^i means that we are omitting the indeterminate x_i^i). Notice that $\mathcal{O}(U_j)[(x_i^j)^{-1}]$ is the superalgebra representing the open subscheme $U_j \cap U_i$ of U_j (and similarly for $\mathcal{O}(U_i)[(x_j^i)^{-1}]$).

Proposition 2.1. *The R -points of $\mathbb{P}^{m|n}$, $R \in (\text{salg})$ are given equivalently by:*

$$(i) \quad \begin{aligned} \mathbb{P}^{m|n}(R) &= \{\alpha : R^{m+1|n} \rightarrow L, R\text{-linear, surjective}\} / \sim, \\ \mathbb{P}^{m|n}(\psi) &: R^{m+1|n} \otimes_R T \rightarrow L \otimes_R T, \end{aligned}$$

where L is locally free of rank $1|0$, $\psi : R \rightarrow T$ and $\alpha : R^{m+1|n} \rightarrow L \sim \alpha' : R^{m+1|n} \rightarrow L'$ if and only if $\ker(\alpha) = \ker(\alpha')$ (or equivalently, $\alpha \sim \alpha'$ if they differ by an automorphism of L by multiplication of an element in R^\times).

$$(ii) \quad \begin{aligned} \mathbb{P}^{m|n}(R) &= \{\alpha : L \hookrightarrow R^{m+1|n} R\text{-linear, injective}\}, \\ \mathbb{P}^{m|n}(\psi) &: L \otimes_R T \rightarrow R^{m+1|n} \otimes_R T, \end{aligned}$$

where L is locally free of rank $1|0$.

Let $\mathcal{O}_S^{m+1|n} = \mathcal{O}_S \otimes k^{m+1|n}$. The S -points of $\mathbb{P}^{m|n}$, $S \in (\text{sschemes})$ are given equivalently by:

$$(a) \quad \begin{aligned} \mathbb{P}^{m|n}(S) &= \{\alpha : \mathcal{O}_S^{m+1|n} \rightarrow \mathcal{L}, \text{surjective}\} / \sim, \\ \mathbb{P}^{m|n}(\psi) &: (\psi^* \mathcal{O}_S)^{m+1|n} \rightarrow \psi^*(\mathcal{L}), \end{aligned}$$

where $\psi : T \rightarrow S$, \mathcal{L} is a line bundle on S (of rank $1|0$) and

$$\alpha : \mathcal{O}_S^{m+1|n} \rightarrow \mathcal{L} \sim \alpha' : \mathcal{O}_S^{m+1|n} \rightarrow \mathcal{L}'$$

if and only if $\ker(\alpha) = \ker(\alpha')$ (or equivalently, $\alpha \sim \alpha'$ if they differ by an automorphism of \mathcal{L} by multiplication of an element in \mathcal{O}_S^\times).

$$(b) \quad \begin{aligned} \mathbb{P}^{m|n}(S) &= \{\alpha : \mathcal{L} \hookrightarrow \mathcal{O}_S^{m+1|n}\}, \\ \mathbb{P}^{m|n}(\psi) &: \psi^* \mathcal{L} \rightarrow (\psi^* \mathcal{O}_S)^{m+1|n}. \end{aligned}$$

Proof. The proof relative to (i) and (a) works as in the ordinary setting and it is detailed in [Carmeli et al. 2011, Chapter 10]. The equivalence with (ii) and (b)

is immediate. The equivalence between (i) and (ii) is essentially the same as in the ordinary setting (see [Eisenbud and Harris 2000, Chapter III, Section 2, Proposition III-40, Corollary III-42]). \square

For every $A \in (\text{salg})$, we denote by $(\text{salg})_A$ the category of superalgebras over A . We will need to consider also $\mathbb{P}_A^{m|n}$, that is, the projective superspace over a base $A \in (\text{salg})$. This means that we are considering the superscheme obtained by patching the affine superspaces $U_i = A[x_j^i, \xi_k^i]$, $i, j = 0, \dots, m, j \neq i, k = 1, \dots, n$ as above. For example, in the second case in Proposition 2.1 each of the T -points, $T \in (\text{salg})_A$, is identified with a morphism $\alpha : L \rightarrow T^{m+1|n}$ of A -modules, where L and $T^{m+1|n}$ are T -modules which become A -modules via the map $\phi : A \rightarrow T$:

$$(2) \quad \mathbb{P}_A^{m|n}(T) = \text{Hom}_{(\text{sschemes})_A}(\underline{\text{Spec}} T, \mathbb{P}_A^{m+1|n}) = \{\alpha : L \hookrightarrow T^{m+1|n}\}.$$

Notice that the functor of points of $\mathbb{P}_A^{m|n}$ is defined on the category of A -superalgebras or equivalently on the category of A -superschemes (that is, superschemes equipped with a morphism to the superscheme $\underline{\text{Spec}} A$ and morphisms compatible with it).

We leave to the reader the generalization of the other cases of Proposition 2.1 since it is straightforward.

We end this section with some observations on line bundles and morphisms on $\mathbb{P}_A^{m|n}$. We start with a result completely similar to the ordinary counterpart, left to the reader as a simple exercise; see also [Carmeli et al. 2011, Chapter 9].

Proposition 2.2. *We have a bijective correspondence between the following:*

- (i) *The set of equivalence classes of $m+n+2$ -tuples $(L, s_0, \dots, s_m, \sigma_1, \dots, \sigma_n)$, where L is a line bundle on $\mathbb{P}_A^{m|n}$ globally generated by the global sections $s_0, \dots, s_m, \sigma_1, \dots, \sigma_n$ of L , under the relation*

$$(L, s_0, \dots, s_m, \sigma_1, \dots, \sigma_n) \sim (L, s'_0, \dots, s'_m, \sigma'_1, \dots, \sigma'_n)$$

if and only if there exists some $c \in \mathcal{O}(\mathbb{P}_A^{m|n})_0^$ such that $s'_i = cs_i$ and $\sigma'_i = c\sigma_i$ for all i .*

- (ii) *The set of A -morphisms $\mathbb{P}_A^{m|n} \rightarrow \mathbb{P}_A^{m|n}$.*

In the ordinary setting we have that a line bundle on \mathbb{P}_A^m is of the form $\mathcal{O}(n) \otimes \mathcal{L}$, where \mathcal{L} is a line bundle on $\underline{\text{Spec}} A$. This nontrivial fact is still true in supergeometry for $\mathbb{P}_A^{m|1}$, and it will turn out to be crucial in our treatment.

Proposition 2.3. *Every line bundle on $\mathbb{P}_A^{m|1}$ is isomorphic to $\mathcal{O}(n) \otimes \mathcal{L}$, where \mathcal{L} is a line bundle on $\underline{\text{Spec}} A$.*

Proof. A line bundle on $\mathbb{P}_A^{m|1}$ is determined once we know its transition functions, say $g_{ij} \in \mathcal{O}_{\mathbb{P}_A^{m|1}}(U_i \cup U_j)_0^*$, which are even. We then need to prove that any such set of transition functions is equivalent, up to a coboundary, to a set of transition

functions for a line bundle of the form $\mathcal{O}(n) \otimes \mathcal{L}$, for \mathcal{L} a line bundle on $\underline{\text{Spec}} A$. In other words we need to show

$$h_i|_{U_i \cap U_j} g_{ij} h_j^{-1}|_{U_i \cap U_j} = (x_j^i)^n, \quad h_i \in \mathcal{O}_{\mathbb{P}_A^{m|1}}(U_i)_0^*.$$

Notice that

$$\mathcal{O}_{\mathbb{P}_A^{m|1}}(U_p)^* = (A[x_k^p, \xi^p])_0^* = (A[\xi^p][x_k^p])_0^*, \quad p = i, j.$$

Since $\phi_{ij}(\xi^j) = \xi^i/x_j^i$, $\phi_{ij}(x_i^j) = 1/x_j^i$ and $\phi_{ij}(x_k^j) = x_k^i/x_j^i$, where ϕ_{ij} is the change of chart as in (1), we can view the restrictions of the h_p 's ($p = i, j$) to $U_i \cap U_j$, through this identification, as both belonging to $(A[\xi^i][x_j^i, (x_j^i)^{-1}])_0^*$. We now apply the classical result and obtain $h'_p \in (A[\xi^i][x_j^i, (x_j^i)^{-1}])_0^*$ such that

$$h'_i g_{ij} (h'_j)^{-1} = (x_j^i)^n.$$

The h'_p 's thus obtained are not yet the sections we want; since the odd dimension is one by hypothesis, the most general possible form for h'_j is

$$h'_j = a_0 + \alpha_0 \xi^i + \sum_K a_K x_K^i (x_j^i)^{-|K|} + \sum_L \alpha_L x_L^i (x_j^i)^{-|L|} \xi^i + \sum_k \beta_k (x_j^i)^{-k} \xi^i,$$

where K and L are multi-indices, $K = (k_1, \dots, k_r)$, $k_l \neq j$ ($r \in \mathbb{N}$) and $x_K^i := x_{k_1}^i \cdots x_{k_r}^i$ (similarly for L).

In order to eliminate the term $\alpha_0 \xi^i$ which is not well defined on U_j , we define:

$$h_i := (a_0 + \alpha_0 \xi^i) h'_i, \quad h_j := (a_0^{-1} - \alpha_0^{-2} \alpha_0 \xi^i) h'_j,$$

and this gives the required sections. \square

Notice that it was absolutely fundamental for our argument that there is only one odd dimension. This calculation will give us key information when we want to determine the automorphism supergroup of the projective linear supergroup.

3. The projective linear supergroup

In this section we want to define the supergroup functor of the projective linear supergroup and to show it is representable by producing an embedding of it as a closed subgroup into the general linear supergroup.

Let $\underline{\mathbf{M}}_{m|n}(R)$ denote the associative superalgebra of supermatrices of order $m|n$ by $m|n$ with entries in a commutative superalgebra R . More intrinsically, $\underline{\mathbf{M}}_{m|n}(R) = \underline{\text{End}}_R(R^{m|n})$.

Definition 3.1. The *automorphism supergroup of supermatrices* is the supergroup functor $\text{Aut}(\underline{\mathbf{M}}_{m|n}) : (\text{salg}) \rightarrow (\text{grps})$,

$$[\text{Aut}(\underline{\mathbf{M}}_{m|n})](R) :=$$

$$\{f : \underline{\mathbf{M}}_{m|n}(R) \rightarrow \underline{\mathbf{M}}_{m|n}(R) \mid f \text{ is an } R\text{-superalgebra automorphism}\}.$$

In analogy with the ordinary setting we also will call this supergroup functor the *projective linear supergroup* and denote it with $\text{PGL}_{m|n}$.

Since $\underline{\mathbf{M}}_{m|n}(R)$ is itself a free R -module of rank $M|N$, where $M = m^2 + n^2$ and $N = 2mn$, $\text{Aut}(\underline{\mathbf{M}}_{m|n})$ is a subfunctor of $\text{GL}_{M|N}$ in a natural way. We want to prove this is the functor of points of a closed subsuperscheme of $\text{GL}_{M|N}$. Before proceeding we need a lemma characterizing the morphisms of the superalgebra of supermatrices.

Lemma 3.2. (i) *An R -linear parity-preserving map $\psi : \underline{\mathbf{M}}_{m|n}(R) \rightarrow \underline{\mathbf{M}}_{m|n}(R)$ is a morphism of the superalgebra of supermatrices $\underline{\mathbf{M}}_{m|n}(R)$ if and only if*

- (a) $\psi(\text{id}) = \text{id}$;
- (b) $\psi(e_{ij})\psi(e_{kl}) = \delta_{kj}\psi(e_{il})$,

where e_{ij} are the elementary matrices in $\underline{\mathbf{M}}_{m|n}(R)$.

(ii) *If R is a local superalgebra, all of the automorphisms of the superalgebra $\underline{\mathbf{M}}_{m|n}(R)$ are of the form*

$$\underline{\mathbf{M}}_{m|n}(R) \rightarrow \underline{\mathbf{M}}_{m|n}(R), (T, X) \mapsto TXT^{-1},$$

for a suitable $T \in \text{GL}_{m|n}(R)$.

(iii) *$\text{Aut}(\underline{\mathbf{M}}_{m|n})$ is a closed subsuperscheme of $\text{GL}_{M|N} = \underline{\text{Spec}} k[x_{ij,kl}][d_1^{-1}, d_2^{-1}]$, $M = m^2 + n^2$ and $N = 2mn$, defined by the equations:*

$$(3) \quad \sum_k x_{ij,kk} = \delta_{ij}, \quad \sum_s x_{rs,ij}x_{st,kl} = \delta_{jk}x_{rt,il},$$

where $\text{GL}_{M|N}(R)$ is identified with the parity-preserving automorphisms of the free R -module $\underline{\mathbf{M}}_{m|n}(R)$.

Proof. (i) If ψ is an R -superalgebra endomorphism of $\underline{\mathbf{M}}_{m|n}(R)$ then the two relations are obviously satisfied and vice versa.

(ii) Now assume ψ is an automorphism of $\underline{\mathbf{M}}_{m|n}(R)$, R local, which satisfies the relations (a) and (b). We need to find $T \in \text{GL}_{m|n}(R)$ such that $\psi(e_{ij}) = Te_{ij}T^{-1}$. This is an application of super Morita theory (see [Kwok 2013]), however we shall recall the main idea to make this proof self-contained. By (a) and (b) we have

$$\sum \psi(e_{ii}) = \text{id}, \quad \psi(e_{ii})^2 = \psi(e_{ii}), \quad \psi(e_{ii})\psi(e_{jj}) = 0, \quad i \neq j,$$

hence we can write

$$R^{m|n} = \oplus \psi(e_{ii})R^{m|n}.$$

Since by (b), $\psi(e_{ji})\psi(e_{ii}) = \psi(e_{ji}) = \psi(e_{jj})\psi(e_{ji})$ we have $\psi(e_{ji}) : \psi(e_{ii})R^{m|n} \rightarrow \psi(e_{jj})R^{m|n}$ (recall that R is local so projective implies free). Hence there exists a basis $\{t_i\}$ of the free module $R^{m|n}$ such that

$$\psi(e_{ii})R^{m|n} = \text{span}_R\{t_i\}$$

and $\psi(e_{ji})t_i = t_j$. Let T be the matrix whose columns are the t_i 's, $T = \sum t_i \otimes e_i^*$, $T^{-1} = \sum e_i \otimes t_i^*$. It is then immediate to verify $\psi(e_{ij}) = T e_{ij} T^{-1}$.

(iii). This is immediate from (i). □

Let us view the multiplicative algebraic supergroup $\mathbb{G}_m^{1|0} : (\text{salg}) \rightarrow (\text{grps})$ as the following subsupergroup of $\text{GL}_{m|n}$:

$$\mathbb{G}_m^{1|0}(R) = \{aI \mid a \in R_0^*\} \subset \text{GL}_{m|n}(R).$$

(Here I denotes the identity matrix).

We do not specify the definition on the arrows whenever it is clear, as in this case.

Definition 3.3. We define the supergroup functor: $\widehat{\text{PGL}}_{m|n} : (\text{salg}) \rightarrow (\text{grps})$,

$$\widehat{\text{PGL}}_{m|n}(R) = \text{GL}_{m|n}(R) / \mathbb{G}_m^{1|0}(R),$$

and we call its sheafification (as customary) $\text{GL}_{m|n} / \mathbb{G}^{1|0}$.

We wish to show that $\text{GL}_{m|n} / \mathbb{G}^{1|0}$ is representable and coincides with the projective linear supergroup, that is, with the automorphism supergroup of supermatrices.

Definition 3.4. We say that a functor $F : (\text{salg}) \rightarrow (\text{grps})$ is *stalky* if for any superalgebra R , the natural map

$$\varinjlim_{f \notin \mathfrak{p}} F(R_f) \rightarrow F(R_{\mathfrak{p}})$$

is an isomorphism for any prime ideal $\mathfrak{p} \in R_0$.

The next two lemmas are standard and their proof is the same as in the ordinary case; see [Sun 2009].

Lemma 3.5. $\text{GL}_{m|n} / \mathbb{G}^{1|0}$ and $\text{Aut}(\underline{\mathbf{M}}_{m|n})$ are stalky.

Lemma 3.6. Let \mathcal{F}, \mathcal{G} be stalky Zariski sheaves $(\text{salg}) \rightarrow (\text{grps})$ and $\alpha : \mathcal{F} \rightarrow \mathcal{G}$ be a morphism. If $\alpha_R : \mathcal{F}(R) \rightarrow \mathcal{G}(R)$ is an isomorphism for all local superarrings R , then α is an isomorphism of sheaves.

Proposition 3.7. The supergroup functor $\text{GL}_{m|n} / \mathbb{G}^{1|0}$ is representable and is realized as the closed subsupergroup $\text{Aut}(\underline{\mathbf{M}}_{m|n})$ of $\text{GL}_{M|N}$ for $M = m^2 + n^2$ and $N = 2mn$.

Proof. We need to establish an isomorphism of sheaves between $\text{GL}_{m|n} / \mathbb{G}^{1|0}$ and a closed subsupergroup of $\text{GL}_{M|N}$. We will first give a morphism of sheaves and then show it is an isomorphism on local superalgebras; since $\text{GL}_{m|n} / \mathbb{G}^{1|0}$ is a stalky sheaf, this will be enough. We start by giving a morphism of presheaves $\widehat{\text{PGL}}_{m|n}$ and $\text{GL}_{M|N}$; since $\text{GL}_{M|N}$ is a sheaf then such a morphism will factor through the sheafification of $\widehat{\text{PGL}}_{m|n}$ thus giving us a sheaf morphism.

Consider the action of $GL_{M|N}$ on supermatrices $\underline{M}_{m|n}$, where $M = m^2 + n^2$, $N = 2mn$:

$$\phi : GL_{m|n}(R) \times \underline{M}_{m|n}(R) \rightarrow \underline{M}_{m|n}(R), \quad (T, X) \mapsto TXT^{-1}.$$

This clearly factors through $\mathbb{G}_m^{1|0}(R)$ and hence gives a well defined action ρ of $\widehat{PGL}_{m|n}$ and then in turn of $GL_{m|n}/\mathbb{G}^{1|0}$ (see comments at the beginning of the proof). Since $X \mapsto TXT^{-1}$ and $T \in (GL_{m|n}/\mathbb{G}^{1|0})(R)$ is a parity-preserving R -superalgebra morphism, it is immediate to verify we have a morphism of sheaves,

$$GL_{m|n}/\mathbb{G}^{1|0} \rightarrow \text{Aut}(\underline{M}_{m|n}).$$

By the first part of [Lemma 3.2](#), we know that $\text{Aut}(\underline{M}_{m|n})$ is represented by the closed subsuperscheme H of $GL_{M|N} = \underline{\text{Spec}} k[x_{ij,kl}][d_1^{-1}, d_2^{-1}]$ defined by the equations

$$(4) \quad \sum_k x_{ij,kk} = \delta_{ij}, \quad \sum_s x_{rs,ij} x_{st,kl} = \delta_{jk} x_{rt,il}.$$

(Here d_i denotes as usual the determinants of the diagonal blocks of indeterminates). We want to show that the group homomorphism $(GL_{m|n}/\mathbb{G}^{1|0})(R) \rightarrow [\text{Aut}(\underline{M}_{m|n})](R)$ is an isomorphism for R local. The automorphism $\psi \in GL_{M|N}(R)$ belongs to $H(R)$ if and only if its entries $\psi(e_{ij})_{kl}$ satisfy the above relations (4) (where in our convention $x_{ij,kl}$ corresponds to $\psi(e_{ij})_{kl}$). Hence by [Lemma 3.2](#) we have the result for R local. By [Lemmas 3.5](#) and [3.6](#), it is true for any superalgebra R and this concludes the proof. \square

Remark 3.8. The projective linear supergroup may also be obtained through the Chevalley supergroup recipe as detailed in [\[Fioresi and Gavarini 2011; 2012; 2013\]](#). It corresponds to the choice of the adjoint action of the Lie superalgebra $\mathfrak{sl}_{m|n}$. In fact one may readily check that the Lie superalgebra of $PGL_{m|n}$ is indeed $\mathfrak{sl}_{m|n}$ and $(PGL_{m|n})_0 = PGL_m \times PGL_n \times k^\times$.

4. The automorphisms of the projective superspace

We want to define the automorphism supergroup of the superscheme $\mathbb{P}^{m|n}$.

Definition 4.1. We define the supergroup functor of *automorphisms of the projective superspace*:

$$\text{Aut}(\mathbb{P}^{m|n})(A) := \text{Aut}_A(\mathbb{P}^{m|n} \times \underline{\text{Spec}} A) = \text{Aut}_A \mathbb{P}_A^{m|n}, \quad A \in (\text{salg}).$$

$\text{Aut}(\mathbb{P}^{m|n})$ is defined in an obvious way on the morphisms.

The equality in the definition is straightforward, noticing that we can identify the T -points of $\mathbb{P}^{m|n} \times \underline{\text{Spec}} A$ and of $\mathbb{P}_A^{m|n}$. In fact, a T -point of $\mathbb{P}^{m|n} \times \underline{\text{Spec}} A$ is a morphism $\phi : A \rightarrow T$ and a morphism $L \rightarrow T^{m|n}$ of A -modules via ϕ . This is exactly an element of $\mathbb{P}_A^{m|n}(T)$ and vice versa.

An automorphism $\psi \in \text{Aut}_A \mathbb{P}_A^{m|n}$ is a family of automorphisms ψ_T for all $T \in (\text{salg})_A$, which is functorial in T . The automorphism $\psi_T : \mathbb{P}_A^{m|n}(T) \rightarrow \mathbb{P}_A^{m|n}(T)$ must assign to a T -point of $\mathbb{P}_A^{m|n}(T)$, that is, a morphism $\alpha : L \rightarrow T^{m|n}$, another morphism $\alpha' : L' \rightarrow T^{m|n}$, where L and L' are projective rank $1|0$ T -modules, where the morphisms are interpreted as A -module morphisms. Similarly for the other characterizations of T -points as in [Proposition 2.1](#).

We are now ready to relate the supergroup scheme $\text{PGL}_{m|n}$ with the automorphisms of $\mathbb{P}^{m-1|n}$.

Proposition 4.2. *There is an embedding of supergroup functors*

$$\text{PGL}_{m|n} \hookrightarrow \text{Aut}(\mathbb{P}^{m-1|n}).$$

Proof. We first establish a morphism $\phi' : \text{GL}_{m|n} \rightarrow \text{Aut}(\mathbb{P}^{m-1|n})$. If $X \in \text{GL}_{m|n}(A)$ and $\alpha \in \mathbb{P}_A^{m-1|n}(T) = \{T^{m|n} \rightarrow L\} / \sim$, $\psi : A \rightarrow T$ we define

$$\phi'(X) = \alpha \circ \text{GL}_{m|n}(\psi)(X).$$

Clearly ϕ' factors through $\mathbb{G}_m(A)$. Since $\text{Aut}(\mathbb{P}^{m-1|1})$ is a sheaf, we have defined a morphism

$$\phi : \text{PGL}_{m|n} \rightarrow \text{Aut}(\mathbb{P}^{m-1|n}).$$

The injectivity is clear. □

Remark 4.3. In general we cannot expect to get an isomorphism between $\text{PGL}_{m|n}$ and $\text{Aut}(\mathbb{P}^{m-1|n})$ for $n > 1$ and this is because of the peculiarity of the odd elements. Let us see this in a simple example, $\mathbb{P}^{1|2}$. Consider the morphism $\phi \in \mathbb{P}_A^{1|2}$ given on the affine pieces $U_0 = \text{Spec } A[u, \mu_1, \mu_2]$ and $U_1 = A[v, \nu_1, \nu_2]$ by

$$\phi|_{U_0}(u, \mu_1, \mu_2) = (u + \mu_1\mu_2, \mu_1, \mu_2), \quad \phi|_{U_1}(v, \nu_1, \nu_2) = (v - \nu_1\nu_2, \nu_1, \nu_2).$$

As ϕ is invertible, $\phi \in \text{Aut}(\mathbb{P}^{m|n})(A)$, but it is not obtained through an element of $\text{PGL}_{2|2}(A)$. In fact the coefficient in $\phi|_{U_0}$ of $\mu_1\mu_2$ in an automorphism induced by a $\text{PGL}_{2|2}(A)$ transformation must be nilpotent. Hence $\phi \notin \text{PGL}_{2|2}(A)$.

We now want to show that we have an isomorphism between the projective linear supergroup and the automorphism of the super projective when $n = 1$. The argument we give follows along the lines of the calculation of $\text{Aut}(\mathbb{P}^n)$ given in [\[Hartshorne 1977, Chapter 2, Section 7\]](#).

Proposition 4.4. *We have an isomorphism of supergroup functors:*

$$\text{PGL}_{m+1|1} \cong \text{Aut}(\mathbb{P}^{m|1}).$$

In particular, $\text{Aut}(\mathbb{P}^{m|1})$ is a supergroup scheme.

Proof. [Proposition 4.2](#) gives us an embedding of supergroup functors $\text{PGL}_{m+1|1} \hookrightarrow \text{Aut}(\mathbb{P}^{m|1})$. Now let $f \in \text{Aut}(\mathbb{P}_A^{m|1})$ and let g be its inverse. We want to show $f \in \text{PGL}_{m+1|1}(A)$. The automorphism f induces the two line bundle morphisms

$f^*\mathcal{O}_A(1) \rightarrow \mathcal{O}_A(1)$ and $g^*\mathcal{O}_A(1) \rightarrow \mathcal{O}_A(1)$, where $\mathcal{O}_A(1) := p_1^*(\mathcal{O}(1))$, with $p_1 : \mathbb{P}_A^{m|1} \rightarrow \mathbb{P}^{m|1}$ being the natural projection. By Proposition 2.3, we know that $f^*\mathcal{O}_A(1) = \mathcal{O}(k) \otimes \mathcal{L}_f$ and $g^*\mathcal{O}_A(1) = \mathcal{O}(l) \otimes \mathcal{L}_g$. Let us choose a suitable open cover of A in which both \mathcal{L}_f and \mathcal{L}_g are trivial. By a common abuse of notation we shall still write A to denote the ring of global sections of an element of the open cover, so we in fact are replacing A with its localization. With such a choice we have $f^*\mathcal{O}_A(1) \cong \mathcal{O}_A(k)$ and $g^*\mathcal{O}_A(1) \cong \mathcal{O}_A(l)$. Since f and g are mutually inverse, we have

$$\mathcal{O}_A(1) = (f^* \circ g^*)(\mathcal{O}_A(1)) = f^*(g^*(\mathcal{O}_A(1))) = f^*(\mathcal{O}_A(l)) = \mathcal{O}_A(kl).$$

Hence $kl = 1$, whence $k = l = 1$, because for $k = l = -1$ we do not have global sections.

So $f^*(\mathcal{O}(1)) \cong \mathcal{O}(1)$, and choosing an isomorphism $F : f^*(\mathcal{O}(1)) \rightarrow \mathcal{O}(1)$ yields an isomorphism of the global sections $\Gamma(\mathbb{P}^m, f^*\mathcal{O}_A(1)) \cong \Gamma(\mathbb{P}^m, \mathcal{O}_A(1))$. By composing such an isomorphism with the natural isomorphism

$$f^* : \Gamma(\mathbb{P}^m, \mathcal{O}_A(1)) \rightarrow \Gamma(\mathbb{P}^m, f^*\mathcal{O}_A(1))$$

we obtain an A -linear automorphism,

$$T_F : \Gamma(\mathbb{P}^m, \mathcal{O}_A(1)) \rightarrow \Gamma(\mathbb{P}^m, \mathcal{O}_A(1)),$$

and identifying $\Gamma(\mathbb{P}^m, \mathcal{O}_A(1))$ with $A^{m+1|1}$ we see that $T_F \in \text{GL}_{m+1|1}(A)$. However, T_F depends on F . Suppose $G : f^*(\mathcal{O}(1)) \rightarrow \mathcal{O}(1)$ is another isomorphism, then $F^{-1} \circ G$ is an automorphism of $\mathcal{O}(1)$. Since $\underline{\text{Hom}}(L, L) = L^* \otimes L = \mathcal{O}$ for any line bundle L , we see that an automorphism of $\mathcal{O}(1)$ is the same thing as an invertible even function on $\mathbb{P}_A^{m|1}$, and F and G differ by composing with multiplication by such a function.

Therefore f determines T_F only up to multiplication by an invertible even function, i.e., f uniquely determines an element $T := [T_F]$ of $\text{PGL}_{m+1|1}(A)$.

Now in suitable coordinates we have that T induces (up to scalar multiplication) an automorphism of the \mathbb{Z} -graded superalgebra $A[z_0, \dots, z_m, \zeta]$. We leave to the reader the check that $\phi(T)$ is indeed f . □

5. The SUSY-preserving automorphisms of $\mathbb{P}_k^{1|1}$

In this section we want to consider those automorphisms of $\mathbb{P}_k^{1|1}$ which preserve its unique (up to isomorphism) SUSY structure. For all of the standard notation of supergeometry refer to [Carmeli et al. 2011].

Let k be our ground field, $\text{char}(k) \neq 2$, k algebraically closed. All algebraic supergroups discussed below will be algebraic supergroups over k .

We recall that if X is a smooth algebraic supervariety over k of dimension $1|1$,

we define a *SUSY structure* on X as a $0|1$ distribution \mathcal{D} on X such that the Frobenius map

$$\mathcal{D} \otimes \mathcal{D} \rightarrow TX/\mathcal{D}, \quad Y \otimes Z \mapsto [Y, Z] \bmod \mathcal{D}$$

is an isomorphism (see, for example, [Manin 1991] for the definition of a SUSY structure in the complex analytic case). If $X \rightarrow S$ is a smooth family of algebraic supervarieties of relative dimension $1|1$ over an algebraic k -supervariety S , then the notion of relative SUSY structure may be defined in the analogous way, as a relative distribution in the relative tangent sheaf TX/S . In this case we say that $X \rightarrow S$ is a *relative SUSY family*.

Our discussion is based on [Witten 2012].

We start by interpreting $\mathbb{P}_k^{1|1}$ as a homogeneous superspace. Let $\underline{k}^{2|1} = (k^2, \mathcal{O}_{\underline{k}^{2|1}})$ denote the affine superspace canonically associated to the k -super vector space $k^{2|1}$. Let us consider the action of the algebraic group \underline{k}^\times on $\underline{k}^{2|1} \setminus \{0\}$, given in the functor of points notation by

$$t \cdot (z_0, z_1, \zeta) = (tz_0, tz_1, t\zeta).$$

Consider the projection (as topological map)

$$\pi : k^2 \setminus \{0\} \rightarrow k^2 \setminus \{0\}/k^\times \cong \mathbb{P}^1.$$

Define the sheaf on the topological space \mathbb{P}_k^1 consisting of the \underline{k}^\times -invariant sections

$$\mathcal{F}(U) := \mathcal{O}_{\underline{k}^{2|1}}(\pi^{-1}(U))^{\underline{k}^\times}.$$

One can readily check that $(\mathbb{P}_k^1, \mathcal{F})$ is the superscheme $\mathbb{P}_k^{1|1}$ as defined in Section 2.

Let z_0, z_1, ζ be global coordinates on $\underline{k}^{2|1}$. We now consider the Euler vector field $E = z_0\partial_{z_0} + z_1\partial_{z_1} + \zeta\partial_\zeta$, which represents (in the chosen coordinates) the infinitesimal generator for the \underline{k}^\times action on $\underline{k}^{2|1} \setminus \{0\}$. Since E is everywhere nonsingular, it generates a trivial $1|0$ line bundle. As in the classical case, we have the Euler exact sequence of vector bundles on $\mathbb{P}_k^{1|1}$:

$$(5) \quad 0 \rightarrow \mathcal{O}^{1|0} \xrightarrow{i} \mathcal{O}(1) \otimes \text{Der}(S) \xrightarrow{j} T\mathbb{P}_k^{1|1} \rightarrow 0,$$

where i is the inclusion of the trivial $1|0$ line bundle $\langle E \rangle$ with global basis the Euler vector field. Here $\text{Der}(S)$ is the k -super vector space of k -linear derivations on $S := \underline{\text{Sym}}((k^{2|1})^*)$; it has as basis the derivations $\partial_{z_i}, \partial_\zeta$. Thus $\mathcal{O}(1) \otimes \text{Der}(S)$ is the sheaf whose sections on U are the linear vector fields on $\pi^{-1}(U)$. Any local section of $\mathcal{O}(1) \otimes \text{Der}(S)$ induces a corresponding local k -linear derivation on $\mathcal{O}_{\mathbb{P}_k^{1|1}}$ by restricting it to act on \underline{k}^\times -invariant functions; this defines j . Injectivity of i and the inclusion $\text{im}(i) \subseteq \ker(j)$ follow from the fact that E is nonsingular and the infinitesimal generator for the \underline{k}^\times -action; a standard calculation in the usual affine cells shows that $\ker(j) \subseteq \text{im}(i)$ and that j is surjective. Note that the sequence continues to remain exact on $\mathbb{P}_A^{1|1}$ after base change to any affine k -supervariety

$\underline{\text{Spec}}(A)$, with $T^{\mathbb{P}_k^{1|1}}$ replaced by the relative tangent bundle $T^{\mathbb{P}_A^{1|1}}/\underline{\text{Spec}}(A)$. We will denote the A -superalgebra $S \otimes_k A$ by S_A .

We now come to the SUSY structure.

Definition 5.1. Let $(X \rightarrow S, \mathcal{D})$ be a relative SUSY family. An S -automorphism $f : X \rightarrow X$ is *SUSY structure-preserving* (or simply *SUSY-preserving*) if and only if $(df_p)(\mathcal{D}_p) = \mathcal{D}_{f(p)}$ for any $p \in X$.

We will consider SUSY structures given by sections of $\mathcal{O}_A(1) \otimes \Omega_{S/A}$. Here $\Omega_{S/A}$ denotes the A -module of Kähler differentials on S_A , i.e., the A -dual to $\text{Der}(S_A)$; it has as basis the differentials $dz_i, d\zeta$. When we speak of the kernel of a section ω of $\mathcal{O}_A(1) \otimes \Omega_{S/A}$, we mean the kernel of ω when ω is interpreted as a morphism of sheaves of $\mathcal{O}_{\mathbb{P}_A^{1|1}}$ -modules from $\mathcal{O}_A(1) \otimes \text{Der}(S_A) \rightarrow \mathcal{O}_A(2)$.

Proposition 5.2. *Let $s := z_1 dz_0 - z_0 dz_1 - \zeta d\zeta$. Then the image of $\ker(s)$ under j is a SUSY structure on $\mathbb{P}_k^{1|1}$.*

Proof. In the affine open subsupervariety $U_1 := \{z_1 \neq 0\} \subset \mathbb{P}_k^{1|1}$, one calculates that the Euler vector field E and the linear vector field $\widehat{Z}_1 = \zeta \partial_{z_0} + z_1 \partial_\zeta$ lie in $\ker(s)$ and are linearly independent. At any point $p \in \mathbb{P}_k^{1|1}$, s induces a linear map of super vector spaces, $s_p : [\mathcal{O}(1) \otimes \text{Der}(S)]_p \rightarrow [\mathcal{O}(2)]_p$, on the fibers. It is clear that s is a basepoint-free section, hence s_p is always surjective. By linear algebra, $\ker(s_p)$ is $1|1$ dimensional and hence E_p and $\widehat{Z}_{1,p}$ span $\ker(s_p)$. By the super Nakayama’s lemma, E and \widehat{Z}_1 span $\ker(s)$ near p . Since p was arbitrary, E and \widehat{Z}_1 form a basis for $\ker(s)$ in U_1 .

One sees that $Z_1 := j(\widehat{Z}_1) = \partial_\eta + \eta \partial_w$, where $w = z_0/z_1$ and $\eta = \zeta/z_1$ are the usual affine coordinates in U_1 . $Z_1^2 = \partial_w$ and so Z_1 defines a SUSY structure in U_1 . A similar calculation with the linear vector field $\widehat{Z}_0 := -\zeta \partial_{z_1} + z_0 \partial_\zeta$ shows that $j(\ker(s))$ defines a SUSY structure on $U_0 = \{z_0 \neq 0\}$, hence the image of $\ker(s)$ under j defines a SUSY structure on $\mathbb{P}_k^{1|1}$. \square

We note that by the considerations of [Fioresi and Lledó 2015], this is the unique SUSY structure on $\mathbb{P}_k^{1|1}$, up to SUSY-isomorphism.

We now need the following proposition. The proof is completely similar to the one in [Fioresi and Lledó 2015, Proposition 5.2], however since the context here is more general, we include it for completeness.

Lemma 5.3. *Let A be an affine k -superalgebra. Let ω, ω' be two global sections of $\mathcal{O}_A(1) \otimes \Omega_{S/A}$ such that $\mathcal{D} := j(\ker(\omega))$ and $\mathcal{D}' := j(\ker(\omega'))$ are $0|1$ distributions on $\mathbb{P}_A^{1|1}$. Suppose $\mathcal{D} = \mathcal{D}'$. Then $\omega' = h\omega$ for some even invertible function h on $\mathbb{P}_A^{1|1}$.*

Proof. Let $p \in \mathbb{P}_A^{1|1}$ be a point. \mathcal{D} is locally a direct summand of $T^{\mathbb{P}_A^{1|1}}/\underline{\text{Spec}}(A)$, so we have a local splitting $\mathcal{D}|_U \oplus \mathcal{E} = (T^{\mathbb{P}_A^{1|1}}/\underline{\text{Spec}}(A))|_U$ in some neighborhood $U \ni p$. Via the Euler exact sequence (base changed to $\text{Spec}(A)$), we may lift $\mathcal{D}|_U$ (resp. \mathcal{E}) uniquely to a rank $1|1$ (resp. $2|0$) submodule $\widehat{\mathcal{D}}$ (resp. $\widehat{\mathcal{E}}$) of $[\mathcal{O}_A(1) \otimes \text{Der}(S_A)]|_U$

containing the $1|0$ line bundle $\langle E \rangle$ spanned by the Euler vector field, such that $\widehat{\mathcal{D}} \cap \widehat{\mathcal{E}} = \langle E \rangle$. We may therefore find local sections \widehat{Z} (resp. \widehat{X}) of $\widehat{\mathcal{D}}$ (resp. $\widehat{\mathcal{E}}$) such that \widehat{Z}, E (resp. \widehat{X}, E) form a basis for $\widehat{\mathcal{D}}$ (resp. $\widehat{\mathcal{E}}$). Note that the condition $\widehat{\mathcal{D}} \cap \widehat{\mathcal{E}} = \langle E \rangle$ implies $\widehat{X}, \widehat{Z}, E$ form a basis of $[\mathcal{O}_A(1) \otimes \text{Der}(S_A)]|_U$.

Viewing $\omega|_U$ as an $\mathcal{O}_{\mathbb{P}_A^{1|1}}$ -linear map from $[\mathcal{O}_A(1) \otimes \text{Der}(S_A)]|_U$ to $\mathcal{O}_A(2)|_U$, we have an induced linear map of super vector spaces,

$$\omega_p : (\mathcal{O}_A(1) \otimes \text{Der}(S_A))_p \rightarrow (\mathcal{O}_A(2))_p.$$

As $\ker(\omega_p) = \text{span}\{\widehat{Z}_p, E_p\}$, we see by linear algebra that ω_p is a surjection, and that $\omega_p(\widehat{X}_p)$ is a basis for $(\mathcal{O}_A(2))_p$; the analogous conclusion holds for ω'_p and $\omega'_p(\widehat{X}_p)$. Hence by the super Nakayama's lemma, $\omega(\widehat{X})$ is a basis for $\mathcal{O}_A(2)|_U$, and the same is true of $\omega'(\widehat{X})$ (shrinking U if necessary). Hence $\omega'(\widehat{X})/\omega(\widehat{X})$ is an invertible even function on U ; let us denote it by h .

To show that h is independent of the local complement \mathcal{E} and the choice of basis element \widehat{X} , suppose \mathcal{E}' is another local complement to \mathcal{D} on U , and let \widehat{X}', E be a basis of the lift $\widehat{\mathcal{E}}'$ of \mathcal{E}' . Then we have $\widehat{X}' = a\widehat{X} + bE + \alpha\widehat{Z}$ for some $a, b, \alpha \in \mathcal{O}_{\mathbb{P}_A^{1|1}}(U)$, with a, b even and α odd. As $\widehat{X}, E, \widehat{Z}$ and $\widehat{X}', E, \widehat{Z}'$ are both local bases for $\mathcal{O}_A(1) \otimes \text{Der}(S_A)$, a must be a unit.

Then we have

$$\omega'(\widehat{X}')/\omega(\widehat{X}') = \omega'(a\widehat{X} + bE + \alpha\widehat{Z})/\omega(a\widehat{X} + bE + \alpha\widehat{Z}) = \omega'(\widehat{X})/\omega(\widehat{X}),$$

since ω, ω' both annihilate E and \widehat{Z} . This proves that the expression $\omega'(\widehat{X})/\omega(\widehat{X})$ is independent of all choices and hence h is a well-defined function on all of $\mathbb{P}_A^{1|1}$. The equality $\omega' = h\omega$ clearly holds locally, and since h is now known to be globally defined, it holds globally. \square

Proposition 5.4. *Let f be an automorphism of $\mathbb{P}_A^{1|1}$. Then f preserves the SUSY structure defined by s if and only if for some (hence every) lift \tilde{f} of f to $\text{GL}_{2|1}(A)$, $\tilde{f}^*(s) = ts$ for some invertible function t .*

Proof. We begin by noting that $\text{GL}_{2|1}(A)$ preserves A_0^* -invariant open subsets of $\mathbb{A}_A^{2|1} \setminus \{0\}$, hence it acts naturally by pullback of functions on $\mathcal{O}_A(1) \otimes \text{Der}(S_A)$, where we interpret the latter as the sheaf assigning to any open subset $U \subseteq \mathbb{P}_A^{1|1}$ the linear vector fields on $\pi^{-1}(U) \subseteq \mathbb{A}_A^{2|1} \setminus \{0\}$.

The supersubgroup of invertible scalar matrices $\{cI : c \in A_0^*\}$ is central in $\text{GL}_{2|1}(A)$, hence this $\text{GL}_{2|1}(A)$ -action preserves the subalgebra of A_0^* -invariant functions on any A_0^* -invariant open subset of $\mathbb{A}_A^{2|1} \setminus \{0\}$. Hence we have an induced $\text{GL}_{2|1}(A)$ -action on the sheaf $\mathcal{O}_{\mathbb{P}_A^{1|1}}$. Clearly, invertible scalar matrices act trivially on $\mathcal{O}_{\mathbb{P}_A^{1|1}}$, thus the $\text{GL}_{2|1}(A)$ -action on $\mathcal{O}_{\mathbb{P}_A^{1|1}}$ factors through $\text{PGL}_{2|1}(A)$.

We see from the above that the action of $\text{GL}_{2|1}(A)$ on $\mathcal{O}_A(1) \otimes \text{Der}(S_A)$ by pullback of functions induces naturally a $\text{PGL}_{2|1}(A)$ -action on $\mathcal{O}_{\mathbb{P}_A^{1|1}}$, hence on

$T\mathbb{P}_A^{1|1}/\underline{\text{Spec}}(A)$, also given by pullback of functions. But this is precisely the $\text{PGL}_{2|1}(A)$ -action on $T\mathbb{P}_A^{1|1}/\underline{\text{Spec}}(A)$ induced by the action of $\text{PGL}_{2|1}(A)$ on $\mathbb{P}_A^{1|1}$ by automorphisms.

Since the sheaf morphism $j : \mathcal{O}_A(1) \otimes \text{Der}(S_A) \rightarrow T\mathbb{P}_A^{1|1}/\underline{\text{Spec}}(A)$ is just given by restricting a linear vector field to act on A_0^* -invariant functions, we see j is equivariant with respect to the $\text{GL}_{2|1}(A)$ - and $\text{PGL}_{2|1}(A)$ -actions previously defined.

We also have a $\text{GL}_{2|1}(A)$ -action on $\mathcal{O}_A(1) \otimes \Omega_{S/A}$ by the natural action on both factors, and for $\omega \in \Gamma(\mathcal{O}_A(1) \otimes \Omega_{S/A}) = \Gamma(\mathcal{O}_A(1)) \otimes \Omega_{S/A}$, we write $g^*(\omega)$ for $g \cdot \omega$.

Since the action of $\text{GL}_{2|1}(A)$ on $\mathcal{O}_A(1) \otimes \text{Der}(S_A)$ is the same as the natural action on the individual factors, and the $\text{GL}_{2|1}(A)$ -action on $\Omega_{S/A}$ is dual to that on $\text{Der}(S_A)$, it follows that the evaluation pairing

$$[\mathcal{O}_A(1) \otimes \text{Der}(S_A)] \otimes [\mathcal{O}_A(1) \otimes \Omega_{S/A}] \rightarrow \mathcal{O}_A(2)$$

is $\text{GL}_{2|1}(A)$ -equivariant, where $\mathcal{O}_A(2)$ is endowed with the natural $\text{GL}_{2|1}(A)$ -action.

From the preceding discussion, we see that f is SUSY-preserving if and only if $j[\ker(\omega)]_p = j[\ker(\tilde{f}^*(\omega))]_p$ for any point p .

We claim this is true if and only if $j[\ker(\omega)] = j[\ker(\tilde{f}^*(\omega))]$. One direction is clear. For the other, suppose $j[\ker(\omega)]_p = j[\ker(\tilde{f}^*(\omega))]_p$ for any point p . Then by the super Nakayama’s lemma $j[\ker(\omega)] = j[\ker(\tilde{f}^*(\omega))]$ in a neighborhood of p , hence globally. The claim then follows from [Lemma 5.3](#). □

In order to determine the supergroup of SUSY-preserving automorphisms of $\mathbb{P}_k^{1|1}$ we must discuss various other supergroups. We follow closely the discussion in [\[Manin 1991\]](#).

Definition 5.5. The 2|1-dimensional *conformal symplectic-orthogonal supergroup* $C_{2|1}$ is the subfunctor of $\text{GL}_{2|1}$ that preserves, up to multiplication by an even invertible constant, the split nondegenerate supersymplectic form on $k^{2|1}$ given by $(v, w) = v^t H w$, where

$$(6) \quad H := \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix},$$

and t denotes the super transpose of a matrix. More precisely, for every k -superalgebra A , $C_{2|1}$ is the functor $(\text{salg})_k \rightarrow (\text{grps})$ given by

$$(7) \quad C_{2|1}(A) := \{B \in \text{GL}_{2|1}(A) : B^t H B = Z(B) H\},$$

where $Z : \text{GL}_{2|1} \rightarrow \mathbb{G}_m^{1|0}$ is a fixed homomorphism.

The 2|1-dimensional *projective conformal symplectic-orthogonal supergroup* $\text{PC}_{2|1}$ is the image of $C_{2|1}$ in $\text{PGL}_{2|1}$, i.e, it is the sheafification of the group-valued functor $A \rightarrow C_{2|1}(A)/\{aI : a \in A_0^*\}$.

Proposition 5.6. $C_{2|1}$ and $PC_{2|1}$ are representable.

Proof. Taking the Berezinian of both sides of (7), one sees that $Z(B) = \text{Ber}(B)^2$. Thus, given

$$B = \begin{pmatrix} a & b & \alpha \\ c & d & \beta \\ \gamma & \delta & e \end{pmatrix} \in \text{GL}_{2|1}(A),$$

a direct calculation shows that B satisfies (7) if and only if the following equations hold: $e^2 + 2\alpha\beta = \text{Ber}(B)^2$, $a\beta - c\alpha - e\gamma = 0$, $ad - bc - \gamma\delta = \text{Ber}(B)^2$, $b\beta - d\alpha - e\delta = 0$. Thus these equations define $C_{2|1}$ as a closed affine algebraic subsupergroup of $\text{GL}_{2|1}$.

To prove that $PC_{2|1}$ is representable, we use the trick of [Manin 1991]. Let $SC_{2|1}$ denote the functor $(\text{salg})_k \rightarrow (\text{grps})$ given by

$$SC_{2|1}(A) := \{B \in C_{2|1}(A) : \text{Ber}(B) = 1\}.$$

Since its defining equations are those of $C_{2|1}$ together with $\text{Ber}(B) = 1$, $SC_{2|1}$ is a closed affine algebraic subsupergroup of $\text{GL}_{2|1}$. There is a short exact sequence of supergroups,

$$(8) \quad 0 \rightarrow SC_{2|1} \rightarrow C_{2|1} \xrightarrow{\text{Ber}} \mathbb{G}_m^{1|0} \rightarrow 0.$$

There is a splitting of this sequence, given on A -points by sending $a \in A_0^*$ to aI , and the image of $\mathbb{G}_m^{1|0}$ under the splitting is clearly normal in $C_{2|1}$, hence $C_{2|1}$ is the internal direct product of $SC_{2|1}$ and the subsupergroup $\{aI : a \in A_0^*\}$. This direct product decomposition allows us to naturally identify the functor $PC_{2|1}$ with the functor of points of $SC_{2|1}$; in particular, we see $PC_{2|1}$ is an affine algebraic supergroup, isomorphic to $SC_{2|1}$. \square

Definition 5.7. The $2|1$ -dimensional *symplectic-orthogonal supergroup* $\text{SpO}_{2|1}$ is the functor $(\text{salg})_k \rightarrow (\text{grps})$,

$$(9) \quad \text{SpO}_{2|1}(A) := \{B \in \text{GL}_{2|1}(A) : B^t H B = H\}.$$

Remark 5.8. $\text{SpO}_{2|1}$ is well known to be representable; the reader may readily write down defining equations for $\text{SpO}_{2|1}$, completely analogous to those for $C_{2|1}$, which show that $\text{SpO}_{2|1}$ is a closed affine algebraic subsupergroup of $\text{GL}_{2|1}$.

Proposition 5.9. $PC_{2|1}$ is isomorphic to the irreducible component $(\text{SpO}_{2|1})^0$ of $\text{SpO}_{2|1}$ containing the identity.

Proof. Taking the Berezinian of both sides of (9) shows that $\text{Ber}(B) = \pm 1$ for any $B \in \text{SpO}_{2|1}(A)$. This yields a short exact sequence of supergroups

$$(10) \quad 0 \rightarrow SC_{2|1} \rightarrow \text{SpO}_{2|1} \xrightarrow{\text{Ber}} \{\pm 1\} \rightarrow 0,$$

which is split by the morphism $\pm 1 \mapsto \pm I$ and $\{\pm I\}$ is obviously normal in $\mathrm{SpO}_{2|1}$. Thus $\mathrm{SpO}_{2|1}$ is the internal direct product of $\{\pm I\}$ and $\mathrm{SC}_{2|1}$. Note that $\mathrm{SC}_{2|1}$ is irreducible (one sees from its defining equations that its reduced algebraic group is SL_2 , which is known to be irreducible). Let $(\mathrm{SpO}_{2|1})^0$ denote the irreducible component of $\mathrm{SpO}_{2|1}$ that contains the identity. We claim $\mathrm{SC}_{2|1} = (\mathrm{SpO}_{2|1})^0$. Since $I \in \mathrm{SC}_{2|1} \cap (\mathrm{SpO}_{2|1})^0$, it is clear $\mathrm{SC}_{2|1} \subseteq (\mathrm{SpO}_{2|1})^0$. Conversely, we see that $(\mathrm{SpO}_{2|1})^0 \subseteq \mathrm{SC}_{2|1}$: the restriction of the morphism Ber to the irreducible supervariety $(\mathrm{SpO}_{2|1})^0$ must be constant, hence equal to 1. Since we previously showed $\mathrm{PC}_{2|1}$ is isomorphic to $\mathrm{SC}_{2|1}$, the proposition is proven. \square

Theorem 5.10. *The algebraic supergroup $\mathrm{Aut}_{\mathrm{SUSY}}(\mathbb{P}_k^{1|1})$ of SUSY-preserving automorphisms of $\mathbb{P}_k^{1|1}$ is isomorphic to $(\mathrm{SpO}_{2|1})^0$.*

Proof. As $\mathrm{Aut}_{\mathrm{SUSY}}(\mathbb{P}_k^{1|1})$ is a sheaf, the theorem reduces to the case of calculating $\mathrm{Aut}_{\mathrm{SUSY}}(\mathbb{P}_k^{1|1})(A)$ where A is a k -superalgebra. For this, we note that $\mathbb{P}_A^{1|1}$ has the SUSY structure over A induced by base change from $\mathbb{P}_k^{1|1}$, given by s .

Let $g \in \mathrm{PGL}_{2|1}(A)$ be an automorphism of $\mathbb{P}_A^{1|1}$, and \hat{g} a lift of g to $\mathrm{GL}_{2|1}(A)$. Recall that we have a natural action of the group of A -points of $\mathrm{GL}_{2|1}(A)$ on $\Gamma(\mathcal{O}_A(1) \otimes \Omega_{S/A})$. More concretely, in the given coordinates we have for any matrix $\hat{g} \in \mathrm{GL}_{2|1}(A)$,

$$\hat{g} \cdot \begin{pmatrix} z_0 \\ z_1 \\ \zeta \end{pmatrix} = \hat{g} \begin{pmatrix} z_0 \\ z_1 \\ \zeta \end{pmatrix}, \quad \hat{g} \cdot \begin{pmatrix} dz_0 \\ dz_1 \\ d\zeta \end{pmatrix} = \hat{g} \begin{pmatrix} dz_0 \\ dz_1 \\ d\zeta \end{pmatrix},$$

where we write z_i for $z_i \otimes 1$ and so on.

By [Lemma 5.3](#), g is SUSY-preserving if and only if \hat{g} sends

$$s = z_1 dz_0 - z_0 dz_1 - \zeta d\zeta = \begin{pmatrix} z_0 & z_1 & \zeta \end{pmatrix} H \begin{pmatrix} dz_0 \\ dz_1 \\ d\zeta \end{pmatrix}, \quad H = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix},$$

to a multiple of s by an invertible even function. Hence

$$\begin{pmatrix} z_0 & z_1 & \zeta \end{pmatrix} \hat{g}^t H \hat{g} \begin{pmatrix} dz_0 \\ dz_1 \\ d\zeta \end{pmatrix} = \begin{pmatrix} z_0 & z_1 & \zeta \end{pmatrix} Z(\hat{g}) H \begin{pmatrix} dz_0 \\ dz_1 \\ d\zeta \end{pmatrix},$$

i.e., $\hat{g} \in \mathrm{C}_{2|1}(A)$. It follows from [\(8\)](#) that g lies in $\mathrm{PC}_{2|1}(A)$, which is naturally identified with $(\mathrm{SpO}_{2|1})^0(A)$ by [Proposition 5.9](#). \square

Acknowledgements. We are indebted to Prof. D. Gaitsgory for clarifying to us the structure of line bundles over \mathbb{P}_A^n in the ordinary setting. We also thank Prof. L. Migliorini for helpful discussions. We are also grateful to the anonymous referee for suggestions and remarks on the paper.

References

- [Carmeli et al. 2011] C. Carmeli, L. Caston, and R. Fiorese, *Mathematical foundations of supersymmetry*, European Mathematical Society, Zürich, 2011. [MR](#) [Zbl](#)
- [Deligne and Morgan 1999] P. Deligne and J. W. Morgan, “Notes on supersymmetry (following Joseph Bernstein)”, pp. 41–97 in *Quantum fields and strings: a course for mathematicians, I* (Princeton, 1996/1997), edited by P. Deligne et al., American Mathematical Society, Providence, RI, 1999. [MR](#) [Zbl](#)
- [Donagi and Witten 2015] R. Donagi and E. Witten, “Supermoduli space is not projected”, pp. 19–71 in *String-math 2012*, edited by R. Donagi et al., Proc. Sympos. Pure Math. **90**, American Mathematical Society, Providence, RI, 2015. [MR](#) [Zbl](#)
- [Eisenbud and Harris 2000] D. Eisenbud and J. Harris, *The geometry of schemes*, Graduate Texts in Mathematics **197**, Springer, 2000. [MR](#) [Zbl](#)
- [Fiorese and Gavarini 2011] R. Fiorese and F. Gavarini, “On the construction of Chevalley supergroups”, pp. 101–123 in *Supersymmetry in mathematics and physics* (Los Angeles, 2010), edited by S. Ferrara et al., Lecture Notes in Math. **2027**, Springer, 2011. [MR](#) [Zbl](#)
- [Fiorese and Gavarini 2012] R. Fiorese and F. Gavarini, *Chevalley supergroups*, Mem. Amer. Math. Soc. **1014**, American Mathematical Society, Providence, RI, 2012. [MR](#) [Zbl](#)
- [Fiorese and Gavarini 2013] R. Fiorese and F. Gavarini, “Algebraic supergroups with Lie superalgebras of classical type”, *J. Lie Theory* **23**:1 (2013), 143–158. [MR](#) [Zbl](#)
- [Fiorese and Lledó 2015] R. Fiorese and M. A. Lledó, *The Minkowski and conformal superspaces*, World Sci., Hackensack, NJ, 2015. [MR](#) [Zbl](#)
- [Hartshorne 1977] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics **52**, Springer, 1977. [MR](#) [Zbl](#)
- [Kwok 2013] S. D. Kwok, “Super Morita theory”, submitted, 2013. [arXiv](#)
- [Manin 1988] Y. I. Manin, *Gauge field theory and complex geometry*, Grundlehren der Math. Wissenschaften **289**, Springer, 1988. [MR](#) [Zbl](#)
- [Manin 1991] Y. I. Manin, *Topics in noncommutative geometry*, Princeton Univ. Press, 1991. [MR](#) [Zbl](#)
- [Sun 2009] Y. Sun, “PGL_n and automorphisms of projective space”, presentation notes, Harvard University, 2009, Available at <https://tinyurl.com/yinotespdf>.
- [Witten 2012] E. Witten, “Notes on super Riemann surfaces and their moduli”, preprint, 2012. [arXiv](#)

Received July 14, 2017.

RITA FIORESI
DIPARTIMENTO DI MATEMATICA
UNIVERSITÀ DI BOLOGNA
BOLOGNA
ITALY
rita.fioresi@unibo.it

STEPHEN D. KWOK
MATHEMATICS RESEARCH UNIT
UNIVERSITY OF LUXEMBOURG
LUXEMBOURG
stephen.kwok@uni.lu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 295 No. 2 August 2018

| | |
|-----------------------------------------------------------------------------------------------------------|-----|
| Nonsmooth convex caustics for Birkhoff billiards | 257 |
| MAXIM ARNOLD and MISHA BIALY | |
| Certain character sums and hypergeometric series | 271 |
| RUPAM BARMAN and NEELAM SAIKIA | |
| On the structure of holomorphic isometric embeddings of complex unit balls into bounded symmetric domains | 291 |
| SHAN TAI CHAN | |
| Hamiltonian stationary cones with isotropic links | 317 |
| JINGYI CHEN and YU YUAN | |
| Quandle theory and the optimistic limits of the representations of link groups | 329 |
| JINSEOK CHO | |
| Classification of positive smooth solutions to third-order PDEs involving fractional Laplacians | 367 |
| WEI DAI and GUOLIN QIN | |
| The projective linear supergroup and the SUSY-preserving automorphisms of $\mathbb{P}^{1 1}$ | 385 |
| RITA FIORESI and STEPHEN D. KWOK | |
| The Gromov width of coadjoint orbits of the symplectic group | 403 |
| IVA HALACHEVA and MILENA PABINIAK | |
| Minimal braid representatives of quasipositive links | 421 |
| KYLE HAYDEN | |
| Four-dimensional static and related critical spaces with harmonic curvature | 429 |
| JONGSU KIM and JINWOO SHIN | |
| Boundary Schwarz lemma for nonequidimensional holomorphic mappings and its application | 463 |
| YANG LIU, ZHIHUA CHEN and YIFEI PAN | |
| Theta correspondence and the Prasad conjecture for $SL(2)$ | 477 |
| HENGFEI LU | |
| Convexity of level sets and a two-point function | 499 |
| BEN WEINKOVE | |