

*Pacific
Journal of
Mathematics*

**MINIMAL BRAID REPRESENTATIVES
OF QUASIPOSITIVE LINKS**

KYLE HAYDEN

MINIMAL BRAID REPRESENTATIVES OF QUASIPOSITIVE LINKS

KYLE HAYDEN

We show that every quasipositive link has a quasipositive minimal braid representative, partially resolving a question posed by Orevkov. These quasipositive minimal braids are used to show that the maximal self-linking number of a quasipositive link is bounded below by the negative of the minimal braid index, with equality if and only if the link is an unlink. This implies that the only amphichiral quasipositive links are the unlinks, answering a question of Rudolph's.

1. Introduction

Quasipositive links in S^3 were introduced by Rudolph [1983] and defined in terms of special braid diagrams, the details of which we recall below. These links possess a variety of noteworthy features. Perhaps most strikingly, results from [Rudolph 1983; Boileau and Orevkov 2001] show that quasipositive links are precisely those links which arise as transverse intersections of the unit sphere $S^3 \subset \mathbb{C}^2$ with complex plane curves $\Sigma \subset \mathbb{C}^2$. The hierarchy of braid-positive, positive, strongly quasipositive, and quasipositive links interacts in compelling ways with conditions such as fiberedness [Etnyre and Van Horn-Morris 2011; Hedden 2010], sliceness [Rudolph 1993], homogeneity [Baader 2005], and symplectic or Lagrangian fillability [Boileau and Orevkov 2001; Hayden and Sabloff 2015]. Quasipositive links also have well-understood behavior with respect to invariants such as the four-ball genus, the maximal self-linking number, and the Ozsváth–Szabó concordance invariant τ [Hedden 2010]. For a different perspective, we can view quasipositive braids as a monoid in the mapping class group of a disk with marked points, where they lie inside the contact-geometrically important monoid of right-veering diffeomorphisms; see [Etnyre and Van Horn-Morris 2015] for more details.

The braid-theoretic description of quasipositivity is as follows: A braid is called *quasipositive* if it is the closure of a word

$$\prod_i \omega_i \sigma_{j_i} \omega_i^{-1},$$

MSC2010: primary 57M25; secondary 57R17.

Keywords: quasipositive links, braid index, self-linking number, amphichirality.

where ω_i is any word in the braid group and σ_{j_i} is a positive standard generator. A link is then called *quasipositive* if it has a quasipositive braid representative. However, an arbitrary braid representative of a quasipositive link need not be a quasipositive braid. Along these lines, Orevkov [2000] posed the following question:

Question 1.1 (Orevkov). Let \mathcal{L} be a quasipositive link and β a minimal braid index representative of \mathcal{L} . Is β quasipositive?

Partial resolutions to this question have appeared in [Etnyre and Van Horn-Morris 2011; Feller and Kratovich 2017]. The first of these showed that the answer to Question 1.1 is “yes” for fibered strongly quasipositive links. (In contrast, the answer to the analogue of Question 1.1 for positive braids is “no”, as Stoimenow [2002] has provided examples of braid positive knots that have no positive minimal braid representatives. See also [Stoimenow 2006, §1].) The main purpose of this note is to provide another partial answer to Question 1.1.

Theorem 1.2. *Every quasipositive link has a quasipositive minimal braid index representative.*

This claim follows quickly from the proof of the generalized Jones conjecture in [LaFountain and Menasco 2014] — a substantial result in the theory of braid foliations. Our method of proof is similar to that of [Etnyre and Van Horn-Morris 2011; 2015].

A few simple consequences follow from Theorem 1.2. First, by considering the self-linking number of a quasipositive minimal braid index representative of a quasipositive link, we obtain a lower bound on the maximal self-linking number $\overline{\text{sl}}$ in terms of the minimal braid index b :

Theorem 1.3. *If \mathcal{L} is a quasipositive link, then*

$$\overline{\text{sl}}(\mathcal{L}) \geq -b(\mathcal{L}),$$

with equality if and only if \mathcal{L} is an unlink.

The calculation underlying Theorem 1.3 also lets us resolve an earlier question of Rudolph’s from [Morton 1988, Problem 9.2]:

Question 1.4 (Rudolph). Are there any amphichiral quasipositive links other than the unlinks?

At the time this question was asked, it was already known that nontrivial strongly quasipositive knots were chiral; see [Rudolph 1999, Remark 4] for a discussion of precedent results. Additional evidence for a negative answer came in the form of strong constraints on invariants of amphichiral quasipositive links (including their being slice [Wu 2011]). We confirm that the answer to Rudolph’s question is “no”.

Corollary 1.5. *If a link \mathcal{L} and its mirror $m(\mathcal{L})$ are both quasipositive, then \mathcal{L} is an unlink. In particular, the unlinks are the only amphichiral quasipositive links.*

After recalling the necessary background in Section 2, we supply proofs for the above results in Section 3.

2. Background

The generalized Jones conjecture, first confirmed by Dynnikov and Prasolov [2013], relates the writhe w and braid index n of braids with a given link type.

Theorem 2.1 [Dynnikov and Prasolov 2013, generalized Jones conjecture]. *Let β and β_0 be closed braids with the same link type \mathcal{L} , where $n(\beta_0)$ is minimal for \mathcal{L} . Then there is an inequality*

$$|w(\beta) - w(\beta_0)| \leq n(\beta) - n(\beta_0).$$

Recall Bennequin’s formula for the self-linking number of a braid β :

$$sl(\beta) = w(\beta) - n(\beta).$$

It follows from the generalized Jones conjecture, Bennequin’s formula, and the transverse Alexander theorem that a minimal braid index representative of \mathcal{L} achieves the maximal self-linking number among all transverse representatives of \mathcal{L} , denoted $\overline{sl}(\mathcal{L})$. For any braid β representing a link type \mathcal{L} , we can plot the pair $(w(\beta), n(\beta))$ in a plane. The cone of β is the collection of all pairs (w, n) realized by braids which are stabilizations of β ; see Figure 1 for an example. If β_0 is a minimal braid index representative of \mathcal{L} , we see that the right edge of its cone consists of all pairs (w, n) corresponding to braids achieving the maximal self-linking number of \mathcal{L} .

The other tool central to the proof of Theorem 1.2 is due to Orevkov and concerns braid moves that preserve quasipositivity.

Theorem 2.2 [Orevkov 2000]. *Suppose the braids β and β' are related by positive (de)stabilization. Then β is quasipositive if and only if β' is quasipositive.*




Figure 1. The cone of a braid β with $(w(\beta), n(\beta)) = (4, 2)$.

Remark 2.3. In [Orevkov 2000], an n -stranded braid is viewed as an isotopy class of n -valued functions $f : [0, 1] \rightarrow \mathbb{C}$ where $f(0)$ and $f(1)$ equal $\{1, 2, \dots, n\} \subset \mathbb{C}$. A braid is then quasipositive if one of its representatives can be expressed as a product of conjugates of the standard generators. For us, it is more convenient to study *closed* braids (up to isotopy through closed braids). Two closed braids are equivalent if and only if they can be expressed as closures of conjugate open braids. Since quasipositivity is a property of conjugacy classes of open braids, [Theorem 2.2](#) holds equally well for closed braids.

3. Quasipositive minimal braids

We proceed to the proof of the of the main result, namely that every quasipositive link has a quasipositive minimal braid representative.

Proof of [Theorem 1.2](#). Let \mathcal{L} be a quasipositive link with a minimal braid index representative β_0 and a quasipositive braid representative β_+ . Since the slice-Bennequin inequality is sharp for quasipositive links [[Rudolph 1993](#); [Hedden 2010](#)], β_+ achieves the maximal self-linking number for \mathcal{L} . As noted above, it follows that $(w(\beta_+), n(\beta_+))$ lies along the right edge of the cone of β_0 . The braids β_0 and β_+ have the same link type, so [[LaFountain and Menasco 2014](#), Proposition 1.1] implies that there are braids β'_0 and β'_+ obtained from β_0 and β_+ by negative and positive stabilization, respectively, such that β'_0 and β'_+ cobound embedded annuli. Note that β'_0 and β'_+ lie along the left and right edges of the cone, respectively, as depicted on the left side of [Figure 2](#). We also note that β'_+ is quasipositive since it is obtained from β_+ by positive stabilization.

Next, as in the proof of [[LaFountain and Menasco 2014](#), Proposition 3.2], we can find braids β''_0 and β''_+ obtained from β'_0 and β'_+ by braid isotopy, destabilization, and exchange moves such that $w(\beta''_+) = w(\beta''_0)$ and $n(\beta''_+) = n(\beta''_0)$. We claim that β''_+ has minimal braid index (as does β''_0). Indeed, since β'_0 and β'_+ lie on the left and right edges of the cone of β_0 , the destabilizations applied to them must be negative and positive, respectively. Given this and the fact that exchange moves preserve writhe and braid index, we see that β''_0 and β''_+ must also lie on the left and right edges of the cone of β_0 , respectively. But since these braids occupy the same (w, n) -point, they must lie where the edges of the cone meet. As depicted on the right side of [Figure 2](#), this implies that β''_0 and β''_+ have minimal braid index.

Finally, we show that the braid β''_+ is quasipositive. As noted above, any destabilizations of β'_+ must be positive, and these preserve quasipositivity by [Theorem 2.2](#). An exchange move also preserves quasipositivity, since it can be expressed as a combination of one positive stabilization, one positive destabilization, and a number of conjugations; see [[Birman and Wrinkle 2000](#), Figure 8]. \square




Figure 2. On the left, β'_0 and β'_+ are obtained from β_0 and β_+ by negative and positive stabilization, respectively. Then, on the right, β''_0 and β''_+ are obtained from β'_0 and β'_+ by negative and positive destabilization, respectively.

Remark 3.1. The question of whether or not *all* minimal braid index representatives of a quasipositive link are quasipositive remains open. The answer is seen to be “yes” for transversely simple link types: beginning with a quasipositive braid representative of a transversely simple link, the transverse Markov theorem implies that any minimal braid index representative can be related to it by positive (de)stabilization, which preserves quasipositivity. By the same reasoning, the answer to [Question 1.1](#) is “yes” for any link type that has a unique transverse class achieving its maximal self-linking number (but is not necessarily transversely simple). This is the case for fibered strongly quasipositive links, as shown by Etnyre and Van Horn-Morris. But it fails to hold even for nonfibered strongly quasipositive links; as pointed out by Etnyre and Van Horn-Morris, there are infinite families of 3-braids found by Birman and Menasco [2006] which are (strongly) quasipositive and of minimal braid index but not transversely isotopic.

Remark 3.2. As pointed out by Eli Grigsby, the proof of [Theorem 1.2](#) can be mirrored to show that any property of closed braids that is

- (1) preserved under transverse isotopy, and
- (2) satisfied by at least one braid representative of \mathcal{L} with maximal self-linking number

is also satisfied by at least one minimal braid index representative of \mathcal{L} .

Now we obtain the lower bound in [Theorem 1.3](#) by applying Bennequin’s formula to a quasipositive minimal braid.

Proof of Theorem 1.3. Recall that a quasipositive braid always achieves the maximal self-linking number of its link type. Thus if β is a quasipositive minimal braid index representative of \mathcal{L} , we have

$$\overline{\text{sl}}(\mathcal{L}) = \text{sl}(\beta) = w(\beta) - n(\beta) = w(\beta) - b(\mathcal{L}).$$

The desired inequality now follows from the fact that the writhe of a quasipositive braid is nonnegative, vanishing if and only if the braid is trivial. \square

Finally, we prove the corollary that resolves [Question 1.4](#).

Proof of Corollary 1.5. Observe that if β is a minimal braid index representative of \mathcal{L} , then its mirror $m(\beta)$ is minimal for $m(\mathcal{L})$. Now suppose \mathcal{L} and $m(\mathcal{L})$ are both quasipositive. The preceding proof implies that $w(\beta)$ and $w(m(\beta)) = -w(\beta)$ are both nonnegative, so $w(\beta)$ must be zero. Since we can choose the braid β to be quasipositive, the vanishing of its writhe implies that the braid itself is trivial. \square

Acknowledgements

The author thanks John Baldwin, Peter Feller, and Eli Grigsby for several stimulating conversations and for introducing him to LaFountain and Menasco's proof of the generalized Jones conjecture.

References

- [Baader 2005] S. Baader, “[Quasipositivity and homogeneity](#)”, *Math. Proc. Cambridge Philos. Soc.* **139**:2 (2005), 287–290. [MR](#) [Zbl](#)
- [Birman and Menasco 2006] J. S. Birman and W. W. Menasco, “[Stabilization in the braid groups, II: Transversal simplicity of knots](#)”, *Geom. Topol.* **10** (2006), 1425–1452. [MR](#) [Zbl](#)
- [Birman and Wrinkle 2000] J. S. Birman and N. C. Wrinkle, “[On transversally simple knots](#)”, *J. Differential Geom.* **55**:2 (2000), 325–354. [MR](#) [Zbl](#)
- [Boileau and Orevkov 2001] M. Boileau and S. Orevkov, “[Quasi-positivité d’une courbe analytique dans une boule pseudo-convexe](#)”, *C. R. Acad. Sci. Paris Sér. I Math.* **332**:9 (2001), 825–830. [MR](#) [Zbl](#)
- [Dynnikov and Prasolov 2013] I. A. Dynnikov and M. V. Prasolov, “[Bypasses for rectangular diagrams. A proof of the Jones conjecture and related questions](#)”, *Trans. Moscow Math. Soc.* (2013), 97–144. [MR](#) [Zbl](#)
- [Etnyre and Van Horn-Morris 2011] J. B. Etnyre and J. Van Horn-Morris, “[Fibered transverse knots and the Bennequin bound](#)”, *Int. Math. Res. Not.* **2011**:7 (2011), 1483–1509. [MR](#) [Zbl](#)
- [Etnyre and Van Horn-Morris 2015] J. B. Etnyre and J. Van Horn-Morris, “[Monoids in the mapping class group](#)”, pp. 319–365 in *Interactions between low-dimensional topology and mapping class groups*, edited by R. I. Bakur et al., *Geom. Topol. Monogr.* **19**, 2015. [MR](#) [Zbl](#)
- [Feller and Krcatovich 2017] P. Feller and D. Krcatovich, “[On cobordisms between knots, braid index, and the Upsilon-invariant](#)”, *Math. Ann.* **369**:1-2 (2017), 301–329. [MR](#) [Zbl](#)
- [Hayden and Sabloff 2015] K. Hayden and J. M. Sabloff, “[Positive knots and Lagrangian fillability](#)”, *Proc. Amer. Math. Soc.* **143**:4 (2015), 1813–1821. [MR](#) [Zbl](#)
- [Hedden 2010] M. Hedden, “[Notions of positivity and the Ozsváth–Szabó concordance invariant](#)”, *J. Knot Theory Ramifications* **19**:5 (2010), 617–629. [MR](#) [Zbl](#)
- [LaFountain and Menasco 2014] D. J. LaFountain and W. W. Menasco, “[Embedded annuli and Jones’ conjecture](#)”, *Algebr. Geom. Topol.* **14**:6 (2014), 3589–3601. [MR](#) [Zbl](#)
- [Morton 1988] H. R. Morton, “[Problems](#)”, pp. 557–574 in *Braids* (Santa Cruz, CA, 1986), edited by J. Birman and A. Libgober, *Contemp. Math.* **78**, Amer. Math. Soc., Providence, RI, 1988. [MR](#) [Zbl](#)

- [Orevkov 2000] S. Y. Orevkov, “Markov moves for quasipositive braids”, *C. R. Acad. Sci. Paris Sér. I Math.* **331**:7 (2000), 557–562. [MR](#) [Zbl](#)
- [Rudolph 1983] L. Rudolph, “Algebraic functions and closed braids”, *Topology* **22**:2 (1983), 191–202. [MR](#) [Zbl](#)
- [Rudolph 1993] L. Rudolph, “Quasipositivity as an obstruction to sliceness”, *Bull. Amer. Math. Soc.* **29**:1 (1993), 51–59. [MR](#) [Zbl](#)
- [Rudolph 1999] L. Rudolph, “Positive links are strongly quasipositive”, pp. 555–562 in *Proceedings of the Kirbyfest* (Berkeley, 1998), edited by J. Hass and M. Scharlemann, Geom. Topol. Monogr. **2**, 1999. [MR](#) [Zbl](#)
- [Stoimenow 2002] A. Stoimenow, “On the crossing number of positive knots and braids and braid index criteria of Jones and Morton–Williams–Franks”, *Trans. Amer. Math. Soc.* **354**:10 (2002), 3927–3954. [MR](#) [Zbl](#)
- [Stoimenow 2006] A. Stoimenow, “Properties of closed 3-braids and other link braid representations”, preprint, 2006. [arXiv](#)
- [Wu 2011] H. Wu, “Generic deformations of the colored $\mathfrak{sl}(N)$ -homology for links”, *Algebr. Geom. Topol.* **11**:4 (2011), 2037–2106. [MR](#) [Zbl](#)

Received April 21, 2017.

KYLE HAYDEN
DEPARTMENT OF MATHEMATICS
BOSTON COLLEGE
CHESTNUT HILL, MA
UNITED STATES
kyle.hayden@bc.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 295 No. 2 August 2018

Nonsmooth convex caustics for Birkhoff billiards	257
MAXIM ARNOLD and MISHA BIALY	
Certain character sums and hypergeometric series	271
RUPAM BARMAN and NEELAM SAIKIA	
On the structure of holomorphic isometric embeddings of complex unit balls into bounded symmetric domains	291
SHAN TAI CHAN	
Hamiltonian stationary cones with isotropic links	317
JINGYI CHEN and YU YUAN	
Quandle theory and the optimistic limits of the representations of link groups	329
JINSEOK CHO	
Classification of positive smooth solutions to third-order PDEs involving fractional Laplacians	367
WEI DAI and GUOLIN QIN	
The projective linear supergroup and the SUSY-preserving automorphisms of $\mathbb{P}^{1 1}$	385
RITA FIORESI and STEPHEN D. KWOK	
The Gromov width of coadjoint orbits of the symplectic group	403
IVA HALACHEVA and MILENA PABINIAK	
Minimal braid representatives of quasipositive links	421
KYLE HAYDEN	
Four-dimensional static and related critical spaces with harmonic curvature	429
JONGSU KIM and JINWOO SHIN	
Boundary Schwarz lemma for nonequidimensional holomorphic mappings and its application	463
YANG LIU, ZHIHUA CHEN and YIFEI PAN	
Theta correspondence and the Prasad conjecture for $SL(2)$	477
HENGFEI LU	
Convexity of level sets and a two-point function	499
BEN WEINKOVE	