

*Pacific
Journal of
Mathematics*

TWO APPLICATIONS OF THE SCHWARZ LEMMA

BINGYUAN LIU

TWO APPLICATIONS OF THE SCHWARZ LEMMA

BINGYUAN LIU

We exhibit two applications of Schwarz lemmas in several complex variables. The first application extends Fornæss and Stout's theorem on monotone unions of balls to monotone unions of ellipsoids. The second application extends Yang's theorem on bidiscs to the generalized bidisc defined by the author in his previous work. These applications reveal a connection between the geometry of domains and their curvatures. The proof contains a careful study of biholomorphisms, a detailed analysis on convergences, and a modified argument of Yang.

Introduction

The most striking and influential result in complex analysis of one variable is the Riemann mapping theorem. It asserts that all proper simply connected open subsets in \mathbb{C} are biholomorphic onto the unit disc. Thus, it was hoped that a similar theorem could be proved in \mathbb{C}^n for higher dimensions $n > 1$. In 1906, Poincaré showed the bidisc $\mathbb{D}^2 = \{(z, w) : |z| < 1 \text{ and } |w| < 1\}$ is not biholomorphic to the ball $\mathbb{B}^2 = \{(z, w) : |z|^2 + |w|^2 < 1\}$. This negated the expectation and motivated a new study on biholomorphism in several complex variables.

On the other hand, Fornæss and Stout [1977] showed that a Kobayashi hyperbolic manifold M is biholomorphic onto the unit ball \mathbb{B}^m , provided that M admits a monotone union of \mathbb{B}^m . Their theorem gives a version of the Riemann mapping theorem in high dimensions under some circumstances. In this paper, we follow this fashion and exhibit a theorem about monotone unions of ellipsoids $E_n := \{(z, w) \in \mathbb{C}^2 : |z|^2 + |w|^{2n} < 1\}$. More precisely, we obtain the following theorem.

Theorem 1. *Let M be a two-dimensional Hermitian manifold with a real bisectional curvature bounded from above by a negative number $-K$, and assume M is a monotone union of ellipsoids E_n for some $n \in \mathbb{Z}^+$. Then M is either biholomorphic onto E_n or onto the unit ball \mathbb{B}^2 .*

This theorem generalizes Fornæss and Stout's theorem on monotone unions of balls to monotone unions of ellipsoids in dimension 2. We remark that Fornæss

MSC2010: 32M17, 32Q05, 32Q15.

Keywords: the Schwarz lemma, applications, the generalized bidisc, ellipsoids.

and Stout's original proof is hard to be adapted into our theorem. Among other difficulties, the situation that biholomorphisms converge to a constant map is hard to be excluded. This difficulty is easy to be resolved in Fornæss and Stout's theorem because of symmetries of balls. However, the shape of ellipsoids is more irregular than balls. Hence, in order to resolve this difficulty we make local estimates around accumulation points and use the estimates to reconstruct biholomorphisms. This new technique is exhibited in [Section 2](#).

The readers are reminded that this theorem does not belong to a classical topic on automorphism groups. For the classical topics on automorphism groups, readers are referred to [[Bedford and Pinchuk 1991](#); [1998](#); [Greene and Krantz 1991](#); [1993](#); [Wong 1977](#)].

The other application of Schwarz lemmas in this article is about curvature bounds. In the 1970s, Yang [[1976](#)] showed that on polydiscs, there do not exist complete Kähler metrics with bounded holomorphic bisectional curvatures. Yang's discovery was recently generalized to product manifolds by Seshadri and Zheng [[2008](#)] and Seo [[2012](#)]. On the other hand, the author introduced a new type of domains called the generalized bidiscs in [[Liu 2017](#)]. It is known that some generalized bidiscs are biholomorphic to bounded domains in \mathbb{C}^2 . The generalized bidiscs are defined to be $\mathbb{D} \times_{\theta} \mathbb{H}^+ := \{(z, w) : z \in \mathbb{D} \text{ and } w \in e^{i\theta(z)}\mathbb{H}^+\}$. Here \mathbb{D} denotes the unit disc, \mathbb{H}^+ denotes the upper half plane, θ denotes a continuous real function depending on z , and $e^{i\theta(z)}\mathbb{H}^+$ denotes the upper half plane rotated by the angle $\theta(z)$. The generalized bidiscs are, in general, not product manifolds. However, in this paper, we show they share similar geometric properties with bidiscs. That is, some generalized bidiscs do not admit complete Kähler metrics with bounded negative holomorphic bisectional curvatures. More precisely, we show:

Theorem 2. *Let $k \in (0, \pi)$ and $\theta(z) \in [0, k)$ for all $z \in \mathbb{D}$. Then there do not exist two numbers $d > c > 0$ and a complete Kähler metric on $\mathbb{D} \times_{\theta} \mathbb{H}^+$ such that the holomorphic bisectional curvature is between $-d$ and $-c$.*

These results about curvature bounds are discussed in [Section 3](#).

1. Preliminary and fundamental facts

Let $n \in \mathbb{Z}^+$. It is classical to define ellipsoids $E_n \subset \mathbb{C}^2$ by

$$E_n = \{(z, w) : |z|^2 + |w|^{2n} < 1\}.$$

Let M be a manifold with dimension m . In this paper, we say M is a monotone union of ellipsoids E_n via f_j if

- (1) there exists a sequence of open subsets $M_j \subset M$ so that $M_j \Subset M_{j+1}$,
- (2) each M_j is biholomorphic, by f_j , onto the ellipsoids E_n , and
- (3) $M = \bigcup_j M_j$.

Remark 1.1. We sometimes omit “via f_j ” and only say “ M is a monotone union of ellipsoids E_n ”.

Remark 1.2. Similarly, one can define a monotone union of Ω for an arbitrary domain Ω .

We also recall some terminologies on Kähler manifolds. Let (M, J, h) be a Kähler manifold M of dimension m with a Kähler metric h and a complex structure J . The curvature tensor R on (M, J, h) is given by

$$R_{i\bar{j}k\bar{l}} = \frac{\partial^2 h_{i\bar{j}}}{\partial z_k \partial \bar{z}_l} - \sum_{\alpha, \beta=1}^m h^{\alpha\bar{\beta}} \frac{\partial h_{i\bar{\beta}}}{\partial z_k} \frac{\partial h_{\alpha\bar{j}}}{\partial \bar{z}_l}$$

in local coordinates (z_1, \dots, z_n) . The holomorphic bisectional curvature for $X \in T_p M$ at $p \in M$ is given by

$$B(X, Y) = -\frac{\sum_{i,j,k,l=1}^m R_{i\bar{j}k\bar{l}} X_i \bar{X}_j Y_k \bar{Y}_l}{\left(\sum_{i,j=1}^m h_{i\bar{j}} X_i \bar{X}_j\right) \left(\sum_{i,j=1}^m h_{i\bar{j}} Y_i \bar{Y}_j\right)},$$

where

$$X = \sum_{j=1}^m X_j \frac{\partial}{\partial z_j} + \sum_{j=1}^m \bar{X}_j \frac{\partial}{\partial \bar{z}_j}, \quad Y = \sum_{j=1}^m Y_j \frac{\partial}{\partial z_j} + \sum_{j=1}^m \bar{Y}_j \frac{\partial}{\partial \bar{z}_j}.$$

We are going to remind readers with backgrounds on Schwarz lemmas and almost maximal principles.

Theorem 1.3 (the Schwarz lemma of [Yau 1978]). *Let $f : M \rightarrow N$ be a holomorphic mapping from a complete Kähler manifold (M, g) with its Ricci curvature bounded from below by a negative constant $-k$ into a Hermitian manifold (N, h) with its holomorphic bisectional curvature bounded from above by a negative constant $-K$. Then*

$$f^*h \leq \frac{k}{K}g.$$

Theorem 1.4 (the almost maximal principles of [Yau 1978]; see also [Kim and Lee 2011]). *Let M be a complete Riemannian manifold M with the Ricci curvature bounded from below. Then for any C^2 smooth function $f : M \rightarrow \mathbb{R}$ that is bounded from above, there exists a sequence $\{p_k\}$ such that*

$$\lim_{k \rightarrow \infty} |\nabla T(p_k)| = 0, \quad \limsup_{k \rightarrow \infty} \Delta T(p_k) \leq 0, \quad \lim_{k \rightarrow \infty} T(p_k) = \sup_M T.$$

Recently Yang and Zheng [2016] defined a new notion of curvatures on Hermitian manifolds called real bisectional curvature. We will briefly give the definition and discuss a version of Schwarz lemma in terms of real bisectional curvature as follows.

Definition 1.5 [Yang and Zheng 2016]. Let (M^n, g) be a Hermitian manifold, and denote by R the curvature tensor of the Chern connection. We say the real bisectional curvature of M is bounded from above by a constant C if

$$\sum_{i,j,k,l} R_{i\bar{j}k\bar{l}} \xi_{ij} \xi_{kl} \leq C \operatorname{tr}(\xi^2),$$

for all nontrivial, nonnegative, Hermitian $n \times n$ matrices ξ .

Observe Theorem 4.5 in [Yang and Zheng 2016], and use the identity $\Delta v = 2\Box v$ for Kähler manifolds (here Δ is the regular Laplacian, \Box is the complex Laplacian, and v is an arbitrary smooth function). One can easily obtain the following Schwarz lemma as a corollary of Theorem 4.5 in [Yang and Zheng 2016].

Theorem 1.6 (the Schwarz lemma of [Yang and Zheng 2016]). *Let $f : M \rightarrow N$ be a holomorphic mapping from a complete Kähler manifold (M, g) with its Ricci curvature bounded from below by a negative constant $-k$ into a Hermitian manifold (N, h) with its real bisectional curvature bounded from above by a negative constant $-K$. If v is the maximal rank of the map f , then*

$$f^*h \leq \frac{kv}{K}g.$$

2. Monotone unions of ellipsoids

We discuss monotone unions of ellipsoids $E_n := \{(z, w) : |z|^2 + |w|^2 < 1\}$ in \mathbb{C}^2 in this section.

Let M be an m -dimensional complex manifold which is a monotone union of Ω via f_j . Take an arbitrary point $q \in M$, and let $j \rightarrow \infty$; then $\{f_j(q)\}_{j=1}^\infty$ has a limit point, possibly after passing to a subsequence, because of the boundedness of Ω . Then the location of limit point of $\{f_j(q)\}_{j=1}^\infty$ has two possibilities. The limit point of $\{f_j(q)\}_{j=1}^\infty$ can be either an interior point of Ω or a boundary point at $\partial\Omega$.

The following lemma settles the case that the limit of $f_j(q)$ is an interior point of Ω . From now on, we will not distinguish between the convergence of sequences and the convergence after passing to subsequences.

Lemma 2.1. *Let M be a m -dimensional Hermitian manifold with a real bisectional curvature bounded from above by a negative number $-K$. Assume M is a monotone union of $\Omega \subset \mathbb{C}^m$ via f_j where Ω is a bounded domain in \mathbb{C}^m with a complete Kähler metric of which the Ricci curvature is bounded from below by a negative number $-k$. We also assume there exists an interior point $q \in M$ so that $f_j(q) \rightarrow p \in \Omega$. Then M is biholomorphic onto Ω .*

Proof. Since Ω is bounded, f_j is a normal family of biholomorphisms. Let f_j converge to a holomorphic map F . Considering the inverses $\{f_j^{-1}\}_{j=1}^\infty$, we want

to show they are locally bounded in a small geodesic ball B_p centered $p \in \Omega$ with radius $\epsilon > 0$. Let d_M be the Hermitian metric of M and d_Ω be the Kähler metric of Ω . Indeed, by [Theorem 1.6](#),

$$(f_j^{-1})^* d_M \leq C d_\Omega$$

for each $j > 0$, where $C = km/K$. Let $N > 0$ be so that $f_j(q) \in B_p$ for all $j > N$. Considering arbitrarily $w \in B_p$, we have

$$(1) \quad d_M(q, f_j^{-1}(w)) \leq C d_\Omega(f_j(q), w) < 2C\epsilon,$$

for $j > N$. This means f_j^{-1} is locally bounded (hence a normal sequence) in B_p . We denote the limit of $\{f_j^{-1}\}_{j=1}^\infty$ by G . One can see that $F \circ G(w) = w$ in B_p because $\{f_j\}_{j=1}^\infty$ is uniformly convergent on compact subsets of M and $\{f_j^{-1}\}_{j=1}^\infty$ is uniformly convergent on compact subsets of B_p .

More generally, $\{f_j^{-1}\}_{j=1}^\infty$ is locally bounded on Ω . Indeed, we consider two interior points $w', w'' \in M$ and use [Theorem 1.6](#) for f_j^{-1} again:

$$(2) \quad d_M(f_j^{-1}(w'), f_j^{-1}(w'')) \leq C d_\Omega(w', w'').$$

From this, we can see that $f_j \circ G$ is well defined everywhere in Ω . Hence, $F \circ G$ is well defined on Ω . Since $F \circ G(w) = w$ for $w \in B_p$, and $F \circ G$ is a holomorphic map, by the identity theorem, we obtain that $F \circ G(w) = w$ for all $w \in \Omega$. This implies F is surjective.

Since $f_j(q) \rightarrow p \in \Omega$ as $j \rightarrow \infty$, it follows that $\det Jf_j(q) \not\rightarrow 0$ by Cartan's theorem. We claim the limit of $(\det Jf_j)(z)$ is nowhere vanishing for arbitrary $z \in M$, where J denotes the Jacobian. The reason is as follows. By the Cauchy estimates, the fact that $\{f_j\}_{j=1}^\infty$ is normal implies that $\{\det Jf_j\}_{j=1}^\infty$ is also normal. But $\{\det Jf_j\}_{j=1}^\infty$ is nowhere zero for each $j > 0$ because f_j is a biholomorphism and then by Hurwitz's theorem, $\det JF$ is a zero function or nowhere zero. And the claim follows by the fact that $\det Jf_j(q) \not\rightarrow 0$. Now $\det Jf_j(z) \not\rightarrow 0$ for all $z \in M$, and hence, $\det JF(z)$ is nonzero everywhere. This also implies $F(M)$ is open by the open mapping theorem.

We are going to show F is 1-1. For this, we consider two interior points $z', z'' \in M$ and use [Theorem 1.6](#) for f_j^{-1} for each $j > 0$ again:

$$(3) \quad d_M(z', z'') \leq C d_\Omega(f_j(z'), f_j(z'')).$$

Since $\det Jf_j(z)$ does not approach zero for all $z \in M$, $f_j(z)$ does not approach the boundary $\partial\Omega$ for fixed $z \in M$. In particular, $f_j(z')$ and $f_j(z'')$ do not approach the boundary $\partial\Omega$ where the Kähler metric d_Ω blows up. From this, $F(z') = F(z'')$ implies $d_\Omega(f_j(z'), f_j(z'')) \rightarrow 0$. By (3), we obtain that $z' = z''$. Consequently, F is 1-1.

Hence, M is biholomorphic onto Ω via bijective F . \square

By a similar argument, we can verify the following corollary. Instead of looking at only the exhaustive subsets of M in the previous lemma, the following corollary considers both exhaustive subsets of M and Ω .

Corollary 2.2. *Let M be an m -dimensional Hermitian manifold with a real bisectional curvature bounded from above by a negative number $-K$. Assume $M = \bigcup_j M_j$ where $M_j \subset M_{j+1}$ and f_j is a biholomorphism from M_j onto $\Omega_j \subset \Omega \subset \mathbb{C}^m$. Suppose Ω is a bounded domain in \mathbb{C}^m and Ω_j is a complete Kähler manifold with the Ricci curvatures bounded from below by a same negative number $-k$ (independent with j). We also assume there exists a point $q \in M$ so that $\det Jf_j(q) \not\rightarrow 0$. Then F is 1-1, and hence, M is taut.*

For the sake of completeness, we also include a short outline of the proof.

Outline of proof. Since Ω is bounded, $\Omega_j \subset \Omega$ is bounded too for each $j > 0$. Hence, $\{f_j\}_{j=1}^\infty$ is still normal. By $\det Jf_j(q) \not\rightarrow 0$, we can see $\det Jf_j(z) \not\rightarrow 0$ everywhere for $z \in M$, where $\{\det Jf_j(z)\}_{j=1}^\infty$ is normal because of the Cauchy estimates. This means, for any $z \in M$, $f_j(z)$ does not approach $\partial\Omega$. So by [Theorem 1.6](#), we find the limit F of f_j is 1-1. Moreover, this means M is taut. □

[Lemma 2.1](#) and [Corollary 2.2](#) tell us that if there exists one point q such that $f_j(q) \rightarrow p \in \Omega$, then for any point $z \in M$, we have $f_j(z)$ approaching an interior point of Ω . Furthermore, the limit of f_j forms a biholomorphism. However, this is not the only case. Indeed, sometimes $f_j(q)$ can approach a boundary point of Ω , and this brings trouble for getting the biholomorphism. For example, the image of $F = \lim_{j \rightarrow \infty} f_j$ might be just a constant map into a boundary point under some circumstances. The constant map of course cannot be a biholomorphism. What's behind this phenomenon is that under this situation $\det Jf_j(q) \rightarrow 0$ as $j \rightarrow \infty$. Thus, we need to compose each f_j with a biholomorphic map ϕ_j so that the resulting map $\det J\phi_j \circ f_j$ has a nonzero limit. To find the appropriate ϕ_j we need to estimate the speed of decay for $\det Jf_j(q)$. It appears the speed of decay can be arbitrary, but indeed, the decay is constrained by the location of $f_j(q)$ due to an application of the Schwarz lemma as follows. The following proposition is one of our main techniques.

Proposition 2.3. *Let M be an m -dimensional Hermitian manifold with a real bisectional curvature bounded from above by a negative number $-K$. Assume M is a monotone union of $\Omega \subset \mathbb{C}^m$ via f_j where Ω is a bounded domain in \mathbb{C}^m with a complete Bergman metric of which the Ricci curvature is bounded from below by a negative number $-k$. We also assume there exists a point $q \in M$ so that $f_j(q) \rightarrow p \in \partial\Omega$ where p is strongly pseudoconvex. Then $|\det Jf_j(q)|/\delta(f_j(q))^{(m+1)/2} \gtrsim \eta$ for some $\eta > 0$, where δ is the Euclidean distance function of Ω , i.e., $\delta(z) = \text{dist}(z, \partial\Omega)$.*

Proof. Applying [Theorem 1.6](#) for f_j^{-1} , we have $(f_j^{-1})^*g_M \leq Cg_\Omega$ for some $C > 0$ where g_M is the metric on M and g_Ω is the Bergman metric of Ω . In local coordinates, we have for any tangent vector $X_o \in T_o\Omega$ at $o \in \Omega$

$$((f_j^{-1})^*X_o)'G_M(f_j^{-1}(o))(f_j^{-1})^*X_o \leq CX_o'G_\Omega(o)X_o,$$

where we denote the conjugate transpose by $'$ and matrices of g_M and g_Ω by G_M and G_Ω , respectively. For each $j > 0$, we let $o = f_j(q)$ and have

$$((f_j^{-1})^*X_{f_j(q)})'G_M(f_j^{-1}(f_j(q)))(f_j^{-1})^*X_{f_j(q)} \leq CX_{f_j(q)}'G_\Omega(f_j(q))X_{f_j(q)}.$$

Without loss of generality, we pick up the coordinates on M at q so that G_Ω is the identity matrix at q . Hence, $(Jf_j^{-1}(f_j(q)))'Jf_j^{-1}(f_j(q)) \leq CG_\Omega(f_j(q))$ and by the Minkowski determinant theorem, we also have

$$(4) \quad |\det Jf_j^{-1}(f_j(q))|^2 \leq C|\det G_\Omega(f_j(q))|.$$

But G_Ω is a metric around a strongly pseudoconvex point p , so by [\[Fefferman 1974\]](#), it is equivalent to the $\partial\bar{\partial}(\log \delta)$ up to nonzero constant. Moreover, by computing the second-order Taylor expansion of δ at p , we also have

$$|\det G_\Omega(o)| \leq \frac{c_0}{\delta(o)^{m+1}}$$

for some $c_0 > 0$, when o is close to p . Again, putting $o = f_j(q)$, we have

$$(5) \quad |\det G_\Omega(f_j(q))| \leq \frac{c_0}{\delta(f_j(q))^{m+1}}$$

for sufficiently big $j > 0$. Since $\det Jf_j^{-1}(f_j(q)) \cdot \det Jf_j(q) = 1$, we have, by (4) and (5), that $|\det Jf_j(q)|/\delta(f_j(q))^{(m+1)/2} > 1/\sqrt{c_0C}$ for sufficient $j > 0$. We let $\eta = 1/\sqrt{c_0C}$, and thus get the desired result. \square

Another technique in this section was motivated by a simple observation in one variable.

Lemma 2.4. *Suppose there is a family of Möbius transforms on the unit disc $\psi_j(z) = (z + \alpha_j)/(1 + \bar{\alpha}_j z)$ where $\alpha_j \in \mathbb{R}$ and $\alpha_j \rightarrow 1$. Fixing $s \in (0, 1)$, we define the disc contained in \mathbb{D} :*

$$\mathfrak{D}_s := \{z \in \mathbb{C} : |z - b| < 1 - b\}$$

where $s = 1 - b$. Then $\psi_j^{-1}(\mathfrak{D}_s) \rightarrow \mathbb{D}$ as $j \rightarrow \infty$ in the sense of convergence in increasing subsets.

Proof. We compute the preimage $\psi_j^{-1}(\mathfrak{D}_s)$. By calculation, we see that

$$\left| \frac{z + \alpha_j}{1 + \bar{\alpha}_j z} - b \right| < 1 - b$$

is equivalent to the inequality

$$\begin{aligned} & \left| z + \frac{(\alpha_j - b)(1 - \alpha_j b) - (1 - b)^2 \alpha_j}{|1 - \bar{\alpha}_j b|^2 - (1 - b)^2 |\alpha_j|^2} \right|^2 \\ & < \frac{|1 - b|^2 - |\alpha_j - b|^2}{|1 - \bar{\alpha}_j b|^2 - (1 - b)^2 |\alpha_j|^2} + \frac{|(\alpha_j - b)(1 - \alpha_j b) - (1 - b)^2 \alpha_j|^2}{(|1 - \bar{\alpha}_j b|^2 - (1 - b)^2 |\alpha_j|^2)^2}. \end{aligned}$$

This is a disc centered at

$$o_j = -\frac{(\alpha_j - b)(1 - \alpha_j b) - (1 - b)^2 \alpha_j}{|1 - \bar{\alpha}_j b|^2 - (1 - b)^2 |\alpha_j|^2}$$

with radius

$$r_j = \sqrt{\frac{|1 - b|^2 - |\alpha_j - b|^2}{|1 - \bar{\alpha}_j b|^2 - (1 - b)^2 |\alpha_j|^2} + \frac{|(\alpha_j - b)(1 - \alpha_j b) - (1 - b)^2 \alpha_j|^2}{(|1 - \bar{\alpha}_j b|^2 - (1 - b)^2 |\alpha_j|^2)^2}}.$$

By a straightforward calculation, $\lim_{j \rightarrow \infty} o_j = 0$ and $\lim_{j \rightarrow \infty} r_j \rightarrow 1$. \square

The imitation to balls is also available.

Lemma 2.5. *Suppose there is a family of automorphisms*

$$\psi_j(z, w) = \left(\frac{z + a_j}{1 + \bar{a}_j z}, \frac{\sqrt{1 - |a_j|^2}}{1 + \bar{a}_j z} w \right)$$

of the unit ball \mathbb{B}^2 , where $a_j \in \mathbb{R}$ and $a_j \rightarrow 1$. Fixing $s \in (0, 1)$, we define a ball contained in \mathbb{B}^2 :

$$\mathcal{B}_s := \{(z, w) \in \mathbb{C} : |(z, w) - (b, 0)| < 1 - b\},$$

where $s = 1 - b$. Then $\psi_j^{-1}(\mathcal{B}_s) \rightarrow \mathbb{B}^m$ as $j \rightarrow \infty$ in the sense of convergence in increasing subsets.

Proof. We want to compute the preimage $\psi_j^{-1}(\mathcal{B}_s)$. For this, we need to calculate the $(z, w) \in \mathbb{C}^2$, such that $|\psi_j^{-1}(z, w) - (b, 0)| < 1 - b$. By calculation, this is equivalent to the inequality

$$\begin{aligned} (6) \quad & \left| z + \frac{(a_j - b)(1 - a_j b) - (1 - b)^2 a_j}{|1 - \bar{a}_j b|^2 - (1 - b)^2 |a_j|^2} \right|^2 + \left| \frac{\sqrt{1 - |a_j|^2}}{\sqrt{|1 - \bar{a}_j b|^2 - (1 - b)^2 |a_j|^2}} w \right|^2 \\ & < \frac{(1 - b)^2 - |a_j - b|^2}{|1 - \bar{a}_j b|^2 - (1 - b)^2 |a_j|^2} + \frac{|(a_j - b)(1 - a_j b) - (1 - b)^2 a_j|^2}{(|1 - \bar{a}_j b|^2 - (1 - b)^2 |a_j|^2)^2}. \end{aligned}$$

Again, as in the previous lemma, one can see the formula in (6) approaches

$$|z|^2 + |w|^2 < 1. \quad \square$$

Due to symmetries of balls, one can see the following lemma is also true.

Lemma 2.6. *Suppose there is a family of automorphisms*

$$\psi_j(z, w) = \left(\frac{\sqrt{1 - |a_j|^2}}{1 + \bar{a}_j w} z, \frac{w + a_j}{1 + \bar{a}_j w} \right)$$

of the unit ball \mathbb{B}^2 where $a_j \in \mathbb{R}$ and $a_j \rightarrow -1$. Fixing $s \in (0, 1)$, we define a ball contained in \mathbb{B}^2 :

$$\mathcal{B}_s := \{(z, w) \in \mathbb{C} : |(z, w) - (0, b)| < 1 + b\}$$

where $s = 1 + b$. Then $\psi_j^{-1}(\mathcal{B}_s) \rightarrow \mathbb{B}^m$ as $j \rightarrow \infty$ in the sense of convergence in increasing subsets.

Proof of Theorem 1. Let $q \in M$ and $f_j(q) \rightarrow p$ as $j \rightarrow \infty$. There are two possibilities for the location of q : $q \in E_n$ or $q \in \partial E_n$.

If $p \in E_n$, then by Lemma 2.1, M is biholomorphic to E_n . Now we analyze the cases that $p \in \partial E_n$. Suppose that $f_j(q) = (a_j, b_j)$. We define

$$\psi_j(z, w) = \left(\frac{z - a_j}{1 - \bar{a}_j z}, e^{-i\theta_j} \frac{\sqrt{1 - |a_j|^2}}{\sqrt{1 - \bar{a}_j z}} w \right).$$

Here ψ_j is a family of automorphisms of E_n and θ_j is defined so that $\psi_j \circ f_j(p) = (0, b'_j)$ with $b'_j \in \mathbb{R}$. Since $(0, b'_j) \in E_n$, by the boundedness we have that $(0, b'_j) \rightarrow (0, b'_0)$, where $-1 \leq b'_0 \leq 1$. If $b'_0 \in (-1, 1)$, then $(0, b'_0) \in E_n$. And then by Lemma 2.1 for $\psi_j \circ f_j$, we know that M is biholomorphic to E_n . If $b'_0 = 1$ or -1 , we discuss it as follows.

Without loss of generality, we now assume $b'_0 = -1$. This means it approaches a strongly pseudoconvex point $p_0 = (0, -1)$. The ellipsoid E_n , by translation, has a defining function

$$\rho(z, w) = |w - 1|^2 - \sqrt{1 - |z|^2} = -2 \operatorname{Re} w + |w|^2 + \frac{1}{n} |z|^2 + o(|z|^2).$$

Here, the point p_0 has been translated to $(0, 0)$.

On the other hand, we define $\mathcal{B}_s := \{(z, w) \in \mathbb{C}^2 : |z|^2 + |w|^2 - 2 \operatorname{Re} w < 0\}$. It is not hard to see \mathcal{B}_s is a ball centered at $(0, 1)$ with radius 1. We also define $\mathcal{B}_l := \{(z, w) \in \mathbb{C}^2 : (1/n)|z|^2 + (1/n)|w|^2 - 2 \operatorname{Re} w < 0\}$. We can see that \mathcal{B}_l is a ball centered at $(0, n)$ with radius n . So $\mathcal{B}_s \subset E_n \subset \mathcal{B}_l$, and they are tangent to each other at $(0, 0)$. We translate and rescale \mathcal{B}_l , \mathcal{B}_s , and E_n so that $\mathcal{B}_l = \mathbb{B}^2$. This setup is good for applying Lemma 2.6 to our situation. Due to the translation and rescaling, $\psi_j \circ f_j(p)$ becomes $(0, (b'_j + 1 - n)/n)$ and p_0 is once again relocated at $(0, -1)$. Since p_0 is strongly pseudoconvex, by Proposition 2.3, we see there exists $\eta > 0$ so that

$$|J(\psi_j \circ f_j)(q)| \gtrsim \eta \delta(\psi_j \circ f_j(q))^{3/2}.$$

This implies that

$$|J(\psi_j \circ f_j)(q)| \gtrsim \frac{\eta}{n} (1 - |b'_j|)^{3/2},$$

where the n is due to the rescaling \mathcal{B}_l into \mathbb{B}^2 . We define a family of automorphisms of $\mathcal{B}_l = \mathbb{B}^2$,

$$\phi_j = \left(\frac{\sqrt{1 - (b'_j)^2}}{1 + b'_j w} z, \frac{w + b'_j}{1 + b'_j w} \right).$$

and consequently, their inverses are

$$\phi_j^{-1} = \left(\frac{\sqrt{1 - (b'_j)^2}}{1 - b'_j w} z, \frac{w - b'_j}{1 - b'_j w} \right).$$

Consider

$$J\phi_j^{-1}(z, w) = \frac{(1 - (b'_j)^2)^{3/2}}{(1 - b'_j w)^3}, \quad (J\phi_j^{-1})(\psi \circ f_j(q)) = \frac{(1 - (b'_j)^2)^{3/2}}{(1 - b'_j(b'_j + 1 - n)/n)^3}.$$

We see that

$$\begin{aligned} \det(J(\phi_j^{-1} \circ \psi_j \circ f_j)(q)) &= \det((J\phi_j^{-1})(\psi_j \circ f_j(q))) \det(J(\psi_j \circ f_j)(q)) \\ &\gtrsim \frac{\eta}{n} \frac{(1 - (b'_j)^2)^3}{(1 - b'_j(b'_j + 1 - n)/n)^3}, \end{aligned}$$

where the last term is bounded below by a positive number. This can be seen by a calculation using l'Hôpital's rule on $x = b'_j \rightarrow -1$.

Thus, the limit F of $\phi_j^{-1} \circ \psi_j \circ f_j$ has nontrivial image. Moreover, the image of F is \mathbb{B}^2 because by [Lemma 2.6](#), $\phi_j^{-1}(\mathcal{B}_s) \subset \phi_j^{-1}(E_n)$ and $\phi_j^{-1}(\mathcal{B}_s)$ grows up to $\mathcal{B}_l = \mathbb{B}^2$.

Finally, we check the injectivity of F . The readers are reminded that the Bergman metric on E_n is invariant under ϕ_j^{-1} . Suppose there are $z', z'' \in M$ so that

$$\lim_{j \rightarrow \infty} \phi_j^{-1} \circ \psi_j \circ f_j(z') = \lim_{j \rightarrow \infty} \phi_j^{-1} \circ \psi_j \circ f_j(z'').$$

We can find big $N > 0$ so that for all $j > N$,

$$\phi_j^{-1} \circ \psi_j \circ f_j(z') \in \phi_N^{-1}(E_n), \quad \phi_j^{-1} \circ \psi_j \circ f_j(z'') \in \phi_N^{-1}(E_n).$$

Consequently, by [Theorem 1.6](#), we have that

$$d_M(z', z'') \leq C d_{\phi_N^{-1}(E_n)}(\phi_j^{-1} \circ \psi_j \circ f_j(z'), \phi_j^{-1} \circ \psi_j \circ f_j(z'')).$$

The assumption that $\lim_{j \rightarrow \infty} \phi_j^{-1} \circ \psi_j \circ f_j(z') = \lim_{j \rightarrow \infty} \phi_j^{-1} \circ \psi_j \circ f_j(z'')$ implies $z' = z''$. This proves the injectivity of F . \square

Without much effort, one can show the following corollary.

Corollary 2.7. *Let M be an m -dimensional Hermitian manifold with a real bisectional curvature bounded from above by a negative number $-K$, and assume M is a monotone union of balls with the same dimension. Then M is biholomorphic onto \mathbb{B}^m .*

3. An application to the generalized bidiscs

In [Liu 2017], the author defined a generalized bidisc $\mathbb{D} \rtimes_{\theta} \mathbb{H}^+ := \{(z, w) : z \in \mathbb{D}, w \in e^{i\theta(z)}\mathbb{H}^+\}$. It has a noncompact automorphism group and shares some properties with the bidisc. Indeed, when $\theta(z)$ is a zero function, $\mathbb{D} \rtimes_{\theta} \mathbb{H}^+$ is biholomorphic to a bidisc.

In this section, we prove that the generalized bidisc cannot have a complete Kähler metric with holomorphic bisectional curvature bounded by two negative numbers. This is a result of Yang type. Recall that Yang's theorem [1976] on bidiscs has certain requirements on both variables of the bidisc, but in the proof, we show that it is possible to relax the requirement for one of them. Of course similar results for higher dimensions hold for the same reason. But we will not discuss them here. Our proof is modified from [Seo 2012].

Proof of Theorem 2. We assume the conclusion is not true. Let us denote the Poincaré metric of \mathbb{D} by g and the complete Kähler metric on $\mathbb{D} \rtimes_{\theta} \mathbb{H}^+$ by h . For each z , we define $i_z(w) = (z, ie^{i\theta(z)}(1+w)/(1-w))$ from \mathbb{D} onto $e^{i\theta(z)}\mathbb{H}^+$. We get $i^*h \leq (4/c)g$ by the Schwarz lemma of Yau [1978] because the Ricci curvature of \mathbb{D} is -4 . Thus,

$$(7) \quad \begin{pmatrix} 0 & \frac{2ie^{i\theta(z)}}{(1-w)^2} \end{pmatrix} \begin{pmatrix} h_{1\bar{1}}\left(z, ie^{i\theta}\frac{1+w}{1-w}\right) & h_{1\bar{2}}\left(z, ie^{i\theta}\frac{1+w}{1-w}\right) \\ h_{2\bar{1}}\left(z, ie^{i\theta}\frac{1+w}{1-w}\right) & h_{2\bar{2}}\left(z, ie^{i\theta}\frac{1+w}{1-w}\right) \end{pmatrix} \begin{pmatrix} 0 \\ \frac{-2ie^{-i\theta(z)}}{(1-\bar{w})^2} \end{pmatrix} \\ = h_{2\bar{2}}\left(z, ie^{i\theta(z)}\frac{1+w}{1-w}\right) \frac{4}{|1-w|^4} \leq \frac{4}{c(1-|w|^2)^2}.$$

The last inequality gives

$$h_{2\bar{2}}\left(z, ie^{i\theta(z)}\frac{1+w}{1-w}\right) \leq \frac{|1-w|^4}{c(1-|w|^2)^2} \leq \frac{16}{c(1-|w|^2)^2}.$$

Since $k < \pi$, there exists $\epsilon > 0$ such that $k + \epsilon < \pi$. Because of $0 \leq \theta(z) < k$, the following is true: $(z, e^{i(k+\epsilon/2)}) \in \mathbb{D} \rtimes_{\theta} \mathbb{H}^+$. We also have, for all $z \in \mathbb{D}$,

$$(8) \quad \frac{\epsilon}{2} < k + \frac{\epsilon}{2} - \theta(z) < k + \frac{\epsilon}{2} < k + \epsilon < \pi.$$

We fix $w_0 = (e^{i(k+\epsilon/2)-\theta(z)} - i)/(e^{i(k+\epsilon/2-\theta(z))} + i)$ for all $z \in \mathbb{D}$, and by the inequality (8), we can see $|1 - |w_0|| > \eta > 0$ for some positive number η depending on ϵ .

Also by the inequality (7), we have

$$h_{2\bar{2}}\left(z, ie^{i\theta(z)}\frac{1+w_0}{1-w_0}\right) = h_{2\bar{2}}(z, e^{i\theta(z)}e^{i((k+\epsilon/2)-\theta(z))}) = h_{2\bar{2}}(z, e^{i(k+\epsilon/2)}) \leq \frac{16}{c\eta^2}.$$

Let $F(z) := h_{2\bar{2}}(z, e^{i(k+\epsilon/2)})$. We see F is a real bounded positive function on \mathbb{D} . We check its Laplacian with respect to Poincaré metric on \mathbb{D} ; we have (considering the bound of $R_{2\bar{2}1\bar{1}}$)

$$\begin{aligned} \Delta_g F(z) &= (1-|z|^2)^2 \frac{\partial^2 F}{\partial z \partial \bar{z}}(z) \\ &= (1-|z|^2)^2 \left(R_{2\bar{2}1\bar{1}}(z, e^{i(k+\epsilon/2)}) + \sum_{\alpha, \beta=1}^2 h^{\alpha\bar{\beta}} \frac{\partial h_{2\bar{\beta}}}{\partial z} \frac{\partial h_{\alpha\bar{2}}}{\partial \bar{z}} \right) \\ &\geq c(1-|z|^2)^2 h_{2\bar{2}}(z, e^{i(k+\epsilon/2)}) h_{1\bar{1}}(z, e^{i(k+\epsilon/2)}) \\ &= cF(z)(1-|z|^2)^2 h_{1\bar{1}}(z, e^{i(k+\epsilon/2)}), \end{aligned}$$

because $\sum_{\alpha, \beta=1}^2 h^{\alpha\bar{\beta}} (\partial h_{2\bar{\beta}} / \partial z) (\partial h_{\alpha\bar{2}} / \partial \bar{z})$ is nonnegative. Let $\pi : \mathbb{D} \times_{\theta} \mathbb{H}^+ \rightarrow \mathbb{D}$, $\pi(z, w) = z$. We also have $\pi^*g \leq (d/4)h$, which is $(1-|z|^2)^2 h_{1\bar{1}}(z, w) \leq 4/d$. Hence, $\Delta_g F(z) \geq (c/d)F$. Calculate

$$\Delta_g \log F(z) = \frac{\Delta_g F(z)}{F(z)} - \frac{|\nabla_g F(z)|^2}{F(z)^2} \geq \frac{2c}{d} - \frac{|\nabla_g F(z)|^2}{F(z)^2}.$$

By Theorem 1.4, a real function T bounded from above on a complete Riemannian manifold M with Ricci curvature bounded below admits a sequence $\{p_k\}_{k=0}^{\infty} \subset M$ such that

$$\lim_{k \rightarrow \infty} |\nabla T(p_k)| = 0, \quad \limsup_{k \rightarrow \infty} \Delta T(p_k) \leq 0, \quad \lim_{k \rightarrow \infty} T(p_k) = \sup_M T.$$

Although $\log F(z)$ is a real function bounded from above on \mathbb{D} , it can not have such sequence $\{p_k\}_{k=0}^{\infty} \subset \mathbb{D}$. This contradiction completes the proof. \square

A natural question is if we can relax the restriction for $\theta(z)$ in the theorem above.

Acknowledgments

I greatly thank my advisor Professor Steven Krantz for always being patient to answer my questions. I thank Professor Quo-Shin Chi, who taught me various geometries and spent much time with me. I also appreciate Professor Edward Wilson for his kindness and consistent support. Last but not least, I thank Kai Tang for his suggestion and (very) fruitful discussion.

References

- [Bedford and Pinchuk 1991] E. Bedford and S. Pinchuk, “Domains in \mathbb{C}^{n+1} with noncompact automorphism group”, *J. Geom. Anal.* **1**:3 (1991), 165–191. [MR](#) [Zbl](#)
- [Bedford and Pinchuk 1998] E. Bedford and S. Pinchuk, “Domains in \mathbb{C}^2 with noncompact automorphism groups”, *Indiana Univ. Math. J.* **47**:1 (1998), 199–222. [MR](#) [Zbl](#)
- [Fefferman 1974] C. Fefferman, “The Bergman kernel and biholomorphic mappings of pseudoconvex domains”, *Invent. Math.* **26**:1 (1974), 1–65. [MR](#) [Zbl](#)
- [Fornæss and Stout 1977] J. E. Fornæss and E. L. Stout, “Polydiscs in complex manifolds”, *Math. Ann.* **227**:2 (1977), 145–153. [MR](#) [Zbl](#)
- [Greene and Krantz 1991] R. E. Greene and S. G. Krantz, “Invariants of Bergman geometry and the automorphism groups of domains in \mathbb{C}^n ”, pp. 107–136 in *Geometrical and algebraical aspects in several complex variables* (Cetraro, 1989), edited by C. A. Berenstein and D. C. Struppa, Sem. Conf. **8**, EditEl, Rende, 1991. [MR](#) [Zbl](#)
- [Greene and Krantz 1993] R. E. Greene and S. G. Krantz, “Techniques for studying automorphisms of weakly pseudoconvex domains”, pp. 389–410 in *Several complex variables* (Stockholm, 1987/1988), edited by J. E. Fornæss, Math. Notes **38**, Princeton Univ. Press, 1993. [MR](#) [Zbl](#)
- [Kim and Lee 2011] K.-T. Kim and H. Lee, *Schwarz’s lemma from a differential geometric viewpoint*, IISc Lecture Notes Series **2**, IISc Press, Bangalore, 2011. [MR](#) [Zbl](#)
- [Liu 2017] B. Liu, “Analysis of orbit accumulation points and the Greene–Krantz conjecture”, *J. Geom. Anal.* **27**:1 (2017), 726–745. [MR](#) [Zbl](#)
- [Seo 2012] A. Seo, “On a theorem of Paul Yang on negatively pinched bisectional curvature”, *Pacific J. Math.* **256**:1 (2012), 201–209. [MR](#) [Zbl](#)
- [Seshadri and Zheng 2008] H. Seshadri and F. Zheng, “Complex product manifolds cannot be negatively curved”, *Asian J. Math.* **12**:1 (2008), 145–149. [MR](#) [Zbl](#)
- [Wong 1977] B. Wong, “Characterization of the unit ball in \mathbb{C}^n by its automorphism group”, *Invent. Math.* **41**:3 (1977), 253–257. [MR](#) [Zbl](#)
- [Yang 1976] P. C. Yang, “On Kähler manifolds with negative holomorphic bisectional curvature”, *Duke Math. J.* **43**:4 (1976), 871–874. [MR](#) [Zbl](#)
- [Yang and Zheng 2016] X. Yang and F. Zheng, “On real bisectional curvature for Hermitian manifolds”, preprint, 2016. To appear in *Trans. Amer. Math. Soc.* [arXiv](#)
- [Yau 1978] S.-T. Yau, “A general Schwarz lemma for Kähler manifolds”, *Amer. J. Math.* **100**:1 (1978), 197–203. [MR](#) [Zbl](#)

Received February 15, 2017. Revised June 4, 2017.

BINGYUAN LIU
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF CALIFORNIA, RIVERSIDE
RIVERSIDE, CA
UNITED STATES
bingyuan@ucr.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 296 No. 1 September 2018

Monotonicity of eigenvalues of geometric operators along the Ricci–Bourguignon flow	1
BIN CHEN, QUN HE and FANQI ZENG	
Composition series of a class of induced representations, a case of one half cuspidal reducibility	21
IGOR CIGANOVIĆ	
Higgs bundles over cell complexes and representations of finitely presented groups	31
GEORGIOS DASKALOPOULOS, CHIKAKO MESE and GRAEME WILKIN	
Besov-weak-Herz spaces and global solutions for Navier–Stokes equations	57
LUCAS C. F. FERREIRA and JHEAN E. PÉREZ-LÓPEZ	
Four-manifolds with positive Yamabe constant	79
HAI-PING FU	
On the structure of cyclotomic nilHecke algebras	105
JUN HU and XINFENG LIANG	
Two applications of the Schwarz lemma	141
BINGYUAN LIU	
Monads on projective varieties	155
SIMONE MARCHESI, PEDRO MACIAS MARQUES and HELENA SOARES	
Minimal regularity solutions of semilinear generalized Tricomi equations	181
ZHUOPING RUAN, INGO WITT and HUICHENG YIN	
Temperedness of measures defined by polynomial equations over local fields	227
DAVID TAYLOR, V. S. VARADARAJAN, JUKKA VIRTANEN and DAVID WEISBART	