

*Pacific  
Journal of  
Mathematics*

**LENGTH SPECTRA OF SUB-RIEMANNIAN METRICS ON  
COMPACT LIE GROUPS**

ANDRÁS DOMOKOS, MATTHEW KRAUEL, VINCENT PIGNO,  
COREY SHANBROM AND MICHAEL VANVALKENBURGH



## LENGTH SPECTRA OF SUB-RIEMANNIAN METRICS ON COMPACT LIE GROUPS

ANDRÁS DOMOKOS, MATTHEW KRAUEL, VINCENT PIGNO,  
COREY SHANBROM AND MICHAEL VANVALKENBURGH

**Length spectra for Riemannian metrics have been well studied, while sub-Riemannian length spectra remain largely unexplored. Here we give the length spectrum for a canonical sub-Riemannian structure attached to any compact Lie group by restricting its Killing form to the sum of the root spaces. Surprisingly, the shortest loops are the same in both the Riemannian and sub-Riemannian cases. We provide specific calculations for  $SU(2)$  and  $SU(3)$ .**

### 1. Introduction

While much is known about the existence and geometric properties of closed geodesics on Riemannian manifolds in general [Klingenberg 1978], and Lie groups in particular, we cannot say the same thing about their connection with the algebraic structure of Lie groups. Moreover, the sub-Riemannian setting has been mostly neglected.

In the case of simple, simply connected, compact Lie groups, Helgason [2001, Proposition 11.9] obtained the length of the shortest Riemannian geodesic loop in terms of the length of the highest root. We expand upon Helgason's work using more algebraic methods, obtaining the sub-Riemannian and Riemannian geodesic loop length spectra. The sub-Riemannian structure consists of the horizontal distribution defined by the orthogonal complement of a Cartan subalgebra and the restriction of the bi-invariant metric defined by the Killing form. To our knowledge, nothing was previously known about the connection between root systems and lengths of sub-Riemannian geodesic loops.

In Section 2 we provide the background for the root space decomposition of semisimple, compact Lie algebras and prove Theorem 2.7, which shows that all sub-Riemannian geodesics are normal. In Section 3 we work in a simple, simply connected, compact Lie group. We find connections between the algebraic information encoded in the root system of the Lie algebra and properties of Riemannian and sub-Riemannian geodesic loops. In Theorems 3.3 and 3.6 we describe the entire

---

*MSC2010:* primary 53C17, 53C22; secondary 22E30, 51N30.

*Keywords:* sub-Riemannian geometry, geodesics, root systems, compact Lie groups.

length spectra of the Riemannian and certain sub-Riemannian geodesic loops. In Theorem 3.9 we find properties that help describe the remaining sub-Riemannian geodesic loops. In Theorem 3.7, we compute the lengths of the shortest Riemannian and sub-Riemannian loops, which unexpectedly turn out to be equal. Further, in Corollary 3.8 we derive a purely algebraic formula for the length of the highest root. In Sections 4 and 5 we provide relevant examples in  $SU(2)$  and  $SU(3)$ .

Note that the terms *length spectrum* and *geodesic* have varying definitions in the literature. By length spectrum, we mean the set of lengths of all primitive geodesic loops. A sub-Riemannian geodesic is defined as in [Montgomery 2002] as a locally length minimizing curve. While in general such curves may not satisfy the geodesic equations, in our setting we show that the two notions coincide (see Theorem 2.7).

## 2. General results

In this section, we assume that  $\mathbb{G}$  is a semisimple, connected, compact matrix Lie group. This assumption is suited to present and prove some general results about sub-Riemannian geodesics, and we will use the more restrictive simple and simply connected assumptions in the following sections, where we prove results about sub-Riemannian geodesic loops. Our notation and definitions will be geared toward the presentation of the sub-Riemannian geometry, rather than the algebraic theory of Lie groups.

The Lie algebra of  $\mathbb{G}$  can be defined in terms of the matrix exponential:

$$\mathcal{G} = \{X \in \mathcal{M}_n : e^{tX} \in \mathbb{G} \text{ for all } t \in \mathbb{R}\},$$

where  $\mathcal{M}_n$  is the linear space of  $n \times n$  real or complex matrices in which  $\mathbb{G}$  is included. Then  $\mathcal{G}$  is a real Lie algebra endowed with the commutator operator

$$[X, Y] = XY - YX.$$

A Lie algebra is called simple if it is noncommutative and does not have any nontrivial ideals, and it is called semisimple if it is the direct sum of simple Lie algebras. A Lie group is simple or semisimple if its Lie algebra has the corresponding property.

The adjoint representation of  $\mathbb{G}$  is the group homomorphism

$$\text{Ad} : \mathbb{G} \rightarrow \text{Aut}(\mathcal{G}), \quad \text{Ad}(g)(X) = gXg^{-1},$$

while its differential at the identity is the adjoint representation of its Lie algebra

$$\text{ad} : \mathcal{G} \rightarrow \text{End}(\mathcal{G}), \quad \text{ad } X(Y) = [X, Y].$$

Note that, among semisimple Lie algebras,  $\text{Ad}$  is an irreducible representation of  $\mathbb{G}$  if and only if  $\mathbb{G}$  is simple.

The Killing form,

$$K(X, Y) = \text{trace}(\text{ad } X \cdot \text{ad } Y),$$

is negative definite and nondegenerate on the Lie algebra of a semisimple, compact Lie group, and hence we can define an inner product on  $\mathcal{G}$  as

$$(2-1) \quad \langle X, Y \rangle = -\rho K(X, Y),$$

where  $\rho > 0$  is a constant which can be adjusted according to our normalization preferences. The inner product (2-1) generates a bi-invariant metric on  $\mathbb{G}$ . The Killing form is Ad-invariant, so  $\text{Ad}(g)$  is a unitary linear transformation of  $\mathcal{G}$  for all  $g \in \mathbb{G}$  and  $\text{ad } X$  is skew-symmetric for all  $X \in \mathcal{G}$ .

Let  $\mathbb{T}$  be a maximal torus in  $\mathbb{G}$  and  $\mathcal{T}$  be its Lie algebra. In this case,  $\mathcal{T}$  is a maximal commutative subalgebra of  $\mathcal{G}$  called the Cartan subalgebra. Its dimension is called the rank of  $\mathcal{G}$ , and also the rank of  $\mathbb{G}$ . Consider an orthonormal basis  $\mathcal{B}_{\mathcal{T}} = \{T_1, \dots, T_r\}$  of  $\mathcal{T}$ , which will be fixed throughout the paper.

We extend the inner product (2-1) on  $\mathcal{G}$  bilinearly to the complexified Lie algebra  $\mathcal{G}_{\mathbb{C}} = \mathcal{G} \oplus i\mathcal{G}$ . The mappings  $\text{ad } T : \mathcal{G}_{\mathbb{C}} \rightarrow \mathcal{G}_{\mathbb{C}}$ ,  $T \in \mathcal{T}$ , commute and are skew-symmetric, so they share eigenspaces and have purely imaginary eigenvalues.

Once we fix the orthonormal basis  $\mathcal{B}_{\mathcal{T}}$ , we can identify  $\mathcal{T}^*$  with  $\mathcal{T}$  and define the roots as elements of the Cartan subalgebra, as in [Domokos 2015].

**Definition 2.1.** We define  $R \in \mathcal{T}$  to be a root if  $R \neq 0$  and the root space  $\mathcal{G}_R \neq \{0\}$ , where

$$\mathcal{G}_R = \{Z \in \mathcal{G}_{\mathbb{C}} : \text{ad } T(Z) = i \langle R, T \rangle Z \text{ for all } T \in \mathcal{T}\}.$$

Additionally, we use the notation  $\mathcal{G}_0 = \mathcal{T}_{\mathbb{C}} = \mathcal{T} \oplus i\mathcal{T}$ .

Let  $\mathcal{R}$  be the set of all roots, which will be partially ordered by the relation  $R_1 > R_2$  if the first nonzero coordinate of  $R_1 - R_2$  relative to the ordered basis  $\mathcal{B}_{\mathcal{T}}$  is positive. We call a root positive if its first nonzero coordinate is positive and let  $\mathcal{R}^+$  denote the set of all positive roots. For the most important properties of  $\mathcal{G}_R$  we quote [Duistermaat and Kolk 2000; Knapp 1986]:

- (i)  $\dim_{\mathbb{C}} \mathcal{G}_R = 1$ .
- (ii) If  $R \in \mathcal{R}$  then  $-R \in \mathcal{R}$ .
- (iii)  $\mathcal{G}_{-R} = \{X - iY : X + iY \in \mathcal{G}_R\}$ .
- (iv)  $\langle \mathcal{G}_{R_1}, \mathcal{G}_{R_2} \rangle = 0$  if  $R_1, R_2 \in \mathcal{R} \cup \{0\}$ ,  $R_1 \neq \pm R_2$ .
- (v)  $[\mathcal{G}_{R_1}, \mathcal{G}_{R_2}] \begin{cases} = \mathcal{G}_{R_1+R_2} & \text{if } R_1 + R_2 \in \mathcal{R} \\ = \{0\} & \text{if } R_1 + R_2 \notin \mathcal{R} \text{ and } R_1 + R_2 \neq 0 \\ \subset i\mathcal{T} & \text{if } R_1 + R_2 = 0. \end{cases}$
- (vi) If  $Z_R \in \mathcal{G}_R$  and  $Z_{-R} \in \mathcal{G}_{-R}$  then  $[Z_R, Z_{-R}] = i \langle Z_R, Z_{-R} \rangle R$ .

The above properties of  $\mathcal{G}_R$  and the real root space decomposition

$$\mathcal{G} = \mathcal{T} \oplus \mathcal{H},$$

where

$$(2-2) \quad \mathcal{H} = \mathcal{T}^\perp = \bigoplus_{R \in \mathcal{R}^+} (\mathcal{G}_R \oplus \mathcal{G}_{-R}) \cap \mathcal{G},$$

allow us to choose an orthonormal basis of  $\mathcal{H}$ ,

$$(2-3) \quad \mathcal{B}_{\mathcal{H}} = \{X_1, \dots, X_k, Y_1, \dots, Y_k\},$$

with the following properties:

- (i) For all  $1 \leq j \leq k$  there exists  $R_j \in \mathcal{R}^+$  such that  $\{X_j, Y_j\} \subset (\mathcal{G}_{R_j} \oplus \mathcal{G}_{-R_j}) \cap \mathcal{G}$ .
- (ii)  $E_{\pm j} = X_j \pm iY_j \in \mathcal{G}_{\pm R_j}$ .
- (iii)  $\langle E_j, E_{-j} \rangle = 2$ .
- (iv)  $[X_j, Y_j] = -R_j$ .

Notice that  $\{(g, \mathcal{H}_g) : g \in \mathbb{G}\}$ , where  $\mathcal{H}_g = g\mathcal{H}$ , forms a sub-bundle of the tangent bundle of  $\mathbb{G}$ , which we call the horizontal sub-bundle. The property  $\mathcal{T} \subset [\mathcal{H}, \mathcal{H}]$  shows that this horizontal sub-bundle is bracket-generating, hence its choice defines a sub-Riemannian metric on  $\mathbb{G}$  in the following way (see [Montgomery 2002]).

We call an absolutely continuous curve  $\gamma : [a, b] \rightarrow \mathbb{G}$  horizontal if  $\gamma'(t) \in \mathcal{H}_{\gamma(t)}$  for every  $t \in [a, b]$  where  $\gamma'(t)$  exists. The length of a horizontal curve is defined as

$$(2-4) \quad \text{Length}(\gamma) = \int_a^b \|\gamma'(t)\| dt.$$

The bracket-generating property implies that any two points can be connected by horizontal curves and therefore we can define a sub-Riemannian (also called Carnot–Carathéodory) distance as

$$d(x, y) = \inf\{\text{Length}(\gamma) : \gamma \text{ is a horizontal curve connecting } x \text{ and } y\}.$$

We say that a horizontal curve  $\gamma$  is a sub-Riemannian geodesic if locally it is a length minimizer. We call a sub-Riemannian geodesic  $\gamma : [0, 1] \rightarrow \mathbb{G}$  a sub-Riemannian geodesic loop if  $\gamma(0) = \gamma(1) = I$  and  $\gamma(t) \neq I$  for all  $t \in (0, 1)$ . Here,  $I$  denotes the identity matrix.

If we do not restrict the curve  $\gamma$  to be horizontal, then similar definitions lead to Riemannian geodesics and geodesic loops. With the choice of the bi-invariant inner product (2-1), the Riemannian geodesics through the identity and in the direction of an arbitrary  $X \in \mathcal{G}$  have the form

$$\gamma(t) = e^{tX};$$

see [Arvanitoyeorgos 2003, Chapter 3].

**Remark 2.2.** With our assumptions on  $\mathbb{G}$  and  $\mathcal{H}$ , all sub-Riemannian geodesics are smooth [Montgomery 1994, Theorem 3]. Moreover, as the inner product on  $\mathcal{H}$  is the restriction of the inner product (2-1) defined on  $\mathcal{G}$ , a sub-Riemannian geodesic is also a smooth curve of equal Riemannian length.

Sub-Riemannian geodesics can be characterized in various ways. We follow the description from [Montgomery 1994; 1995; 2002], but also see [Agrachev and Sarychev 1999; Boscain et al. 2002]. If a sub-Riemannian geodesic is a projection to  $\mathbb{G}$  of a solution to Hamilton’s equations for the sub-Riemannian Hamiltonian, then we call it normal, otherwise we call it abnormal. If a sub-Riemannian geodesic is a critical point of the endpoint map, then we call it singular, otherwise we call it regular [Montgomery 1994]. The following implications hold.

**Proposition 2.3** [Montgomery 2002, Theorem 5.8]. *All regular sub-Riemannian geodesics are normal and, therefore, all abnormal geodesics are singular.*

If the horizontal distribution is fat, which means that for all  $X \in \mathcal{H}$

$$\mathcal{H} + [X, \mathcal{H}] = \mathcal{G},$$

then all sub-Riemannian geodesics are normal [Montgomery 1995, Proposition 4]. For example, the horizontal distribution is fat in the case of  $SU(2)$ , but not in the case of  $SU(3)$ .

Regarding the form of the normal geodesics we have the following result, which is [Montgomery 2002, Theorem 11.8] adapted to our setting. See also [Boscain et al. 2002].

**Proposition 2.4.** *Consider a semisimple, connected, compact Lie group  $\mathbb{G}$  endowed with horizontal distribution defined by the orthogonal complement  $\mathcal{H}$  of a Cartan subalgebra  $\mathcal{T}$ , and inner product (2-1). Then the normal sub-Riemannian geodesics through the identity are of the form*

$$(2-5) \quad \gamma(t) = e^{tX} \cdot e^{-tX^\perp},$$

where  $X$  is any element of  $\mathcal{G}$  and  $X^\perp$  is the orthogonal projection of  $X$  onto  $\mathcal{T}$ .

**Definition 2.5.** If  $X \in \mathcal{H}$ , then we call  $\gamma_X(t) = e^{tX}$  a horizontal Riemannian geodesic.

These are precisely the Riemannian geodesics which are also sub-Riemannian. As we will see, they can be regular or singular.

If  $R \in \mathcal{R}^+$ , then let us use the notation  $\mathcal{H}_R = (\mathcal{G}_R \oplus \mathcal{G}_{-R}) \cap \mathcal{G}$ . With this notation we can rewrite (2-2) as

$$(2-6) \quad \mathcal{H} = \mathcal{T}^\perp = \bigoplus_{R \in \mathcal{R}^+} \mathcal{H}_R.$$

From the relations

$$[\mathcal{H}_R, \mathcal{H}_R] = \text{span}\{R\} \quad \text{and} \quad [R, \mathcal{H}_R] = \mathcal{H}_R,$$

we conclude that

$$(2-7) \quad \text{su}(2)_R = \mathcal{H}_R \oplus \text{span}\{R\}$$

is a subalgebra of  $\mathcal{G}$ , isomorphic to  $\text{su}(2)$ .

For each  $T \in \mathcal{T}$  let

$$\mathcal{R}^T = \{R \in \mathcal{R}^+ : \langle R, T \rangle = 0\}$$

and

$$(2-8) \quad \mathcal{G}^T = \bigoplus_{R \in \mathcal{R}^T} \text{su}(2)_R.$$

If  $\mathcal{R}^T \neq \emptyset$ , then  $\mathcal{G}^T$  is a nontrivial Lie subalgebra of  $\mathcal{G}$  and therefore we can find a closed, connected subgroup  $\mathbb{G}^T$  of  $\mathbb{G}$ , which has  $\mathcal{G}^T$  as its Lie algebra. Note that  $\mathbb{G}^T$  carries a sub-Riemannian geometry, for which the horizontal distribution is

$$(2-9) \quad \mathcal{H}^T = \bigoplus_{R \in \mathcal{R}^T} \mathcal{H}_R.$$

Therefore, horizontal curves in  $\mathbb{G}^T$  are also horizontal in  $\mathbb{G}$  and if a normal sub-Riemannian geodesic of  $\mathbb{G}$  lies in  $\mathbb{G}^T$ , then it is a normal sub-Riemannian geodesic of  $\mathbb{G}^T$  too.

A characteristic subgroup for a singular sub-Riemannian geodesic  $\gamma$  is a closed connected subgroup within which  $\gamma$  is regular.

**Proposition 2.6** [Montgomery 1994, Theorem 2]. *Every singular sub-Riemannian geodesic of  $\mathbb{G}$  lies in some characteristic subgroup  $\mathbb{G}^T$  with dimension strictly less than the dimension of  $\mathbb{G}$ .*

Propositions 2.3 and 2.6 allow us to give a simple algebraic proof of the following result, which is also proved using control theoretic methods, including generalized Maslov index theory, in [Boscain et al. 2002].

**Theorem 2.7.** *Consider a semisimple, connected, compact Lie group  $\mathbb{G}$  endowed with the horizontal distribution defined by the orthogonal complement  $\mathcal{H}$  of a Cartan subalgebra  $\mathcal{T}$ , and inner product (2-1). Then we have the following results.*

- (i) *All sub-Riemannian geodesics are normal.*
- (ii) *All sub-Riemannian geodesics through the identity have the form*

$$\gamma(t) = e^{tX} \cdot e^{-tX^\perp},$$

where  $X \in \mathcal{G}$  and  $X^\perp$  is the orthogonal projection of  $X$  onto  $\mathcal{T}$ .



*Proof.* Let us assume that  $\gamma$  is an abnormal sub-Riemannian geodesic of  $\mathbb{G}$ . Then, by Proposition 2.3,  $\gamma$  is singular and by Proposition 2.6, there exists  $T \in \mathcal{T}$  such that  $\gamma$  lies in a characteristic subgroup  $\mathbb{G}^T$ . But, as  $\gamma$  is regular in  $\mathbb{G}^T$ , by Proposition 2.3 it is also normal in  $\mathbb{G}^T$ . Hence,  $\gamma$  must have the form (2-5) in  $\mathbb{G}^T$ , which, by (2-6)–(2-9), gives a normal sub-Riemannian geodesic of  $\mathbb{G}$ .

Once all sub-Riemannian geodesics are normal, part (ii) is a direct consequence of Proposition 2.4. □

### 3. Lengths of sub-Riemannian geodesic loops

In this section we assume that  $\mathbb{G}$  is a simple, simply connected, compact matrix Lie group.

For each root  $R \in \mathcal{R}$  and  $n \in \mathbb{Z}$  we define the hyperplane in  $\mathcal{T}$ :

$$P(R, 2\pi n) = \{T \in \mathcal{T} : \langle R, T \rangle = 2\pi n\}.$$

The reflections in  $\mathcal{T}$  across the hyperplanes  $P(R, 0)$  will be denoted by  $r_R$ . Note that

$$r_R(T) = T - \frac{2\langle R, T \rangle}{\|R\|^2} R.$$

The Weyl group of  $\mathbb{G}$  can be defined as the group  $W$  generated by the reflections  $\{r_R : R \in \mathcal{R}\}$ .

The set

$$\mathcal{T} \setminus \bigcup_{R \in \mathcal{R}} P(R, 0)$$

is a union of disjoint, open cones, called Weyl chambers. The Weyl group acts transitively on the Weyl chambers. We define the positive Weyl chamber by

$$C = \{T \in \mathcal{T} : \langle R, T \rangle > 0, \text{ for all } R \in \mathcal{R}^+\},$$

and let  $\bar{C}$  denote its closure.

Let us choose the simple roots  $\mathcal{R}_s = \{R_1, \dots, R_m\}$ . In the case of a simple Lie algebra, the root system is irreducible and the length of the roots can take at most 2 values, which implies that the entries of the Cartan matrix,

$$N(R_j, R_k) = \frac{2\langle R_j, R_k \rangle}{\|R_k\|^2},$$

can take only the following values:

$$N(R_j, R_k) = \begin{cases} 2 & \text{if } j = k, \\ 0, -1, -2, -3 & \text{if } j \neq k, \end{cases}$$

where at most one of  $-2$  or  $-3$  can appear in the matrix. For each  $R \in \mathcal{R}$  we denote by

$$(3-1) \quad P_R = \frac{2\pi R}{\|R\|^2}$$

the orthogonal projection of the origin onto the hyperplane  $P(R, 2\pi)$ . It is known from [Helgason 2001, Chapter 7, Lemma 7.6] that

$$(3-2) \quad e^{2P_R} = I, \text{ for all } R \in \mathcal{R}.$$

The unit lattice in  $\mathcal{T}$  is defined by

$$\mathcal{L}_{\mathcal{T}} = \{T \in \mathcal{T} : e^T = I\},$$

and let us also set

$$\mathcal{Z}_{\mathcal{T}} = \{n_1 2P_{R_1} + \dots + n_m 2P_{R_m} : n_1, \dots, n_m \in \mathbb{Z}\}.$$

By the commutativity of  $\mathcal{T}$ , it is evident that  $\mathcal{Z}_{\mathcal{T}} \subset \mathcal{L}_{\mathcal{T}}$ . By [Simon 1996, Theorem IX.1.6] we know that  $\mathcal{L}_{\mathcal{T}}/\mathcal{Z}_{\mathcal{T}} \cong \pi_1(\mathbb{G})$ , where  $\pi_1(\mathbb{G})$  is the fundamental group of  $\mathbb{G}$ . Since  $\mathbb{G}$  is simply connected, it follows that

$$(3-3) \quad \mathcal{L}_{\mathcal{T}} = \mathcal{Z}_{\mathcal{T}}.$$

From [Simon 1996, Theorem IX.1.4], it is also known that

$$(3-4) \quad \mathcal{L}_{\mathcal{T}} \subseteq \{T \in \mathcal{T} : \langle R, T \rangle \in 2\pi\mathbb{Z} \text{ for all } R \in \mathcal{R}\},$$

and the two sets in (3-4) are equal only if the center of  $\mathbb{G}$  equals  $\{I\}$ .

**Definition 3.1.** We call the numbers  $n_1, \dots, n_m \in \mathbb{N} \cup \{0\}$  relatively prime if at least one of the numbers is nonzero and the greatest common factor of the nonzero numbers is 1. In particular, if we have only one nonzero number, then it must be 1.

**Remark 3.2.** By (3-3), if the numbers  $n_1, \dots, n_m \in \mathbb{N} \cup \{0\}$  are relatively prime, then the line segment joining the origin to  $n_1 2P_{R_1} + \dots + n_m 2P_{R_m}$  intersects  $\mathcal{L}_{\mathcal{T}}$  only at the endpoints.

**Theorem 3.3.** *Let  $\mathbb{G}$  be a simple, simply connected, compact Lie group endowed with the bi-invariant inner product (2-1).*

- (a) *If the numbers  $n_1, \dots, n_m \in \mathbb{N} \cup \{0\}$  are relatively prime and*

$$T = n_1 2P_{R_1} + \dots + n_m 2P_{R_m},$$

*then  $\gamma_T(t) = e^{tT}$ ,  $0 \leq t \leq 1$ , is a Riemannian geodesic loop with length*

$$\|n_1 2P_{R_1} + \dots + n_m 2P_{R_m}\|.$$

- (b) *All Riemannian geodesic loops in  $\mathbb{G}$  have lengths*

$$\|n_1 2P_{R_1} + \dots + n_m 2P_{R_m}\|,$$

*where  $n_1, \dots, n_m \in \mathbb{N} \cup \{0\}$  are relatively prime.*

*Proof.* (a) If the rank of  $\mathbb{G}$  is 1, then  $\mathbb{T} = \text{U}(1)$  and any geodesic loop in  $\mathbb{T}$  has length  $2\pi$ . Now suppose the rank of  $\mathbb{G}$  is greater than or equal to 2. Let  $T = n_1 2P_{R_1} + \dots + n_m 2P_{R_m}$ , where  $n_1, \dots, n_m \in \mathbb{N} \cup \{0\}$  are relatively prime and  $\gamma_T(t) = e^{tT}$ . By the commutativity of the elements of  $\mathcal{T}$  we know that  $\gamma_T(1) = I$ . If, for some  $0 < t < 1$ , we have  $\gamma_T(t) = I$ , then  $t(n_1 2P_{R_1} + \dots + n_m 2P_{R_m}) \in \mathcal{L}_{\mathcal{T}}$  and, by Remark 3.2, this contradicts the fact that  $n_1, \dots, n_m$  are relatively prime. Hence, the length of one loop described by  $\gamma_T$  is

$$(3-5) \quad \text{Length}(\gamma_T) = \int_0^1 \|T\| dt = \|n_1 2P_{R_1} + \dots + n_m 2P_{R_m}\|.$$

(b) Let  $X \in \mathcal{G}$  and  $\gamma_X(t) = e^{tX}$ . Assume that  $\gamma_X(1) = I$  and  $\gamma_X(t) \neq I$  if  $0 < t < 1$ . Since  $\text{Ad}(\mathbb{G})(X) \cap \mathcal{T}$  is nonempty and finite, and the Weyl group acts transitively on the Weyl chambers, and each element of the Weyl group can be written as  $\text{Ad}(g)$  for some  $g \in \mathbb{G}$ , it follows that there exists  $g \in \mathbb{G}$  such that  $T_X = \text{Ad}(g)X \in \overline{\mathcal{C}}$ . Hence,  $e^{T_X} = g e^X g^{-1} = I$  and therefore  $T_X \in \mathcal{L}_{\mathcal{T}}$ . By (3-3) we have that  $T_X = n_1 2P_{R_1} + \dots + n_m 2P_{R_m}$ , where  $n_1, \dots, n_m \in \mathbb{N} \cup \{0\}$  are relatively prime. Using the fact that  $\|T_X\| = \|X\|$  we find that

$$\text{Length}(\gamma_X) = \int_0^1 \|T_X\| dt = \|n_1 2P_{R_1} + \dots + n_m 2P_{R_m}\|. \quad \square$$

**Remark 3.4.** Moreover, for any  $0 \neq T = n_1 2P_{R_1} + \dots + n_m 2P_{R_m}$  we have that  $\text{Ad}(\mathbb{G})(T) \cap (\mathcal{G} \setminus \mathcal{T}) \neq \emptyset$ , so there exists  $X \notin \mathcal{T}$  in the same conjugacy class with  $T$ . Hence we have a Riemannian geodesic loop outside of  $\mathbb{T}$ , corresponding to  $X$ , which has length equal to  $\|T\|$  in (3-5).

We need the following lemma to generalize Theorem 3.3 to the case of horizontal Riemannian geodesic loops (see Definition 2.5).

**Lemma 3.5.** *For any  $T \in \mathcal{T}$  we have  $\text{Ad}(\mathbb{G})(T) \cap \mathcal{H} \neq \emptyset$ .*

*Proof.* By [D’Andrea and Maffei 2016, Lemma 2.2], given  $\mathcal{T}$ , we can construct another Cartan subalgebra  $\mathcal{T}'$  which is orthogonal to  $\mathcal{T}$ . Hence,  $\mathcal{T}' \subset \mathcal{H}$  and, as any two Cartan subalgebras are conjugate, there exists some  $g \in \mathbb{G}$  such that  $\text{Ad}(g)\mathcal{T} = \mathcal{T}'$ . Hence, we conclude that for any  $T \in \mathcal{L}_{\mathcal{T}}$  we have that  $\text{Ad}(\mathbb{G})(T) \cap \mathcal{H} \neq \emptyset$ .  $\square$

**Theorem 3.6.** *Consider a simple, simply connected, compact Lie group  $\mathbb{G}$  endowed with horizontal distribution defined by the orthogonal complement  $\mathcal{H}$  of a Cartan subalgebra  $\mathcal{T}$ , and inner product (2-1). Then the horizontal Riemannian geodesic loops have lengths*

$$\|n_1 2P_{R_1} + \dots + n_m 2P_{R_m}\|,$$

where  $n_1, \dots, n_m \in \mathbb{N} \cup \{0\}$  are relatively prime.

*Proof.* Let  $X \in \mathcal{H}$  and  $\gamma_X(t) = e^{tX}$ . If  $\gamma_X(1) = I$  and  $\gamma_X(t) \neq I$  for all  $0 < t < 1$ , then we can follow the proof of Theorem 3.3(b), to conclude that there exist  $n_1, \dots, n_m \in \mathbb{N} \cup \{0\}$  relatively prime such that

$$\text{Length}(\gamma_X) = \|n_1 P_{R_1} + \dots + n_m P_{R_m}\|.$$

By Lemma 3.5, the entire length spectrum of  $\|n_1 P_{R_1} + \dots + n_m P_{R_m}\|$ , where  $n_1, \dots, n_m \in \mathbb{N} \cup \{0\}$  are relatively prime, is covered, and this finishes the proof.  $\square$

One might expect the shortest sub-Riemannian geodesic loops to be longer than their Riemannian counterparts. Surprisingly, the following result, which generalizes the Riemannian case of [Helgason 2001, Chapter 7, Proposition 11.9], proves otherwise.

**Theorem 3.7.** *The shortest sub-Riemannian geodesic loops are also the shortest Riemannian geodesic loops. Their common length is  $4\pi/\|R^*\|$ , where  $R^*$  is the highest root.*

*Proof.* We first consider the Riemannian case. Without loss of generality we can assume that the rank of  $\mathcal{G}$  is greater than 1. Let  $\gamma^*(t) = e^{t2P_{R^*}}$ . By (3-2) we know that  $\gamma^*(1) = I$ . Moreover, there exists  $R \in \mathcal{R}^+$  such that

$$N(R, R^*) = 2 \frac{\langle R, R^* \rangle}{\|R^*\|^2} = 1.$$

Therefore, for any  $0 < t < 1$  we have

$$\langle R, t2P_{R^*} \rangle = 2\pi t,$$

which, by (3-4), implies that  $\gamma^*(t) \neq I$  if  $0 < t < 1$ . Hence, the length of one loop described by  $\gamma^*$  is

$$\text{Length}(\gamma^*) = \int_0^1 \|2P_{R^*}\| dt = \frac{4\pi}{\|R^*\|}.$$

Let  $T = n_1 2P_{R_1} + \dots + n_m 2P_{R_m}$ , where  $n_1, \dots, n_m \in \mathbb{N} \cup \{0\}$  are relatively prime and let  $\gamma_T(t) = e^{tT}$ . Assume that  $\gamma_T(t) \neq I$  if  $0 < t < 1$  and

$$\text{Length}(\gamma_T) \leq \text{Length}(\gamma^*).$$

Hence,

$$\|T\| \leq \frac{4\pi}{\|R^*\|} = \|2P_{R^*}\|.$$

As in the proof of Theorem 3.3, by the fact that the Weyl group acts transitively on the Weyl chambers, there exist  $g \in \mathbb{G}$  and  $T_1 \in \bar{C}$  such that  $T_1 = \text{Ad}(g)T$ . Therefore,  $e^{T_1} = I$  and hence  $\langle R^*, T_1 \rangle = 2\pi n$  for some  $n \in \mathbb{N}$ . By [Helgason 2001, Chapter 7, Theorem 6.1],

$$P(R^*, 2\pi) \cap \bar{C} \cap \mathcal{L}_T = \emptyset,$$

which implies that  $n \neq 1$ . On the other hand,  $\|T_1\| = \|T\| \leq \|2P_{R^*}\|$ , which is the shortest distance from the origin to  $P(R^*, 4\pi)$ . Therefore,  $n = 2$  and this implies that  $T_1 = 2P_{R^*}$ . In conclusion, we have  $\text{Length}(\gamma_T) = 4\pi/\|R^*\|$ , which establishes the length of the shortest Riemannian geodesic loops. Note that this slight generalization of [Helgason 2001, Chapter 7, Proposition 11.9] is proved differently here.

We now consider the sub-Riemannian case. Theorem 3.6 implies that the shortest horizontal Riemannian geodesic loops have length equal to  $4\pi/\|R^*\|$ , which equals the length of the shortest Riemannian geodesic loops by the argument above. By Remark 2.2, every sub-Riemannian geodesic is a smooth Riemannian curve of equal length, so we conclude that  $4\pi/\|R^*\|$  is the shortest length for any sub-Riemannian geodesic loop.  $\square$

Theorem 3.7 implies the following result concerning the highest root.

**Corollary 3.8.** *We have*

$$\|R^*\| = \max \frac{4\pi}{\|n_1 2P_{R_1} + \dots + n_m P_{R_m}\|},$$

where  $n_1, \dots, n_m \in \mathbb{N} \cup \{0\}$  are relatively prime.

Regarding the sub-Riemannian geodesic loops which are not necessarily horizontal Riemannian, we have the following result.

**Theorem 3.9.** *Let  $X = H + X^\perp$  be such that  $H \in \mathcal{H}$  and  $X^\perp \in \mathcal{T}$ . Consider  $\gamma(t) = e^{tX} \cdot e^{-tX^\perp}$  and assume that  $\gamma(t) \neq I$  if  $0 < t < 1$  and  $\gamma(1) = I$ . Then:*

(a) *The length of  $\gamma$  satisfies*

$$\text{Length}(\gamma) = \|H\| \geq \frac{4\pi}{\|R^*\|},$$

*and there is an  $X = H + X^\perp$  for which  $4\pi/\|R^*\|$  is attained.*

(b) *We have*

$$(3-6) \quad H = e^{X^\perp} H e^{-X^\perp}.$$

(c) *If*

$$\text{Ad}(\mathbb{G})(X^\perp) \cap \mathcal{T} = \{S_1, \dots, S_l\},$$

*then for all  $1 \leq j \leq l$  there exist  $L_j \in \mathcal{L}_\mathcal{T}$  such that*

$$\text{Ad}(\mathbb{G})(X) \cap \mathcal{T} = \{S_1 + L_1, \dots, S_l + L_l\}.$$

*Proof.* (a) Note that  $\gamma(1) = I$  implies that  $e^X = e^{X^\perp}$ . Then,

$$(3-7) \quad \gamma'(t) = e^{tX} \cdot X \cdot e^{-tX^\perp} - e^{tX} \cdot X^\perp \cdot e^{-tX^\perp} = e^{tX}(X - X^\perp)e^{-tX^\perp},$$

and

$$\|\gamma'(t)\|_{\gamma(t)} = \|\gamma(t)^{-1} \cdot \gamma'(t)\|_I = \|e^{tX^\perp} \cdot (X - X^\perp) \cdot e^{-tX^\perp}\| = \|X - X^\perp\|.$$

Hence, the length of  $\gamma$  is  $\|X - X^\perp\| = \|H\|$ . The fact that  $\|H\|$  is at least  $4\pi/\|R^*\|$  is an immediate consequence of Theorem 3.6.

Consider the simply connected Lie subgroup  $SU(2)_{R^*}$  of  $\mathbb{G}$  which has its Lie algebra equal to  $\mathfrak{su}(2)_{R^*}$ , and denote by  $X^*$  and  $Y^*$  those elements of (2-3) which, together with  $R^*$ , generate  $\mathfrak{su}(2)_{R^*}$ . The relations

$$[R^*, X^*] = -\|R^*\|^2 Y^* \quad \text{and} \quad [R^*, Y^*] = \|R^*\|^2 X^*$$

show that the only positive root, and hence the highest root in  $\mathfrak{su}(2)_{R^*}$ , is  $R^*$ . In a similar way to the proof of (4-2), we can obtain a sub-Riemannian geodesic loop in  $SU(2)_{R^*}$  whose length is  $4\pi/\|R^*\|$ .

(b) We claim that  $\gamma'(0) = \gamma'(1)$ . This information can be found in [Helgason 2001, page 148, Exercise 3] and its proof is based on the fact that for all  $t \in \mathbb{R}$ ,

$$\begin{aligned} \gamma(t+1) &= e^{(t+1)X} e^{-(t+1)X^\perp} = e^{tX} e^X e^{-tX^\perp} e^{-X^\perp} \\ &= e^{tX} e^X e^{-X^\perp} e^{-tX^\perp} = e^{tX} e^{-tX^\perp} = \gamma(t). \end{aligned}$$

By (3-7), it follows that

$$X - X^\perp = e^X (X - X^\perp) e^{-X^\perp},$$

which clearly implies (3-6).

(c) By the properties of the adjoint representation there exist  $S_1, \dots, S_l \in \mathcal{T}$  and  $S'_1, \dots, S'_l \in \mathcal{T}$ , where  $l$  is the number of Weyl chambers, such that

$$(3-8) \quad \text{Ad}(\mathbb{G})(X^\perp) \cap \mathcal{T} = \{S_1, \dots, S_l\}$$

and

$$(3-9) \quad \text{Ad}(\mathbb{G})(X) \cap \mathcal{T} = \{S'_1, \dots, S'_l\}.$$

Note that in (3-8) and (3-9) some of the  $S_j$  and  $S'_j$  might be repeated if they belong to one of the hyperplanes  $P(R, 0)$  for  $R \in \mathcal{R}$ .

Therefore,

$$(3-10) \quad \text{Ad}(\mathbb{G})(e^{X^\perp}) \cap \mathbb{T} = \{e^{S_1}, \dots, e^{S_l}\},$$

and

$$(3-11) \quad \text{Ad}(\mathbb{G})(e^X) \cap \mathbb{T} = \{e^{S'_1}, \dots, e^{S'_l}\}.$$

The fact that  $e^{X^\perp} = e^X$  implies that the sets in (3-10) and (3-11) must coincide. Therefore, by rearranging the elements if necessary, we can suppose that for all  $1 \leq j \leq l$ , we have  $e^{S_j} = e^{S'_j}$ , which immediately implies the existence of  $L_j \in \mathcal{L}_\mathcal{T}$  such that  $S'_j = S_j + L_j$ . □

Since  $X^\perp \in \mathcal{T}$ , we can see that one of  $S_1, \dots, S_l$  in (3-8) must be  $X^\perp$ . Therefore, we have the following corollary.

**Corollary 3.10.** *Under the assumptions of Theorem 3.9, there exist  $g \in \mathbb{G}$  and  $L \in \mathcal{L}_{\mathcal{T}}$  such that  $X = \text{Ad}(g)(X^{\perp} + L)$ .*

**4. The case of SU(2)**

The special unitary group of  $2 \times 2$  complex matrices is

$$\text{SU}(2) = \left\{ g = \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} : \alpha, \beta \in \mathbb{C}, |\alpha|^2 + |\beta|^2 = 1 \right\}.$$

Its Lie algebra is the three dimensional real Lie algebra

$$\text{su}(2) = \left\{ X = \begin{pmatrix} ix_1 & x_2 + ix_3 \\ -x_2 + ix_3 & -ix_1 \end{pmatrix} : x_1, x_2, x_3 \in \mathbb{R} \right\}.$$

The Killing form of  $\text{su}(2)$  is

$$K(X, Y) = 4 \text{trace}(XY),$$

while the inner product (2-1) is defined as

$$\langle X, Y \rangle = -\frac{1}{2} \text{trace}(XY).$$

The Cartan subalgebra  $\mathcal{T}$  is spanned by the unit vector

$$T_1 = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}$$

and the orthonormal basis of  $\mathcal{H}$  is formed by

$$X_1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \text{ and } Y_1 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}.$$

The exponential map  $\exp : \text{su}(2) \rightarrow \text{SU}(2)$  has the following simple form:

$$\exp(X) = e^X = \cos(\|X\|) I + \frac{\sin(\|X\|)}{\|X\|} X.$$

Consider  $X = aX_1 + bY_1 + cT_1$ . Then, for  $0 \leq t \leq 1$ ,

$$e^{tX} = \cos(t\sqrt{a^2 + b^2 + c^2}) I + \frac{\sin(t\sqrt{a^2 + b^2 + c^2})}{\sqrt{a^2 + b^2 + c^2}} X,$$

and

$$e^{tX^{\perp}} = \cos(tc) I + \sin(tc) T_1.$$

If we have  $\sqrt{a^2 + b^2 + c^2} = 2\pi$ , the Riemannian geodesic  $e^{tX}$  closes the first time at  $t = 1$ , which shows that the Riemannian length spectrum equals  $\{2\pi\}$ . For the sub-Riemannian geodesic  $\gamma(t) = e^{tX} e^{-tX^{\perp}}$ , the condition  $e^X = e^{X^{\perp}}$  implies that

$$(4-1) \quad \sqrt{a^2 + b^2 + c^2} = n\pi \quad \text{and} \quad |c| = m\pi,$$

where  $m, n \in \mathbb{N} \cup \{0\}$ ,  $m \leq n$ , are both even or both odd. To ensure  $\gamma(t) \neq I$  for all  $0 < t < 1$ , we require that  $m, n \in \mathbb{N} \cup \{0\}$  are both odd and relatively prime or both even and  $\frac{m}{2}, \frac{n}{2}$  are relatively prime with one of them odd and the other even.

Notice that, as the only positive root of  $SU(2)$  is  $R_1 = 2T_1$ , we have  $\|R^*\| = \|2T_1\| = 2$ , and therefore

$$(4-2) \quad \|H\| = \|X - X^\perp\| = \pi\sqrt{n^2 - m^2} = \frac{2\pi\sqrt{n^2 - m^2}}{\|R^*\|}.$$

The same result can be obtained by Theorem 3.9. In  $SU(2)$  the unit lattice is  $\mathcal{L}_{\mathcal{T}} = \{2k\pi T_1 : k \in \mathbb{Z}\}$ . The formula (3-6) implies that

$$c = m\pi \in \pi\mathbb{Z}.$$

The matrices  $S'_1$  and  $S'_2$  from (3-9) are diagonal with entries consisting of the eigenvalues of  $X$ . Thus, Theorem 3.9 implies that there exists some  $k \in \mathbb{N}$  such that

$$\sqrt{a^2 + b^2 + m^2\pi^2} - m\pi = 2k\pi,$$

and this implies (4-1).

We have therefore presented two algebraic proofs of the following proposition, which is a special case of Theorems 3.3, 3.6, and 3.7, and which extends the results from [Chang et al. 2011; Klapheck and VanValkenburgh 2019].

**Proposition 4.1.** *In  $SU(2)$  the following properties hold.*

- (a) *The Riemannian geodesic loops have length equal to  $2\pi$ .*
- (b) *The horizontal Riemannian geodesic loops have length equal to  $2\pi$ .*
- (c) *The shortest sub-Riemannian geodesic loops have length equal to  $2\pi$ .*
- (d) *The sub-Riemannian geodesic loops have lengths equal to  $\pi\sqrt{n^2 - m^2}$ , where  $m, n \in \mathbb{N} \cup \{0\}$  are odd and relatively prime or even and  $\frac{m}{2}, \frac{n}{2}$  are relatively prime with one of them odd and the other even.*

**Remark 4.2.** As an introduction to the next section, let us show that we can use Viète’s formulas to get the result of Proposition 4.1(d). Indeed, the characteristic polynomial of  $X$  is

$$P(\lambda) = \lambda^2 + (a^2 + b^2 + c^2),$$

and by Theorem 3.9 and the first Viète formula, the eigenvalues of  $X$  must be of the form  $\lambda_1 = -ci - 2k\pi i$  and  $\lambda_2 = ci + 2k\pi i$ , where  $k \in \mathbb{N}$ . The second Viète formula gives

$$\lambda_1\lambda_2 = (c + 2k\pi)^2 = a^2 + b^2 + c^2,$$

which leads to (4-1).



**Remark 4.3.** For comparison with the case of  $SU(3)$  in the next section, note that in  $SU(2)$  the sub-Riemannian geodesic loops have the form

$$(4-3) \quad \gamma(t) = e^{t(a_1 X_1 + b_1 Y_1 + \frac{m}{2} R_1)} e^{-t \frac{m}{2} R_1},$$

where  $a, b, c, m$  satisfy (4-1).

### 5. The case of $SU(3)$

Consider the special unitary group of  $3 \times 3$  complex matrices

$$SU(3) = \{g \in GL(3, \mathbb{C}) : g \cdot g^* = I, \det g = 1\},$$

and its Lie algebra

$$\mathfrak{su}(3) = \{X \in \mathfrak{gl}(3, \mathbb{C}) : X + X^* = 0, \text{trace } X = 0\}.$$

The inner product is defined by

$$\langle X, Y \rangle = -\frac{1}{2} \text{trace}(XY).$$

We consider the maximal torus

$$\mathbb{T} = \left\{ \begin{pmatrix} e^{ia_1} & 0 & 0 \\ 0 & e^{ia_2} & 0 \\ 0 & 0 & e^{ia_3} \end{pmatrix} : a_1, a_2, a_3 \in \mathbb{R}, a_1 + a_2 + a_3 = 0 \right\}$$

and its Lie algebra

$$\mathcal{T} = \left\{ \begin{pmatrix} ia_1 & 0 & 0 \\ 0 & ia_2 & 0 \\ 0 & 0 & ia_3 \end{pmatrix} : a_1, a_2, a_3 \in \mathbb{R}, a_1 + a_2 + a_3 = 0 \right\},$$

which is our choice for the Cartan subalgebra. The following are the Gell-Mann matrices, which form an orthonormal basis of  $\mathfrak{su}(3)$  and satisfy the relations in (2-2), (2-3), and (i)–(iv) on page 324:

$$T_1 = \begin{pmatrix} -i & 0 & 0 \\ 0 & i & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad T_2 = \begin{pmatrix} \frac{-i}{\sqrt{3}} & 0 & 0 \\ 0 & \frac{-i}{\sqrt{3}} & 0 \\ 0 & 0 & \frac{2i}{\sqrt{3}} \end{pmatrix}, \quad X_1 = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad Y_1 = \begin{pmatrix} 0 & i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

$$X_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \quad Y_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & -i & 0 \end{pmatrix}, \quad X_3 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad Y_3 = \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}.$$

The positive roots are

$$R_1 = \begin{pmatrix} -2i & 0 & 0 \\ 0 & 2i & 0 \\ 0 & 0 & 0 \end{pmatrix} = 2T_1,$$

$$R_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 2i & 0 \\ 0 & 0 & -2i \end{pmatrix} = T_1 - \sqrt{3}T_2,$$

$$R_3 = \begin{pmatrix} -2i & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 2i \end{pmatrix} = T_1 + \sqrt{3}T_2.$$

The highest root is  $R^* = R_1$ , while the two simple roots are  $R_2$  and  $R_3$ . The unit lattice is

$$\mathcal{L}_{\mathcal{T}} = \{n\pi R_2 + m\pi R_3 : n, m \in \mathbb{Z}\},$$

and observe that

$$(5-1) \quad e^{\pi R_1} = e^{\pi R_2} = e^{\pi R_3} = I.$$

For  $k = 1, 2, 3$ , the projections of the origin onto the hyperplanes  $P(R_k, 2\pi)$  are

$$P_{R_k} = \frac{\pi}{2} R_k,$$

and, indeed, (5-1) is equivalent to

$$e^{2P_{R_k}} = I, \quad k = 1, 2, 3.$$

Observing that

$$\|n\pi R_2 + m\pi R_3\| = 2\pi\sqrt{n^2 - nm + m^2},$$

we conclude that, in  $SU(3)$ , Theorems 3.3, 3.6, and 3.7 have the following special form.

**Proposition 5.1.** *In  $SU(3)$  the following properties hold.*

(a) *The Riemannian geodesic loops have lengths equal to*

$$2\pi\sqrt{n^2 - nm + m^2},$$

*where  $n, m \in \mathbb{N} \cup \{0\}$  are relatively prime.*

(b) *The horizontal Riemannian geodesic loops have lengths equal to*

$$2\pi\sqrt{n^2 - nm + m^2},$$

*where  $n, m \in \mathbb{N} \cup \{0\}$  are relatively prime.*

(c) *The shortest sub-Riemannian geodesic loops have length equal to  $2\pi$ .*

To obtain information about the full sub-Riemannian length spectrum in  $SU(3)$ , consider

$$H = a_1 X_1 + b_1 Y_1 + a_2 X_2 + b_2 Y_2 + a_3 X_3 + b_3 Y_3,$$

$$X^\perp = \frac{c_1}{2} R_3 + \frac{c_2}{2} R_2 = \begin{pmatrix} -c_1 i & 0 & 0 \\ 0 & c_2 i & 0 \\ 0 & 0 & (c_1 - c_2) i \end{pmatrix},$$

$$X = H + X^\perp, \text{ and } \gamma(t) = e^{tX} e^{-tX^\perp}.$$

The characteristic polynomial of  $X$  is

$$P(\lambda) = -\lambda^3 - p\lambda + qi,$$

where

$$p = \sum_{j=1}^3 (a_j^2 + b_j^2) + c_1^2 + c_2^2 - c_1 c_2,$$

and

$$q = c_2(a_3^2 + b_3^2 - a_1^2 - b_1^2 + c_1^2) - c_1(a_2^2 + b_2^2 - a_1^2 - b_1^2 + c_2^2) + 2(a_1 a_2 b_3 + a_1 b_2 a_3 - b_1 a_2 a_3 + b_1 b_2 b_3).$$

Note that  $p = \|H\|^2 + \|X^\perp\|^2$ . Formula (3-6) gives

$$(5-2) \quad \begin{aligned} c_1 + c_2 &\in 2\pi\mathbb{Z} && \text{if } a_1 + b_1 i \neq 0, \\ c_1 - 2c_2 &\in 2\pi\mathbb{Z} && \text{if } a_2 + b_2 i \neq 0, \\ 2c_1 - c_2 &\in 2\pi\mathbb{Z} && \text{if } a_3 + b_3 i \neq 0. \end{aligned}$$

To see the connection with the case of  $SU(2)$ , let us start with the following simple cases.

**Case 1:** Consider  $c_2 = -c_1$ ,  $a_2 = b_2 = a_3 = b_3 = 0$ . This corresponds to the case of  $SU(2)$  from the previous section and these geodesics are singular in  $SU(3)$ . Therefore the sub-Riemannian geodesics have the form (4-3) and the lengths  $\pi\sqrt{n^2 - m^2}$  from Proposition 4.1(d).

**Case 2:** Consider  $c_2 = 0$ ,  $a_2 = b_2 = a_3 = b_3 = 0$ . These geodesics are not contained in any copy of  $SU(2)$  and are regular in  $SU(3)$ . Here,  $c_1 = 2m\pi$ ,  $m \in \mathbb{Z}$ , and the eigenvalues of  $X$  are

$$-2m\pi i \quad \text{and} \quad \left(m\pi \pm \sqrt{a_1^2 + b_1^2 + m^2\pi^2}\right)i.$$

By Theorem 3.9(c) we have that

$$(5-3) \quad |c_1| = 2m\pi \quad \text{and} \quad a_1^2 + b_1^2 + m^2\pi^2 = n^2\pi^2,$$

where  $m, n \in \mathbb{N} \cup \{0\}$ ,  $m \leq n$ , are both odd or even. Therefore, the sub-Riemannian geodesic loop corresponding to (5-3) is

$$\gamma(t) = e^{t(a_1 X_1 + b_1 Y_1 + m R_3)} e^{-tm R_3},$$

and its length is  $\pi \sqrt{n^2 - m^2}$ .

**Case 3:** If at least two of  $a_j + b_j i$ ,  $j = 1, 2, 3$ , are not zero, then

$$c_1 = \frac{4n + 2m}{3} \pi, \quad c_2 = \frac{2n - 2m}{3} \pi,$$

where  $n, m \in \mathbb{Z}$ . Theorem 3.9(c) and the first Viète formula for the characteristic polynomial imply that the eigenvalues of  $X$  must have the form

$$\begin{aligned} \lambda_1 &= (-c_1 - 2k\pi)i, \\ \lambda_2 &= (c_2 + 2l\pi)i, \\ \lambda_3 &= (c_1 - c_2 + 2(k - l)\pi)i. \end{aligned} \tag{5-4}$$

The second Viète formula gives

$$\|H\| = \sqrt{c_1(4k - 2l)\pi + c_2(4l - 2k)\pi + 4(k^2 + l^2 - kl)\pi^2}. \tag{5-5}$$

From the third Viète formula we find

$$\begin{aligned} &4c_1 c_2 (k - l)\pi + 4c_1 l (2k - l)\pi^2 + 4c_2 k (k - 2l)\pi^2 + 2c_1^2 l\pi - 2c_2^2 k\pi + 8kl(k - l)\pi^3 \\ &= (a_3^2 + b_3^2 - a_1^2 - b_1^2)c_2 - (a_2^2 + b_2^2 - a_1^2 - b_1^2)c_1 \\ &\quad + 2(a_1 a_2 b_3 + a_1 b_2 a_3 - b_1 a_2 a_3 + b_1 b_2 b_3). \end{aligned}$$

The complexity of this formula hides its true geometric meaning; however, in the case when  $q = 0$ , it reduces to  $0 = 0$ , and we have the following eigenvalues for  $X$ :

$$0 \quad \text{and} \quad \pm \sqrt{\|H\|^2 + \|X^\perp\|^2} i.$$

Without loss of generality we can assume that  $\lambda_1 = 0$ . Then  $c_1 = 2k\pi$ , which implies that  $m = 3k - 2n$  and  $c_2 = 2(n - k)$ . From (5-5) it follows that

$$\|H\| = 2\pi \sqrt{(2k - l)^2 - nk + 2nl},$$

which in the case of  $k = l$  reduces to

$$\|H\| = 2\pi \sqrt{k^2 + nk} = 2\pi \sqrt{\left(k + \frac{n}{2}\right)^2 - \frac{n^2}{4}}.$$

This shows that, as expected, formula (5-5) includes the sub-Riemannian geodesic loop length spectrum of  $SU(2)$ .

Note that  $q = 0$  is satisfied if  $c_2 = 0$ ,  $a_1^2 + b_1^2 = a_2^2 + b_2^2$  and  $a_3 = b_3 = 0$ . As a numerical example we can give the sub-Riemannian geodesic loop of length  $8\pi$  described by

$$\gamma(t) = e^{\pi(5X_1 + \sqrt{7}Y_1 + 5X_2 + \sqrt{7}Y_2 + 3R_3)t} e^{-3\pi R_3 t}.$$

In conclusion, we have the following result.

**Proposition 5.2.** *In  $SU(3)$  the sub-Riemannian geodesic loops have lengths equal to*

$$2\pi \sqrt{\left( \frac{(2n+m)(2k-l)}{3} + \frac{(n-m)(2l-k)}{3} \right) + (k^2 + l^2 - kl)},$$

where  $m, n, k, l \in \mathbb{Z}$ .

### Acknowledgements

The authors are grateful to Richard Montgomery for helpful discussions and advice on exposition. We also thank the referees for valuable suggestions.

### References

- [Agrachev and Sarychev 1999] A. A. Agrachev and A. V. Sarychev, “Sub-Riemannian metrics: minimality of abnormal geodesics versus subanalyticity”, *ESAIM Control Optim. Calc. Var.* **4** (1999), 377–403. MR Zbl
- [Arvanitoyeorgos 2003] A. Arvanitoyeorgos, *An introduction to Lie groups and the geometry of homogeneous spaces*, Student Mathematical Library **22**, American Mathematical Society, Providence, RI, 2003. MR Zbl
- [Boscain et al. 2002] U. Boscain, T. Chambrion, and J.-P. Gauthier, “On the  $K + P$  problem for a three-level quantum system: optimality implies resonance”, *J. Dynam. Control Systems* **8**:4 (2002), 547–572. MR Zbl
- [Chang et al. 2011] D.-C. Chang, I. Markina, and A. Vasil’ev, “Hopf fibration: geodesics and distances”, *J. Geom. Phys.* **61**:6 (2011), 986–1000. MR Zbl
- [D’Andrea and Maffei 2016] A. D’Andrea and A. Maffei, “Commutators of small elements in compact semisimple groups and Lie algebras”, *J. Lie Theory* **26**:3 (2016), 683–690. MR
- [Domokos 2015] A. Domokos, “Subelliptic Peter–Weyl and Plancherel theorems on compact, connected, semisimple Lie groups”, *Nonlinear Anal.* **126** (2015), 131–142. MR Zbl
- [Duistermaat and Kolk 2000] J. J. Duistermaat and J. A. C. Kolk, *Lie groups*, Springer, 2000. MR Zbl
- [Helgason 2001] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Graduate Studies in Mathematics **34**, American Mathematical Society, Providence, RI, 2001. MR Zbl
- [Klapheck and VanValkenburgh 2019] D. Klapheck and M. VanValkenburgh, “The length spectrum of the sub-Riemannian three-sphere”, *Involve* **12**:1 (2019), 45–61.
- [Klingenberg 1978] W. Klingenberg, *Lectures on closed geodesics*, Grundlehren der Math. Wissenschaften **230**, Springer, 1978. MR Zbl
- [Knapp 1986] A. W. Knap, *Representation theory of semisimple groups*, Princeton Mathematical Series **36**, Princeton University Press, 1986. MR Zbl

[Montgomery 1994] R. Montgomery, “Singular extremals on Lie groups”, *Math. Control Signals Systems* **7**:3 (1994), 217–234. MR Zbl

[Montgomery 1995] R. Montgomery, “A survey of singular curves in sub-Riemannian geometry”, *J. Dynam. Control Systems* **1**:1 (1995), 49–90. MR Zbl

[Montgomery 2002] R. Montgomery, *A tour of sub-Riemannian geometries, their geodesics and applications*, Mathematical Surveys and Monographs **91**, American Mathematical Society, Providence, RI, 2002. MR Zbl

[Simon 1996] B. Simon, *Representations of finite and compact groups*, Graduate Studies in Mathematics **10**, American Mathematical Society, Providence, RI, 1996. MR Zbl

Received April 7, 2017. Revised December 11, 2017.

ANDRÁS DOMOKOS

domokos@csus.edu

MATTHEW KRAUEL

krauel@csus.edu

VINCENT PIGNO

vincent.pigno@csus.edu

COREY SHANBROM

corey.shanbrom@csus.edu

MICHAEL VANVALKENBURGH

mjv@csus.edu

.

(all authors)

DEPARTMENT OF MATHEMATICS AND STATISTICS  
CALIFORNIA STATE UNIVERSITY, SACRAMENTO  
SACRAMENTO, CA  
UNITED STATES

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Matthias Aschenbrenner  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[matthias@math.ucla.edu](mailto:matthias@math.ucla.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Wee Teck Gan  
Mathematics Department  
National University of Singapore  
Singapore 119076  
[matgwt@nus.edu.sg](mailto:matgwt@nus.edu.sg)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 296 No. 2 October 2018

---

Congruence subgroups and super-modular categories	257
PARSA BONDERSON, ERIC C. ROWELL, ZHENGHAN WANG and QING ZHANG	
On the Chow ring of the stack of truncated Barsotti–Tate Groups	271
DENNIS BROKEMPER	
Hyperbolic manifolds containing high topological index surfaces	305
MARION CAMPISI and MATT RATHBUN	
Length spectra of sub-Riemannian metrics on compact Lie groups	321
ANDRÁS DOMOKOS, MATTHEW KRAUEL, VINCENT PIGNO, COREY SHANBROM and MICHAEL VANVALKENBURGH	
The action of the Hecke operators on the component groups of modular Jacobian varieties	341
TAEKYUNG KIM and HWAJONG YOO	
Liouville theorems, volume growth, and volume comparison for Ricci shrinkers	357
LI MA	
Presentations of generalisations of Thompson’s group $V$	371
CONCHITA MARTÍNEZ-PÉREZ, FRANCESCO MATUCCI and BRITA NUCINKIS	
Localization functors and cosupport in derived categories of commutative Noetherian rings	405
TSUTOMU NAKAMURA and YUJI YOSHINO	
Zeta integrals for $\mathrm{GSp}(4)$ via Bessel models	437
RALF SCHMIDT and LONG TRAN	
Toric surfaces over an arbitrary field	481
FEI XIE	
Erratum for the article The Johnson–Morita theory for the rings of Fricke characters of free groups	509
TAKAO SATOH	



0030-8730(201810)296:2;1-S