

*Pacific  
Journal of  
Mathematics*

**LIUVILLE THEOREMS, VOLUME GROWTH, AND  
VOLUME COMPARISON FOR RICCI SHRINKERS**

LI MA

# LIUVILLE THEOREMS, VOLUME GROWTH, AND VOLUME COMPARISON FOR RICCI SHRINKERS

LI MA

**We study volume growth, a Liouville theorem for  $f$ -harmonic functions, and a volume comparison property of unit balls in complete noncompact gradient Ricci shrinkers and gradient steady Ricci solitons. We also study integral properties of  $f$ -harmonic functions and harmonic functions on complete manifolds, such as the Ricci–Einstein solitons.**

## 1. Introduction

In the study of Ricci flow on a compact Riemannian manifold, because of its complicated nonlinearity, one meets singularities of the flow in finite time. After blowing up, one expects to get self-similar Ricci shrinkers or steady Ricci solitons [Hamilton 1995]. In the case of a type-I singularity in dimension three, one gets nontrivial gradient Ricci shrinkers via the use of the Ivey–Hamilton pinching estimate, and the classification of this type of self-similar Ricci shrinker was done by G. Perelman [2000]. In the case of a type-I singularity of dimension four, A. Naber [2010] showed that one gets a gradient Ricci shrinker, and nontrivial properties of this Ricci shrinker have been studied by others. These solitons can be considered as special examples of weighted Riemannian manifolds or metric measure spaces [Bakry and Émery 1985; Wei and Wylie 2009; Lott and Villani 2009; Chen 2009; Chow et al. 2006; Lichnerowicz 1970; Lott 2003; Ma 2013; Sturm 2006a; 2006b; Yang 2009; Munteanu and Wang 2011; 2014; 2015]. Because of the importance of four-dimensional Ricci shrinkers, many people study various properties of them; see for example [Ma 2013; Cao and Zhou 2010; Cao 2007; 2010; Carrillo and Ni 2009; Munteanu and Wang 2015; Haslhofer and Müller 2011]. In this paper, we examine three questions about Ricci shrinking and steady solitons, in particular for Ricci shrinkers which are the complete Riemannian manifold  $(M, g)$  such that  $\text{Ric}_f := \text{Ric} + D^2 f = \lambda g$  on  $M$ , where  $\text{Ric}$  is the Ricci curvature of  $(M, g)$ ,  $f : M \rightarrow \mathbb{R}$  is a smooth function in  $M$ , and  $\lambda > 0$  is a constant. One question under

---

This research is partially supported by the National Natural Science Foundation of China, grant nos. 11771124 and 11271111.

MSC2010: 53C21, 53C44, 53C25.

Keywords: Ricci shrinkers,  $f$ -harmonic functions, Liouville theorem, volume comparison.

consideration in this paper is about the volume comparison of unit balls on Ricci shrinkers. The other two are about  $f$ -harmonic functions and harmonic functions with finite energy. The volume comparison of unit balls is an important step to understanding the volume growth of geodesic balls in gradient Ricci shrinkers. The Liouville theorems for  $f$ -harmonic and harmonic functions with finite energy are important to understanding the connectivity at infinity about gradient Ricci solitons; see [Munteanu and Wang 2015].

We have the following new results. The first is the volume comparison of unit balls at any point  $x \in M$  with a unit ball at a fixed point  $p \in M$ , which deals with the injectivity radius decay from the point  $p$  to the point  $x$  and is important to understanding the topology of the underlying manifold at infinity.

**Theorem 1.** *On the complete noncompact gradient Ricci shrinker  $(M, g, f)$  with Ricci curvature bounded by  $-(n - 1)k^2$  for some constant  $k \geq 0$ , we have*

$$\text{Vol}(B_x(1)) \geq \exp(-\sqrt{c_0(n - 1)R}) \text{Vol}(B_p(1)),$$

where  $c_0$  is a uniform constant which does not depend on  $x$  and  $R = d(p, x) > R_0$  for some uniform constant  $R_0 > 1$ .

The above result is motivated by the work of Ovidiu Munteanu and Jiaping Wang [2014] and the result is not sharp, as pointed out by the referee. In fact, if the Ricci curvature is bounded from below, the logarithmic Sobolev inequality of [Carrillo and Ni 2009] implies a uniform lower bound of the volume of unit balls,  $\text{Vol}(B_x(1)) \geq \frac{1}{c} \text{Vol}(B_p(1))$ . However, from our argument, our result is true on a complete smooth measure space of dimension  $n$  with  $\text{Ric}_f \geq \frac{1}{2}g$  and  $|\nabla f|^2 \leq f$  and with Ricci curvature bounded below by  $-(n - 1)k^2$  for some constant  $k \geq 0$  on  $M$ . Notice that with the extra bound on Ricci curvature, our result improves the result of Lemma 5.2 in [Munteanu and Wang 2014]. We remark that the above result is still true for steady Ricci solitons. See Theorem 5 in Section 2.

We want to understand the topology and geometry from properties of (super- or sub-) harmonic functions and  $f$ -harmonic functions on Ricci solitons. By a well-known argument, we know that there is no nontrivial positive  $f$ -harmonic function on a gradient Ricci shrinker. In fact there is no nonconstant positive  $f$ -superharmonic function  $u$  ( $\Delta_f u := \Delta u - \nabla f \cdot \nabla u \leq 0$ ) on the complete Riemannian manifold  $(M, g)$  with  $\text{Ric}_f \geq \frac{1}{2}g$  on  $M$ . The process of proving this is below. Recall that the weighted volume of  $(M, g)$  is finite [Morgan 2005]; i.e.,  $V_f(M) := \int_M e^{-f} dv_g < \infty$ . Assume  $u > 0$  is such a positive  $f$ -superharmonic function on  $M$ . Let  $v = \log u$ . Then we have

$$\Delta_f v = \frac{\Delta_f u}{u} - |\nabla v|^2 \leq -|\nabla v|^2.$$

Thus, for any cut-off function  $\phi \geq 0$  on the ball  $B_p(2R) \subset M$  with  $\phi = 1$  in  $B_p(R)$ , we have

$$\int_M |\nabla v|^2 \phi^2 e^{-f} dv_g \leq 2 \int_M \phi \nabla \phi \cdot \nabla v e^{-f} dv_g.$$

By the Cauchy–Schwarz inequality we get

$$\int_M |\nabla v|^2 \phi^2 e^{-f} dv_g \leq 4 \int_M |\nabla \phi|^2 e^{-f} dv_g \leq \frac{16}{R^2} V_f(M) \rightarrow 0$$

as  $R \rightarrow \infty$ . Hence,  $|\nabla v|^2 = 0$  in  $M$ , which implies that  $u$  is a constant on  $M$ .

Although this result is known to experts, we cannot find it in the literature. We formulate it as below.

**Proposition 2.** *Let  $(M, g, f)$  be a complete Riemannian manifold  $(M, g)$  with potential function  $f$  satisfying  $\text{Ric}_f \geq \frac{1}{2}g$  on  $M$ . Then there is no nonconstant positive  $f$ -superharmonic function on  $(M, g)$ .*

We are now trying to find another kind of Liouville theorem for an  $f$ -harmonic function with weighted finite energy on a gradient Ricci shrinker. We show that as a direct consequence of a Bochner-type formula, see [Ma and Du 2010], we have the following Liouville-type theorem.

**Theorem 3.** *Let  $(M, g)$  be a complete noncompact Riemannian manifold such that  $\text{Ric}_f \geq h(x)g$  for some potential functions  $f(x)$  and some nontrivial nonnegative function  $h(x)$  in  $M$ . There is no nontrivial  $f$ -harmonic function  $u$  defined in  $(M, g)$  with weighted finite energy; i.e.,*

$$\int_M |\nabla u|^2 e^{-f} dv_g < \infty.$$

The proof of this result is given in Section 3. We remark that when one studies self-similar solutions to the Ricci–Einstein-type flow

$$\partial_t g = -2 \text{Rc}(g) + \gamma Rg,$$

where  $\gamma$  is a physical constant and  $R = R(g)$  is the scalar curvature of the manifold  $(M, g)$ , one gets the soliton equation of it in the form

$$\text{Ric}_f = \gamma Rg + \lambda g,$$

which gives what may be called Ricci–Einstein solitons. Theorem 3 can be used for these solitons.

**Proposition 4.** *Fix any  $p \in M$ . Assume that the complete noncompact Riemannian manifold  $(M, g)$  satisfies  $\text{Ric}_f \geq h(x)g$  for some potential functions  $f(x)$  and some smooth function  $h(x)$  with*

$$|\nabla f(x)| \leq \alpha d(x, p) + b$$

for some uniform constants  $\alpha \geq 0$  and  $b > 0$  in  $M$ . Then for any harmonic function  $u$  with finite energy, i.e.,

$$\int_M |\nabla u|^2 < \infty,$$

we have the integral inequality

$$\int_M |\nabla^2 u|^2 + \int_M h(x) |\nabla u|^2 \leq \frac{1}{2} \int_M \Delta f |\nabla u|^2.$$

As a consequence, when  $\Delta f(x) \leq 2h(x)$  on  $M$ , we have  $D^2u = 0$  in  $M$ ; i.e.,  $\nabla u$  is a parallel vector field on  $M$ .

We now give a few remarks related to this result.

(1) Naber [2010] proved that for the weighted smooth metric space  $(M, g, f)$  satisfying  $\text{Ric}_f \leq \frac{1}{2}g$  and  $|\text{Ric}| \leq C$ , there exists  $\alpha > 0$  such that if  $\Delta_f u := \Delta u - \nabla f \cdot \nabla u = 0$  on  $M$  with  $|u(x)| \leq A \exp(\alpha d(x, p)^2)$  for some  $A > 0$  and  $p \in M$ , then  $u$  is a constant.

(2) Munteanu and Sesum [2013] proved that for gradient shrinking Kähler–Ricci solitons, if the harmonic function  $u$  has finite energy, i.e.,  $\int_M |\nabla u|^2 < \infty$ , then  $u$  is a constant. As a consequence of this result, they showed that such a manifold has at most one nonparabolic end; see [Munteanu and Sesum 2013] for the definition of nonparabolic end. In the earlier work [Munteanu and Wang 2011], they proved that on a weighted smooth metric space  $(M, g, f)$  satisfying  $\text{Ric}_f \geq 0$  and  $f$  is a bounded function, any sublinear growth  $f$ -harmonic function on  $M$  must be a constant.

(3) Some consequences of Proposition 4 are given in Section 3.

Here is the plan of the paper. We study the volume comparison of unit balls of gradient Ricci shrinkers and gradient steady Ricci solitons in Section 2. We prove Proposition 2 and Theorem 3 in Section 3. In the last section, we consider integral properties for harmonic functions on Ricci solitons and prove Proposition 4.

## 2. Volume comparison of unit balls

We now prove Theorems 1 and 5. We use an idea similar to the proof of Theorem 2.3 in [Munteanu and Wang 2014].

We first give a proof of an improved volume comparison of unit balls on the weighted Riemannian manifold of shrinking type.

*Proof of Theorem 1.* Again, we take any point  $x \in M$  and express the volume form in the geodesic polar coordinates centered at  $x$  as

$$dV|_{\exp_x(r\xi)} = J(x, r, \xi) dr d\xi$$

for  $r > 0$  and  $\xi \in S_x M$  a unit tangent vector at  $x$ . We let  $R = d(p, x)$  and omit the dependence of the geometric quantities on  $\xi$ . Let  $R = d(p, x) > 2$ . Let  $\gamma(s)$  be the minimizing geodesic starting from  $x$  connecting  $\gamma(0) = x$  to any point  $\gamma(T) \in B_p(1)$ . By the triangle inequality we know that  $T \in [R - 1, R + 1]$ . It is well known [Li 1993] that along the minimizing geodesic curve  $\gamma$ , we have

$$m'(r) + \frac{1}{n-1}m^2(r) + \text{Ric}(\partial_r, \partial_r) \leq 0,$$

where  $r > 0$  and  $m = m(r) = \frac{d}{dr}(\log J)(r)$ . Using the Ricci soliton equation  $\text{Ric}_f = \frac{1}{2}g$  we immediately obtain

$$m'(r) + \frac{1}{n-1}m^2(r) \leq -\frac{1}{2} + f''(r).$$

Integrating this relation, we get for  $r > 1$ ,

$$m(r) + \frac{1}{n-1} \int_1^r m^2(t) dt \leq -\frac{r-1}{2} + f'(r) - f'(1) + m(1).$$

Recall the well-known fact (see (2.3) and (2.8) in [Cao and Zhou 2010] in different notation) that for any  $T > \tau > 0$ , we have

$$(1) \quad -\frac{1}{2}(T - \tau) - c \leq f'(\tau) \leq -\frac{1}{2}(T - \tau) + c$$

and  $|f'(T)| \leq c$  since  $\gamma(T) \in B_p(1)$ . Here and everywhere in the proofs,  $c$  denotes a constant depending only on the dimension  $n$  and  $f(p)$ . By this we know

$$-\frac{r-1}{2} + f'(r) - f'(1) \leq c_0$$

for some uniform constant  $c_0 > 0$ .

Using the Ricci curvature lower bound, a standard argument shows, see (1.1.8) in [Schoen and Yau 1994], that there is a uniform constant  $c_1 > 0$  such that  $m(s) \leq c_1$  for  $s \geq \frac{1}{2}$ . Then we have for another uniform constant  $c_0 > 0$  and for any  $r > 1$ ,

$$m(r) + \frac{1}{n-1} \int_1^r m^2(t) dt \leq c_0.$$

By the Cauchy–Schwarz inequality we obtain

$$(2) \quad m(r) + \frac{1}{(n-1)r} \left( \int_1^r m(t) dt \right)^2 < c$$

for any  $c > c_0$  and  $r > 1$ .

**Claim.** For any  $r > 1$ ,

$$(3) \quad \int_1^r m(t) dt \leq \sqrt{c(n-1)r}.$$

In fact, let

$$v(t) = \sqrt{c(n-1)t} - \int_1^t m(r) dr.$$

Then

$$v'(t) = \frac{\sqrt{c(n-1)}}{2\sqrt{t}} - m(t).$$

Clearly  $v(1) > 0$  by the fact that  $c > c_0$ . Suppose that  $v$  is negative somewhere for  $t > 1$ . Let  $d > 1$  be the first zero point of  $v$ ; i.e.,  $v(d) = 0$ . Then by the choice of  $d$ , we have  $v'(d) \leq 0$ . That is,

$$\int_1^d m(t) dt = \sqrt{c(n-1)d}$$

and

$$m(d) \geq \frac{\sqrt{c(n-1)}}{2\sqrt{d}}.$$

By direct computation we know

$$m(d) + \frac{1}{(n-1)d} \left( \int_1^d m(t) dt \right)^2 \geq \frac{\sqrt{c(n-1)}}{2\sqrt{d}} + c,$$

which is a contradiction with (2) at  $r = d$ .

The relation (3) implies

$$\log J(x, r, \xi) / J(x, 1, \xi) \leq \sqrt{c(n-1)r}$$

for any  $r > 1$  and we have at  $r = R$ ,

$$J(x, 1, \xi) \geq \exp(-\sqrt{c(n-1)R}) J(x, R, \xi).$$

Integrating over the unit tangent vectors  $\xi$  we get

$$\text{Area}(\partial B_x(1)) \geq \exp(-\sqrt{c(n-1)R}) \text{Vol}(B_p(1)),$$

where  $R = d(p, x) > 2$ . Similarly we have

$$\text{Area}(\partial B_x(s)) \geq \exp(-\sqrt{c(n-1)R}) \text{Vol}(B_p(1))$$

for any  $s \in [\frac{1}{2}, 1]$ . Hence, we have

$$\text{Vol}(B_x(1)) \geq \exp(-\sqrt{c(n-1)R}) \text{Vol}(B_p(1)). \quad \square$$

A similar argument gives us the result below for gradient steady Ricci solitons.

**Theorem 5.** *On the complete noncompact Riemannian manifold  $(M, g, f)$  with Ricci curvature bounded below by  $-(n-1)k^2$  for some constant  $k \geq 0$  and  $\text{Rc}_f \geq 0$  on  $M$  and  $|\nabla f| \leq C$  on  $M$  for some uniform constant  $C > 0$ , we have*

$$\text{Vol}(B_x(1)) \geq \exp(-\sqrt{c_0(n-1)R}) \text{Vol}(B_p(1)),$$

where  $c_0$  is a uniform constant which does not depends on  $p$  and  $R = d(p, x) > R_0$  for some uniform constant  $R_0 > 1$ .

The proof of [Theorem 5](#) is almost the same as the previous proof. So we only present the necessary modification and use the same notation as above. Again we have along the minimizing geodesic curve  $\gamma$ ,

$$m'(r) + \frac{1}{n-1}m^2(r) + \text{Ric}(\partial_r, \partial_r) \leq 0.$$

Using the relation  $\text{Ric}_f \geq 0$  we immediately obtain

$$m'(r) + \frac{1}{n-1}m^2(r) \leq f''(r).$$

Integrating this relation we get for  $t > 1$  and  $s \in [\frac{1}{2}, 1]$ ,

$$m(t) + \frac{1}{n-1} \int_s^t m^2(r) dr \leq f'(t) - f'(1) + m(s).$$

This means that for any  $t > 1$  and  $s = 1$ ,

$$m(t) + \frac{1}{n-1} \int_1^t m^2(r) dr \leq C_0$$

for some uniform constant  $C_0$ . The remaining part of the proof is the same as the proof of [Theorem 1](#) and we omit the details.

We now consider the volume growth of geodesic balls in manifolds with density and we show that for  $(M, g, e^{-f} dv)$  a complete smooth metric measure space of dimension  $n$  with  $\text{Ric}_f \geq \frac{1}{2}$ ,  $|\nabla f|^2 \leq f$ , and also with both Ricci curvature and  $\Delta f$  bounded from above, the volume growth of geodesic balls is in polynomial order (which may be smaller than  $n$ ).

**Proposition 6.** *Let  $(M, g, e^{-f} dv)$  be a complete smooth metric measure space of dimension  $n$ . Assume that  $\text{Ric}_f \geq \frac{1}{2}$ ,  $|\nabla f|^2 \leq f$ . Assume further that  $\Delta f \leq K$  and  $\text{Ric} \leq K_1$  for some constants  $K > 0$  and  $K_1 > 0$ . Then for some  $p \in M$ , the volume growth of geodesic balls  $B_p(r)$  is of polynomial order; i.e., there is a uniform constant  $C > 0$  such that*

$$V(r) \leq Cr^{2K}.$$

*Proof.* Recall that under the conditions  $\text{Ric}_f \geq \frac{1}{2}$  and  $|\nabla f|^2 \leq f$ , there are a point  $p \in M$ , two constants  $r_0 > 0$  and  $a$  depending only on  $n$  and  $f(p)$  such that

$$(4) \quad \left(\frac{1}{2}d(x, p) - a\right)^2 \leq f(x) \leq \left(\frac{1}{2}d(x, p) + a\right)^2.$$

This is from Proposition 4.2 in the interesting paper [\[Munteanu and Wang 2014\]](#). By this we know that  $|\nabla f(x)| \leq \frac{1}{2}d(x, p) + a$ . We may assume that  $d(x, p) > 2$ . Consider any minimizing normal geodesic  $\gamma(s)$ ,  $0 \leq s \leq r := d(x, p)$ , starting from



the point  $\gamma(0) = p$  to the point  $\gamma(r) = x$ . Let  $X = \dot{\gamma}(s)$ . By the second variation formula of arc length we know

$$\int_0^r \phi^2 \operatorname{Ric}(X, X) ds \leq (n - 1) \int_0^r |\dot{\phi}(s)|^2 ds$$

for any  $\phi \in C_0^{1-}([0, r])$ . Let  $\phi(s) = s$  on  $[0, 1]$ ,  $\phi(s) = r - s$  on  $[r - 1, r]$ , and  $\phi(s) = 1$  on  $[1, r - 1]$ . Then we have

$$\int_0^r \operatorname{Ric}(X, X) ds = \int_0^r \phi^2 \operatorname{Ric}(X, X) ds + \int_0^r (1 - \phi^2) \operatorname{Ric}(X, X) ds.$$

We derive via the use of  $\operatorname{Ric} \leq K_1$ , and by an argument similar to the proof before (2.8) in [Cao and Zhou 2010], that

$$\int_0^r \operatorname{Ric}(X, X) ds \leq 2(n - 1) + 2K_1.$$

Since

$$\nabla_X \dot{f} = \nabla^2 f(X, X) \geq \frac{1}{2} - \operatorname{Ric}(X, X),$$

integrating it from 0 to  $r$ , we get

$$(5) \quad \dot{f}(r) = \frac{1}{2}r - \int_0^r \operatorname{Ric}(X, X) ds \geq \frac{1}{2}r - c$$

for some constant  $c$  depending only on  $K_1, n$  and  $f(p)$ . Hence,

$$|\nabla f|(x) \geq \dot{f}(r) \geq \frac{1}{2}d(x, p) - c.$$

Define

$$\rho(x) = 2\sqrt{f(x)}.$$

Then,

$$|\nabla \rho| = \frac{|\nabla f|}{\sqrt{f}} \leq 1.$$

Let, for  $r > 0$  large,

$$D(r) = \{x \in M; \rho(x) \leq r\}, \quad V(r) = \operatorname{Vol}(D(r)).$$

As in [Cao and Zhou 2010], by using the coarea formula we have

$$V(r) = \int_0^r ds \int_{\partial D(r)} \frac{1}{|\nabla \rho|} dA,$$

$$V'(r) = \int_{\partial D(r)} \frac{1}{|\nabla \rho|} dA = \frac{r}{2} \int_{\partial D(r)} \frac{1}{|\nabla f|} dA.$$

By the assumption  $\Delta f \leq K$  and the divergence theorem we have

$$2KV(r) \geq 2 \int_{D(r)} \Delta f = 2 \int_{\partial D(r)} |\nabla f| dA.$$

By (5) we know that on  $\partial D(r)$ , there is a uniform constant  $C > 2$  such that for  $r \geq 2C$ ,

$$|\nabla f|^2 \geq f - C.$$

Then we have

$$2 \int_{\partial D(r)} |\nabla f| \, dA \geq 2 \int_{\partial D(r)} \frac{f - C}{|\nabla f|} \, dA.$$

The right side of above inequality is greater than or equal to

$$(r - 2)V'(r).$$

Hence we have

$$2K V(r) \geq (r - 2)V'(r),$$

which then implies

$$V(r) \leq V(2C)r^{2K}$$

for  $r > 2C$ . □

We remark that the above argument is motivated by the proof of the volume growth estimate in [Cao and Zhou 2010]. Our result is different from the deep result Theorem 1.4 in [Munteanu and Wang 2014] in the case when the constant  $2K$  is smaller than  $n$ .

### 3. Harmonic and $f$ -harmonic functions on Ricci–Einstein solitons

We now prove Theorem 3. We wish that we could use the Caccioppoli argument (see the proof of Proposition 8.1 in [Naber 2010], with the use of Lemma 2.2 replaced by Proposition 4.2 in [Munteanu and Wang 2014]) to conclude that with some decay assumption such as finite energy, an  $f$ -harmonic function  $u$  is a constant function on  $M$ . However, we have a simpler proof of this result below.

*Proof of Theorem 3.* Recall the Bochner formula for the  $f$ -harmonic function  $u : M \rightarrow \mathbb{R}$ ,

$$\frac{1}{2} \Delta_f |\nabla u|^2 = |\nabla^2 u|^2 + \text{Rc}_f(\nabla u, \nabla u).$$

By our assumption that  $\text{Ric}_f \geq h(x)g$ , we know

$$\frac{1}{2} \Delta_f |\nabla u|^2 \geq |\nabla^2 u|^2 + h(x)|\nabla u|^2.$$

Let  $\phi$  be the standard cut-off function on  $B_p(2r)$  and let  $dm = \exp(-f) \, dv_g$ . Then we have

$$\int_M (|\nabla^2 u|^2 + h(x)|\nabla u|^2) \phi \, dm \leq \int_M \left(\frac{1}{2} \Delta_f \phi\right) |\nabla u|^2 \, dm.$$

The right side goes to zero as  $r \rightarrow \infty$ . Hence we have

$$\int_M (|\nabla^2 u|^2 + h(x)|\nabla u|^2) dm = 0,$$

which implies that  $u$  is a constant. □

Let  $(M, g)$  be a complete noncompact Riemannian manifold of dimension  $n$ . Fix  $p \in M$ . In this section we always assume that  $(M, g)$  satisfies  $\text{Ric}_f \geq h(x)g$  for some function  $h(x)$  and  $|\nabla f| \leq \alpha d(x, p) + b$ . Then we have  $R + \Delta f \geq nh(x)$  on  $M$ . We study the  $L^2$  estimate for the Hessian matrix for harmonic functions with finite energy.

*Proof of Proposition 4.* Let  $u : M \rightarrow R$  be a harmonic function on  $(M, g, f)$  with finite energy

$$\int_M |\nabla u|^2 < \infty.$$

Recall the Bochner formula for the harmonic function  $u : M \rightarrow R$ ,

$$\frac{1}{2} \Delta |\nabla u|^2 = |\nabla^2 u|^2 + \text{Rc}(\nabla u, \nabla u).$$

Using the assumption  $\text{Ric}_f \geq h(x)g$  we have

$$(6) \quad \frac{1}{2} \Delta |\nabla u|^2 \geq |\nabla^2 u|^2 + h(x)|\nabla u|^2 - \nabla^2 f(\nabla u, \nabla u).$$

Recall the Hessian matrix  $\nabla^2 f = (f_{ij})$  in local coordinates  $(x_i)$  in  $M$ .

Let  $\phi = \phi_r$  be the cut-off function on  $B_{2r}(p)$ . We write by  $o(1)$  the quantities such that  $o(1) \rightarrow 0$  as  $r \rightarrow \infty$ . Then, we have

$$(7) \quad \int_M (|\nabla^2 u|^2 + h(x)|\nabla u|^2) \phi^2 \leq \int_M (\frac{1}{2} \Delta |\nabla u|^2 + \nabla^2 f(\nabla u, \nabla u)) \phi^2.$$

By direct computation we have, for  $\epsilon > 0$  small,

$$\int_M (\frac{1}{2} \Delta |\nabla u|^2) \phi^2 = -2 \int_M \phi D^2 u(\nabla u, \nabla \phi) \leq \epsilon \int_M |\nabla^2 u|^2 \phi^2 + o(1).$$

Using  $|\nabla f| \leq \alpha d(x, p) + b$  and integrating by parts, we obtain

$$\int_M f_{ij} u_i u_j \phi^2 = - \int f_i u_{ij} u_j \phi^2 + o(1).$$

Furthermore, we have

$$\int_M f_{ij} u_i u_j \phi^2 = \frac{1}{2} \int_M \Delta f |\nabla u|^2 \phi^2 + o(1).$$

Hence by (6) we have

$$\int_M ((1 - \epsilon)|\nabla^2 u|^2 + h(x)|\nabla u|^2) \phi^2 - \frac{1}{2} \int_M \Delta f |\nabla u|^2 \phi^2 \leq o(1).$$

That is,

$$\int_M (1 - \epsilon)|\nabla^2 u|^2 \phi^2 + \int_M h(x)|\nabla u|^2 \phi^2 \leq \frac{1}{2} \int_M \Delta f |\nabla u|^2 \phi^2 + o(1).$$

Sending  $r \rightarrow \infty$  and letting  $\epsilon \rightarrow 0$ , we obtain

$$(8) \quad \int_M |\nabla^2 u|^2 + \int_M h(x)|\nabla u|^2 \leq \frac{1}{2} \int_M \Delta f |\nabla u|^2.$$

Note that when  $\Delta f \leq 2h(x)$  on  $M$ , by (8) we have  $D^2u = 0$  in  $M$ ; i.e.,  $\nabla u$  is a parallel vector field on  $M$ . □

We remark that when  $\text{Ric}_f = \lambda g$  (and  $R + \Delta f = n\lambda$ ) with  $\lambda$  being a constant, by the proof of Proposition 4 we have

$$\int_M |\nabla^2 u|^2 + \frac{1}{2} \int_M R |\nabla u|^2 \leq \int_M \frac{(n-2)\lambda}{2} |\nabla u|^2.$$

Note that the assumption about the potential function  $f$  in Proposition 4 is true on the steady soliton  $(M, g)$ ; see [Hamilton 1995]. We now give an application of this integral inequality (8). A special case of the Liouville-type theorem below, due to Munteanu and Sesum [2013, Theorem 4.1], can be derived from the integral estimate (8).

**Proposition 7.** *Let  $n = 2$ . Assume that the complete noncompact surface  $(M, g, f)$  satisfies  $\text{Ric}_f = h(x)g$  on  $M$  with  $R \geq 0$ , and  $|\nabla f| \leq b$  for some  $b > 0$  in  $M$ . Then there is no nontrivial harmonic function on  $(M, g)$  with finite energy on  $(M, g)$ .*

*Proof.* Note that  $R + \Delta f = nh(x) = 2h(x)$  in  $M$ . Then  $\Delta f = 2h(x) - R$  in  $M$ . By (8) we have

$$\int_M |\nabla^2 u|^2 + \frac{1}{2} \int_M R |\nabla u|^2 \leq 0.$$

If  $R = 0$ , then  $(M, g)$  is flat and the result follows from Theorem 4.1 in [Munteanu and Sesum 2013].

We may assume  $R > 0$  in  $M$ . We argue by contradiction. Assume that there is a nontrivial harmonic function with finite energy on  $(M, g)$ . By (8) we know

$$\int_M |\nabla^2 u|^2 + \frac{1}{2} \int_M R |\nabla u|^2 = 0.$$

Hence  $\nabla u$  is a parallel vector field on  $M$  and  $R = 0$ , a contradiction with  $R > 0$ . □

We remark that we can give a new proof of Theorem 4.1 in [Munteanu and Sesum 2013], which is on a gradient steady Ricci soliton. It says that there is no nontrivial harmonic function with finite energy on the steady Ricci soliton  $(M, g)$ . The proof is below. We may assume that  $(M, g, f)$  is a nontrivial steady Ricci

soliton. Recall that it is well known that either  $R > 0$  or  $R = 0$  on  $M$ . By (8), we have  $R = 0$ , and then

$$\Delta_f R = -2|\text{Ric}|^2,$$

and we know that  $\text{Ric} = 0$  on  $M$ . By [Schoen and Yau 1994], we know that there is no nontrivial harmonic function with finite energy.

### Acknowledgements

The author thanks the anonymous referees very much for useful suggestions. This work was done when the author was visiting the University of Minnesota in September 2016; he would like to thank Prof. Jiaping Wang for helpful discussions and the hospitality of the School of Mathematics.

### References

- [Bakry and Émery 1985] D. Bakry and M. Émery, “Diffusions hypercontractives”, pp. 177–206 in *Séminaire de Probabilités XIX* (Strasbourg, 1983/84), edited by J. Azéma and M. Yor, Lecture Notes in Math. **1123**, Springer, 1985. [MR](#) [Zbl](#)
- [Cao 2007] X. Cao, “Compact gradient shrinking Ricci solitons with positive curvature operator”, *J. Geom. Anal.* **17**:3 (2007), 425–433. [MR](#) [Zbl](#)
- [Cao 2010] H.-D. Cao, “Recent progress on Ricci solitons”, pp. 1–38 in *Recent advances in geometric analysis*, edited by Y.-I. Lee et al., Adv. Lect. Math. (ALM) **11**, International, Somerville, MA, 2010. [MR](#) [Zbl](#)
- [Cao and Zhou 2010] H.-D. Cao and D. Zhou, “On complete gradient shrinking Ricci solitons”, *J. Differential Geom.* **85**:2 (2010), 175–185. [MR](#) [Zbl](#)
- [Carrillo and Ni 2009] J. A. Carrillo and L. Ni, “Sharp logarithmic Sobolev inequalities on gradient solitons and applications”, *Comm. Anal. Geom.* **17**:4 (2009), 721–753. [MR](#) [Zbl](#)
- [Chen 2009] B.-L. Chen, “Strong uniqueness of the Ricci flow”, *J. Differential Geom.* **82**:2 (2009), 363–382. [MR](#) [Zbl](#)
- [Chow et al. 2006] B. Chow, P. Lu, and L. Ni, *Hamilton’s Ricci flow*, Graduate Studies in Mathematics **77**, American Mathematical Society, Providence, RI, 2006. [MR](#) [Zbl](#)
- [Hamilton 1995] R. S. Hamilton, “The formation of singularities in the Ricci flow”, pp. 7–136 in *Surveys in differential geometry, II* (Cambridge, MA, 1993), edited by S.-T. Yau, International, Cambridge, MA, 1995. [MR](#) [Zbl](#)
- [Haslhofer and Müller 2011] R. Haslhofer and R. Müller, “A compactness theorem for complete Ricci shrinkers”, *Geom. Funct. Anal.* **21**:5 (2011), 1091–1116. [MR](#) [Zbl](#)
- [Li 1993] P. Li, *Lecture notes on geometric analysis*, Lecture Notes Series **6**, Seoul National University, Seoul, 1993. [MR](#) [Zbl](#)
- [Lichnerowicz 1970] A. Lichnerowicz, “Variétés riemanniennes à tenseur C non négatif”, *C. R. Acad. Sci. Paris Sér. A-B* **271** (1970), A650–A653. [MR](#) [Zbl](#)
- [Lott 2003] J. Lott, “Some geometric properties of the Bakry–Émery–Ricci tensor”, *Comment. Math. Helv.* **78**:4 (2003), 865–883. [MR](#) [Zbl](#)
- [Lott and Villani 2009] J. Lott and C. Villani, “Ricci curvature for metric-measure spaces via optimal transport”, *Ann. of Math. (2)* **169**:3 (2009), 903–991. [MR](#) [Zbl](#)

- [Ma 2013] L. Ma, “Remarks on compact shrinking Ricci solitons of dimension four”, *C. R. Math. Acad. Sci. Paris* **351**:21-22 (2013), 817–823. [MR](#) [Zbl](#)
- [Ma and Du 2010] L. Ma and S.-H. Du, “Extension of Reilly formula with applications to eigenvalue estimates for drifting Laplacians”, *C. R. Math. Acad. Sci. Paris* **348**:21-22 (2010), 1203–1206. [MR](#) [Zbl](#)
- [Morgan 2005] F. Morgan, “Manifolds with density”, *Notices Amer. Math. Soc.* **52**:8 (2005), 853–858. [MR](#) [Zbl](#)
- [Munteanu and Sesum 2013] O. Munteanu and N. Sesum, “On gradient Ricci solitons”, *J. Geom. Anal.* **23**:2 (2013), 539–561. [MR](#) [Zbl](#)
- [Munteanu and Wang 2011] O. Munteanu and J. Wang, “Smooth metric measure spaces with non-negative curvature”, *Comm. Anal. Geom.* **19**:3 (2011), 451–486. [MR](#) [Zbl](#)
- [Munteanu and Wang 2014] O. Munteanu and J. Wang, “Geometry of manifolds with densities”, *Adv. Math.* **259** (2014), 269–305. [MR](#) [Zbl](#)
- [Munteanu and Wang 2015] O. Munteanu and J. Wang, “Geometry of shrinking Ricci solitons”, *Compos. Math.* **151**:12 (2015), 2273–2300. [MR](#) [Zbl](#)
- [Naber 2010] A. Naber, “Noncompact shrinking four solitons with nonnegative curvature”, *J. Reine Angew. Math.* **645** (2010), 125–153. [MR](#) [Zbl](#)
- [Perelman 2000] G. Perelman, “The entropy formula for the Ricci flow and its geometric applications”, preprint, 2000. [Zbl](#) [arXiv](#)
- [Schoen and Yau 1994] R. Schoen and S.-T. Yau, *Lectures on differential geometry*, Conference Proceedings and Lecture Notes in Geometry and Topology **1**, International, Cambridge, MA, 1994. [MR](#) [Zbl](#)
- [Sturm 2006a] K.-T. Sturm, “On the geometry of metric measure spaces, I”, *Acta Math.* **196**:1 (2006), 65–131. [MR](#) [Zbl](#)
- [Sturm 2006b] K.-T. Sturm, “On the geometry of metric measure spaces, II”, *Acta Math.* **196**:1 (2006), 133–177. [MR](#) [Zbl](#)
- [Wei and Wylie 2009] G. Wei and W. Wylie, “Comparison geometry for the Bakry–Émery Ricci tensor”, *J. Differential Geom.* **83**:2 (2009), 377–405. [MR](#) [Zbl](#)
- [Yang 2009] N. Yang, “A note on nonnegative Bakry–Émery Ricci curvature”, *Arch. Math. (Basel)* **93**:5 (2009), 491–496. [MR](#) [Zbl](#)

Received January 3, 2017. Revised December 5, 2017.

LI MA  
SCHOOL OF MATHEMATICS AND PHYSICS  
UNIVERSITY OF SCIENCE AND TECHNOLOGY BEIJING  
BEIJING  
CHINA

and

DEPARTMENT OF MATHEMATICS  
HENAN NORMAL UNIVERSITY  
XINXIANG  
CHINA

[lma@tsinghua.edu.cn](mailto:lma@tsinghua.edu.cn)

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Wee Teck Gan  
Mathematics Department  
National University of Singapore  
Singapore 119076  
[matgwt@nus.edu.sg](mailto:matgwt@nus.edu.sg)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Matthias Aschenbrenner  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[matthias@math.ucla.edu](mailto:matthias@math.ucla.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 296    No. 2    October 2018

---

Congruence subgroups and super-modular categories	257
PARSA BONDERSON, ERIC C. ROWELL, ZHENGHAN WANG and QING ZHANG	
On the Chow ring of the stack of truncated Barsotti–Tate Groups	271
DENNIS BROKEMPER	
Hyperbolic manifolds containing high topological index surfaces	305
MARION CAMPISI and MATT RATHBUN	
Length spectra of sub-Riemannian metrics on compact Lie groups	321
ANDRÁS DOMOKOS, MATTHEW KRAUEL, VINCENT PIGNO, COREY SHANBROM and MICHAEL VANVALKENBURGH	
The action of the Hecke operators on the component groups of modular Jacobian varieties	341
TAEKYUNG KIM and HWAJONG YOO	
Liouville theorems, volume growth, and volume comparison for Ricci shrinkers	357
LI MA	
Presentations of generalisations of Thompson’s group $V$	371
CONCHITA MARTÍNEZ-PÉREZ, FRANCESCO MATUCCI and BRITA NUCINKIS	
Localization functors and cosupport in derived categories of commutative Noetherian rings	405
TSUTOMU NAKAMURA and YUJI YOSHINO	
Zeta integrals for $\mathrm{GSp}(4)$ via Bessel models	437
RALF SCHMIDT and LONG TRAN	
Toric surfaces over an arbitrary field	481
FEI XIE	
Erratum for the article The Johnson–Morita theory for the rings of Fricke characters of free groups	509
TAKAO SATOH	