

*Pacific
Journal of
Mathematics*

**A RESTRICTION ON THE ALEXANDER POLYNOMIALS
OF L -SPACE KNOTS**

DAVID KRCA TOVICH

A RESTRICTION ON THE ALEXANDER POLYNOMIALS OF L -SPACE KNOTS

DAVID KRCA TOVICH

Using an invariant defined by Rasmussen, we extend an argument given by Hedden and Watson which further restricts which Alexander polynomials can be realized by L -space knots.

1. Introduction

In [Ozsváth and Szabó 2008], it is shown how the filtered chain homotopy type of the knot Floer complex $\text{CFK}^-(K)$ can be used to compute the Heegaard Floer homology of $S_n^3(K)$, the rational homology sphere obtained by performing Dehn surgery along $K \subset S^3$ with slope n . In [Ozsváth and Szabó 2005], this relationship is used to investigate which knots admit lens space surgeries, using the fact that if Y is a lens space, Y has the “smallest possible” Heegaard Floer homology:

$$(1) \quad \text{rank } \widehat{\text{HF}}(Y) = |H_1(Y; \mathbb{Z})|.$$

More generally, a rational homology sphere which satisfies condition (1) is called an L -space. So, from a Heegaard–Floer perspective, a natural extension of the question “which knots admit lens space surgeries?” is “which knots admit L -space surgeries?”.

Letting $A(x)$ denote the Alexander grading of a homogeneous element x in CFK^- , the following proposition is a straightforward consequence of [Ozsváth and Szabó 2005, Theorem 1.2]; see [Hom 2014, Remark 6.6].

Proposition 1.1. *Suppose $K \subset S^3$ is a knot on which some positive integral surgery yields an L -space. Then $\text{CFK}^-(K)$ has a basis $\{x_{-k}, \dots, x_k\}$ with the following properties:*

- $A(x_i) = n_i$, where $n_{-k} < n_{-k+1} < \dots < n_{k-1} < n_k$.
- $n_i = -n_{-i}$.
- If $i \equiv k \pmod{2}$, then $\partial(x_i) = 0$.
- If $i \equiv k + 1 \pmod{2}$, then $\partial(x_i) = x_{i-1} + U^{n_{i+1}-n_i} x_{i+1}$. □

MSC2010: primary 57M25; secondary 57M27.

Keywords: L -space knot, Alexander polynomial, lens space surgery.

Notice that x_k is in the kernel of ∂ and not in the image. The complex $\widehat{\text{CFK}}(K)$ is the quotient complex of $\text{CFK}^-(K)$ obtained by setting $U = 0$ (we refer the reader to [Ozsváth and Szabó 2004b] for details on knot Floer homology). It is a Heegaard Floer complex $\widehat{\text{CF}}(S^3)$, with an additional filtration induced by K . After setting $U = 0$, we still observe that

$$(2) \quad x_k \text{ generates } H_*(\widehat{\text{CFK}}(K), \partial) \cong \widehat{\text{HF}}(S^3) \cong \mathbb{F}.$$

By convention then, $M(x_k) = 0$ (where M is the Maslov grading). Since U decreases M by 2, and ∂ decreases M by 1, this determines the Maslov grading on all homogeneous elements of $\text{CFK}^-(K)$.

Ozsváth and Szabó [2004b] also showed that the graded Euler characteristic of $\widehat{\text{CFK}}(K)$ is the symmetrized Alexander polynomial of K ,

$$(3) \quad \sum_i \chi(\widehat{\text{CFK}}(K, i)) \cdot T^i = \Delta_K(T),$$

so a corollary to Proposition 1.1 is the following:

Corollary 1.2 [Ozsváth and Szabó 2005, Corollary 1.3]. *If $K \subset S^3$ is a knot which admits an L -space surgery, then*

$$(4) \quad \Delta_K(T) = \sum_{i=-k}^k (-1)^{k+i} T^{n_i}$$

for some sequence of integers $n_{-k} < n_{-k+1} < \dots < n_{k-1} < n_k$ satisfying $n_{-i} = n_i$.

Remark 1.3. Although Proposition 1.1 only applies to knots which have positive L -space surgeries, a knot K has a negative L -space surgery if and only if its mirror image \bar{K} has a positive L -space surgery. Corollary 1.2 then follows in this generality because $\Delta_K(T) = \Delta_{\bar{K}}(T)$.

In particular, all of the nonzero coefficients of $\Delta_K(T)$ are ± 1 . Note that

$$(5) \quad n_k = g(K) = |\tau(K)| = g_4(K),$$

where g is the Seifert genus, τ is the Ozsváth–Szabó concordance invariant defined in [Ozsváth and Szabó 2003], and g_4 is the smooth four-genus of K . The first equality follows from the knot Floer homology detection of genus [Ozsváth and Szabó 2004a], the second follows from (2), and the third follows from the fact shown in [Ozsváth and Szabó 2003] that, for any knot K ,

$$|\tau(K)| \leq g_4(K) \leq g(K).$$

This was the most general restriction on the Alexander polynomials of knots admitting L -space surgeries in the literature until Hedden and Watson showed the following proposition.

Proposition 1.4 [Hedden and Watson 2018, Corollary 9]. *If $K \subset S^3$ is a knot which admits an L -space surgery, then $\Delta_K(T)$ is as described in Corollary 1.2, and further, $n_k - n_{k-1} = 1$.*

Originally stated by Hedden and Watson as a corollary to a more general restriction on knot Floer complexes, this particular result was already known by Rasmussen, as conveyed to the author in a private communication. The proof hinges on an invariant defined by Rasmussen, and a particular inequality which it satisfies. Roughly, if large n -surgery is done on an unknot and on a knot K , the differences in the d -invariants (defined in (8)) of the resulting manifolds are bounded above by numbers depending on $g_4(K)$. Proposition 1.4 is then proved by showing that if a complex has a basis as in Proposition 1.1 and $n_k - n_{k-1} > 1$, Rasmussen’s inequality is violated, and therefore this complex cannot be the knot Floer complex of any knot.

Our aim here is to extend this argument. We will introduce Rasmussen’s invariant and inequality in Section 2. In Section 3, we will show how to compute the invariant for L -space knots from their Alexander polynomials (that is, from the sequence of the n_i). We will then see that Rasmussen’s inequality places further restrictions on the n_i , analogous to the restriction $n_k - n_{k-1} = 1$. As a result, it will be shown that certain symmetric Laurent polynomials satisfying Proposition 1.4 cannot be the Alexander polynomial of any L -space knot.

Theorem 1.5. *Suppose $K \subset S^3$ is a knot which admits an L -space surgery. Then its symmetrized Alexander polynomial can be written as*

$$\Delta_K(T) = \sum_{i=-k}^k (-1)^{k+i} T^{n_i},$$

for some sequence of integers $n_{-k} < n_{-k+1} < \dots < n_{k-1} < n_k$ satisfying the following:

- $n_i = -n_{-i}$,
- if we let $r_i = n_{k+2-2i} - n_{k+1-2i}$, then $r_1 = 1$, and for any $j \leq k$,

$$(6) \quad \sum_{i=2}^j r_i \leq \sum_{i=k-j+2}^k r_i.$$

As we will explain in Section 3, the restriction is more easily stated in terms of a modified version of the Alexander polynomial,

$$\tilde{\Delta}_K(T) := \frac{\Delta_K(T)}{1 - T^{-1}}.$$

It follows from [Corollary 1.2](#) that when K is a knot which admits an L -space surgery,

$$\tilde{\Delta}_K(T) = \sum_{i=0}^{\infty} T^{a_i}$$

for some sequence of integers satisfying

- $a_0 = g(K)$,
- $a_{i+1} < a_i$,
- $a_i = -i$ for $i \geq g(K)$.

We can then rephrase [Theorem 1.5](#) as follows.

Theorem 1.6 (restatement of [Theorem 1.5](#) in terms of $\tilde{\Delta}$). *Suppose $K \subset S^3$ is a knot which admits an L -space surgery and $\{a_i\}$ is the decreasing sequence of integers such that*

$$\tilde{\Delta}_K(T) = \sum_{i=0}^{\infty} T^{a_i}.$$

Then, for all $0 \leq i \leq g(K)$,

$$(7) \quad a_i \leq g(K) - 2i.$$

To see the preceding theorems as generalizations of [Proposition 1.4](#), note that in the language of [Theorem 1.5](#), [Proposition 1.4](#) translates to the statement $r_1 = 1$; in the language of [Theorem 1.6](#), it translates to $a_1 \leq g(K) - 2$.

As a concrete example, there does not exist a knot in S^3 which admits an L -space surgery and has Alexander polynomial

$$-1 + (T^2 + T^{-2}) - (T^3 + T^{-3}) + (T^4 + T^{-4}).$$

Correspondingly, there does not exist a knot in S^3 with knot Floer complex as shown in [Figure 1](#). As demonstrated after the proof of [Theorem 1.5](#), this settles the question of which polynomials belong to knots of genus 4 or less which admit L -space surgeries.

2. The invariant $h_m(K)$

A useful feature of Heegaard Floer theory is that its groups satisfy surgery exact triangles; for example, a long exact sequence between Heegaard Floer homology groups of manifolds which are 0-, ∞ - and n -framed surgery along the same knot K [[Ozsváth and Szabó 2004c](#), Section 9]. Rasmussen [[2003](#), Definition 7.1] defines an invariant $h_m(K)$ as the rank of a particular map in such a sequence (cf. [[Frøyshov 2002](#)], where an instanton-Floer invariant h is introduced).




Figure 1. A bifertered chain complex which cannot be the knot Floer complex of any knot in S^3 .

Recall that if (Y, \mathfrak{t}) is a spin^c rational homology sphere, Ozsváth and Szabó define the d -invariant of (Y, \mathfrak{t}) as

$$(8) \quad d(Y, \mathfrak{t}) = \min\{M(x) \mid x \in \text{Im}(\pi_* : \text{HF}^\infty(Y, \mathfrak{t}) \rightarrow \text{HF}^+(Y, \mathfrak{t}))\}.$$

Rasmussen [2004, Section 2.2] shows that, in the case where $S^3_{-n}(K)$ is an L -space, the invariant $d(S^3_{-n}(K), \mathfrak{s}_m)$ is equal to twice $h_m(K)$, up to a shift which is independent of K . In particular, since $h_m(\text{unknot}) = 0$ for all m , we have

$$(9) \quad h_m(K) = \frac{1}{2}(d(S^3_{-n}(K), \mathfrak{s}_m) - d(S^3_{-n}(\text{unknot}), \mathfrak{s}_m)).$$

The key to obtaining restrictions on L -space knots is the following inequality, analogous to an inequality in instanton Floer homology proved by Frøyshov [2004].

Proposition 2.1 [Rasmussen 2004, Theorem 2.3]. *Let K be a knot in S^3 and let $g_4(K)$ be its slice genus. Then $h_m(K) = 0$ for $|m| > g_4(K)$, while for $|m| \leq g_4(K)$,*

$$(10) \quad h_m(K) \leq \lceil \frac{1}{2}(g_4(K) - |m|) \rceil.$$

Note that for a knot admitting an L -space surgery, due to (5), we can replace $g_4(K)$ with $g(K)$ and obtain

$$(11) \quad h_m(K) \leq \lceil \frac{1}{2}(g(K) - |m|) \rceil.$$

While the term L -space knot refers to a knot which admits an L -space surgery, different conventions are used regarding the restrictions on the slope of the L -space surgeries. Here it will be convenient to adopt the definition that an L -space knot in

S^3 is one with a *positive* L -space surgery. This is opposite Rasmussen’s point of view [2004], but note that K admits a positive L -space surgery if and only if its mirror image \bar{K} admits a negative L -space surgery. Accordingly, we follow Hedden and Watson in defining

$$(12) \quad \bar{h}_m(K) := \frac{1}{2}(d(S_n^3(\text{unknot}), \mathfrak{s}_m) - d(S_n^3(K), \mathfrak{s}_m))$$

and recall their observation that $\bar{h}_m(K) = h_m(\bar{K})$. Finally, we should note that $g(K) = g(\bar{K})$, so \bar{h}_m satisfies the same inequality which h_m does for knots admitting L -space surgeries: for $|m| \leq g(K)$,

$$(13) \quad \bar{h}_m(K) \leq \lceil \frac{1}{2}(g(K) - |m|) \rceil.$$

3. Values of \bar{h}_m for L -space knots

Next, we recall how to compute d -invariants, and therefore \bar{h}_m , from CFK^- . It was shown independently by Ozsváth and Szabó [2004b] and Rasmussen [2003] that for large n -surgery (that is, for $n \geq 2g(K) - 1$), the Heegaard Floer homology groups $\text{HF}^-(S_n^3(K))$ are the homology groups of certain subcomplexes of $\text{CFK}^-(K)$, up to a shift in Maslov grading which is independent of K . In particular, if we let A_m denote the subcomplex consisting of elements with Alexander grading less than or equal to m , then

$$\text{HF}^-(S_n^3(K), \mathfrak{s}_m) \cong H_*(A_m),$$

up to a shift in grading.¹ It follows that

$$d(S_n^3(K), \mathfrak{s}_m) = \max\{M(x) \mid x \text{ a nontorsion generator of } H_*(A_m)\} + c,$$

where c is a constant which depends on n , but not on K . Therefore, the “shifted” d -invariant

$$(14) \quad \tilde{d}(K, m) := \max\{M(x) \mid x \text{ a nontorsion generator of } H_*(A_m)\}$$

is well-defined, and satisfies

$$(15) \quad d(S_n^3(\text{unknot}), \mathfrak{s}_m) - d(S_n^3(K), \mathfrak{s}_m) = \tilde{d}(\text{unknot}, m) - \tilde{d}(K, m)$$

for any sufficiently large n . For the unknot, we have the complex

$$\text{CFK}^-(\text{unknot}) \cong \mathbb{F}[U],$$

where the generator has Maslov grading and Alexander grading equal to zero. Since multiplication by U lowers the Alexander grading by 1 and the Maslov grading by 2,

$$\tilde{d}(\text{unknot}, m) = m - |m|.$$

¹Here we are adopting the convention that CF^- and CFK^- contain the element 1 in $\mathbb{F}[U]$.

Therefore, we can rewrite inequality (13) using (15) and the above: if $K \subset S^3$ is an L -space knot, then for $|m| \leq g(K)$,

$$(16) \quad \begin{aligned} \bar{h}_m(K) &= \frac{1}{2}(\tilde{d}(\text{unknot}, m) - \tilde{d}(K, m)) \leq \left\lceil \frac{1}{2}(g(K) - |m|) \right\rceil, \\ -\frac{1}{2}\tilde{d}(K, m) &\leq \left\lceil \frac{1}{2}(g(K) - m) \right\rceil. \end{aligned}$$

With inequality (16) in hand, it remains to see how the values of \tilde{d} are determined by the Alexander polynomial of an L -space knot.

Recall that the Alexander polynomial is the graded Euler characteristic of $\widehat{\text{CFK}}(K)$:

$$\sum_i \chi(\widehat{\text{CFK}}(K, i)) \cdot T^i = \Delta_K(T).$$

Further, CFK^- is generated by the same set as $\widehat{\text{CFK}}$, over $\mathbb{F}[U]$ rather than \mathbb{F} . Since U lowers the Alexander grading by 1 and preserves the parity of the Maslov grading,

$$(17) \quad \begin{aligned} \sum_i \chi(\text{CFK}^-(K, i)) \cdot T^i &= \sum_i \chi(\widehat{\text{CFK}}(K, i)) \cdot T^i \cdot (1 + T^{-1} + T^{-2} + \dots) \\ &= \frac{\Delta_K(T)}{1 - T^{-1}} =: \tilde{\Delta}_K(T). \end{aligned}$$

In other words, $\tilde{\Delta}_K(T)$ is the graded Euler characteristic of $\text{CFK}^-(K)$.

Remark 3.1. If $K \subset S^3$ is a knot for which $\Delta_K(T)$ is of the type described in Corollary 1.2, then

$$\tilde{\Delta}_K(T) = \sum_{i=0}^{\infty} T^{a_i},$$

where

- $a_0 = g(K)$,
- $a_{i+1} < a_i$, and
- $a_i = -i$ for all $i \geq g(K)$.

In [Krcatovich 2015], a reduced complex $\underline{\text{CFK}}^-$ was defined, and it was shown that for an L -space knot,

$$(18) \quad \underline{\text{CFK}}^-(K) \cong \mathbb{F}[U],$$

supported in Maslov grading zero [Krcatovich 2015, Corollary 4.2]. Roughly speaking, the complex CFK^- has a filtration induced by U , and a filtration induced the knot; ignoring the knot filtration, one recovers $\text{CF}^-(S^3)$, whereas ignoring the U -filtration, one gets a “reduced” knot Floer complex. We refer the reader to [Krcatovich 2015] for a precise statement, and here simply remark that the structure of CFK^- for an L -space knot, as described in Proposition 1.1, is what makes its reduced complex have a simple form.

Since the reduced complex is filtered chain homotopy equivalent to $\text{CFK}^-(K)$ (with respect to the knot filtration), they have the same Euler characteristic. In particular, (18) says that every generator has even Maslov grading, so each contributes a positive term to the Euler characteristic. In other words, if

$$\tilde{\Delta}_K(T) = \sum_{i=0}^{\infty} T^{a_i},$$

then $\text{CFK}^-(K)$ has one generator with Alexander grading a_i , for each $i \geq 0$. Since multiplication by U is a filtered map (i.e., it never increases the Alexander grading), then necessarily

$$M(a_i) = -2i.$$

Figure 2 gives an illustration for the case of the (3, 4)-torus knot,² where

$$(19) \quad \Delta_{T_{3,4}}(T) = 1 - (T^2 + T^{-2}) + (T^3 + T^{-3}),$$

and therefore

$$(20) \quad \tilde{\Delta}_{T_{3,4}}(T) = T^3 + 1 + T^{-1} + T^{-3} + T^{-4} + \dots$$

Proof of Theorem 1.6. First note that it is sufficient to prove the proposition for positive surgeries (see Remark 1.3).

So, let K be an L -space knot in S^3 , so that

$$\tilde{\Delta}_K(T) = \sum_{i=0}^{\infty} T^{a_i}.$$

Then the reduced complex $\text{CFK}^-(K)$ consists of a single generator of Alexander grading a_i and Maslov grading $-2i$, for each $i \geq 0$. Since the a_i are strictly decreasing, it follows that \tilde{d} , as defined in (14), is given by

$$\tilde{d}(K, m) = -2 \min\{i \mid a_i \leq m\},$$

and therefore

$$(21) \quad \tilde{d}(K, a_i - 1) = -2(i + 1).$$

Substituting these values into inequality (16), we obtain

$$(22) \quad i + 1 \leq \left\lceil \frac{1}{2}(g(K) - (a_i - 1)) \right\rceil,$$

so

$$a_i \leq g(K) - 2i. \quad \square$$

²It was shown by Moser [1971] that torus knots admit lens space (hence L -space) surgeries.




Figure 2. To the left is the knot Floer complex CFK^- for the $(3,4)$ -torus knot, which, by Proposition 1.1, is determined by its Alexander polynomial. To the right is the reduced complex, which, for any L -space knot, is isomorphic to $\mathbb{F}[U]$, supported in Maslov grading zero. Note that the reduced complex on the right has a generator for each \circ generator on the left, the bottom-most generator of each staircase summand.

Proof of Theorem 1.5. Let K be an L -space knot, so that

$$\Delta_K(T) = \sum_{i=-k}^k (-1)^{k+i} T^{n_i}.$$

We have introduced the variables

$$\vec{r} = (r_1, \dots, r_k)$$

as the “gaps” in the Alexander polynomial (the difference in the exponents of consecutive nonzero terms),

$$r_i = n_{k+2-2i} - n_{k+1-2i}.$$

While \vec{r} records only every second gap, by the symmetry of $\Delta(T)$, this determines the polynomial uniquely. Diagrammatically, notice that \vec{r} is simply the list of horizontal lengths of a staircase summand of CFK^- , in order from left to right. See Figure 3 for the example of the $(4, 5)$ -torus knot, which has

$$\Delta_{T(4,5)}(T) = -1 + (T^2 + T^{-2}) - (T^5 + T^{-5}) + (T^6 + T^{-6}).$$




Figure 3. The complex CFK^- for the $(4, 5)$ -torus knot, and its reduced form \underline{CFK}^- . Note that the integers r_1, r_2 and r_3 are the horizontal lengths of each staircase, from left to right (and by symmetry, the vertical lengths, from bottom to top). This figure illustrates how the m_j — the Alexander gradings at which the reduced complex “jumps” — are determined by the r_i , and further, how the values of $\tilde{d}(K, m_j)$, given in parentheses to the right, are determined by the r_i .

Next we observe how, given \vec{r} , to compute both sides of inequality (16) for any m , with Figure 3 as a guide. We focus on the values labeled m_j in Figure 3; in other words, the values where we have the “jumps” in the reduced complex. More precisely, if we let

$$m_j = g(K) - \left(\sum_{i=1}^j r_i + \sum_{i=k-j+2}^k r_i \right),$$

then we have that

$$\tilde{d}(K, m_j) = -2 \sum_{i=1}^j r_i.$$

Substituting these values into inequality (16) when $m = m_j$ gives

$$(23) \quad \sum_{i=1}^j r_i \leq \left\lceil \frac{g(K) - (g(K) - (\sum_{i=1}^j r_i + \sum_{i=k-j+2}^k r_i))}{2} \right\rceil.$$

The case $j = 1$ gives $r_1 \leq \lceil \frac{1}{2}r_1 \rceil$, so, since each r_i is a positive integer, r_1 must equal 1. Substituting this into (23) gives

$$1 + \sum_{i=2}^j r_i \leq \left\lceil \frac{1 + \sum_{i=2}^j r_i + \sum_{i=k-j+2}^k r_i}{2} \right\rceil,$$

$$\sum_{i=2}^j r_i \leq \left\lceil \frac{-1 + \sum_{i=2}^j r_i + \sum_{i=k-j+2}^k r_i}{2} \right\rceil,$$

from which it follows that

$$\sum_{i=2}^j r_i \leq \sum_{i=k-j+2}^k r_i.$$

This is sufficient to prove the claim. We could similarly obtain inequalities by considering values of m different from the m_j , but those would be no stronger, and therefore provide no more restrictions on \vec{r} . \square

As an example, consider a knot K with

$$\Delta_K(T) = -1 + (T^2 + T^{-2}) - (T^3 + T^{-3}) + (T^4 + T^{-4}),$$

so that

$$\tilde{\Delta}_K(T) = T^4 + T^2 + T + T^{-2} + T^{-4} + T^{-5} + \dots$$

This polynomial satisfies the restriction of Proposition 1.4, but if K were an L -space knot, we would have $g(K) = 4$, and $a_2 = 1$. This violates inequality (7), so K (and its mirror image) cannot admit an L -space surgery. Alternatively, this polynomial has gaps $\vec{r} = (1, 2, 1)$, and since $r_2 \not\leq r_3$, this violates inequality (6).

In fact, this completely determines which Alexander polynomials are realized by L -space knots of genus less than or equal to 4. All other polynomials satisfying Proposition 1.4 are realized by known L -space knots. For knots of genus 5, Theorem 1.5 eliminates the polynomials corresponding to $\vec{r} = (1, 2, 1, 1)$ and $\vec{r} = (1, 3, 1)$, but there are still three more which are not realized by any L -space knot known to the author (corresponding to $(1, 1, 2, 1)$, $(1, 2, 2)$ and $(1, 4)$).

\vec{r}	L -space knot with corresponding $\Delta(T)$
(1)	$T(2, 3)$
(1,1)	$T(2, 5)$
(1,1,1)	$T(2, 7)$
(1,2)	$T(3, 4)$, (2, 3)-cable of $T(2, 3)$
(1,1,1,1)	$T(2, 9)$
(1,1,2)	$T(3, 5)$
(1,2,1)	excluded by Theorem 1.5
(1,3)	(2, 5)-cable of $T(2, 3)$

Finally, we should point out the relation between the sequence of the a_i for an L -space knot and the gap function defined by Borodzik and Livingston [2014, Definition 2.6]; namely,

$$a_i = \min_m \{I_K(m) = i\} - g(K).$$

As pointed out to the author by Borodzik, their restrictions given in [Borodzik and Livingston 2016, Theorem 2.14] can be reinterpreted in terms of the a_i . Informally, if two L -space knots are related by a small number of crossing changes, they have similar Alexander polynomials. More precisely, if a_i and a'_i are the exponents of $\tilde{\Delta}$ for two L -space knots which differ by changing p positive crossings to negative crossings, then

$$|a_i - a'_i| \leq p \quad \text{for all } i.$$

Acknowledgments

The author would like to thank his advisor, Matt Hedden, for explaining the proof of [Proposition 1.4](#), and for suggesting the restatement of [Theorem 1.5](#) in terms of the polynomial $\tilde{\Delta}$, and Maciej Borodzik for pointing out the relation to the gap function defined in [Borodzik and Livingston 2014]. The author also thanks the anonymous referee for suggesting improvements to an earlier draft. The author was partially supported by NSF grant DMS-1150872.

References

- [Borodzik and Livingston 2014] M. Borodzik and C. Livingston, “Heegaard Floer homology and rational cuspidal curves”, *Forum Math. Sigma* **2** (2014), art. id. e28. [MR](#) [Zbl](#)
- [Borodzik and Livingston 2016] M. Borodzik and C. Livingston, “Semigroups, d -invariants and deformations of cuspidal singular points of plane curves”, *J. Lond. Math. Soc.* (2) **93**:2 (2016), 439–463. [MR](#) [Zbl](#)
- [Frøyshov 2002] K. A. Frøyshov, “Equivariant aspects of Yang–Mills Floer theory”, *Topology* **41**:3 (2002), 525–552. [MR](#) [Zbl](#)

- [Frøyshov 2004] K. A. Frøyshov, “An inequality for the h -invariant in instanton Floer theory”, *Topology* **43**:2 (2004), 407–432. [MR](#) [Zbl](#)
- [Hedden and Watson 2018] M. Hedden and L. Watson, “On the geography and botany of knot Floer homology”, *Selecta Math. (N.S.)* **24**:2 (2018), 997–1037. [MR](#) [Zbl](#)
- [Hom 2014] J. Hom, “The knot Floer complex and the smooth concordance group”, *Comment. Math. Helv.* **89**:3 (2014), 537–570. [MR](#) [Zbl](#)
- [Krcatovich 2015] D. Krcatovich, “The reduced knot Floer complex”, *Topology Appl.* **194** (2015), 171–201. [MR](#) [Zbl](#)
- [Moser 1971] L. Moser, “Elementary surgery along a torus knot”, *Pacific J. Math.* **38** (1971), 737–745. [MR](#) [Zbl](#)
- [Ozsváth and Szabó 2003] P. Ozsváth and Z. Szabó, “Knot Floer homology and the four-ball genus”, *Geom. Topol.* **7** (2003), 615–639. [MR](#) [Zbl](#)
- [Ozsváth and Szabó 2004a] P. Ozsváth and Z. Szabó, “Holomorphic disks and genus bounds”, *Geom. Topol.* **8** (2004), 311–334. [MR](#) [Zbl](#)
- [Ozsváth and Szabó 2004b] P. Ozsváth and Z. Szabó, “Holomorphic disks and knot invariants”, *Adv. Math.* **186**:1 (2004), 58–116. [MR](#) [Zbl](#)
- [Ozsváth and Szabó 2004c] P. Ozsváth and Z. Szabó, “Holomorphic disks and three-manifold invariants: properties and applications”, *Ann. of Math. (2)* **159**:3 (2004), 1159–1245. [MR](#) [Zbl](#)
- [Ozsváth and Szabó 2005] P. Ozsváth and Z. Szabó, “On knot Floer homology and lens space surgeries”, *Topology* **44**:6 (2005), 1281–1300. [MR](#) [Zbl](#)
- [Ozsváth and Szabó 2008] P. S. Ozsváth and Z. Szabó, “Knot Floer homology and integer surgeries”, *Algebr. Geom. Topol.* **8**:1 (2008), 101–153. [MR](#) [Zbl](#)
- [Rasmussen 2003] J. A. Rasmussen, *Floer homology and knot complements*, Ph.D. thesis, Harvard University, 2003, available at <https://search.proquest.com/docview/305332635>.
- [Rasmussen 2004] J. Rasmussen, “Lens space surgeries and a conjecture of Goda and Teragaito”, *Geom. Topol.* **8** (2004), 1013–1031. [MR](#) [Zbl](#)

Received August 30, 2016. Revised October 4, 2017.

DAVID KRCA TOVICH
DEPARTMENT OF MATHEMATICS
RICE UNIVERSITY
HOUSTON, TX
UNITED STATES
david.krcatovich@gmail.com

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 297 No. 1 November 2018

On Legendre curves in normed planes	1
VITOR BALESTRO, HORST MARTINI and RALPH TEIXEIRA	
Remarks on critical metrics of the scalar curvature and volume functionals on compact manifolds with boundary	29
HALYSON BALTAZAR and ERNANI RIBEIRO, JR.	
Cherlin's conjecture for sporadic simple groups	47
FRANCESCA DALLA VOLTA, NICK GILL and PABLO SPIGA	
A characterization of round spheres in space forms	67
FRANCISCO FONTENELE and ROBERTO ALONSO NÚÑEZ	
A non-strictly pseudoconvex domain for which the squeezing function tends to 1 towards the boundary	79
JOHN ERIK FORNÆSS and ERLEND FORNÆSS WOLD	
An Amir–Cambern theorem for quasi-isometries of $C_0(K, X)$ spaces	87
ELÓI MEDINA GALEGO and ANDRÉ LUIS PORTO DA SILVA	
Weak amenability of Lie groups made discrete	101
SØREN KNUDBY	
A restriction on the Alexander polynomials of L -space knots	117
DAVID KRATOVICH	
Stability of capillary hypersurfaces in a Euclidean ball	131
HAIZHONG LI and CHANGWEI XIONG	
Non-minimality of certain irregular coherent preminimal affinizations	147
ADRIANO MOURA and FERNANDA PEREIRA	
Interior gradient estimates for weak solutions of quasilinear p -Laplacian type equations	195
TUOC PHAN	
Local unitary periods and relative discrete series	225
JERROD MANFORD SMITH	