

*Pacific  
Journal of  
Mathematics*

**YAMABE FLOW WITH PRESCRIBED SCALAR CURVATURE**

INAS AMACHA AND RACHID REGBAOUI

Volume 297 No. 2

December 2018



# YAMABE FLOW WITH PRESCRIBED SCALAR CURVATURE

INAS AMACHA AND RACHID REGBAOUI

**We study the Yamabe flow corresponding to the prescribed scalar curvature problem on compact Riemannian manifolds with negative scalar curvature. The long time existence and convergence of the flow are proved under appropriate conditions on the prescribed scalar curvature function.**

## 1. Introduction

The prescribed scalar curvature problem on a compact Riemannian manifold  $(M, g_0)$  of dimension  $n \geq 3$ , consists of finding a conformal metric  $g$  to  $g_0$  whose scalar curvature  $R_g$  is equal to a given function  $f \in C^\infty(M)$ . If we set  $g = u^{4/(n-2)}g_0$ , where  $0 < u \in C^\infty(M)$ , then we have

$$R_g = u^{-\frac{n+2}{n-2}}(-c_n \Delta u + R_0 u),$$

where  $\Delta$  is the Laplace operator associated with  $g_0$ ,  $R_0$  is the scalar curvature of  $g_0$  and  $c_n = 4\frac{n-1}{n-2}$ .

Then the prescribed scalar curvature problem,

$$R_g = f,$$

is equivalent to solving the nonlinear PDE

$$(1-1) \quad -c_n \Delta u + R_0 u = f u^{\frac{n+2}{n-2}}$$

on the space of smooth positive functions on  $M$ . The solvability of this equation depends on  $R_0$  and the prescribed function  $f$ . When  $f$  is constant, (1-1) becomes the famous Yamabe equation whose resolution has been a challenging problem in geometric analysis for a long time. See [Aubin 1976; Hebey and Vaugon 1993; Lee and Parker 1987; Schoen 1984; 1991] for more details on the Yamabe problem, and [Ambrosetti and Malchiodi 1999; Bismuth 2000; Escobar and Schoen 1986; Kazdan and Warner 1975; Rauzy 1995; Vázquez and Véron 1991], concerning the prescribed scalar curvature problem.

By changing  $g_0$  conformally if necessary, we may always assume that  $R_0$  satisfies one of the conditions,  $R_0 > 0$ ,  $R_0 = 0$  or  $R_0 < 0$  everywhere on  $M$ . Equation (1-1)

*MSC2010:* 35K55, 53A30, 58J35.

*Keywords:* conformal metric, scalar curvature, Yamabe flow.

has a variational structure since there are different functionals whose Euler–Lagrange equations are equivalent to (1-1). When  $R_0 < 0$ , the following functional seems more appropriate to handle the prescribed scalar curvature problem:

$$(1-2) \quad \mathcal{E}(g) = \int_M R_g dV_g - \frac{n-2}{n} \int_M f dV_g,$$

where  $g = u^{4/(n-2)}g_0$  belongs to the conformal class  $[g_0]$  of  $g_0$ ,  $R_g$  is the scalar curvature of  $g$  and  $dV_g = u^{2n/(n-2)}dV_{g_0}$  is the volume element of  $g$ .

Simple computations ([Besse 1987]) show the  $L^2$ -gradient of  $\mathcal{E}$  is  $\frac{n-2}{2n}(R_g - f)g$ , and then, after changing time by a constant scale, the associated negative gradient flow equation is

$$(1-3) \quad \begin{cases} \partial_t g = -(R_g - f)g, \\ g(0) = g^0, \end{cases}$$

where  $g^0 = u_0^{4/(n-2)}g_0$  is a given metric in the conformal class of  $g_0$ .

Since (1-3) preserves the conformal structure of  $M$ , then any smooth solution of (1-3) is of the form  $g(t) = u(t)^{4/(n-2)}g_0$ , where  $0 < u(t) \in C^\infty(M)$ . For simplicity we have used the notation  $u(t) := u(t, \cdot)$ ,  $t \in I$  for any function  $u$  defined on  $I \times M$ , where  $I$  is a subset of  $\mathbb{R}$ . In terms of  $u(t)$ , the flow (1-3) may be written in the equivalent form:

$$(1-4) \quad \begin{cases} \partial_t u^N = \frac{n+2}{4}(c_n \Delta u - R_0 u + f u^N), \\ u(0) = u_0 \in C^\infty(M), \quad u_0 > 0, \end{cases}$$

where  $N = \frac{n+2}{n-2}$ .

Our aim in this paper is to investigate this gradient flow by proving its longtime existence and analysing its asymptotic behaviour when  $t \rightarrow +\infty$ .

Our first result is the following existence theorem:

**Theorem 1.1.** *Suppose that  $R_0 < 0$  and let  $f \in C^\infty(M)$ . Then for any  $g^0 = u_0^{4/(n-2)}g_0$  with  $0 < u_0 \in C^\infty(M)$ , there exists a unique solution  $g(t) = u(t)^{4/(n-2)}g_0$  of (1-3) defined on  $[0, +\infty)$ , where  $0 < u \in C^\infty([0, +\infty) \times M)$ . Moreover, the functional  $\mathcal{E}$  is decreasing along the solution  $g(t)$ , that is,*

$$\frac{d}{dt} \mathcal{E}(g(t)) \leq 0 \quad \text{for all } t \in [0, +\infty).$$

We note here that apart from the smoothness of  $f$ , no further assumptions on the function  $f$  are needed in Theorem 1.1. However, for the longtime behaviour, it is necessary to assume additional conditions in order to get the convergence of the flow. Indeed, if  $f \geq 0$ , by applying the maximum principle to (1-4), we can easily check that

$$u(t) \geq \left( \min_M u_0^{4/(n-2)} + \min_M |R_0|t \right)^{\frac{n-2}{4}} \rightarrow +\infty \quad \text{as } t \rightarrow +\infty.$$

So if we want to get the convergence of the flow, it is necessary to assume at least

that  $f$  is negative somewhere on  $M$ . We note that this last condition is also necessary for the resolution of (1-1) since it is well known that if the negative gradient flow associated with a functional  $\mathcal{F}$  converges (in some sense), then its limit is a critical point of  $\mathcal{F}$ .

Before giving conditions on  $f$  ensuring the convergence of the flow, let us fix some notation: if  $\Omega \subset M$  is an open set, we denote by  $\lambda_\Omega$  the first eigenvalue of the conformal Laplacian  $L = -c_n \Delta + R_0$  on  $\Omega$  with zero Dirichlet boundary conditions, that is,

$$\lambda_\Omega = \inf_{0 \neq u \in H_0^1(\Omega)} \frac{\int_M (c_n |\nabla u|^2 + R_0 u^2) dV_{g_0}}{\int_M u^2 dV_{g_0}}.$$

We then assume the following conditions on  $f$ :

There exists an open set  $\Omega \subset M$  such that

**(H1)**  $\lambda_\Omega > 0$  and  $f < 0$  on  $M \setminus \Omega$

and

**(H2)**  $\sup_{x \in \Omega} f(x) \leq C_\Omega \inf_{x \in M \setminus \Omega} |f(x)|,$

where  $C_\Omega$  is a positive constant depending only on  $\Omega$ .

We then have the following result:

**Theorem 1.2.** *Suppose that  $R_0 < 0$  and that  $f \in C^\infty(M)$  satisfies conditions **(H1)** and **(H2)**. Then there exists a function  $0 < \bar{u} \in C^\infty(M)$  such that for any smooth metric  $g^0 = u_0^{4/(n-2)} g_0$  with  $0 < u_0 \leq \bar{u}$ , the flow  $g(t) = u(t)^{4/(n-2)} g_0$  given by Theorem 1.1 converges in the  $C^\infty$ -topology to a conformal metric  $g_\infty = u_\infty^{4/(n-2)} g_0$  whose scalar curvature is  $f$ , that is,  $R_{g_\infty} = f$ .*

A particular interesting case is when the function  $f$  satisfies  $f(x) < 0$  for almost all  $x \in M$ . In this case conditions **(H1)** and **(H2)** are automatically satisfied and then we have the following corollary:

**Corollary 1.3.** *Suppose that  $R_0 < 0$  and  $f \in C^\infty(M)$  such that  $f < 0$  almost everywhere on  $M$ . Then there exists a function  $0 < \bar{u} \in C^\infty(M)$  such that for any smooth metric  $g^0 = u_0^{4/(n-2)} g_0$  with  $0 < u_0 \leq \bar{u}$ , the flow  $g(t) = u(t)^{4/(n-2)} g_0$  given by Theorem 1.1 converges in the  $C^\infty$ -topology to a conformal metric  $g_\infty = u_\infty^{4/(n-2)} g_0$  whose scalar curvature is  $f$ , that is,  $R_{g_\infty} = f$ .*

It is natural to ask if conditions **(H1)** and **(H2)** in Theorem 1.2 are necessary. The following theorem gives a partial answer to this question:

**Theorem 1.4.** *Suppose that  $R_0 < 0$  and let  $f \in C^\infty(M)$  such that condition **(H1)** is not satisfied, that is, for any open set  $\Omega \subset M$  such that  $f > 0$  on  $M \setminus \Omega$ , we*

suppose  $\lambda_\Omega \leq 0$ . Then for any  $0 < u_0 \in C^\infty(M)$ , the solution  $u(t)$  of (1-4) satisfies, for some constant  $C > 0$  depending only on  $u_0, g_0, f$ ,

$$\max_{x \in M} u(t, x) \geq Ct^{\frac{n-2}{n+2}} \rightarrow +\infty \text{ as } t \rightarrow +\infty.$$

We note here that condition **(H1)** is conformally invariant. Similar conditions to **(H1)** and **(H2)** were found by many authors to solve (1-1) by the direct method of elliptic PDEs; see [Bismuth 2000; Rauzy 1995; Vázquez and Véron 1991] for more details. To our knowledge, the only known results on Yamabe type flow on dimension  $n \geq 3$  concern the case where  $f$  is constant or  $M = \mathbb{S}^n$ . The Yamabe flow was first introduced by Hamilton [1988] and has been the subject of several studies; see [Brendle 2005; 2007; Chow 1992; Schwetlick and Struwe 2003; Ye 1994]. For the case when  $f$  is nonconstant, we mention the work of Struwe [2005] about the Nirenberg’s problem on the sphere  $\mathbb{S}^2$ , and the results of Chen and Xu [2012] concerning  $\mathbb{S}^n, n \geq 3$ . A general evolution problem related to the prescribed Gauss curvature on surfaces was studied by Baird, Fardoun and Regbaoui [Baird et al. 2004].

The paper is organized as follows. In Section 2 we prove the global existence of the flow by establishing local  $C^k$ -estimates on the solution  $u$  of (1-4). In Section 3, we study the asymptotic behaviour of the flow when  $t \rightarrow +\infty$ . In particular we prove uniform  $C^k$ -estimates on  $u$  which are necessary to get the convergence of the flow.

### 2. Global existence of the flow

In this section we shall establish some estimates on the solution  $u$  of (1-4) which will be an important tool in proving that the flow  $g(t)$  is globally defined on  $[0, +\infty)$ . In this section we suppose that  $R_0 < 0$  and  $f \in C^\infty(M)$ .

As already mentioned in the previous section, (1-3) is equivalent to (1-4), so it suffices to prove the existence of a solution  $u(t)$  of (1-4) defined on  $[0, +\infty)$  to obtain a metric  $g(t)$  solution of (1-3) defined on  $[0, +\infty)$ . Since (1-4) is a parabolic equation (on the set of smooth positive functions on  $[0, T) \times M$ , for any  $T > 0$ ), there exists a smooth solution  $u(t)$  of (1-4) defined on a maximal interval  $[0, T^*)$  satisfying  $u(t) > 0$  on  $[0, T^*)$ . Thus we have a solution  $g(t) = u^{4/(n-2)}g_0$  of (1-3) defined on a maximal interval  $[0, T^*)$ . For simplicity, we shall write  $u$  instead of  $u(t)$  and  $g$  instead of  $g(t)$ .

Now, we derive some properties of  $g$  which will be important later. One can check by using (1-4) that the scalar curvature  $R_g$  satisfies the equation

$$(2-1) \quad \partial_t R_g = (n - 1)\Delta_g(R_g - f) + R_g(R_g - f),$$

where  $\Delta_g$  is the Laplacian associated with  $g(t)$ .

A simple computation using (2-1) gives

$$(2-2) \quad \frac{d}{dt} \mathcal{E}(g) = -\frac{n-2}{2} \int_M (R_g - f)^2 dV_g,$$

so the functional  $\mathcal{E}$  is decreasing along the flow  $g(t)$ . If we set

$$E(u) := \mathcal{E}(g) = \mathcal{E}(u^{\frac{4}{n-2}} g_0) = \int_M \left( c_n |\nabla u|^2 + R_0 u^2 - \frac{n-2}{n} f u^{\frac{2n}{n-2}} \right) dV_{g_0},$$

then (2-2) can be written in terms of  $u$ :

$$(2-3) \quad \frac{d}{dt} E(u) = -\frac{8}{n-2} \int_M |\partial_t u|^2 u^{\frac{4}{n-2}} dV_{g_0} \leq 0.$$

The following lemma will be very useful to prove integral estimates on the solution  $g$ .

**Lemma 2.1.** *We have for any  $p > 1$ ,*

$$\begin{aligned} \frac{d}{dt} \int_M |R_g - f|^p dV_g &= -\frac{4(n-1)(p-1)}{p} \int_M |\nabla_g |R_g - f|^{\frac{p}{2}}|_g^2 dV_g \\ &\quad + \left(p - \frac{n}{2}\right) \int_M (R_g - f) |R_g - f|^p dV_g + p \int_M f |R_g - f|^p dV_g, \end{aligned}$$

where  $\nabla_g$  is the gradient with respect to the metric  $g$  and  $|\cdot|_g$  is the Riemannian norm with respect to  $g$ .

*Proof.* We have for any  $p \geq 1$

$$\frac{d}{dt} \int_M |R_g - f|^p dV_g = p \int_M |R_g - f|^{p-2} (R_g - f) \partial_t R_g dV_g + \frac{1}{2} \int_M R_g \operatorname{tr}_g (\partial_t g) dV_g.$$

Using (1-3) and (2-1), it follows that

$$\begin{aligned} \frac{d}{dt} \int_M |R_g - f|^p dV_g &= (n-1)p \int_M |R_g - f|^{p-2} (R_g - f) \Delta_g (R_g - f) \\ &\quad + p \int_M R_g |R_g - f|^p - \frac{n}{2} \int_M |R_g - f|^p (R_g - f) dV_g \\ &= -\frac{4(n-1)(p-1)}{p} \int_M |\nabla_g |R_g - f|^{\frac{p}{2}}|_g^2 dV_g \\ &\quad + \left(p - \frac{n}{2}\right) \int_M (R_g - f) |R_g - f|^p dV_g + p \int_M f |R_g - f|^p dV_g. \quad \square \end{aligned}$$

In order to prove that the solution  $g(t) = u(t)^{4/(n-2)} g_0$  is globally defined on  $[0, +\infty)$ , we need upper and lower bounds on  $u(t)$ .

**Proposition 2.2.** *Let  $g(t) = u(t)^{4/(n-2)} g_0$  be the solution of (1-3) defined on a maximal interval  $[0, T^*)$ . Then we have for any  $t \in [0, T^*)$ ,*

$$(2-4) \quad \min(C_0, \min_M u_0) \leq u(t) \leq \max(1, \max_M u_0) e^{C_1 t},$$

where

$$C_0 = \left(\min_M |R_0| / \max_M |f|\right)^{\frac{n-2}{4}}, \quad C_1 = \frac{n-2}{4} \left(\max_M |R_0| + \max_M |f|\right).$$

*Proof.* The proof uses an elementary maximum principal argument. Indeed, fix  $t \in [0, T)$  and let  $(t_0, x_0) \in [0, t] \times M$  such that  $u(t_0, x_0) = \min_{[0,t] \times M} u$ . If  $t_0 = 0$ ,

$$\min_{[0,t] \times M} u = \min_M u_0,$$

so the first inequality in (2-4) is proved in this case. Now suppose that  $t_0 > 0$ . We have then  $\partial_t u(t_0, x_0) \leq 0$  and  $\Delta u(t_0, x_0) \geq 0$ . Thus after substituting in (1-4) we obtain that

$$0 \geq -R_0(x_0)u(t_0, x_0) + f(x_0)u^N(t_0, x_0)$$

which implies

$$u(t_0, x_0) \geq \left(\min_M |R_0| / \max_M |f|\right)^{\frac{1}{N-1}},$$

where  $N = \frac{n+2}{n-2}$ . This proves the first inequality in (2-4). In order to prove the second inequality we set  $v = e^{-C_1 t} u$  instead of  $u$ , where  $C_1 = \frac{4}{n-2} (\max_M |R_0| + \max_M |f|)$ . As above, fix  $t \in [0, T)$  and let  $(t_0, x_0) \in [0, t] \times M$  such that  $v(t_0, x_0) = \max_{[0,t] \times M} v$ . If  $t_0 = 0$ , then  $\max_{[0,t] \times M} v = \max_M u_0$ , which implies

$$\max_{[0,t] \times M} u \leq \max_M u_0 e^{C_1 t},$$

so the second inequality in (2-4) is proved in this case. Now suppose that  $t_0 > 0$ . We have then  $\partial_t v(t_0, x_0) \geq 0$  and  $\Delta v(t_0, x_0) \leq 0$ , that is,  $\partial_t u(t_0, x_0) \geq C_1 u(t_0, x_0)$  and  $\Delta u(t_0, x_0) \leq 0$ . We obtain after substituting in (1-4) that

$$NC_1 u^N(t_0, x_0) \leq \frac{n+2}{4} (-R_0(x_0)u(t_0, x_0) + f(x_0)u^N(t_0, x_0))$$

which implies that

$$(2-5) \quad u(t_0, x_0) \leq 1$$

since  $NC_1 = \frac{n+2}{4} (\max_M |R_0| + \max_M |f|)$ . It is clear that (2-5) implies that

$$\max_{[0,t] \times M} u \leq e^{C_1 t}.$$

The proof of Proposition 2.2 is then complete. □

Now we prove integral estimates on  $R_g$  which will imply estimates on  $\partial_t u$ :

**Proposition 2.3.** *Let  $g(t)$  be the solution of (1.3) defined on a maximal interval  $[0, T^*)$ . Then we have for any  $t \in [0, T^*)$ ,*

$$(2-6) \quad \int_M |R_{g(t)} - f|^p dV_{g(t)} \leq C e^{Ct}$$

where  $p = \frac{n^2}{2(n-2)}$  and  $C$  is a positive constant depending only on  $f, g_0, u_0$ .



*Proof.* In what follows  $C$  denotes a positive constant depending on  $f, g_0, u_0$ , whose value may change from line to line.

We have by Lemma 2.1 for any  $t \in [0, T^*)$

$$(2-7) \quad \frac{d}{dt} \int_M |R_g - f|^p dV_g = -\frac{4(n-1)(p-1)}{p} \int_M |\nabla_g |R_g - f|^{\frac{p}{2}}|_g^2 dV_g + \left(p - \frac{n}{2}\right) \int_M (R_g - f) |R_g - f|^p dV_g + p \int_M f |R_g - f|^p dV_g,$$

where  $\nabla_g$  is the gradient with respect to the metric  $g$  and  $|\cdot|_g$  is the Riemannian norm with respect to  $g$ . It follows from (2-7) that

$$(2-8) \quad \frac{d}{dt} \int_M |R_g - f|^p dV_g + \frac{4(n-1)(p-1)}{p} \int_M |\nabla_g |R_g - f|^{\frac{p}{2}}|_g^2 dV_g \leq \left|p - \frac{n}{2}\right| \int_M |R_g - f|^{p+1} dV_g + C \int_M |R_g - f|^p dV_g.$$

By (2-4) we have

$$(2-9) \quad \int_M |\nabla_g |R_g - f|^{\frac{p}{2}}|_g^2 dV_g = \int_M |\nabla |R_g - f|^{\frac{p}{2}}|^2 u^2 dV_{g_0} \geq C \int_M |\nabla |R_g - f|^{\frac{p}{2}}|^2 dV_{g_0}$$

and

$$(2-10) \quad \int_M |R_g - f|^p dV_g = \int_M |R_g - f|^p u^{\frac{2n}{n-2}} dV_{g_0} \geq C \int_M |R_g - f|^p dV_{g_0}.$$

By Sobolev's inequality we have

$$\left(\int_M |R_g - f|^{\frac{pn}{n-2}} dV_{g_0}\right)^{\frac{n-2}{n}} \leq C \left(\int_M |\nabla |R_g - f|^{\frac{p}{2}}|^2 dV_{g_0} + \int_M |R_g - f|^p dV_{g_0}\right)$$

which, using (2-4), gives

$$(2-11) \quad \left(\int_M |R_g - f|^{\frac{pn}{n-2}} dV_g\right)^{\frac{n-2}{n}} \leq C e^{Ct} \left(\int_M |\nabla |R_g - f|^{\frac{p}{2}}|^2 dV_{g_0} + \int_M |R_g - f|^p dV_{g_0}\right).$$

It follows from (2-8), (2-9), (2-10) and (2-11) that

$$(2-12) \quad \frac{d}{dt} \int_M |R_g - f|^p dV_g + C^{-1} e^{-Ct} \left(\int_M |R_g - f|^{\frac{pn}{n-2}} dV_g\right)^{\frac{n-2}{n}} \leq \left(p - \frac{n}{2}\right) \int_M |R_g - f|^{p+1} dV_g + C \int_M |R_g - f|^p dV_g.$$

By taking  $p = \frac{n}{2}$  in (2-12) we get

$$\frac{d}{dt} \int_M |R_g - f|^p dV_g \leq C \int_M |R_g - f|^p dV_g,$$

which implies that

$$(2-13) \quad \int_M |R_g - f|^{\frac{n}{2}} dV_g \leq C e^{Ct}.$$

Now taking again  $p = \frac{n}{2}$  in (2-12) and integrating on  $[0, t], t \in [0, T^*)$ , by using (2-13) we obtain

$$(2-14) \quad \int_0^t \left( \int_M |R_{g(s)} - f|^{\frac{n^2}{2(n-2)}} dV_{g(s)} \right)^{\frac{n-2}{n}} ds \leq C e^{Ct}.$$

We have by Hölder’s inequality and Young’s inequality, for any  $\varepsilon > 0$  and  $p > \frac{n}{2}$ ,

$$(2-15) \quad \int_M |R_g - f|^{p+1} dV_g \leq \varepsilon \left( \int_M |R_g - f|^{\frac{pn}{n-2}} dV_g \right)^{\frac{n-2}{n}} + \varepsilon^{-\frac{n}{2p-n}} \left( \int_M |R_g - f|^p dV_g \right)^{\frac{2p-n+2}{2p-n}}.$$

If we combine (2-15) with (2-12) and taking  $\varepsilon = (p - \frac{n}{2})^{-1} C^{-1} e^{-Ct}$ , we get

$$\frac{d}{dt} \int_M |R_g - f|^p dV_g \leq C e^{Ct} \left( \int_M |R_g - f|^p dV_g \right)^{\frac{2p-n+2}{2p-n}} + C \int_M |R_g - f|^p dV_g,$$

that is,

$$\frac{d}{dt} \log \left( \int_M |R_g - f|^p dV_g \right) \leq C \left( e^{Ct} \left( \int_M |R_g - f|^p dV_g \right)^{\frac{2}{2p-n}} + 1 \right)$$

In particular by choosing  $p = \frac{n^2}{2(n-2)}$  and integrating on  $[0, t], t \in [0, T^*)$ , we obtain

$$\begin{aligned} \log \left( \int_M |R_{g(t)} - f|^{\frac{n^2}{2(n-2)}} dV_{g(t)} \right) &\leq \log \left( \int_M |R_{g(0)} - f|^{\frac{n^2}{2(n-2)}} dV_{g(0)} \right) \\ &\quad + C e^{Ct} \int_0^t \left( \int_M |R_{g(s)} - f|^{\frac{n^2}{2(n-2)}} dV_{g(s)} \right)^{\frac{n-2}{n}} ds + Ct \end{aligned}$$

which by using (2-14) gives

$$\log \left( \int_M |R_{g(t)} - f|^{\frac{n^2}{2(n-2)}} dV_{g(t)} \right) \leq C e^t.$$

This proves Proposition 2.3. □

With the estimates of Proposition 2.2 we would like to apply the classical Schauder estimates for parabolic equations. To this end we need  $C^\alpha$ -estimates:

**Proposition 2.4.** *Let  $g(t) = u(t)^{4/(n-2)}g_0$  be the solution of (1-3) defined on a maximal interval  $[0, T^*)$ . Then we have for some  $\alpha \in (0, 1)$  and any  $T \in [0, T^*)$*

$$\|u\|_{C^\alpha([0,T] \times M)} \leq Ce^{CT},$$

where  $C$  is a positive constant depending only on  $u_0, g_0$  and  $f$ .

*Proof.* By using Propositions 2.2 and 2.3, the proof is identical to that of Proposition 2.6 in Brendle [2005]. □

*Proof of Theorem 1.1.* Let  $g(t) = u(t)^{4/(n-2)}g_0$  be the solution of (1-3) defined on a maximal interval  $[0, T^*)$ . Assume by contradiction that  $T^* < +\infty$ . Then by using Propositions 2.2 and 2.4 we have

$$\|u\|_{C^\alpha([0,T^*) \times M)} \leq Ce^{CT^*} \quad \text{and} \quad \min_{[0,T^*) \times M} u \geq \min(C_0, \min_M u_0)$$

for some  $\alpha \in (0, 1)$ , where  $C$  is a positive constant depending  $u_0, f, g_0$ . The classical theory of linear parabolic equations applied to (1-4) implies that  $u$  is bounded in  $C^k([0, T^*) \times M)$  for any  $k \in \mathbb{N}$ , that is,

$$(2-16) \quad \|u\|_{C^k([0,T^*) \times M)} \leq C_k,$$

where  $C_k$  is a positive constant depending only on  $u_0, g_0, f$  and  $k$ . It is clear that (2-16) allows us to extend the solution beyond  $T^*$  contradicting thus the maximality of  $T^*$ . We see from (2-2) that the functional  $\mathcal{E}$  is decreasing along the flow. The proof of Theorem 1.1 is then complete. □

### 3. Long Time behaviour of the flow

In this section we study the asymptotic behaviour of the flow  $g(t)$  when  $t \rightarrow +\infty$ . First we prove the following proposition which gives a super solution of (1-1) when conditions **(H1)** and **(H2)** are satisfied.

**Proposition 3.1.** *Suppose that there exists an open set  $\Omega \subset M$  such that conditions **(H1)** and **(H2)** are satisfied. Then there exists a conformal metric  $\bar{g} = \bar{u}^{4/(n-2)}g_0$ ,  $0 < \bar{u} \in C^\infty(M)$ , satisfying*

$$(3-1) \quad R_{\bar{g}} - f \geq 0$$

or equivalently

$$(3-2) \quad -c_n \Delta \bar{u} + R_0 \bar{u} - f \bar{u}^N \geq 0, \quad N = \frac{n+2}{n-2}.$$

*Proof.* By hypothesis, there is an open set  $\Omega \subset M$  satisfying **(H1)** and **(H2)**, that is,

$$(H1) \quad \lambda_\Omega > 0 \text{ and } f < 0 \text{ on } M \setminus \Omega$$

and

$$(H2) \quad \sup_{x \in \Omega} f(x) \leq C_\Omega \inf_{x \in M \setminus \Omega} |f(x)|,$$

where  $C_\Omega$  is a positive constant depending only on  $\Omega$ .

Let  $\varepsilon > 0$  and set

$$\Omega_\varepsilon = \{x \in M : d(x, \Omega) < \varepsilon\}.$$

For  $\varepsilon > 0$  sufficiently small we have from **(H1)** that  $\lambda_{\Omega_\varepsilon} > 0$ , where  $\lambda_{\Omega_\varepsilon}$  is the first eigenvalue of the operator  $-c_n \Delta + R_0$  on  $\Omega_\varepsilon$  with zero Dirichlet boundary conditions. Let  $D \subset M$  be an open set of smooth boundary such that  $\bar{\Omega} \subset D \subset \Omega_\varepsilon$ . Then we have  $\lambda_D \geq \lambda_{\Omega_\varepsilon} > 0$ . Let  $\varphi_0$  an eigenfunction associated with  $\lambda_D$ , that is,

$$-c_n \Delta \varphi_0 + R_0 \varphi_0 = \lambda_D \varphi_0.$$

Then we have that  $\varphi_0 \in C^\infty(\bar{D})$  and using the maximum principle of elliptic equations we have  $\varphi_0 > 0$  on  $D$ . By normalising if necessary, we may suppose that

$$(3-3) \quad 0 < \varphi_0 \leq 1 \text{ on } D.$$

Let  $\chi \in C_0^\infty(D)$  such that  $0 \leq \chi \leq 1$  and  $\chi = 1$  on  $\bar{\Omega}$ . We define the function  $\bar{u} \in C^\infty(M)$  by setting

$$\bar{u} = \delta(\chi \varphi_0 + 1 - \chi),$$

where  $\delta > 0$  will be chosen later. By (3-3) and the definition of  $\chi$  it is easy to check

$$m_0 := \inf_M (\chi \varphi_0 + 1 - \chi) > 0,$$

so

$$(3-4) \quad \bar{u} \geq \delta m_0.$$

Now let us prove that  $\bar{u}$  satisfies (3-2). If we set

$$\mathcal{L}(\bar{u}) = -c_n \Delta \bar{u} + R_0 \bar{u} - f \bar{u}^{\frac{n+2}{n-2}},$$

then (3-2) is equivalent to  $\mathcal{L}(\bar{u}) \geq 0$ .

A simple computation shows that we have on  $\Omega$  (using the fact that  $\chi = 1$  on  $\bar{\Omega}$ ):

$$\mathcal{L}(\bar{u}) = \lambda_D \delta \varphi_0 - f \delta^N \varphi_0^N = \delta \varphi_0 (\lambda_D - \delta^{N-1} f \varphi_0^{N-1})$$

and by using (3-3) it follows that

$$(3-5) \quad \mathcal{L}(\bar{u}) \geq \delta \varphi_0 (\lambda_D - \delta^{N-1} \sup_{x \in \Omega} f(x)).$$

It follows from (3-5) that if we want  $\mathcal{L}(\bar{u}) \geq 0$  on  $\Omega$ , we have to choose  $\delta > 0$  satisfying

$$(3-6) \quad \delta^{N-1} \sup_{x \in \Omega} f(x) \leq \lambda_D.$$

Now we examine the sign of  $\mathcal{L}(\bar{u})$  on  $M \setminus \Omega$ . We have from the definition of  $\bar{u}$  that

$$(3-7) \quad \mathcal{L}(\bar{u}) = \delta(-c_n \Delta + R_0)(\chi \varphi_0 + 1 - \chi) - f \bar{u}^N.$$

By using (3-4) and the fact that  $f < 0$  on  $M \setminus \Omega$ , it follows from (3-7) that

$$\mathcal{L}(\bar{u}) \geq -\delta m_1 + \delta^N m_0^N \inf_{x \in M \setminus \Omega} |f(x)|,$$

where

$$m_1 = \sup_M |(-c_n \Delta + R_0)(\chi \varphi_0 + 1 - \chi)|.$$

Thus, if we want  $\mathcal{L}(\bar{u}) \geq 0$  on  $M \setminus \Omega$ , we have to assume

$$-m_1 + \delta^{N-1} m_0^N \inf_{x \in M \setminus \Omega} |f(x)| \geq 0,$$

that is,

$$(3-8) \quad \delta^{N-1} \inf_{x \in M \setminus \Omega} |f(x)| \geq m_1 m_0^{-N}.$$

It is clear that the existence of  $\delta > 0$  satisfying both (3-6) and (3-8) is equivalent to condition **(H2)** with  $C_\Omega = \lambda_D m_0^N / m_1$ . □

Proposition 3.1 allows us to prove uniform  $L^\infty$ -estimates on the flow.

**Proposition 3.2.** *Let  $0 < u_0 \in C^\infty(M)$  such that  $u_0 \leq \bar{u}$  where  $\bar{u}$  is given by Proposition 3.1. Then the solution  $u$  of (1-4) satisfies, for any  $(t, x) \in [0, +\infty) \times M$ ,*

$$(3-9) \quad \min(C_0, \min_M u_0) \leq u(t, x) \leq \max_M \bar{u},$$

where

$$C_0 = \left( \min_M |R_0| / \max_M |f| \right)^{\frac{n-2}{4}}.$$

*Proof.* First observe that the first inequality in (3-9) has already been proved in Proposition 2.2. It remains then to prove the second inequality, that is,

$$u(t, x) \leq \max_M \bar{u}.$$

Let  $v = \bar{u} - u$ . Since  $u$  satisfies (1-4) and  $\bar{u}$  satisfies (3-2), we have

$$(3-10) \quad \partial_t(\bar{u}^N - u^N) \geq \frac{n+2}{4}(c_n \Delta v - R_0 v + f(\bar{u}^N - u^N)).$$

We have  $\bar{u}^N - u^N = av$ , where

$$a(t, x) = N \int_0^1 (s\bar{u}(t, x) + (1-s)u(t, x))^{N-1} ds,$$

so it follows from (3-10) that

$$(3-11) \quad \partial_t(av) \geq \frac{n+2}{4}(c_n \Delta v - R_0 v + afv).$$

Since  $v(0, x) = \bar{u}(x) - u_0(x) \geq 0$ , by applying the maximum principle to (3-11) we get  $v(t, x) \geq 0$  for any  $t \geq 0$ , that is,

$$u(t, x) \leq \bar{u}(x). \quad \square$$

Now we prove that the integral estimate (2-6) in Proposition 2.3 can be improved when  $t \rightarrow +\infty$ . More precisely, we have:

**Proposition 3.3.** *Let  $0 < u_0 \in C^\infty(M)$  such that  $u_0 \leq \bar{u}$  where  $\bar{u}$  is given by Proposition 3.1. Let  $g(t)$  be the solution of (1-3) given by Theorem 1.1 such that  $g(0) = u_0^{4/(n-2)} g_0$ . Then we have, for any  $p \geq 1$ ,*

$$(3-12) \quad \lim_{t \rightarrow +\infty} \int_M |R_{g(t)} - f|^p dV_{g(t)} = 0.$$

*Proof.* In what follows  $C$  denotes a positive constant depending only on  $u_0, g_0, f, p$ , and its value may change from line to line.

We have by (2-2) for any  $t \geq 0$ ,

$$(3-13) \quad \frac{n-2}{2} \int_0^t \int_M |R_g - f|^2 dV_g = \mathcal{E}(g(0)) - \mathcal{E}(g(t)).$$

On the other hand, we have

$$\mathcal{E}(g(t)) = \int_M \left( c_n |\nabla u|^2 + R_0 u^2 - \frac{n-2}{n} f u^{\frac{2n}{n-2}} \right) dV_{g_0},$$

and since  $u$  is uniformly bounded by Proposition 3.2, we have  $\mathcal{E}(g(t)) \geq -C$ . So it follows from (3-13) that

$$(3-14) \quad \int_0^{+\infty} \int_M |R_{g(t)} - f|^2 dV_{g(t)} \leq C.$$

Since by Proposition 3.2 the volume of  $g(t)$  is uniformly bounded, it suffices to prove (3-12) for a sequence  $p_k \rightarrow +\infty$ . We shall prove (3-12) by induction when  $p = p_k$ , where

$$p_k := \frac{n}{2} \left( \frac{n}{n-2} \right)^k, \quad k \in \mathbb{N}.$$

First we prove (3-12) for  $p_0 = \frac{n}{2}$ . As in the proof of Proposition 2.3, if we use Lemma 2.1 and the fact that  $u$  is uniformly bounded by Proposition 3.2, then we

have for any  $p > 1$ :

$$(3-15) \quad \frac{d}{dt} \int_M |R_g - f|^p dV_g + C^{-1} \left( \int_M |R_g - f|^{\frac{pn}{n-2}} dV_g \right)^{\frac{n-2}{n}} \leq C \int_M |R_g - f|^p dV_g + \left( p - \frac{n}{2} \right) \int_M |R_g - f|^{p+1} dV_g.$$

Set

$$\phi_p(t) = \int_M |R_g - f|^p dV_g.$$

If  $p_0 < 2$ , then by using Hölder’s inequality and the fact that  $u$  is uniformly bounded, we have

$$(3-16) \quad \phi_{p_0} \leq C \phi_2^{p_0/2}.$$

So it follows from (3-15) by taking  $p = p_0 = \frac{n}{2}$  that

$$(3-17) \quad \frac{d}{dt} \phi_{p_0}^{2/p_0} \leq C \phi_2.$$

By (3-14) there is a sequence  $t_\nu \rightarrow +\infty$  such that  $\int_{t_\nu}^{+\infty} \phi_2(s) ds \rightarrow 0$  and  $\phi_2(t_\nu) \rightarrow 0$ . So by integrating (3-17) on  $[t_\nu, t]$  and using (3-16) we get

$$\phi_{p_0}^{2/p_0}(t) \leq \phi_{p_0}^{2/p_0}(t_\nu) + C \int_{t_\nu}^t \phi_2(s) ds \leq C \phi_2(t_\nu) + C \int_{t_\nu}^t \phi_2(s) ds$$

Letting  $t \rightarrow +\infty$  and  $\nu \rightarrow +\infty$  we obtain  $\phi_{p_0}(t) \rightarrow 0$  as  $t \rightarrow +\infty$ .

If  $p_0 \geq 2$ , by using Hölder’s inequality and Young’s inequality we have, for any  $\varepsilon > 0$ ,

$$\begin{aligned} & \int_M |R_g - f|^{p_0} dV_g \\ & \leq \varepsilon \left( \int_M |R_g - f|^{p_0 n/n-2} dV_g \right)^{\frac{n-2}{n}} + \varepsilon^{-n(p_0-2)/4} \left( \int_M |R_g - f|^2 dV_g \right)^{p_0/2}. \end{aligned}$$

By taking  $\varepsilon = \frac{1}{2} C^{-1}$ , where  $C^{-1}$  is the constant appearing in (3-15), we obtain from (3-15)(where we take  $p = p_0 = \frac{n}{2}$ ),

$$(3-18) \quad \frac{d}{dt} \phi_{p_0} + C^{-1} \phi_{p_0 n/(n-2)}^{(n-2)/n} \leq C \phi_2^{p_0/2}.$$

But by Hölder’s inequality, since the volume of  $g$  is uniformly bounded, we have

$$\phi_2 \leq C \phi_{p_0}^{2/p_0} \quad \text{and} \quad \phi_{p_0} \leq C \phi_{p_0 n/(n-2)}^{(n-2)/n}.$$

Thus it follows from (3-18) that

$$(3-19) \quad \frac{d}{dt} \phi_{p_0}^{2/p_0} + C^{-1} \phi_{p_0}^{2/p_0} \leq C \phi_2.$$

If we integrate (3-19) on  $[0, t]$  and use (3-14) we get

$$\int_0^t \phi_{p_0}^{2/p_0}(s) ds \leq C,$$

which implies, since  $t \geq 0$  is arbitrary,

$$\int_0^{+\infty} \phi_{p_0}^{2/p_0}(s) ds \leq C.$$

Thus there exists a sequence  $t_\nu \rightarrow +\infty$  such that  $\phi_{p_0}^{2/p_0}(t_\nu) \rightarrow 0$  as  $\nu \rightarrow +\infty$ . If we integrate again (3-19) on  $[t_\nu, t]$ , we obtain

$$\phi_{p_0}^{2/p_0}(t) \leq \phi_{p_0}^{2/p_0}(t_\nu) + C \int_{t_\nu}^t \phi_2(s) ds.$$

By using (3-14), it follow that  $\phi_{p_0}^{2/p_0}(t) \rightarrow 0$  as  $t \rightarrow +\infty$ .

Now suppose by induction that

$$(3-20) \quad \lim_{t \rightarrow +\infty} \phi_{p_k}(t) = 0.$$

First let us prove that

$$(3-21) \quad \lim_{t \rightarrow +\infty} \int_t^{t+1} \phi_{p_{k+1}}^{(n-2)/n}(s) ds = 0.$$

We may suppose  $k \geq 1$ . Indeed, if  $k = 0$  (that is,  $p_k = p_0 = \frac{n}{2}$ ), then (3-21) follows directly from (3-15) (with  $p = \frac{n}{2}$ ) by integrating on  $[t, t + 1]$  and using (3-20). Thus let us prove (3-21) when  $k \geq 1$ .

By using Hölder's inequality and Young's inequality we have for any  $p > \frac{n}{2}$  and  $\varepsilon > 0$ ,

$$(3-22) \quad \int_M |R_g - f|^{p+1} dV_g \leq \varepsilon \left( \int_M |R_g - f|^{\frac{pn}{n-2}} dV_g \right)^{\frac{n-2}{n}} + \varepsilon^{-\frac{n}{2p-n}} \left( \int_M |R_g - f|^p dV_g \right)^{1+\frac{2}{2p-n}}$$

By taking  $p = p_k$ ,  $\varepsilon = \frac{1}{2}C^{-1}$ , where  $C$  is the constant appearing in (3-15), from (3-15) we obtain

$$\frac{d}{dt} \phi_{p_k} + \frac{1}{2}C^{-1} \phi_{p_{k+1}}^{(n-2)/n} \leq C \phi_{p_k}^{1+2/(2p_k-n)} + C \phi_{p_k}.$$

Then (3-21) follows by integrating on  $[t, t + 1]$  and using (3-20).

Now if we apply (3-22) by taking  $p = p_{k+1}$  and  $\varepsilon = C^{-1}/(p_{k+1} - n/2)$ , where  $C$  is the constant appearing in (3-15), we obtain from (3-15) (where we take  $p = p_{k+1}$ ),

$$\frac{d}{dt} \phi_{p_{k+1}} \leq C \phi_{p_{k+1}}^{1+\alpha_k} + C \phi_{p_{k+1}},$$



where  $\alpha_k = \frac{2}{2p_{k+1}} - n$ . The last inequality is equivalent to

$$(3-23) \quad \frac{d}{dt} \log \phi_{p_{k+1}} \leq C(\phi_{p_{k+1}}^{\alpha_k} + 1).$$

By (3-21) there is a sequence  $t_\nu \rightarrow +\infty$  such that  $\nu \leq t_\nu \leq \nu + 1$  satisfying  $\phi_{p_{k+1}}(t_\nu) \rightarrow 0$  as  $\nu \rightarrow +\infty$ . If we integrate (3-23) on  $[t_\nu, t]$  where  $t \in [\nu, \nu + 1]$ , we obtain

$$(3-24) \quad \log \frac{\phi_{p_{k+1}}(t)}{\phi_{p_{k+1}}(t_\nu)} \leq C \left( \int_\nu^{t_\nu} \phi_{p_{k+1}}^{\alpha_k}(s) ds + 1 \right).$$

We note here that  $\alpha_k \leq \frac{n-2}{n}$ , so by Hölder's inequality we have

$$\int_\nu^{t_\nu} \phi_{p_{k+1}}^{\alpha_k}(s) ds \leq \left( \int_\nu^{t_\nu} \phi_{p_{k+1}}^{(n-2)/n}(s) ds \right)^{\frac{n\alpha_k}{n-2}} \rightarrow 0 \text{ as } \nu \rightarrow +\infty$$

by (3-21). Thus it follows from (3-24) that

$$\log \frac{\phi_{p_{k+1}}(t)}{\phi_{p_{k+1}}(t_\nu)} \leq C,$$

which implies that  $\phi_{p_{k+1}}(t) \rightarrow 0$  as  $t \rightarrow +\infty$ . The proof of Proposition 3.3 is then complete. □

Now we can prove uniform  $C^\alpha$ -estimates on the solution.

**Proposition 3.4.** *Let  $0 < u_0 \in C^\infty(M)$  such that  $u_0 \leq \bar{u}$  where  $\bar{u}$  is given by Proposition 3.1. Then the solution  $u$  of (1-4) satisfies, for some  $\alpha \in (0, 1)$ ,*

$$\|u\|_{C^\alpha([0, +\infty) \times M)} \leq C,$$

where  $C$  is a positive constant depending only on  $u_0, g_0$  and  $f$ .

*Proof.* By using Propositions 3.2 and 3.3, the proof is identical to that of Proposition 2.6 in Brendle [2005]. □

Now we are in position to prove Theorem 1.2.

*Proof of Theorem 1.2.* Let  $g = u^{4/(n-2)}g_0$  be the solution of (1-3) given by Theorem 1.1. By Proposition 3.2 we have that  $u$  is bounded from below and above uniformly on  $[0, +\infty)$ . As in the proof of Theorem 1.1, this implies that (1-4) is uniformly parabolic and by Proposition 3.4 we have a uniform  $C^\alpha$ -bound on the solution  $u$  on  $[0, +\infty) \times M$ . We then apply the classical regularity theory of linear parabolic equations to obtain uniform  $C^k$ -bound for any  $k \in \mathbb{N}$ , that is,

$$(3-25) \quad \|u(t)\|_{C^k(M)} \leq C_k,$$

for some constant  $C_k$  independent of  $t$ . It follows from (3-25) that there is a sequence  $t_\nu \rightarrow +\infty$  such that  $u(t_\nu)$  converges in  $C^k(M)$  for any  $k \in \mathbb{N}$ , to some function

$u_\infty \in C^\infty(M)$ . Since  $u(t)$  is uniformly bounded from below by Proposition 3.2, then we have  $u_\infty > 0$ . By using Proposition 3.3 and passing to the limit when  $\nu \rightarrow \infty$ , we see that  $R_{g_\infty} = f$ , where  $g_\infty = u_\infty^{4/(n-2)}g_0$ , that is,  $f$  is the scalar curvature of  $g_\infty$ . By the general result of Simon [1983] on evolution equations,  $u_\infty$  is the unique limit of  $u(t)$  when  $t \rightarrow +\infty$ .  $\square$

*Proof of Corollary 1.3.* Since  $f < 0$  almost everywhere on  $M$ , then for  $\varepsilon > 0$  small enough, the open set

$$\Omega_\varepsilon = \{x \in M : f(x) > -\varepsilon\}$$

has arbitrary small volume. This implies that the first eigenvalue  $\mu_{\Omega_\varepsilon}$  of  $-c_n \Delta$  on  $\Omega_\varepsilon$  with zero Dirichlet conditions is arbitrarily large if  $\varepsilon$  is small enough. But since

$$\lambda_{\Omega_\varepsilon} \geq \mu_{\Omega_\varepsilon} + \min_M R_0,$$

we have  $\lambda_{\Omega_\varepsilon} > 0$  if  $\varepsilon$  is small enough. Thus the condition **(H1)** is satisfied with  $\Omega = \Omega_\varepsilon$ . Condition **(H2)** is also satisfied since by continuity of  $f$  we have  $f \leq 0$  everywhere on  $M$ .  $\square$

*Proof of Theorem 1.4.* Suppose that condition **(H1)** is not satisfied, that is, for any open set  $\Omega \subset M$  such that  $f < 0$  on  $M \setminus \Omega$ , we suppose  $\lambda_\Omega \leq 0$ . For  $\varepsilon > 0$ , consider the following family of open sets:

$$\Omega_\varepsilon = \{x \in M : f(x) > -\varepsilon\}.$$

For simplicity of notation we set  $\lambda_\varepsilon = \lambda_{\Omega_\varepsilon}$ . According to our hypothesis we have

$$(3-26) \quad \lambda_\varepsilon \leq 0 \quad \text{for all } \varepsilon > 0.$$

By using Sard's theorem, there exists a sequence  $\varepsilon_n \rightarrow 0$  such that  $\varepsilon_n$  is a regular value of  $f$  and then  $\Omega_{\varepsilon_n}$  has a smooth boundary

$$\partial\Omega_{\varepsilon_n} = \{x \in M : f(x) = -\varepsilon_n\}.$$

Let  $\varphi_n$  an eigenfunction of  $-c_n \Delta + R_0$  associated with  $\lambda_{\varepsilon_n}$ . As already mentioned in the proof of Proposition 3.2, we have by the maximum principle that

$$(3-27) \quad \varphi_n > 0 \text{ on } \Omega_{\varepsilon_n} \quad \text{and} \quad \frac{\partial\varphi_n}{\partial\nu} \leq 0 \text{ on } \partial\Omega_{\varepsilon_n},$$

where  $\nu$  is the outer normal vector to  $\partial\Omega_{\varepsilon_n}$ . By normalising if necessary, we may assume that

$$(3-28) \quad \int_{\Omega_{\varepsilon_n}} \varphi_n dV_{g_0} = 1.$$

If we multiply (1-4) by  $\varphi_n$  and integrate on  $\Omega_{\varepsilon_n}$ , we have

$$(3-29) \quad \begin{aligned} \frac{d}{dt} \int_{\Omega_{\varepsilon_n}} u^N \varphi_n dV_{g_0} \\ = \frac{n+2}{4} \int_{\Omega_{\varepsilon_n}} (c_n \Delta u - R_0 u) \varphi_n dV_{g_0} + \frac{n+2}{4} \int_{\Omega_{\varepsilon_n}} f u^N \varphi_n dV_{g_0}. \end{aligned}$$

An integration by parts gives

$$\int_{\Omega_{\varepsilon_n}} (c_n \Delta u - R_0 u) \varphi_n dV_{g_0} = -\lambda_{\varepsilon_n} \int_{\Omega_{\varepsilon_n}} u \varphi_n dV_{g_0} - c_n \int_{\partial\Omega_{\varepsilon_n}} \frac{\partial \varphi_n}{\partial \nu} u dV_{g_0}.$$

Since  $\lambda_{\varepsilon_n} \leq 0$ , by using (3-27) and (3-28) we then obtain

$$(3-30) \quad \int_{\Omega_{\varepsilon_n}} (c_n \Delta u - R_0 u) \varphi_n dV_{g_0} \geq -\lambda_{\varepsilon_n} \inf_M u - c_n \inf_M u \int_{\partial\Omega_{\varepsilon_n}} \frac{\partial \varphi_n}{\partial \nu} dV_{g_0}.$$

On the other hand we have

$$c_n \int_{\partial\Omega_{\varepsilon_n}} \frac{\partial \varphi_n}{\partial \nu} dV_{g_0} = c_n \int_{\Omega_{\varepsilon_n}} \Delta \varphi_n dV_{g_0} = \int_{\Omega_{\varepsilon_n}} (-\lambda_{\varepsilon_n} + R_0) \varphi_n dV_{g_0}$$

and by using (3-28) we get, since  $R_0 < 0$ ,

$$(3-31) \quad -c_n \int_{\partial\Omega_{\varepsilon_n}} \frac{\partial \varphi_n}{\partial \nu} dV_{g_0} \geq \lambda_{\varepsilon_n} + \inf_M |R_0|.$$

Combining (3-30) and (3-31) we obtain

$$\int_{\Omega_{\varepsilon_n}} (c_n \Delta u - R_0 u) \varphi_n dV_{g_0} \geq \inf_M |R_0| \inf_M u.$$

If we substitute in (3-29) we get

$$(3-32) \quad \frac{d}{dt} \int_{\Omega_{\varepsilon_n}} u^N \varphi_n dV_{g_0} \geq \frac{n+2}{4} \inf_M |R_0| \inf_M u + \frac{n+2}{4} \int_{\Omega_{\varepsilon_n}} f u^N \varphi_n dV_{g_0}.$$

By Proposition 3.2 we have  $u \geq C_0$ , where  $C_0$  is a positive constant depending only on  $u_0, g_0$  and  $f$ . Using the fact that  $f > -\varepsilon_n$  on  $\Omega_{\varepsilon_n}$ , it follows from (3-32) that

$$\frac{d}{dt} \int_{\Omega_{\varepsilon_n}} u^N \varphi_n dV_{g_0} \geq C - \frac{n+2}{4} \varepsilon_n \int_{\Omega_{\varepsilon_n}} u^N \varphi_n dV_{g_0},$$

where  $C$  is a positive constant depending only on  $u_0, g_0$  and  $f$ . By integrating this differential inequality on  $[0, t]$ , we get

$$\int_{\Omega_{\varepsilon_n}} u^N(t) \varphi_n dV_{g_0} \geq \int_{\Omega_{\varepsilon_n}} u_0^N \varphi_n dV_{g_0} + Ct - \frac{n+2}{4} \varepsilon_n \int_0^t \int_{\Omega_{\varepsilon_n}} u^N(s) \varphi_n dV_{g_0} ds,$$

which, using (3-28), implies

$$\max_{x \in M} u^N(t, x) \geq Ct - \frac{n+2}{4} \varepsilon_n \int_0^t \max_{x \in M} u^N(s, x) ds.$$

Letting  $n \rightarrow +\infty$ , we obtain

$$\max_{x \in M} u^N(t, x) \geq Ct.$$

The proof of Theorem 1.4 is complete. □

### References

- [Ambrosetti and Malchiodi 1999] A. Ambrosetti and A. Malchiodi, “A multiplicity result for the Yamabe problem on  $S^n$ ”, *J. Funct. Anal.* **168**:2 (1999), 529–561. MR Zbl
- [Aubin 1976] T. Aubin, “Équations différentielles non linéaires et problème de Yamabe concernant la courbure scalaire”, *J. Math. Pures Appl.* (9) **55**:3 (1976), 269–296. MR Zbl
- [Baird et al. 2004] P. Baird, A. Fardoun, and R. Regbaoui, “The evolution of the scalar curvature of a surface to a prescribed function”, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) **3**:1 (2004), 17–38. MR Zbl
- [Besse 1987] A. L. Besse, *Einstein manifolds*, Ergebnisse der Mathematik (3) **10**, Springer, 1987. MR Zbl
- [Bismuth 2000] S. Bismuth, “Prescribed scalar curvature on a complete Riemannian manifold in the negative case”, *J. Math. Pures Appl.* (9) **79**:9 (2000), 941–951. MR Zbl
- [Brendle 2005] S. Brendle, “Convergence of the Yamabe flow for arbitrary initial energy”, *J. Differential Geom.* **69**:2 (2005), 217–278. MR Zbl
- [Brendle 2007] S. Brendle, “Convergence of the Yamabe flow in dimension 6 and higher”, *Invent. Math.* **170**:3 (2007), 541–576. MR Zbl
- [Chen and Xu 2012] X. Chen and X. Xu, “The scalar curvature flow on  $S^n$ -perturbation theorem revisited”, *Invent. Math.* **187**:2 (2012), 395–506. Erratum on pp. 507–509. MR Zbl
- [Chow 1992] B. Chow, “The Yamabe flow on locally conformally flat manifolds with positive Ricci curvature”, *Comm. Pure Appl. Math.* **45**:8 (1992), 1003–1014. MR Zbl
- [Escobar and Schoen 1986] J. F. Escobar and R. M. Schoen, “Conformal metrics with prescribed scalar curvature”, *Invent. Math.* **86**:2 (1986), 243–254. MR Zbl
- [Hamilton 1988] R. S. Hamilton, “The Ricci flow on surfaces”, pp. 237–262 in *Mathematics and general relativity* (Santa Cruz, 1986), edited by J. A. Isenberg, Contemp. Math. **71**, Amer. Math. Soc., Providence, RI, 1988. MR Zbl
- [Hebey and Vaugon 1993] E. Hebey and M. Vaugon, “Le problème de Yamabe équivariant”, *Bull. Sci. Math.* **117**:2 (1993), 241–286. MR Zbl
- [Kazdan and Warner 1975] J. L. Kazdan and F. W. Warner, “Existence and conformal deformation of metrics with prescribed Gaussian and scalar curvatures”, *Ann. of Math.* (2) **101** (1975), 317–331. MR Zbl
- [Lee and Parker 1987] J. M. Lee and T. H. Parker, “The Yamabe problem”, *Bull. Amer. Math. Soc. (N.S.)* **17**:1 (1987), 37–91. MR Zbl
- [Rauzy 1995] A. Rauzy, “Courbures scalaires des variétés d’invariant conforme négatif”, *Trans. Amer. Math. Soc.* **347**:12 (1995), 4729–4745. MR Zbl

- [Schoen 1984] R. Schoen, “Conformal deformation of a Riemannian metric to constant scalar curvature”, *J. Differential Geom.* **20**:2 (1984), 479–495. MR Zbl
- [Schoen 1991] R. M. Schoen, “A report on some recent progress on nonlinear problems in geometry”, pp. 201–241 in *Surveys in differential geometry* (Cambridge, 1990), edited by C. L. Terng and K. Uhlenbeck, Lehigh Univ., Bethlehem, PA, 1991. MR Zbl
- [Schwetlick and Struwe 2003] H. Schwetlick and M. Struwe, “Convergence of the Yamabe flow for ‘large’ energies”, *J. Reine Angew. Math.* **562** (2003), 59–100. MR Zbl
- [Simon 1983] L. Simon, “Asymptotics for a class of nonlinear evolution equations, with applications to geometric problems”, *Ann. of Math. (2)* **118**:3 (1983), 525–571. MR Zbl
- [Struwe 2005] M. Struwe, “A flow approach to Nirenberg’s problem”, *Duke Math. J.* **128**:1 (2005), 19–64. MR Zbl
- [Vázquez and Véron 1991] J. L. Vázquez and L. Véron, “Solutions positives d’équations elliptiques semi-linéaires sur des variétés riemanniennes compactes”, *C. R. Acad. Sci. Paris Sér. I Math.* **312**:11 (1991), 811–815. MR Zbl
- [Ye 1994] R. Ye, “Global existence and convergence of Yamabe flow”, *J. Differential Geom.* **39**:1 (1994), 35–50. MR Zbl

Received May 22, 2017. Revised April 26, 2018.

INAS AMACHA  
UNIVERSITÉ OF BREST  
BREST  
FRANCE

inas.amacha@univ-brest.fr

*Current address:*

LABORATOIRE DE MATHÉMATIQUES, UMR 6205 CNRS  
UNIVERSITÉ DE BRETAGNE OCCIDENTALE  
29238 BREST CEDEX 3  
FRANCE

RACHID REGBAOUI  
LABORATOIRE DE MATHÉMATIQUES, UMR 6205 CNRS  
UNIVERSITÉ DE BRETAGNE OCCIDENTALE  
29238 BREST CEDEX 3  
FRANCE

rachid.regbaoui@univ-brest.fr



# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Matthias Aschenbrenner  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[matthias@math.ucla.edu](mailto:matthias@math.ucla.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Wee Teck Gan  
Mathematics Department  
National University of Singapore  
Singapore 119076  
[matgwt@nus.edu.sg](mailto:matgwt@nus.edu.sg)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 297 No. 2 December 2018

---

Yamabe flow with prescribed scalar curvature INAS AMACHA and RACHID REGBAOUI	257
Rigidity of proper holomorphic mappings between generalized Fock–Bargmann–Hartogs domains ENCHAO BI and ZHENHAN TU	277
Galoisian methods for testing irreducibility of order two nonlinear differential equations GUY CASALE and JACQUES-ARTHUR WEIL	299
Combinatorial classification of quantum lens spaces PETER LUNDING JENSEN, FREDERIK RAVN KLAUSEN and PETER M. R. RASMUSSEN	339
On generic quadratic forms NIKITA A. KARPENKO	367
Rankin–Cohen brackets and identities among eigenforms ARVIND KUMAR and JABAN MEHER	381
Duality for differential operators of Lie–Rinehart algebras THIERRY LAMBRE and PATRICK LE MEUR	405
Nondegeneracy of the Gauss curvature equation with negative conic singularity JUNCHENG WEI and LEI ZHANG	455
Extensions of almost faithful prime ideals in virtually nilpotent mod- $p$ Iwasawa algebras WILLIAM WOODS	477



0030-8730(201812)297:2;1-P