

*Pacific
Journal of
Mathematics*

**COMBINATORIAL CLASSIFICATION
OF QUANTUM LENS SPACES**

PETER LUNDING JENSEN, FREDERIK RAVN KLAUSEN
AND PETER M. R. RASMUSSEN

COMBINATORIAL CLASSIFICATION OF QUANTUM LENS SPACES

PETER LUNDING JENSEN, FREDERIK RAVN KLAUSEN
AND PETER M. R. RASMUSSEN

We answer the question of how large the dimension of a quantum lens space must be, compared to the primary parameter r , for the isomorphism class to depend on the secondary parameters. Since classification results in C^* -algebra theory reduce this question to one concerning a certain kind of SL-equivalence of integer matrices of a special form, our approach is entirely combinatorial and based on the counting of certain paths in the graphs shown by Hong and Szymański to describe the quantum lens spaces.

1. Introduction

In a seminal paper by Hong and Szymański [2003], an important class of *quantum lens spaces* $C(L_q(r; (m_1, \dots, m_n)))$ was given a description as C^* -algebras arising from certain graphs—or their adjacency matrices—in the vein of Cuntz and Krieger [1980]. These graphs can be read off directly from the data $(r; (m_1, \dots, m_n))$ determining the quantum lens space, where $r > 2$ are integers and m_i are units of $\mathbb{Z}/r\mathbb{Z}$. Using this characterisation, it is easy to see that $C(L_q(r; (m_1, \dots, m_n)))$ can only be isomorphic to $C(L_q(r'; (m'_1, \dots, m'_{n'})))$ when $r = r'$ and $n = n'$, and this raises the important question of to what extent the choice of the units can influence the C^* -algebras.

To answer such questions, one appeals naturally to the classification theory for C^* -algebras by K -theory, as indeed a large class of Cuntz–Krieger algebras were classified by Restorff [2006]. Unfortunately, the quantum lens spaces fall outside this class, and indeed, outside any class considered at the time [Hong and Szymański 2003] was written. Thus, apart from noting that the m_i can obviously not influence the C^* -algebras when $n \leq 3$, Hong and Szymański left the question open.

Quantum lens spaces are still a subject of interest, however, see for instance Arici, Brain, and Landi [Arici et al. 2015] and Brzeziński and Szymański [2018], and using recent classification results obtained for Cuntz–Krieger algebras with

MSC2010: primary 46L35; secondary 05C30.

Keywords: quantum lens space, C^* -algebra, SL equivalence.

uncountably many ideals, Eilers, Restorff, Ruiz, and Sørensen in [Eilers et al. 2018] managed to reduce this question to elementary matrix algebra and to prove that when $n = 4$ there are precisely two different $C(L_q(r; (m_1, \dots, m_n)))$ when r is a multiple of 3, and only one when r is not.

Søren Eilers conducted computer experiments for other r and n which suggested that the quantum lens spaces are unique when $n < s$ for s the smallest even number strictly larger than the smallest divisor of r which is not 2, and that at least two choices of m_i give different C^* -algebras when $n \geq s$. It is the aim of the paper at hand to provide the combinatorial insight needed to prove that this in fact is the case, and to study the number of different C^* -algebras that can be obtained by varying the m_i .

We will not work directly on questions of isomorphism of the C^* -algebras, and hence, *no prior knowledge on C^* -algebras or their classification theory is required*. Instead we study the equivalent notion of SL equivalence of the graphs associated to the given data. Indeed, a result of [Eilers et al. 2018] states that the following are equivalent:

- $C(L_q(r; (m_1, \dots, m_n))) \otimes \mathbb{K} \simeq C(L_q(r; (m'_1, \dots, m'_n))) \otimes \mathbb{K}$.
- There exist integer matrices U, V both of the form

$$\begin{pmatrix} 1 & * & * & \cdots & * \\ & 1 & * & & \\ & & \ddots & \ddots & \vdots \\ & & & 1 & * \\ & & & & 1 \end{pmatrix}$$

so that $U(A_{(r; (m_1, \dots, m_n))} - I) = (A_{(r; (m'_1, \dots, m'_n))} - I)V$.

The exact notation and definitions will be given in Section 2 together with the rudimentary results needed for our classification. Section 3 handles the most general case, basically establishing the influence of the odd prime divisors of the parameter r on the number of C^* -algebras emerging by varying the m_i . A lower bound on the number of such C^* -algebras is found and for $4 \nmid r$ the exact s such that the C^* -algebra is unique for $n < s$ is determined. The special case of finding s when $4 \mid r$ is then dealt with in Section 4.

The main result of the paper is Theorem 5.1 which combines the results of Sections 3 and 4 to find for every $r > 2$ the s such that the C^* -algebra is unique for every $n < s$. The other major achievement is Theorem 3.9 which bounds the number of different quantum lens spaces arising for some $r > 2$ and $n \in \mathbb{N}$. Based on computer experiments, we conjecture that this bound is in fact an equality when $4 \nmid r$ (Conjecture 5.3).

2. Preliminaries

We dedicate this section to setting the stage. We establish notation, definitions, and find initial results that will assist in showing the later sections' classification results.

Number theoretical notation.

Definition 2.1. We let Z_n denote the multiplicative group of integers modulo n . That is $Z_n = (\mathbb{Z}/n\mathbb{Z})^*$.

Notation 2.2. We write $p^k \parallel n$ if $p^k \mid n$ and $p^{k+1} \nmid n$, i.e., k is the greatest power of p dividing n .

Notation 2.3. To ease notation, we write the reduction of an integer a calculated modulo r as $[a]_r$, i.e., we always have $0 \leq [a]_r \leq r - 1$.

The graph. This section will introduce a definition of the graph $M_{(r;(m_1,\dots,m_n))}$, arising from the quantum lens space $C(L_q(r; (m_1, \dots, m_n)))$ as defined in [Hong and Szymański 2003]. Further, we introduce another graph $N_{(r;(m_1,\dots,m_n))}$, which is easier to work with in the combinatorial setting, but has similar properties in a sense that will be made clear.

Definition 2.4. Let $r > 2$ and $\bar{m} = (m_1, \dots, m_n) \in (\mathbb{Z}_r)^n$ for some $n \in \mathbb{N}$. Then we define a directed graph $M_{(r;\bar{m})}$ in the following way:

- For every pair s, t with $1 \leq s \leq n$ and $0 \leq t < r$ there is a vertex $g_{s,t}$.
- There is a directed edge from g_{s_1,t_1} to g_{s_2,t_2} if and only if $s_1 \leq s_2$ and $t_2 = [t_1 + m_{s_1}]_r$.

For every $s \in \mathbb{N}$ we will call the subgraph consisting of the vertices $\{g_{s,x} \mid 0 \leq x < r\}$ the s -th subgraph of $M_{(r;\bar{m})}$, and we will call a vertex of the form $g_{s,c}$ a c -vertex.

An example of the graph $M_{(5;(1,2,1))}$ is sketched in Figure 1.




Figure 1. Example of $M_{(r;\bar{m})}$ with $n = 3$, $r = 5$ and $m = (1, 2, 1)$. The bold 3 in a darker circle denotes the vertex $g_{2,3}$.




Figure 2. Example of $N_{(r;\bar{m})}$ where $n = 3$, $r = 5$, and $m = (1, 2, 1)$.

Definition 2.5. Let $r > 2$ and $\bar{m} = (m_1, \dots, m_n) \in (\mathbb{Z}_r)^n$ for some $n \in \mathbb{N}$. Then we define a directed graph $N_{(r;\bar{m})}$ in the following way:

- For every pair s, t with $1 \leq s \leq n$ and $0 \leq t < r$ there is a vertex $c_{s,t}$.
- There is a directed edge from c_{s_1,t_1} to c_{s_2,t_2} in the following two cases:
 - ◊ $s_1 + 1 = s_2$ and $t_2 = t_1$.
 - ◊ $s_1 = s_2$ and $t_2 = [t_1 + m_{s_1}]_r$.

For every s we will call the subgraph consisting of the vertices $\{c_{s,x} \mid 0 \leq x < r\}$ the s -th subgraph of $N_{(r;\bar{m})}$, and we will call a vertex of the form $c_{s,t}$ a t -vertex.

Here is the graph we would rather look at. Instead of having edges from a subgraph to all the subgraphs after it, it only has edges to the one just after it. This edge will always go from $c_{s,t}$ to $c_{s+1,t}$. We show an example of the graph on Figure 2.

Definition 2.6. Let $r > 2$ and $\bar{m} = (m_1, \dots, m_n) \in (\mathbb{Z}_r)^n$ for some $n \in \mathbb{N}$. Then we let $A_{(r;\bar{m})}$ be the matrix such that $A_{(r;\bar{m})}(i, j)$ is the number of directed paths in $M_{(r;\bar{m})}$ from the 0-vertex of the i -th subgraph to the 0-vertex of the j -th subgraph that does not pass through the 0-vertex of any other subgraph. We call a path that satisfies these criteria *legal*.

Definition 2.7. Let $r > 2$ and $\bar{m} = (m_1, \dots, m_n) \in (\mathbb{Z}_r)^n$ for some $n \in \mathbb{N}$. Then we let $B_{(r;\bar{m})}$ be the matrix such that $B_{(r;\bar{m})}(i, j)$ is the number of directed paths on $N_{(r;\bar{m})}$ from the 0-vertex of the i -th subgraph to the 0-vertex of the j -th subgraph which do not exclusively visit 0-vertices and which do not visit the 0-vertex of any other subgraph except if all the following vertices of the path are 0-vertices. We will call a path that satisfies these criteria *legal*.

We introduce this new graph definition $N_{(r;\bar{m})}$ because it is easier to work with than $M_{(r;\bar{m})}$. Note we always calculate indices in subgraphs modulo r .

Lemma 2.8. Let $r > 2$ and $\bar{m} \in (\mathbb{Z}_r)^n$ be given. Then $A_{(r;\bar{m})} = B_{(r;\bar{m})}$.

Proof. There is a bijection between the edges of $M_{(r;\bar{m})}$ and paths of $N_{(r;\bar{m})}$ as follows. The edge $g_{s_1,t_1} \rightarrow g_{s_2,t_1+m_{s_1}}$ of $M_{(r;\bar{m})}$ corresponds to the path

$$c_{s_1,t_1} \rightarrow c_{s_1,t_1+m_{s_1}} \rightarrow c_{s_1+1,t_1+m_{s_1}} \rightarrow \cdots \rightarrow c_{s_2,t_1+m_{s_1}}$$

on $N_{(r;\bar{m})}$. That this is a bijection follows immediately from the fact that the edge and path are both uniquely determined by s_1 , s_2 , and t_1 .

Now, we need to establish a bijection between the legal paths on $M_{(r;\bar{m})}$ and the legal paths on $N_{(r;\bar{m})}$. This happens naturally by translating any edge in a legal path on $M_{(r;\bar{m})}$ into a subpath of the form above of a legal path on $N_{(r;\bar{m})}$. That this map has an inverse follows easily since any legal path in $N_{(r;\bar{m})}$ consists of subpaths of the above form where a new subpath starts whenever we stay in the same subgraph. Further, we have that the constraint of Definition 2.6 translates into the constraint of Definition 2.7: an edge from the t -th subgraph to the 0-vertex of the n -th subgraph in Definition 2.6 corresponds to going to the 0-vertex in the t -th subgraph and then visiting 0-vertices exclusively until reaching the 0-vertex of the n -th subgraph in Definition 2.7. □

Equivalence classes. The overall aim of the article is to classify the quantum lens spaces, which is a problem that Theorem 7.1 of Section 7.2 of [Eilers et al. 2018] reduces to a question of SL equivalence, hence elementary matrix algebra.

Theorem 2.9 (Eilers, Restorff, Ruiz, and Sørensen). *Let $r > 2$ and $\bar{m}, \bar{m}' \in (\mathbb{Z}_r)^n$ be given. The following are equivalent:*

- $C(L_q(r; (m_1, \dots, m_n))) \otimes \mathbb{K} \simeq C(L_q(r; (m'_1, \dots, m'_n))) \otimes \mathbb{K}$.
- *There exist matrices U, V both of the form*

$$\begin{pmatrix} 1 & * & * & \cdots & * \\ & 1 & * & & \\ & & \ddots & \ddots & \vdots \\ & & & 1 & * \\ & & & & 1 \end{pmatrix}$$

so that $U(A_{(r;(m_1,\dots,m_n))} - I) = (A_{(r;(m'_1,\dots,m'_n))} - I)V$.

Thus, determining whether two quantum lens spaces, $C(L_q(r; (m'_1, \dots, m'_n)))$ and $C(L_q(r; (m_1, \dots, m_n)))$, are isomorphic comes down to whether or not the matrices $A_{(r;\bar{m})}$ and $A_{(r;\bar{m}')}$ (or $B_{(r;\bar{m})}$ and $B_{(r;\bar{m}')}$ by Lemma 2.8) are equivalent with respect to the equivalence relation, \sim , defined below.

Definition 2.10. We will say that two matrices C and D are upper triangular equivalent, written $C \cong D$, if there exist upper triangular matrices, X, Y , with 1 in every entry of the diagonal such that $XC = DY$.

Equivalently, the matrices C and D are upper triangular equivalent, if there is a series of pivots transforming C into D with the restrictions that

- (1) a multiple of row k can only be added to row l if $k > l$,
- (2) a multiple of column k can only be added to column l if $k < l$.

Note that this is clearly an equivalence relation since such upper triangular matrices are invertible.

Definition 2.11. We say that two matrices, A and C , are equivalent, if

$$A - I \cong C - I.$$

In that case we write $A \sim C$.

In particular, we are interested in efficiently deciding the number of equivalence classes given n and $r > 2$ and deciding whether or not two graphs belong to the same equivalence class.

Definition 2.12. Let $r > 2$ and $n \in \mathbb{N}$ be given. Then we define

$$S_{r,n} = \{A_{(r;\bar{m})} \mid \bar{m} \in (Z_r)^n\} = \{B_{(r;\bar{m})} \mid \bar{m} \in (Z_r)^n\}$$

as the set of all matrices produced by vectors of length n with parameter r .

Definition 2.13. Let $r > 2$ and $n \in \mathbb{N}$ be given. Then $\varphi_r(n)$ denotes the number of elements of $S_{r,n}/\sim$ and $\tilde{\varphi}(r)$ denotes the least n such that $\varphi_r(n) > 1$.

Thus, our goal in this paper is to find a bound for φ_r given r and to express $\tilde{\varphi}$ in closed form.

Invariants. In this section we establish some invariants and properties in relation to changes to the vector \bar{m} in $N_{(r;\bar{m})}$.

Lemma 2.14. *The matrix $B_{(r;\bar{m})}$ does not depend on the choice of m_1 and m_n .*

Proof. If $n = 1$ this is obvious, so assume $n > 1$. Consider legal paths in $N_{(r;(m_1, \dots, m_n))}$ from the 0-vertex of the first subgraph to the 0-vertex of the j -th subgraph for $j > 1$. No matter what m_1 is there is exactly one way to reach any of the vertices of the second subgraph from the 0-vertex of the first subgraph. Thus, the number of such directed paths is independent of m_1 and the first part follows.

Now, consider the last subgraph. Once it is reached, there is exactly one way to reach the 0-vertex, once it is reached, so this does not depend on m_n . \square

Lemma 2.15. *Let $r > 2$, $\bar{m} \in (Z_r)^n$, and $b \in Z_r$. Then $B_{(r;\bar{m})} = B_{(r;b\bar{m})}$.*

Proof. We will show that there is a bijection between the legal paths of $B_{(r;\bar{m})}$ and $B_{(r;b\bar{m})}$ as follows. Let γ be a legal path

$$c_{s_1,0} = c_{s_1,t_1} \rightarrow c_{s_2,t_2} \rightarrow \cdots \rightarrow c_{s_q,t_q} = c_{s_q,0}$$

on $N_{(r;\bar{m})}$. Our bijection sends the legal γ to the path ω on $N_{(r;(b\cdot\bar{m}))}$ given by

$$c_{s_1,0} = c_{s_1,[b\cdot t_1]_r} \rightarrow c_{s_2,[b\cdot t_2]_r} \rightarrow \dots \rightarrow c_{s_q,[b\cdot t_q]_r} = c_{s_q,0}.$$

That the map is injective follows since multiplication by $b \in Z_r$ is an injection $Z_r \rightarrow Z_r$. Further, it is easy to see that all legal paths on $N_{(r;\bar{m})}$ will be mapped to legal paths on $N_{(r;(b\cdot\bar{m}))}$ since multiplication by b does not change the positions of the 0-vertices in a path. Thus, there is an injection from the legal paths on $N_{(r;\bar{m})}$ to the legal paths on $N_{(r;(b\cdot\bar{m}))}$ and by the same argument there must be an injection from the legal paths on $N_{(r;(b\cdot\bar{m}))}$ to the legal paths on $N_{(r;\bar{m})}$. It follows that said map is a bijection and we are done. \square

Corollary 2.16. *Let $\bar{m} = (m_1, m_2, \dots, m_{n-1}, m_n) \in (Z_r)^n$. Then there exists an $\bar{m}' \in (Z_r)^n$ with 1 in the first, last and k -th index, i.e.,*

$$\bar{m}' = (1, m'_2, \dots, m'_{k-1}, 1, m'_{k+1}, \dots, m'_{n-1}, 1),$$

such that $B_{(r;\bar{m})} = B_{(r;\bar{m}')}.$

Proof. Take b to be the inverse in Z_r of m_k in Lemma 2.15. Then

$$B_{(r;\bar{m})} = B_{(r;m_k^{-1}\cdot\bar{m})} = B_{(r;\bar{m}')}.$$

where the last equality follows from Lemma 2.14. \square

Entry specific properties and formulae. To proceed with any further results we need some combinatorial formulae and properties to be in place.

Theorem 2.17. *Let $\bar{1} = (1, \dots, 1)$. Then $B_{(r;\bar{1})}\langle i, j \rangle = \binom{r-1+(j-i)}{j-i}$.*

Proof. Since every directed edge in $N_{(r;\bar{1})}$ either goes from $c_{s,t}$ to $c_{s+1,t}$ or from $c_{s,t}$ to $c_{s,t+1}$, we can characterise any directed path from the 0-vertex of the i -th subgraph to the 0-vertex of the j -th subgraph satisfying Definition 2.7 by a $j - i + 1$ -tuple (a_0, \dots, a_{j-i}) , such that a_s is the number of edges of the form $c_{s,t} \rightarrow c_{s,t+1}$ that occur in the path. A necessary and sufficient condition for such a tuple to characterise a directed path of the desired form is that $\sum_{l=0}^{j-i} a_l = r$, $a_s \geq 0$ for all $s > 0$, and $a_0 > 0$.

Thus, $B_{(r;\bar{1})}\langle i, j \rangle$ is equal to the number of ways that r can be written as the sum of $j - i + 1$ nonnegative integers, where the first one has to be at least 1. This is equivalent to the number of ways to write $r - 1$ as the sum of $j - i + 1$ nonnegative integers. The latter being a known combinatorial problem, we get

$$B_{(r;\bar{1})}\langle i, j \rangle = \binom{r-1+(j-i)}{j-i}. \quad \square$$

Corollary 2.18. *Let $r > 2$ and $\bar{m} \in (Z_r)^n$. Then $B_{(r;\bar{m})}\langle i, i \rangle = 1$, $B_{(r;\bar{m})}\langle i, i+1 \rangle = r$, and $B_{(r;\bar{m})}\langle i, i+2 \rangle = \frac{r(r+1)}{2}$ for all i .*

Proof. When we consider only $B_{(r;\bar{m})}(i, i)$, $B_{(r;\bar{m})}(i, i + 2)$, and $B_{(r;\bar{m})}(i, i + 2)$, their values depend solely on the vector (m_i, m_{i+1}, m_{i+2}) , so we can assume by Corollary 2.16 that

$$m_i = m_{i+1} = m_{i+2} = 1.$$

The conclusion then follows trivially from Theorem 2.17 □

From the corollary we immediately obtain the following result, which also appeared in [Eilers et al. 2018] and we will note for future use:

Corollary 2.19. *Let $r > 2$. Then $\tilde{\varphi}(r) \geq 4$.*

Proof. By Corollary 2.18 we have that for $n \leq 3$ the matrices $B_{(m;\bar{r})}$ do not depend on \bar{m} . Thus, they are all equal when \bar{m} varies and we can only have one equivalence class. □

In fact, Eilers et al. [2018] established that $\tilde{\varphi}(r) = 4$ if and only if $3 \mid r$. As stated earlier, we shall see a general closed expression for $\tilde{\varphi}$ in a later section.

Equivalence of matrices. To show equivalence of matrices we need to do some manipulations with matrices that might be a bit technical. So the following lemma simply establishes the equivalence of two matrices where every entry except for the diagonal is divisible by either r or $\frac{r}{2}$ when r is even.

Lemma 2.20. *Let $r > 2$ be given such that $r = 2^t s$ for some $t \in \{0, 1\}$ and odd $s \in \mathbb{N}$. Suppose that the two $n \times n$ upper triangular integer matrices A, B have 1's in their diagonal, r on the diagonal from $\langle 1, 2 \rangle$ to $\langle n - 1, n \rangle$, and $\frac{r(r+1)}{2}$ on the diagonal from $\langle 1, 3 \rangle$ to $\langle n - 2, n \rangle$. Further, suppose s divides every entry of $A - I$ and $B - I$. Then $A \sim B$.*

Proof. We need to show that we can transform the matrix $A - I$ into $B - I$ by integer row and column operations. If r is odd, every entry of $A - I$ and $B - I$ is divisible by r by assumption, and the matrices are of the form

$$r \begin{pmatrix} 0 & 1 & & & \\ 0 & 0 & 1 & & \\ \vdots & \vdots & & \ddots & \\ 0 & 0 & \dots & 0 & 1 \\ 0 & 0 & \dots & 0 & 0 \end{pmatrix}$$

with integer entries in the upper right corner. All such matrices can easily be transformed into an upper triangular matrix with zeros everywhere except for the diagonal from $\langle 1, 2 \rangle$ to $\langle n - 1, n \rangle$ by row and column operations, so since \sim is an equivalence relation, we get $A \sim B$.

Now, assume that $2 \mid r$, but $4 \nmid r$. Then the matrices $A - I$ and $B - I$ are of the form

$$A - I = \frac{r}{2} \begin{pmatrix} 0 & 2 & r+1 & & & & \\ 0 & 0 & 2 & r+1 & & & \\ \vdots & \vdots & & \ddots & \ddots & & \\ 0 & 0 & \dots & 0 & 2 & r+1 & \\ 0 & 0 & \dots & 0 & 0 & 2 & \\ 0 & 0 & \dots & 0 & 0 & 0 & \end{pmatrix}$$

with integer entries in the upper-right corner. We show that such a matrix can be transformed by row and column operations into the matrix

$$C - I := \frac{r}{2} \begin{pmatrix} 0 & 2 & r+1 & 0 & \dots & \dots & 0 \\ 0 & 0 & 2 & r+1 & 0 & \dots & 0 \\ \vdots & \vdots & & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & 0 & 2 & r+1 & 0 \\ 0 & 0 & \dots & 0 & 0 & 2 & r+1 \\ 0 & 0 & \dots & 0 & 0 & 0 & 2 \\ 0 & 0 & \dots & 0 & 0 & 0 & 0 \end{pmatrix},$$

which by transitivity shows $A \sim B$.

We proceed by induction on n . For $n = 1, 2, 3$ all matrices on the described form will be identical and thus equivalent to C .

Now, assume that for $k < n$ every matrix on the described form is equivalent to C , and consider the $n \times n$ matrix A on that form. By the induction hypothesis and considering A as an $(n - 1) \times (n - 1)$ matrix with an added row and column, we can reduce A by row and column operations to a matrix with diagonals like $C - I$ and zeroes everywhere else except for in the rightmost column.

Using column operations we can now make every entry of that rightmost column (except for the $r - 1$ -entry) even without changing the rest of the matrix. If an entry of the column is odd, we can just subtract $r - 1$ from it by subtracting the appropriate column.

Having made all those entries even they can all be eliminated by subtracting the 2 in the $(n - 1)$ -st row an appropriate amount of times. Then the matrix $C - I$ is achieved, which concludes the proof. \square

Another useful result on when matrices are not equivalent is the following:

Lemma 2.21. *Let A and B be $n \times n$ upper-rectangular matrices with 1 in their diagonal. If every entry of $A - I$ and $B - I$ except for the entry $\langle 1, n \rangle$ is divisible by $k \in \mathbb{N}$ and $(A - I)\langle 1, n \rangle \not\equiv (B - I)\langle 1, n \rangle \pmod{k}$, then $A \not\sim B$.*

Proof. Since every entry of $A - I$ and $B - I$ except the upper right is divisible by k , the upper right entry is invariant modulo k under row and column operations. \square

3. The general case

In general it is very difficult to find an explicit formula for $B_{(r;\bar{m})}\langle i, j \rangle$ given arbitrary r and \bar{m} . However, for the purpose of bounding $\varphi_r(n)$ from below and deciding $\tilde{\varphi}(r)$ in the case where $4 \nmid r$, it turns out to be sufficient to be able to compute $B_{(r;\bar{m})}\langle i, j \rangle$ modulo r .

Thus, this section aims to develop techniques for assessing $B_{(r;\bar{m})}\langle i, j \rangle$ modulo r . The main technical result is Theorem 3.2 from which the exact value of $\tilde{\varphi}(r)$ follows for $4 \nmid r$ and a lower bound on $\varphi_r(n)$, which appears to be an equality when $4 \nmid r$ (see Conjecture 5.3). Throughout the section, we will define $0^0 = 1$ and $0! = 1$ for the sake of simplicity.

We start with the following lemma, which formally captures the technique which will be used multiple times in the proof of Theorem 3.2:

Lemma 3.1. *Suppose D is a finite set, p is a prime, $s, b : D \rightarrow \mathbb{Z}$ are functions, $k, j \in \mathbb{N}$, and $a : \mathbb{N}_0 \times \mathbb{N}_0 \rightarrow \mathbb{Z}$ satisfies $\gcd(a(m, m), p) = 1$ for all $m \leq j$. Define the function*

$$w(l) = \sum_{d \in D} s(d) \sum_{t=0}^l a(t, l) b(d)^t$$

and assume that $p^k \mid w(l)$ for $0 \leq l < j$. Then

$$w(j) \equiv \sum_{d \in D} a(j, j) s(d) b(d)^j \pmod{p^k}.$$

Proof. First, we show by strong induction over t that $p^k \mid \sum_{d \in D} s(d) b(d)^t$ for all $t < j$. For $t = 0$ we have $p^k \mid \sum_{d \in D} s(d) a(0, 0)$ and since $\gcd(p^k, a(0, 0)) = 1$ we get $p^k \mid \sum_{d \in D} s(d)$. Now, assume that $p^k \mid \sum_{d \in D} s(d) b(d)^t$ for all t satisfying $0 \leq t < m$ for some $m < j$. Then,

$$0 \equiv w(m) = \sum_{d \in D} s(d) \sum_{t=0}^m a(t, m) b(d)^t \equiv \sum_{d \in D} a(m, m) s(d) b(d)^m \pmod{p^k},$$

so $p^k \mid \sum_{d \in D} s(d) b(d)^m$ as $\gcd(a(m, m), p^k) = 1$.

Second, the fact that $p^k \mid \sum_{d \in D} s(d) b(d)^t$ for all $t < j$ yields

$$w(j) = \sum_{d \in D} s(d) \sum_{t=0}^j a(t, j) b(d)^t \equiv \sum_{d \in D} a(j, j) s(d) b(d)^j \pmod{p^k}. \quad \square$$

Having proved the lemma we now turn to the main technical theorem of the section from which the remaining results follow naturally.

Theorem 3.2. *Let p be an odd prime, $r > 2$ and $n \leq p + 1$ be given, and $\bar{m} = (m_1, \dots, m_n)$. Suppose that $p^k \mid r$ and $p^k \mid B_{(r;\bar{s})}\langle 1, a \rangle$ for every $\bar{s} \in (Z_r)^{p+1}$ and*

every $a < n$. Then

$$B_{(r;\bar{m})}(1, n) \equiv \binom{r+n-2}{n-1} \prod_{k=2}^{n-1} m_k^{-1} \pmod{p^k}.$$

Proof. For every vector \bar{m} we can reduce the problem to considering a vector \bar{m}' which satisfies $m'_1 = m'_2 = m'_n = 1$ as follows. First, recall that no matter what vector we consider, we can always assume without loss of generality that its first and last entry is 1 since it does not affect any of the sides of the above expression by Lemma 2.14. Second, as in the proof of Corollary 2.16 we can multiply \bar{m} by a m_2^{-1} to get $\bar{m}' = m_2^{-1} \cdot \bar{m}$, which means that $m'_2 = 1$. Then the left hand side will not change since $B_{(r;\bar{m})}(1, n) = B_{(r;\bar{m}')} (1, n)$ by Lemma 2.15 and the right hand side will satisfy

$$\binom{r+n-2}{n-1} \prod_{k=2}^{n-1} m_k^{-1} \equiv \binom{r+n-2}{n-1} \prod_{k=2}^{n-1} b m_k^{-1} \pmod{p^k}$$

since for $n < p + 1$, $p^k \mid \binom{r+n-2}{n-1}$, and for $n = p + 1$, $b^{n-2} \equiv 1 \pmod{p}$ and $p^{k-1} \mid \binom{r+n-2}{n-1}$. Now, assuming $m'_1 = m'_n = 1$ yields the above. Thus, for the remaining proof we will assume that $m_1 = m_2 = m_n = 1$.

Now, let \bar{n}_j denote the vector $(m_1, \dots, m_{n-j}, \mathbf{1}_j)$, where $\mathbf{1}_j$ is the vector of 1's of length j , and note that with the above assumption, $\bar{n}_1 = \bar{m}$ and $\bar{n}_{n-2} = \bar{1}$. Our approach will be to show that for all $1 \leq j < n - 1$ we have

$$(1) \quad B_{(r;\bar{n}_j)}(1, n) \equiv m_{n-j}^{-1} g_j(m_1, \dots, m_{n-j-1}) \pmod{p^k}$$

for some integer function $g_j: (\mathbb{Z}_r)^{n-j-1} \rightarrow \mathbb{Z}$ which is independent of m_{n-j} and where m_{n-j}^{-1} is the inverse of m_{n-j} modulo p^k . Noting that (1) yields $B_{(r;\bar{n}_{j+1})}(1, n) \equiv g(j, m_1, \dots, m_{n-j-1}) \pmod{p^k}$, we get

$$B_{(r;\bar{n}_j)}(1, n) \equiv B_{(r;\bar{n}_{j+1})}(1, n) m_{n-j}^{-1} \pmod{p^k},$$

and applying this together with Theorem 2.17 and $m_2 = 1$ gives us

$$\begin{aligned} B_{(r;\bar{m})}(1, n) &= B_{(r;\bar{n}_1)}(1, n) \equiv B_{(r;\bar{n}_2)}(1, n) m_{n-1}^{-1} \\ &\quad \vdots \\ &\equiv B_{(r;\bar{n}_{n-2})}(1, n) \prod_{k=3}^{n-1} m_k^{-1} = B_{(r;\bar{1})}(1, n) \prod_{k=3}^{n-1} m_k^{-1} \\ &= \binom{r+n-2}{n-1} \prod_{k=2}^{n-1} m_k^{-1} \pmod{p^k}. \end{aligned}$$

Thus, all we need to do is prove that we can indeed write an expression for $B_{(r;\bar{n}_j)}(1, n)$ of the form (1).

To do so, fix a j with $1 \leq j < n - 1$ and consider the graph $N_{(r;\overline{n_j})}$. We may write

$$(2) \quad \mathbb{B}_{(r;\overline{n_j})}\langle 1, n \rangle = \sum_{q=0}^{r-1} L_j(q) S_j(q),$$

where $S_j(q)$ denotes the number of paths on $N_{(r;\overline{n_j})}$ from $c_{1,0}$ to $c_{n-j,q}$ that are subpaths of a legal path from $c_{1,0}$ to $c_{n,0}$ and that do not traverse any edges in the $(n-j)$ -th subgraph, and similarly $L_j(q)$ is the number of paths on $N_{(r;\overline{n_j})}$ from $c_{n-j,q}$ to $c_{n,0}$ that are subpaths of a legal path from $c_{1,0}$ to $c_{n,0}$.

We start our analysis by finding a formula for $L_j(q)$. First, we consider the $(n-j+1)$ -th subgraph of $N_{(r;\overline{n_j})}$ and count the number of paths from $c_{n-j+1,i}$ to $c_{n,0}$ on $N_{(r;\overline{n_j})}$ that are subpaths of a legal path from $c_{1,0}$ to $c_{n,0}$ for each $0 \leq i < r$. As in the proof of Theorem 2.17 one can see choosing such a path as choosing a partition of $[r-i]_r$ into a sum of $j-1$ nonnegative integers since $m_{n-j+1} = m_{n-j+2} = \dots = m_n = 1$. Thus, the number of such paths equals $\binom{[r-i]_r + j - 1}{j-1}$.

Second, there are three cases to consider. When $i, q > 0$ there is exactly one path from $c_{n-j,q}$ to $c_{n-j+1,i}$ not traversing any edges in the $(n-j+1)$ -th subgraph that is a subpath of a legal path from $c_{1,0}$ to $c_{n,0}$ if and only if $[m_{n-j}^{-1}q]_r \leq [m_{n-j}^{-1}i]_r$. Otherwise there are none. This is clear since such a path would be of the form

$$c_{n-j,q} \rightarrow c_{n-j,[q+m_{n-j}]_r} \rightarrow \dots \rightarrow c_{n-j,i} \rightarrow c_{n-j+1,i}$$

and zero is not a member of $\{q, q+m_{n-j}, \dots, i\}$ if and only if $[m_{n-j}^{-1}q]_r \leq [m_{n-j}^{-1}i]_r$. For $i = 0$ there is exactly one such subpath for every q and for $q = 0$ there is exactly one such subpath if and only if $i = 0$. Thus, for $q > 0$,

$$\begin{aligned} L_j(q) &= \sum_{i=0}^{r-1} (1_{\{[m_{n-j}^{-1}q]_r \leq [m_{n-j}^{-1}i]_r\}} + 1_{\{i=0\}}) \binom{[r-i]_r + j - 1}{j-1} \\ &= \sum_{i=[m_{n-j}^{-1}q]_r}^r \binom{[r-i]_r + j - 1}{j-1}, \end{aligned}$$

where 1_{boolean} is an indicator function assuming the value 1 if true and 0 otherwise, and where we changed $i = 0$ terms into $i = r$ terms. Introducing the new variable $\sigma = r - i$ we rewrite the sum as

$$(3) \quad L_j(q) = \sum_{\sigma=0}^{[m_{n-j}^{-1}q]_r} \binom{[\sigma m_{n-j}]_r + j - 1}{j-1}.$$

Since evidently $L_j(0) = 1$, this formula holds even for $q = 0$ and thus for all $0 \leq q < r$.

For $j = 1$, (3) yields $L_j(q) = [-m_{n-1}^{-1}q]_r$ and inserting this into (2) yields

$$B_{(r; \bar{n}_1)} = \sum_{q=0}^{r-1} [-m_{n-1}^{-1}q]_r S_j(q) \equiv -m_{n-1}^{-1} \sum_{q=0}^{r-1} S_j(q)q \pmod{p^k}.$$

Since $S_j(q)q$ only depends on m_1, m_2, \dots, m_{n-2} , it follows that we can write $B_{(r; \bar{n}_1)}$ in the form (1).

So let us consider the case when $j > 1$. Inserting the expression (3) into (2) and substituting $d = r - q$ and noting that the $d = 0$ is equal to the $d = r$ term yields

$$(j - 1)! B_{(r; \bar{m})} \langle 1, n \rangle = \sum_{d=0}^{r-1} \sum_{\sigma=0}^{[m_{n-j}^{-1}d]_r} S_j([r - d]_r) \prod_{i=1}^{j-1} ([\sigma m_{n-j}]_r + i).$$

Expanding the product and introducing $s(d) = S_j([r - d]_r)$ we get the sum

$$(j - 1)! B_{(r; \bar{n}_j)} \langle 1, n \rangle = \sum_{d=0}^{r-1} \sum_{\sigma=0}^{[m_{n-j}^{-1}d]_r} \sum_{t=0}^{j-1} a(t, j - 1) ([\sigma m_{n-j}]_r)^t s(d)$$

for an integer function $a: \mathbb{N}_0 \times \mathbb{N}_0 \rightarrow \mathbb{Z}$, where $a(j - 1, j - 1) = 1$. Now, note that by the same reasoning we must also have for every $0 \leq l < j - 1$ that

$$\begin{aligned} w(l) &:= l! B_{(r; \bar{n}_j)} \langle 1, n - j + l + 1 \rangle \\ &= \sum_{d=0}^{r-1} \sum_{\sigma=0}^{[m_{n-j}^{-1}d]_r} s(d) \sum_{t=0}^l a(t, l) ([\sigma m_{n-j}]_r)^t, \end{aligned}$$

where $a(l, l) = 1$. By assumption, p^k divides $B_{(r; \bar{n}_j)} \langle 1, n - j + l + 1 \rangle$ for every $l < j - 1$. Hence, $p^k \mid w(l)$ for $0 \leq l < j - 1$. Applying Lemma 3.1 with the functions $s, a, b(\sigma) := [\sigma m_{n-j}]_r$, prime p , and exponent t , we get

$$\begin{aligned} (4) \quad (j - 1)! B_{(r; \bar{n}_j)} \langle 1, n \rangle &= w(j - 1) \\ &\equiv \sum_{d=0}^{r-1} \sum_{\sigma=0}^{[m_{n-j}^{-1}d]_r} a(j - 1, j - 1) s(d) ([\sigma m_{n-j}]_r)^{j-1} \\ &\equiv m_{n-j}^{j-1} \sum_{d=0}^{r-1} \sum_{\sigma=0}^{[m_{n-j}^{-1}d]_r} s(d) \sigma^{j-1} \pmod{p^k}. \end{aligned}$$

By Faulhaber’s formula ([Graham et al. 1989, Chapter 6.5]) in the convention $B_1 = \frac{1}{2}$, we can write

$$\sum_{\sigma=0}^{[m_{n-j}^{-1}d]_r} \sigma^{j-1} = \frac{1}{j} \sum_{t=0}^{j-1} \binom{j}{j - t - 1} B_{j-t-1} ([m_{n-j}^{-1}d]_r)^{t+1},$$

where B_n is the n -th Bernoulli number. Inserting this into (4), noting that $p^k \mid r$ and multiplying both sides by j , we find that

$$j! B_{(r; \overline{n_j})} \langle 1, n \rangle \equiv m_{n-j}^{j-2} \sum_{d=0}^{r-1} s(d) d \sum_{t=0}^{j-1} \binom{j}{j-t-1} B_{j-t-1} (m_{n-j}^{-1} d)^t \pmod{p^k}.$$

As it is a well-known fact that $j! B_l$, $l < j$, is an integer, we multiply by $j! (m_{n-j}^{j-2})^{-1}$ to ensure that each factor of each term is an integer

$$(m_{n-j}^{j-2})^{-1} j!^2 B_{(r; \overline{n_j})} \langle 1, n \rangle \equiv \sum_{d=0}^{r-1} s(d) d \sum_{t=0}^{j-1} \binom{j}{j-t-1} j! B_{j-t-1} (m_{n-j}^{-1} d)^t \pmod{p^k}.$$

To apply Lemma 3.1 again we write

$$\begin{aligned} \tilde{s}(d) &:= s(d) d, \\ \tilde{a}(t, l) &:= \binom{l+1}{l-t} (l+1)! B_{l-t}, \\ \tilde{b}(d) &:= [m_{n-j}^{-1} d]_r, \end{aligned}$$

and considering the vectors $\overline{v_{l+1}} = (m_1, m_2, \dots, m_{n-j}, \mathbf{1}_{l+1})$ one finds:

$$\begin{aligned} \tilde{w}(l) &:= \sum_{d=0}^{r-1} \tilde{s}(d) \sum_{t=0}^l \tilde{a}(t, l) \tilde{b}(d)^t \\ &\equiv (m_{n-j}^{l-1})^{-1} (l+1)!^2 B_{(r; \overline{v_{l+1}})} \langle 1, n-j+l+1 \rangle \pmod{p^k}. \\ &\equiv (m_{n-j}^{l-1})^{-1} (l+1)!^2 B_{(r; \overline{n_j})} \langle 1, n-j+l+1 \rangle \pmod{p^k}. \end{aligned}$$

Now,

$$\tilde{a}(l, l) = \binom{l+1}{0} (l+1)! B_0 = (l+1)!$$

so for $0 \leq l < j-1 < n-1 \leq p$ we have $\gcd(\tilde{a}(l, l), p) = 1$. Further, by assumption, $p^k \mid B_{(r; \overline{n_j})} \langle 1, n-j+l+1 \rangle$ for $0 \leq l < j-1$, so $p^k \mid \tilde{w}(l)$ for $0 \leq l < j-1$. Thus, Lemma 3.1 yields

$$\begin{aligned} (m_{n-j}^{j-2})^{-1} j!^2 B_{(r; \overline{n_j})} \langle 1, n \rangle &= \tilde{w}(j-1) \\ &\equiv \sum_{q=0}^{r-1} s(d) d \tilde{a}(j-1, j-1) (m_{n-j}^{-1} d)^{j-1} \\ &\equiv m_{n-j}^{1-j} \sum_{q=0}^{r-1} j! s(d) d^j \pmod{p^k}. \end{aligned}$$

This means that

$$B_{(r;\bar{n}_j)}\langle 1, n \rangle \equiv m_{n-j}^{-1} \sum_{q=0}^{r-1} j!^{-1} s(d) d^j \pmod{p^k},$$

where we note that $j!^{-1}$ is well defined because $j < n - 1 \leq p$ so $\gcd(j!, p) = 1$. Since $s(d)$ only depends on m_1, \dots, m_{n-j-1} , it is clear that we can find g_j satisfying (1) and we are done. \square

Having proved the above theorem we can apply it to find $\tilde{\varphi}(r)$ whenever $4 \nmid r$. We first use the theorem to prove the following lemma, which will give the first half of the proof.

Lemma 3.3. *Let $r > 2$, p be an odd prime, and $p^k \parallel r$ for some $k \in \mathbb{N}$. For every vector \bar{m} with entries in Z_r and every pair a, b satisfying $0 < b - a < p$, we have*

$$p^k \mid B_{(r;\bar{m})}\langle a, b \rangle.$$

Proof. We proceed by induction on the difference $n = b - a$. When $n = 1$ we have $p^k \mid B_{(r;\bar{m})}\langle a, b \rangle = r$. Now, suppose that $p^k \mid B_{(r;\bar{m})}\langle a', b' \rangle$ for every a', b' satisfying $0 < b' - a' < n$ for some n with $1 < n < p$ and let $b - a = n$. Then we can apply Theorem 3.2 with the indices $\langle 1, n \rangle$ shifted to $\langle a, b \rangle$ to get

$$\begin{aligned} B_{(r;\bar{m})}\langle a, b \rangle &\equiv \binom{r-1+(b-a)}{b-a} \prod_{k=a+1}^{b-1} m_k^{-1} \\ &\equiv \frac{r \cdots (r-1+(b-a))}{(b-a)!} \prod_{k=a+1}^{b-1} m_k^{-1} \\ &\equiv 0 \pmod{p^k}, \end{aligned}$$

where the last equivalence follows since p^k divides

$$\frac{r \cdots (r-1+(b-a))}{(b-a)!}$$

because $b - a < p$ and r divides the numerator. \square

Now, using the previous lemma and Theorem 3.2 we obtain an upper bound on $\tilde{\varphi}$ simply by pointing to two graphs that are not equivalent. In Theorem 3.9 below we will establish a lower bound for the number of equivalence classes from which the result will follow. But for clarity we now give a short independent proof.

Theorem 3.4. *Let $r > 2$ be given and let p be the smallest odd prime dividing r . Then $\tilde{\varphi}(r) \leq p + 1$.*

Proof. Let k be such that $p^k \parallel r$, set

$$\bar{a} = (\mathbf{1}_{p+1}) \text{ and } \bar{b} = (1, -1, \mathbf{1}_{p-1}),$$

and consider the matrices $A = B_{(r;\bar{a})}$, $B = B_{(r;\bar{b})}$. Then by Lemma 3.3 we have $p^k \mid A\langle a, b \rangle$ and $p^k \mid B\langle a, b \rangle$ for $a < b$ and $\langle a, b \rangle \neq \langle 1, p + 1 \rangle$. Using Theorem 3.2 twice and noting that $(r + 1) \cdots (r + p - 1) \equiv (p - 1)! \pmod{p^k}$, we get

$$\begin{aligned} A\langle 1, p + 1 \rangle &= \binom{r + p - 1}{p} \prod_{k=2}^p a_k^{-1} \\ &= \frac{r}{p} \pmod{p^k}, \end{aligned}$$

and

$$\begin{aligned} B\langle 1, p + 1 \rangle &\equiv \binom{r + p - 1}{p} \prod_{k=2}^p b_k^{-1} \\ &\equiv -\frac{r}{p} \pmod{p^k}, \end{aligned}$$

since $b_2 = -1$. It follows that p^k divides every entry of $A - I$ and $B - I$ except for the entry $\langle 1, p + 1 \rangle$. Applying Lemma 2.21 we get $B_{(r;\bar{a})} \not\sim B_{(r;\bar{b})}$ implying $\varphi_r(p + 1) > 1$ and the conclusion follows. \square

Now, using Theorem 3.4, we determine $\tilde{\varphi}(r)$ whenever $4 \nmid r$.

Theorem 3.5. *Let $r > 2$ be given such that $4 \nmid r$ and let p be the smallest odd prime dividing r . Then $\tilde{\varphi}(r) = p + 1$.*

Proof. It follows from Lemmas 2.20 and 3.3 that for every $n \leq p$ and every $\bar{m} \in (Z_r)^n$ we have $B_{(r;\bar{m})} \sim B_{(r;\bar{1})}$, so $\varphi_r(n) = 1$ for $n \leq p$. Thus, $\varphi(r) > p$. The conclusion now follows from Theorem 3.4. \square

The remainder of this section deals with the number of equivalence classes, $\varphi_r(n)$.

Notation 3.6. Let $A = (a_{ij})$ be a matrix. Then we denote by $A[c, d]$ the partial square matrix

$$\begin{pmatrix} a_{cc} & \cdots & a_{cd} \\ \vdots & & \vdots \\ a_{dc} & \cdots & a_{dd} \end{pmatrix}.$$

Lemma 3.7. *Let A, B be upper triangular matrices with $A \sim B$. Then $A[b, b + c] \sim B[b, b + c]$ for $b, c \in \mathbb{N}$ whenever the partial matrices are well defined.*

Proof. By definition, we have $A \sim B$ if and only if $A - I$ can be transformed into $B - I$ by pivots where a row can only be added to a row above it and a column can only be added to a column on its right. Noting that any such series of pivots on A will act on the submatrix $(A - I)[b, b + c]$ as though they were simply pivots carried out on $(A - I)[b, b + c]$ as an independent matrix, it follows that $(A - I)[b, b + c] = A[b, b + c] - I$ can be transformed into $B[b, b + c] - I$ with pivots as described in our definition and the result follows. \square

We introduce a necessary condition for two vectors \bar{m} and \bar{n} to have graphs with equivalent matrices.

Theorem 3.8. *Let $r > 2$ have prime factorisation $r = 2^j p_1^{\alpha_1} \cdots p_k^{\alpha_k}$, $j \in \mathbb{N}_0$, for distinct odd primes p_i . Further, let $\bar{m}, \bar{m}' \in (\mathbb{Z}_r)^n$ be given such that $B_{(r;\bar{m})} \sim B_{(r;\bar{m}')}$. Then for every i with $1 \leq i \leq k$ and every t with $1 \leq t \leq n - p_i$ we have*

$$\prod_{l=t+1}^{t+p_i-1} m_l \equiv \prod_{l=t+1}^{t+p_i-1} m'_l \pmod{p_i}.$$

Proof. Assume for contradiction that for some i, t we have

$$\prod_{l=t+1}^{t+p_i-1} m_l \not\equiv \prod_{l=t+1}^{t+p_i-1} m'_l \pmod{p_i}$$

and consider the matrices

$$A = B_{(r;\bar{m})}[t, t + p_i][t, t + p_i] \quad \text{and} \quad B = B_{(r;\bar{m}')}[t, t + p_i][t, t + p_i].$$

By Lemma 3.7 we must have $A \sim B$ and by Lemma 3.3, $p_i^{\alpha_i}$ divides every entry of $A - I$ and $B - I$ except the entry $\langle 1, p_i \rangle$. For the entry $\langle 1, p_i \rangle$ note that $p^{\alpha_i-1} \parallel \binom{r+p_i-1}{p_i}$ and that given integers a, b, c such that $a \not\equiv b \pmod{p}$ and $p^{\alpha-1} \parallel c$ for a prime p , then $ac \not\equiv bc \pmod{p^\alpha}$. Combining these two observations yields

$$\begin{aligned} A\langle 1, p_i \rangle &\equiv \binom{r+p_i-1}{p_i} \prod_{l=t+1}^{t+p_i-1} m_l^{-1} \\ &\not\equiv \binom{r+p_i-1}{p_i} \prod_{l=t+1}^{t+p_i-1} m_l'^{-1} \\ &\equiv B\langle 1, p_i \rangle \pmod{p_i^{\alpha_i}}. \end{aligned}$$

Thus, by Lemma 2.21 we have $A \not\sim B$, which is a contradiction. □

This necessary condition on equivalence translates directly into a lower bound on the number of equivalence classes, $\varphi_r(n)$.

Theorem 3.9. *Let $r > 2$ have prime factorisation $r = 2^j p_1^{\alpha_1} \cdots p_k^{\alpha_k}$, $j \in \mathbb{N}_0$, for odd distinct primes p_i . Then*

$$\varphi_r(n) \geq \prod_{i=1}^k \lceil (p_i - 1)^{n-p_i} \rceil.$$

Proof. For every $1 \leq i \leq k$ we define a function $T_i: (Z_{p_i})^n \rightarrow (Z_{p_i})^{n-p_i}$ given by

$$T_i(\bar{m}) = \left(\left[\prod_{l=t+1}^{t+p_i-1} m_l \right]_{p_i} \right)_{t=1}^{n-p_i}.$$

In the case $n \leq p_i$, T_i is simply the function $T_i: (Z_{p_i})^n \rightarrow \{1\}$. To show that each T_i is surjective, let $\bar{m}' \in (Z_{p_i})^{n-p_i}$ be a vector and define $\bar{m} \in (Z_{p_i})^n$ as follows:

$$m_l = \begin{cases} 1, & l < p_i \\ m'_{l-p_i+1} \left[\prod_{q=l-p_i+2}^{l-1} m_q^{-1} \right]_{p_i}, & l \geq p_i. \end{cases}$$

Since the t -th entry of $T_i(\bar{m})$ is given by

$$\left[\prod_{l=t+1}^{t+p_i-1} m_l \right]_{p_i} = \left[m_{t+p_i-1} \prod_{l=t+1}^{t+p_i-2} m_l \right]_{p_i} = m'_t,$$

it follows that $\bar{m}' \in T_i((Z_{p_i})^n)$ and thus, T_i is surjective.

Now, define the map $T: (Z_r)^n \rightarrow (Z_{p_1})^{n-p_1} \times \dots \times (Z_{p_k})^{n-p_k}$ by $T(\bar{m}) = (T_1(\bar{m}), T_2(\bar{m}), \dots, T_k(\bar{m}))$ in the natural way. Since each T_i is surjective on $(Z_{p_i})^n \rightarrow (Z_{p_i})^{n-p_i}$ it follows by the Chinese remainder theorem that T is also surjective. Now, for any two vectors $\bar{m}, \bar{n} \in (Z_r)^n$ such that $B_{(r;\bar{m})} \sim B_{(r;\bar{n})}$ we must have $T(\bar{m}) = T(\bar{n})$ by Theorem 3.8. Thus, T is an invariant of the equivalence relation \sim , it is surjective, and its codomain has $\prod_{i=1}^k \lceil (p_i - 1)^{n-p_i} \rceil$ elements and it follows that, indeed,

$$\varphi_r(n) \geq \prod_{i=1}^k \lceil (p_i - 1)^{n-p_i} \rceil. \quad \square$$

By Theorem 3.9, we now have a lower bound on the number of equivalence classes, but we conjecture that the condition in Theorem 3.8 is actually sufficient whenever $4 \nmid r$. This would then result in equality in Theorem 3.9; see Conjectures 5.3 and 5.2. Note further that using the inequality we can obtain Theorems 3.4 and 3.5 since when $n = p + 1$, where p is the least odd prime dividing r , we will get at least $(p - 1)$ classes.

4. The case of multiples of 4

Until now we have not determined $\tilde{\varphi}(r)$ in the special case where 4 divides r . This section will show that for $4 \mid r$ we have $\tilde{\varphi}(r) \leq 6$ with equality if and only if $3 \nmid r$. To this end, we start with a few lemmas regarding specific entries of $B_{(r;\bar{m})}$. Throughout the section we will change our notation slightly to make our calculations more natural, identifying the r -th vertex of any subgraph of $N_{(r;\bar{m})}$ with the 0-th.

Lemma 4.1. *Let $r > 2$ be given with $2^t \mid r$, $t > 1$, and let $\bar{m} \in (Z_r)^4$. Then*

$$2^t \mid B_{(r;\bar{m})}\langle 1, 4 \rangle.$$

Proof. By Corollary 2.16 we can assume without loss of generality that $\bar{m} = (1, m_2, 1, 1)$ for some $m_2 \in Z_r$. We calculate $B_{(r;\bar{m})}\langle 1, 4 \rangle$ by counting the number of legal paths from $c_{1,0}$ to $c_{4,0}$. We will sum over the last vertex q , $1 \leq q \leq r$, of the second subgraph that each path visits. Denote by $S_2(q)$ the number of paths from $c_{1,0}$ to $c_{2,q}$ that are subpaths of a legal path from $c_{1,0}$ to $c_{4,0}$ and similarly, let $L_2(q)$ denote the number of paths from $c_{2,q}$ to $c_{4,0}$ that do not traverse any edges of the second subgraphs and that are subpaths of a legal path from $c_{1,0}$ to $c_{4,0}$. Then

$$B_{(r;\bar{m})}\langle 1, 4 \rangle = \sum_{q=1}^r S_2(q)L_2(q).$$

First, it is not hard to see that $L_2(q) = r - q + 1$ as $m_3 = 1$. Second, if we write $q = [tm_2]_r$, $1 \leq t \leq r$ we can see that for every subpath ϕ counted by $S_2(q)$ there must be a first vertex $c_{2,v}$ of the second subgraph that it visits. We must have $v \in \{[wm_2]_r \mid 1 \leq w \leq t\}$ for else ϕ could never legally visit $c_{2,q}$. Further, there is exactly one subpath ϕ going through $c_{2,v}$ as specified, the path

$$c_{1,0} \rightarrow c_{1,1} \rightarrow \cdots \rightarrow c_{1,v} \rightarrow c_{2,v} \rightarrow c_{2,[v+m_2]_r} \rightarrow c_{2,t}.$$

It follows that

$$S_2(q) = |\{[wm_2]_r \mid 1 \leq w \leq t\}| = t \equiv qm_2^{-1} \pmod{r},$$

so we can calculate

$$B_{(r;\bar{m})}\langle 1, 4 \rangle \equiv \sum_{q=1}^r qm_2^{-1}(r - q + 1) \equiv m_2^{-1} \sum_{q=1}^r q(r - q + 1) \pmod{r}.$$

By noting that $B_{(r;\bar{1})}\langle 1, 4 \rangle \equiv \sum_{q=1}^r q(r - q + 1) \pmod{r}$, it follows that

$$B_{(r;\bar{m})}\langle 1, 4 \rangle \equiv m_2^{-1} B_{(r;\bar{1})}\langle 1, 4 \rangle \equiv m_2^{-1} \binom{r+2}{3} \equiv 0 \pmod{2^t}$$

by use of Theorem 2.17. □

Lemma 4.2. *Let $r > 2$ be given and assume that $2^t \parallel r$ for a $t > 1$ and let $\bar{m} \in (Z_r)^5$. Then*

$$2^{t-2} \parallel B_{(r;\bar{m})}\langle 1, 5 \rangle.$$

Proof. By Corollary 2.16 we can assume without loss of generality that $\bar{m} = (1, m_2, 1, m_4, 1)$. We calculate $B_{(r;\bar{m})}\langle 1, 4 \rangle$ by counting the number of legal paths from $c_{1,0}$ to $c_{5,0}$. We will sum over the last vertex q , $1 \leq q \leq r$, of the second subgraph that each path visits. Denote by $S_2(q)$ be the number of paths from $c_{1,0}$

to $c_{2,q}$ that are subpaths of a legal path from $c_{1,0}$ to $c_{5,0}$ and similarly, let $L_2(q)$ denote the number of paths from $c_{2,q}$ to $c_{5,0}$ that do not traverse any edges of the second subgraph and are subpaths of a legal path from $c_{1,0}$ to $c_{5,0}$. Then,

$$B_{(r;\bar{m})}\langle 1, 5 \rangle = \sum_{q=1}^r S_2(q)L_2(q).$$

As in the proof of Lemma 4.1, $S_2(q) \equiv qm_2^{-1} \pmod{r}$. It follows that

$$(5) \quad B_{(r;\bar{m})}\langle 1, 5 \rangle \equiv \sum_{q=1}^r qm_2^{-1}L_2(q) \equiv m_2^{-1} B_{(r;(1,1,1,m_4,1))}\langle 1, 5 \rangle \pmod{r}.$$

We proceed to calculate $B_{(r;(1,1,1,m_4,1))}\langle 1, 5 \rangle$ by almost the same approach as before. Write

$$B_{(r;(1,1,1,m_4,1))}\langle 1, 5 \rangle = \sum_{q=1}^r S_3(q)L_4(q),$$

where $S_3(q)$ is the number of paths on $N_{(r;((1,1,1,m_4,1)))}$ from $c_{1,0}$ to $c_{3,q}$ that are subpaths of a legal path from $c_{1,0}$ to $c_{5,0}$. Further, $L_4(q)$ is the number of paths from $c_{3,q}$ to $c_{5,0}$ that do not traverse any edge of the third subgraph and are subpaths of a legal path from $c_{1,0}$ to $c_{5,0}$. Let ϕ be a path counted by $L_3(q)$ and let $c_{4,v}$ be the last vertex of the fourth subgraph that ϕ visits. By $L_3(q, v)$ we count the number of such ϕ . Then,

$$\begin{aligned} B_{(r;(1,1,1,m_4,1))}\langle 1, 5 \rangle &= \sum_{q=1}^r S_3(q) \sum_{v=1}^r L_3(q, v) = \sum_{v=1}^r \sum_{q=1}^r S_3(q)L_3(q, v) \\ &= \sum_{q=1}^r S_3(q)L_3(q, r) + \sum_{v=1}^{r-1} \sum_{q=1}^r S_3(q)L_3(q, v). \end{aligned}$$

Since $\sum_{q=1}^r S_3(q)L_3(q, r)$ simply counts the number of legal paths from $c_{1,0}$ to $c_{4,r} = c_{4,0}$ that are subpaths of a legal path from $c_{1,0}$ to $c_{5,0}$, we have

$$\sum_{q=1}^r S_3(q)L_3(q, r) = B_{(r;(1,1,1,m_4,1))}\langle 1, 4 \rangle = \binom{r+2}{3} \equiv 0 \pmod{2^t}$$

by Theorem 2.17 and Lemma 2.14. Considering the case $1 \leq v < r$ yields that $L_3(q, v) = 0$ if $[qm_4^{-1}]_r > [vm_4^{-1}]_r$ since there is no legal path from $c_{4,q}$ to $c_{4,v}$ because such a path would visit $c_{4,0}$ and $v \neq 0$. Further, if $[qm_4^{-1}]_r \leq [vm_4^{-1}]_r$ we have $L_3(q, v) = 1$ since only the path

$$c_{3,q} \rightarrow c_{4,q} \rightarrow c_{4,q+m_4} \rightarrow \cdots \rightarrow c_{4,v} \rightarrow c_{5,v} \rightarrow \cdots \rightarrow c_{5,0}$$

satisfies the criteria. It follows that

$$\begin{aligned} B_{(r;(1,1,1,m_4,1))} \langle 1, 5 \rangle &\equiv \sum_{v=1}^{r-1} \sum_{q=1}^r S_3(q) L_3(q, v) \\ &\equiv \sum_{q=1}^r \sum_{\substack{1 \leq v < r \\ [qm_4^{-1}]_r \leq [vm_4^{-1}]_r}} S_3(q) \\ &= \sum_{q=1}^r [r - qm_4^{-1}]_r S_3(q) \pmod{2^t}, \end{aligned}$$

where the last equality follows since multiplying by m_4^{-1} modulo r induces a bijection on the set $\{1, \dots, r - 1\}$, yielding

$$|\{v \mid 1 \leq v < r \wedge [qm_4^{-1}]_r \leq [vm_4^{-1}]_r\}| = [r - qm_4^{-1}]_r.$$

So we get

$$\begin{aligned} B_{(r;(1,1,1,m_4,1))} \langle 1, 5 \rangle &\equiv \sum_{q=1}^r [r - qm_4^{-1}]_r S_3(q) \\ &\equiv m_4^{-1} \sum_{q=1}^r -q S_3(q) \\ &\equiv m_4^{-1} B_{(r;\bar{1})} \langle 1, 5 \rangle \pmod{2^t}. \end{aligned}$$

Inserting this into (5) then finally yields

$$\begin{aligned} B_{(r;\bar{m})} \langle 1, 5 \rangle &\equiv m_2^{-1} B_{(r;(1,1,1,m_4,1))} \langle 1, 5 \rangle \\ &\equiv m_2^{-1} m_4^{-1} B_{(r;\bar{1})} \langle 1, 5 \rangle \\ &\equiv m_2^{-1} m_4^{-1} \binom{r+3}{4} \\ &\equiv s 2^{t-2} \pmod{2^t} \end{aligned}$$

for some odd integer s since $2^{t-2} \parallel \binom{r+3}{4}$ as $4 \mid r$. □

Lemma 4.3. *Let $r > 2$ be given. Then*

$$B_{(r;(1,1,-1,1,1,1))} \langle 1, 6 \rangle = \frac{11}{20}r + \frac{3}{8}r^2 - \frac{1}{8}r^3 + \frac{1}{8}r^4 + \frac{3}{40}r^5$$

Proof. In the graph $N_{(r;(1,1,-1,1,1,1))}$, again let $S_3(q)$ be the number of paths from vertex $c_{1,0}$ to $c_{3,q}$ that are subpaths of a legal path from $c_{1,0}$ to $c_{6,0}$ such that $c_{3,q}$ is the last vertex visited in the third subgraph and let $L_3(q)$ be the number of paths from $c_{3,q}$ to $c_{6,0}$ that does not traverse any edges of the third subgraph and are

subpaths of a legal path from $c_{1,0}$ to $c_{6,0}$. We will find a closed form for each function.

First, let $0 < q < r$. Counting the paths of $S_3(q)$, we notice that there is exactly one path from $c_{1,0}$ to $c_{3,q}$ for every path from $c_{1,0}$ to $c_{2,i}$ for $p > i \geq q$. This is the path

$$c_{1,0} \rightarrow \cdots \rightarrow c_{2,i} \rightarrow c_{3,i} \rightarrow c_{3,i-1} \rightarrow \cdots \rightarrow c_{3,q}.$$

Since $m_1 = m_2 = 1$ in this case, the number of paths from $c_{1,0}$ to $c_{2,i}$ that are part of a legal path from $c_{1,0}$ to $c_{6,0}$ is i . Thus,

$$S_3(q) = \sum_{i=q}^{r-1} i = \frac{(r-q)(r+q-1)}{2}, \quad 0 < q < r.$$

The function $L_3(q)$ is only counting paths that are traversing subgraphs with parameter $m_i = 1$. We see by Corollary 2.18 that

$$L_3(q) = B_{(r-q+1; \bar{1})} \langle 1, 3 \rangle = \frac{(r-q+1)(r-q+2)}{2}.$$

Second, for $q = 0$ we have $S_3(0) = \frac{r(r+1)}{2}$ by Corollary 2.18 since this is simply $B_{(r; (1,1, -1,1,1))} \langle 1, 3 \rangle$. Further, there is only one legal subpath from $c_{4,0}$ to $c_{6,0}$ of a legal path from $c_{1,0}$ to $c_{6,0}$ so $L_3(0) = 1$.

Thus, we have

$$\begin{aligned} B_{(r; (1,1, -1,1,1))} \langle 1, 6 \rangle &= \sum_{q=0}^{r-1} S_3(q) L_3(q) \\ &= \frac{r(r+1)}{2} + \sum_{q=1}^{r-1} \frac{(r-q)(r-q+1)(r-q+2)(r+q-1)}{4} \\ &= \frac{11}{20}r + \frac{3}{8}r^2 - \frac{1}{8}r^3 + \frac{1}{8}r^4 + \frac{3}{40}r^5, \end{aligned}$$

where the last equality follows by writing out the expression and applying Faulhaber’s formula ([Graham et al. 1989, Chapter 6.5]). □

Theorem 4.4. *Let $r > 2$ be given such that $4 \mid r$. Then $\tilde{\varphi}(r) \leq 6$ with equality if and only if $3 \nmid r$.*

Proof. If $3 \mid r$, we have $\tilde{\varphi}(r) \leq 4$ by Theorem 3.4, so we will now only consider the case when $3 \nmid r$.

First, we show that $\tilde{\varphi}(r) > 5$. Let $\bar{m}, \bar{m}' \in (Z_r)^5$ be given and let $X = A_{(r; \bar{m})}$ and $Y = A_{(r; \bar{m}'})$. We will demonstrate that $X \sim Y$, proving that $\varphi_r(5) = 1$. Since $3 \nmid r$ it follows from Lemma 3.3 that if $r = s2^t$, $2 \nmid s$, then s will divide every entry of $B_{(r; \bar{m})}$ and $B_{(r; \bar{m}'})$ except for the diagonal. Thus, by Lemmas 4.1 and 4.2 the

matrices are of the form

$$X - I = \begin{pmatrix} 0 & r & \frac{1}{2}r(r+1) & x_1r & \frac{1}{4}x_2r \\ 0 & 0 & r & \frac{1}{2}r(r+1) & x_3r \\ 0 & 0 & 0 & r & \frac{1}{2}r(r+1) \\ 0 & 0 & 0 & 0 & r \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$Y - I = \begin{pmatrix} 0 & r & \frac{1}{2}r(r+1) & y_1r & \frac{1}{4}y_2r \\ 0 & 0 & r & \frac{1}{2}r(r+1) & y_3r \\ 0 & 0 & 0 & r & \frac{1}{2}r(r+1) \\ 0 & 0 & 0 & 0 & r \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

for integers $x_1, x_2, x_3, y_1, y_2, y_3$, where $2 \nmid x_2, y_2$. Now, reducing according to Definition 2.11 in a number of steps, we get

$$X - I \stackrel{1}{\cong} \begin{pmatrix} 0 & r & \frac{1}{2}r & 0 & \frac{1}{4}x_2r \\ 0 & 0 & r & \frac{1}{2}r(r+1) & 0 \\ 0 & 0 & 0 & r & \frac{1}{2}r \\ 0 & 0 & 0 & 0 & r \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\stackrel{2}{\cong} \begin{pmatrix} 0 & r & \frac{1}{2}r & 0 & \frac{1}{4}y_2r \\ 0 & 0 & r & \frac{1}{2}r(r+1) & 0 \\ 0 & 0 & 0 & r & \frac{1}{2}r \\ 0 & 0 & 0 & 0 & r \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \stackrel{3}{\cong} Y - I.$$

Step 1 reduces the entries of the first row and last column of $X - I$ modulo r by subtracting the fourth row and second column from the others. Step 2 adds the third column to the last column $\frac{1}{2}(y_2 - x_2)$ times and then subtracts the fourth row from the second $\frac{1}{2}(y_2 - x_2)$ times. Step 3 is simply the reverse of Step 1 except with $Y - I$ instead of $X - I$. It follows that $X \sim Y$.

Second, we show that $\tilde{\varphi}(r) \leq 6$, which completes the proof. Suppose that $5 \mid r$. Then it follows by Theorem 3.4 that $\tilde{\varphi}(r) \leq 6$. So assume that $3, 5 \nmid r$. Now, since $4 \mid r$, Theorem 2.17 yields

$$r \mid B_{(r;\bar{1})}(1, 6) = \binom{r+4}{5}.$$

Using Lemmas 4.1, 4.2, and 4.3 and noting that since $4 \mid r$ we have

$$\frac{11}{20}r + \frac{3}{8}r^2 - \frac{1}{8}r^3 + \frac{1}{8}r^4 + \frac{3}{40}r^5 \equiv \pm \frac{1}{4}r \pmod{r}$$

we get by Lemma 4.3 that

$$\begin{aligned}
 B_{(r;(1,1,-1,1,1,1))} - I &\stackrel{1}{\cong} \begin{pmatrix} 0 & r & \frac{1}{2}r & 0 & \frac{1}{4}x_1r & \pm\frac{1}{4}r \\ 0 & 0 & r & \frac{1}{2}r(r+1) & x_2r & \frac{1}{4}x_3r \\ 0 & 0 & 0 & r & \frac{1}{2}r(r+1) & 0 \\ 0 & 0 & 0 & 0 & r & \frac{1}{2}r \\ 0 & 0 & 0 & 0 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \\
 &\stackrel{2}{\cong} \begin{pmatrix} 0 & r & \frac{1}{2}r & 0 & \frac{1}{4}x_1r & \frac{1}{4}r \\ 0 & 0 & r & \frac{1}{2}r & x_2r & \frac{1}{4}x_3r \\ 0 & 0 & 0 & r & \frac{1}{2}r & 0 \\ 0 & 0 & 0 & 0 & r & \frac{1}{2}r \\ 0 & 0 & 0 & 0 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \stackrel{3}{\cong} \begin{pmatrix} 0 & r & \frac{1}{2}r & 0 & \frac{1}{4}r & \frac{1}{4}r \\ 0 & 0 & r & \frac{1}{2}r & 0 & \frac{1}{4}r \\ 0 & 0 & 0 & r & \frac{1}{2}r & 0 \\ 0 & 0 & 0 & 0 & r & \frac{1}{2}r \\ 0 & 0 & 0 & 0 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.
 \end{aligned}$$

and

$$\begin{aligned}
 B_{(r;(1,1,1,1,1,1))} - I &\stackrel{1}{\cong} \begin{pmatrix} 0 & r & \frac{1}{2}r & 0 & \frac{1}{4}y_1r & 0 \\ 0 & 0 & r & \frac{1}{2}r(r+1) & y_2r & \frac{1}{4}y_3r \\ 0 & 0 & 0 & r & \frac{1}{2}r(r+1) & 0 \\ 0 & 0 & 0 & 0 & r & \frac{1}{2}r \\ 0 & 0 & 0 & 0 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \\
 &\stackrel{2}{\cong} \begin{pmatrix} 0 & r & \frac{1}{2}r & 0 & \frac{1}{4}y_1r & 0 \\ 0 & 0 & r & \frac{1}{2}r & y_2r & \frac{1}{4}y_3r \\ 0 & 0 & 0 & r & \frac{1}{2}r & 0 \\ 0 & 0 & 0 & 0 & r & \frac{1}{2}r \\ 0 & 0 & 0 & 0 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \stackrel{3}{\cong} \begin{pmatrix} 0 & r & \frac{1}{2}r & 0 & \frac{1}{4}r & 0 \\ 0 & 0 & r & \frac{1}{2}r & 0 & \frac{1}{4}r \\ 0 & 0 & 0 & r & \frac{1}{2}r & 0 \\ 0 & 0 & 0 & 0 & r & \frac{1}{2}r \\ 0 & 0 & 0 & 0 & 0 & r \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.
 \end{aligned}$$

for odd $x_1, x_2, x_3, y_1, y_2, y_3$ by the following steps. Step 1 reduces the first row and last column modulo r by subtracting the second column and fifth row repeatedly from the other columns and rows. Step 2 subtracts the third column (fourth row) $\frac{r}{2}$ times from the fourth column (third row) and adds the second column (fifth row) $\frac{r}{4}$ times to the fourth column (third row). Step 3 reduces the entries $\langle 1, 5 \rangle, \langle 2, 5 \rangle,$ and $\langle 2, 6 \rangle$ modulo $\frac{r}{2}$ by subtracting the fourth column and third row repeatedly from the fifth and sixth column and first and second row repeatedly. Note that the changes to entries $\langle 4, 1 \rangle, \langle 4, 2 \rangle, \langle 5, 3 \rangle,$ and $\langle 6, 3 \rangle$ can be inverted by adding the second and third column to the fourth column and by adding the fourth and fifth row to the third row.

Now, dividing every entry by $\frac{r}{4}$, it follows that we have $B_{(r;\bar{1})} \sim B_{(r;(1,1,-1,1,1,1))}$ if and only if

$$\begin{pmatrix} 0 & 4 & 2 & 0 & 1 & \pm 1 \\ 0 & 0 & 4 & 2 & 0 & 1 \\ 0 & 0 & 0 & 4 & 2 & 0 \\ 0 & 0 & 0 & 0 & 4 & 2 \\ 0 & 0 & 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \cong \begin{pmatrix} 0 & 4 & 2 & 0 & 1 & 0 \\ 0 & 0 & 4 & 2 & 0 & 1 \\ 0 & 0 & 0 & 4 & 2 & 0 \\ 0 & 0 & 0 & 0 & 4 & 2 \\ 0 & 0 & 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

However, this can be checked to not be the case simply by solving the system of linear equations induced by Definition 2.11 and finding that there are no solutions. Our conclusion follows. □

5. Concluding remarks

Combining the results of the previous sections, we arrive at our main result, which answers the question of for which parameters n and r there only is a single, unique quantum lens space.

Theorem 5.1. *Let $r > 2$ and let p be the smallest odd prime dividing r . Then*

$$\tilde{\varphi}(r) = \begin{cases} p + 1, & 4 \nmid r \\ \min\{6, p + 1\}, & 4 \mid r. \end{cases}$$

Proof. For $4 \nmid r$ this follows directly from Theorem 3.5. Thus, let $4 \mid r$. By Corollary 2.19, $\tilde{\varphi}(r) \geq 4$, and it follows that if $p = 3$ we have $\tilde{\varphi}(r) = 4$ by Theorem 3.4 and if $p \neq 3$ we have $\tilde{\varphi}(r) = 6$ by Theorem 4.4. □

We recall that $\tilde{\varphi}(r)$ is the minimum n for which there exists an m such that $C(L_q(r, \bar{1})) \otimes \mathbb{K} \not\cong C(L_q(r, \bar{m})) \otimes \mathbb{K}$ so that our result explains exactly how to find the smallest dimension where the m -vector influences the stable isomorphism class of the quantum lens space for any fixed r . In fact, using Proposition 14.5 in [Eilers et al. 2016] we get that $\tilde{\varphi}(r)$ is the minimum n for which there is an m such that $C(L_q(r, \bar{1})) \not\cong C(L_q(r, \bar{m}))$.

Further, for the case when the quantum lens space is not uniquely given, we studied the number of equivalence classes arising by varying the parameter $\bar{m} \in (Z_r)^n$. In Theorem 3.8, a lower bound on the number of such equivalence classes was found by giving a necessary condition for two quantum lens spaces to be isomorphic. However, computer experiments suggest that this necessary condition is in fact even sufficient when $4 \nmid r$. We thus conjecture the following which we have confirmed by computer experiments for $r \in \{3, 5, 6, 9\}$ and $n \leq 8$, and for $r \in \{10, 15, 21\}$ and $n \leq 7$.

Conjecture 5.2. Let $r = 2^t \cdot p_1^{\alpha_1} \cdots p_k^{\alpha_k}$, $t \in \{0, 1\}$. Further, let $\bar{m}, \bar{m}' \in (\mathbb{Z}_r)^n$ be given. Then $B_{(r;\bar{m})} \sim B_{(r;\bar{m}')}$ if and only if for every $1 \leq i \leq k$ and $1 \leq t \leq n - p_i$,

$$\prod_{l=t+1}^{t+p_i-1} m_l \equiv \prod_{l=t+1}^{t+p_i-1} m'_l \pmod{p_i}.$$

This conjecture is true if and only if we have equality in Theorem 3.9 when $4 \nmid r$, so an equivalent conjecture is the following.

Conjecture 5.3. Let $r > 2$ have the prime factorisation $r = 2^t \cdot p_1^{\alpha_1} \cdots p_k^{\alpha_k}$, $t \in \{0, 1\}$. Then

$$\varphi_r(n) = \prod_{i=1}^k \lceil (p_i - 1)^{n-p_i} \rceil.$$

Proving these conjectures seems hard to do using the methods of this paper, however, since determining equivalence of matrices once they become sufficiently large is a complex task unless one can find better invariants to rely on. Worth noting is that proving Conjectures 5.2 and 5.3 would yield the following satisfactory result, which resounds well with the overall findings of this paper.

Conjecture 5.4. The equivalence classes of $S_{r,n} / \sim$ all have the same number of members.

Acknowledgement

The authors wish to thank Prof. Søren Eilers for his guidance throughout this entire process. From posing the problem to providing final edits, his help has been invaluable in presenting these results.

References

- [Arici et al. 2015] F. Arici, S. Brain, and G. Landi, “The Gysin sequence for quantum lens spaces”, *J. Noncommut. Geom.* **9**:4 (2015), 1077–1111. MR Zbl
- [Brzeziński and Szymański 2018] T. Brzeziński and W. Szymański, “The C^* -algebras of quantum lens and weighted projective spaces”, *J. Noncommut. Geom.* **12**:1 (2018), 195–215. MR Zbl
- [Cuntz and Krieger 1980] J. Cuntz and W. Krieger, “A class of C^* -algebras and topological Markov chains”, *Invent. Math.* **56**:3 (1980), 251–268. MR Zbl
- [Eilers et al. 2016] S. Eilers, G. Restorff, E. Ruiz, and A. P. W. Sørensen, “The complete classification of unital graph C^* -algebras: geometric and strong”, preprint, 2016. arXiv
- [Eilers et al. 2018] S. Eilers, G. Restorff, E. Ruiz, and A. P. W. Sørensen, “Geometric classification of graph C^* -algebras over finite graphs”, *Canad. J. Math.* **70**:2 (2018), 294–353. MR Zbl
- [Graham et al. 1989] R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete mathematics*, Addison-Wesley, Reading, MA, 1989. MR Zbl
- [Hong and Szymański 2003] J. H. Hong and W. Szymański, “Quantum lens spaces and graph algebras”, *Pacific J. Math.* **211**:2 (2003), 249–263. MR Zbl

[Restorff 2006] G. Restorff, “Classification of Cuntz–Krieger algebras up to stable isomorphism”, *J. Reine Angew. Math.* **598** (2006), 185–210. MR Zbl

Received January 16, 2017. Revised April 19, 2018.

PETER LUNDING JENSEN
DEPARTMENT OF MATHEMATICAL SCIENCES
UNIVERSITY OF COPENHAGEN
DENMARK
dvh408@alumni.ku.dk

FREDERIK RAVN KLAUSEN
DEPARTMENT OF MATHEMATICAL SCIENCES
UNIVERSITY OF COPENHAGEN
DENMARK
tlk870@alumni.ku.dk

PETER M. R. RASMUSSEN
DEPARTMENT OF MATHEMATICAL SCIENCES
UNIVERSITY OF COPENHAGEN
DENMARK
pmrr@di.ku.dk

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2018 is US \$475/year for the electronic version, and \$640/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 297 No. 2 December 2018

Yamabe flow with prescribed scalar curvature INAS AMACHA and RACHID REGBAOUI	257
Rigidity of proper holomorphic mappings between generalized Fock–Bargmann–Hartogs domains ENCHAO BI and ZHENHAN TU	277
Galoisian methods for testing irreducibility of order two nonlinear differential equations GUY CASALE and JACQUES-ARTHUR WEIL	299
Combinatorial classification of quantum lens spaces PETER LUNDING JENSEN, FREDERIK RAVN KLAUSEN and PETER M. R. RASMUSSEN	339
On generic quadratic forms NIKITA A. KARPENKO	367
Rankin–Cohen brackets and identities among eigenforms ARVIND KUMAR and JABAN MEHER	381
Duality for differential operators of Lie–Rinehart algebras THIERRY LAMBRE and PATRICK LE MEUR	405
Nondegeneracy of the Gauss curvature equation with negative conic singularity JUNCHENG WEI and LEI ZHANG	455
Extensions of almost faithful prime ideals in virtually nilpotent mod- p Iwasawa algebras WILLIAM WOODS	477



0030-8730(201812)297:2;1-P