

*Pacific
Journal of
Mathematics*

THE GENERAL LINEAR 2-GROUPOID

MATÍAS DEL HOYO AND DAVIDE STEFANI

Volume 298 No. 1

January 2019

THE GENERAL LINEAR 2-GROUPOID

MATÍAS DEL HOYO AND DAVIDE STEFANI

We deal with the symmetries of a (2-term) graded vector space or bundle. Our first theorem shows that they define a (strict) Lie 2-groupoid in a natural way. Our second theorem explores the construction of nerves for Lie 2-categories, showing that it yields simplicial manifolds if the 2-cells are invertible. Finally, our third and main theorem shows that smooth pseudo-functors into our general linear 2-groupoid classify 2-term representations up to homotopy of Lie groupoids.

1. Introduction	33
2. Basics on 2-categories	36
3. The nerve of a 2-category	39
4. Defining Lie 2-groupoids	43
5. The general linear 2-groupoid	45
6. The nerve of a Lie 2-category	49
7. Representations as pseudofunctors	52
Acknowledgements	56
References	56

1. Introduction

A Lie group G can be thought of as a smooth collection of symmetries of an abstract object. A linear representation $G \curvearrowright V$ is therefore a way to realize these symmetries on a concrete vector space V , which we will assume to be finite-dimensional and real. Such a representation can be defined either as a smooth map $\rho : G \times V \rightarrow V$ satisfying $\rho^h \rho^g = \rho^{hg}$ and $\rho^1 = \text{id}$, or as a Lie group morphism $G \rightarrow \text{GL}(V)$ into the general linear group. We can then study the group G by looking at its representations $G \curvearrowright V$, and this approach turns out to be very profitable.

Following the previous philosophy, a Lie groupoid $G \rightrightarrows M$ should be thought of as a smooth collection of symmetries of an abstract family parametrized by M . Lie groupoids have received much attention lately, as they provide a unifying framework for classic geometries, and also serve as models for spaces with singularities such as orbifolds and, more generally, differentiable stacks. The infinitesimal versions

MSC2010: 18G30, 22A22, 57R22.

Keywords: Lie 2-groupoids, nerve, simplicial manifolds, representation up to homotopy.

of Lie groupoids are Lie algebroids, geometric objects intertwining Lie algebra bundles and (singular) foliations. Differentiation and integration set up a fruitful interaction between the two theories.

A linear representation $(G \rightrightarrows M) \curvearrowright (V \rightarrow M)$ of a Lie groupoid over a vector bundle associates to each arrow $y \xleftarrow{g} x$ a linear isomorphism $\rho^g : V^x \rightarrow V^y$ between the corresponding fibers, in a way compatible with identities and compositions. It can be presented either as a partially defined map $G \times V \rightarrow V$ or as a Lie groupoid map $G \rightarrow \text{GL}(V)$ into the general linear groupoid [del Hoyo 2013]. The problem with Lie groupoid representations is that they are rather scarce, they impose strong conditions on V , and they do not provide us with enough information on $G \rightrightarrows M$. This reflects in the lack of an adjoint representation, or in the limitations when establishing a Tannaka duality result for Lie groupoids (see [Trentinaglia 2010]).

A solution for these problems, proposed by C. Arias Abad and M. Crainic [2013], involves representations up to homotopy $G \curvearrowright V$ of a Lie groupoid over a graded vector bundle. Such objects can be easily defined as differentials on certain bigraded algebras of sections; alternatively, they can be regarded as a sequence of tensors: a differential ∂ on V , followed by chain maps $\rho^g : V^x \rightarrow V^y$ between the fibers, then chain homotopies $\gamma^{h,g}$ relating ρ^{hg} and $\rho^h \rho^g$, etc. Representation up to homotopy has proved to be a useful concept in dealing, for instance, with cohomology theory [loc. cit.], deformations [Crainic et al. 2018] and Morita equivalences [del Hoyo and Ortiz 2018].

When $V = V_1 \oplus V_0$ is a 2-term graded vector bundle, a representation up to homotopy $G \curvearrowright V$ leads to a VB-groupoid, a double structure mixing Lie groupoids and vector bundles, via a semidirect product construction $G \ltimes V \rightarrow G$. It turns out that any VB-groupoid can be split as a semidirect product, by choosing a horizontal lift of arrows, as proven first in [Gracia-Saz and Mehta 2017]. This yields a one-to-one correspondence between VB-groupoids and 2-term representations up to homotopy, which can be extended to maps, and respect equivalence classes [del Hoyo and Ortiz 2018]. Prominent examples of VB-groupoids are the tangent and cotangent constructions. They encode the adjoint and coadjoint representations, respectively.

A VB-groupoid is an instance of a fibration of groupoids, and according to classic Grothendieck correspondence, after choosing a horizontal lift of arrows, a groupoid fibration $E \rightarrow G$ is the same as a pseudofunctor $G \dashrightarrow \{\text{Groupoids}\}$ (see [SGA 1 1971]). It follows that 2-term representations up to homotopy should, in some sense, be the same as pseudofunctors. The main purpose of the present paper is to shed light on this. To take care of the smooth and the linear structure, we are led to fix a 2-term graded vector bundle V and restrict our attention to pseudofunctors involving the several fibers of V . The resulting $G \dashrightarrow \text{GL}(V)$ is a suitable generalization of the classification map $G \rightarrow \text{GL}(V)$ for actual representations.

Given a graded vector bundle $V = V_1 \oplus V_0 \rightarrow M$, we construct a general linear

2-groupoid $GL(V)$, consisting of differentials on the fibers, quasi-isomorphisms between them, and chain homotopies. There are several nonequivalent notions of Lie 2-groupoids in the literature, some of them too strict and some others too lax for our purposes. After discussing some variants, we introduce a notion of Lie 2-groupoid, and prove our Theorem 5.5, asserting that $GL(V)$ is indeed a Lie 2-groupoid. It is remarkable that even for a 2-term graded vector space V , its general linear 2-groupoid $GL(V)$ is not a 2-group; it has more than one object, so groupoids arise naturally.

In the set-theoretic context there is a nerve for 2-categories that relates lax functors with simplicial maps [Bullejos et al. 2005; Lack 2010]. We develop the smooth version of it, and our Theorem 6.3 shows that, even though NC is not always a simplicial manifold, it is so when the Lie 2-category C has invertible 2-arrows, in particular for a Lie 2-groupoid. This nerve construction relates our notion of Lie 2-groupoids with the simplicial approach to Lie 2-groupoids, based on the horn-filling condition, which has received much attention lately. This can be seen as a piece of evidence supporting our definitions for Lie 2-groupoids and smooth pseudofunctors. We also compare our construction with that of [Mehta and Tang 2011].

Building on the previous results, which we believe are of interest in their own right, we finally establish our Theorem 7.7, setting an equivalence of categories between 2-term representations up to homotopy $G \curvearrowright V$ and pseudofunctors $G \dashrightarrow GL(V)$ commuting with basic projections. Combining this with the main theorem of [Gracia-Saz and Mehta 2017], and its extension in [del Hoyo and Ortiz 2018], we get what we might call a smooth linear variant of Grothendieck correspondence (see Remark 2.5):

$$\left\{ \begin{array}{c} \text{VB-groupoids} \\ \Gamma \rightarrow G \end{array} \right\} \rightleftharpoons \left\{ \begin{array}{c} \text{2-term representations up to homotopy} \\ G \curvearrowright V_1 \oplus V_0 \end{array} \right\} \rightleftharpoons \left\{ \begin{array}{c} \text{pseudofunctors} \\ G \dashrightarrow GL(V) \end{array} \right\}.$$

It seems natural to extend this result for higher degrees, relating positively graded representations up to homotopy and maps into a general linear ∞ -groupoid. Also, as potential applications of our theorem, we believe it is possible to relate our correspondence with the infinitesimal version announced in [Mehta 2014], and to frame the main theorem from [del Hoyo and Ortiz 2018] as a result about maps between differentiable 2-stacks. These problems will be explored elsewhere.

Organization. In Sections 2 and 3 we quickly review 2-categories and their nerves, to fix notation and provide a reference for the tools needed later. Section 4 introduces our notion of a Lie 2-groupoid and compares it with other important ones found in the literature. In Section 5 we prove our first theorem, which constructs the fundamental example: the general linear 2-groupoid. Section 6 explores the combinatorics behind the nerve of 2-categories, and exploits it to establish our second theorem: the nerve

of a Lie 2-category whose 2-cells are invertible is a simplicial manifold. In Section 7 we prove our main theorem, realizing representations up to homotopies as maps, and we discuss further questions and applications.

2. Basics on 2-categories

We review here definitions and basic facts on set-theoretic 2-categories that are fundamental for the rest of the paper. We give a definition of 2-groupoid, compare it with others in the literature, and discuss the notion of lax functors. We refer to [Borceux 1994; Lack 2010; Mac Lane 1998] for further details. The material here is preparatory, to set notation and conventions and to serve as a quick reference.

A 2-category C is a category enriched over the category of small categories. It has three levels of structure: objects, arrows between objects, and arrows between arrows or 2-cells, whose collections we denote by C_0, C_1, C_2 , respectively. We use letters x, y, \dots for objects, f, g, \dots for arrows, and α, β, \dots for 2-cells:

$$\begin{array}{ccc} & f & \\ y & \xleftarrow{\quad} & x \\ & \Downarrow \alpha & \\ & g & \end{array}$$

The arrows and 2-cells between two fixed objects x, y form a category $C(y, x)$, whose composition we denote by \bullet . For each triple x, y, z there is a composition functor $C(z, y) \times C(y, x) \xrightarrow{\circ} C(z, x)$ and a unit $\text{id}_x \in C(x, x)$ satisfying the axioms encoded in the following commutative diagrams:

$$\begin{array}{ccccc} & & C(w, z) \times C(z, y) \times C(y, x) & & \\ & \swarrow & \circ \times \text{id} & \searrow & \text{id} \times \circ \\ C(w, y) \times C(y, x) & & & & C(w, z) \times C(z, x) \\ & \searrow & \circ & \swarrow & \circ \\ & & C(w, x) & & \end{array}$$

$$\begin{array}{ccccc} & & C(y, x) & & \\ & \swarrow & \text{id}_y \times \text{id} & \searrow & \text{id} \times \text{id}_x \\ C(y, y) \times C(y, x) & & & & C(y, x) \times C(x, x) \\ & \searrow & \circ & \swarrow & \circ \\ & & C(y, x) & & \end{array}$$

Example 2.1. The paradigmatic example of a 2-category is that of small categories, functors and natural transformations. Another basic example is that of spaces, continuous maps and (homotopy classes of) homotopies.

We are interested in 2-groupoids. For us, a 2-groupoid G is a 2-category such that (i) it is small, in the sense that G_0 is a set, (ii) every 2-cell is invertible, and (iii) every arrow $y \xleftarrow{f} x$ is invertible up to homotopy, namely, there exist $x \xleftarrow{g} y$ and 2-cells $fg \cong \text{id}_y$ and $gf \cong \text{id}_x$. Some references demand the arrows be invertible on the nose. We call such 2-groupoids *strict*. Let us remark that our fundamental example, that of the general linear 2-groupoid, is not strict.

Example 2.2. A topological space X yields a 2-groupoid $\pi_2(X)$ whose objects are the points of X , whose arrows are the continuous paths $I \rightarrow X$, and whose 2-cells are (homotopy classes of) path homotopies. Composition is given by juxtaposition, moving through each path at double speed. A nonconstant path is only invertible up to homotopy, hence $\pi_2(X)$ is not strict.

A simple characterization of (small) 2-categories and strict 2-groupoids is by using *double structures*, namely diagrams of compatible structures as below, where compatible means that the horizontal structural maps are functorial with respect to the vertical structures:

$$\begin{array}{ccc} G_2 & \rightrightarrows & G_0 \\ \Downarrow & & \Downarrow \\ G_1 & \rightrightarrows & G_0 \end{array}$$

However, our notion of 2-groupoid does not benefit much from this perspective. The following lemma, which is automatic for strict groupoids but works in general, will be useful later.

Lemma 2.3. *If G is a 2-groupoid and $y \xleftarrow{f} x$ is an arrow in G , then the right multiplication functor $R_f : G(z, y) \rightarrow G(z, x)$ is an equivalence of categories for any z . The same holds for left multiplication.*

Proof. A 2-cell $\alpha : f \Rightarrow g$ defines a natural isomorphism $R_f \Rightarrow R_g$, for the 2-cells are invertible. Then, given an arbitrary f , and picking g a quasi-inverse, we have $\text{id}_{G(x,x)} = R_{\text{id}_x} \cong R_g R_f$ and analogously $\text{id}_{G(y,y)} = R_{\text{id}_y} \cong R_f R_g$. \square

A functor $\phi : C \rightarrow D$ between 2-categories consists of functions $\phi_i : C_i \rightarrow D_i$ preserving all the structure *on the nose*. This notion is sometimes too rigid, for it involves many identities between functors. A useful variant is that of a (normal) *lax functor* $\phi : C \dashrightarrow D$, which consists of three maps $\phi_i : C_i \rightarrow D_i$ preserving source, target, units and the composition \bullet , but only preserving \circ up to a given natural transformation. More precisely, also given is a map

$$\phi_{1,1} : C_1 \times_{C_0} C_1 \rightarrow D_2, \quad \phi_{1,1}(g, f) : \phi_1(gf) \Rightarrow \phi_1(g) \circ \phi_1(f),$$

ruling the failure of associativity of \circ and satisfying these coherence axioms:

- (i) $\phi_{1,1}(\text{id}_y, f) = \text{id}_f = \phi_{1,1}(f, \text{id}_x)$, where $y \xleftarrow{f} x$ (normality).

(ii) $(\phi_2(\beta) \circ \phi_2(\alpha)) \bullet \phi_{1,1}(g, f) = \phi_{1,1}(g', f') \bullet \phi_2(\beta \circ \alpha)$, where

$$z \begin{array}{c} \xleftarrow{g} \\ \Downarrow \beta \\ \xleftarrow{g'} \end{array} y \begin{array}{c} \xleftarrow{f} \\ \Downarrow \alpha \\ \xleftarrow{f'} \end{array} x .$$

(iii) $(\phi_{1,1}(h, g) \circ \phi_1(f)) \bullet \phi_{1,1}(hg, f) = (\phi_1(h) \circ \phi_{1,1}(g, f)) \bullet \phi_{1,1}(h, gf)$, where

$$w \xleftarrow{h} z \xleftarrow{g} y \xleftarrow{f} x .$$

When the structure 2-cells $\phi_{1,1}(g, f)$ are invertibles the lax functor is called a *pseudofunctor*. These notions are very interesting even when C is a usual category, viewed as a 2-category with only identity 2-cells. To ease the notation we will often write ϕ instead of ϕ_i , etc.

Example 2.4. Given an epimorphism of groups $\pi : G \rightarrow H$, a set-theoretic section $\sigma : H \rightarrow G$, $\sigma(1_H) = 1_G$, leads to a pseudofunctor $\phi : H \dashrightarrow \{\text{Groups}\}$, where G is viewed as a 2-groupoid with one object and only identity 2-cells, and $\{\text{Groups}\}$ is the 2-category of groups, morphisms, and inner automorphisms as 2-cells. Here $\phi(*) = K$ is the kernel of π , $\phi(h)$ is given by conjugation by $\sigma(h)$, and $\phi(h', h)$ is the conjugation by $\sigma(h')\sigma(h)\sigma(h'h)^{-1}$. The lax functor is an actual functor if and only if σ is a morphism.

We also need to deal with morphisms between lax functors (see [Borceux 1994]). Given lax functors $\phi, \psi : C \dashrightarrow D$ between 2-categories, a *lax transformation* $H : \phi \Rightarrow \psi$ associates to each $x \in C_0$ an arrow $H_x : \phi(x) \rightarrow \psi(x)$ and to each arrow $f : x \rightarrow y$ a 2-cell $H_f : H_y\phi(f) \Rightarrow \psi(f)H_x$ satisfying these conditions:

- (i) $H_{\text{id}_x} = \text{id}_{H_x}$ (normality),
- (ii) $(\psi(\alpha) \circ \text{id}_{H_x}) \bullet H_f = H_g \bullet (\text{id}_{H_y} \circ \phi(\alpha))$, where

$$y \begin{array}{c} \xleftarrow{f} \\ \Downarrow \alpha \\ \xleftarrow{g} \end{array} x .$$

- (iii) For each pair of composable arrows $z \xleftarrow{g} y \xleftarrow{f} x$ there is a commutative prism with vertical faces H_g, H_f, H_{gf} and horizontal faces given by the structural 2-cells of ϕ, ψ :

$$\begin{array}{ccccc} \phi(z) & \xleftarrow{\quad} & \phi(x) & & \\ & \swarrow & \searrow & & \\ & & \phi(y) & & \\ & \swarrow & \searrow & & \\ \psi(z) & \xleftarrow{\quad} & \psi(x) & & \\ & \swarrow & \searrow & & \\ & & \psi(y) & & \end{array}$$

Such an H is a *lax equivalence* if the H_x are invertible up to a 2-cell and the H_f are invertible.

Remark 2.5. Example 2.4 can be easily extended to suitable epimorphisms between categories, known as *fibered categories* [Borceux 1994; SGA 1 1971]. The outcome is the *Grothendieck correspondence* between equivalence classes of fibered categories $E \rightarrow C$ and pseudofunctors $C \dashrightarrow \{\text{Categories}\}$. This is the first and most important example of lax functors. The main goal of the present paper can be considered to be a smooth linear variant of this correspondence.

3. The nerve of a 2-category

After reviewing the classic nerve construction, we discuss here the nerve for 2-categories and 2-groupoids. We explain its behavior with respect to lax functors, and we use it to relate 2-groupoids with the weak approach to higher categories based on the horn-filling condition. Some references for this are [Bullejos and Cegarra 2003; Bullejos et al. 2005; Henriques 2008; Lack 2010].

As usual, let $[n] = \{n, n-1, \dots, 1, 0\}$ denote the ordinal of cardinality $n+1$, and let Δ be the category of finite ordinals and order-preserving maps, spanned by the elementary maps

$$d^i : [n-1] \rightarrow [n], \quad d^i(k) = \begin{cases} k & \text{if } k < i, \\ k+1 & \text{if } k \geq i, \end{cases}$$

$$s^j : [n+1] \rightarrow [n], \quad s^j(k) = \begin{cases} k & \text{if } k \leq j, \\ k-1 & \text{if } k > j, \end{cases}$$

which satisfy the so-called simplicial identities. Then a *simplicial set* is a contravariant functor $X : \Delta^\circ \rightarrow \{\text{Sets}\}$. It can be described as a sequence of sets $X_n = X([n])$ and a collection of *face* $d_i = X(d^i)$ and *degeneracy* $s_j = X(s^j)$ operators satisfying the (dual) simplicial identities. Maps of simplicial sets are natural transformations, or equivalently, sequences of maps compatible with the faces and degeneracies. Simplicial objects on a category \mathcal{C} are defined analogously.

Example 3.1. A simple but fundamental example is the *n-simplex* Δ^n . From the functorial viewpoint, it is the one represented by the ordinal $[n]$. Thinking of Δ^n as a graded set with further structure, it is freely generated by an element of type $[n]$, namely $\text{id}_{[n]}$. By Yoneda's lemma, a map $\Delta^n \rightarrow X$ is the same as an element in X_n . The *border* $\partial\Delta^n \subset \Delta^n$ is spanned by all the faces of the generator, and the *horn* $\Lambda_k^n \subset \Delta^n$ by all the faces but the k -th.

Given a category \mathcal{C} and a covariant functor $\phi : \Delta \rightarrow \mathcal{C}$, which should be thought of as a model for simplices in \mathcal{C} , we can define a *singular functor*

$$\phi^* : \mathcal{C} \rightarrow \{\text{Simplicial sets}\}$$

that associates to each object $X \in \mathcal{C}$ a simplicial set by the formula $(\phi^*X)_n = \text{hom}_{\mathcal{C}}(\phi([n]), X)$. In other words, ϕ^*X is the restriction of the contravariant functor represented by X to Δ via ϕ .

Example 3.2. When \mathcal{C} is the category of topological spaces and $\phi([n])$ is the topological n -simplex, then $\phi^*X = SX$ is the *singular simplicial set* associated to X , used to define its homology. When \mathcal{C} is the category of (small) categories and $\phi([n]) = [n]$, where we see an ordinal as a category by setting an arrow $i \rightarrow j$ if $i \leq j$, then $\phi^*C = \text{NC}$ is the *nerve* of the category, whose n -simplices are chains of n composable arrows and whose faces and degeneracies are given by dropping an extremal arrow, composing two consecutive ones, or inserting an identity.

We are concerned with the nerve construction for 2-categories, namely the singular functor defined when \mathcal{C} is the category of 2-categories and lax functors, and $\phi([n]) = [n]$ is viewed as a 2-category with only identity 2-cells. Thus, if C is a 2-category, then its *nerve* NC has as n -simplices the lax functors $u : [n] \dashrightarrow C$, and its simplicial operators are given by precomposition. Note that $\text{NC}_0 = C_0$ and $\text{NC}_1 = C_1$ consist of the objects and arrows of C , respectively, and NC_2 consists of triangles that are commutative up to a given 2-cell:

$$\begin{array}{ccc} & y & \\ g \swarrow & & \nwarrow f \\ z & \xleftarrow{h} & x \end{array} \quad \begin{array}{c} \uparrow \alpha \\ \uparrow \end{array}$$

To describe the higher simplices, note that a lax functor $u : [n] \dashrightarrow C$ can be thought of as a labeling in an abstract n -simplex, where u_i are objects at the vertices, $u_{j,i}$ are arrows at its edges, and $u_{k,j,i}$ are 2-cells corresponding to each triangle. For each tetrahedron on the simplex the following equation among 2-cells must hold:

$$\begin{array}{ccc} & u_{l,i} & \\ u_{l,k,i} \swarrow & & \searrow u_{l,j,i} \\ u_{l,k} u_{k,i} & & u_{l,j} u_{j,i} \\ u_{l,k} u_{k,j,i} \swarrow & & \searrow u_{l,k,j} u_{j,i} \\ & u_{l,k} u_{k,j} u_{j,i} & \end{array}$$

The above data completely determines the nerve NC in the sense that it is 3-*coskeletal*, that is, for $k > 3$ we have


$$\text{NC}_k = \{\partial \Delta^k \rightarrow \text{NC}\}.$$

A fundamental feature of the classic nerve for 1-categories is that it defines a fully faithful functor; it embeds the category of (small) categories into that of simplicial

sets. Extending this, there is the following proposition for the nerve of 2-categories, which also provides information about the 2-cells. Here, by a *simplicial homotopy* we mean a simplicial map $X \times \Delta^1 \rightarrow Y$.


Proposition 3.3 [Bullejos et al. 2005]. *The nerve $C \mapsto \text{NC}$ defines a fully faithful functor from the category of (small) 2-categories and (normal) lax functors to the category of simplicial sets. Moreover, if $\phi, \psi : C \dashrightarrow D$ are lax functors and every 2-cell in D is invertible, then there is a lax transformation $H : \phi \Rightarrow \psi$ if and only if there is a simplicial homotopy $\tilde{H} : N\phi \cong N\psi$.*

Sketch of proof. Given a simplicial map $\tilde{\phi} : \text{NC} \rightarrow \text{ND}$, we can define a lax functor $\phi : C \dashrightarrow D$ such that $N\phi = \tilde{\phi}$ by setting $\phi_0 = \tilde{\phi}_0$, $\phi_1 = \tilde{\phi}_1$, and defining ϕ_2 and $\phi_{1,1}$ as restrictions of $\tilde{\phi}_2$ to the following types of triangles:



The simplicial identities on $\tilde{\phi}$ imply the axioms of a lax functor on ϕ , and that $N\phi = \tilde{\phi}$, proving the first assertion.

Regarding the second triangle, given lax functors $\phi, \psi : C \dashrightarrow D$, while a lax transformation $H : \phi \cong \psi$ associates to an arrow $y \xleftarrow{f} x$ a 2-cell filling a commutative square, a simplicial homotopy $\tilde{H} : N\phi \cong N\psi$ should provide a triangulation of that square:



where $\tilde{H}_{f,0}$ and $\tilde{H}_{f,1}$ are short for $\tilde{H}(s_1(f), s_0(\text{id}_{[1]}))$ and $\tilde{H}(s_0(f), s_1(\text{id}_{[1]}))$. The lax transformation H induces a simplicial homotopy \tilde{H} by setting $\tilde{H}_{f,0} = \text{id}$ and $\tilde{H}_{f,1} = H_f$. Conversely, if every 2-cell on D is invertible, we can define an H out of \tilde{H} by setting

$$H_f = \tilde{H}_{f,1} \bullet (\tilde{H}_{f,0})^{-1}. \quad \square$$

Another fundamental feature of the classic nerve is the following characterization of its image: a simplicial set is the nerve of a category if and only if every inner horn ($0 < k < n$) admits a filling, and this filling is unique for $n > 1$. Similarly, it is

the nerve of a groupoid if and only if the same holds for every horn, inner or not:


$$\begin{array}{ccc}
 \Lambda_k^n & \xrightarrow{\forall} & X \\
 \downarrow & \nearrow \exists(!) & \\
 \Delta^n & &
 \end{array}$$

This motivates an approach to higher category theory that has received much attention lately. A simplicial set X is then a *weak m -category* if every inner horn in X admits a filling, and the filling is unique for $n > m$, and X is a *weak m -groupoid* if the same holds for every horn, inner or not. The missing face of the horn, provided by the filling, should be thought of as a *composition*, defined up to homotopy, of the remaining faces. The next proposition relates 2-groupoids with weak 2-groupoids via the nerve functor. Similar results are discussed in [Duskin 2002].

Proposition 3.4. *Let C be a 2-category. NC is a weak 2-category if and only if every 2-cell of C is invertible, and NC is a weak 2-groupoid if and only if C is a 2-groupoid.*

Proof. Since NC is 3-coskeletal, every (n, k) -horn has a unique filling for $n \geq 5$. For $n = 2$ the horizontal composition of arrows provides inner horn-fillings, and the fillings of the outer horns correspond to the existence of quasi-inverses. So let us study the cases $n = 3, 4$.

For $n = 3$, given a 2-cell $\alpha : f \Rightarrow g : x \rightarrow y$, we can build a $(3, 1)$ -horn with faces thus:




and the remaining face of a filling will give a right inverse $\beta : g \Rightarrow f$ to α , showing that inner-horn filling implies that every 2-cell is invertible. Conversely, a horn gives three 2-cells, which correspond to three sides of this square:

$$\begin{array}{ccc}
 u_{3,0} & \Longrightarrow & u_{3,1}u_{1,0} \\
 \Downarrow & & \Downarrow \\
 u_{3,2}u_{2,0} & \Longrightarrow & u_{3,2}u_{2,1}u_{1,0}
 \end{array}$$

In an inner horn, either the 2-cell on the top or in the left is missing, but since every 2-cell is invertible, we can fill the square by taking the obvious composition.

In an outer horn, either the 2-cell on the bottom or on the right is missing, and assuming C is a 2-groupoid, we can get the missing face by factoring the triple composition by either $u_{3,2}$ or $u_{1,0}$ as it follows from Lemma 2.3.

For $n = 4$, the 2-skeleton of a 4-simplex u gives the edges of a cube as below:



Each face of the 4-simplex corresponds to the commutativity of the corresponding face of the cube. The bottom face commutes because of the compatibility between horizontal and vertical composition. Since every 2-cell is invertible, five commuting faces on the cube imply that the other is commutative as well, thus every horn admits a unique filling, concluding the proof. \square

Remark 3.5. Other ways to associate a simplicial set to a 2-category C are by regarding it as a double category with a trivial side, applying twice the classic nerve, or reducing the resulting bisimplicial set by using the *diagonal* d and the *total functor* T , also known as the bar or codiagonal:

$$2\text{-categories} \xrightarrow{N^2} \text{bisimplicial sets} \xrightarrow{d, T} \text{simplicial sets}.$$

It is shown in [Bullejos and Cegarra 2003] that TN^2C and dN^2C are equivalent to NC from a homotopy viewpoint. We remark here that, when C is a strict 2-groupoid there is actually an isomorphism $TN^2C \cong NC$, which is completely determined by the following formula for 2-cells:

$$\begin{array}{ccc}
 z \xleftarrow{g} y & \xleftarrow{h} & x \\
 \parallel & \uparrow \alpha & \parallel \\
 y \xleftarrow{f} x & & x \\
 & & \parallel \\
 & & x
 \end{array}
 \mapsto
 \begin{array}{ccc}
 & y & \\
 g \swarrow & & \nwarrow f \\
 z & \xleftarrow{gh} & x \\
 & \uparrow g\alpha^{-1} &
 \end{array}$$

4. Defining Lie 2-groupoids

We discuss here the smooth versions of 2-categories and 2-groupoids we will work with, provide some examples, and discuss other uses for those terms in the literature.

A *Lie 2-category* C is, roughly speaking, a 2-category internal to the category of

smooth manifolds. It consists of a (small) 2-category as defined before, on which (i) the sets of objects C_0 , arrows C_1 and 2-cells C_2 are equipped with manifold structures; (ii) the source and target maps $s, t : C_i \rightarrow C_{i-1}$ of 2-cells and arrows are surjective submersions, and (iii) the units $u : C_{i-1} \rightarrow C_i$ and the multiplications \circ and \bullet are smooth. Functors $\phi : C \rightarrow D$ between Lie 2-categories are easy to define, as 2-functors for which the three maps $\phi_i : C_i \rightarrow D_i$ are smooth.

Example 4.1. Let (\mathbb{R}, \cdot) be the multiplicative monoid of real numbers, viewed as a Lie 2-category with a single object, space of arrows \mathbb{R} , and both horizontal and vertical composition equal to the multiplication. This is a Lie 2-category on which not every 2-cell is invertible.

Let G be a Lie 2-category that, from the set-theoretic viewpoint, is also a 2-groupoid, as defined in the previous sections. In order to define when G is a Lie 2-groupoid we have to make sense of smooth inversions. For 2-cells this is clear, for there is an inversion map $i : G_2 \rightarrow G_2$, and we can require it to be smooth. For arrows this is less clear, for inversion is only defined up to homotopy: there is no inversion map in general. Note that, since source and target $G_2 \rightarrow G_1$ are surjective submersions, the sets of 2-horns $N_{2,i}G = \text{hom}(\Lambda_i^2, \text{NG})$ define manifolds:

$$N_{2,0}G = \left\{ \begin{array}{ccc} & y & f \\ & \swarrow & \searrow \\ z & \xleftarrow{h} & x \end{array} \right\}, \quad N_{2,1}G = \left\{ \begin{array}{ccc} & g & y & f \\ & \swarrow & \swarrow & \searrow \\ z & & & x \end{array} \right\}, \quad N_{2,2}G = \left\{ \begin{array}{ccc} & g & y \\ & \swarrow & \searrow \\ z & \xleftarrow{h} & x \end{array} \right\}.$$

We will discuss a smooth structure on the whole nerve NG in the following sections. For now, we just endow N_2G with a manifold structure using the fibered product

$$\begin{array}{ccc} N_2G & \longrightarrow & N_{2,1}G \\ \downarrow & & \downarrow m \\ G_2 & \xrightarrow{t} & G_1 \end{array}$$

We define G to be a *Lie 2-groupoid* if, besides being a Lie 2-category and a 2-groupoid, (i) the inversion of 2-cells $i : G_2 \rightarrow G_2$ is smooth, and (ii) the following restriction maps are surjective submersions:

$$d_{2,0} : N_2G \rightarrow N_{2,0}G, \quad d_{2,2} : N_2G \rightarrow N_{2,2}G.$$

We say that the Lie 2-groupoid is *strict* if it is set-theoretic strict and the inversion of arrows $i : G_1 \rightarrow G_1$ is smooth. The smooth structure on N_2G also allow us to make sense of lax functors in the smooth setting. We define a *smooth lax functor* between Lie 2-categories $\phi : C \dashrightarrow D$ as a lax functor such that ϕ_0, ϕ_1 and the map $(\phi_2, \phi_{1,1}) : N_2C \rightarrow N_2D$ are smooth. A *smooth lax transformations* $H : \phi \Rightarrow \psi$ is one on which the maps $C_0 \rightarrow D_1, C_1 \rightarrow D_2$ are smooth.

Example 4.2. Given an abelian Lie group K , we can see it as the 2-cells of a Lie 2-category with one object and one arrow, and where both multiplications \bullet and \circ agree with that of K . The resulting 2-category $K \rightrightarrows * \rightrightarrows *$ is in fact a Lie 2-groupoid. A similar thing can be done with a bundle of abelian Lie groups $G \rightrightarrows M$, such as a torus bundle. This *delooping* construction stays within the finite-dimensional setting and plays a key role for instance in the theory of *gerbes*.

We would like to quickly review the *Dold–Kan construction*. When \mathcal{C} is an abelian category, e.g., that of vector spaces, then a simplicial object $X : \Delta^\circ \rightarrow \mathcal{C}$ gives rise to a chain complex (X'_n, ∂) by defining $X'_n = \bigcap_{i>0} \ker(d_i : X_n \rightarrow X_{n-1})$ and $\partial = d_0$. It turns out that this construction yields an equivalence of categories between simplicial objects and positively graded chain complexes. The horn-filling condition translates into the abelian setting, in such a way that categories and groupoids both correspond to 2-term complexes, and linear natural transformations correspond to chain homotopies.

Example 4.3. By a *linear 2-category* we mean a Lie 2-category V on which the V_i are (real finite-dimensional) vector spaces and the structure maps are linear. They are examples of Lie 2-groupoids. Viewing them as double linear categories, and applying Dold–Kan correspondence both horizontally and vertically, we encode such a V into a 3-term complex thus:

$$\begin{array}{ccc} V'_2 & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ V'_1 & \longrightarrow & V_0 \end{array}$$

Remark 4.4. The term “Lie 2-groupoid” is used in the literature in senses other than the one we have introduced, which is suitable for our fundamental example. In [Mehta and Tang 2011] and other references, it refers to what we called a strict Lie 2-groupoid and presupposes the existence of inverse arrows, whereas our notion is more general. In [Zhu 2009] and other references, a Lie 2-groupoid is defined as a smooth version of a weak 2-groupoid; the existence of a well-defined composition is not required. We will see later that a smooth version of the nerve functor for Lie 2-categories allows us to regard our Lie 2-groupoids as examples of them.

5. The general linear 2-groupoid

Here we show our first main theorem, asserting that the symmetries of a (2-term) graded vector space or bundle can be endowed with the structure of a Lie 2-groupoid, which we call the general linear 2-groupoid. This construction extends the general linear groupoid of a vector bundle without a grading (see, e.g., [del Hoyo 2013]).

Throughout this section, let $V = V_1 \oplus V_0 \rightarrow M$ be a graded vector bundle over a smooth manifold. We will first describe the set-theoretic structure of its general linear 2-groupoid $\mathrm{GL}(V)$ and then take care of the smoothness. From the set-theoretic viewpoint we have:

- (i) An object $\partial^x \in \mathrm{GL}(V)_0$ is a differential $\partial^x : V_1^x \rightarrow V_0^x$ on the fiber $V^x = V_0^x \oplus V_1^x$.
- (ii) An arrow $\alpha : \partial^x \rightarrow \partial^y \in \mathrm{GL}(V)_1$ is a couple of linear maps $\alpha_1 : V_1^x \rightarrow V_1^y$, $\alpha_0 : V_0^x \rightarrow V_0^y$, defining a quasi-isomorphism between V^x and V^y :

$$\begin{array}{ccc} V_1^x & \xrightarrow{\alpha_1} & V_1^y \\ \partial^x \downarrow & & \downarrow \partial^y \\ V_0^x & \xrightarrow{\alpha_0} & V_0^y \end{array}$$

- (iii) A 2-cell $R : \alpha \rightarrow \alpha' : \partial^x \rightarrow \partial^y$ on $\mathrm{GL}(V)_2$ is a chain homotopy, given by a linear map $R : V_0^x \rightarrow V_1^y$ such that $R\partial^x = \alpha_1 - \alpha'_1$ and $\partial^y R = \alpha_0 - \alpha'_0$.

$$\begin{array}{ccc} V_1^x & \xrightarrow{\alpha_1} & V_1^y \\ \partial^x \downarrow & \nearrow R & \downarrow \partial^y \\ V_0^x & \xrightarrow{\alpha_0} & V_0^y \end{array}$$

The multiplication \circ in $\mathrm{GL}(V)$ is the composition of maps, and the multiplication \bullet is the composition of chain homotopies, which is just the sum of the corresponding maps R . Every 2-cell is invertible, and every arrow is invertible up to a 2-cell. Thus we have a well-defined 2-groupoid $\mathrm{GL}(V)$. Via Dold–Kan we can embed it into the 2-category of linear categories.

Remark 5.1. Even when $M = *$ our construction $\mathrm{GL}(V)$ yields a 2-groupoid and not what one might call a 2-group, for there are many objects and not just one. Fixing an object ∂ on $\mathrm{GL}(V)$, its isotropy 2-groupoid $\mathrm{GL}(V)_\partial$ can be compared with the construction studied in [Sheng and Zhu 2012].

Next we show that $\mathrm{GL}(V)$ inherits a smooth structure from certain vector bundles. To ease the notation, given vector bundles $A, B \rightarrow M$, we write $[A, B] \rightarrow M$ for the inner-hom vector bundle. Then:

- (i) $\mathrm{GL}(V)_0$ identifies with the total space of $[V_1, V_0] \rightarrow M$.
- (ii) $\mathrm{GL}(V)_1$ is a subspace of

$$E = [\pi_1^* V_1, \pi_1^* V_0] \oplus [\pi_2^* V_1, \pi_2^* V_0] \oplus [\pi_1^* V_1, \pi_2^* V_1] \oplus [\pi_1^* V_0, \pi_2^* V_0],$$

a vector bundle over $M \times M$, where $\pi_i : M \times M \rightarrow M$ are the obvious projections.

(iii) $\mathrm{GL}(V)_2$ is the set-theoretic fiber product $\mathrm{GL}(V)_1 \times_{M \times M} [\pi_1^* V_0, \pi_2^* V_1]$.

The issue here is to show that $\mathrm{GL}(V)_1 \subset E$ is a submanifold. Then $\mathrm{GL}(V)_2$ will identify with a fibered product along a submersion, in fact with a pullback vector bundle. This issue is rather subtle and will require a careful analysis. The first step in our argument is to provide a simple system of equations describing $\mathrm{GL}(V)_1 \subset E$.

Lemma 5.2. *We can write $\mathrm{GL}(V)_1 = F \cap U_1 \cap U_0$, where*

$$\begin{aligned} F &= \{(\partial^x, \partial^y, \alpha_0, \alpha_1) \in E : \alpha_0 \partial^x = \partial^y \alpha_1\}, \\ U_1 &= \{(\partial^x, \partial^y, \alpha_0, \alpha_1) \in E : \ker(\partial^x) \cap \ker(\alpha_1) = 0\}, \\ U_0 &= \{(\partial^x, \partial^y, \alpha_0, \alpha_1) \in E : \mathrm{im}(\partial^y) + \mathrm{im}(\alpha_0) = V_0^y\}. \end{aligned}$$

Proof. An element $(\partial^x, \partial^y, \alpha_0, \alpha_1)$ belongs to F if and only if the corresponding square of vector space maps commutes, it belongs to U_1 if and only if the morphism between the fibers is injective in degree 1 homology, and belongs to U_0 if and only if it is surjective in degree 0 homology. Since both fibers V^x, V^y , as 2-term complexes, have the same Euler characteristic $\dim V_0 - \dim V_1$, then so do their homologies, and therefore the two inequalities $\dim H_1(V^x) \leq \dim H_1(V^y)$ and $\dim H_0(V^x) \geq \dim H_0(V^y)$ imply that α is in fact a quasi-isomorphism. \square

The subset F can be seen as the preimage of the zero section of the following map between the total space of vector bundles over $M \times M$, where $E' = [\pi_1^* V_1, \pi_2^* V_0]$:

$$\phi : E \rightarrow E', \quad \phi(\partial^x, \partial^y, \rho_1, \alpha_0) = \alpha_0 \partial^x - \partial^y \alpha_1.$$

This map is quadratic and its rank is not constant in general, as the next example shows.

Example 5.3. Let $M = *$ and $V_0 = V_1 = \mathbb{R}$. Then $\mathrm{GL}(V)_0 \cong \mathbb{R}$, $E \cong \mathbb{R}^4$ and F identifies with $\{(x, y, z, w) \in \mathbb{R}^4 : xy - zw = 0\}$, which is not a submanifold of \mathbb{R}^4 . This example shows that if we define the general linear 2-category $\mathrm{gl}(V)$ as we have defined $\mathrm{GL}(V)$, but without imposing the quasi-isomorphism axiom, then $\mathrm{gl}(V)$ cannot be made a Lie 2-category in a reasonable way.

Next we show that the map ϕ above has maximal rank over the opens U_i , and since the zero section $0_{M \times M} \subset E'$ is closed embedded, the same holds for $\mathrm{GL}(V)_1$.

Proposition 5.4. *The map $\phi : E \rightarrow E'$ has maximal rank over the opens U_i .*

Proof. Let $p = (\partial^x, \partial^y, \alpha_1, \alpha_0) \in U_1$ and let $q = \phi(p) = \alpha_0 \partial^x - \partial^y \alpha_1$. To show that $d\phi_p : T_p E \rightarrow T_q E'$ is surjective we argue by realizing vectors as 1-jets of curves. Given $\gamma(t) \in E'$, $\gamma(0) = q$, we want to lift the curve γ to a curve on E through p . By using local trivializations of V we can assume that $x(t) = x$ and

$y(t) = y$ are fixed. Let us suppose that $p \in U_1$; the other case is analogous. Since $\ker \partial^x \cap \ker \alpha_1 = 0$, the linear map $(\partial^x, \alpha_1) : V_1^x \rightarrow V_0^x \oplus V_1^y$ is a monomorphism, and therefore it admits a linear retraction $(\tilde{\partial}^x, \tilde{\alpha}_1) : V_0^x \oplus V_1^y \rightarrow V_1^x$. Then the curve $\tilde{\gamma}(t) = (\partial^x, \gamma(t)\tilde{\alpha}_1, \alpha_1, \gamma(t)\tilde{\partial}^x) \in E$ is a lift as required. \square

Theorem 5.5. *Given a graded vector bundle $V = V_1 \oplus V_0$, the general linear 2-groupoid $\text{GL}(V)$ inherits a natural structure of a Lie 2-groupoid.*

Proof. As we have already discussed, $\text{GL}(V)_0$ identifies $[V_1, V_0]$, $\text{GL}(V)_1 \subset E$ with the preimage of a closed embedded submanifold along a maximal rank map, and $\text{GL}(V)_2$ is a fiber product along a submersion. It is straightforward to check that with these definitions the structure maps of $\text{GL}(V)$ are smooth, including the inversion of 2-cells. It only remains to show that the following restriction maps are surjective submersions:

$$d_{2,0} : N_2G \rightarrow N_{2,0}G, \quad d_{2,2} : N_2G \rightarrow N_{2,2}G.$$

Let us show it for $d_{2,0}$, the other case is analogous. We argue again by lifting curves. We start with $\alpha(t) : \partial^{x(t)} \rightarrow \partial^{y(t)}$ and $\gamma(t) : \partial^{x(t)} \rightarrow \partial^{z(t)} \in \text{GL}(V)_1$, defining a curve on $N_{(2,0)}G$, and in order to lift it to N_2G , we want to define $\beta(t) : \partial^{y(t)} \rightarrow \partial^{z(t)}$ and $R(t) : \gamma(t) \Rightarrow \beta(t)\alpha(t)$. Working locally we can again assume $x = x(t)$, $y = y(t)$, $z = z(t)$ are fixed. The monomorphism $(\alpha_1(t), \partial^x(t)) : V_1^x \rightarrow V_1^y \oplus V_0^y$ admits a retraction $\tilde{\alpha}_1(t), \tilde{\partial}^x(t)$, and by basic arguments on linear algebra, we can take it smooth on t . Then the short exact sequence

$$0 \rightarrow V_1^x \xrightarrow{(\alpha_1(t), \partial^x(t))} V_1^y \oplus V_0^y \xrightarrow{(\tilde{\partial}^y(t), \alpha_0(t))} V_0^y \rightarrow 0$$

splits smoothly and we gain a section $(\tilde{\partial}^y(t), \tilde{\alpha}_0(t))$. We can then define

$$\beta_i(t) = \gamma_i(t)\tilde{\alpha}_i(t), \quad R(t) = \gamma_1(t)\tilde{\partial}^x. \quad \square$$

Remark 5.6. Let us denote by $\text{GL}'(V) \subset \text{GL}(V)$ the open Lie 2-groupoid with the same objects, arrows the invertible chain maps, and 2-cells the chain homotopies. This is a strict Lie 2-groupoid, somehow simpler than our version, and both agree around the units, thus both should behave in the same way with respect to *differentiation*, even though this process is not yet clear. See [Sheng and Zhu 2012] for a related discussion. But regarding our purposes, this simpler construction $\text{GL}(V)'$ is not satisfactory; there are representations up to homotopy of Lie groupoids that cannot be invertible. An example is the adjoint representation of the pair groupoid of the sphere $\text{Pair}(S^2)$, or of any other nonparallelizable manifold. We will come back to this later.

6. The nerve of a Lie 2-category

We deal here with the problem of endowing the nerve NC of a Lie 2-category C with a reasonable smooth structure. We show with a simple example that for general C this may not be possible. Our second main theorem shows that if every 2-cell is invertible then NC is indeed a simplicial manifold, and this happens for instance if C is a Lie 2-groupoid.

Given a Lie 2-category C , we define its *ambient* simplicial manifold AC for the nerve NC , roughly speaking, by considering arbitrary collections $\{u_{k,j,i}\}$ of 2-cells and disregarding any compatibility. More precisely, we define AC by

$$A_n C = \prod_{[2] \xrightarrow{a} [n]} C_2, \quad u \in A_n C, \quad b : [m] \rightarrow [n] \Rightarrow b^*(u)_a = u_{b \circ a} \in A_m C$$

This way AC is a well-defined simplicial manifold, and every face map is a surjective submersion, for it is just the projection onto some of the coordinates. There is a *canonical inclusion* $\phi : \text{NC} \rightarrow \text{AC}$ defined by the formula $\phi(u)_a = (u \circ a)_{2,1,0}$, where $u \in N_n C$, $u : [n] \dashrightarrow C$, and $a : [2] \rightarrow [n]$. In other words, $\phi(u)$ keeps track of the 2-cells corresponding to each triangle, and by means of the identities, the arrows on the edges and the objects on the vertices. Since every simplex in NC is determined by its 2-skeleton, the map ϕ is injective. We are concerned with the question of whether $\phi(N_n C) \subset A_n C$ is a submanifold, which is not the case in general.

Example 6.1. Let (\mathbb{R}, \cdot) be the multiplicative monoid viewed as a Lie 2-category as described in Example 4.1. Then $N_0 C = \{*\}$, $N_1 C = \{\text{id}_*\}$, and $N_2 C = \mathbb{R}$, but $N_3 C \subset A_3 C$ is not a submanifold. Disregarding the degenerate coordinates, we can identify $N_3 C$ with tuples $(x, y, z, w) \in \mathbb{R}^4$ such that $xy = zw$, the equation corresponding to the commutativity of the tetrahedron.

For C a 1-category, a simplex $u \in N_n C$ is the same as a chain of n composable arrows, so we can write $N_n C$ as an iterated fiber product, and use this to define a smooth structure on it. Next we develop a similar combinatorial description for simplices $u \in N_n C$, where C is a 2-category whose 2-cells are invertible.

We see Δ^{n-1} inside Δ^n by using the face d_n , and define a decreasing filtration

$$\Delta^n = F_0 \Delta^n \supset F_1 \Delta^n \supset \dots \supset F_{n-1} \Delta^n \supset \Delta^{n-1}$$

by setting $F_k \Delta^n = \{a : [m] \rightarrow [n] / a(m) < n \text{ or } a(0) \geq k\}$, namely $F_k \Delta^n$ is the union of Δ^{n-1} with the last face of dimension k . As an example, Figure 1 depicts the filtration for $n = 3$.

Define $N_n^k C = \{F_k \Delta^n \rightarrow \text{NC}\}$. Note that $N_n^0 C = N_n C$, that we have projections $N_n^k C \rightarrow N_n^{k+1} C$, and that $N_n^{n-1} C = N_{n-1} \times_{C_0} C_1$ is the set-theoretic fiber product over $u \mapsto u_n$ and s .




Figure 1. From left to right: $F_0\Delta^3$, $F_1\Delta^3$, $F_2\Delta^3$.

Proposition 6.2. *If every 2-cell of C is invertible then there are set-theoretic fiber products:*

$$\begin{array}{ccc} N_n^{k-1}C & \longrightarrow & C_2 \\ \downarrow & & \downarrow t \\ N_n^k C & \xrightarrow{\phi_n^k} & C_1 \end{array}, \quad \phi_n^k(u) = u_{n,k} \circ u_{k,k-1}.$$

Proof. The inclusion $F_{k+1}\Delta^n \rightarrow F_k\Delta^n$ has all the vertices on its image, all the edges except for (n, k) , and all the triangles except for (n, l, k) , with $k < l < n$. Thus, given $u : F_k\Delta^n \rightarrow \text{NC}$, if we know its restriction u' to $F_{k+1}\Delta^n$ and the 2-cell α corresponding to the triangle $(n, k+1, k)$, then we have all the vertices, we recover the edge (n, k) as the source of α , and we recover the 2-cells corresponding to the triangles (n, l, k) inductively on $l - k$ by means of the equation

$$u_{n,l,k} = (u_{n,l} \circ u_{l,k+1,k})^{-1} \bullet (u_{n,l,k+1} \circ u_{k+1,k}) \bullet u_{n,k+1,k}.$$

This shows that the map $N_n^k C \rightarrow N_n^{k+1} C \times_{C_1} C_2$ is injective.

To see that it is also surjective, we need to check that, given $u' : F_{k+1}\Delta^n \rightarrow \text{NC}$ and given $\alpha : u'_{n,k} \Rightarrow u'_{n,k+1} u'_{k+1,k}$, the above equations can be used to define a simplicial map $u : F_k\Delta^n \rightarrow \text{NC}$. This translates into showing that for every tetrahedron (l, k, j, i) the above equation holds. The only tetrahedrons that deserve an explanation are those of the type (n, l', l, k) with $k < l < l' < n$. Moreover, if $l = k + 1$ then the equation holds by the construction of u . So let us assume that $k + 1 < l$. The 4-simplex corresponding to $(n, l', l, k + 1, k)$ yields a cube:

$$\begin{array}{ccccc} & & u_{n,k} & & \\ & \swarrow & \Downarrow & \searrow & \\ u_{n,k+1}u_{k+1,k} & & u_{n,l}u_{l,k} & & u_{n,l'}u_{l',k} \\ \Downarrow & \swarrow & \Downarrow & \searrow & \Downarrow \\ u_{n,l}u_{l,k+1}u_{k+1,k} & & u_{n,l'}u_{l',k+1}u_{k+1,k} & & u_{n,l'}u_{l',l}u_{l,k} \\ & \swarrow & \Downarrow & \searrow & \\ & & u_{n,l'}u_{l',l}u_{l,k+1}u_{k+1,k} & & \end{array}$$

We want to see that the back right face commutes. But we know that: the back left face commutes by definition of $u_{n,l,k}$; the upper face commutes by definition of $u_{n,l',k}$; the left front face commutes for it factors through $u_{k+1,k}$; the right front face commutes for it factors through $u_{n,l'}$; and the bottom face commutes for \circ and \bullet are mutually distributive. Hence the result. \square

We can now prove our second main theorem.

Theorem 6.3. *Given a Lie 2-category C whose 2-arrows are (smoothly) invertible, the nerve NC is naturally a simplicial manifold.*

Proof. We endow each $N_n C$ with a smooth structure inductively. For $n = 0, 1$ we do this by means of the obvious identifications $N_0 C = C_0$ and $N_1 C = C_1$. For larger n we use the filtration and fiber products of the previous proposition, noting that one of the maps is always a surjective submersion, and using the standard transversality criterion. Hence $N_n C$ is a closed embedded submanifold of the product

$$N_n C \subset N_{n-1} C \times \prod_{(i+1,i)} C_1 \times \prod_{(n,i+1,i)} C_2.$$

We will prove that, for these smooth structures, the canonical inclusion $\phi : N_n C \rightarrow A_n C$ into the ambient is a closed embedding. This implies that (i) the smooth structures that we have defined on $N_n C$ do not depend on the particular filtration we have used, and that (ii) the simplicial maps on NC are smooth and NC is a simplicial manifold.

For each triple (k, j, i) , we have to show that the composition

$$\phi_{k,j,i} = \pi_{k,j,i} \phi : N_n C \rightarrow A_n C \rightarrow C_2$$

is smooth. By projecting on the first coordinate of the above product, and using an inductive argument, we solve the case $n > k$. By projecting on the other coordinates we solve the cases $(n, i + 1, i)$. It remains to study the other projections $\phi_{n,j,i}$. But such a projection can be written as an expression involving the other coordinates and the multiplications \circ and \bullet , which are smooth. A similar argument applies also to the degenerate coordinates. \square

It follows from our theorem that the nerve of a Lie 2-groupoid is a simplicial manifold, and that a smooth pseudofunctor $\phi : G \dashrightarrow G'$ is the same as a simplicial smooth map $\phi : NG \rightarrow NG'$. Next we present a less immediate corollary.

Corollary 6.4. *With the above hypothesis, the face maps $d_i : N_n C \rightarrow N_{n-1} C$ are surjective submersions.*

Proof. This is more a corollary of the proof rather than of the statement. When $i = n$ it follows by factoring d_n through the filtration, for each projection $N_n^k C \rightarrow N_n^{k+1} C$ is the base-change of a surjective submersion, as well as $N_n^{n-1} C \rightarrow N_{n-1} C$. When

$i \neq n$ we can argue similarly, but now using a different filtration of Δ^n , by complexes containing the face $d_i(\Delta^{n-1})$. \square

We finish this section by developing a smooth version of Proposition 3.4, setting a bridge between our theory and that of weak Lie 2-categories and weak Lie 2-groupoids, as defined in [Henriques 2008; Zhu 2009]. A simplicial manifold X is a *weak Lie m -category* or a *weak Lie m -groupoid* if the corresponding restriction maps $X_n \rightarrow X_{n,k}$ are surjective submersions, for some reasonable smooth structure on the space of (n, k) -horns. The space of horns $X_{n,k}$ can be expressed as an equalizer

$$\prod_{i \neq k} X_{n-1} \rightrightarrows \prod_{i, j \neq k} X_{n-1},$$

which may not exist in the category of manifolds. In general this is proved by an inductive argument. In our case, when $X = \text{NC}$ is the nerve of a Lie 2-category with invertible 2-arrows, it follows from our construction that $X_n \rightarrow \prod_{i \neq k} X_{n-1}$ is a closed embedded submanifold for $n > 3$ and for $n = 3$, $k = 2$. The case $n = 3$, $k = 1$, follows by using a symmetric filtration on the simplex. Therefore, since X_n is also a set-theoretic equalizer, we conclude that $X_n \cong X_{n,k}$ is a diffeomorphism in these cases. The case $n = 2$ is easy, and therefore we can conclude:

Proposition 6.5. *Let C be a Lie 2-category on which every 2-arrow is invertible. Then NC is a weak Lie 2-category. Moreover, NC is a weak Lie 2-groupoid if and only if C is a Lie 2-groupoid.*

Remark 6.6. The main theorem on [Mehta and Tang 2011] shows that if G is a strict Lie 2-groupoid then TN^2G is a weak Lie 2-groupoid. Thus, in light of the isomorphism described in Remark 3.5, our theorem can be regarded as an extension of that to a nonstrict Lie 2-groupoid. This is crucial for us, for our fundamental example $\text{GL}(V)$ is not strict.

7. Representations as pseudofunctors

In this section we review the notion of representation up to homotopy $G \curvearrowright V$ of a Lie groupoid G , the particular case of 2-term vector bundles $V = V_1 \oplus V_0$, and present our main theorem, stating a one-to-one correspondence between representations $G \curvearrowright V$ and pseudofunctors $G \dashrightarrow \text{GL}(V)$.

Given a Lie groupoid $G \rightrightarrows M$ and a vector bundle $E \rightarrow M$, a *representation* $G \curvearrowright E$ can be defined as a map $\rho : G \times_M E \rightarrow E$, $\rho(y \xleftarrow{g} x, e) = \rho_g(e)$, such that (i) $\rho_g : E_x \rightarrow E_y$ is linear, (ii) $\rho_{\text{id}} = \text{id}$, and (iii) $\rho_h \rho_g = \rho_{hg}$. A *pseudorepresentation* is a sort of nonassociative action; it is defined analogously but just requiring (i) and (ii).

Example 7.1. If $G \rightrightarrows *$ is a Lie group, viewed as a Lie groupoid with a single object, then its representations are the usual ones. If $M \rightrightarrows M$ is a manifold, viewed

as a Lie groupoid with only identities arrows, then its representations are the vector bundles over M . More generally, if $G \times M \rightrightarrows M$ is the groupoid arising from a Lie group action $G \curvearrowright M$, then a representation $(G \times M) \curvearrowright E$ is the same as an equivariant vector bundle.

Example 7.2. Given a manifold M , a representation $\text{Pair}(M) \curvearrowright E$ of its pair groupoid is the same as a trivialization of E . Given a surjective submersion $q : M \rightarrow N$, a representation $M \times_N M \curvearrowright E$ of the submersion groupoid (see [del Hoyo 2013]) is the same as an isomorphism $E \cong q^*E'$ with a pullback vector bundle. This can be further generalized to a foliation $F \subset \text{TM}$, which yields a holonomy groupoid $\text{Hol}(F) \rightrightarrows M$, whose representations are the same as foliated bundles.

Example 7.3. Let P^2 denote the real projective plane, and let $E \rightarrow P^2$ be its tautological line bundle. Since it is not trivial there cannot be a representation of the pair groupoid $\text{Pair}(P^2) \curvearrowright E$. Still, we can define a pseudorepresentation $\text{Pair}(P^2) \curvearrowright E$, by defining for instance $\rho_{(\ell, \ell')}(v)$ as the orthogonal projection of $v \in \ell$ over ℓ' .

By means of the exponential law, a Lie groupoid representation can be described as a Lie groupoid morphism into the *general linear groupoid* (see, e.g., [del Hoyo 2013])

$$\rho^\# : (G \rightrightarrows M) \rightarrow (\text{GL}(E) \rightrightarrows M), \quad \rho^\#(g) = \rho_g,$$

whose objects are the fibers of $E \rightarrow M$ and whose arrows are isomorphisms between fibers. In the case of a pseudorepresentation we still have a smooth map $G \rightarrow \text{GL}(E)$ between the arrow spaces, compatible with source and target but that may fail to preserve the multiplication. This viewpoint allows one to treat representations as maps, and it is especially useful when dealing with differentiation and integration.

Lie groupoid representations turn out to be very restrictive. A convenient generalization, is that of a *representation up to homotopy* of a Lie groupoid G over a graded vector bundle $V = \bigoplus V_i$. It is defined as a degree 1 differential D on a space of sections $\Gamma(\text{NG}, V)$ of V over the nerve of G inducing a graded module structure. By decomposing $D = \bigoplus D_i$ into bihomogeneous components, we can reinterpret D as a pseudorepresentation over a complex (V, ∂) with homotopies controlling its associativity. See [Arias Abad and Crainic 2013; del Hoyo and Ortiz 2018; Mehta and Tang 2011] for further details. We recall here the 2-term case, the simplest new case, using an homological convention.

Proposition 7.4 [del Hoyo and Ortiz 2018; Gracia-Saz and Mehta 2017]. *If $V = V_1 \oplus V_0$, then a representation up to homotopy $G \curvearrowright V$ is the same as a tuple $(\partial, \rho_1, \rho_0, \gamma)$, where $\partial : V_1 \rightarrow V_0$ is a linear map, $\rho_i : G \curvearrowright V_i$ are pseudorepresentations commuting with ∂ , and*

$$\gamma : (z \xrightarrow{h} y \xrightarrow{g} x) \mapsto (\gamma^{h,g} : \rho^{hg} \Rightarrow \rho^h \rho^g)$$

is a curvature tensor satisfying

$$\rho_1^{g_3} \circ \gamma^{g_2, g_1} - \gamma^{g_3 g_2, g_1} + \gamma^{g_3, g_2 g_1} - \gamma^{g_3, g_2} \circ \rho_0^{g_1} = 0.$$

A morphism $\theta : V \rightarrow V'$ is the same as a triple $(\theta_1, \theta_0, \mu)$ where $\theta = (\theta_1, \theta_0) : V \rightarrow V'$ is a vector bundle chain map and $\mu : (y \xrightarrow{g} x) \mapsto (\mu^g : V_0^x \rightarrow V_1^y)$ is a tensor satisfying $\rho' \theta - \theta \rho = \partial' \mu + \mu \partial$, and

$$\theta_1^z \gamma^{h, g} + \mu^h \rho_0^g + \rho_1^h \mu_g - \mu^{hg} - \gamma^{h, g} \theta_0^x = 0.$$

The point-wise homology of a 2-term representation $G \curvearrowright V$ consists of $H_1^x(V) = \ker \partial^x$ and $H_0^x(V) = \operatorname{coker} \partial^x$. If the rank of ∂ is constant then $H_1(V)$ and $H_0(V)$ are vector bundles and there is an induced representation over them. A representation up to homotopy V whose point-wise homology vanishes is called *acyclic*. A morphism $\theta : V \rightarrow W$ of 2-term representations up to homotopy inducing isomorphisms on the point-wise homology is called a *quasi-isomorphism*.

Example 7.5. for $\rho : \operatorname{Pair}(P^2) \curvearrowright E$ the pseudorepresentation discussed before, we can define an acyclic representation up to homotopy $\operatorname{Pair}(P^2) \curvearrowright E \oplus E$ by setting $\partial = \operatorname{id}$, $\rho_1 = \rho_0 = \rho$ and $\gamma = \rho - \rho\rho$. The same can be done for any pseudorepresentation.

Example 7.6. Given a Lie groupoid $G \rightrightarrows M$ endowed with a connection σ , namely a section of $s : TG \rightarrow s^* \operatorname{TM}$, the *adjoint representation* $G \curvearrowright (A \oplus \operatorname{TM})$ has ∂ equal to the anchor map and ρ_0 given by $t\sigma$. The equivalence class does not depend on σ . This generalizes the classical adjoint representation of Lie groups and plays a role in the deformation theory of groupoids. The *coadjoint representation* $G \curvearrowright T^*M \oplus A^*$ is defined by duality.

We are now ready to present our main theorem. Given a Lie groupoid $G \rightrightarrows M$ we have a canonical projection $\pi_G : G \rightarrow \operatorname{Pair}(M)$ into the pair groupoid that just remembers the source and target of an arrow. Given a 2-term vector bundle $V \rightarrow M$, we have a canonical projection $\pi_V : \operatorname{GL}(V) \rightarrow \operatorname{Pair}(M)$ that only remembers the base-points on the vector bundle.

Theorem 7.7. *Given a Lie groupoid $G \rightrightarrows M$ and a graded vector bundle $V = V_1 \oplus V_0 \rightarrow M$, there is an equivalence between the category of representations up to homotopy $\rho : G \curvearrowright V$ and quasi-isomorphisms and the category of pseudofunctors $\phi : G \dashrightarrow \operatorname{GL}(V)$ satisfying $\pi_V \phi = \pi_G$ and smooth lax equivalences.*

This result is truly a generalization of the situation for ordinary representations. That is, when V is only in degree 0, then $\operatorname{GL}(V)$ is the usual general linear groupoid, and the pseudofunctors $G \dashrightarrow \operatorname{GL}(V)$ are just morphisms of Lie groupoids.

Proof. This is a direct consequence of the constructions and results collected during our work. In light of the set-theoretical simplicial interpretation in Proposition 3.3,

our construction of the general linear 2-groupoid in Theorem 5.5, and our characterization for smooth nerve in Theorem 6.3, a smooth pseudofunctor $\phi : G \dashrightarrow \mathrm{GL}(V)$ is the same as a simplicial map $\phi : \mathrm{NG} \rightarrow \mathrm{NGL}(V)$. The degree 0 component ϕ_0 is the same as a differential ∂ on V , the degree 1 component ϕ_1 gives a pseudo-representation ρ on V compatible with ∂ , and the degree 2 component ϕ_2 yields a curvature tensor

$$\gamma : (z \xleftarrow{h} y \xleftarrow{g} x) \mapsto (\gamma^{h,g} : \rho^{hg} \Rightarrow \rho^h \rho^g),$$

defining a 2-term representation up to homotopy, as characterized in Proposition 7.4. Similarly, a smooth lax equivalence $H : \phi \Rightarrow \psi : G \dashrightarrow \mathrm{GL}(V)$ consists of smooth maps $M \rightarrow \mathrm{GL}(V)_1$, $G \rightarrow \mathrm{GL}(V)_2$, corresponding to the components θ and μ of a quasi-isomorphism (see Proposition 7.4). It is straightforward to check that these correspondences between objects and arrows are functorial. \square

There are some remarks to be made regarding functoriality. Firstly, even though a quasi-isomorphism $\theta : V \rightarrow V$ of representations up to homotopy gives a simplicial homotopy $\mathrm{NG} \times I \rightarrow \mathrm{NGL}(V)$, not every such homotopy arises in this way, as can be seen in the proof of Proposition 3.3. Secondly, if we want to consider morphisms $V \rightarrow V$ that are not quasi-isomorphisms, then the corresponding lax transformations would involve chain maps that are not within $\mathrm{GL}(V)$. Lastly, since the construction $V \mapsto \mathrm{GL}(V)$ is not functorial, it makes little sense to frame the noninvertible morphism $V \rightarrow V'$ between different vector bundles within our theory.

We close this paper by outlining three different problems related to our results, the first related to the infinitesimal picture, the second to the theory of 2-stacks, and the third to higher versions of our results.

Remark 7.8. In [Mehta 2014], an infinitesimal analog to our main theorem was announced. It is commonly accepted that weak higher Lie groupoids and higher Lie algebroids are related by a theory of differentiation and integration, though the details of such a theory are yet to be understood. Within this context, we expect that the differentiation of our general linear 2-groupoid is the object $\mathrm{gl}(V)$ introduced there, and that the differentiation and integration of maps will provide an alternative approach to the integration of 2-term representations up to homotopy, other than that of [Bursztyn et al. 2016].

Remark 7.9. In [del Hoyo and Ortiz 2018], the Morita equivalences of VB-groupoids are discussed. It is proved there that the derived category of VB-groupoids $\mathrm{VB}[G]$ over a fixed base is a Morita invariant, and consequently, the same holds for 2-term representations up to homotopy. This result, from our framework, admits the following interpretation. Our general linear 2-groupoid $\mathrm{GL}(V)$ represents a *differentiable 2-stack*, and the maps into it classify certain VB-groupoids, with prescribed side and core bundle. This should be thought of as an incarnation of

the 2-stack Perf_2 appearing in algebraic geometry. Further details demand a better understanding of differentiable 2-stacks, and are postponed to be studied elsewhere.

Remark 7.10. It is natural to expect our results to remain valid on higher degrees. The construction of the general linear groupoid seems suitable to be generalized for more general graded vector bundles. The understanding of pseudofunctors within this context seems to be less clear, though a complete immersion into the simplicial approach would solve this issue. Related to this, a realization of more general representations up to homotopy as higher VB-groupoids is currently being studied [del Hoyo and Trentinaglia \geq 2019]. Expectations here should be curbed, for even disregarding the smooth and linear structures, such a higher analog for Grothendieck correspondence is still unknown.

Acknowledgements

The authors thank H. Bursztyn, G. Ginot, A. Cabrera, E. Dubuc and M. Anel for many useful conversations that shaped their view on the topic. This collaboration took place during a series of visits of Stefani to IMPA, Rio de Janeiro, where del Hoyo worked as a visiting professor. The authors thank IMPA for this opportunity, and Capes Cofecub and Projeto P.V.E. 88881.030367/2013-01 (CAPES/Brazil) for partial financial support. Del Hoyo was partially supported by grant PQ 303034/2017-3 from National Council for Scientific and Technological Development – CNPq.

References

- [Arias Abad and Crainic 2013] C. Arias Abad and M. Crainic, “Representations up to homotopy and Bott’s spectral sequence for Lie groupoids”, *Adv. Math.* **248** (2013), 416–452. MR Zbl
- [Borceux 1994] F. Borceux, *Handbook of categorical algebra, I: Basic category theory*, Encyclopedia of Math. and Its Applications **50**, Cambridge Univ. Press, 1994. MR Zbl
- [Bullejos and Cegarra 2003] M. Bullejos and A. M. Cegarra, “On the geometry of 2-categories and their classifying spaces”, *K-Theory* **29**:3 (2003), 211–229. MR Zbl
- [Bullejos et al. 2005] M. Bullejos, E. Faro, and V. Blanco, “A full and faithful nerve for 2-categories”, *Appl. Categ. Structures* **13**:3 (2005), 223–233. MR Zbl
- [Bursztyn et al. 2016] H. Bursztyn, A. Cabrera, and M. del Hoyo, “Vector bundles over Lie groupoids and algebroids”, *Adv. Math.* **290** (2016), 163–207. MR Zbl
- [Crainic et al. 2018] M. Crainic, J. a. N. Mestre, and I. Struchiner, “Deformations of Lie groupoids”, *Int. Math. Res. Not.* (published online September 2018). arXiv
- [Duskin 2002] J. W. Duskin, “Simplicial matrices and the nerves of weak n -categories, I: Nerves of bicategories”, *Theory Appl. Categ.* **9**:10 (2002), 198–308. MR Zbl
- [Gracia-Saz and Mehta 2017] A. Gracia-Saz and R. A. Mehta, “ $\mathcal{V}\mathcal{B}$ -groupoids and representation theory of Lie groupoids”, *J. Symplectic Geom.* **15**:3 (2017), 741–783. MR Zbl
- [Henriques 2008] A. Henriques, “Integrating L_∞ -algebras”, *Compos. Math.* **144**:4 (2008), 1017–1045. MR Zbl

- [del Hoyo 2013] M. L. del Hoyo, “Lie groupoids and their orbispaces”, *Port. Math.* **70**:2 (2013), 161–209. MR Zbl
- [del Hoyo and Ortiz 2018] M. del Hoyo and C. Ortiz, “Morita equivalences of vector bundles”, *Int. Math. Res. Not.* (published online June 2018).
- [del Hoyo and Trentinaglia \geq 2019] M. del Hoyo and G. Trentinaglia, “Simplicial vector bundles over Lie groupoids”, work in progress.
- [Lack 2010] S. Lack, “A 2-categories companion”, pp. 105–191 in *Towards higher categories*, edited by J. C. Baez and J. P. May, IMA Vol. Math. Appl. **152**, Springer, 2010. MR Zbl
- [Mac Lane 1998] S. Mac Lane, *Categories for the working mathematician*, 2nd ed., Graduate Texts in Math. **5**, Springer, 1998. MR Zbl
- [Mehta 2014] R. Mehta, “Representing representations up to homotopy”, slides from lecture, Univ. California, Berkeley, 2014, Available at <https://tinyurl.com/mehtaueb>.
- [Mehta and Tang 2011] R. A. Mehta and X. Tang, “From double Lie groupoids to local Lie 2-groupoids”, *Bull. Braz. Math. Soc. (N.S.)* **42**:4 (2011), 651–681. MR Zbl
- [SGA 1 1971] A. Grothendieck, *Revêtements étales et groupe fondamental* (Séminaire de Géométrie Algébrique du Bois Marie 1960–1961), Lecture Notes in Math. **224**, Springer, 1971. MR Zbl
- [Sheng and Zhu 2012] Y. Sheng and C. Zhu, “Integration of semidirect product Lie 2-algebras”, *Int. J. Geom. Methods Mod. Phys.* **9**:5 (2012), art. id. 1250043. MR Zbl
- [Trentinaglia 2010] G. Trentinaglia, “Tannaka duality for proper Lie groupoids”, *J. Pure Appl. Algebra* **214**:6 (2010), 750–768. MR Zbl
- [Zhu 2009] C. Zhu, “ n -groupoids and stacky groupoids”, *Int. Math. Res. Not.* **2009**:21 (2009), 4087–4141. MR Zbl

Received July 13, 2017. Revised June 3, 2018.

MATÍAS DEL HOYO
UNIVERSIDADE FEDERAL FLUMINENSE
NITERÓI, RJ
BRAZIL
mldelhoyo@id.uff.br

DAVIDE STEFANI
UNIVERSITÉ PIERRE ET MARIE CURIE
PLACE JUSSIEU
PARIS
FRANCE
davide.stefani@imj-prg.fr

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2019 is US \$490/year for the electronic version, and \$665/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 298 No. 1 January 2019

On some refinements of the embedding of critical Sobolev spaces into BMO ALMAZ BUTAEV	1
A counterexample to the easy direction of the geometric Gersten conjecture. DAVID BRUCE COHEN	27
The general linear 2-groupoid MATÍAS DEL HOYO and DAVIDE STEFANI	33
Equivariant formality of Hamiltonian transversely symplectic foliations YI LIN and XIANGDONG YANG	59
Heegaard Floer homology of L -space links with two components BEIBEI LIU	83
On the Σ -invariants of wreath products LUIS AUGUSTO DE MENDONÇA	113
Enhanced adjoint actions and their orbits for the general linear group KYO NISHIYAMA and TAKUYA OHTA	141
Revisiting the saddle-point method of Perron CORMAC O'SULLIVAN	157
The Gauss–Bonnet–Chern mass of higher-codimension graphs ALEXANDRE DE SOUSA and FREDERICO GIRÃO	201
The asymptotic bounds of viscosity solutions of the Cauchy problem for Hamilton–Jacobi equations KAIZHI WANG	217
Global well-posedness for the 2D fractional Boussinesq Equations in the subcritical case DAOGUO ZHOU, ZILAI LI, HAIFENG SHANG, JIAHONG WU, BAOQUAN YUAN and JIEFENG ZHAO	233



0030-8730(201901)298:1;1-S