

Pacific Journal of Mathematics

**UNIQUENESS OF TANGENT CONES FOR BIHARMONIC MAPS
WITH ISOLATED SINGULARITIES**

YOU MIN CHEN AND HAO YIN

Volume 299 No. 2

April 2019

UNIQUENESS OF TANGENT CONES FOR BIHARMONIC MAPS WITH ISOLATED SINGULARITIES

YOU MIN CHEN AND HAO YIN

We study the problem of uniqueness of a tangent cone for minimizing extrinsic biharmonic maps. Following the celebrated result of Simon, we prove that if the target manifold is a compact analytic submanifold in $\mathbb{R}^p$ and if there is one tangent map whose singularity set consists of the origin only, then this tangent map is unique.

1. Introduction

In this paper, we prove the biharmonic map version of the celebrated result of Simon [1983]. Here we restrict ourselves to the case of extrinsic biharmonic maps. Let $B \subset \mathbb{R}^m$ be the unit ball around the origin and N be a closed Riemannian manifold isometrically embedded in $\mathbb{R}^p$. A map $u \in W^{2,2}(B, \mathbb{R}^p)$ into N is called a (extrinsic) biharmonic map if and only if it is the critical point of the energy

$$(1) \quad E(u) = \int_B |\Delta u|^2 dx.$$

It is called a minimizing (biharmonic) map if for any $B_r(x) \subset B$ and $W^{2,2}$ maps v with $v \equiv u$ on $B \setminus B_r(x)$, we have

$$E(v) \geq E(u).$$

Since the pioneering work of Chang, Wang, and Yang [Chang et al. 1999], many authors studied the regularity problem of biharmonic maps; see [Strzelecki 2003; Wang 2004a; 2004b; 2004c; Hong and Wang 2005; Lamm and Rivi re 2008]. Roughly speaking, stationary biharmonic maps are regular away from a singularity set of codimension 4. For minimizing maps, one expects better regularity since it was proved by Schoen and Uhlenbeck [1982] that minimizing harmonic maps are regular away from a singularity of codimension 3. Moreover, Luckhaus [1988] proved the compactness of minimizing harmonic maps using a lemma which was later named after him. This compactness is crucial to the theory of singularity set

The research work of Yin is supported by NSFC 11471300.

MSC2010: primary 58E20; secondary 35J30.

Keywords: biharmonic map, tangent map.

of minimizing harmonic maps. We refer the readers to the book of Simon [1996] for a nice presentation of this deep theory. The limit of a sequence of minimizing biharmonic maps was studied by Scheven [2008]. Instead of proving the direct analogue of Luckhaus lemma, the author studied the defect measure after Lin [1999]. In particular, it was shown that the limit is a stationary biharmonic map, which implies that the singularity set of minimizing biharmonic maps is of codimension 5. The interesting problem of whether this limit is minimizing remains open.

Thanks to the result of Scheven, we may study the tangent map at a singular point of a minimizing biharmonic map. The problem of uniqueness of such tangent maps is usually very difficult. Simon [1983] set up a general framework to attack such problem under a set of assumptions. The argument has been adapted to many different problems, for example, to minimal submanifolds [Simon 1983], Yang–Mills fields [Yang 2003], and Einstein metrics [Colding and Minicozzi 2014; Cheeger and Tian 1994]. To the best of our knowledge, all such generalizations are about the isolated singularity of solutions to some second order partial differential equation. It is the purpose of this paper to show that this argument also works in the case of the fourth-order problem. More precisely, we prove

Theorem 1.1. *Suppose N is an analytic submanifold of the Euclidean space $\mathbb{R}^p$ and $u : B \rightarrow N$ is a minimizing biharmonic map (with finite energy), where $B \subset \mathbb{R}^m$ ($m \geq 5$) is the unit ball. If 0 is a singularity of u and one of the tangent maps of u at 0 is of the form $\varphi(x/|x|)$ for some smooth $\varphi : S^{m-1} \rightarrow N$, then this tangent map is the unique tangent map at 0 .*

Suppose that (r, θ) is the polar coordinates in B and that $t = -\log r$. Then the theorem claims that $\lim_{t \rightarrow \infty} u(t)$ exists (and therefore is unique). As is well known, this is related to an estimate on the speed of convergence of $\partial_t u$ to zero when $t \rightarrow +\infty$. It is not hard to derive from the monotonicity formula (see [Scheven 2008, (2.4)] and (38)) that

$$(2) \quad \int_1^{+\infty} \|\partial_t u\|_{L^2(S^{m-1})}^2 < +\infty.$$

Here S^{m-1} is the unit sphere in $\mathbb{R}^m$. If we can show

$$(3) \quad \int_1^{+\infty} \|\partial_t u\|_{L^2(S^{m-1})} < +\infty,$$

then we know at least $u(t)$ converges to a unique limit in the sense of L^2 norm. However, in general, (3) does not follow from (2).

Simon [1983] observed that an infinite-dimensional version of the Łojasiewicz inequality is helpful here. In the case of a harmonic map, $u(t)$ is regarded as a

family of maps from S^{m-1} into N evolving by some second-order (abstract) ODE:

$$(4) \quad u'' - (m-2)u' = \nabla \mathcal{E}_{S^{m-1}}(v) + R,$$

where $u' = \partial_t u$, $\mathcal{E}_{S^{m-1}}$ is the harmonic map energy on S^{m-1} and R is some small perturbation term. A stationary point of this ODE (i.e., a solution independent of t) is the smooth map φ in the assumptions of the theorem (in the harmonic map case). With the help of the Łojasiewicz inequality, he studied the dynamics of this second-order ODE in a small neighborhood of φ . More precisely, he proved [Simon 1996, Chapter 3]

$$(5) \quad \left(\int_{t-1}^{+\infty} \|\partial_t u\|_{L^2(S^{m-1})}^2 \right)^{2-\alpha} \leq C \int_{t-1}^{t+1} \|\partial_t u\|_{L^2(S^{m-1})}^2$$

for any t and some $\alpha \in (0, 1)$. This amounts to (up to technical issue) an ordinary differential inequality of $h(t) = \int_t^{+\infty} \|\partial_t u\|_{L^2(S^{m-1})}^2$,

$$h(t)^{2-\alpha} \leq C(-h'(t)).$$

From this inequality, it is easy to derive some decay estimate that implies (3).

To generalize this argument to the biharmonic map case, we found that the Łojasiewicz inequality is not a problem because it is a general property of analytic functions, and the Lyapunov–Schmidt reduction works as long as the gradient of the functional is elliptic. The difficulty is to find the correct counterpart of (5). We will eventually prove a discrete version of ordinary differential inequality with time delay (see (44)). Fortunately, we can still derive the decay estimate we need from it.

The paper is organized as follows. We recall some basic properties of biharmonic maps in Section 2. In particular, we prove an improved ε -regularity lemma of Schoen and Uhlenbeck type (see [Schoen and Uhlenbeck 1982, Proposition 4.5]). In Section 3, we prove the Łojasiewicz inequality (following [Simon 1996]). Section 4 is the most important part of this paper, which contains the derivation of our analogue of (5). Finally, we give the proof of Theorem 1.1 in Section 5 following the framework of Simon [1996].

2. Preliminaries on biharmonic maps

In this section, we collect a few results, mainly PDE estimates, that are needed for the proof of our main theorem.

We start by introducing the Euler–Lagrange equation for extrinsic biharmonic energy $E(u)$ (see [Wang 2004a, Proposition 2.2]),

$$(6) \quad \Delta^2 u = \Delta(A(u)(\nabla u, \nabla u)) + 2\nabla \cdot \langle \Delta u, \nabla(P(u)) \rangle - \langle \Delta(P(u)), \Delta u \rangle.$$

Here A is the second fundamental form of N in $\mathbb{R}^p$ and $P(u)$ is the projection from $\mathbb{R}^p$ to $T_u N$. When u is a smooth (extrinsic) biharmonic map, this is equivalent to the statement that $\Delta^2 u$ is perpendicular to $T_u N$ in $\mathbb{R}^p$. Often in the following discussion, this simpler form is good enough.

An improved ε -regularity. The famous ε -regularity theorem for stationary harmonic maps requires that the (rescaled) energy is small on a ball. It has a biharmonic map analogue as follows:

Lemma 2.1 [Wang 2004a; Struwe 2008; Scheven 2008]. *There exist $\varepsilon_1 > 0$ and constants $C(k)$ only depending on N such that if u is a stationary (extrinsic) biharmonic map on $B_r(x) \subset \mathbb{R}^m$ ($m \geq 5$) satisfying*

$$(7) \quad r^{4-m} \int_{B_r(x)} (|\nabla^2 u|^2 + r^{-2} |\nabla u|^2) dx \leq \varepsilon_1,$$

then

$$\sup_{B_{r/2}(x)} r^k |\nabla^k u| \leq C(k) \quad \forall k \in \mathbb{N}.$$

Remark 2.2. Here and throughout the paper, $B_r(x)$ means the ball of radius r centered at x , which is usually omitted if $x = 0$. Also the subscript r is omitted if $r = 1$.

For minimizing harmonic maps, this result can be improved in the sense that a smallness condition on $\int_{B_r(x)} |u - u^*|^2 dx$ replaces (7), where u^* is the average of u on $B_r(x)$ (see [Schoen and Uhlenbeck 1982, Proposition 4.5]). The improved version plays an important role in the analysis of minimal tangent maps and the uniqueness of tangent cones of harmonic maps. Therefore, we also need a biharmonic map version of it.

Since the extension lemmas in [Schoen and Uhlenbeck 1982; Luckhaus 1988] are not available for biharmonic maps, the original proof in [Schoen and Uhlenbeck 1982] does not work here. Fortunately, Scheven [2008, Theorem 1.5] proved that if u_i is a sequence of minimizing biharmonic maps with bounded total energy, then there is a subsequence converging *strongly* to a stationary biharmonic map. More precisely, we have

Lemma 2.3 [Scheven 2008, Proposition 1.5]. *Suppose that $u_i : B_2 \rightarrow N$ is a sequence of minimizing biharmonic maps with bounded energy. Then there is a subsequence u_{i_j} that converges strongly to a stationary biharmonic map on B_1 .*

Proof. Since we have assumed that N is compact, then u_i are uniformly bounded on B_2 . The energy bound then implies that $\|u_i\|_{W^{2,2}(B_{3/2})}$ is bounded, so that we can use [Scheven 2008, Proposition 1.5] to get a subsequence converging to u strongly in $W^{2,2}$; u is stationary because the minimizers are stationary and the property of being stationary is preserved in strong limit. $\square$

We then combine Lemma 2.3 and Lemma 2.1 to get the biharmonic version of [Schoen and Uhlenbeck 1982, Proposition 4.5].

Lemma 2.4 (biharmonic map version of [Schoen and Uhlenbeck 1982, Proposition 4.5]). *For $\Lambda > 0$ fixed, there is $\varepsilon_2 = \varepsilon_2(N, \Lambda) > 0$ such that the following holds: Suppose that $u : B_2 \rightarrow N$ is a minimizing biharmonic map with $E(u) < \Lambda$ and $B_r(x) \subset B_1$. If*

$$r^{-m} \int_{B_r(x)} |u - q|^2 dx \leq \varepsilon_2$$

for some $q \in N$, then

$$\sup_{B_{r/4}(x)} r^k |\nabla^k u| \leq C(k) \quad \forall k \in \mathbb{N}$$

for some $C(k) > 0$.

Proof. By Lemma 2.1, it suffices to show

$$\left(\frac{1}{2}r\right)^{4-m} \int_{B_{r/2}(x)} |\nabla^2 u|^2 + \left(\frac{1}{2}r\right)^{-2} |\nabla u|^2 dx \leq \varepsilon_1.$$

If otherwise, we have a sequence of minimizing biharmonic maps $u_i : B_2 \rightarrow N$ with $E(u_i) < \Lambda$ such that for some $B_{r_i}(x_i) \subset B_1$, we have

$$(8) \quad r_i^{-m} \int_{B_{r_i}(x_i)} |u_i - q_i|^2 dx \rightarrow 0$$

and

$$(9) \quad \left(\frac{1}{2}r_2\right)^{4-m} \int_{B_{r_i/2}(x)} |\nabla^2 u_i|^2 + \left(\frac{1}{2}r_2\right)^{-2} |\nabla u_i|^2 dx \geq \varepsilon_1.$$

Note that r_i may converge to zero. Let $v_i(x) = u_i(r_i x + x_i)$. The monotonicity formula (see [Wang 2004a, Lemma 5.3]) tells us that

$$\int_{B_2(0)} |\nabla^2 v_i|^2 + |\nabla v_i|^2 dx \leq C(\Lambda).$$

By Lemma 2.3, taking subsequence if necessary, v_i converges to some stationary biharmonic map v strongly in $W^{2,2}(B_1)$, which must be the trivial map due to (8). Since the convergence is strong in $W^{2,2}$, we know that

$$\int_{B_1(0)} |\nabla^2 v_i|^2 + |\nabla v_i|^2 dx \rightarrow 0,$$

for i sufficiently large. This is a contradiction with (9) and therefore the lemma is proved. $\square$

Section of a tangent cone. Let u be the minimizing biharmonic map in Theorem 1.1. By the assumptions of the theorem, there is some sequence $r_i \rightarrow 0$ such that $u(r_i x)$ converges to a homogenous tangent map (which is biharmonic)

$$(10) \quad \tilde{\varphi} := \varphi\left(\frac{x}{|x|}\right)$$

and φ is a smooth map from S^{m-1} to N . It follows from Lemma 2.3 and Lemma 2.1 that this convergence is in fact smooth convergence away from the origin.

Recall that in the harmonic map case, $\tilde{\varphi}$ is a harmonic map if and only if so is φ . Here for the biharmonic maps, the situation is somewhat different and it is the purpose of this subsection to characterize φ that appears as the section of homogeneous biharmonic maps.

Let (r, θ) be the polar coordinates of $\mathbb{R}^m$. A direct computation shows

$$\Delta^2 \tilde{\varphi} = \left(\frac{\partial^2}{\partial r^2} + \frac{m-1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \Delta_{S^{m-1}} \right)^2 \tilde{\varphi} = r^{-4} (\Delta_{S^{m-1}}^2 \varphi - (2m-8) \Delta_{S^{m-1}} \varphi).$$

If $\tilde{\varphi}$ is a biharmonic map, then $\Delta^2 \tilde{\varphi} \perp T_{\tilde{\varphi}} N$, which is equivalent to

$$(11) \quad \Delta_{S^{m-1}}^2 \varphi - (2m-8) \Delta_{S^{m-1}} \varphi \perp T_{\varphi} N.$$

Instead of working out the explicit formula of (11), it suffices for our purpose to note that it is the Euler–Lagrange equation of the energy functional

$$(12) \quad F(\varphi) := \int_{S^{m-1}} |\Delta_{S^{m-1}} \varphi|^2 + (2m-8) |\nabla_{S^{m-1}} \varphi|^2 d\theta.$$

Here we write $d\theta$ for the volume element on S^{m-1} and φ is a map from S^{m-1} to N .

L^2 closeness implies C^5 closeness. Let φ be the smooth section in Theorem 1.1, which is a smooth critical map of F . We define

$$\mathcal{O}_{L^2}(\sigma) = \{\psi : S^{m-1} \rightarrow N \mid \|\psi - \varphi\|_{L^2(S^{m-1})} < \sigma\}$$

and

$$\mathcal{O}_{C^5}(\sigma) = \{\psi : S^{m-1} \rightarrow N \mid \|\psi - \varphi\|_{C^5(S^{m-1})} < \sigma\}.$$

Let u be a smooth biharmonic map defined on $B \setminus \{0\}$ and (t, θ) be the cylinder coordinates. In this paper, we often regard $u(t)$ as a family of maps from S^{m-1} to N . In the proof of our main theorem, these $u(t)$ are often close to φ in various sense. The next theorem roughly says that L^2 -closeness (of $u(t)$) to φ on some t -interval implies C^5 -closeness in a smaller t -interval.

Lemma 2.5. *For any $\sigma_1 > 0$, there is $\sigma_2 > 0$ (depending on σ_1 , φ , and N) such that the following is true. Let $u(t, \theta)$ be as above. If*

$$u(s) \in \mathcal{O}_{L^2}(\sigma_2) \quad \forall s \in (t_0 - 2, t_0 + 2),$$

then

$$(13) \quad u(s) \in \mathcal{O}_{C^5}(\sigma_1) \quad \forall s \in (t_0 - 1, t_0 + 1).$$

Moreover, for some $C > 0$ (depending on σ_1 , φ , and N),

$$(14) \quad \sum_{k=1,2,3,4} \sum_{j=0}^{4-k} |\partial_t^k \nabla_{S^{m-1}}^j u|(s, \theta) \leq C \quad \forall s \in (t_0 - 1, t_0 + 1).$$

Remark 2.6. It is clear from the proof below that the lemma is still true for any C^k neighborhood of φ instead of C^5 .

Proof. Although the lemma is stated in terms of (t, θ) coordinates, the proof is more clearly presented in the (r, θ) coordinates. By the scaling invariance of (6), we may assume that $t_0 = 2$ and study (6) on $B_1 \setminus B_{e^{-4}}$. By abuse of notation, we also write φ for the function

$$\varphi(r, \theta) = \varphi(\theta).$$

The assumption that $u(s) \in \mathcal{O}_{L^2}(\sigma_2)$ implies that there is a constant $C(\sigma_2)$ (satisfying $\lim_{\sigma_2 \rightarrow 0} C(\sigma_2) = 0$) such that

$$(15) \quad \int_{B_1 \setminus B_{e^{-4}}} |u - \varphi|^2 dx \leq C(\sigma_2).$$

Since φ is smooth, there is some constant C_φ depending only on φ such that

$$(16) \quad |\varphi(x) - \varphi(y)| \leq C_\varphi |x - y|$$

for any $x, y \in B_1 \setminus B_{e^{-4}}$. For some $y \in B_{e^{-1}} \setminus B_{e^{-3}}$, consider the ball $B_\sigma(y) \subset B_1 \setminus B_{e^{-4}}$ for some $\sigma > 0$ to be determined later. By (15) and (16), we have

$$\begin{aligned} \sigma^{-m} \int_{B_\sigma(y)} |u(x) - \varphi(y)|^2 dx &\leq 2\sigma^{-m} \int_{B_\sigma(y)} |u(x) - \varphi(x)|^2 dx + 2\sigma^{-m} \int_{B_\sigma(y)} |\varphi(x) - \varphi(y)|^2 dx \\ &\leq 2\sigma^{-m} C(\sigma_2) + 2|B|C_\varphi^2 \sigma^2. \end{aligned}$$

Here, $|B|$ is the volume of the unit ball in $\mathbb{R}^m$.

Let ε_2 be the constant in Lemma 2.4. We first take σ small with $2|B|C_\varphi^2 \sigma^2 < \frac{1}{2}\varepsilon_2$ and then choose σ_2 sufficiently small so that $2\sigma^{-m} C(\sigma_2) < \frac{1}{2}\varepsilon_2$. Hence, Lemma 2.4 gives

$$(17) \quad \|u\|_{C^6(B_{e^{-1}} \setminus B_{e^{-3}})} \leq C,$$

from which (14) follows. Equation (13) can be proved by interpolation between the C^6 bound (17) and the L^2 bound (15), if we choose σ_2 smaller. $\square$

Estimates of $\partial_t u$. Since being biharmonic is invariant under scaling and the group of scaling is generated by the vector field $r\partial_r = \partial_t$, if u is a biharmonic map, then $\partial_t u$ satisfies the linearization equation of (6), which is a homogeneous linear elliptic system whose coefficients depend on u . Using this equation, we can prove

Lemma 2.7. *If u satisfies (13) and (14) for $s \in (t_0 - 1, t_0 + 1)$, then we have*

$$(18) \quad \sum_{k=1,2,3,4} \sum_{j=0}^{4-k} |\partial_t^k \nabla_{S^{m-1}}^j u|^2(t_0, \theta) \leq \tilde{C} \int_{t_0-1}^{t_0+1} \int_{S^{m-1}} |\partial_t u|^2 d\theta dt,$$

for some constant $\tilde{C}$ depending only on σ_1 (in (13)), C (in (14)), and the target manifold N .

Proof. The proof is an interior estimate of an elliptic system. By scaling invariance of (6), we may assume that $t_0 = 2$. Hence to show (18), it suffices to prove

$$\|\partial_r u\|_{C^3(B_{e^{-3/2}} \setminus B_{e^{-5/2}})} \leq \tilde{C} \|\partial_r u\|_{L^2(B_{e^{-1}} \setminus B_{e^{-3}})}.$$

The observation is that if we compute the homogeneous elliptic system of $r\partial_r u$ mentioned above, the Hölder norm of all coefficients are bounded due to (13) and (14). $\square$

3. The Łojasiewicz–Simon inequality

The main purpose of this section is to prove the Łojasiewicz–Simon inequality for F defined by (12):

$$F(\psi) = \int_{S^{m-1}} |\Delta_{S^{m-1}} \psi|^2 + (2m - 8) |\nabla_{S^{m-1}} \psi|^2 d\theta.$$

Lemma 3.1. *Let φ be a smooth critical point of $F(\psi)$. Then there are $\varepsilon > 0$, $\alpha \in (0, 1]$ and $C > 0$ depending on φ such that for all $\psi : S^{m-1} \rightarrow N$ with*

$$\|\psi - \varphi\|_{C^5(S^{m-1})} \leq \varepsilon,$$

we have

$$(19) \quad |F(\psi) - F(\varphi)|^{1-\alpha/2} \leq C \|\mathcal{M}_F(\psi)\|_{L^2(S^{m-1})},$$

where $\mathcal{M}_F(\psi)$ is the Euler–Lagrange operator of F .

An equivalent form. Since φ is smooth, there is a natural correspondence between the maps that are close to φ (in C^5 topology) and the small (in C^5 norm) sections of the pull-back bundle $V := \varphi^*TN$. More precisely, we embed N isometrically as a submanifold in $\mathbb{R}^p$ and identify a section u of φ^*TN with a map

$$u : S^{m-1} \rightarrow \mathbb{R}^p \quad \text{satisfying} \quad u(\omega) \in T_{\varphi(\omega)}N \subset \mathbb{R}^p.$$

Via the nearest point projection Π defined in a tubular neighborhood of N , for any ψ close to φ , we define u by

$$(20) \quad \psi(\omega) = \Pi(\varphi(\omega) + u(\omega)).$$

This is well defined because for each ω , Π is a diffeomorphism between a neighborhood of $\varphi(\omega)$ in N and a neighborhood of 0 in $T_{\varphi(\omega)}N \subset \mathbb{R}^p$.

Since φ is a fixed smooth map, the $C^{k,\beta}$ norm of u as a section of V defined by the induced pull-back connection is equivalent to the $C^{k,\beta}$ norm of u as a map from S^{m-1} to $\mathbb{R}^p$ (with restrictions to the image). The same applies to the Sobolev norms as well. While the intrinsic role of u as a section is enough for the argument (the Lyapunov–Schmidt reduction), the extrinsic role of u as a map is important in using the analyticity assumption. (See Appendix A.)

With the above identification in mind, define

$$\tilde{F}(u) = F(\psi) - F(\varphi).$$

Then $\tilde{F}(0) = 0$ and $u = 0$ is a critical point of $\tilde{F}$. Let $\mathcal{M}_{\tilde{F}}(u)$ be the Euler–Lagrange operator of $\tilde{F}$ at u . Since the L^2 inner product of V that we use to compute $\mathcal{M}_{\tilde{F}}$ is not identical to the L^2 inner product used for the computation of $\mathcal{M}_F$, $\mathcal{M}_F(\psi)$ is not trivially the same as $\mathcal{M}_{\tilde{F}}(u)$ with u and ψ related by (20). However, we have

Lemma 3.2. *Let $\tilde{F}$ be defined as above. If ψ is a map from S^{m-1} to N with $\|\psi - \varphi\|_{C^{4,\beta}} < \delta$ for sufficiently small $\delta > 0$ and u is defined by (20), then*

$$(21) \quad (1 - C\delta)|\mathcal{M}_F(\psi)| \leq |\mathcal{M}_{\tilde{F}}(u)| \leq |\mathcal{M}_F(\psi)|.$$

The proof follows trivially from the equation (whose derivation is given in Appendix A; see (47))

$$\mathcal{M}_{\tilde{F}}(u) = P_\varphi \mathcal{M}_F(\psi)$$

and the fact that the tangent space $T_\psi N$ is close to $T_\varphi N$ since φ is close to ψ .

Given Lemma 3.2, Lemma 3.1 is reduced to

Lemma 3.3. *There are $\varepsilon > 0$, $\alpha \in (0, 1]$, and $C > 0$ depending on φ such that for all $u \in C^5(V)$ with*

$$\|u\|_{C^5(V)} \leq \varepsilon,$$

we have

$$(22) \quad |\tilde{F}(u)|^{1-\alpha/2} \leq C \|\mathcal{M}_{\tilde{F}}(u)\|_{L^2(V)}.$$

The Lyapunov–Schmidt reduction. The proof of Lemma 3.3 is an application of the Lyapunov–Schmidt reduction argument. The local behavior of $\tilde{F}$ near $u = 0$ is related to an analytic function defined on the finite dimensional kernel of an elliptic operator. More precisely, let $\mathcal{L}_{\tilde{F}}$ be the linearization of $\mathcal{M}_{\tilde{F}}$ at $u = 0$, which is an

elliptic operator from $C^{4,\beta}(V)$ to $C^{0,\beta}(V)$. By the theory of elliptic operators, the kernel of $\mathcal{L}_{\tilde{F}}$ is a finite dimensional space, denoted by K . Let P_K be the orthogonal projection of $L^2(V)$ onto K .

Setting

$$\mathcal{N}(u) = P_K u + \mathcal{M}_{\tilde{F}}(u),$$

we find that $\mathcal{N}(0) = 0$ and the linearization of $\mathcal{N}$ at $u = 0$ is given by

$$P_K + \mathcal{L}_{\tilde{F}},$$

which is an isomorphism between $C^{4,\beta}(V)$ onto $C^{0,\beta}(V)$ because it is self-adjoint with trivial kernel. The inverse function theorem then gives an inverse $\Psi = \mathcal{N}^{-1}$ from a neighborhood of $0 \in C^{0,\beta}(V)$ to $C^{4,\beta}(V)$.

Remark 3.4. (1) For the ellipticity and self-adjointness of $\mathcal{L}_{\tilde{F}}$, see Section A.

(2) The inverse function here actually appears as the real part of a complexified inverse function, which we need to justify the analyticity of f in (23) below.

Moreover, we have the following estimate for Ψ :

Lemma 3.5 (L^2 estimate). *There is a neighborhood W of 0 in $C^{0,\beta}(V)$ and a constant C , depending only on $\tilde{F}$, such that*

$$\|\Psi(f_1) - \Psi(f_2)\|_{W^{4,2}(V)} \leq C \|f_1 - f_2\|_{L^2(V)} \quad \text{for any } f_1, f_2 \in W.$$

We refer to Appendix B for the proof.

With the help of Ψ , we define

$$(23) \quad f(\xi) = \tilde{F}\left(\Psi\left(\sum_{j=1}^l \xi^j \varphi_j\right)\right)$$

for $|\xi|$ small, where $l = \dim K$ and $\{\varphi_j\}$ is a basis of K with respect to the L^2 inner product.

The key to the proof of Lemma 3.3 and hence Lemma 3.1 is the fact that f is real analytic in a neighborhood of 0. The proof relies on an analytic version of inverse function theorem for maps between complex Banach spaces and finally depends on the assumption about the analyticity of N in Theorem 1.1. It takes some effort to be precise in tracing the use of this assumption and the details of this argument are given in Appendix A.

For now, we take the analyticity of f near 0 for granted. Therefore, it follows from the classical Lojasiewicz inequality that there are constants $\alpha \in (0, 1]$, C , and $\sigma > 0$ such that

$$(24) \quad |f(\xi)|^{(1-\alpha/2)} \leq C |\nabla f(\xi)| \quad \text{for } \xi \in B_\sigma(0).$$

For the proof of Lemma 3.3, we need the following:

Lemma 3.6. *When $\|u\|_{C^{4,\beta}(V)}$ is sufficiently small and hence $\xi^j = (u, \varphi_j)_{L^2}$ is small, we have*

$$(25) \quad |\tilde{F}(u) - f(\xi)| \leq C \|\mathcal{M}_{\tilde{F}}(u)\|_{L^2}^2$$

and

$$(26) \quad \frac{1}{2} |\nabla f|(\xi) \leq \left\| \mathcal{M}_{\tilde{F}} \left(\Psi \left(\sum_{j=1}^l \xi^j \varphi_j \right) \right) \right\|_{L^2} \leq 2 |\nabla f|(\xi).$$

Before the proof of Lemma 3.6, we show how Lemma 3.3 follows from it and (24).

In fact, by plugging (25) and (26) directly into (24), we get

$$(27) \quad \begin{aligned} |\tilde{F}(u)|^{1-\alpha/2} &\leq C \left(\left\| \mathcal{M}_{\tilde{F}} \left(\Psi \left(\sum \xi^j \varphi_j \right) \right) \right\|_{L^2} + \|\mathcal{M}_{\tilde{F}}(u)\|_{L^2}^{2-\alpha} \right) \\ &\leq C \left(\left\| \mathcal{M}_{\tilde{F}} \left(\Psi \left(\sum \xi^j \varphi_j \right) \right) \right\|_{L^2} + \|\mathcal{M}_{\tilde{F}}(u)\|_{L^2} \right). \end{aligned}$$

Here in the last line above, we use the facts that $2 - \alpha \geq 1$ and that $\|\mathcal{M}_{\tilde{F}}(u)\|$ is bounded for u in the lemma. The first term in the right-hand side of (27) is dominated by the second, because

$$(28) \quad \begin{aligned} \left\| \mathcal{M}_{\tilde{F}} \left(\Psi \left(\sum \xi^j \varphi_j \right) \right) - \mathcal{M}_{\tilde{F}}(u) \right\|_{L^2} &\leq C \left\| \Psi \left(\sum \xi^j \varphi_j \right) - u \right\|_{W^{4,2}} \\ &\leq C \left\| \sum \xi^j \varphi_j - \Psi^{-1} u \right\|_{L^2} \\ &\leq C \|\mathcal{M}_{\tilde{F}}(u)\|_{L^2}. \end{aligned}$$

Here for the first line above, we noticed that $\mathcal{M}_{\tilde{F}}$ is a (nonlinear) fourth-order differential operator (see (47)) and both the $C^{4,\beta}$ norms of $\Psi(\sum \xi^j \varphi_j)$ and u are bounded; for the third line above, we used Lemma 3.5; for the last line, we used the definition of $\mathcal{N} = \Psi^{-1}$ and $P_K u = \sum \xi^j \varphi_j$. Now, Lemma 3.3 is a consequence of (27) and (28).

The rest of this section is devoted to the proof of Lemma 3.6.

The proof of Lemma 3.6. By the definition of f (see (23)) and ξ (in the assumption of Lemma 3.6), $f(\xi) = \tilde{F}(\Psi(P_K u))$. Hence, to prove (25), we compute

$$\begin{aligned} |\tilde{F}(u) - \tilde{F}(\Psi(P_K u))| &= \left| \int_0^1 \frac{d}{ds} \tilde{F}(u + s(\Psi(P_K u) - u)) ds \right| \\ &= \left| \int_0^1 (\mathcal{M}_{\tilde{F}}(u + s(\Psi(P_K u) - u)), \Psi(P_K u) - u)_{L^2} ds \right|. \end{aligned}$$

Again, by the facts that $\mathcal{M}_{\tilde{F}}$ is a fourth-order operator and that $C^{4,\beta}$ norms of u and $u + s(\Psi(P_K u) - u)$ are bounded for any $s \in [0, 1]$, we have

$$\|\mathcal{M}_{\tilde{F}}(u + s(\Psi(P_K u) - u)) - \mathcal{M}_{\tilde{F}}(u)\|_{L^2} \leq C\|\Psi(P_K u) - u\|_{W^{4,2}},$$

which implies that

$$\begin{aligned} |\tilde{F}(u) - \tilde{F}(\Psi(P_K u))| &\leq C\|\Psi(P_K u) - u\|_{L^2}(\|\mathcal{M}_{\tilde{F}}(u)\|_{L^2} + \|\Psi(P_K u) - u\|_{W^{4,2}}) \\ &\leq C\|\mathcal{M}_{\tilde{F}}(u)\|_{L^2}^2. \end{aligned}$$

Here in the last line above, we used

$$\|\Psi(P_K u) - u\|_{W^{4,2}} \leq C\|\mathcal{M}_{\tilde{F}}(u)\|_{L^2},$$

which appeared as a part of (28) and was proved there. This concludes the proof of (25).

For the proof of (26), we compute using (23) and the chain rule to get

$$(29) \quad (\eta, \nabla f(\xi))_{\mathbb{R}^l} = \left(\mathcal{M}_{\tilde{F}}\left(\Psi\left(\sum \xi^j \varphi_j\right)\right), d\Psi|_{\sum \xi^j \varphi_j}\left(\sum \eta^j \varphi_j\right) \right)_{L^2}$$

for some $\eta \in \mathbb{R}^l$ with $|\eta| = 1$.

Notice that $d\Psi|_{\sum \xi^j \varphi_j}$ depends smoothly on ξ in a compact neighborhood of $\xi = 0$, hence there is $C > 0$ such that

$$(30) \quad \|d\Psi|_{\sum \xi^j \varphi_j} - d\Psi|_0\| \leq C|\xi| \quad \text{for small } |\xi|.$$

Remark 3.7. (1) For the smooth dependence in ξ , we shall prove in Appendix A that Ψ has a complexification that is analytic (hence smooth by Theorem A.2).

(2) The norm in (30) should be the norm of bounded linear operator from $C^\beta(V)$ to $C^{4,\beta}(V)$, according to our discussion in the appendix. What we need here is the inequality

$$\left\| (d\Psi|_{\sum \xi^j \varphi_j} - d\Psi|_0)\left(\sum \eta^j \varphi_j\right) \right\|_{L^2} \leq C|\xi| \left\| \sum \eta^j \varphi_j \right\|_{L^2}.$$

This is true because $\sum \eta^j \varphi_j$ lies in K and when restricted to the finite dimensional space K , L^2 norm is equivalent to C^β norm.

On the other hand,

$$(31) \quad d\Psi|_0\left(\sum \eta^j \varphi_j\right) = \sum \eta^j \varphi_j \quad \text{for any } \eta \in \mathbb{R}^l,$$

because $d\Psi|_0 = (d\mathcal{N}|_0)^{-1} = (P_K + \mathcal{L}_{\tilde{F}})^{-1}$, and $\sum \eta^j \varphi_j$ is in K , the kernel of $\mathcal{L}_{\tilde{F}}$.

By (30) and (31), (29) implies that

$$(32) \quad \left| (\eta, \nabla f(\xi))_{\mathbb{R}^l} - \left(\mathcal{M}_{\tilde{F}} \left(\Psi \left(\sum \xi^j \varphi_j \right) \right), \sum \eta^j \varphi_j \right)_{L^2} \right| \\ \leq C|\xi| \|\mathcal{M}_{\tilde{F}} \left(\Psi \left(\sum \xi^j \varphi_j \right) \right)\|_{L^2}.$$

Now, in (32), if we choose η parallel to $\nabla f(\xi)$ in $\mathbb{R}^l$, we obtain

$$|\nabla f| \leq (1 + C|\xi|) \left\| \mathcal{M}_{\tilde{F}} \left(\Psi \left(\sum \xi^j \varphi_j \right) \right) \right\|_{L^2};$$

if we choose η so that $\sum \eta^j \varphi_j$ is parallel to $\mathcal{M}_{\tilde{F}}(\Psi(\sum \xi^j \varphi_j))$ (which is in K), then we get

$$(1 - C|\xi|) \|\mathcal{M}_{\tilde{F}} \left(\Psi \left(\sum \xi^j \varphi_j \right) \right)\|_{L^2} \leq |\nabla f|.$$

This finishes the proof of (26) and hence Lemma 3.6 if ξ is small.

4. Dynamics near a critical point of F

Let u be the minimizing biharmonic map given in Theorem 1.1. Recall that (r, θ) are the polar coordinates and that $t = -\log r$. By the assumptions of the theorem, there exists $t_i \rightarrow \infty$ such that $u(t_i, \theta)$ as maps on S^{m-1} converge smoothly to a critical point φ of F . (See the discussion in Section 2.)

Therefore, for i sufficiently large, $u(t_i, \theta)$ is very close in C^5 topology to the critical point φ of F . Since u is a biharmonic map, the biharmonic map equation determines how $u(t)$ should change as a map on S^{m-1} . In this section, we study these dynamics of $u(t)$ in a very small neighborhood of φ . More precisely, we are interested in the speed of decay of

$$\int_t^\infty \int_{S^{m-1}} |\partial_t u|^2 d\theta dt$$

as explained in the introduction. In fact, we shall control the decay of a larger quantity, namely,

$$(33) \quad G(t) = \int_t^\infty \int_{S^{m-1}} (2m-8) |\partial_t^2 u|^2 + (2m-8) |\partial_t \nabla_{S^{m-1}} u|^2 \\ + (2m-8)(m-2) |\partial_t u|^2 d\theta dt.$$

Lemma 4.1. *Suppose φ is a smooth critical point of F . There is some constant $\sigma > 0$ (depending on φ) such that if $u(t, \theta)$ (cylinder coordinates) is a smooth biharmonic map satisfying*

$$\|u(s) - \varphi\|_{C^5(S^{m-1})} \leq \sigma \quad \text{for } s \in [t-3, t+3],$$

then there exist $C' > 0$ and $\theta \in (0, 1)$ such that

$$(34) \quad G(s-1)^\theta - G(s+1)^\theta \geq C'(G(s-1) - G(s+1))^{1/2}.$$

Before we start the proof, we rewrite $\Delta^2 u$ in (t, θ) coordinates and split it into two parts. Since

$$\Delta u = e^{2t}(\partial_t^2 - (m-2)\partial_t + \Delta_{S^{m-1}})u,$$

we have

$$\begin{aligned} \Delta^2 u &= e^{4t}(\partial_t^2 + \Delta_{S^{m-1}} - (m-6)\partial_t + (8-2m))(\partial_t^2 - (m-2)\partial_t + \Delta_{S^{m-1}})u \\ &:= e^{4t}(I_a + I_b), \end{aligned}$$

where

$$\begin{aligned} I_a &= \partial_t^4 u + 2\partial_t^2 \Delta_{S^{m-1}} u - (2m-8)\partial_t^3 u - (2m-8)\partial_t \Delta_{S^{m-1}} u \\ &\quad + (m^2 - 10m + 20)\partial_t^2 u + (2m-8)(m-2)\partial_t u \end{aligned}$$

and

$$I_b = \Delta_{S^{m-1}}^2 u + (8-2m)\Delta_{S^{m-1}} u.$$

The idea behind this splitting is that we put everything involving ∂_t in I_a and the rest in I_b . An easy observation is that I_b is almost (up to a projection) the gradient of F discussed in Section 2, namely,

$$\begin{aligned} (35) \quad \partial_t F(u(t)) &= 2 \int_{S^{m-1}} \Delta_{S^{m-1}} u \Delta_{S^{m-1}} \partial_t u + (2m-8) \nabla_{S^{m-1}} u \cdot \nabla_{S^{m-1}} \partial_t u \, d\theta \\ &= 2 \int_{S^{m-1}} I_b \cdot \partial_t u \, d\theta. \end{aligned}$$

The way we use the biharmonic map equation has nothing to do with the right-hand side of (6). We multiply the equation by $\partial_t u$ and integrate over S^{m-1} to obtain

$$(36) \quad 0 = \int_{S^{m-1}} \Delta^2 u \cdot \partial_t u \, d\theta = \int_{S^{m-1}} (I_a + I_b) \cdot \partial_t u \, d\theta.$$

While $\int_{S^{m-1}} I_b \cdot \partial_t u \, d\theta$ is known in (35), the structure of $\int_{S^{m-1}} I_a \cdot \partial_t u \, d\theta$ is still complicated. There is some positivity hidden in it. To reveal it, we use the elementary equalities

$$\partial_t^4 u \cdot \partial_t u = \partial_t (\partial_t^3 u \partial_t u - \frac{1}{2} |\partial_t^2 u|^2)$$

and

$$\partial_t^3 u \cdot \partial_t u = \partial_t (\partial_t^2 u \cdot \partial_t u) - |\partial_t^2 u|^2$$

to get

$$\begin{aligned}
& \int_{S^{m-1}} I_a \cdot \partial_t u \, d\theta \\
&= \partial_t \left(\int_{S^{m-1}} \partial_t^3 u \partial_t u - \frac{1}{2} |\partial_t^2 u|^2 \right. \\
&\quad \left. - |\partial_t \nabla_{S^{m-1}} u|^2 - (2m-8) \partial_t^2 u \partial_t u + \frac{1}{2} (m^2 - 10m + 20) |\partial_t u|^2 \, d\theta \right) \\
&\quad + \left(\int_{S^{m-1}} (2m-8) |\partial_t^2 u|^2 + (2m-8) |\partial_t \nabla_{S^{m-1}} u|^2 \right. \\
&\quad \left. + (2m-8)(m-2) |\partial_t u|^2 \, d\theta \right) \\
&:= \partial_t \left(\int_{S^{m-1}} II_a \, d\theta \right) + \int_{S^{m-1}} II_b \, d\theta.
\end{aligned}$$

Notice that II_b is nonnegative and this is how we obtain the definition of $G(t)$ in (33), i.e.,

$$G(t) = \int_t^\infty \int_{S^{m-1}} II_b \, d\theta \, dt.$$

By (36) and (35), we have

$$(37) \quad \frac{1}{2} \partial_t F(u(t)) = - \int_{S^{m-1}} I_a \cdot \partial_t u \, d\theta.$$

Let t_i be the sequence mentioned in the beginning of this section such that $u(t_i)$ converges smoothly to the smooth section map φ . Moreover, $u(t + t_i)$ regarded as a map defined on $[-1, 1] \times S^{m-1}$ converges smoothly to $\tilde{\varphi}(t, \theta) = \varphi(\theta)$. This implies that

$$\lim_{i \rightarrow \infty} \int_{S^{m-1}} II_a(t_i) \, d\theta = 0,$$

so that if we integrate (37) from s to t_i and take the limit $i \rightarrow \infty$, we obtain

$$(38) \quad \frac{1}{2} (F(\varphi) - F(u(s))) = \int_{S^{m-1}} II_a(s) \, d\theta - \int_s^{+\infty} \int_{S^{m-1}} II_b \, d\theta.$$

As a by-product of the above computation, $G(t)$ is a finite number, which is the biharmonic counterpart of (2).

We may choose σ small so that for u in the lemma and $s \in [t-3, t+3]$, $\|u(s) - \varphi\|_{C^5(S^{m-1})}$ is small and hence $u(s)$ satisfies the assumption of Lemma 3.1. The Łojasiewicz–Simon inequality (in Lemma 3.1) and (38) imply that

$$(39) \quad - \int_{S^{m-1}} II_a(s) \, d\theta + \int_s^{+\infty} \int_{S^{m-1}} II_b \, d\theta \, dt \leq C \|\mathcal{M}_F(u(s))\|_{L^2(S^{m-1})}^{2/(2-\alpha)}$$

for some $\alpha \in (0, 1]$.

Next, we show that the right-hand side and the first term in the left-hand side of (39) are controlled by $\int_{s-1}^{s+1} |II_b|^2 d\theta$. To see this, recall that by (35), $\mathcal{M}_F(u(s))$ is the projection of $I_b(s)$ onto the tangent bundle of TN at $u(s)$. If we denote this projection from $\mathbb{R}^p$ onto $T_u N$ by Π ,

$$(40) \quad \mathcal{M}_F(u(s)) = 2\Pi(I_b(s)).$$

On the other hand, since u is an extrinsic biharmonic map, the Euler–Lagrange equation reads

$$(41) \quad \Pi(\Delta^2 u) = \Pi(I_a + I_b) = 0.$$

Combining (40) and (41), we get

$$(42) \quad \|\mathcal{M}_F(u(s))\|_{L^2(S^{m-1})} \leq 2\|I_a(s)\|_{L^2(S^{m-1})}.$$

Notice that the integrands of both $I_a(s)$ and $II_a(s)$ involve $\partial_t u$ and its derivatives, which are estimated in Section 2. More precisely, by taking σ small, we may apply Lemma 2.5 first to get (14) and then Lemma 2.7 to see

$$(43) \quad \|I_a(s)\|_{L^2(S^{m-1})}^2 + \int_{S^{m-1}} |II_a(s)| d\theta \leq C \int_{s-1}^{s+1} \int_{S^{m-1}} |II_b|^2 d\theta dt.$$

By the definition of $G(t)$ in (33), equations (39), (42), and (43) imply

$$-C(G(s-1) - G(s+1)) + G(s) \leq C(G(s-1) - G(s+1))^{1/(2-\alpha)}.$$

Since $G(s-1) - G(s+1)$ is bounded and $1/(2-\alpha) \leq 1$, the first term can be absorbed into the left-hand side. In fact, in the proof that follows, we shall require $G(s)$ to be very small (see the definition of η in the next section). By the monotonicity of G , the above inequality is further simplified to

$$(44) \quad G(s+1) \leq C(G(s-1) - G(s+1))^{1/(2-\alpha)}.$$

Here is a lemma similar to [Simon 1996, (9), §3.15].

Lemma 4.2. *Suppose that $\theta \in (0, \frac{1}{2}]$. If for some positive C and any $a, b \in (0, 1)$ satisfying $b < a$,*

$$(45) \quad b \leq C(a-b)^{1/(2-2\theta)},$$

then there is another C' depending only on C and θ such that

$$a^\theta - b^\theta \geq C'(a-b)^{1/2}.$$

Proof. The proof is an elementary discussion.

Case 1: $b < \frac{1}{2}a$. Noticing that $\theta \leq \frac{1}{2}$ and $a < 1$, we have

$$a^\theta - b^\theta \geq \left(1 - \frac{1}{2^\theta}\right)a^\theta \geq \left(1 - \frac{1}{2^\theta}\right)a^{1/2} \geq \left(1 - \frac{1}{2^\theta}\right)(a - b)^{1/2}.$$

Case 2: $b \geq \frac{1}{2}a$. Equation (45) gives

$$\frac{a}{2C} \leq (a - b)^{1/(2(1-\theta))},$$

which is

$$(46) \quad a^{1-\theta} \leq (2C)^{1-\theta}(a - b)^{1/2}.$$

Therefore,

$$a^\theta - b^\theta \geq \theta a^{\theta-1}(a - b) \geq \frac{\theta}{(2C)^{1-\theta}}(a - b)^{1/2}.$$

Here in the above line we have used the mean value theorem for the first inequality and (46) for the second.

In either case, the lemma is proved by taking C' to be $\min\{1 - 1/2^\theta, \theta/(2C)^{1-\theta}\}$.

□

5. A stability argument and the proof of Theorem 1.1

In this section, we prove Theorem 1.1 by using a routine stability argument. We shall define two neighborhoods of φ : a larger one (see $\mathcal{O}_{C^5}(\sigma_1)$ below) in which the results in Section 3 and Section 4 hold and a smaller one (see $\mathcal{O}_{L^2}(\eta)$ below) such that if $u(t_i)$ lies in the smaller neighborhood for sufficiently large i , then $u(t)$ will stay in the larger neighborhood forever and converge to the unique limit claimed in Theorem 1.1.

We choose σ_1 so that it is smaller than both the ε in Lemma 3.1 and the σ in Lemma 4.1. For some $\eta > 0$ small (to be determined later), by the definition of φ as the section of a tangent map, we can choose (and fix) t_i large such that

- (1) for all $t \in (t_i - 3, t_i + 3)$, $u(t) \in \mathcal{O}_{C^5}(\sigma_1)$;
- (2) $u(t_i) \in \mathcal{O}_{L^2}(\eta)$;
- (3) $G(t_i) \leq \eta^2$, because $G(t)$ is finite and decreases down to zero.

Set

$$T = \sup_t \{t \mid \text{for any } s \in [t_i, t), u(s) \in \mathcal{O}_{C^5}(\sigma_1)\}.$$

By (1) above, we know $T \geq t_i + 3$. Now we claim that T is infinity. If otherwise, we want to find a contradiction by showing $u(T) \in \mathcal{O}_{C^5}(\frac{1}{2}\sigma_1)$. Thanks to Lemma 2.5,

there is $\sigma_2 > 0$ depending on $\frac{1}{2}\sigma_1$ such that it suffices to show for any $s \in (t_i, T + 2)$, we have $u(s) \in \mathcal{O}_{L^2}(\sigma_2)$. Let k be the largest integer with $t_i + 2k \leq s$. Hence,

$$\begin{aligned}
 & \int_{t_i}^s \|\partial_t u\|_{L^2(S^{m-1})} \\
 & \leq \sum_{j=1}^k \int_{t_i+2(j-1)}^{t_i+2j} \|\partial_t u\|_{L^2(S^{m-1})} + \int_{t_i+2k}^s \|\partial_t u\|_{L^2(S^{m-1})} \\
 & \leq C \sum_{j=1}^k \left(\int_{t_i+2(j-1)}^{t_i+2j} \|\partial_t u\|_{L^2(S^{m-1})}^2 \right)^{1/2} + C \left(\int_{t_i+2k}^s \|\partial_t u\|_{L^2(S^{m-1})}^2 \right)^{1/2} \\
 & \leq C \sum_{j=1}^k \left(\int_{t_i+2(j-1)}^{t_i+2j} \|\partial_t u\|_{L^2(S^{m-1})}^2 \right)^{1/2} + C\eta.
 \end{aligned}$$

Here in the third line above, we used Hölder inequality and in the last line, we used (3).

By the definition of G , we have

$$\int_{t_i+2(j-1)}^{t_i+2j} \|\partial_t u\|_{L^2(S^{m-1})}^2 \leq G(t_i + 2(j-1)) - G(t_i + 2j).$$

We can apply Lemma 4.2 with $a = G(t_i + 2j)$ and $b = G(t_i + 2(j-1))$ to get

$$\begin{aligned}
 \int_{t_i}^s \|\partial_t u\|_{L^2(S^{m-1})} & \leq C \sum_{j=1}^k (G(t_i + 2(j-1))^\theta - G(t_i + 2j)^\theta) + C\eta \\
 & \leq C \cdot G(t_i)^\theta + C\eta \leq C\eta^{2\theta} + C\eta.
 \end{aligned}$$

If we choose η small, we can have for any $s \in (t_i, T + 2)$,

$$\|u(s) - \varphi\|_{L^2(S^{m-1})} \leq \|u(t_i) - \varphi\|_{L^2(S^{m-1})} + \int_{t_i}^s \|\partial_t u\|_{L^2(S^{m-1})} \leq \frac{1}{2}\sigma_2.$$

Lemma 2.5 gives the contradiction and proves that $T = \infty$.

We can repeat the above computation with $k = \infty$ to get

$$\int_{t_i}^{+\infty} \|\partial_t u\|_{L^2(S^{m-1})} \leq C\eta^{2\theta} + C\eta < \infty,$$

which shows that

$$\lim_{t \rightarrow \infty} \|u(t) - \varphi\|_{L^2(S^{m-1})} = 0.$$

As in Remark 2.6, we have u bounded in any $C^{k+1}(S^{m-1})$ norm. By interpolation, we know

$$\lim_{t \rightarrow \infty} \|u(t) - \varphi\|_{C^k(S^{m-1})} = 0.$$

Appendix A: The assumption of analyticity

The purpose of this section is to justify (see Lemma A.7) the use of the classical Łojasiewicz inequality to the function f (see (23)) that arises in the Lyapunov–Schmidt reduction in Section 3. Indeed, we shall show how the analyticity assumption of N in Theorem 1.1 carries on step by step to that of f . These arguments, independent from the rest of the proof, are technical and hence presented in the appendix.

Analytic function between Banach spaces. For completeness, we collect a few basic definitions and properties of analytic functions between abstract (complex) Banach spaces. We refer to [Taylor 1937] for proofs and more detailed discussions.

Let E , E' and E'' be complex Banach spaces.

Definition A.1. (1) Let $f(x)$ be a function on E to E' , defined in the neighborhood of $x_0 \in E$. If for each $y \in E$, the limit

$$\lim_{\tau \rightarrow 0} \frac{f(x_0 + \tau y) - f(x_0)}{\tau}$$

exists (for $\tau \in \mathbb{C}$), then it is called the *Gateaux* differential, denoted by $\delta f(x_0; y)$.

(2) A function $f(x)$ on a domain D of E to E' is said to be *analytic* in D if it is continuous and has a Gateaux differential at each point of D . A function is said to be analytic at a point x_0 , if it is analytic in some neighborhood of x_0 .

Recall that the *Fréchet* differential is defined to be the bounded linear map $Df(x_0)$ from E to E' such that

$$f(x_0 + h) = f(x_0) + Df(x_0)h + o(\|h\|_E).$$

While the existence of the Fréchet differential is obviously stronger than the Gateaux differential, Taylor proved the following:

Theorem A.2 [Taylor 1937, Theorem 3, Theorem 12]. *If f is analytic at x_0 , then it admits Fréchet differentials of all orders in the neighborhood of that point. Moreover, the Fréchet differential and the Gateaux differential are equal.*

With the equivalence in mind, we recall a version of the inverse function theorem, which follows from [Dieudonné 1960, (10.2.5)] (see also [Nirenberg 1974, §2.7]).

Theorem A.3. *Let E and E' be two complex Banach spaces, f an analytic function from a neighborhood V of $x_0 \in E$ to E' . If $Df(x_0)$ is a linear homeomorphism of E onto E' , there exists an open neighborhood $U \subset V$ of x_0 such that the restriction of f to U is a homeomorphism of U onto an open neighborhood of $y_0 = f(x_0)$. Moreover, the inverse is analytic.*

Complexification and analyticity. In Section 3, we have defined the functional $\tilde{F} : C^{4,\beta}(V) \rightarrow \mathbb{R}$ where V is the pullback bundle φ^*TN and a map $\mathcal{N}$ from $C^{4,\beta}(V)$ to $C^{0,\beta}(V)$. Instead of claiming the analyticity of $\tilde{F}$ and $\mathcal{N}$ directly, we consider its complexification.

$C^{4,\beta}(V) \otimes \mathbb{C}$ is understood to be the set of $u + iv$, where $u, v \in C^{4,\beta}(V)$, with a naturally defined norm. The same applies to $C^{0,\beta}(V) \otimes \mathbb{C}$. Obviously, they are complex Banach spaces.

A complexification of a map f from a Banach space E_1 to another Banach space E_2 is some map $\tilde{f}$ from $E_1 \otimes \mathbb{C}$ to $E_2 \otimes \mathbb{C}$ such that f is the real part of $\tilde{f}$ when restricted to (some open set of) E_1 . Such complexifications are usually not unique. We are interested in analytic ones, that we define below (making use of special properties of f).

The complexification of $\tilde{F}$ and $\mathcal{N}$ relies on some particular form of the maps themselves. More precisely, we need the definition of $\tilde{F}(u)$ and $\mathcal{N}(u)$ to be given by a converging series. For this purpose, we start with an extrinsic point of view of V .

Since N is embedded in $\mathbb{R}^p$, we regard $T_y N$ as a subspace (not the affine space passing y) of $\mathbb{R}^p$. Hence, the pullback bundle V is the disjoint union of $V_\omega := T_{\varphi(\omega)} N$ and a section u of V is a map from S^{m-1} to $\mathbb{R}^p$ satisfying

$$u(\omega) \in T_{\varphi(\omega)} N \subset \mathbb{R}^p.$$

For a fixed smooth φ , the $C^{k,\beta}$ norm of u as a map into $\mathbb{R}^p$ agrees with the $C^{k,\beta}$ norm defined intrinsically using the pullback connection of φ^*TN . The same holds for various Sobolev norms.

For the complexification of $\tilde{F}$, we regard it as the composition of

$$C^{2,\beta}(V) \xrightarrow{\mathcal{F}} C^\beta(S^{m-1}, \mathbb{R}) \xrightarrow{\mathcal{I}} \mathbb{R},$$

where

$$\mathcal{F}(u) = |\Delta_{S^{m-1}} \Pi(\varphi + u)|^2 + (2m - 8) |\nabla_{S^{m-1}} \Pi(\varphi + u)|^2$$

and

$$\mathcal{I}(h) = \int_{S^{m-1}} h \, d\theta.$$

Recall that Π is the nearest-point-projection of N and the discussion works only for u with small C^0 norm.

We claim that there exists an analytic map $\tilde{F}_C$ from $C^{2,\beta}(V) \otimes \mathbb{C}$ to $\mathbb{C}$ with $\tilde{F}$ as its real part.

The proof of the claim is the combination of the following facts:

- (F1) The $\Delta_{S^{m-1}}$ from $C^{2,\beta}(V)$ to $C^{0,\beta}(V)$, $\nabla_{S^{m-1}}$ from $C^{2,\beta}(V)$ to $C^{1,\beta}(V)$, and $\mathcal{I}$ are bounded linear maps. Their complexifications, obtained by linear extension, are naturally bounded linear maps and hence analytic.

(F2) Let $\mathcal{F}_1$ be the map from $C^{0,\beta}(S^{m-1}, \mathbb{R}^p)$ to $C^{0,\beta}(S^{m-1}, \mathbb{R})$ given by $u \mapsto |u|^2$. Its complexification $\mathcal{F}_{C,1}$ is given by

$$\mathcal{F}_{C,1}(u + iv) = (u + iv) \cdot (u + iv).$$

It is analytic.

(F3) If (as assumed in Theorem 1.1) $\Pi(\varphi + \cdot)$ is an analytic map from $B_r(0) \subset \mathbb{R}^p$ to $\mathbb{R}^p$, then the map

$$u \mapsto \Pi(\varphi + u)$$

has an analytic extension from $C^{2,\beta}(S^{m-1}, \mathbb{C}^p)$ to itself. To see this, one first expands $\Pi(\varphi + u)$ into converging power series of u and then replace u by $u + iv$. It is then an exercise to check that the map thus obtained are analytic in the sense of Definition A.1.

For the complexification of $\mathcal{N}$, it suffices to consider $\mathcal{M}_{\tilde{F}}(u)$. For u and v in $C^4(V)$, setting $\psi = \Pi(\varphi + u)$, we compute

$$\begin{aligned} & \left. \frac{d}{dt} \right|_{t=0} \tilde{F}(u + tv) \\ &= \left. \frac{d}{dt} \right|_{t=0} \int_{S^{m-1}} |\Delta_{S^{m-1}} \Pi(\varphi + u + tv)|^2 \\ & \quad + (2m - 8) |\nabla_{S^{m-1}} \Pi(\varphi + u + tv)|^2 d\theta \\ &= 2 \int_{S^{m-1}} \Delta_{S^{m-1}} \psi \Delta_{S^{m-1}} D\Pi_{\varphi+u} v + (2m - 8) \nabla_{S^{m-1}} \psi \nabla_{S^{m-1}} D\Pi_{\varphi+u} v d\theta \\ &= 2 \int_{S^{m-1}} (\Delta_{S^{m-1}}^2 \psi - (2m - 8) \Delta_{S^{m-1}} \psi) D\Pi_{\varphi+u} v d\theta \\ &= 2 \int_{S^{m-1}} P_\psi (\Delta_{S^{m-1}}^2 \psi - (2m - 8) \Delta_{S^{m-1}} \psi) v d\theta. \end{aligned}$$

Here in the last line above, we used the fact that $D\Pi_{\varphi+u} v$ is nothing but the orthogonal projection from $\mathbb{R}^p$ onto $T_\psi N$, which we denote by P_ψ .

Similar to the (bi)harmonic map case, $P_\psi (\Delta_{S^{m-1}}^2 \psi - (2m - 8) \Delta_{S^{m-1}} \psi)$ is the Euler–Lagrange operator of $F(\psi)$, denoted by $\mathcal{M}_F(\psi)$. For each $\omega \in S^{m-1}$, $\mathcal{M}_F(\psi)(\omega)$ lies in $T_\psi N \subset \mathbb{R}^p$, while $v(\omega)$ is in $T_\varphi N$. Therefore,

$$(47) \quad \mathcal{M}_{\tilde{F}}(u) = P_\varphi \mathcal{M}_F(\psi),$$

where $\psi = \Pi(\varphi + u)$.

Since the projection P_φ is a linear map that does not depend on u , the complexification of $\mathcal{M}_{\tilde{F}}(u)$ is reduced to that of $\mathcal{M}_F(\Pi(\varphi + u))$, which we regard as the composition of the following:

(M1) the map $\Pi(\varphi + \cdot)$ from $C^{4,\beta}(V)$ to $C^{4,\beta}(S^{m-1}, \mathbb{R}^p)$, which has been discussed in (F3) above;

- (M2) the map $\Delta_{S^{m-1}}^2 \psi - (2m-8) \Delta_{S^{m-1}} \psi$ from $C^{4,\beta}(S^{m-1}, \mathbb{R}^p)$ to $C^{0,\beta}(S^{m-1}, \mathbb{R}^p)$, which has been discussed in (F1) above;
- (M3) the projection P_ψ is a p by p matrix that depends analytically on ψ , since N is an analytic submanifold. Keeping in mind that $\psi = \Pi(\varphi + u)$ is known (see (M1) above) to be analytic map in u , the complexification of P_ψ is given by expanding the analytic (matrix-valued) map $P_\psi = P_{\Pi(\varphi+u)}$ as a converging power series of u and then replacing u by $u + iv$ as in (F3).

Properties of the complexification. Let's denote the complexification of $\mathcal{M}_{\tilde{F}}$ by $\mathcal{M}_{\tilde{F},C}$. In this section, we study the ellipticity of $\mathcal{M}_{\tilde{F},C}$ and the self-adjointness of its linearization at 0. Please notice that although the ellipticity of $\mathcal{M}_{\tilde{F}}$ is quite natural, the ellipticity of $\mathcal{M}_{\tilde{F},C}$ as an operator between the complexified Banach spaces is not true in general. Fortunately, we have the following:

Lemma A.4. *The linearizations of both $\mathcal{M}_{\tilde{F}}$ and $\mathcal{M}_{\tilde{F},C}$ at $u = 0$ are elliptic.*

Remark A.5. In fact, as the following proof shows, $\mathcal{M}_{\tilde{F}}$ is elliptic for small u such that it is defined and $\mathcal{M}_{\tilde{F},C}$ is elliptic at $u + iv \in C^{4,\beta}(V) \otimes \mathbb{C}$ if $v = 0$.

Proof. Neglecting the lower order part, it suffices to compute the linearization of

$$P_\varphi P_\psi \Delta_{S^{m-1}}^2 \Pi(\varphi + u),$$

where $\psi = \Pi(\varphi + u)$. If we do the computation at $u \in C^{4,\beta}(V)$ with infinitesimal increment h and neglect all lower order terms, we get

$$(48) \quad P_\varphi P_\psi \Delta_{S^{m-1}}^2 h,$$

whose symbol is for any $\xi \in T_\omega^* S^{m-1}$,

$$(49) \quad \xi \mapsto P_\varphi P_\psi |\xi|^4 h.$$

If ξ is not zero, then this is clearly a linear isomorphism from the sections of V onto itself, because ψ is close to φ .

Now, for $\mathcal{M}_{\tilde{F},C}$, we denote the complexification of P_ψ ($\Pi(\varphi + u)$) by $P_{\psi,C}$ ($\Pi_C(\varphi + u)$ respectively). Although we do not know any exact formula for them, it suffices for us to note that when computing (48),

- (1) the contribution of Π_C goes to the lower order terms and does not matter;
- (2) since we have assumed that $u \in C^{4,\beta}(V)$, by the definition of complexification, $P_{\psi,C} = P_\psi$. Therefore, we get the same symbol as in (49), which is now an isomorphism from the sections of complexified- V onto itself. $\square$

If we denote the linearizations of $\mathcal{M}_{\tilde{F}}$ and $\mathcal{M}_{\tilde{F},C}$ at $u = 0$ by $\mathcal{L}_{\tilde{F}}$ and $\mathcal{L}_{\tilde{F},C}$, then

Lemma A.6. *For any $u, v \in C^{4,\beta}(V)$,*

$$(50) \quad \mathcal{L}_{\tilde{F},C}(u + iv) = \mathcal{L}_{\tilde{F}}(u) + i\mathcal{L}_{\tilde{F}}(v).$$

In particular, $\mathcal{L}_{\tilde{F},C}$ is an elliptic and self-adjoint operator from $C^{4,\beta}(V) \otimes \mathbb{C}$ to $C^{0,\beta}(V) \otimes \mathbb{C}$.

Proof. By definition, $\mathcal{L}_{\tilde{F},C}(u) = d/dt|_{t=0}\mathcal{M}_{\tilde{F},C}(tu) = d/dt|_{t=0}\mathcal{M}_{\tilde{F}}(tu) = \mathcal{L}_{\tilde{F}}(u)$. Hence, it suffices to show

$$\tilde{\mathcal{L}}_{\tilde{F},C}(iv) = i\mathcal{L}_{\tilde{F}}(v).$$

Since $\mathcal{M}_{\tilde{F}}$ is a composition of P_φ , P_ψ , $\Delta_{S^{m-1}}$, $\nabla_{S^{m-1}}$, and $\Pi(\varphi + \cdot)$, it suffices to show that (50) holds for (the linearization of) each one of them. This is trivial for P_φ , $\Delta_{S^{m-1}}$, and $\nabla_{S^{m-1}}$, because they are linear operators and (50) is exactly how their complexification is defined.

For $\Pi(\varphi + \cdot)$, we recall that

$$\Pi_C(\varphi + (u + iv)) = \sum_k a_k(u + iv)^k$$

and the series converges for small u and v . Equation (50) then follows from direct computation. The same argument works for P_ψ .

The self-adjointness of $\mathcal{L}_{\tilde{F}}$ follows from expanding the following identity:

$$\left. \frac{d}{ds} \right|_{s=0} \left. \frac{d}{dt} \right|_{t=0} \tilde{F}(tu + sv) = \left. \frac{d}{dt} \right|_{t=0} \left. \frac{d}{ds} \right|_{s=0} \tilde{F}(tu + sv).$$

The self-adjointness of $\mathcal{L}_{\tilde{F},C}$ is then a consequence of (50). $\square$

Now, we state the result that motivates the discussion in this section.

Lemma A.7. *For f defined (23), it is an analytic function of ξ in a neighborhood of $0 \in \mathbb{R}^l$.*

Proof. Let $\mathcal{N}_C$ be the complexification of $\mathcal{N}$ defined in Section A. Its linearization at $u = 0$ is given by

$$P_K + \tilde{\mathcal{L}}.$$

By the results above, this $\tilde{\mathcal{L}}$ is elliptic and self-adjoint with trivial kernel. Hence, the inverse function theorem (Theorem A.3) gives an inverse map Ψ_C , which is analytic, from a neighborhood of 0 in $C^{0,\beta}(V) \otimes \mathbb{C}$ to a neighborhood of 0 in $C^{4,\beta}(V) \otimes \mathbb{C}$. If $\tilde{F}_C$ is the complexification of $\tilde{F}$ given in Section A, then f in (23) is the restriction (to the real part of $(z_1, \dots, z_l)$) of

$$f_C(z_1, \dots, z_l) := \tilde{F}_C \left(\Psi_C \left(\sum_{i=1}^l z_i \varphi_i \right) \right),$$

which is analytic in (a neighborhood of 0 in) $\mathbb{C}^l$. $\square$

Appendix B: Proof of Lemma 3.5

For some $\delta > 0$ to be determined, we will take $W = \{f \mid \|f\|_{C^\beta(V)} < \delta\}$. For any f_1, f_2 in W , we have

$$\Psi(f_1) - \Psi(f_2) = \int_0^1 \frac{d}{dt} \Psi(tf_1 + (1-t)f_2) dt = \int_0^1 D\Psi|_{tf_1+(1-t)f_2}(f_1 - f_2) dt.$$

Hence, it suffices to show that for any $f \in W$, the linearization of Ψ at f , $D\Psi|_f$ is a uniformly bounded linear operator from $L^2(V)$ to $W^{4,2}(V)$. More precisely, we need to find $\delta > 0$ and $C > 0$ such that

$$\sup_{f \in W} \|D\Psi|_f\|_{L(L^2(V), W^{4,2}(V))} \leq C.$$

Here $\|\cdot\|_{L(L^2(V), W^{4,2}(V))}$ is the norm of linear operators.

Since Ψ is the inverse of $\mathcal{N}$, it suffices to show that there exist $\delta' > 0$ and $C > 0$ such that if $W' = \{u \in C^{4,\beta}(V) \mid \|u\|_{C^{4,\beta}(V)} < \delta'\}$,

$$(51) \quad \inf_{u \in W'} \|D\mathcal{N}|_u\|_{L(W^{4,2}, L^2)} \geq C > 0.$$

The proof of (51) consists of two steps. First, we show that

$$(52) \quad \|D\mathcal{N}|_0\|_{L(W^{4,2}, L^2)} \geq C > 0.$$

Recall that $\mathcal{N} = P_K + \mathcal{M}_{\tilde{F}}$, where K is the kernel of $D\mathcal{M}_{\tilde{F}}|_0 = \mathcal{L}_{\tilde{F}}$. For any $h \in W^{4,2}(V)$, we denote $h - P_K h$ by $h^\perp$. Since $\mathcal{L}_{\tilde{F}}$ is an elliptic operator with trivial kernel in the complement space of K , there is a constant depending only on φ such that

$$(53) \quad \|h^\perp\|_{W^{4,2}} \leq C_0 \|\mathcal{L}_{\tilde{F}} h^\perp\|_{L^2}.$$

Since K is a finite-dimensional space, there is $C_1 > 0$ such that

$$(54) \quad \|P_K h\|_{W^{4,2}} \leq C_1 \|P_K h\|_{L^2}.$$

Combining (53) and (54) and noticing that the image of $\mathcal{L}_{\tilde{F}}$ is normal to K in L^2 , we get $C_2 > 0$ such that

$$(55) \quad \|h\|_{W^{4,2}} \leq C_2 \|D\mathcal{N}|_0 h\|_{L^2},$$

which implies (52).

The second step is to show that for $u \in W'$,

$$(56) \quad \|(D\mathcal{N}|_u - D\mathcal{N}|_0)h\|_{L^2} \leq C(\delta') \|h\|_{W^{4,2}}$$

for some $C(\delta')$ satisfying $\lim_{\delta' \rightarrow 0} C(\delta') = 0$. Before the proof of (56), we notice that if δ' is small, (56) and (55) imply that

$$\|h\|_{W^{4,2}} \leq C \|D\mathcal{N}|_u h\|_{L^2},$$

which finishes the proof of (51) and hence the proof of Lemma 3.5.

For (56), we notice that the contribution of $P_K h$ cancels out and it suffices to bound

$$(57) \quad \|(D\mathcal{M}_{\tilde{F}}|_u - D\mathcal{M}_{\tilde{F}}|_0)h\|_{L^2}.$$

Recalling that $\mathcal{M}_{\tilde{F}}(u) = P_\varphi P_\psi (\Delta_{S^{m-1}}^2 \psi - (2m-8)\Delta_{S^{m-1}} \psi)$ with $\psi = \Pi(\varphi + u)$, we get

$$(58) \quad D\mathcal{M}_{\tilde{F}}|_u h = P_\varphi (DP)_\psi (\Delta_{S^{m-1}}^2 \psi - (2m-8)\Delta_{S^{m-1}} \psi) h \\ + P_\varphi P_\psi (\Delta_{S^{m-1}}^2 h - (2m-8)\Delta_{S^{m-1}} h)$$

and

$$(59) \quad D\mathcal{M}_{\tilde{F}}|_0 h = P_\varphi (DP)_\varphi (\Delta_{S^{m-1}}^2 \varphi - (2m-8)\Delta_{S^{m-1}} \varphi) h \\ + P_\varphi (\Delta_{S^{m-1}}^2 h - (2m-8)\Delta_{S^{m-1}} h).$$

Notice that (58) and (59) are fourth-order linear operators of h and if we subtract them, the difference of the corresponding coefficients are bounded by using

$$\|\psi - \varphi\|_{C^{4,\beta}} \leq C \|u\|_{C^{4,\beta}(V)}.$$

References

- [Chang et al. 1999] S.-Y. A. Chang, L. Wang, and P. C. Yang, “A regularity theory of biharmonic maps”, *Comm. Pure Appl. Math.* **52**:9 (1999), 1113–1137. MR Zbl
- [Cheeger and Tian 1994] J. Cheeger and G. Tian, “On the cone structure at infinity of Ricci flat manifolds with Euclidean volume growth and quadratic curvature decay”, *Invent. Math.* **118**:3 (1994), 493–571. MR Zbl
- [Colding and Minicozzi 2014] T. H. Colding and W. P. Minicozzi, II, “On uniqueness of tangent cones for Einstein manifolds”, *Invent. Math.* **196**:3 (2014), 515–588. MR Zbl
- [Dieudonné 1960] J. Dieudonné, *Foundations of modern analysis*, Pure and Applied Math. **10**, Academic Press, New York, 1960. MR Zbl
- [Hong and Wang 2005] M.-C. Hong and C. Wang, “Regularity and relaxed problems of minimizing biharmonic maps into spheres”, *Calc. Var. Partial Differential Equations* **23**:4 (2005), 425–450. MR Zbl
- [Lamm and Rivière 2008] T. Lamm and T. Rivière, “Conservation laws for fourth order systems in four dimensions”, *Comm. Partial Differential Equations* **33**:1-3 (2008), 245–262. MR Zbl
- [Lin 1999] F.-H. Lin, “Gradient estimates and blow-up analysis for stationary harmonic maps”, *Ann. of Math.* (2) **149**:3 (1999), 785–829. MR Zbl
- [Luckhaus 1988] S. Luckhaus, “Partial Hölder continuity for minima of certain energies among maps into a Riemannian manifold”, *Indiana Univ. Math. J.* **37**:2 (1988), 349–367. MR Zbl

- [Nirenberg 1974] L. Nirenberg, *Topics in nonlinear functional analysis*, Courant Inst. Math. Sci., New York, 1974. MR Zbl
- [Scheven 2008] C. Scheven, “Dimension reduction for the singular set of biharmonic maps”, *Adv. Calc. Var.* **1**:1 (2008), 53–91. MR Zbl
- [Schoen and Uhlenbeck 1982] R. Schoen and K. Uhlenbeck, “A regularity theory for harmonic maps”, *J. Differential Geom.* **17**:2 (1982), 307–335. Correction in **18**:2 (1983), 329. MR Zbl
- [Simon 1983] L. Simon, “Asymptotics for a class of nonlinear evolution equations, with applications to geometric problems”, *Ann. of Math. (2)* **118**:3 (1983), 525–571. MR Zbl
- [Simon 1996] L. Simon, *Theorems on regularity and singularity of energy minimizing maps*, Birkhäuser, Basel, 1996. MR Zbl
- [Struwe 2008] M. Struwe, “Partial regularity for biharmonic maps, revisited”, *Calc. Var. Partial Differential Equations* **33**:2 (2008), 249–262. MR Zbl
- [Strzelecki 2003] P. Strzelecki, “On biharmonic maps and their generalizations”, *Calc. Var. Partial Differential Equations* **18**:4 (2003), 401–432. MR Zbl
- [Taylor 1937] A. E. Taylor, “Analytic functions in general analysis”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (2)* **6**:3-4 (1937), 277–292. MR Zbl
- [Wang 2004a] C. Wang, “Stationary biharmonic maps from $\mathbb{R}^n$ into a Riemannian manifold”, *Comm. Pure Appl. Math.* **57**:4 (2004), 419–444. MR Zbl
- [Wang 2004b] C. Wang, “Biharmonic maps from $\mathbb{R}^4$ into a Riemannian manifold”, *Math. Z.* **247**:1 (2004), 65–87. MR Zbl
- [Wang 2004c] C. Wang, “Remarks on biharmonic maps into spheres”, *Calc. Var. Partial Differential Equations* **21**:3 (2004), 221–242. MR Zbl
- [Yang 2003] B. Yang, “The uniqueness of tangent cones for Yang–Mills connections with isolated singularities”, *Adv. Math.* **180**:2 (2003), 648–691. MR Zbl

Received May 5, 2017. Revised May 15, 2018.

YOU MIN CHEN
SCHOOL OF MATHEMATICAL SCIENCES
UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA
HEFEI
CHINA
chym189@mail.ustc.edu.cn

HAO YIN
SCHOOL OF MATHEMATICAL SCIENCES
UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA
HEFEI
CHINA
haoyin@ustc.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2019 is US \$490/year for the electronic version, and \$665/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 299 No. 2 April 2019

- Hierarchically hyperbolic spaces II: Combination theorems and the distance formula 257
JASON BEHRSTOCK, MARK HAGEN and ALESSANDRO SISTO
- The weighted σ_k -curvature of a smooth metric measure space 339
JEFFREY S. CASE
- Uniqueness of tangent cones for biharmonic maps with isolated singularities 401
YOU MIN CHEN and HAO YIN
- Hamiltonian unknottedness of certain monotone Lagrangian tori in $S^2 \times S^2$ 427
KAI CIELIEBAK and MARTIN SCHWINGENHEUER
- Chord shortening flow and a theorem of Lusternik and Schnirelmann 469
MARTIN MAN-CHUN LI
- Rigidity theorems of hypersurfaces with free boundary in a wedge in a space form 489
JUNCHEOL PYO



0030-8730(201904)299:2;1-N