

Pacific Journal of Mathematics

**SYMMETRY AND MONOTONICITY OF
POSITIVE SOLUTIONS FOR AN INTEGRAL SYSTEM
WITH NEGATIVE EXPONENTS**

ZHAO LIU

SYMMETRY AND MONOTONICITY OF POSITIVE SOLUTIONS FOR AN INTEGRAL SYSTEM WITH NEGATIVE EXPONENTS

ZHAO LIU

We study the following integral system with negative exponents:

$$\begin{cases} u(x) = \int_{\mathbb{R}^n} |x - y|^{\alpha-n} v^{-p}(y) dy, \\ v(x) = \int_{\mathbb{R}^n} |x - y|^{\alpha-n} u^{-q}(y) dy, \end{cases}$$

with $\alpha > n$, $p, q > 0$ and $\frac{1}{p-1} + \frac{1}{q-1} = \frac{\alpha-n}{n}$. Such a nonlinear integral system is related to the study of the best constant of the reversed Hardy–Littlewood–Sobolev type inequality. Motivated by work of Dou, Guo and Zhu (*Adv. Math.* **312** (2017) 1–45) where they used the improved method of moving planes, we prove that each pair of positive measurable solutions is radially symmetric and monotonic increasing about some point. Our result is an extension of the work for $\alpha < n$, $p, q < 0$ of Chen, Li and Ou (*Comm. Partial Differential Equations* **30**:1–3 (2005), 59–65).

1. Introduction

Let $0 < \gamma < n$, and let $s_1, s_2 > 1$ such that

$$\frac{1}{s_1} + \frac{1}{s_2} = \frac{n + \gamma}{n}.$$

The well-known Hardy–Littlewood–Sobolev inequality states that

$$(1) \quad \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} f(x) |x - y|^{\gamma-n} g(y) dx dy \leq C(n, \gamma, s_1) \|f\|_{L^{s_1}(\mathbb{R}^n)} \|g\|_{L^{s_2}(\mathbb{R}^n)},$$

for any $f \in L^{s_1}(\mathbb{R}^n)$ and $g \in L^{s_2}(\mathbb{R}^n)$.

It is well known that the Hardy–Littlewood–Sobolev inequality has many important applications in partial differential equations and geometry, as well as in quantum field theory. It is also known that the Hardy–Littlewood–Sobolev inequality is closely related to the Sobolev inequality in Euclidean spaces and Moser–Onofri–Beckner type inequalities on spheres (see Beckner [1993; 2008]). For more research

The author was supported by the NSFS (No. 11801237).

MSC2010: 45G15, 46E30.

Keywords: Hardy–Littlewood–Sobolev inequality, method of moving plane, negative exponent.

results with integral equations related to the Hardy–Littlewood–Sobolev inequality, we refer the reader to [Beckner 1993; Brascamp and Lieb 1976; Chen et al. 2005a; 2016; Chen and Li 2008; Lu and Zhu 2011]. For Hardy–Littlewood–Sobolev inequalities on the Riemannian manifolds and the upper half space $\mathbb{R}_+^n$, see, e.g., [Dou 2016; Dou and Zhu 2015b; Han and Zhu 2016; Ngô and Nguyen 2017b].

To find the best constant $C(n, \gamma, s_1)$ in (1), we can maximize the functional

$$(2) \quad J(f, g) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} f(x) |x - y|^{\gamma-n} g(y) dx dy,$$

under the constraints

$$(3) \quad \|f\|_{L^{s_1}(\mathbb{R}^n)} = \|g\|_{L^{s_2}(\mathbb{R}^n)} = 1.$$

Let (f, g) be a minimizer, or more generally a critical point of (2) under the constraints (3). Letting $u = a_1 f^{s_1-1}$, $v = a_2 g^{s_2-1}$, $p_1 = \frac{1}{s_1-1}$, $p_2 = \frac{1}{s_2-1}$, and by a proper choice of constants a_1 and a_2 , we can see that (u, v) satisfies the following integral system in $\mathbb{R}^n$:

$$(4) \quad \begin{cases} u(x) = \int_{\mathbb{R}^n} |x - y|^{\gamma-n} v^{p_2}(y) dy, \\ v(x) = \int_{\mathbb{R}^n} |x - y|^{\gamma-n} u^{p_1}(y) dy, \end{cases}$$

with $\frac{1}{p_1+1} + \frac{1}{p_2+1} = \frac{n-\gamma}{n}$.

Under the natural integrability conditions $u \in L^{p_1}(\mathbb{R}^n)$ and $v \in L^{p_2}(\mathbb{R}^n)$, Chen, Li and Ou [Chen et al. 2005b] proved that all the solutions are radially symmetric and monotonic decreasing about some point.

Suppose that $\alpha > n$, and $0 < r_1, r_2 < 1$ satisfying

$$\frac{1}{r_1} + \frac{1}{r_2} = \frac{n+\alpha}{n}.$$

Dou and Zhu [2015a] established the following reversed Hardy–Littlewood–Sobolev inequality (see also Ngô and Nguyen [2017a]), which can be seen as an extension of (1), which states

$$(5) \quad \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} f(x) |x - y|^{\alpha-n} g(y) dx dy \geq C(n, \alpha, r_1) \|f\|_{L^{r_1}(\mathbb{R}^n)} \|g\|_{L^{r_2}(\mathbb{R}^n)},$$

for any $f \in L^{r_1}(\mathbb{R}^n)$ and $g \in L^{r_2}(\mathbb{R}^n)$.

The corresponding Euler–Lagrange equation becomes

$$(6) \quad \begin{cases} u(x) = \int_{\mathbb{R}^n} |x - y|^{\alpha-n} v^{-p}(y) dy, \\ v(x) = \int_{\mathbb{R}^n} |x - y|^{\alpha-n} u^{-q}(y) dy, \end{cases}$$

with $\alpha > n$, $p, q > 0$ and,

$$(7) \quad \frac{1}{p-1} + \frac{1}{q-1} = \frac{\alpha-n}{n}.$$

The study of system (6), especially the classification result, is a crucial step in finding the best constant in the reversed Hardy–Littlewood–Sobolev inequality. Dou and Zhu [2015a] classified the positive solutions of (6) with critical exponents $p = q = \frac{n+\alpha}{\alpha-n}$ and gave its best constant. Recently, Lei [2015] considered system (6) and obtained that assumption (7) was the necessary condition for the existence of C^1 positive entire solutions. In fact, (7) is the necessary and sufficient condition for the existence of C^1 positive entire solutions, since the sufficient condition was proved by Ngô and Nguyen [2017a]. However, as far as we know, there have been few results about positive solutions of (6) with exponent condition (7) except the case of both critical exponents $p = q = \frac{n+\alpha}{\alpha-n}$. This is a problem well worth studying.

Motivated by the work of Dou, Guo and Zhu [Dou et al. 2017; Dou and Zhu 2015a], we investigate the above problem and prove the following results.

Theorem 1. *If (u, v) is a pair of positive measurable solutions of (6), then u and v are radially symmetric and monotonic increasing about some point x_0 .*

In particular, if we set $u = v$, $\alpha - n = \nu$, and $p = q$, the integral system (4) will be reduced to a single equation

$$(8) \quad u(x) = \int_{\mathbb{R}^n} |x-y|^\nu u^{-p}(y) dy.$$

The negative exponent of integral equation (8) was studied by Li [2004]. Li classified the positive solutions of (8) for $p = \frac{2n+\nu}{\nu}$. It turns out that for $n = 3$ and $\nu = 1$, integral equation (8) is associated with some fourth order conformal covariant operator on three-dimensional compact Riemannian manifolds, arising from the study of conformal geometry. See, e.g., [Xu and Yang 2002] and [Chang and Yang 2002]. Xu [2005] proved that if $n = 3$ and $\nu = 1$, then $p = 7$ and u must take the form $u(x) = c(1 + |x|^2)^{\frac{1}{2}}$ up to dilation and translation. Furthermore, Xu [2007] proved that (8) has a C^1 positive solution if and only if $p = \frac{2n+\nu}{\nu}$. For more related results about integral systems with negative exponents, we refer the reader to [Choi and Xu 2009; Dou et al. 2016; Guo and Wei 2014; Liu et al. 2018; Ma and Wei 2008].

To prove the radial symmetry and monotonicity of the solutions, we use the improved method of moving planes introduced by Dou, Guo and Zhu [2017], which is slightly different from that introduced by Chen, Li and Ou [2006] because they rely on iteration of Hardy–Littlewood–Sobolev inequality to start the moving plane process. In this paper, by the asymptotic behavior and regularity of the solutions, we are able to establish the symmetry and monotonicity results.

2. The proof of radial symmetry and monotonicity

In this section, we use the method of moving planes to prove [Theorem 1](#). For a given real number λ , define

$$T_\lambda = \{x \in \mathbb{R}^n \mid x_1 = \lambda\}, \quad \Sigma_\lambda = \{x \in \mathbb{R}^n \mid x_1 \leq \lambda\}.$$

Let $x_\lambda = \{2\lambda - x_1, x_2, \dots, x_n\}$, $u_\lambda(x) = u(x_\lambda)$ and $v_\lambda(x) = v(x_\lambda)$.

The following lemma was proved in [\[Li 2004\]](#) (see also [\[Dou and Zhu 2015a\]](#)).

Lemma 2. *Let (u, v) be a pair of positive measurable solutions of (6). Then:*

$$(i) \int_{\mathbb{R}^n} (1 + |y|^{\alpha-n}) u^{-q}(y) dy < \infty, \quad \int_{\mathbb{R}^n} (1 + |y|^{\alpha-n}) v^{-p}(y) dy < \infty.$$

$$(ii) 0 < a := \lim_{|x| \rightarrow \infty} |x|^{n-\alpha} u(x) = \int_{\mathbb{R}^n} v^{-p}(y) dx < \infty,$$

$$0 < b := \lim_{|x| \rightarrow \infty} |x|^{n-\alpha} v(x) = \int_{\mathbb{R}^n} u^{-q}(y) dx < \infty.$$

(iii) *There exist some constants $C_1, C_2 > 0$ such that*

$$\frac{1 + |x|^{\alpha-n}}{C_1} \leq u(x) \leq C_1(1 + |x|^{\alpha-n}),$$

and

$$\frac{1 + |x|^{\alpha-n}}{C_2} \leq v(x) \leq C_2(1 + |x|^{\alpha-n}).$$

(iv) $u \in C^\infty(\mathbb{R}^n)$ and $v \in C^\infty(\mathbb{R}^n)$.

From the above lemma, we know that $p > \frac{\alpha}{\alpha-n}$ and $q > \frac{\alpha}{\alpha-n}$.

Lemma 3. *Let (u, v) be a pair of positive measurable solutions of (6). Then*

$$(9) \quad u(x) - u_\lambda(x) = \int_{\Sigma_\lambda} K(\lambda, x, y) (v_\lambda^{-p}(y) - v^{-p}(y)) dy$$

and

$$(10) \quad v(x) - v_\lambda(x) = \int_{\Sigma_\lambda} K(\lambda, x, y) (u_\lambda^{-q}(y) - u^{-q}(y)) dy,$$

where $K(\lambda, x, y) = |x_\lambda - y|^{\alpha-n} - |x - y|^{\alpha-n}$.

Proof. We only prove (9); by (6), we have,

$$\begin{aligned} u(x) &= \int_{\Sigma_\lambda} |x - y|^{\alpha-n} v^{-p}(y) dy + \int_{\mathbb{R}^n \setminus \Sigma_\lambda} |x - y|^{\alpha-n} v^{-p}(y) dy \\ &= \int_{\Sigma_\lambda} |x - y|^{\alpha-n} v^{-p}(y) dy + \int_{\Sigma_\lambda} |x - y_\lambda|^{\alpha-n} v_\lambda^{-p}(y) dy \end{aligned}$$

and

$$\begin{aligned} u_\lambda(x) &= \int_{\Sigma_\lambda} |x - y|^{\alpha-n} v^{-p}(y) dy + \int_{\mathbb{R}^n \setminus \Sigma_\lambda} |x - y|^{\alpha-n} v^{-p}(y) dy \\ &= \int_{\Sigma_\lambda} |x_\lambda - y|^{\alpha-n} v^{-p}(y) dy + \int_{\Sigma_\lambda} |x_\lambda - y_\lambda|^{\alpha-n} v_\lambda^{-p}(y) dy. \end{aligned}$$

Combining with $|x_\lambda - y| = |x - y_\lambda|$ and $|x_\lambda - y_\lambda| = |x - y|$, we obtain (9). $\square$

To prove [Theorem 1](#), we compare $u(x)$ with $u_\lambda(x)$ and $v(x)$ with $v_\lambda(x)$ on Σ_λ . The proof consists of two steps. In Step 1, we show that there exists an $N < 0$ such that for $\lambda \leq N$,

$$(11) \quad u(x) \geq u_\lambda(x), \quad v(x) \geq v_\lambda(x), \quad \text{for all } x \in \Sigma_\lambda.$$

Thus we can start moving the plane continuously from $\lambda \leq N$ to the right as long as (11) holds. In Step 2, we show that if the plane stops at $x_1 = \lambda_0$ for some $\lambda_0 < 0$, then $u(x)$ and $v(x)$ must be symmetric and monotonic increasing about the plane $x_1 = \lambda_0$. Otherwise, we can move the plane all the way to x_1 . Since the direction of x_1 can be chosen arbitrarily, we deduce that $u(x)$ and $v(x)$ must be radially symmetric and monotonic increasing about some point.

Step 1. We first prove that there exists sufficiently negative $N < 0$ such that

$$(12) \quad \nabla(|x|^{\frac{n-\alpha}{2}} u(x)) \cdot x = |x|^{\frac{n-\alpha}{2}} \left(\nabla u(x) \cdot x + \frac{n-\alpha}{2} u(x) \right) > 0, \quad \text{for all } x_1 < N.$$

To obtain (12), it is sufficient to prove that

$$(13) \quad \lim_{|x| \rightarrow \infty} |x|^{n-\alpha} \left(\nabla u(x) \cdot x + \frac{n-\alpha}{2} u(x) \right) > 0.$$

We only prove (13) for $i = 1$. Fix $x_2, x_3, \dots, x_n$. By [Lemma 2](#), $u(x)$ is differentiable in x_1 ; then, we have

$$\begin{aligned} \left| \frac{\partial u(x)}{\partial x_1} \right| &= \left| (\alpha - n) \int_{\mathbb{R}^n} |x - y|^{\alpha-n-2} (x_1 - y_1) v^{-p}(y) dy \right| \\ &\leq (\alpha - n) \int_{\mathbb{R}^n} |x - y|^{\alpha-n-1} v^{-p}(y) dy. \end{aligned}$$

Now we carry out the proof in two cases.

Case (i): If $\alpha - n \geq 1$, then by [Lemma 2](#), letting $|x_1| \geq 1$, we have

$$\begin{aligned} \frac{\left| \frac{\partial u(x)}{\partial x_1} \right|}{|x_1|^{\alpha-n-1}} &\leq \left| (\alpha - n) \int_{\mathbb{R}^n} (1 + |y|^{\alpha-n-1}) v^{-p}(y) dy \right| \\ &\leq C \int_{\mathbb{R}^n} (1 + |y|^{\alpha-n}) v^{-p}(y) dy. \end{aligned}$$

Thus, by using Lebesgue's dominated convergence theorem, we have

$$\lim_{|x_1| \rightarrow \infty} \frac{\left| \frac{\partial u(x)}{\partial x_1} \right|}{|x_1|^{\alpha-n-1}} = a(\alpha-n).$$

Case (ii): If $0 < \alpha - n < 1$, we can calculate

$$\begin{aligned} \frac{\left| \frac{\partial u(x)}{\partial x_1} \right|}{|x|^{\alpha-n-1}} &\leq (\alpha-n) \int_{\mathbb{R}^n} \frac{|x|^{n-\alpha+1}}{|x-y|^{n-\alpha+1}} v^{-p}(y) dy \\ &= (\alpha-n) \int_{\mathbb{R}^n \setminus B(x, \frac{|x|}{2})} \frac{|x|^{n-\alpha+1}}{|x-y|^{n-\alpha+1}} v^{-p}(y) dy \\ &\quad + (\alpha-n) \int_{B(x, \frac{|x|}{2})} \frac{|x|^{n-\alpha+1}}{|x-y|^{n-\alpha+1}} v^{-p}(y) dy \\ &:= (\alpha-n) J_1(x) + (\alpha-n) J_2(x). \end{aligned}$$

For $J_1(x)$, since $|x-y| > \frac{|x|}{2}$, we have

$$(14) \quad J_1(x) \leq C \int_{\mathbb{R}^n \setminus B(x, \frac{|x|}{2})} \frac{|x|^{n-\alpha+1}}{|x-y|^{n-\alpha+1}} v^{-p}(y) dy \leq C \int_{\mathbb{R}^n} v^{-p}(y) dy.$$

For $J_2(x)$, by [Lemma 2](#),

$$\begin{aligned} J_2(x) &= C \int_{B(x, \frac{|x|}{2})} \frac{|x|^{n-\alpha+1}}{|z|^{n-\alpha+1}} v^{-p}(x+z) dz \\ &\leq C \int_{B(0, \frac{|x|}{2})} \frac{|x|^{n-\alpha+1}}{|z|^{n-\alpha+1}} (1+|x+z|^{\alpha-n})^{-p} dz. \end{aligned}$$

Notice that for $|x|$ large enough and $z \in B(0, \frac{|x|}{2})$, it is easy to check that $|x+z| \geq \frac{|x|}{2} \geq \frac{|z|}{2}$. Therefore,

$$\begin{aligned} J_2(x) &\leq C \int_{B(0, \frac{|x|}{2})} \frac{|x|^{n-\alpha+1}}{|z|^{n-\alpha+1}} \cdot \frac{(1+|x+z|^{\alpha-n})^{-p+\frac{n-\alpha+1}{\alpha-n}}}{(1+|x+z|^{\alpha-n})^{\frac{n-\alpha+1}{\alpha-n}}} dz \\ &\leq C \int_{B(0, \frac{|x|}{2})} \frac{|x|^{n-\alpha+1}}{|z|^{n-\alpha+1}} \cdot \frac{(1+|z|^{\alpha-n})^{-p+\frac{n-\alpha+1}{\alpha-n}}}{(1+|x|^{\alpha-n})^{\frac{n-\alpha+1}{\alpha-n}}} dz \\ &\leq C \int_{\mathbb{R}^n} \frac{(1+|z|^{\alpha-n})^{-p+\frac{n-\alpha+1}{\alpha-n}}}{|z|^{n-\alpha+1}} dz \\ &= C \int_{\mathbb{R}^n \setminus B(0,1)} \frac{(1+|z|^{\alpha-n})^{-p+\frac{n-\alpha+1}{\alpha-n}}}{|z|^{n-\alpha+1}} dz + C \int_{B(0,1)} \frac{(1+|z|^{\alpha-n})^{-p+\frac{n-\alpha+1}{\alpha-n}}}{|z|^{n-\alpha+1}} dz \\ &:= C J_{21}(x) + C J_{22}(x). \end{aligned}$$

Since $0 < \alpha - n < 1$,

$$(15) \quad J_{22}(x) \leq C \int_{B(0,1)} \frac{1}{|z|^{n-\alpha+1}} dz < \infty,$$

and

$$(16) \quad J_{21}(x) \leq C \int_{\mathbb{R}^n \setminus B(0,1)} \frac{|z|^{-p(\alpha-n)+n-\alpha+1}}{|z|^{n-\alpha+1}} dz \leq C \int_{\mathbb{R}^n \setminus B(0,1)} \frac{1}{|z|^{p(\alpha-n)}} dz < \infty,$$

where we use $p > \frac{\alpha}{\alpha-n}$.

Combining with (14), (15), (16) and using Lebesgue's dominated convergence theorem, we have

$$\lim_{|x_1| \rightarrow \infty} \frac{\left| \frac{\partial u(x)}{\partial x_1} \right|}{|x_1|^{\alpha-n-1}} = a(\alpha - n).$$

Recall that $a = \lim_{|x| \rightarrow \infty} |x|^{n-\alpha} u(x) = \int_{\mathbb{R}^n} v^{-p}(y) dx < \infty$; thus, we obtain (13).

If $|x_1| \geq -N$ and $|2\lambda - x_1| > -N$, by (12), we have

$$|x|^{\frac{n-\alpha}{2}} u(x) > |x_\lambda|^{\frac{n-\alpha}{2}} u_\lambda(x), \quad x \in \Sigma_\lambda.$$

Then we obtain

$$u(x) \geq u_\lambda(x).$$

If $|2\lambda - x_1| \leq -N$, by Lemma 2, we can take λ large enough such that

$$u(x) \geq \frac{1 + |x_1|^{\alpha-n}}{C_1} \geq \frac{1 + |\lambda|^{\alpha-n}}{C_1} \geq C_1(1 + |N|^{\alpha-n}) \geq C_1(1 + |2\lambda - x_1|^{\alpha-n}) \geq u_\lambda(x).$$

Therefore (11) holds. This completes Step 1.

Step 2. For $x \in \mathbb{R}^n$, define

$$\lambda_0 = \sup\{\mu < 0 \mid u(x) \geq u_\lambda(x), v(x) \geq v_\lambda(x) \text{ for all } \lambda < \mu, x_1 \leq \lambda\}.$$

It is easy to check that $|x_{\lambda_0} - y|^{\alpha-n} - |x - y|^{\alpha-n} > 0$ for any $y_1, x_1 < \lambda_0$. So by (9) and (10), for all $x \in \Sigma_{\lambda_0}$, there are the following four cases:

Case (i): $u(x) = u_{\lambda_0}(x), v(x) = v_{\lambda_0}(x)$.

Case (ii): $u(x) > u_{\lambda_0}(x), v(x) = v_{\lambda_0}(x)$.

Case (iii): $u(x) = u_{\lambda_0}(x), v(x) > v_{\lambda_0}(x)$.

Case (iv): $u(x) > u_{\lambda_0}(x), v(x) > v_{\lambda_0}(x)$.

For Case (i), we are done. Case (ii) and Case (iii) are impossible according to (9) and (10). Thus it suffices to show that Case (iv) is also impossible. We carry out

the argument by contradiction. Suppose that Case (iv) is valid. We will show that there exists an $\varepsilon > 0$ such that

$$(17) \quad u(x) \geq u_\lambda(x), \quad v(x) \geq v_\lambda(x), \quad \text{for all } \lambda_0 < \lambda < \lambda_0 + \varepsilon, \quad x \in \Sigma_\lambda.$$

By (9) and (10) and Fatou's lemma, for all $x_1 \leq \lambda_0 - 1$, we can calculate

$$\begin{aligned} & \lim_{|x| \rightarrow \infty} \inf |x|^{n-\alpha} (u(x) - u_{\lambda_0}(x)) \\ &= \int_{\Sigma_{\lambda_0}} \lim_{|x| \rightarrow \infty} \inf |x|^{n-\alpha} (|x_{\lambda_0} - y|^{\alpha-n} - |x - y|^{\alpha-n}) (v_{\lambda_0}^{-p}(y) - v^{-p}(y)) dy > 0 \end{aligned}$$

and

$$\begin{aligned} & \lim_{|x| \rightarrow \infty} \inf |x|^{n-\alpha} (v(x) - v_{\lambda_0}(x)) \\ &= \int_{\Sigma_{\lambda_0}} \lim_{|x| \rightarrow \infty} \inf |x|^{n-\alpha} (|x_{\lambda_0} - y|^{\alpha-n} - |x - y|^{\alpha-n}) (u_{\lambda_0}^{-q}(y) - u^{-q}(y)) dy > 0. \end{aligned}$$

Then, for any $x_1 \leq \lambda_0 - 1$, there exists $\varepsilon_1 \in (0, 1)$ such that

$$u(x) - u_{\lambda_0}(x) \geq \varepsilon_1 |x|^{\alpha-n}, \quad v(x) - v_{\lambda_0}(x) \geq \varepsilon_1 |x|^{\alpha-n}.$$

We can use continuity of u and v with respect to the variable λ to obtain that

$$\begin{aligned} (18) \quad u(x) - u_\lambda(x) &\geq \varepsilon_1 |x|^{\alpha-n} + (u_\lambda(x) - u_{\lambda_0}(x)) \\ &\geq \frac{\varepsilon_1}{2} |x|^{\alpha-n}, \quad \text{for all } x_1 \leq \lambda_0 - 1, \quad \lambda_0 \leq \lambda \leq \lambda_0 + \varepsilon_2 \end{aligned}$$

and

$$\begin{aligned} (19) \quad v(x) - v_\lambda(x) &\geq \varepsilon_1 |x|^{\alpha-n} + (v_\lambda(x) - v_{\lambda_0}(x)) \\ &\geq \frac{\varepsilon_1}{2} |x|^{\alpha-n}, \quad \text{for all } x_1 \leq \lambda_0 - 1, \quad \lambda_0 \leq \lambda \leq \lambda_0 + \varepsilon_2 \end{aligned}$$

for sufficiently small $\varepsilon_2 > 0$. Thus, for $\varepsilon \in (0, \varepsilon_2)$, which we choose below, it suffices to verify that for $\lambda_0 - 1 \leq x_1 \leq \lambda$,

$$u(x) \geq u_\lambda(x), \quad v(x) \geq v_\lambda(x), \quad \text{for all } \lambda_0 \leq \lambda \leq \lambda_0 + \varepsilon.$$

We only show that

$$u(x) \geq u_\lambda(x), \quad \text{for all } \lambda_0 \leq \lambda \leq \lambda_0 + \varepsilon, \quad \lambda_0 - 1 \leq x_1 \leq \lambda.$$

By (13), there exists $R_1 \geq 4(|\lambda_0| + 1)$ large enough such that, for any $x \in (\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}) \setminus B(0, R_1/2)$,

$$\nabla(|x|^{\frac{n-\alpha}{2}} \cdot x) = |x|^{\frac{n-\alpha}{2}} \left(\nabla u(x) \cdot x + \frac{n-\alpha}{2} u(x) \right) > 0.$$

We know that if $|x| \geq R_1$, then $|x_\lambda| \geq \frac{R_1}{2}$; thus

$$|x|^{\frac{n-\alpha}{2}} u(x) > |x_\lambda|^{\frac{n-\alpha}{2}} u_\lambda(x).$$

Therefore,

$$u(x) \geq u_\lambda(x), \quad \text{for } |x| \geq R_1.$$

Now we fix R_1 and consider

$$x \in (\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}) \cap B(0, R_1).$$

By [Lemma 3](#), we can write

$$\begin{aligned} u(x) - u_\lambda(x) &= \int_{\Sigma_\lambda} K(\lambda, x, y)(v_\lambda^{-p}(y) - v^{-p}(y)) dy \\ &\geq \int_{\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}} K(\lambda, x, y)(v_\lambda^{-p}(y) - v^{-p}(y)) dy \\ &\quad + \int_{\Sigma_{\lambda_0-2} \setminus \Sigma_{\lambda_0-3}} K(\lambda, x, y)(v_\lambda^{-p}(y) - v^{-p}(y)) dy \\ &\geq \int_{(\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}) \setminus B(0, R_0)} K(\lambda, x, y)(v_\lambda^{-p}(y) - v_{\lambda_0}^{-p}(y)) dy \\ &\quad + \int_{(\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}) \cap B(0, R_0)} K(\lambda, x, y)(v_\lambda^{-p}(y) - v_{\lambda_0}^{-p}(y)) dy \\ &\quad + \int_{\Sigma_{\lambda_0-2} \setminus \Sigma_{\lambda_0-3}} K(\lambda, x, y)(v_\lambda^{-p}(y) - v^{-p}(y)) dy \\ &:= I_1(x) + I_2(x) + I_3(x). \end{aligned}$$

We first estimate $I_1(x)$. Take $R_0 \geq 2R_1$ large enough such that

$$\frac{1}{R_0^{p(\alpha-n)-\alpha+1}} \leq \varepsilon,$$

where we use $p > \frac{\alpha}{\alpha-n}$. Since

$$x \in (\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}) \cap B(0, R_1) \quad \text{and} \quad y \in (\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}) \setminus B(0, R_0),$$

it easy to derive that $|x - y| \sim |x_\lambda - y| \sim |y|$; then

$$\begin{aligned} K(\lambda, x, y) &= |x - y_\lambda|^{\alpha-n} - |x - y|^{\alpha-n} \\ &\leq C \max\{|x - y_\lambda|^{\alpha-n-1}, |x - y|^{\alpha-n-1}\} ||x - y_\lambda| - |x - y|| \\ &\leq C |y|^{\alpha-n-1} (\lambda - x_1). \end{aligned}$$

Thus, by using [Lemma 2](#),

$$\begin{aligned}
 (20) \quad & \left| \int_{(\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}) \setminus B(0, R_0)} K(\lambda, x, y)(v_\lambda^{-p}(y) - v_{\lambda_0}^{-p}(y)) dy \right| \\
 & \leq C(\lambda - x_1) \int_{(\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}) \setminus B(0, R_0)} |y|^{\alpha-n-1} |y|^{-p(\alpha-n)} dy \\
 & \leq C(\lambda - x_1) \int_{\mathbb{R}^n \setminus B(0, R_0)} |y|^{\alpha-n-1-p(\alpha-n)} dy \\
 & = C(\lambda - x_1) \int_{R_0}^{\infty} r^{\alpha-2-p(\alpha-n)} dr \\
 & = C(\lambda - x_1) R_0^{\alpha-1-p(\alpha-n)} \\
 & \leq C\varepsilon(\lambda - x_1).
 \end{aligned}$$

Now, we estimate $I_2(x)$. We can take $0 < \varepsilon_3 < \varepsilon$ small enough such that for any $\lambda_0 \leq \lambda \leq \lambda_0 + \varepsilon_3$ and $y \in (\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}) \cap B(0, R_0)$,

$$|v_\lambda^{-p}(y) - v_{\lambda_0}^{-p}(y)| \leq C_1 \varepsilon.$$

Then,

$$\begin{aligned}
 (21) \quad |I_2(x)| & \leq \int_{(\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}) \cap B(0, R_0)} |K(\lambda, x, y)(v_\lambda^{-p}(y) - v_{\lambda_0}^{-p}(y))| dy \\
 & \leq C(R_0, R_1) |x_\lambda - x| \int_{(\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}) \cap B(0, R_0)} |v_\lambda^{-p}(y) - v_{\lambda_0}^{-p}(y)| dy \\
 & \leq C\varepsilon(\lambda - x_1).
 \end{aligned}$$

For $I_3(x)$, we can assume $\lambda_0 \leq \lambda \leq \lambda_0 + \varepsilon_3$. By [\(18\)](#) and [\(19\)](#), there exists $\delta_1 > 0$ which is independent of ε_3 such that

$$(22) \quad v_\lambda^{-p}(y) - v^{-p}(y) \geq \delta_1, \quad \text{for any } y \in \Sigma_{\lambda_0-2} \setminus \Sigma_{\lambda_0-3}.$$

Notice that

$$\frac{\partial K(\lambda, x, y)}{\partial x_1} \cdot x_1|_{x_1=\lambda} = 2(\alpha - n)|x - y|^{\alpha-n-2}(y_1 - \lambda)x_1 > 0, \quad y_1 < \lambda.$$

Since $K(\lambda, x, y)|_{x_1=\lambda} = 0$, for any

$$x \in (\Sigma_\lambda \setminus \Sigma_{\lambda_0-1}) \cap B(0, R_1) \quad \text{and} \quad y \in (\Sigma_{\lambda_0-2} \setminus \Sigma_{\lambda_0-3}),$$

there exists $\delta_2 > 0$ which are independent of $\varepsilon_3, \varepsilon$, such that

$$(23) \quad |K(\lambda, x, y)| \geq \delta_2(\lambda - x_1).$$

By (20), (21), (22), and (23), for sufficiently small $\varepsilon > 0$, we conclude that

$$u(x) - u_\lambda(x) \geq (\delta_1 \delta_2 - C_1 C_2 \varepsilon)(\lambda - x_1) \geq 0.$$

Combining with (18) and (19), we can derive a contradiction with the definition of λ_0 . This completes the proof of Step 2. Thus, we have finished the proof of the main theorem.

Acknowledgements

The author would like to express his deep appreciation to Prof. Jingbo Dou for many fruitful discussions. He also thanks the anonymous referee for his/her valuable comments and suggestions.

References

- [Beckner 1993] W. Beckner, “Sharp Sobolev inequalities on the sphere and the Moser–Trudinger inequality”, *Ann. of Math.* (2) **138**:1 (1993), 213–242. [MR](#) [Zbl](#)
- [Beckner 2008] W. Beckner, “Weighted inequalities and Stein–Weiss potentials”, *Forum Math.* **20**:4 (2008), 587–606. [MR](#) [Zbl](#)
- [Brascamp and Lieb 1976] H. J. Brascamp and E. H. Lieb, “Best constants in Young’s inequality, its converse, and its generalization to more than three functions”, *Adv. Math.* **20**:2 (1976), 151–173. [MR](#) [Zbl](#)
- [Chang and Yang 2002] S.-Y. A. Chang and P. C. Yang, “Non-linear partial differential equations in conformal geometry”, pp. 189–207 in *Proceedings of the International Congress of Mathematicians, I* (Beijing, 2002), edited by T. Li, Higher Ed. Press, Beijing, 2002. [MR](#) [arXiv](#)
- [Chen and Li 2008] W. Chen and C. Li, “The best constant in a weighted Hardy–Littlewood–Sobolev inequality”, *Proc. Amer. Math. Soc.* **136**:3 (2008), 955–962. [MR](#) [Zbl](#)
- [Chen et al. 2005a] W. Chen, C. Jin, C. Li, and J. Lim, “Weighted Hardy–Littlewood–Sobolev inequalities and systems of integral equations”, *Discrete Contin. Dyn. Syst.* **2005**:suppl. (2005), 164–172. [MR](#) [Zbl](#)
- [Chen et al. 2005b] W. Chen, C. Li, and B. Ou, “Classification of solutions for a system of integral equations”, *Comm. Partial Differential Equations* **30**:1-3 (2005), 59–65. [MR](#) [Zbl](#)
- [Chen et al. 2006] W. Chen, C. Li, and B. Ou, “Classification of solutions for an integral equation”, *Comm. Pure Appl. Math.* **59**:3 (2006), 330–343. Correction in **59**:7 (2006), 1064. [MR](#) [Zbl](#)
- [Chen et al. 2016] L. Chen, Z. Liu, and G. Lu, “Symmetry and regularity of solutions to the weighted Hardy–Sobolev type system”, *Adv. Nonlinear Stud.* **16**:1 (2016), 1–13. [MR](#) [Zbl](#)
- [Choi and Xu 2009] Y. S. Choi and X. Xu, “Nonlinear biharmonic equations with negative exponents”, *J. Differential Equations* **246**:1 (2009), 216–234. [MR](#) [Zbl](#)
- [Dou 2016] J. Dou, “Weighted Hardy–Littlewood–Sobolev inequalities on the upper half space”, *Commun. Contemp. Math.* **18**:5 (2016), art. id. 1550067. [MR](#) [Zbl](#)
- [Dou and Zhu 2015a] J. Dou and M. Zhu, “Reversed Hardy–Littlewood [sic]–Sobolev inequality”, *Int. Math. Res. Not.* **2015**:19 (2015), 9696–9726. [MR](#) [Zbl](#)
- [Dou and Zhu 2015b] J. Dou and M. Zhu, “Sharp Hardy–Littlewood–Sobolev inequality on the upper half space”, *Int. Math. Res. Not.* **2015**:3 (2015), 651–687. [MR](#) [Zbl](#)

- [Dou et al. 2016] J. Dou, F. Ren, and J. Villavert, “Classification of positive solutions to a Lane–Emden type integral system with negative exponents”, *Discrete Contin. Dyn. Syst.* **36**:12 (2016), 6767–6780. [MR](#) [Zbl](#)
- [Dou et al. 2017] J. Dou, Q. Guo, and M. Zhu, “Subcritical approach to sharp Hardy–Littlewood–Sobolev type inequalities on the upper half space”, *Adv. Math.* **312** (2017), 1–45. [MR](#) [Zbl](#)
- [Guo and Wei 2014] Z. Guo and J. Wei, “Liouville type results and regularity of the extremal solutions of biharmonic equation with negative exponents”, *Discrete Contin. Dyn. Syst.* **34**:6 (2014), 2561–2580. [MR](#) [Zbl](#)
- [Han and Zhu 2016] Y. Han and M. Zhu, “Hardy–Littlewood–Sobolev inequalities on compact Riemannian manifolds and applications”, *J. Differential Equations* **260**:1 (2016), 1–25. [MR](#) [Zbl](#)
- [Lei 2015] Y. Lei, “On the integral systems with negative exponents”, *Discrete Contin. Dyn. Syst.* **35**:3 (2015), 1039–1057. [MR](#) [Zbl](#)
- [Li 2004] Y. Y. Li, “Remark on some conformally invariant integral equations: the method of moving spheres”, *J. Eur. Math. Soc.* **6**:2 (2004), 153–180. [MR](#) [Zbl](#)
- [Liu et al. 2018] Z. Liu, L. Chen, and X. Wang, “Liouville type theorems for general integral system with negative exponents”, *Taiwanese J. Math.* **22**:3 (2018), 661–675. [MR](#)
- [Lu and Zhu 2011] G. Lu and J. Zhu, “Symmetry and regularity of extremals of an integral equation related to the Hardy–Sobolev inequality”, *Calc. Var. Partial Differential Equations* **42**:3-4 (2011), 563–577. [MR](#) [Zbl](#)
- [Ma and Wei 2008] L. Ma and J. C. Wei, “Properties of positive solutions to an elliptic equation with negative exponent”, *J. Funct. Anal.* **254**:4 (2008), 1058–1087. [MR](#) [Zbl](#)
- [Ngô and Nguyen 2017a] Q. A. Ngô and V. H. Nguyen, “Sharp reversed Hardy–Littlewood–Sobolev inequality on $\mathbb{R}^n$ ”, *Israel J. Math.* **220**:1 (2017), 189–223. [MR](#) [Zbl](#)
- [Ngô and Nguyen 2017b] Q. A. Ngô and V. H. Nguyen, “Sharp reversed Hardy–Littlewood–Sobolev inequality on the half space $\mathbb{R}_+^n$ ”, *Int. Math. Res. Not.* **2017**:20 (2017), 6187–6230. [MR](#) [Zbl](#)
- [Xu 2005] X. Xu, “Exact solutions of nonlinear conformally invariant integral equations in $\mathbb{R}^3$ ”, *Adv. Math.* **194**:2 (2005), 485–503. [MR](#) [Zbl](#)
- [Xu 2007] X. Xu, “Uniqueness theorem for integral equations and its application”, *J. Funct. Anal.* **247**:1 (2007), 95–109. [MR](#) [Zbl](#)
- [Xu and Yang 2002] X. Xu and P. C. Yang, “On a fourth order equation in 3-D”, *ESAIM Control Optim. Calc. Var.* **8** (2002), 1029–1042. [MR](#) [Zbl](#)

Received February 26, 2018. Revised September 11, 2018.

ZHAO LIU

SCHOOL OF MATHEMATICS AND COMPUTER SCIENCE

JIANGXI SCIENCE AND TECHNOLOGY NORMAL UNIVERSITY

NANCHANG

CHINA

liuzhao2008tj@sina.com

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2019 is US \$490/year for the electronic version, and \$665/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY



mathematical sciences publishers

nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 300 No. 2 June 2019

Hochschild cochain spectral sequence and the Beilinson residue	257
OLIVER BRAUNLING and JESSE WOLFSON	
The graph Laplacian and Morse inequalities	331
IVAN CONTRERAS and BOYAN XU	
Counting using Hall algebras III: Quivers with potentials	347
JIARUI FEI	
J -invariant of hermitian forms over quadratic extensions	375
RAPHAËL FINO	
On the determinants and permanents of matrices with restricted entries over prime fields	405
DOOWON KOH, THANG PHAM, CHUN-YEN SHEN and LE ANH VINH	
Symmetry and monotonicity of positive solutions for an integral system with negative exponents	419
ZHAO LIU	
On the Braverman–Kazhdan proposal for local factors: spherical case	431
ZHILIN LUO	
Fields of character values for finite special unitary groups	473
A. A. SCHAEFFER FRY and C. RYAN VINROOT	
Sharp quantization for Lane–Emden problems in dimension two	491
PIERRE-DAMIEN THIZY	
Towards a sharp converse of Wall’s theorem on arithmetic progressions	499
JOSEPH VANDEHEY	