

Pacific Journal of Mathematics

**UNIFORM BOUNDS OF THE PILTZ DIVISOR PROBLEM
OVER NUMBER FIELDS**

WATARU TAKEDA

Volume 301 No. 2

August 2019

UNIFORM BOUNDS OF THE PILTZ DIVISOR PROBLEM OVER NUMBER FIELDS

WATARU TAKEDA

We consider the upper bound of the Piltz divisor problem over number fields. The Piltz divisor problem is known as a generalization of the Dirichlet divisor problem. We deal with this problem over number fields and improve the error term of this function for many cases. Our proof uses the estimate of exponential sums. We also show uniform results for the ideal counting function and relatively r -prime lattice points as one of its applications.

1. Introduction

The behavior of arithmetic functions has long been studied, and it is one of the most important areas of research in analytic number theory. But many arithmetic functions $f(n)$ fluctuate as n increases, and it becomes difficult to deal with them. Thus, many authors study partial sums $\sum_{n \leq x} f(n)$ to obtain some information about arithmetic functions $f(n)$. In this paper we consider the Piltz divisor function $I_K^m(x)$ over a number field. Let K be a number field with extension degree $[K : \mathbb{Q}] = n$, and let \mathbb{O}_K be its ring of integers. Let D_K be the absolute value of the discriminant of K . Then the Piltz divisor function $I_K^m(x)$ counts the number of m -tuples of ideals $(\mathfrak{a}_1, \mathfrak{a}_2, \dots, \mathfrak{a}_m)$ such that the product of their ideal norms $\mathfrak{N}\mathfrak{a}_1 \cdots \mathfrak{N}\mathfrak{a}_m \leq x$. It is known that

$$(1-1) \quad I_K^m(x) \sim \operatorname{Res}_{s=1} \left(\zeta_K(s)^m \frac{x^s}{s} \right).$$

We denote by $\Delta_K^m(x)$ the error term of $I_K^m(x)$, that is, $I_K^m(x) - \operatorname{Res}_{s=1} \left(\zeta_K(s)^m \frac{x^s}{s} \right)$.

In the case $m = 1$, this function is the ordinary ideal counting function over K . For simplicity we substitute $I_K(x)$ and $\Delta_K(x)$ for $I_K^1(x)$ and $\Delta_K^1(x)$, respectively. There are many results about $I_K(x)$ from the 1900s. In the case $K = \mathbb{Q}$, integer ideals of \mathbb{Z} and positive integers are in one-to-one correspondence, so $I_{\mathbb{Q}}(x) = [x]$, where $[\cdot]$ is the Gauss symbol. For the general case, the best estimate of $\Delta_K(x)$ hitherto is the following theorem.

MSC2010: primary 11N45; secondary 11R42, 11H06, 11P21.

Keywords: ideal counting function, exponential sum, Piltz divisor problem.

Theorem 1-2. *The following estimates hold for all $\varepsilon > 0$:*

$n = [K : \mathbb{Q}]$	$\Delta_K(x)$	
2	$O(x^{\frac{131}{416}}(\log x)^{\frac{18627}{8320}})$	[Huxley 2002]
3	$O(x^{\frac{43}{96}+\varepsilon})$	[Müller 1988]
4	$O(x^{\frac{41}{72}+\varepsilon})$	[Bordellès 2015]
$5 \leq n \leq 10$	$O(x^{1-\frac{4}{2n+1}+\varepsilon})$	[Bordellès 2015]
$11 \leq n$	$O(x^{1-\frac{3}{n+6}+\varepsilon})$	[Lao 2010]

There are also many results about $I_{\mathbb{Q}}^m$ from the 1800s. In 1849 Dirichlet showed that

$$I_{\mathbb{Q}}^2(x) = x \log x + (2\gamma - 1)x + O(x^{\frac{1}{2}}),$$

where γ is the Euler constant, defined by

$$\gamma = \lim_{n \rightarrow \infty} \left(\sum_{k=1}^n \frac{1}{k} - \log n \right).$$

The O -term has been improved by many researchers many times; the best estimate hitherto is $x^{\frac{517}{1648}+\varepsilon}$ [Bourgain and Watt 2017].

As we have mentioned above, there exist many results about other divisor problems, but it seems that there are not many results about the Piltz divisor problem over number fields.

Theorem 1-3 [Nowak 1993]. *When $n = [K : \mathbb{Q}] \geq 2$, then we get*

$$\Delta_K^m(x) = \begin{cases} O_K(x^{1-\frac{2}{mn}+\frac{8}{mn(5mn+2)}}(\log x)^{m-1-\frac{10(m-2)}{5n+2}}) & \text{for } 3 \leq mn \leq 6, \\ O_K(x^{1-\frac{2}{mn}+\frac{3}{2m^2n^2}}(\log x)^{m-1-\frac{2(m-2)}{mn}}) & \text{for } mn \geq 7. \end{cases}$$

For the estimate of the lower bound, Girstmair, Kühleitner, Müller, and Nowak obtain the following Ω -results:

Theorem 1-4 [Girstmair et al. 2005]. *For any fixed number field K with $n = [K : \mathbb{Q}] \geq 2$,*

$$(1-5) \quad \Delta_K^m(x) = \Omega(x^{\frac{1}{2}-\frac{1}{2mn}}(\log x)^{\frac{1}{2}-\frac{1}{2mn}}(\log \log x)^{\kappa}(\log \log \log x)^{-\lambda}),$$

where κ and λ are constants depending on K . To be more precise, let K^{gal} be the Galois closure of K/\mathbb{Q} , $G = \text{Gal}(K^{\text{gal}}/\mathbb{Q})$ its Galois group, and $H = \text{Gal}(K^{\text{gal}}/K)$ the subgroup of G corresponding to K . Then

$$\kappa = \frac{mn+1}{2mn} \left(\sum_{v=1}^n \delta_v v^{\frac{2mn}{mn+1}} - 1 \right) \quad \text{and} \quad \lambda = \frac{mn+1}{4mn} R + \frac{mn-1}{2mn},$$

where

$$\delta_v = \frac{|\{\tau \in G \mid |\{\sigma \in G \mid \tau \in \sigma H \sigma^{-1}\}| = v|H|\}|}{|G|}$$

and R is the number of $1 \leq v \leq n$ with $\delta_v > 0$.

We know the following conditional result. If we assume the Lindelöf hypothesis for the Dedekind zeta function, it holds that for all $\varepsilon > 0$, for all K , and for all m ,

$$(1-6) \quad \Delta_K^m(x) = O_\varepsilon(x^{\frac{1}{2}+\varepsilon} D_K^\varepsilon).$$

In this paper we estimate the error term of $\Delta_K^m(x)$ by using exponential sums. In [Nowak 1993; Girstmair et al. 2005], they use other approaches, so we expect new development for the Piltz divisor problem over number fields. As a results, we improve the estimate of upper bound of $\Delta_K^m(x)$ for many K and many m .

In Section 2, we show some auxiliary theorems to consider the upper bound of the error term $\Delta_K^m(x)$. First we give a review of the convexity bound for the Dedekind zeta function and generalized Atkinson lemma [1941]. Next we show Proposition 2-6, which reduces an ideal counting problem to a problem of exponential sums. This proposition plays a crucial role in our computing $\Delta_K^m(x)$.

In Section 3, we prove the following theorem about the error term $\Delta_K^m(x)$ by using estimates of exponential sums.

Theorem 1-7. *For every $\varepsilon > 0$ the following estimate holds. When $mn \geq 4$, then*

$$\Delta_K^m(x) = O_{n,m,\varepsilon}(x^{\frac{2mn-3}{2mn+1}+\varepsilon} D_K^{\frac{2m}{2mn+1}+\varepsilon}).$$

This theorem gives improvement of the upper bound of $\Delta_K^m(x)$ for $mn \geq 4$.

In Section 4, we give some applications. First we give a uniform estimate for the ideal counting function over number fields. Second we show a good uniform upper bound of the distribution of relatively r -prime lattice points over number fields as a corollary of the first application.

In Section 5, we consider a conjecture about estimates for the Piltz divisor functions over number fields. It is proposed that for all number fields K and for all m the best upper bound of the error term is better than that on the assumption of the Lindelöf hypothesis (1-6).

2. Auxiliary theorem

In this section, we show some important lemmas for our argument. Let $s = \sigma + it$ and $n = [K : \mathbb{Q}]$. We use the convexity bound of the Dedekind zeta function to obtain an upper bound of the error term of the Piltz divisor function $\Delta_K^m(x)$.

It is well known that the Dedekind zeta function satisfies the functional equation

$$(2-1) \quad \zeta_K(1-s) = D_K^{s-\frac{1}{2}} 2^{n(1-s)} \pi^{-ns} \Gamma(s)^n \left(\cos \frac{\pi s}{2}\right)^{r_1+r_2} \left(\sin \frac{\pi s}{2}\right)^{r_2} \zeta_K(s),$$

where r_1 is the number of real embeddings of K and r_2 is the number of pairs of complex embeddings.

The Phragmen–Lindelöf principle and (2-1) give the well known convexity bound of the Dedekind zeta function [Rademacher 1959]: for any $\varepsilon > 0$ and $n = [K : \mathbb{Q}]$,

$$(2-2) \quad \zeta_K(\sigma + it) = \begin{cases} O_{n,\varepsilon}(|t|^{\frac{n}{2}-n\sigma+\varepsilon} D_K^{\frac{1}{2}-\sigma+\varepsilon}) & \text{if } \sigma \leq 0, \\ O_{n,\varepsilon}(|t|^{\frac{n(1-\sigma)}{2}+\varepsilon} D_K^{\frac{1-\sigma}{2}+\varepsilon}) & \text{if } 0 \leq \sigma \leq 1, \\ O_{n,\varepsilon}(|t|^\varepsilon D_K^\varepsilon) & \text{if } 1 \leq \sigma \end{cases}$$

as $|t|^n D_K \rightarrow \infty$, where K runs through number fields with $[K : \mathbb{Q}] = n$. In the previous papers, we also use this convexity bound (2-2) to estimate the distribution of ideals. In the following sections, we show some estimates for $\Delta_K^m(x)$ in the similar way to our previous papers.

Lemma 2-3 states the growth of the product of the Gamma function and trigonometric functions in the functional equation (2-1) of the Dedekind zeta function.

Lemma 2-3. *Let $\tau \in \{\cos, \sin\}$ and n be a positive integer. Then*

$$\begin{aligned} \frac{\Gamma(s)^n}{1-s} \left(\cos \frac{\pi s}{2}\right)^{r_1+r_2} \left(\sin \frac{\pi s}{2}\right)^{r_2} \\ = C n^{-ns} \Gamma\left(ns - \frac{n+1}{2}\right) \tau\left(\frac{n\pi s}{2}\right) + O_n(|t|^{-2+n\sigma-\frac{n}{2}}), \end{aligned}$$

where C is a constant and $s = \sigma + it$.

Proof. This lemma is shown from the Stirling formula and estimate for trigonometric functions. \square

Next we introduce the generalized Atkinson lemma. This lemma is quite useful for calculating integrals of the Dedekind zeta function.

Lemma 2-4 [Atkinson 1941]. *Let $y > 0$, $1 < A \leq B$, and $\tau \in \{\cos, \sin\}$, and define*

$$I = \frac{1}{2\pi i} \int_{A-iB}^{A+iB} \Gamma(s) \tau\left(\frac{\pi s}{2}\right) y^{-s} ds.$$

If $y \leq B$, then

$$I = \tau(y) + O\left(y^{-\frac{1}{2}} \min\left(\left(\log \frac{B}{y}\right)^{-1}, B^{\frac{1}{2}}\right) + y^{-A} B^{A-\frac{1}{2}} + y^{-\frac{1}{2}}\right).$$

If $y > B$, then

$$I = O\left(y^{-A} \left(B^{A-\frac{1}{2}} \min\left(\left(\log \frac{y}{B}\right)^{-1}, B^{\frac{1}{2}}\right) + A^{A-\frac{1}{2}}\right)\right).$$

Finally we introduce the following lemma to reduce the ideal counting problem to an exponential sum problem.

Lemma 2-5 [Bordellès 2015]. *Let $1 \leq L \leq R$ be a real number and f be an arithmetical function satisfying $f(m) = O(m^\varepsilon)$, and let $e(x) = \exp(2\pi i x)$ and $F = f * \mu$, where $*$ is the Dirichlet product symbol. For $a \in \mathbb{R} - \{1\}$ and $b, x \in \mathbb{R}$ and for every $\varepsilon > 0$ the following estimate holds:*

$$\sum_{m \leq R} \frac{f(m)}{m^a} \tau(2\pi x m^b) = O_{n,\varepsilon} \left(L^{1-a} + R^\varepsilon \max_{L < S \leq R} S^{-a} \right. \\ \left. \times \max_{S < S_1 \leq 2S} \max_{\substack{M, N \leq S_1 \\ MN \asymp S}} \max_{\substack{M \leq M_1 \leq 2M \\ N \leq N_1 \leq 2N}} \left| \sum_{M < m \leq M_1} F(m) \sum_{N < n \leq N_1} e(x(mn)^b) \right| \right).$$

The next proposition plays a crucial role in our computing $I_K^m(x)$. We consider the distribution of ideals of \mathbb{O}_K , where K runs through extensions with $[K : \mathbb{Q}] = n$ and some conditions. The detail of the conditions will be determined later, but they state the relation of the principal term and the error term.

Proposition 2-6. *Let $F_K = I_K^m * \mu$. For every $\varepsilon > 0$ the following estimate holds:*

$$\Delta_K^m(x) = O_{n,m,\varepsilon} \left(L^{1-\alpha} + x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}} R^\varepsilon \max_{L \leq S \leq R} S^{-\frac{mn+1}{2mn}} \right. \\ \left. \times \max_{S < S_1 \leq 2S} \max_{\substack{M, N \leq S_1 \\ MN \asymp S}} \max_{\substack{M \leq M_1 \leq 2M \\ N \leq N_1 \leq 2N}} \left| \sum_{M < l \leq M_1} F_K(m) \sum_{N < k \leq N_1} e \left(mn \left(\frac{xlk}{D_K} \right)^{\frac{1}{mn}} \right) \right| \right. \\ \left. + x^{\frac{mn-2}{2mn} + \varepsilon} D_K^{\frac{1}{n} + \varepsilon} R^{\frac{mn-2}{2mn} + \varepsilon} + x^{\frac{mn-1}{mn} + \varepsilon} D_K^{\frac{1}{n} + \varepsilon} R^{-\frac{1}{mn} + \varepsilon} \right),$$

where K runs through number fields with $[K : \mathbb{Q}] = n$ and some conditions.

Proof. Let $d_K^m(l)$ be the number of m -tuples of ideals $(\mathfrak{a}_1, \mathfrak{a}_2, \dots, \mathfrak{a}_m)$ such that the product of their ideal norms $\mathfrak{N}\mathfrak{a}_1 \cdots \mathfrak{N}\mathfrak{a}_m = l$. Then one can easily check that

$$(2-7) \quad \zeta_K(s)^m = \sum_{l=1}^{\infty} \frac{d_K^m(l)}{l^s} \quad \text{for } \Re s > 1$$

and


$$I_K^m(x) = \sum_{l \leq x} d_K^m(l).$$

Thus, Perron's formula plays a crucial role in this proof.

We consider the integral

$$\frac{1}{2\pi i} \int_C \zeta_K(s)^m \frac{x^s}{s} ds,$$

where C is the contour $C_1 \cup C_2 \cup C_3 \cup C_4$ shown below:



In a way similar to the well known proof of Perron's formula, we estimate

$$(2-8) \quad \frac{1}{2\pi i} \int_{C_1} \zeta_K(s)^m \frac{x^s}{s} ds = I_K^m(x) + O_\varepsilon\left(\frac{x^{1+\varepsilon}}{T}\right).$$

We can select the large T , so that the O -term in the right-hand side is sufficiently small. For estimating the left-hand side by using estimate (2-2), we divide it into the integrals over C_2 , C_3 , and C_4 .

First we consider the integrals over C_2 and C_4 as

$$\begin{aligned} & \left| \frac{1}{2\pi i} \int_{C_2 \cup C_4} \zeta_K(s)^m \frac{x^s}{s} ds \right| \\ & \leq \frac{1}{2\pi} \int_{-\varepsilon}^{1+\varepsilon} |\zeta_K(\sigma + iT)|^m \frac{x^\sigma}{T} d\sigma + \frac{1}{2\pi} \int_{-\varepsilon}^{1+\varepsilon} |\zeta_K(\sigma - iT)|^m \frac{x^\sigma}{T} d\sigma. \end{aligned}$$

It holds by the convexity bound of the Dedekind zeta function (2-2) that their sum is estimated as

$$\begin{aligned} (2-9) \quad & \left| \frac{1}{2\pi i} \int_{C_2 \cup C_4} \zeta_K(s)^m \frac{x^s}{s} ds \right| = O_{n,m,\varepsilon} \left(\int_{-\varepsilon}^{1+\varepsilon} (T^{mn} D_K^m)^{\frac{1-\sigma}{2} + \varepsilon} \frac{x^\sigma}{T} d\sigma \right) \\ & = O_{n,m,\varepsilon} \left(\frac{x^{1+\varepsilon} D_K^\varepsilon}{T^{1-\varepsilon}} + T^{\frac{mn}{2} - 1 + \varepsilon} D_K^{\frac{m}{2} + \varepsilon} x^{-\varepsilon} \right). \end{aligned}$$

By the Cauchy residue theorem, (2-8), and (2-9) we obtain

$$(2-10) \quad \Delta_K^m(x) = \int_{C_3} \zeta_K(s)^m \frac{x^s}{s} ds + O_{n,m,\varepsilon} \left(\frac{x^{1+\varepsilon} D_K^\varepsilon}{T^{1-\varepsilon}} + T^{\frac{mn}{2} - 1 + \varepsilon} D_K^{\frac{m}{2} + \varepsilon} x^{-\varepsilon} \right).$$

Thus, it suffices to consider the integral over C_3 as

$$\frac{1}{2\pi i} \int_{C_3} \zeta_K(s)^m \frac{x^s}{s} ds = \frac{1}{2\pi i} \int_{-\varepsilon - iT}^{-\varepsilon + iT} \zeta_K(s)^m \frac{x^s}{s} ds.$$

Changing the variable s to $1-s$, we have

$$\frac{1}{2\pi i} \int_{C_3} \zeta_K(s)^m \frac{x^s}{s} ds = \frac{1}{2\pi i} \int_{1+\varepsilon-iT}^{1+\varepsilon+iT} \zeta_K(1-s)^m \frac{x^{1-s}}{1-s} ds.$$

From functional equation (2-1), it holds that

$$\begin{aligned} \frac{1}{2\pi i} \int_{C_3} \zeta_K(s)^m \frac{x^s}{s} ds &= \frac{1}{2\pi i} \int_{1+\varepsilon-iT}^{1+\varepsilon+iT} \left(D_K^{s-\frac{1}{2}} 2^{n(1-s)} \pi^{-ns} \Gamma(s)^n \left(\cos \frac{\pi s}{2} \right)^{r_1+r_2} \right. \\ &\quad \left. \times \left(\sin \frac{\pi s}{2} \right)^{r_2} \zeta_K(s) \right)^m \frac{x^{1-s}}{1-s} ds. \end{aligned}$$

By Lemma 2-3 the integral over C_3 can be expressed as

$$\begin{aligned} &\frac{1}{2\pi i} \int_{C_3} \zeta_K(s)^m \frac{x^s}{s} ds \\ &= \frac{Cx}{2\pi i} \int_{1+\varepsilon-iT}^{1+\varepsilon+iT} D_K^{-\frac{m}{2}} \left(\frac{(2n)^{mn} \pi^{mn} x}{D_K^m} \right)^{-s} \Gamma\left(mns - \frac{mn+1}{2}\right) \tau\left(\frac{mn\pi s}{2}\right) \zeta_K(s) ds \\ &\quad + O_{n,m,\varepsilon}(D_K^{\frac{m}{2}+\varepsilon} T^{\frac{mn}{2}-1+\varepsilon} x^{-\varepsilon}). \end{aligned}$$

Changing the variable $mns - \frac{mn+1}{2}$ to s , we have

$$\begin{aligned} \frac{1}{2\pi i} \int_{C_3} \zeta_K(s)^m \frac{x^s}{s} ds &= \frac{Cx^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}}}{2\pi i} \int_{\frac{mn-1}{2}+mn\varepsilon-mniT}^{\frac{mn-1}{2}+mn\varepsilon+mniT} \left(2mn\pi \left(\frac{x}{D_K^m} \right)^{\frac{1}{mn}} \right)^{-s} \\ &\quad \times \Gamma(s) \tau\left(\frac{\pi s}{2} + \frac{(mn+1)\pi}{4}\right) \zeta_K\left(\frac{s}{mn} + \frac{mn+1}{2mn}\right) ds \\ &\quad + O_{n,m,\varepsilon}(D_K^{\frac{m}{2}+\varepsilon} T^{\frac{mn}{2}-1+\varepsilon} x^{-\varepsilon}). \end{aligned}$$

From (2-7) the function $\zeta_K(s)^m$ can be expressed as a Dirichlet series. It is absolutely and uniformly convergent on compact subsets on $\Re(s) > 1$. Therefore, we can interchange the order of summation and integral. Thus, we obtain

$$\begin{aligned} &\int \left(2mn\pi \left(\frac{x}{D_K^m} \right)^{\frac{1}{mn}} \right)^{-s} \Gamma(s) \tau\left(\frac{\pi s}{2} + \frac{(mn+1)\pi}{4}\right) \zeta_K\left(\frac{s}{mn} + \frac{mn+1}{2mn}\right) ds \\ &= \sum_{l=1}^{\infty} \frac{d_K^m(l)}{l^{\frac{mn+1}{2mn}}} \int \left(2mn\pi \left(\frac{lx}{D_K^m} \right)^{\frac{1}{mn}} \right)^{-s} \Gamma(s) \tau\left(\frac{\pi s}{2} + \frac{(mn+1)\pi}{4}\right) ds, \end{aligned}$$

where the integration is on the vertical line from $\frac{mn-1}{2} + mn\varepsilon - mniT$ to $\frac{mn-1}{2} + mn\varepsilon + mniT$. Properties of trigonometric functions lead to

$$\tau\left(\frac{\pi s}{2} + \frac{(mn+1)\pi}{4}\right) = \pm \begin{cases} \tau\left(\frac{\pi s}{2}\right) & \text{if } mn \text{ is odd,} \\ \frac{1}{\sqrt{2}}(\tau\left(\frac{\pi s}{2}\right) \pm \tau_1\left(\frac{\pi s}{2}\right)) & \text{if } mn \text{ is even,} \end{cases}$$

where $\{\tau, \tau_1\} = \{\sin, \cos\}$. Hence, it holds that

$$\begin{aligned} \frac{1}{2\pi i} \int_{C_3} \zeta_K(s) m \frac{x^s}{s} ds &= \frac{C x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}}}{2\pi i} \sum_{l=1}^{\infty} \frac{d_K^m(l)}{l^{\frac{mn+1}{2mn}}} \\ &\quad \times \int_{\frac{mn-1}{2} + mn\varepsilon - mniT}^{\frac{mn-1}{2} + mn\varepsilon + mniT} \left(2mn\pi \left(\frac{lx}{D_K^m} \right)^{\frac{1}{mn}} \right)^{-s} \Gamma(s) \tau \left(\frac{\pi s}{2} \right) ds \\ &\quad + O_{n,m,\varepsilon} \left(D_K^{\frac{m}{2} + \varepsilon} T^{\frac{mn}{2} - 1 + \varepsilon} x^{-\varepsilon} \right). \end{aligned}$$

Applying Lemma 2-4 to this integral with $y = 2mn\pi \left(\frac{lx}{D_K^m} \right)^{\frac{1}{mn}}$, $A = \frac{mn-1}{2} + mn\varepsilon$, $B = mnT$, and $T = 2\pi \left(\frac{xR}{D_K^m} \right)^{\frac{1}{mn}}$, this becomes

$$\begin{aligned} \frac{1}{2\pi i} \int_{C_3} \zeta_K(s) m \frac{x^s}{s} ds &= \frac{C x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}}}{2\pi i} \sum_{l \leq R} \frac{d_K^m(l)}{l^{\frac{mn+1}{2mn}}} \tau \left(2mn\pi \left(\frac{lx}{D_K^m} \right)^{\frac{1}{mn}} \right) \\ &\quad + O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} \sum_{l \leq R} \frac{d_K^m(l)}{l^{\frac{mn+2}{2mn}}} \min \left\{ \left(\log \frac{R}{l} \right)^{-1}, \left(\frac{Rx}{D_K^m} \right)^{\frac{1}{2mn}} \right\} \right) \\ &\quad + O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} \sum_{l \leq R} \frac{d_K^m(l)}{l^{\frac{mn+2}{2mn}}} \left(\left(\frac{R}{l} \right)^{\frac{mn-2}{2mn}} + 1 \right) \right) \\ &\quad + O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} R^{\frac{mn-2}{2mn} + \varepsilon} \sum_{l > R} \frac{d_K^m(l)}{l^{1+\varepsilon}} \min \left\{ \left(\log \frac{l}{R} \right)^{-1}, \left(\frac{Rx}{D_K^m} \right)^{\frac{1}{2mn}} \right\} \right) \\ &\quad + O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn} + \varepsilon} D_K^{\frac{1}{n} + \varepsilon} R^{\frac{mn-2}{2mn} + \varepsilon} \right). \end{aligned}$$

We evaluate three O -terms as follows. First we consider the first O -term. One can estimate $(\log \frac{R}{l})^{-1} = O(\frac{R}{R-l})$, so we obtain

$$\begin{aligned} &O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} \sum_{l \leq R} \frac{d_K^m(l)}{l^{\frac{mn+2}{2mn}}} \min \left\{ \left(\log \frac{R}{l} \right)^{-1}, \left(\frac{Rx}{D_K^m} \right)^{\frac{1}{2mn}} \right\} \right) \\ &= O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} \sum_{l \leq [R]-1} \frac{d_K^m(l)}{l^{\frac{mn+2}{2mn}}} \left(\log \frac{R}{l} \right)^{-1} \right. \\ &\quad \left. + x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} \sum_{[R] \leq l \leq R} \frac{d_K^m(l)}{l^{\frac{mn+2}{2mn}}} \left(\frac{Rx}{D_K^m} \right)^{\frac{1}{2mn}} \right) \\ &= O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} \sum_{l \leq [R]-1} \frac{d_K^m(l)}{l^{\frac{mn+2}{2mn}}} \frac{R}{R-l} + x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}} R^{\frac{1}{2mn}} \sum_{[R] \leq l \leq R} \frac{d_K^m(l)}{l^{\frac{mn+2}{2mn}}} \right) \\ &= O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} R^{\frac{mn-2}{2mn} + \varepsilon} + x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}} R^{-\frac{mn+1}{2mn}} \right). \end{aligned}$$

Next we calculate the second O -term

$$\begin{aligned} O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} \sum_{l \leq R} \frac{d_K^m(l)}{l^{\frac{mn+2}{2mn}}} \left(\left(\frac{R}{l} \right)^{\frac{mn-2}{2mn}} + 1 \right) \right) \\ = O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} R^{\frac{mn-2}{2mn}} \sum_{l \leq R} \frac{d_K^m(l)}{l} \right). \end{aligned}$$

Since it is well known that $d_K^m(l) = O(l^\varepsilon)$, we get

$$\begin{aligned} O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} \sum_{l \leq R} \frac{d_K^m(l)}{l^{\frac{mn+2}{2mn}}} \left(\left(\frac{R}{l} \right)^{\frac{mn-2}{2mn}} + 1 \right) \right) \\ = O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} R^{\frac{mn-2}{2mn}} \int_1^R \frac{t^\varepsilon}{t} dt \right) \\ = O_{n,m,\varepsilon} (x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} R^{\frac{mn-2}{2mn} + \varepsilon}). \end{aligned}$$

Finally we estimate the third O -term in a similar way to calculate the first O -term. One can estimate $(\log \frac{l}{R})^{-1} = O(\frac{R}{l-R})$, so we obtain

$$\begin{aligned} O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} R^{\frac{mn-2}{2mn} + \varepsilon} \sum_{l > R} \frac{d_K^m(l)}{l^{1+\varepsilon}} \min \left\{ \left(\log \frac{l}{R} \right)^{-1}, \left(\frac{Rx}{D_K^m} \right)^{\frac{1}{2mn}} \right\} \right) \\ = O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} R^{\frac{mn-2}{2mn} + \varepsilon} \left(\sum_{R < l \leq [R]+1} \frac{d_K^m(l)}{l^{1+\varepsilon}} \left(\frac{Rx}{D_K^m} \right)^{\frac{1}{2mn}} \right. \right. \\ \left. \left. + \sum_{[R]+2 \leq l} \frac{d_K^m(l)}{l^{1+\varepsilon}} \left(\log \frac{l}{R} \right)^{-1} \right) \right) \\ = O_{n,m,\varepsilon} \left(x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}} R^{\frac{mn-1}{2mn} + \varepsilon} \sum_{R < l \leq [R]+1} \frac{d_K^m(l)}{l^{1+\varepsilon}} \right. \\ \left. + x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} R^{\frac{mn-2}{2mn} + \varepsilon} \sum_{[R]+2 \leq l} \frac{d_K^m(l)}{l^{1+\varepsilon}} \frac{R}{l-R} \right) \\ = O_{n,m,\varepsilon} (x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}} R^{-\frac{mn+1}{2mn} + \varepsilon} + x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} R^{\frac{mn-2}{2mn} + \varepsilon}). \end{aligned}$$

From above results, we obtain

$$\begin{aligned} (2-11) \quad \frac{1}{2\pi i} \int_{C_3} \zeta_K(s)^m \frac{x^s}{s} ds = \frac{C x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}}}{2\pi i} \sum_{l \leq R} \frac{d_K^m(l)}{l^{\frac{mn+1}{2mn}}} \tau \left(2n\pi \left(\frac{lx}{D_K^m} \right)^{\frac{1}{mn}} \right) \\ + O_{n,m,\varepsilon} (x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}} R^{-\frac{mn+1}{2mn} + \varepsilon} + x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} R^{\frac{mn-2}{2mn} + \varepsilon}). \end{aligned}$$

From estimates (2-10) and (2-11), it is obtained that

$$\begin{aligned} \Delta_K^m(x) &= \frac{C x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}}}{2\pi i} \sum_{l \leq R} \frac{d_K^m(l)}{l^{\frac{mn+1}{2mn}}} \tau \left(2mn\pi \left(\frac{lx}{D_K^m} \right)^{\frac{1}{mn}} \right) \\ &\quad + O_{n,m,\varepsilon} \left(x^{\frac{mn-2}{2mn}+\varepsilon} D_K^{\frac{1}{n}+\varepsilon} R^{\frac{mn-2}{2mn}+\varepsilon} + x^{\frac{mn-1}{mn}+\varepsilon} D_K^{\frac{1}{n}+\varepsilon} R^{-\frac{1}{mn}+\varepsilon} \right). \end{aligned}$$

Next we consider the above sum. Let $F_K = d_K^m * \mu$, where $*$ is the Dirichlet product symbol. From Lemma 2-5 this becomes

$$\begin{aligned} \Delta_K^m(x) &= O_{n,m,\varepsilon} \left(L^{1-\alpha} + x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}} R^\varepsilon \max_{L \leq S \leq R} S^{-\frac{mn+1}{2mn}} \right. \\ &\quad \times \max_{S < S_1 \leq 2S} \max_{\substack{M, N \leq S_1 \\ MN \asymp S}} \max_{\substack{M \leq M_1 \leq 2M \\ N \leq N_1 \leq 2N}} \left| \sum_{M < l \leq M_1} F_K(l) \sum_{N < k \leq N_1} e \left(mn \left(\frac{xlk}{D_K^m} \right)^{\frac{1}{mn}} \right) \right| \\ &\quad \left. + x^{\frac{mn-2}{2mn}+\varepsilon} D_K^{\frac{1}{n}+\varepsilon} R^{\frac{mn-2}{2mn}+\varepsilon} + x^{\frac{mn-1}{mn}+\varepsilon} D_K^{\frac{1}{n}+\varepsilon} R^{-\frac{1}{mn}+\varepsilon} \right). \quad \square \end{aligned}$$

Let $\mathcal{G}_K(x, S)$ be the sum in the O -term, that is,

$$S^{-\frac{mn+1}{2mn}} \max_{S < S_1 \leq 2S} \max_{\substack{M, N \leq S_1 \\ MN \asymp S}} \max_{\substack{M \leq M_1 \leq 2M \\ N \leq N_1 \leq 2N}} \left| \sum_{M < l \leq M_1} F_K(l) \sum_{N < k \leq N_1} e \left(mn \left(\frac{xlk}{D_K^m} \right)^{\frac{1}{mn}} \right) \right|.$$

This proposition reduces the initial problem to an exponential sums problem. There are many results to estimate an exponential sums. In the next section, we estimate the Piltz divisor function by using some results for exponential sums established by many authors.

3. Estimate of counting function

In the last section, we showed that the error term of the Piltz divisor function $\Delta_K^m(x)$ can be expressed as an exponential sum. Let $X > 1$ be a real number, $1 \leq M < M_1 \leq 2M$ and $1 \leq N < N_1 \leq 2N$ be integers, and $(a_m), (b_n) \subset \mathbb{C}$ be sequences of complex numbers, and let $\alpha, \beta \in \mathbb{R}$. Then we define

$$(3-1) \quad \mathcal{G} = \sum_{M < m \leq M_1} a_m \sum_{N < n \leq N_1} b_n e \left(X \left(\frac{m}{M} \right)^\alpha \left(\frac{n}{N} \right)^\beta \right).$$

Lemma 3-2 [Wu 1998]. *Let $\alpha, \beta \in \mathbb{R}$ such that $\alpha\beta(\alpha-1)(\beta-1) \neq 0$, and $|a_m| \leq 1$ and $|b_n| \leq 1$ and $\mathcal{L} = \log(XMN + 2)$. Then*

$$\begin{aligned} \mathcal{L}^{-2\mathcal{G}} &= O \left((XM^3N^4)^{\frac{1}{5}} + (X^4M^{10}N^{11})^{\frac{1}{16}} + (XM^7N^{10})^{\frac{1}{11}} \right. \\ &\quad \left. + MN^{\frac{1}{2}} + (X^{-1}M^{14}N^{23})^{\frac{1}{22}} + X^{-\frac{1}{2}}MN \right). \end{aligned}$$

Next Bordellès also shows this lemma by using estimates for triple exponential sums by Robert and Sargos.

Lemma 3-3 [Bordellès 2015]. *Let $\alpha, \beta \in \mathbb{R}$ such that $\alpha\beta(\alpha-1)(\beta-1) \neq 0$, and $|a_m| \leq 1$ and $|b_n| \leq 1$. If $X = O(M)$, then*

$$(MN)^{-\varepsilon} \mathcal{G} = O\left((XM^5N^7)^{\frac{1}{8}} + N(X^{-2}M^{11})^{\frac{1}{12}} + (X^{-3}M^{21}N^{23})^{\frac{1}{24}} + M^{\frac{3}{4}}N + X^{-\frac{1}{4}}MN\right).$$

The following Srinivasan result is important for our estimating $\Delta_K^m(x)$.

Lemma 3-4 [Srinivasan 1962]. *Let N and P be positive integers and $u_n \geq 0$, $v_p > 0$, A_n , and B_p denote constants for $1 \leq n \leq N$ and $1 \leq p \leq P$. Then there exists q with properties*

$$Q_1 \leq q \leq Q_2$$

and

$$\begin{aligned} \sum_{n=1}^N A_n q^{u_n} + \sum_{p=1}^P B_p q^{-v_p} \\ = O\left(\sum_{n=1}^N \sum_{p=1}^P u_n + v_p \sqrt{A_n^{v_p} B_p^{u_n}} + \sum_{n=1}^N A_n Q_1^{u_n} + \sum_{p=1}^P B_p Q_2^{-v_p}\right). \end{aligned}$$

The constant involved in the O -symbol is less than $N + P$.

Srinivasan [1962] remarks that the inequality in Lemma 3-4 corresponds to the “best possible” choice of q in the range $Q_1 \leq q \leq Q_2$. We apply Lemma 3-4 to improve the error term $\Delta_K^m(x)$.

Theorem 3-5. *For every $\varepsilon > 0$ the following estimate holds. When $mn \geq 4$, then*

$$\Delta_K^m(x) = O_{n,m,\varepsilon}\left(x^{\frac{2mn-3}{2mn+1}+\varepsilon} D_K^{\frac{2m}{2mn+1}+\varepsilon}\right)$$

as x tends to infinity.

Proof. We note that

$$\begin{aligned} \left| \sum_{M < l \leq M_1} F_K(l) \sum_{N < k \leq N_1} e\left(mn \left(\frac{xlk}{D_K^m}\right)^{\frac{1}{mn}}\right) \right| \\ = \left| \sum_{M < l \leq M_1} F_K(l) \sum_{N < k \leq N_1} e\left(mn \left(\frac{xMN}{D_K^m}\right)^{\frac{1}{mn}} \left(\frac{l}{M}\right)^{\frac{1}{mn}} \left(\frac{k}{N}\right)^{\frac{1}{mn}}\right) \right|. \end{aligned}$$

We use the above lemmas with $X = mn \left(\frac{xMN}{D_K^m} \right)^{\frac{1}{mn}} > 0$. Let $0 \leq \alpha \leq \frac{1}{3}$; we consider four cases:

$$\begin{aligned} \text{case 1,} \quad & S^\alpha \ll N \ll S^{\frac{1}{2}}, \\ \text{case 2,} \quad & S^{\frac{1}{2}} \ll N \ll S^{1-\alpha}, \\ \text{case 3,} \quad & S^{1-\alpha} \ll N, \\ \text{case 4,} \quad & N \ll S^\alpha. \end{aligned}$$

When $S^\alpha \ll N \ll S^{\frac{1}{2}}$, we apply Lemma 3-2 and this gives

$$\begin{aligned} (3-6) \quad & S^{-\varepsilon} x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}} \mathcal{J}_K(x, S) \\ &= O_{n,m,\varepsilon} \left(x^{\frac{5mn-3}{10mn}} D_K^{\frac{3}{10n}} R^{\frac{2mn-3}{10mn}} + x^{\frac{2mn-1}{4mn}} D_K^{\frac{1}{4n}} R^{\frac{5mn-8}{32mn}} + x^{\frac{11mn-9}{22mn}} D_K^{\frac{9}{22n}} R^{\frac{6mn-9}{22mn}} \right. \\ &\quad \left. + x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}} R^{\frac{mn-1}{2mn}-\frac{1}{2}\alpha} + x^{\frac{11mn-12}{22mn}} D_K^{\frac{6}{11n}} R^{\frac{15mn-24}{44mn}} + x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} R^{\frac{mn-2}{2mn}} \right). \end{aligned}$$

When $S^{\frac{1}{2}} \ll N \ll S^{1-\alpha}$ we use Lemma 3-2 again reversing the role of M and N . We obtain the same estimate for the case that $S^\alpha \ll N \ll S^{\frac{1}{2}}$. For case 3, we use Lemma 3-3:

$$\begin{aligned} (3-7) \quad & S^{-\varepsilon} x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}} \mathcal{J}_K(x, S) \\ &= O_{n,m,\varepsilon} \left(x^{\frac{4mn-3}{8mn}} D_K^{\frac{3}{8n}} R^{\frac{mn-3}{8mn}+\frac{1}{4}\alpha} + x^{\frac{3mn-4}{6mn}} D_K^{\frac{2}{3n}} R^{\frac{5mn-8}{12mn}+\frac{1}{12}\alpha} \right. \\ &\quad \left. + x^{\frac{4mn-5}{8mn}} D_K^{\frac{5}{8n}} R^{\frac{3mn-5}{8mn}-\frac{1}{12}\alpha} + x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}} R^{\frac{mn-2}{4mn}+\frac{1}{4}\alpha} + x^{\frac{2mn-3}{4mn}} D_K^{\frac{3}{4n}} R^{\frac{2mn-3}{4mn}} \right). \end{aligned}$$

If $x^{\frac{1}{mn(1-\alpha)-1}} D_K^{-\frac{m}{mn(1-\alpha)-1}} \ll S$, the condition of Lemma 3-3, $X = O(N)$, is satisfied. Therefore, it suffices to choose $L = x^{\frac{1}{mn(1-\alpha)-1}} D_K^{-\frac{m}{mn(1-\alpha)-1}}$. For case 4, we use Lemma 3-3 again reversing the role of M and N . We obtain the same estimate for the case that $N \ll S^\alpha$. Combining (3-6) and (3-7) with Proposition 2-6, we obtain

$$\begin{aligned} (3-8) \quad \Delta_K^m(x) &= O_{n,m,\varepsilon} \left(x^{\frac{5mn-3}{10mn}} D_K^{\frac{3}{10n}} R^{\frac{2mn-3}{10mn}+\varepsilon} + x^{\frac{2mn-1}{4mn}} D_K^{\frac{1}{4n}} R^{\frac{5mn-8}{32mn}+\varepsilon} \right. \\ &\quad + x^{\frac{11mn-9}{22mn}} D_K^{\frac{9}{22n}} R^{\frac{6mn-9}{22mn}+\varepsilon} + x^{\frac{mn-1}{2mn}} D_K^{\frac{1}{2n}} R^{\frac{mn-1}{2mn}-\frac{1}{2}\alpha+\varepsilon} \\ &\quad + x^{\frac{11mn-12}{22mn}} D_K^{\frac{6}{11n}} R^{\frac{15mn-24}{44mn}+\varepsilon} + x^{\frac{mn-2}{2mn}} D_K^{\frac{1}{n}} R^{\frac{mn-2}{2mn}+\varepsilon} \\ &\quad + x^{\frac{4mn-3}{8mn}} D_K^{\frac{3}{8n}} R^{\frac{mn-3}{8mn}+\frac{1}{4}\alpha+\varepsilon} + x^{\frac{3mn-4}{6mn}} D_K^{\frac{2}{3n}} R^{\frac{5mn-8}{12mn}+\frac{1}{12}\alpha+\varepsilon} \\ &\quad + x^{\frac{4mn-5}{8mn}} D_K^{\frac{5}{8n}} R^{\frac{3mn-5}{8mn}+\frac{1}{12}\alpha+\varepsilon} + x^{\frac{2mn-3}{4mn}} D_K^{\frac{3}{4n}} R^{\frac{2mn-3}{4mn}+\varepsilon} \\ &\quad \left. + x^{\frac{mn-1}{mn}+\varepsilon} D_K^{\frac{1}{n}+\varepsilon} R^{-\frac{1}{mn}+\varepsilon} + x^{\frac{1-\alpha}{mn(1-\alpha)-1}} D_K^{-\frac{m(1-\alpha)}{mn(1-\alpha)-1}} \right). \end{aligned}$$

By Lemma 3-4 with $x^{\frac{1}{mn(1-\alpha)-1}} D_K^{-\frac{m}{mn(1-\alpha)-1}} \leq R \leq xD$ there exists R such that the error term of estimate (3-8) is much less than

$$\begin{aligned}
 & x^{\frac{2mn}{2mn+7}+\varepsilon} D_K^{\frac{2m}{2mn+7}+\varepsilon} + x^{\frac{5mn+3}{5mn+24}+\varepsilon} D_K^{\frac{5m}{5mn+24}+\varepsilon} + x^{\frac{6mn-4}{6mn+13}+\varepsilon} D_K^{\frac{6m}{6mn+13}+\varepsilon} \\
 & + x^{\frac{(1-\alpha)mn+\alpha-1}{(1-\alpha)mn+1}+\varepsilon} D_K^{\frac{(1-\alpha)m}{(1-\alpha)mn+1}+\varepsilon} + x^{\frac{15mn-17}{15mn+20}+\varepsilon} D_K^{\frac{3m}{3mn+4}+\varepsilon} + x^{\frac{mn-2}{mn}+\varepsilon} D_K^{\frac{1}{n}+\varepsilon} \\
 & + x^{\frac{(2\alpha+1)mn-2\alpha}{(2\alpha+1)mn+5}+\varepsilon} D_K^{\frac{(2\alpha+1)m}{(2\alpha+1)mn+5}+\varepsilon} + x^{\frac{(\alpha+5)mn-\alpha-7}{(\alpha+5)mn+4}+\varepsilon} D_K^{\frac{(\alpha+5)m}{(\alpha+5)mn+4}+\varepsilon} \\
 & + x^{\frac{(2\alpha+9)mn-2\alpha-12}{(2\alpha+9)mn+9}+\varepsilon} D_K^{\frac{(2\alpha+9)m}{(2\alpha+9)mn+9}+\varepsilon} + x^{\frac{2mn-3}{2mn+1}+\varepsilon} D_K^{\frac{2m}{2mn+1}+\varepsilon} \\
 & + x^{\frac{5mn(1-\alpha)-6+3\alpha}{10mn(1-\alpha)-10}+\varepsilon} D_K^{\frac{m-3m\alpha}{10mn(1-\alpha)-10}+\varepsilon} + x^{\frac{16mn(1-\alpha)-19+8\alpha}{32mn(1-\alpha)-32}+\varepsilon} D_K^{\frac{3m-8m\alpha}{32mn(1-\alpha)-32}+\varepsilon} \\
 & + x^{\frac{11mn(1-\alpha)-14+9\alpha}{22mn(1-\alpha)-22}+\varepsilon} D_K^{\frac{3m-9m\alpha}{22mn(1-\alpha)-22}+\varepsilon} + x^{\frac{1}{2}+\varepsilon} \\
 & + x^{\frac{22mn(1-\alpha)-31+24\alpha}{44mn(1-\alpha)-44}+\varepsilon} D_K^{\frac{9m-24m\alpha}{44mn(1-\alpha)-44}+\varepsilon} + x^{\frac{mn(1-\alpha)-2+2\alpha}{2mn(1-\alpha)-2}+\varepsilon} D_K^{\frac{m-2m\alpha}{2mn(1-\alpha)-2}+\varepsilon} \\
 & + x^{\frac{4mn(1-\alpha)-6+5\alpha}{8mn(1-\alpha)-8}+\varepsilon} D_K^{\frac{2m-5m\alpha}{8mn(1-\alpha)-8}+\varepsilon} + x^{\frac{6mn(1-\alpha)-9+9\alpha}{12mn(1-\alpha)-12}+\varepsilon} D_K^{\frac{3m-9m\alpha}{12mn(1-\alpha)-12}+\varepsilon} \\
 & + x^{\frac{12mn(1-\alpha)-18+17\alpha}{24mn(1-\alpha)-24}+\varepsilon} D_K^{\frac{6m-17m\alpha}{24mn(1-\alpha)-24}+\varepsilon} + x^{\frac{2mn(1-\alpha)-3+3\alpha}{4mn(1-\alpha)-4}+\varepsilon} D_K^{\frac{m-3m\alpha}{4mn(1-\alpha)-4}+\varepsilon} \\
 & + x^{\frac{1-\alpha}{mn(1-\alpha)-1}} D_K^{-\frac{m(1-\alpha)}{mn(1-\alpha)-1}}.
 \end{aligned}$$

When $mn \geq 4$ and $\alpha = \frac{mn+3}{7mn-5}$, then we have

$$\Delta_K^m(x) = O_{n,m,\varepsilon}(x^{\frac{2mn-3}{2mn+1}+\varepsilon} D_K^{\frac{2m}{2mn+1}+\varepsilon}). \quad \square$$

For $mn \geq 4$ this theorem gives new results for the Piltz divisor problem over number fields. In particular, if we fix K with $[K : \mathbb{Q}] = 4$, then we improve the estimate for $\Delta_K(x)$ as follows:

Corollary 3-9. *For any number field K with $[K : \mathbb{Q}] = 4$,*

$$\Delta_K(x) = O_{K,\varepsilon}(x^{\frac{5}{9}+\varepsilon}).$$

This result is better than Bordellès' result.

4. Application

In this section we introduce some applications of our theorems. First we obtain a uniform estimate for the ideal counting function $I_K(x)$. From the proof of Theorem 3-5, we obtain the following theorem.

Theorem 4-1. *For all $\varepsilon > 0$ for any fixed $0 \leq \beta \leq \frac{8}{2n+5} - \varepsilon$ and $C > 0$ the following holds. If K runs through number fields with $[K : \mathbb{Q}] \leq n$ and $D_K \leq Cx^\beta$, then*

$$\Delta_K(x) = O_{C,n,\varepsilon}(x^{\frac{2n-3+2\beta}{2n+1}+\varepsilon}).$$

The condition $D_K \leq Cx^\beta$ is caused by the relation between the principal term and the error term. It is well known that $I_K(x)$ is very important to estimate the distribution of relatively r -prime lattice points. We regard an ℓ -tuple of ideals $(\mathfrak{a}_1, \mathfrak{a}_2, \dots, \mathfrak{a}_\ell)$ of \mathbb{O}_K as a lattice point in K^ℓ . We say that a lattice point $(\mathfrak{a}_1, \mathfrak{a}_2, \dots, \mathfrak{a}_\ell)$ is *relatively r -prime* for a positive integer r if there exists no prime ideal \mathfrak{p} such that $\mathfrak{a}_1, \mathfrak{a}_2, \dots, \mathfrak{a}_\ell \subset \mathfrak{p}^r$. Let $V_\ell^r(x, K)$ denote the number of relatively r -prime lattice points $(\mathfrak{a}_1, \mathfrak{a}_2, \dots, \mathfrak{a}_\ell)$ such that their ideal norm $\mathfrak{N}\mathfrak{a}_i \leq x$.

B. D. Sittinger [2010] shows that

$$V_\ell^r(x, K) \sim \frac{\rho_K^\ell}{\zeta_K(r\ell)} x^\ell,$$

where ρ_K is the residue of ζ_K at $s = 1$. It is well known that

$$(4-2) \quad \rho_K = \frac{2^{r_1} (2\pi)^{r_2} h_K R_K}{w_K \sqrt{D_K}},$$

where h_K is the class number of K , R_K is the regulator of K , and w_K is the number of roots of unity in \mathbb{O}_K^* .

After that we show some results for the error term:

$$E_\ell^r(x, K) = V_\ell^r(x, K) - \frac{\rho_K^\ell}{\zeta_K(r\ell)} x^\ell.$$

In [Takeda 2017; Takeda and Koyama 2018] we consider the relation between the relatively r -prime problem and other mathematical problems. If we assume the Lindelöf hypothesis for $\zeta_K(s)$, then it holds that for all $\varepsilon > 0$

$$(4-3) \quad E_\ell^r(x, K) = \begin{cases} O_\varepsilon(x^{\frac{1}{r}(\frac{3}{2}+\varepsilon)}) & \text{if } r\ell = 2, \\ O_\varepsilon(x^{\ell-\frac{1}{2}+\varepsilon}) & \text{otherwise.} \end{cases}$$

From easy calculation, we obtain the following corollary.

Corollary 4-4. *For all $\varepsilon > 0$ and for any fixed $0 \leq \beta \leq \frac{8}{2n+5} - \varepsilon$ and $C > 0$ the following holds. If K runs through number fields with $[K : \mathbb{Q}] \leq n$ and $D_K \leq Cx^\beta$, then*

$$E_\ell^r(x, K) = \begin{cases} O_{C,n,\varepsilon}(x^{\frac{4n-2}{r(2n+1)}+\frac{4}{2n+1}\beta+\varepsilon}) & \text{if } r\ell = 2, \\ O_{C,n,\varepsilon}(x^{\ell-\frac{4}{2n+1}+\frac{2n+5-(2n+1)\ell}{2(2n+1)}\beta+\varepsilon}) & \text{otherwise.} \end{cases}$$

For the proof of this corollary, please see the proof of Theorem 4.1 of [Takeda and Koyama 2018].

5. Conjecture

Theorem 4-1 states good uniform upper bounds. It is proposed that for all number fields K the best uniform upper bound of the error term is better than that on the assumption of the Lindelöf hypothesis (1-6).

Conjecture 5-1. *If K runs through number fields with $D_K < x$, then*

$$\Delta_K^m(x) = o(x^{\frac{1}{2}}).$$

From estimate (1-5), this conjecture may give the best estimate for uniform upper bound of $\Delta_K^m(x)$. As we remarked above (Theorem 1-2) this conjecture is very difficult even when K is fixed and $m = 1$.

References

- [Atkinson 1941] F. V. Atkinson, “A divisor problem”, *Quart. J. Math.* **12** (1941), 193–200. MR Zbl
- [Bordellès 2015] O. Bordellès, “On the ideal theorem for number fields”, *Funct. Approx. Comment. Math.* **53**:1 (2015), 31–45. MR Zbl
- [Bourgain and Watt 2017] J. Bourgain and N. Watt, “Mean square of zeta function, circle problem and divisor problem revisited”, preprint, 2017. arXiv
- [Girstmair et al. 2005] K. Girstmair, M. Kühleitner, W. Müller, and W. G. Nowak, “The Piltz divisor problem in number fields: an improved lower bound by Soundararajan’s method”, *Acta Arith.* **117**:2 (2005), 187–206. MR Zbl
- [Huxley 2002] M. N. Huxley, “Integer points, exponential sums and the Riemann zeta function”, pp. 275–290 in *Number theory for the millennium, II* (Urbana, IL, 2000), edited by M. A. Bennett et al., Peters, Natick, MA, 2002. MR Zbl
- [Lao 2010] H. Lao, “On the distribution of integral ideals and Hecke Grössencharacters”, *Chin. Ann. Math. Ser. B* **31**:3 (2010), 385–392. MR Zbl
- [Müller 1988] W. Müller, “On the distribution of ideals in cubic number fields”, *Monatsh. Math.* **106**:3 (1988), 211–219. MR Zbl
- [Nowak 1993] W. G. Nowak, “On the distribution of integer ideals in algebraic number fields”, *Math. Nachr.* **161** (1993), 59–74. MR Zbl
- [Rademacher 1959] H. Rademacher, “On the Phragmén–Lindelöf theorem and some applications”, *Math. Z.* **72** (1959), 192–204. MR Zbl
- [Sittinger 2010] B. D. Sittinger, “The probability that random algebraic integers are relatively r -prime”, *J. Number Theory* **130**:1 (2010), 164–171. MR Zbl
- [Srinivasan 1962] B. R. Srinivasan, “On van der Corput’s and Nieland’s results on the Dirichlet’s divisor problem and the circle problem”, *Proc. Natl. Inst. Sci. India A* **28**:5 (1962), 732–742. Zbl
- [Takeda 2017] W. Takeda, “Visible lattice points and the extended Lindelöf hypothesis”, *J. Number Theory* **180** (2017), 297–309. MR Zbl
- [Takeda and Koyama 2018] W. Takeda and S.-y. Koyama, “Estimates of lattice points in the discriminant aspect over abelian extension fields”, *Forum Math.* **30**:3 (2018), 767–773. MR Zbl
- [Wu 1998] J. Wu, “On the average number of unitary factors of finite abelian groups”, *Acta Arith.* **84**:1 (1998), 17–29. MR Zbl

Received July 23, 2018. Revised August 29, 2018.

WATARU TAKEDA
DEPARTMENT OF MATHEMATICS
NAGOYA UNIVERSITY
CHIKUSA-KU
NAGOYA
JAPAN
d18002r@math.nagoya-u.ac.jp

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2019 is US \$490/year for the electronic version, and \$665/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 301 No. 2 August 2019

New applications of extremely regular function spaces	385
TROND A. ABRAHAMSEN, OLAV NYGAARD and MÄRT PÖLDVERE	
Regularity and upper semicontinuity of pullback attractors for a class of nonautonomous thermoelastic plate systems	395
FLANK D. M. BEZERRA, VERA L. CARBONE, MARCELO J. D. NASCIMENTO and KARINA SCHIABEL	
Variations of projectivity for C^* -algebras	421
DON HADWIN and TATIANA SHULMAN	
Lower semicontinuity of the ADM mass in dimensions two through seven	441
JEFFREY L. JAUREGUI	
Boundary regularity for asymptotically hyperbolic metrics with smooth Weyl curvature	467
XIAOSHANG JIN	
Geometric transitions and SYZ mirror symmetry	489
ATSUSHI KANAZAWA and SIU-CHEONG LAU	
Self-dual Einstein ACH metrics and CR GJMS operators in dimension three	519
TAIJI MARUGAME	
Double graph complex and characteristic classes of fibrations	547
TAKAHIRO MATSUYUKI	
Integration of modules I: stability	575
DMITRIY RUMYNIN and MATTHEW WESTAWAY	
Uniform bounds of the Piltz divisor problem over number fields	601
WATARU TAKEDA	
Explicit Whittaker data for essentially tame supercuspidal representations	617
GEO KAM-FAI TAM	
K-theory of affine actions	639
JAMES WALDRON	
Optimal decay estimate of strong solutions for the 3D incompressible Oldroyd-B model without damping	667
RENHUI WAN	
Triangulated categories with cluster tilting subcategories	703
WUZHONG YANG, PANYUE ZHOU and BIN ZHU	
Free Rota–Baxter family algebras and (tri)dendriform family algebras	741
YUANYUAN ZHANG and XING GAO	



0030-8730(201908)301:2;1-G