Pacific Journal of Mathematics

A CHARACTERIZATION OF FUCHSIAN ACTIONS BY TOPOLOGICAL RIGIDITY

KATHRYN MANN AND MAXIME WOLFF

Volume 302 No. 1

September 2019

A CHARACTERIZATION OF FUCHSIAN ACTIONS BY TOPOLOGICAL RIGIDITY

KATHRYN MANN AND MAXIME WOLFF

We give a simple proof that any rigid representation of $\pi_1(\Sigma_g)$ in Homeo⁺(S_1) with Euler number at least g is necessarily semiconjugate to a discrete, faithful representation into PSL(2, \mathbb{R}). Combined with earlier work of Matsumoto, this precisely characterizes Fuchsian actions by a topological rigidity property. We have proved this result in greater generality, but with a much more involved proof, in arxiv:1710.04902.

1. Introduction

Let Σ_g be a surface of genus $g \ge 2$, and let $\Gamma_g = \pi_1(\Sigma_g)$. The *representation* space Hom(Γ_g , Homeo⁺(S^1)) is the set of all actions of Γ_g on S^1 by orientationpreserving homeomorphisms, equipped with the compact-open topology. This is also the space of *flat topological circle bundles* over Σ_g , or equivalently, the space of circle bundles with a foliation transverse to the fibers. The *Euler class* of a representation $\rho \in \text{Hom}(\Gamma_g, \text{Homeo}^+(S^1))$ is defined to be the Euler class of the associated bundle, and the *Euler number* $eu(\rho)$ is the integer obtained by pairing the Euler class with the fundamental class of the surface. The classical Milnor–Wood inequality [Milnor 1958; Wood 1971] is the statement that the absolute value of the Euler number of a flat bundle is bounded by the absolute value of the Euler characteristic of the surface.

While the Euler number determines the topological type of a flat S^1 bundle, it does not determine its flat structure — except in the special case where the Euler number is maximal, i.e., equal to $\pm(2g-2)$. In this case, a celebrated result of Matsumoto states that for any representation ρ with $eu(\rho) = \pm(2g-2)$, there is a continuous, degree one, monotone map $h: S^1 \rightarrow S^1$ such that

(1)
$$h \circ \rho = \rho_F \circ h,$$

where ρ_F is *Fuchsian*, meaning a faithful representation of Γ_g onto a cocompact lattice in PSL(2, \mathbb{R}). (We view PSL(2, \mathbb{R}) \subset Homeo⁺(S^1) via the action on $\mathbb{R}P^1 \cong S^1$ by Möbius transformations.)

MSC2010: 20H10, 37E10, 37E45, 57S25, 58D29.

Keywords: rigidity, geometricity, Euler class, surface group actions on the circle.

An important consequence of Matsumoto's theorem is that representations with maximal Euler number are dynamically stable or rigid in the following sense.

Definition 1.1. Let Γ be a discrete group. A representation $\rho : \Gamma \to \text{Homeo}^+(S^1)$ is called *path-rigid* if its path-component in Hom(Γ , Homeo⁺(S^1)) consists of a single semiconjugacy class.

Semiconjugacy is the equivalence relation generated by the property shared by ρ and ρ_F in (1) above; we recall the precise definition in Section 2. As semiconjugacy classes are connected in Hom(Γ_g , Homeo⁺(S^1)), path-rigid representations are precisely those whose path-component is as small as possible.

The purpose of this article is to prove the following converse to Matsumoto's result.

Theorem 1.2. Let $\rho : \Gamma_g \to \text{Homeo}^+(S^1)$ be a path-rigid representation, with $|\text{eu}(\rho)| \ge g$. Then $\text{eu}(\rho)$ is maximal, i.e., $|\text{eu}(\rho)| = 2g - 2$, and ρ is semiconjugate to a discrete, faithful representation into PSL(2, \mathbb{R}).

As shown in [Mann 2015], any 2-fold lift of a Fuchsian representation is pathrigid and has Euler class g - 1; hence the inequality $|eu(\rho)| \ge g$ is optimal for this statement.

A stronger, but equally natural notion of rigidity comes from considering the *character space*, $X(\Gamma_g, \text{Homeo}^+(S^1))$, defined as the largest Hausdorff quotient of the quotient $\text{Hom}(\Gamma_g, \text{Homeo}^+(S^1))/\text{Homeo}^+(S^1)$. We say a representation is *rigid* if its image in $X(\Gamma_g, \text{Homeo}^+(S^1))$ is an isolated point. In [Mann and Wolff 2017], we prove that all rigid representations are semiconjugate to the *k*-fold lift of a Fuchsian representation, for some divisor *k* of 2g - 2; and that the weaker hypothesis of path-rigidity is sufficient provided the Euler class is nonzero. This is a more general statement than Theorem 1.2 here, but the proof in [Mann and Wolff 2017] is long and involved. This article gives a much easier, self-contained proof of this partial result. The assumption $|eu(\rho)| \ge g$ greatly simplifies the situation, as it implies in particular that many elements of the group have north-south dynamics. In fact, our assumption here can be replaced with an a priori strictly weaker assumption on the dynamics of ρ , phrased in terms of rotation numbers of elements, as follows.

Theorem 1.3. Suppose $\rho : \Gamma_g \to \text{Homeo}^+(S^1)$ is path-rigid. If there exist based simple closed curves $a, b \in \Gamma_g$ with intersection number 1 and such that

$$\widetilde{\text{rot}}[\rho(a), \rho(b)] = \pm 1,$$

then $eu(\rho) = \pm (2g - 2)$, and ρ is semiconjugate to a Fuchsian representation.

Commutators of elements of Homeo⁺(S^1) have a well defined translation number, as we will recall in Section 2A. The hypothesis $\tilde{rot}[\rho(a), \rho(b)] = \pm 1$ is equivalent to the statement that the restriction of the representation to the torus defined by *a*

and b is semiconjugate to a standard Fuchsian one (see [Matsumoto 2016]). Thus, one can think of the statement above as a local-to-global result: the local condition that a torus is Fuchsian, together with path-rigidity, implies the global statement that the representation is Fuchsian.

Outline. In Section 2 we recall standard material on dynamics of groups acting on the circle, including rotation numbers and the Euler number for actions of surface groups. We then introduce important tools for the proof of Theorem 1.3, and give a quick proof that Theorem 1.3 implies Theorem 1.2.

Sections 3 through 5 are devoted to the proof of Theorem 1.3. Given a representation ρ satisfying the hypotheses of Theorem 1.3, we proceed as follows:

1. After modifying ρ by a semiconjugacy, we show there exists $a \in \Gamma_g$ represented by a nonseparating simple closed curve such that $\rho(a)$ is *hyperbolic*, meaning that it is conjugate to a hyperbolic element of PSL(2, \mathbb{R}).

2. Using step 1, we show that (again after semiconjugacy of ρ), any $\gamma \in \Gamma_g$ represented by a nonseparating simple closed curve has the property that $\rho(\gamma)$ is hyperbolic. These two first steps are done in Section 3.

3. Next, in Section 4, we start to "reconstruct the surface", showing that the arrangement of attracting and repelling points of hyperbolic elements $\rho(\gamma)$, as γ ranges over simple closed curves, mimics that of a Fuchsian representation.

4. Finally, in Section 5 we show that the restriction of ρ to small subsurfaces is semiconjugate to a Fuchsian representation; this is then improved to a global result by additivity of the *relative Euler class*.

Throughout this paper, whenever we say "deformation", we mean deformation along a continuous path in Hom(Γ_g , Homeo⁺(S^1)).

2. Preliminaries

This section gives a quick review of basic concepts used later in the text. The only material that is not standard is the *based intersection number* discussed in Section 2D.

2A. *Rotation numbers and the Euler number.* Most of the material in Sections 2A and 2B is covered in more detail in [Ghys 2001] and [Mann 2018].

Let $\text{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$ denote the group of homeomorphisms of \mathbb{R} that commute with integer translations; this is a central extension of $\text{Homeo}^+(S^1)$ by \mathbb{Z} . The primary dynamical invariant of such homeomorphisms is the translation or rotation number, whose use can be traced back to work of Poincaré [1885, Chapitre XV]. If $\tilde{g} \in \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$ and $x \in \mathbb{R}$, the *translation number* of \tilde{g} is defined by $\tilde{\text{rot}}(\tilde{g}) := \lim_{n \to \infty} (\tilde{g}^n(x))/n$; this limit exists and does not depend on x. If $g \in \text{Homeo}^+(S^1)$, its *rotation number* is defined by $\text{rot}(g) := \tilde{\text{rot}}(\tilde{g}) \mod \mathbb{Z}$, where \tilde{g} is any lift of g. The translation number is invariant under conjugacy (and under semiconjugacy), and restricts to a morphism on every abelian subgroup of $\text{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$. On the whole group it is a *quasimorphism*, as it satisfies the following inequality.

Lemma 2.1 (see [Calegari and Walker 2011, Theorem 3.9]). Let $f, g \in \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$. Then $|\widetilde{\text{rot}}(fg) - \widetilde{\text{rot}}(f) - \widetilde{\text{rot}}(g)| \le 1$, and $-1 \le \widetilde{\text{rot}}([f, g]) \le 1$.

The second inequality is a direct consequence of the first. This in turn was implicit already in [Wood 1971]. An optimal inequality, which depends on the values of $\tilde{rot}(f)$ and $\tilde{rot}(g)$, is obtained in [Calegari and Walker 2011].

One way of defining the Euler number of a representation is in terms of translation numbers. This was perhaps first observed by Milnor and Wood [1958; 1971], who showed the following. For the purposes of this work, the reader may take this as the definition of the Euler number.

Proposition 2.2. Consider a standard presentation

$$\Gamma_g = \left\langle a_1, b_1, \ldots, a_g, b_g \mid \prod_i [a_i, b_i] \right\rangle.$$

Let $\rho \in \text{Hom}(\Gamma_g, \text{Homeo}^+(S^1))$, and let $\rho(a_i)$ and $\rho(b_i)$ be any lifts of $\rho(a_i)$ and $\rho(b_i)$ to $\text{Homeo}^+_{\mathbb{Z}}(\mathbb{R})$. Then the **Euler number** $\operatorname{eu}(\rho)$ is given by

$$\operatorname{eu}(\rho) = \operatorname{\widetilde{rot}}([\rho(\overline{a_1}), \rho(\overline{b_1})] \cdots [\rho(\overline{a_g}), \rho(\overline{b_g})]).$$

Note that, for any f and g in Homeo⁺(S^1), the value of the commutator $[\tilde{f}, \tilde{g}] \in$ Homeo⁺_Z(\mathbb{R}) is independent of the choice of lifts \tilde{f} and \tilde{g} . Abusing notation slightly, we will often denote its translation number by $\tilde{rot}([f, g])$ (as in the statement of Theorem 1.3). Thus, in the statement above, the translation by an integer, $[\rho(a_1), \rho(b_1)] \cdots [\rho(a_g), \rho(b_g)]$, is independent of the choices of lifts. The Euler number $eu(\rho)$ is then simply the magnitude of this translation.

As remarked in the introduction, the Milnor–Wood inequality is the statement that $|eu(\rho)| \le 2g - 2$; it is a consequence of Lemma 2.1.

Though unimportant in the preceding remarks, in what follows we will need to fix a convention for commutators and group multiplication.

Convention 2.3. We read words in Γ_g from right to left, so that group multiplication coincides with function composition. We set the notation for a commutator as

$$[a, b] := b^{-1}a^{-1}ba.$$

2B. Dynamics of groups acting on S^1 .

Definition 2.4 [Ghys 1987]. Let Γ be a group. Two representations ρ_1 , ρ_2 in Hom(Γ , Homeo⁺(S^1)) are *semiconjugate* if there is a monotone (possibly non-continuous or noninjective) map $\tilde{h} : \mathbb{R} \to \mathbb{R}$ such that $\tilde{h}(x+1) = \tilde{h}(x) + 1$ for

all $x \in \mathbb{R}$, and such that, for all $\gamma \in \Gamma$, there are lifts $\rho_1(\gamma)$ and $\rho_2(\gamma)$ such that $\tilde{h} \circ \rho_1(\gamma) = \rho_2(\gamma) \circ \tilde{h}$.

Ghys gave an (incorrect, as he himself later noted [2001]) version of this definition in the introduction of [Ghys 1987]; but his text becomes correct and consistent upon replacing it by Definition 2.4. He proved that semiconjugacy is an equivalence relation on Hom(Γ , Homeo⁺(S^1)), and it follows from his [1987, Propositions 2.2 and 2.3; 2001, Proposition 5.8] that this is the relation generated by the relationship shared by ρ and ρ_F in (1) of Section 1; this latter equivalence relation was used by other authors as a definition of semiconjugacy (see, e.g., [Calegari 2006]). Historical elements, and more discussion on the theme of semiconjugacy can be found in [Bucher et al. 2016].

The next proposition states a useful dynamical trichotomy for groups acting on the circle, which in particular can be used to explain when a semiconjugacy map can be taken to be continuous. As it is classical, we do not repeat the proof; the reader may refer to [Ghys 2001, Proposition 5.6].

Proposition 2.5. Let $G \subset \text{Homeo}^+(S^1)$. Then exactly one of the following holds:

- (i) G has a finite orbit.
- (ii) G is minimal, meaning that all orbits are dense.
- (iii) There is a unique compact G-invariant subset of S^1 contained in the closure of any orbit, on which G acts minimally. This set is homeomorphic to a Cantor set and called the **exceptional minimal set** for G.

In case (iii), defining h to be a map that collapses each interval in the complement of the exceptional minimal set to a point gives the following (we leave the proof as an exercise; see, e.g., [Ghys 2001, Proposition 5.8; 1987, Proposition 2.2] for more detail).

Proposition 2.6. Let $\rho: G \to \text{Homeo}^+(S^1)$ be a homomorphism such that $\rho(G)$ has an exceptional minimal set. Then ρ is semiconjugate to a homomorphism ν whose image is minimal. Moreover, provided that ν is minimal, any semiconjugacy h to any representation ρ' such that $h \circ \rho' = \nu \circ h$ is necessarily continuous.

We will make frequent use of the following two consequences of Proposition 2.6.

Corollary 2.7. Suppose that ρ and ρ' are semiconjugate representations. If both ρ and ρ' are minimal, then they are **conjugate**.

Corollary 2.8. Let $\rho \in \text{Hom}(\Gamma_g, \text{Homeo}^+(S^1))$ be a path-rigid representation. Then ρ is semiconjugate to a minimal representation.

Proof. Corollary 2.7 follows immediately from Proposition 2.6. We now prove Corollary 2.8. Using Propositions 2.5 and 2.6, it suffices to show that a representation with a finite orbit is not path-rigid. If ρ has a finite orbit, then we

may perform the Alexander trick, replacing the points of the periodic orbit with intervals and collapsing the complementary intervals, to continuously deform ρ into a representation with image in a conjugate *K* of SO(2). As Hom(Γ_g, K) = K^{2g} , the representation ρ can be deformed arbitrarily within this space, in particular to a representation which is not semiconjugate.

Following Corollary 2.8, in the proof of Theorem 1.3 we will occasionally make the (justified) assumption that a path-rigid representation ρ is also minimal.

2C. Deforming actions of surface groups. Let $\gamma \in \Gamma_g$ be a based, simple loop. Cutting Σ_g along γ decomposes Γ_g into an amalgamated product $\Gamma_g = A *_{\langle \gamma \rangle} B$ if γ is separating, and an HNN-extension $A *_{\langle \gamma \rangle}$ if not. In both cases, A and B are free groups. As there is no obstruction to deforming a representation of a free group into any topological group, deforming a representation $\rho : \Gamma_g \to \text{Homeo}^+(S^1)$ amounts to deforming the restriction(s) of ρ on A (and B, if γ separates), subject to the single constraint that these should agree on γ .

The following explicit deformations are analogous to special cases of *bending deformations* from the theory of quasi-Fuchsian and Kleinian groups.

Definition 2.9. (bending deformations) Let $\rho : \Gamma_g \to \text{Homeo}^+(S^1)$.

- (1) Separating curves. Let $\gamma = c \in \Gamma_g$ represent a separating simple closed curve with $\Gamma_g = A *_{\langle c \rangle} B$. Let c_t be a one-parameter group of homeomorphisms commuting with $\rho(c)$. Define ρ_t to agree with ρ on A, and to be equal to $c_t \rho c_t^{-1}$ on B.
- (2) *Nonseparating curves.* Let γ = a be a nonseparating curve, and let b be a nonseparating curve such that a and b are standard generators of a once-holed torus embedded in Σ_g (equivalently, the first two generators of a standard generating set of Γ_g). Let c = [a, b], and let A = ⟨a, b⟩ ⊂ Γ_g; we write again Γ_g = A *_{⟨c⟩} B. Let a_t be a one-parameter group commuting with ρ(a) and define ρ_t to agree with ρ on B and on ⟨a⟩, and define ρ_t(b) = a_tρ(b).

In both cases, we call this deformation of ρ a *bending along* γ .

In particular, if γ_t is a one-parameter group with $\gamma_1 = \rho(\gamma)$, then the deformation given above is the precomposition of ρ with τ_{γ_*} , where τ_{γ} is the Dehn twist along γ . Note that we have made a specific (though arbitrary) choice realizing the Dehn twist as an automorphism of Γ_g . This will allow us to do specific computations, for which having a twist defined only up to inner automorphism would not suffice. (See the discussion on based curves in the next subsection for more along these lines.)

Not every $f \in \text{Homeo}^+(S^1)$ embeds in a one-parameter group. However, every element with at least one fixed point does. Indeed, $S^1 \setminus \text{Fix}(f)$ is then a union of

intervals on which the action of f is conjugated to the map $\mathbb{R} \to \mathbb{R}$, $t \mapsto t+1$ or its inverse, and it is easy to build a one-parameter group out of this observation; see, e.g., [Ghys 2001, Proposition 5.10] for more detail. This is the situation in which we will typically apply bending deformations in this article.

The next corollary is used frequently in the proof of Theorem 1.3.

Corollary 2.10. Suppose that ρ is a path-rigid, minimal representation. Let ρ_t be a bending deformation along a, using a deformation a_t , with $a_1 = \rho(a)^N$ for some $N \in \mathbb{Z}$. Then ρ_1 is conjugate to ρ .

Proof. By the discussion above, ρ_1 agrees with precomposition of ρ with an automorphism of Γ_g , so has the same image. Corollary 2.7 now implies that these are conjugate.

2D. *Based curves, chains, and Fuchsian tori.* If *a* and *b* are simple closed curves on Σ_g , the familiar *geometric intersection number* is the minimum value of $|a' \cap b'|$, where a' and b' are any curves freely homotopic to *a* and *b*, respectively. It is well known that if *a* and *b* are nonseparating simple closed curves with geometric intersection number 1, then there is a subsurface $T \subset \Sigma$ homeomorphic to a torus with one boundary component with fundamental group (freely) generated by *a* and *b*. (See, e.g., [Farb and Margalit 2012, Section 1.2.3])

As mentioned earlier, the fact that we are working with specific representations, rather than conjugacy classes of elements, forces us to take the basepoint and orientation of curves into account. Although our notation $\Gamma_g = \pi_1(\Sigma_g)$ does not mention a basepoint, all elements of $\pi_1(\Sigma_g)$ will henceforth always be assumed based, and we will use the following variation on the standard definition of intersection number.

Definition 2.11 (based intersection number). Let $a, b \in \Gamma_g$. We write i(a, b) = 0 if we can represent a and b by differentiable maps $a, b : [0, 1] \to \Sigma_g$, based at the base point, whose restrictions to [0, 1) are injective, and such that the cyclic order of their tangent vectors at the base point is either (a'(0), -a'(1), b'(0), -b'(1)) or (a'(0), -a'(1), -b'(1), b'(0)), or the reverse of one of these.

If, instead, the cyclic order of tangent vectors is (a'(0), b'(0), -a'(1), -b'(1)) or the reverse, we write i(a, b) = 1 and i(a, b) = -1, respectively.

This is a somewhat ad hoc definition. In particular, i(a, b) is left undefined for many pairs (a, b).

Definition 2.12. A *directed k-chain* in Σ_g is a *k*-tuple $(\gamma_1, \ldots, \gamma_k)$ of elements of Γ_g that can be represented by the images of the edges under an embedding (possibly orientation-reversing, but respecting the orientation of the edges) of the fat graph shown in Figure 1.

In particular, $i(\gamma_i, \gamma_j) = \pm 1$ if |j - i| = 1, and 0 otherwise. Note that we do not



Figure 1. A directed chain of length 5.

require that the embedding be π_1 -injective. For example, whenever $i(\gamma_1, \gamma_2) = 1$, then $(\gamma_1, \gamma_2, \gamma_1^{-1})$ is a (rather degenerate) directed 3-chain.

These *k*-chains will be useful especially to study bending deformations that realize sequences of Dehn twists. Whenever $(\gamma_1, \ldots, \gamma_k)$ is a directed *k*-chain, the Dehn twist along the curve γ_i may be described by an automorphism of Γ_g leaving invariant the elements γ_j for $|j - i| \ge 2$ and j = i, and mapping γ_{i-1} to $\gamma_i^{-1}\gamma_{i-1}$, and γ_{i+1} to $\gamma_{i+1}\gamma_i$.

Notation 2.13. Let $i(a, b) = \pm 1$. Then their commutator [a, b] bounds a genus 1 subsurface (well-defined up to homotopy) containing *a* and *b*. We denote this surface by T(a, b).

Definition 2.14. We call any representation $\rho : \pi_1(T(a, b)) \to \text{PSL}(2, \mathbb{R})$ arising from a complete hyperbolic structure of infinite volume on T(a, b) a *standard Fuchsian representation of a once-punctured torus*. Similarly, we say that ρ : $\Gamma_g \to \text{PSL}(2, \mathbb{R})$ is *standard Fuchsian* if it comes from a hyperbolic structure on Σ_g .

Convention 2.15. We assume Σ_g is oriented; hence standard Fuchsian representations of Γ_g have Euler number -2g + 2, and are all conjugate in Homeo⁺(S^1). Similarly, T(a, b) inherits an orientation, so all its standard Fuchsian representations are conjugate in Homeo⁺(S^1).

Definition 2.16. We say that $\rho : \Gamma_g \to \text{Homeo}^+(S^1)$ has a *Fuchsian torus* if there exist two simple closed curves $a, b \in \Gamma_g$, with $i(a, b) = \pm 1$ and such that $\widetilde{\text{rot}}([\rho(a), \rho(b)]) = \pm 1$.

The terminology "Fuchsian torus" in Definition 2.16 comes from the following observation by Matsumoto.

Observation 2.17 [Matsumoto 1987]. Let α , $\beta \in \text{Homeo}^+(S^1)$ satisfy $\tilde{\text{rot}}([\alpha, \beta]) = \pm 1$. Then α and β generate a free group, and, up to reversing the orientation of S^1 , this group is semiconjugate to a standard Fuchsian representation of a one-holed torus T(a, b) with $\rho(a) = \alpha$ and $\rho(b) = \beta$.

The proof is not difficult; an easily readable sketch is given in [Matsumoto 2016, §3].

The next lemma shows the existence of such a torus is guaranteed, provided the absolute value of the Euler number of a representation is sufficiently high.

Lemma 2.18. If $|eu(\rho)| \ge g$ then ρ has a Fuchsian torus.

Proof. If $eu(\rho) \ge g$, then conjugating ρ by an orientation-reversing homeomorphism of S^1 gives a representation with Euler number at most -g. Thus, we may assume that $eu(\rho) \le -g$. Let $f \in \text{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$. It is an easy consequence of the definition of rot that rot(f) > 0 if and only if f(x) > x for all $x \in \mathbb{R}$. Hence if $f_1, \ldots, f_g \in$ $Homeo_{\mathbb{Z}}^+(\mathbb{R})$ satisfy $rot(f_i) > 0$ for all i, then $rot(f_1 \cdots f_g) > 0$.

By composing such f_i by the translation by -1, which is central in Homeo⁺_Z(\mathbb{R}), we deduce that if $\widetilde{rot}(f_i) > -1$ for all *i* then $\widetilde{rot}(f_1 \cdots f_g) > -g$. Now let ρ be a representation, and let $f_i = [\rho(a_i), \rho(b_i)]$, where a_i, b_i are standard generators for Γ_g . Then the inequality $\operatorname{eu}(\rho) \leq -g$ implies $\widetilde{rot}(f_i) \leq -1$ for some *i*. As the maximum absolute value of the rotation number of a commutator is 1 by Lemma 2.1, we in fact have $\widetilde{rot}(f_i) = -1$ for some *i*.

Lemma 2.18 immediately shows that Theorem 1.3 implies Theorem 1.2. The rest of this work is devoted to the proof of Theorem 1.3.

3. Steps 1 and 2: Existence and abundance of hyperbolic elements

Definition 3.1. We say a homeomorphism $f \in \text{Homeo}^+(S^1)$ is *hyperbolic* if it is conjugate to a hyperbolic element of PSL(2, \mathbb{R}), i.e., it has one *attracting fixed point* $f_+ \in S^1$ and one *repelling fixed point* $f_- \neq f_+$ such that $\lim_{n \to +\infty} f^n(x) = f_+$ for all $x \neq f_-$, and $\lim_{n \to +\infty} f^{-n}(x) = f_-$ for all $x \neq f_+$.

The first step of the proof of Theorem 1.3 is to show that a rigid, minimal representation has very many hyperbolic elements.

Lemma 3.2. Let T(a, b) be a one-holed torus subsurface, and let $A = \pi_1 T(a, b)$. Suppose $\rho : A \to \text{Homeo}^+(S^1)$ is semiconjugate to a standard Fuchsian representation, as in Definition 2.14. Then there exists a continuous deformation ρ_t with $\rho_0 = \rho$ such that

- (i) $\rho_1(a)$ is hyperbolic, and
- (ii) there exists a continuous family of homeomorphisms $f_t \in \text{Homeo}^+(S^1)$ such that $\rho_t([a, b]) = f_t \rho([a, b]) f_t^{-1}$ for all t.

Proof. Let *c* denote the commutator [a, b]. Let $\overline{\rho}$ denote the minimal representation (unique up to conjugacy) that is semiconjugate to ρ . Since ρ is semiconjugate to a standard Fuchsian representation, we may suppose $\overline{\rho}$ is a representation corresponding to a *finite volume* complete hyperbolic structure on T(a, b). By Proposition 2.6, there is a continuous map $h: S^1 \to S^1$, collapsing each component of the exceptional minimal set for ρ to a point, satisfying $h\rho = \overline{\rho}h$. Let x_+ and x_- be the endpoints

of the axis of $\bar{\rho}(a)$, and X_+ and X_- the preimages under *h* of their orbits $\rho(A)x_+$ and $\rho(A)x_-$.

Note that X_+ and X_- are both $\rho(A)$ -invariant sets and their images under h are the attractors (respectively, repellers) of closed curves in T(a, b) conjugate to a. Moreover, for this reason, X_+ and X_- lie in a single connected component of $S^1 \setminus \text{Fix}(\rho(c))$, and the interiors of the intervals that make up X_+ and X_- are disjoint from the exceptional minimal set of ρ .

Define a continuous family of continuous maps $h_t : S^1 \to S^1$, with $h_0 = id$, as follows: We define h_t to be the identity on the complement of the connected component of $S^1 \setminus \text{Fix}(\rho(c))$ containing X_+ and X_- , and for each interval I of X_+ or of X_- , have h_t be a homotopy contracting that interval so that $h_1(I)$ is a point. To make this precise, one needs to fix an identification of the target of h_t with the standard unit circle. Let J be the connected component of $S^1 \setminus \text{Fix}(\rho(c))$ that contains the exceptional minimal set of $\rho(A)$. Define h_t to rescale the length of each connected component of X_+ or X_- by a factor of (1 - t) and rescale the complement of $X_+ \cup X_-$ in J so that the total length of J remains unchanged. This gives us the desired map h_t which is the identity outside of J, and contracts intervals of X_+ and X_- to points.

Now define ρ_t by $h_t \rho(g) h_t^{-1} = \rho_t(g)$ for $t \in [0, 1)$. We claim that there is a unique $\rho_1(g)$ satisfying $h_1\rho(g) = \rho_1(g)h_1$. Indeed, $\rho(g)$ permutes the complementary intervals of the exceptional minimal set for ρ , so letting $h_1^{-1}(x)$ denote the preimage of x by h_1 (which is either a point or an open interval complementary to the exceptional minimal set), $h_1\rho(g)h_1^{-1}(x)$ is always a single point, and $h_1\rho(g)h_1^{-1}$ defines in this way a homeomorphism, which we denote by $\rho_1(g)$. It is easily verified that $\rho_t(g)$ approaches $\rho_1(g)$ as $t \to 1$. By construction, $\rho_1(a)$ is hyperbolic, and $\rho_t(c)$ is conjugate to a translation on the interval J defined above (and hence its restriction to J is conjugate to $\rho(c)|_J$), and $\rho_t(c)$ restricted to $S^1 \setminus J$ agrees with $\rho(c)$. Let $f_t : S^1 \to S^1$ be a continuous family of homeomorphisms supported on J that conjugate the action of $\rho_t([a, b])$ to the action of $\rho(c)$ there. (For the benefit of the reader, justification of this step via a simple construction of such a family is given in Lemma 3.3 below.) Then $\rho_t(c) = f_t \rho(c) f_t^{-1}$, as claimed.

Lemma 3.3. Let g_t be a continuous family (though not necessarily a subgroup) of homeomorphisms of an open interval I, with $\operatorname{Fix}(g_t) \cap I = \emptyset$ for all $t \in [0, 1]$. There exists a continuous family of homeomorphisms f_t such that $f_tg_t f_t^{-1} = g_0$ for all t.

Proof. Fix *x* in the interior of *I*, and let $D_t := [x, g_t(x)]$ be a fundamental domain for the action of g_t . Define the restriction of f_t to D_0 be the (unique) affine homeomorphism $D_0 \rightarrow D_t$, and extend f_t equivariantly to give a homeomorphism of *I*. \Box

Corollary 3.4. Let $\rho : \Gamma_g \to \text{Homeo}^+(S^1)$. Suppose that *a* and *b* are simple closed curves in Γ_g with $i(a, b) = \pm 1$ and $\operatorname{\widetilde{rot}}([\rho(a), \rho(b)]) = \pm 1$. Then there exists a

deformation ρ' of ρ such that $\rho'(a)$ is hyperbolic. If, additionally, ρ is assumed path-rigid and minimal, then $\rho(a)$ is hyperbolic.

Proof. Let *A* denote the subgroup generated by *a* and *b* and let c = [a, b], so $\Gamma_g = A *_{\langle c \rangle} B$. Let $\bar{\rho}$ denote the restriction of ρ to *A*. By Lemma 3.2, there exists a family of representations $\bar{\rho}_t : A \to \text{Homeo}^+(S^1)$ such that $\bar{\rho}_t(c) = f_t \bar{\rho}(c) f_t^{-1}$ for some continuous family $f_t \in \text{Homeo}^+(S^1)$, and such that $\bar{\rho}_1(a)$ is hyperbolic. As in the bending construction, define a deformation of ρ by

$$\rho_t(\gamma) = \begin{cases} \bar{\rho}_t(\gamma) & \text{for } \gamma \in A, \\ f_t \rho(\gamma) f_t^{-1} & \text{for } \gamma \in B. \end{cases}$$

By construction, ρ_t is a well-defined representation, and $\rho_1(a) = \overline{\rho}_1(a)$ is hyperbolic.

If ρ is assumed path-rigid, then this deformation ρ_1 is semiconjugate to ρ . If ρ is additionally known to be minimal, then there is a continuous map h satisfying $h \circ \rho_1 = \rho \circ h$. In particular, this implies that $\operatorname{Fix}(\rho(a)) = h\operatorname{Fix}(\rho_1(a))$, so $\rho(a)$ has at most two fixed points. In this case, if $\rho(a)$ does not have hyperbolic dynamics then it has a lift to $\operatorname{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$ satisfying $|x - \rho(a)(x)| \leq 1$ for all x. However, this easily implies that $|\operatorname{fot}([\rho(a), \rho(b)])| < 1$. (The reader may verify this as an exercise, or see the proof of Theorem 2.2 in [Matsumoto 1987] where this computation is carried out.) We conclude that $\rho(a)$ must be hyperbolic when ρ is path-rigid and minimal.

Having found one hyperbolic element, our next goal is to produce many others. An important tool here, and in what follows, is the following basic observation on dynamics of circle homeomorphisms.

Observation 3.5. Let $f \in \text{Homeo}^+(S^1)$ be hyperbolic, with attracting point f_+ and repelling point f_- , and let $g \in \text{Homeo}^+(S^1)$. For any neighborhoods U_- and U_+ of f_- and f_+ , respectively, and any neighborhoods V_- and V_+ of $g^{-1}(f_-)$ and $g(f_+)$, respectively, there exists $N \in \mathbb{N}$ such that

$$f^N g(S^1 \smallsetminus V_-) \subset U_+$$
 and $gf^N(S^1 \smallsetminus U_-) \subset V_+$.

The proof is a direct consequence of Definition 3.1. Note that, if f is hyperbolic, then f^{-1} is as well (with attracting point f_{-} and repelling point f_{+}), so an analogous statement holds with f^{-1} in place of f and the roles of f_{+} and f_{-} reversed.

We now state two useful consequences of this observation. The proofs are elementary and left to the reader.

Corollary 3.6. Let $f \in \text{Homeo}^+(S^1)$ be hyperbolic, and suppose g does not exchange the fixed points of f. Then for N sufficiently large, either $f^N g$ or $f^{-N} g$ has a fixed point.

Corollary 3.7. Let $f \in \text{Homeo}^+(S^1)$ be hyperbolic, and suppose $g^{-1}(f_-) \neq f_+$. Suppose also that $f^N g$ is known to be hyperbolic for large N. Then as $N \to \infty$, the attracting point of $f^N g$ approaches f_+ and the repelling point approaches $g^{-1}(f_-)$.

With these tools in hand, we can use one hyperbolic element to find others.

Proposition 3.8. Let ρ be path-rigid and minimal, and suppose that $i(a, b) = \pm 1$ and that $\rho(a)$ is hyperbolic. Then $\rho(b)$ is hyperbolic.

Proof. We prove this under the assumption that $\rho(b)$ does not exchange the fixed points of $\rho(a)$. This assumption is justified by Lemma 3.9 below. Assuming $\rho(b)$ does not exchange the points of Fix($\rho(a)$), by Corollary 3.6, there exists some $N \in \mathbb{Z}$ such that $\rho(a^N b)$ has a fixed point. Since $\rho(a)$ is hyperbolic, $\rho(a)^N$ belongs to a one-parameter family of homeomorphisms, and a bending deformation using this family gives a deformation ρ_1 of ρ with $\rho_1(b) = \rho(a^N b)$. By Corollary 2.10, using the fact that ρ is minimal, ρ_1 and ρ are conjugate. Thus, $\rho(b)$ has a fixed point and belongs to a one-parameter group b_t .

Now we can build a bending deformation ρ'_t such that $\rho'_1(b) = \rho(b)$ and $\rho'_1(a) = \rho(ba)$. Thus, $\rho'_1(a^{-1}b) = \rho(a^{-1})$, which is hyperbolic. Since ρ'_1 and ρ are conjugate, this means that $\rho(a^{-1}b)$ is hyperbolic. Similarly, using the fact that *a* belongs to a one-parameter group, there exists a bending deformation ρ''_t with $\rho''_1(a^{-1}b) = \rho(b)$, and such that ρ''_1 is conjugate to ρ . This implies that $\rho(b)$ is hyperbolic.

Lemma 3.9. Let $a, b \in \Gamma_g$ satisfy $i(a, b) = \pm 1$, and let $\rho : \Gamma_g \to \text{Homeo}^+(S^1)$. Suppose that $\rho(a)$ is hyperbolic, and $\rho(b)$ exchanges the fixed points of $\rho(a)$. Then there is a deformation ρ' of ρ which is not semiconjugate to ρ .

Proof. Note first that the property that $\rho(b)$ exchanges the fixed points of $\rho(a)$ implies that $\rho(b^{-1}a^{-1}b)$ is hyperbolic with the same attracting and repelling points as *a*. Hence $[\rho(a), \rho(b)]$ is hyperbolic with the same attracting and repelling points as well. We now produce a deformation ρ_1 of ρ such that $\rho_1(a)$ and $\rho_1(b)$ are in PSL(2, \mathbb{R}), after this we will easily be able to make an explicit further deformation to a representation which is not semiconjugate.

First, conjugate ρ so that $\rho(a) \in PSL(2, \mathbb{R})$ and so that the attracting and repelling fixed points of $\rho(a)$ are at 0 and 1/2 respectively (thinking of S^1 as \mathbb{R}/\mathbb{Z}). Now choose a continuous path b_t from $b_0 = b$ to the order two rotation $b_1 : x \mapsto x + 1/2$, and such that $b_t(0) = 1/2$ and $b_t(1/2) = 0$ for all t. By the observation above, $[\rho(a), b_t]$ is hyperbolic with attracting fixed point 0 and repelling fixed point 1/2 for all t, and so is conjugate to $\rho(a)$. By Lemma 3.3, applied separately to (0, 1/2) and (1/2, 1), there exists a continuous choice of conjugacies f_t such that $f_t[\rho(a), \rho(b)]f_t^{-1} = [\rho(a), b_t]$. Now to define ρ_t , we consider $\Gamma_g = A *_c B$ where $A = \langle a, b \rangle$ and c = [a, b], and set

$$\rho_t(\gamma) = f_t \rho(\gamma) f_t^{-1} \text{ for } \gamma \in B, \qquad \rho_t(a) = \rho(a), \qquad \rho_t(b) = b_t.$$

This gives a continuous family of well-defined representations, with $\rho_1(b)$ the standard order 2 rotation, and $\rho_1(a) \in PSL(2, \mathbb{R})$.

To finish the proof of the lemma, it suffices to note that, for a sufficiently small deformation b'_t of $\rho_1(b)$ in SO(2), the commutator $[\rho_1(a), b'_t]$ will remain hyperbolic, as the set of hyperbolic elements is open in PSL(2, \mathbb{R}). Thus, there is a continuous path of conjugacies in Homeo⁺(S^1) to $[\rho_1(a), b]$. This allows us to build a deformation ρ' of ρ with $\rho'(b) = b'_t \in$ SO(2), using the strategy from Corollary 3.4. Since $\operatorname{rot}(b'_t) \neq \operatorname{rot}(b) = 1/2$, it follows that ρ' and ρ are not semiconjugate.

The following corollary summarizes the results of this section.

Corollary 3.10. Let \sim_i denote the equivalence relation on nonseparating simple closed curves in Σ_g generated by $a \sim_i b$ if $i(a, b) = \pm 1$. Suppose ρ : $\Gamma_g \rightarrow \text{Homeo}^+(S^1)$ is path-rigid, and suppose that there are simple closed curves a, b with $i(a, b) = \pm 1$ such that $\operatorname{rot}[\rho(a), \rho(b)] = \pm 1$. Then ρ is semiconjugate to a (minimal) representation with $\rho(\gamma)$ hyperbolic for all $\gamma \sim_i a$.

Remark 3.11. In fact, the relation \sim_i has only a single equivalence class! This statement of connectedness of a certain complex of *based* curves can be proved using the connectedness of the arc complex of the once-punctured surface Σ_g^1 ; see [Mann and Wolff 2017, Section 2.1] for details. However, we will not need to use this fact here, so to keep the proof as self-contained and short as possible we will not refer to it further.

4. Step 3: Configuration of fixed points

The objective of this section is to organize the fixed points of the hyperbolic elements in a directed 5-chain; we will achieve this gradually by considering first 2-chains, then 3-chains, and finally 5-chains.

As in Definition 3.1, for a hyperbolic element $f \in \text{Homeo}^+(S^1)$ we let f_+ denote the attracting fixed point of f, and f_- the repelling point. By "Fix(f) separates Fix(g)" we mean that g_- and g_+ lie in different connected components of $S^1 \setminus \text{Fix}(f)$. In particular, Fix(f) and Fix(g) are disjoint.

Lemma 4.1. Let ρ be path-rigid and minimal, and let a, b be simple closed curves with $i(a, b) = \pm 1$ and $\rho(a)$ hyperbolic. Then $\rho(b)$ is hyperbolic, and Fix($\rho(a)$) separates Fix($\rho(b)$) in S^1 .

Proof. That $\rho(b)$ is hyperbolic follows from Proposition 3.8 above.

We prove the separation statement. As a first step, let us show that $Fix(\rho(a))$ and $Fix(\rho(b))$ are disjoint. Suppose for contradiction that they are not. Then, (after reversing orientations if needed) we have $\rho(a)_+ = \rho(b)_+$. Let *I* be a neighborhood of $\rho(a)_+$ with closure disjoint from { $\rho(a)_-$, $\rho(b)_-$ }. Then, for N > 0 large enough,

we have $\overline{I} \subset \rho(a^{-N}b)(I)$. Let ρ_t be a bending deformation with $\rho_0 = \rho$, $\rho_t(a) = \rho(a)$ and $\rho_1(b) = \rho(a^{-N}b)$. By Corollary 2.10, $\rho_1(b)$ is hyperbolic. Since $\overline{I} \subset \rho(a^{-N}b)(I)$, its attracting fixed point is outside *I*, and hence $\rho_1(b)_+ \neq \rho_1(a)_+$. But ρ and ρ_1 are conjugate by Corollary 2.10; this is a contradiction.

Now that we know that $\operatorname{Fix}(\rho(a)) \cap \operatorname{Fix}(\rho(b)) = \emptyset$, we will prove that they separate each other. Suppose for contradiction that $\operatorname{Fix}(\rho(a))$ does not separate $\operatorname{Fix}(\rho(b))$. Up to conjugating ρ by an orientation-reversing homeomorphism of S^1 , and up to replacing *b* with b^{-1} , the fixed points of $\rho(a)$ and $\rho(b)$ have cyclic order (a_+, a_-, b_+, b_-) . (For simplicity, we have suppressed the notation ρ .)

Fix $N \in \mathbb{N}$ large, and let ρ' be a bending deformation of ρ so that $\rho'(b) = \rho(a^N)\rho(b)$, and $\rho'(a) = \rho(a)$. It follows from Corollaries 2.10 and 3.7 that, if N is large enough, the points $b'_+ = \rho'(b)_+$ and $b'_- = \rho'(b)_-$ can be taken arbitrarily close, respectively, to a_+ and $\rho(b)^{-1}(a_-)$. Since the cyclic order of fixed points is preserved under deformation, they are also in order (a_+, a_-, b'_+, b'_-) . This is incompatible with the positions of a_+ and $\rho(b)^{-1}(a_-)$, unless perhaps if $\rho(b)^{-1}(a_-) = a_+$. But if $\rho(b)^{-1}(a_-) = a_+$, then $\rho'(b)$ has no fixed point in $(\rho(b)^{-1}(a_+), a_+)$ as this interval is mapped into (a_+, a_-) by $\rho(b')$. This again gives an incompatibility with the cyclic order.

Lemma 4.2. Let ρ be path-rigid and minimal, and let (a, b, c) be a directed 3chain. Suppose that $\rho(a)$ is hyperbolic, and suppose that $\rho(a)$ and $\rho(c)$ do not have a common fixed point. Then $\rho(b)$ and $\rho(c)$ are hyperbolic, and, up to reversing the orientation of S^1 , their fixed points are in the cyclic order

$$(\rho(a)_{-}, \rho(b)_{-}, \rho(a)_{+}, \rho(c)_{-}, \rho(b)_{+}, \rho(c)_{+}).$$

Proof. It follows from Proposition 3.8 that $\rho(b)$ and $\rho(c)$ are hyperbolic, and from Lemma 4.1 that up to reversing orientation, the fixed points of $\rho(a)$ and $\rho(b)$ come in the cyclic order

$$(a_{-}, b_{-}, a_{+}, b_{+}).$$

(For simplicity we drop ρ from the notation for the fixed points.) As mentioned above, the effect of a bending deformation that realizes a power of a Dehn twist along *a* is to leave *a* and *c* invariant and to replace *b* with ba^N . Corollary 2.10 says that the resulting representation is conjugate to ρ . By doing this with N > 0and N < 0 large, we get representations for which $b'_{-} = \rho(ba^N)_{-}$ can be taken arbitrarily close to a_+ , as well as to a_- . This, and Lemma 4.1 applied to the curves (b, c), imply that the intervals (a_+, b_+) and (b_+, a_-) each contain one fixed point of *c*. To prove the lemma, it now suffices to prove the cyclic order of fixed points

$$(a_{-}, b_{-}, a_{+}, c_{+}, b_{+}, c_{-})$$

cannot occur. Suppose for contradiction that this configuration holds, and apply a power of Dehn twist along *b*, replacing *a* with $b^{-N}a$ and *c* with cb^{N} (and leaving

b invariant), for N > 0 large. Denote by c'_+ , c'_- , a'_- and a'_+ the resulting fixed points, i.e., the fixed points of $\rho(cb^N)$ and $\rho(b^{-N}a)$ for N > 0 large. If N is chosen large enough, then c'_+ , c'_- and a'_- are arbitrarily close to $c(b_+)$, b_- and $a^{-1}(b_+)$, respectively. (See Corollary 3.7.) These three points are in the reverse cyclic order as c_+ , c_- and a_- ; hence, the representation ρ' obtained from this Dehn twist cannot be conjugate to ρ . This contradicts Corollary 2.10, and so eliminates the undesirable configuration.

We are now ready to prove the main result of this section.

Proposition 4.3. Let ρ be a path-rigid, minimal representation, and (a, b, c, d, e) be a directed 5-chain in Σ_g . Suppose $\rho(a)$ is hyperbolic. Then, $\rho(b), \ldots, \rho(e)$ are hyperbolic as well, and up to reversing the orientation of the circle, their fixed points are in the following (total) cyclic order:

$$(a_{-}, b_{-}, a_{+}, c_{-}, b_{+}, d_{-}, c_{+}, e_{-}, d_{+}, e_{+}).$$

In particular, these fixed points are all distinct. As before, for simplicity we have dropped ρ from the notation.

Proof. That $\rho(b), \ldots, \rho(e)$ are all hyperbolic follows from Proposition 3.8. Next, using a bending deformation realizing a Dehn twist along d, we may change the action of c into $d^{-N}c$ without changing a, and without changing the conjugacy class of ρ . In particular, such a deformation moves the fixed points of c, so we can ensure that $Fix(\rho(a))$ and $Fix(\rho(c))$ are disjoint.

Similarly, for any two elements in the chain (a, b, c, d, e), there is a third one that intersects one but not the other. Thus, we may apply the same reasoning to show that all these five hyperbolic elements have pairwise disjoint fixed sets. It remains to order these fixed sets. For this, we will apply Lemma 4.2 repeatedly.

First, fix the orientation of S^1 so that, applying Lemma 4.2 to the directed 3-chain (a, b, c), we have the cyclic order of fixed points

$$(a_{-}, b_{-}, a_{+}, c_{-}, b_{+}, c_{+}).$$

Now, Lemma 4.2 applied to the directed 3-chain (b, c, d) implies that d_{-} lies in the interval (b_{+}, c_{+}) and d_{+} in the interval (c_{+}, b_{-}) . Applying the lemma to the directed 3-chain (a, cb, d) implies that d_{+} in fact lies in the interval (c_{+}, a_{-}) .

The same argument using Lemma 4.2 applied to the directed 3-chains (c, d, e) and (a, dcb, e) shows that e_{-} lies in the interval (c_{+}, d_{+}) and e_{+} in the interval (d_{+}, a_{-}) , as desired.

5. Step 4: Maximality of the Euler number

In order to compute the Euler number of ρ , we will decompose Σ_g into subsurfaces and compute the contribution to $eu(\rho)$ from each part. The proper framework for



Figure 2. A pair of pants with standard generators of its fundamental group.

discussing this is the language of bounded cohomology: if Σ is a surface with boundary $\partial \Sigma$, and $\rho : \pi_1(\Sigma) \to \text{Homeo}^+(S^1)$, one obtains a characteristic number by pulling back the *bounded Euler class* in $H_b^2(\text{Homeo}^+(S^1); \mathbb{R})$ to $H_b^2(\Sigma, \partial \Sigma; \mathbb{R})$ and pairing it with the fundamental class $[\Sigma, \partial \Sigma]$. The contribution to the Euler number of $\rho : \Sigma_g \to \text{Homeo}^+(S^1)$ from a subsurface Σ is simply this Euler number for the restriction of ρ to Σ .

However, in order to keep this work self-contained and elementary, we will avoid introducing the language of bounded cohomology, and give definitions in terms of rotation numbers alone. The reader may refer to [Burger et al. 2014, §4.3] for details on the cohomological framework.

Definition 5.1 (Euler number for pants). Let $\rho : \Gamma_g \to \text{Homeo}^+(S^1)$, and let $P \subset \Sigma_g$ be a subsurface homeomorphic to a pair of pants, bounded by curves a, d and $(da)^{-1}$, with orientation induced from the boundary. Let $\rho(a)$ and $\rho(d)$ be any lifts of $\rho(a)$ and $\rho(d)$ to $\text{Homeo}_{\mathbb{Z}}^+(\mathbb{R})$. The *Euler number of* ρ *on* P is the real number

$$\operatorname{eu}_P(\rho) = \widetilde{\operatorname{rot}}(\rho(\widetilde{a})) + \widetilde{\operatorname{rot}}(\rho(\widetilde{d})) - \widetilde{\operatorname{rot}}(\rho(\widetilde{d})\rho(\widetilde{a})).$$

An illustration in the case where P contains the basepoint is given in Figure 2.

Note that the number $eu_P(\rho)$ is independent of the choice of lifts of $\rho(a)$ and $\rho(d)$. We also allow for the possibility that the image of P in Σ_g has two boundary curves identified, and so is a one-holed torus subsurface. In this case, one may choose free generators a, b for the fundamental group, with i(a, b) = -1 so the torus is T(a, b) and the boundary of P is given by the curves b^{-1} , $a^{-1}ba$ and the commutator [a, b]. Then the definition above gives

$$\operatorname{eu}_{P}(\rho) = \operatorname{\widetilde{rot}}[\rho(a), \rho(b)].$$

Now, the following is a restatement of Lemma 2.1 above.

Lemma 5.2. Let *P* be any pants and ρ a representation. Then $|eu_P(\rho)| \le 1$.

More generally, if $S \subset \Sigma_g$ is any subsurface, we define the Euler number $eu_S(\rho)$ to be the sum of relative Euler numbers over all pants in a pants decomposition of *S*. From the perspective of bounded cohomology, it is immediate that this sum does not depend on the pants decomposition; however, since we are intentionally avoiding cohomological language, we give a short stand-alone proof.

Lemma 5.3. For any subsurface $S \subseteq \Sigma_g$, the number $eu_S(\rho)$ is well defined, i.e., independent of the decomposition of S into pants.

Proof. Any two pants decompositions can be joined by a sequence of elementary moves; namely those of types (I) and (IV) as shown in [Hatcher and Thurston 1980]. A type (IV) move takes place within a pants-decomposed one-holed torus P and so does not change the value of eu_P , which is simply the rotation number of the boundary curve, as remarked above. A type (I) move occurs within a four-holed sphere S'; if the boundary of the sphere is given by oriented curves a, b, c, d with dcba = 1, then it consists of replacing the decomposition along da with a decomposition along ab. It is easy to verify by the definition that, in either case, the sum of the Euler numbers of the two pants on S' is given by

$$\widetilde{\mathrm{rot}}(\widetilde{\rho(a)}) + \widetilde{\mathrm{rot}}(\widetilde{\rho(b)}) + \widetilde{\mathrm{rot}}(\widetilde{\rho(c)}) + \widetilde{\mathrm{rot}}(\widetilde{\rho(d)}).$$

Corollary 5.4 (additivity of Euler number). Let \mathcal{P} be any decomposition of Σ into pants. Then

$$\operatorname{eu}(\rho) = \sum_{P \in \mathcal{P}} \operatorname{eu}_P(\rho).$$

Proof. By Lemma 5.3, we may use any pants decomposition to compute the Euler class. By using a standard generating system (a_1, \ldots, b_g) and cutting Σ_g along geodesics freely homotopic to a_i , $c_i = [a_i, b_i]$, for $i = 1, \ldots, g$ and $d_i = c_i \cdots c_1$ for $i = 2, \ldots, g - 1$, we recover the formula taken as a definition in Proposition 2.2. \Box

We now return to our main goal: we prove that maximality of the Euler class holds first on small subsurfaces, then globally on Σ_g .

Proposition 5.5. Let $S \subset \Sigma_g$ be a subsurface homeomorphic to a four-holed sphere. Suppose that none of the boundary components of S is separating in Σ_g , and let ρ be a path-rigid, minimal representation mapping one boundary component of S to a hyperbolic element of Homeo⁺(S^1). Then, ρ maps all four boundary components of S to hyperbolic elements, and the relative Euler class $eu_S(\rho)$ is equal to ± 2 .

In the statement above, we do not require that the boundary components are geodesics for some metric on Σ_g , in particular, two of them may well be freely homotopic.

Proof. Put the base point inside of S. The complement $\Sigma_g \setminus S$ may have one or two connected components, since none of the curves of ∂S are separating in Σ_g . In



Figure 3. A four-holed sphere and two 5-chains.

either case, we may find two based, nonseparating, simple closed curves $u, v \in \Gamma_g$, with i(u, v) = 0, each having nonzero intersection number with exactly two of the boundary components of *S*, as shown in Figure 3. Additionally, we may fix orientations for *u* and *v* and choose four elements *a*, *b*, *c*, *d* $\in \pi_1 S$, each freely homotopic to a different boundary component of *S*, with dcba = 1, and such that $(a, u, d^{-1}a^{-1}, v, d)$ and (c, v, ad, u, b) are directed 5-chains in Σ_g . As we have assumed that the image under ρ of one of *a*, *b*, *c* or *d* is hyperbolic, Proposition 3.8 implies that all the curves appearing in these 5-chains are in fact hyperbolic.

Orient the circle so that $(u_-, (ad)_+, u_+, (ad)_-)$ are in cyclic order (as before, we drop the letter ρ from the notation, for better readability). Then, Proposition 4.3 applied to the two directed 5 chains above gives the cyclic orderings

and

$$(a_{-}, u_{-}, a_{+}, (ad)_{+}, u_{+}, v_{-}, (ad)_{-}, d_{-}, v_{+}, d_{+})$$

 $(c_{-}, v_{-}, c_{+}, (ad)_{-}, v_{+}, u_{-}, (ad)_{+}, b_{-}, u_{+}, b_{+}).$

These two orderings together yield the cyclic ordering

 $((ad)_{-}, d_{-}, d_{+}, a_{-}, a_{+}, (ad)_{+}, b_{-}, b_{+}, c_{-}, c_{+}).$

We now use this ordering to prove maximality of the Euler class. Let α , β , γ and δ , respectively, denote the lifts of $\rho(a)$, $\rho(b)$, $\rho(c)$ and $\rho(d)$ to Homeo⁺_Z(\mathbb{R}) with translation number zero. Let $x = (ad)_{-}$ be the repelling fixed point of *ad*.

Since *x* has a repelling fixed point of *d* immediately to the right, and an attracting fixed point of *d* to the left, we have $\delta(x) < x$. By the same reasoning, if *y* is any point in the interval between consecutive lifts of fixed points a_+ and a_- containing *x*, then $\alpha(y) < y$. Since ad(x) = x, it follows that $\delta(x)$ must lie to the left of the lift of a_+ , and we have $\alpha\delta(x) = x - 1$.

Since cbad = 1, we also have that cb(x) = x. Considering the location of repelling points of *b* and *c* and imitating the argument above, we have again $\beta(x) < x$, and also $\gamma\beta(x) < x$. It follows that $\gamma\beta(x) = x - 1$, hence $\gamma\beta\alpha\delta(x) = x - 2$, and $eu_S(\rho) = -2$.

With this information on subsurfaces, we prove the Euler number of ρ is maximal. **Proposition 5.6.** Let ρ be path-rigid, and suppose that ρ admits a Fuchsian torus. Then ρ has Euler number $\pm (2g - 2)$.

Proof. After semiconjugacy, we may assume that ρ is minimal. Let T(a, b) be a Fuchsian torus for ρ . By Corollary 3.4, we may suppose that $\rho(a)$ is hyperbolic. Ignoring the curve *b*, find a system of simple closed curves $a_1 = a, a_2, \ldots, a_{g-1}$, with each a_i nonseparating, that decomposes Σ_g into a disjoint union of pairs of pants.

The dual graph of such a pants decomposition is connected (because Σ_g is connected), so we may choose a finite path that visits all the vertices. In other words, we may choose a sequence P_1, \ldots, P_N of pants from the decomposition (possibly with repetitions), that contains each of the pants of the decomposition, such that each two consecutive pants P_i and P_{i+1} are distinct, but share a boundary component. Let S_i denote the four-holed sphere obtained by taking the union of P_i and P_{i+1} along a shared boundary curve. (If P_i and P_{i+1} share more than one boundary component, choose only one). We may further assume that *a* is one of the boundary curves of S_1 .

Starting with S_1 as the base case, and applying Proposition 5.5, we inductively conclude that all boundary components of all the S_i are hyperbolic, and that $eu_{S_i}(\rho) = \pm 2$. Thus, the contributions of P_i and P_{i+1} are equal, and equal to ± 1 , for all *i*. It follows that the contributions of all pairs of pants of the decomposition have equal contributions, equal to ± 1 . By definition of the Euler class, we conclude that $eu(\rho) = \pm (2g - 2)$.

The proof of Theorem 1.3 now concludes by citing Matsumoto's result [1987] that such a representation of maximal Euler number is semiconjugate to a Fuchsian representation.

Acknowledgements

This work was started at MSRI during spring 2015 at a program supported by NSF grant 0932078. Both authors also acknowledge support of the U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 "RNMS: Geometric Structures and Representation Varieties" (the GEAR network). Mann was partially supported by NSF grant DMS-1606254, and thanks the Inst. Math. Jussieu and Foundation Sciences Mathématiques de Paris. This work was finished while Wolff was visiting the Universidad de la República, Montevideo, Uruguay.

References

[[]Bucher et al. 2016] M. Bucher, R. Frigerio, and T. Hartnick, "A note on semi-conjugacy for circle actions", *Enseign. Math.* **62**:3-4 (2016), 317–360. MR Zbl

[[]Burger et al. 2014] M. Burger, A. Iozzi, and A. Wienhard, "Higher Teichmüller spaces: from $SL(2, \mathbb{R})$ to other Lie groups", pp. 539–618 in *Handbook of Teichmüller theory, IV*, edited by A. Papadopoulos, IRMA Lect. Math. Theor. Phys. **19**, Eur. Math. Soc., Zürich, 2014. MR Zbl

- [Calegari 2006] D. Calegari, "Dynamical forcing of circular groups", *Trans. Amer. Math. Soc.* **358**:8 (2006), 3473–3491. MR Zbl
- [Calegari and Walker 2011] D. Calegari and A. Walker, "Ziggurats and rotation numbers", *J. Mod. Dyn.* **5**:4 (2011), 711–746. MR Zbl
- [Farb and Margalit 2012] B. Farb and D. Margalit, *A primer on mapping class groups*, Princeton Math. Series **49**, Princeton Univ. Press, 2012. MR Zbl
- [Ghys 1987] É. Ghys, "Groupes d'homéomorphismes du cercle et cohomologie bornée", pp. 81–106 in *The Lefschetz Centennial Conference, III* (Mexico City, 1984), edited by A. Verjovsky, Contemp. Math. **58**, Amer. Math. Soc., Providence, RI, 1987. MR Zbl
- [Ghys 2001] É. Ghys, "Groups acting on the circle", *Enseign. Math.* (2) **47**:3-4 (2001), 329–407. MR Zbl
- [Hatcher and Thurston 1980] A. Hatcher and W. Thurston, "A presentation for the mapping class group of a closed orientable surface", *Topology* **19**:3 (1980), 221–237. MR Zbl
- [Mann 2015] K. Mann, "Spaces of surface group representations", *Invent. Math.* 201:2 (2015), 669–710. MR Zbl
- [Mann 2018] K. Mann, "Rigidity and flexibility of group actions on the circle", pp. 705–752 in *Handbook of group actions, IV*, edited by L. Ji et al., Adv. Lectures in Math. **41**, Int. Press, Somerville, MA, 2018. arXiv
- [Mann and Wolff 2017] K. Mann and M. Wolff, "Rigidity and geometricity for surface group actions on the circle", preprint, 2017. arXiv
- [Matsumoto 1987] S. Matsumoto, "Some remarks on foliated S^1 bundles", *Invent. Math.* **90**:2 (1987), 343–358. MR Zbl
- [Matsumoto 2016] S. Matsumoto, "Basic partitions and combinations of group actions on the circle: a new approach to a theorem of Kathryn Mann", *Enseign. Math.* **62**:1-2 (2016), 15–47. MR Zbl
- [Milnor 1958] J. Milnor, "On the existence of a connection with curvature zero", *Comment. Math. Helv.* **32** (1958), 215–223. MR Zbl
- [Poincaré 1885] H. Poincaré, "Sur les courbes définies par les équations différentielles (III)", *Journal de mathématiques pures et appliquées* **4**:1 (1885), 167–244.
- [Wood 1971] J. W. Wood, "Bundles with totally disconnected structure group", *Comment. Math. Helv.* **46** (1971), 257–273. MR Zbl

Received March 9, 2018. Revised January 16, 2019.

KATHRYN MANN Department of Mathematics Cornell University Ithaca, NY United States

k.mann@cornell.edu

MAXIME WOLFF Sorbonne Université Université Paris Diderot CNRS, Institut de Mathématiques de Jussieu-Paris Rive Gauche, IMJ-PRG Paris France maxime.wolff@imj-prg.fr

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906-1982) and F. Wolf (1904-1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor) Department of Mathematics University of California Los Angeles, CA 90095-1555 blasius@math.ucla.edu

Paul Balmer Department of Mathematics University of California Los Angeles, CA 90095-1555 balmer@math.ucla.edu

Wee Teck Gan Mathematics Department National University of Singapore Singapore 119076 matgwt@nus.edu.sg

Sorin Popa Department of Mathematics University of California Los Angeles, CA 90095-1555 popa@math.ucla.edu

Paul Yang Department of Mathematics Princeton University Princeton NJ 08544-1000 yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI CALIFORNIA INST. OF TECHNOLOGY INST. DE MATEMÁTICA PURA E APLICADA KEIO UNIVERSITY MATH. SCIENCES RESEARCH INSTITUTE NEW MEXICO STATE UNIV. OREGON STATE UNIV.

Matthias Aschenbrenner

Department of Mathematics

University of California

Los Angeles, CA 90095-1555

matthias@math.ucla.edu

Daryl Cooper

Department of Mathematics

University of California

Santa Barbara, CA 93106-3080

cooper@math.ucsb.edu

Jiang-Hua Lu

Department of Mathematics

The University of Hong Kong

Pokfulam Rd., Hong Kong

jhlu@maths.hku.hk

STANFORD UNIVERSITY UNIV. OF BRITISH COLUMBIA UNIV. OF CALIFORNIA, BERKELEY UNIV. OF CALIFORNIA, DAVIS UNIV. OF CALIFORNIA, LOS ANGELES UNIV. OF CALIFORNIA, RIVERSIDE UNIV. OF CALIFORNIA, SAN DIEGO UNIV. OF CALIF., SANTA BARBARA Vyjayanthi Chari Department of Mathematics University of California Riverside, CA 92521-0135 chari@math.ucr.edu

Kefeng Liu Department of Mathematics University of California Los Angeles, CA 90095-1555 liu@math.ucla.edu

Jie Qing Department of Mathematics University of California Santa Cruz, CA 95064 qing@cats.ucsc.edu

UNIV. OF CALIF., SANTA CRUZ UNIV. OF MONTANA UNIV. OF OREGON UNIV. OF SOUTHERN CALIFORNIA UNIV. OF UTAH UNIV. OF WASHINGTON WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2019 is US \$490/year for the electronic version, and \$665/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

mathematical sciences publishers

nonprofit scientific publishing

http://msp.org/ © 2019 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 302 No. 1 September 2019

On masas in <i>q</i> -deformed von Neumann algebras	1
MARTIJN CASPERS, ADAM SKALSKI allu MATEUSZ WASILEWSKI	
The compact picture of symmetry-breaking operators for rank-one orthogonal and unitery groups	23
IAN EDALM and BENT ØDSTED	
On the Londoham sumption of a close of Einstein matrice construct from the	77
navigation problem	//
LIBING HUANG, HUAIFU LIU and XIAOHUAN MO	
Symplectic and odd orthogonal Pfaffian formulas for algebraic cobordism	97
THOMAS HUDSON and TOMOO MATSUMURA	
A compactness theorem on Branson's Q -curvature equation	119
GANG LI	
A characterization of Fuchsian actions by topological rigidity	181
KATHRYN MANN and MAXIME WOLFF	
Fundamental domains and presentations for the Deligne–Mostow lattices with 2-fold symmetry	201
IRENE PASQUINELLI	
Binary quartic forms with bounded invariants and small Galois groups	249
CINDY (SIN YI) TSANG and STANLEY YAO XIAO	
Obstructions to lifting abelian subalgebras of corona algebras	293
Andrea Vaccaro	
Schwarz lemma at the boundary on the classical domain of type \mathcal{FV}	309
JIANFEI WANG, TAISHUN LIU and XIAOMIN TANG	
Cyclic η -parallel shape and Ricci operators on real hypersurfaces in two-dimensional nonflat complex space forms	335
YANING WANG	
Finsler spheres with constant flag curvature and finite orbits of prime closed geodesics	353
Ming Xu	
Degeneracy theorems for two holomorphic curves in $\mathbb{P}^n(\mathbb{C})$ sharing few	371
hypersurfaces	
KAI ZHOU and LU JIN	