

Pacific Journal of Mathematics

**TRANSITIVE TOPOLOGICAL MARKOV CHAINS
OF GIVEN ENTROPY AND PERIOD
WITH OR WITHOUT MEASURE OF MAXIMAL ENTROPY**

SYLVIE RUETTE

TRANSITIVE TOPOLOGICAL MARKOV CHAINS OF GIVEN ENTROPY AND PERIOD WITH OR WITHOUT MEASURE OF MAXIMAL ENTROPY

SYLVIE RUETTE

We show that, for every positive real number h and every positive integer p , there exist oriented graphs G, G' (with countably many vertices) that are strongly connected, of period p , of Gurevich entropy h , and such that G is positive recurrent (thus the topological Markov chain on G admits a measure of maximal entropy) and G' is transient (thus the topological Markov chain on G' admits no measure of maximal entropy).

1. Vere-Jones classification of graphs

In this paper, all the graphs are oriented and have a finite or countable set of vertices, and if u, v are two vertices, there is at most one arrow $u \rightarrow v$. A *path* of length n in the graph G is a sequence of vertices (u_0, u_1, \dots, u_n) such that $u_i \rightarrow u_{i+1}$ in G for all $i \in \llbracket 0, n-1 \rrbracket$. This path is called a *loop* if $u_0 = u_n$.

Definition 1. Let G be an oriented graph, and let u, v be two vertices in G . We define the following quantities:

- $p_{uv}^G(n)$ is the number of paths (u_0, u_1, \dots, u_n) such that $u_0 = u$ and $u_n = v$; $R_{uv}(G)$ is the radius of convergence of the series $\sum p_{uv}^G(n)z^n$.
- $f_{uv}^G(n)$ is the number of paths (u_0, u_1, \dots, u_n) such that $u_0 = u$, $u_n = v$, and $u_i \neq v$ for all $0 < i < n$; $L_{uv}(G)$ is the radius of convergence of the series $\sum f_{uv}^G(n)z^n$.

Definition 2. Let G be an oriented graph and V its set of vertices. The graph G is *strongly connected* if, for all $u, v \in V$, there exists a path from u to v in G . The *period* of a strongly connected graph G is the greatest common divisor of $(p_{uu}^G(n))_{u \in V, n \geq 0}$. The graph G is *aperiodic* if its period is 1.

Proposition 3 [Vere-Jones 1962]. *Let G be an oriented graph. If G is strongly connected, $R_{uv}(G)$ does not depend on u and v ; it is denoted by $R(G)$.*

MSC2010: primary 37B10; secondary 37B40.

Keywords: topological Markov chain, countable oriented graph, topological entropy.

	transient	null recurrent	positive recurrent
$\sum_{n>0} f_{uu}^G(n) R^n$	< 1	1	1
$\sum_{n>0} n f_{uu}^G(n) R^n$	$\leq +\infty$	$+\infty$	$< +\infty$
$\sum_{n\geq 0} p_{uv}^G(n) R^n$	$< +\infty$	$+\infty$	$+\infty$
$\lim_{n \rightarrow +\infty} p_{uv}^G(n) R^n$	0	0	$\lambda_{uv} > 0$
	$R = L_{uu}$	$R = L_{uu}$	$R \leq L_{uu}$

Table 1. Properties of the series associated to a transient, null recurrent or positive recurrent graph G (G is strongly connected); these properties do not depend on the vertices u, v .

If there is no confusion, $R(G)$ and $L_{uv}(G)$ will be written R and L_{uv} .

Vere-Jones [1962] gives a classification of strongly connected graphs as transient, null recurrent, or positive recurrent. These definitions are lines 1 and 2 in Table 1. The other lines of Table 1 state properties of the series $\sum p_{uv}^G(n)z^n$, which give alternative definitions (lines 3 and 4 are in [Vere-Jones 1962], and the last line is Proposition 4).

Proposition 4 [Salama 1992]. *Let G be a strongly connected oriented graph. If G is transient or null recurrent, then $R = L_{uu}$ for all vertices u . Equivalently, if there exists a vertex u such that $R < L_{uu}$, then G is positive recurrent.*

2. Topological Markov chains and Gurevich entropy

Let G be an oriented graph and V its set of vertices. We define Γ_G as the set of two-sided infinite paths in G , that is,

$$\Gamma_G := \{(v_n)_{n \in \mathbb{Z}} \mid \text{for all } n \in \mathbb{Z}, v_n \rightarrow v_{n+1} \text{ in } G\} \subset V^{\mathbb{Z}}.$$

The map σ is the shift on Γ_G . The *topological Markov chain* on the graph G is the dynamical system (Γ_G, σ) .

The set V is endowed with the discrete topology, and Γ_G is endowed with the induced topology of $V^{\mathbb{Z}}$. The space Γ_G is not compact unless G is finite.

The topological Markov chain (Γ_G, σ) is transitive if and only if the graph G is strongly connected. It is topologically mixing if and only if the graph G is strongly connected and aperiodic.

If G is a finite graph, Γ_G is compact and the topological entropy $h_{\text{top}}(\Gamma_G, \sigma)$ is well defined (see, e.g., [Denker et al. 1976] for the definition of the topological entropy). If G is a countable graph, the *Gurevich entropy* [1969] of the graph G (or of the topological Markov chain Γ_G) is given by

$$h(G) := \sup\{h_{\text{top}}(\Gamma_H, \sigma) \mid H \subset G, H \text{ finite}\}.$$

This entropy can also be computed in a combinatorial way, as the exponential growth of the number of paths with fixed endpoints.

Proposition 5 [Gurevich 1970]. *Let G be a strongly connected oriented graph. Then for all vertices u, v ,*

$$h(G) = \lim_{n \rightarrow +\infty} \frac{1}{n} \log p_{uv}^G(n) = -\log R(G).$$

Moreover, the variational principle is still valid for topological Markov chains.

Theorem 6 [Gurevich 1969]. *Let G be an oriented graph. Then*

$$h(G) = \sup\{h_\mu(\Gamma_G) \mid \mu \text{ } \sigma\text{-invariant probability measure}\}.$$

In this variational principle, the supremum is not necessarily reached. The next theorem gives a necessary and sufficient condition for the existence of a measure of maximal entropy (that is, a probability measure μ such that $h(G) = h_\mu(\Gamma_G)$) when the graph is strongly connected.

Theorem 7 [Gurevich 1970]. *Let G be a strongly connected oriented graph of finite positive entropy. Then the topological Markov chain on G admits a measure of maximal entropy if and only if the graph G is positive recurrent. Moreover, such a measure is unique if it exists.*

3. Construction of graphs of given entropy and given period that are either positive recurrent or transient

Lemma 8. *Let $\beta \in (1, +\infty)$. There exist a sequence of nonnegative integers $(a(n))_{n \geq 1}$ and positive constants c, M such that*

- $a(1) = 1$,
- $\sum_{n \geq 1} a(n)(1/\beta^n) = 1$,
- for all $n \geq 2$, $c \cdot \beta^{n^2-n} \leq a(n^2) \leq c \cdot \beta^{n^2-n} + M$,
- for all $n \geq 1$, $0 \leq a(n) \leq M$ if n is not a square.

These properties imply that the radius of convergence of $\sum_{n \geq 1} a(n)z^n$ is $L = 1/\beta$ and that $\sum_{n \geq 1} na(n)L^n < +\infty$.

Proof. First we look for a constant $c > 0$ such that

$$(1) \quad \frac{1}{\beta} + c \sum_{n \geq 2} \beta^{n^2-n} \frac{1}{\beta^{n^2}} = 1.$$

We have

$$\sum_{n \geq 2} \beta^{n^2-n} \frac{1}{\beta^{n^2}} = \sum_{n \geq 2} \beta^{-n} = \frac{1}{\beta(\beta-1)}.$$

Thus,

$$(1) \iff \frac{1}{\beta} + \frac{c}{\beta(\beta-1)} = 1 \iff c = (\beta-1)^2.$$

Since $\beta > 1$, the constant $c := (\beta-1)^2$ is positive. We define the sequence $(b(n))_{n \geq 1}$ by

- $b(1) := 1$,
- $b(n^2) := \lfloor c\beta^{n^2-n} \rfloor$ for all $n \geq 2$,
- $b(n) := 0$ for all $n \geq 2$ such that n is not a square.

Then

$$\sum_{n \geq 1} b(n) \frac{1}{\beta^n} \leq \frac{1}{\beta} + c \sum_{n \geq 2} \beta^{n^2-n} \frac{1}{\beta^{n^2}} = 1.$$

We set $\delta := 1 - \sum_{n \geq 1} b(n)(1/\beta^n) \in [0, 1)$ and $k := \lfloor \beta^2 \delta \rfloor$. Then $k \leq \beta^2 \delta < k+1 < k+\beta$, which implies that $0 \leq \delta - k/\beta^2 < 1/\beta$. We write the β -expansion of $\delta - k/\beta^2$ (see, e.g., [Dajani and Kraaikamp 2002, p. 51] for the definition): there exist integers $d(n) \in \{0, \dots, \lfloor \beta \rfloor\}$ such that $\delta - k/\beta^2 = \sum_{n \geq 1} d(n)(1/\beta^n)$. Moreover, $d(1) = 0$ because $\delta - k/\beta^2 < 1/\beta$. Thus, we can write

$$\delta = \sum_{n \geq 2} d'(n) \frac{1}{\beta^n}$$

where $d'(2) := d(2) + k$ and $d'(n) := d(n)$ for all $n \geq 3$.

We set $a(1) := b(1)$ and $a(n) := b(n) + d'(n)$ for all $n \geq 2$. Let $M := \beta + k$. We then have

- $a(1) = 1$,
- $\sum_{n \geq 1} a(n)(1/\beta^n) = 1$,
- for all $n \geq 2$, $c \cdot \beta^{n^2-n} \leq a(n^2) \leq c \cdot \beta^{n^2-n} + \beta \leq c \cdot \beta^{n^2-n} + M$,
- $0 \leq a(2) \leq \beta + k = M$,
- for all $n \geq 3$, $0 \leq a(n) \leq \beta \leq M$ if n is not a square.

The radius of convergence L of $\sum_{n \geq 1} a(n)z^n$ satisfies

$$-\log L = \limsup_{n \rightarrow +\infty} \frac{1}{n} \log a(n) = \lim_{n \rightarrow +\infty} \frac{1}{n^2} \log a(n^2) = \log \beta$$

because $a(n^2) \sim c\beta^{n^2-n}$.

Thus, $L = 1/\beta$. Moreover,

$$\sum_{n \geq 1} na(n) \frac{1}{\beta^n} \leq M \sum_{n \geq 1} n \frac{1}{\beta^n} + c \sum_{n \geq 1} n^2 \beta^{n^2-n} \frac{1}{\beta^{n^2}} = M \sum_{n \geq 1} \frac{n}{\beta^n} + c \sum_{n \geq 1} \frac{n^2}{\beta^n} < +\infty. \quad \square$$




Figure 1. The graphs $G(\beta)$ and $G'(\beta)$; the bold loop belongs to $G(\beta)$ and not to $G'(\beta)$, otherwise the two graphs coincide.

Lemma 9 [Ruelle 2003, Lemma 2.4]. *Let G be a strongly connected oriented graph and u a vertex.*

- (i) $R < L_{uu}$ if and only if $\sum_{n \geq 1} f_{uu}^G(n) L_{uu}^n > 1$.
- (ii) If G is recurrent, then R is the unique positive number x such that

$$\sum_{n \geq 1} f_{uu}^G(n) x^n = 1.$$

Proof. For (i) and (ii), use Table 1 and the fact that $F(x) = \sum_{n \geq 1} f_{uu}^G(n) x^n$ is increasing for $x \in [0, +\infty)$. \square

Proposition 10. *Let $\beta \in (1, +\infty)$. There exist aperiodic strongly connected graphs $G'(\beta) \subset G(\beta)$ such that $h(G(\beta)) = h(G'(\beta)) = \log \beta$, $G(\beta)$ is positive recurrent, and $G'(\beta)$ is transient.*

Remark. Salama [1988, Theorem 3.9] proved the part of this proposition concerning positive recurrent graphs.

Proof. This is a variant of the proof of [Ruelle 2003, Example 2.9].

Let u be a vertex, and let $(a(n))_{n \geq 1}$ be the sequence given by Lemma 8 for β . The graph $G(\beta)$ is composed of $a(n)$ loops of length n based at the vertex u for all $n \geq 1$ (see Figure 1). More precisely, define the set of vertices of $G(\beta)$ as

$$V := \{u\} \cup \bigcup_{n=1}^{+\infty} \{v_k^{n,i} \mid i \in \llbracket 1, a(n) \rrbracket, k \in \llbracket 1, n-1 \rrbracket\},$$

where the vertices $v_k^{n,i}$ above are distinct. Let $v_0^{n,i} = v_n^{n,i} = u$ for all $i \in \llbracket 1, a(n) \rrbracket$. There is an arrow $v_k^{n,i} \rightarrow v_{k+1}^{n,i}$ for all $k \in \llbracket 0, n-1 \rrbracket$, $i \in \llbracket 1, a(n) \rrbracket$, and $n \geq 2$; there is an arrow $u \rightarrow u$; and there is no other arrow in $G(\beta)$. The graph $G(\beta)$ is strongly connected, and $f_{uu}^{G(\beta)}(n) = a(n)$ for all $n \geq 1$.

By Lemma 8, the sequence $(a(n))_{n \geq 1}$ is defined such that $L = 1/\beta$ and

$$(2) \quad \sum_{n \geq 1} a(n) L^n = 1,$$

where $L = L_{uu}(G(\beta))$ is the radius of convergence of the series $\sum a(n)z^n$. If $G(\beta)$ is transient, then $R(G(\beta)) = L_{uu}(G(\beta))$ by Proposition 4. But (2) contradicts the definition of transient (see the first line of Table 1). Thus, $G(\beta)$ is recurrent, and $R(G(\beta)) = L$ by (2) and Lemma 9(ii). Moreover,

$$\sum_{n \geq 1} na(n)L^n < +\infty$$

by Lemma 8, and thus the graph $G(\beta)$ is positive recurrent (see Table 1). By Proposition 5, $h(G(\beta)) = -\log R(G(\beta)) = \log \beta$.

The graph $G'(\beta)$ is obtained from $G(\beta)$ by deleting a loop starting at u of length n_0 for some $n_0 \geq 2$ such that $a(n_0) \geq 1$ (such an integer n_0 exists because $L < +\infty$). Obviously one has $L_{uu}(G'(\beta)) = L$ and

$$\sum_{n \geq 1} f_{uu}^{G'(\beta)}(n)L^n = 1 - L^{n_0} < 1.$$

Since $R(G'(\beta)) \leq L_{uu}(G'(\beta))$, this implies that $G'(\beta)$ is transient. Moreover, $R(G'(\beta)) = L_{uu}(G'(\beta))$ by Proposition 4, so $R(G'(\beta)) = R(G(\beta))$, and hence $h(G'(\beta)) = h(G(\beta))$ by Proposition 5. Finally, both $G(\beta)$ and $G'(\beta)$ are of period 1 because of the arrow $u \rightarrow u$. \square

Corollary 11. *Let p be a positive integer and $h \in (0, +\infty)$. There exist strongly connected graphs G, G' of period p such that $h(G) = h(G') = h$, G is positive recurrent, and G' is transient.*

Proof. For G , we start from the graph $G(\beta)$ given by Proposition 10 with $\beta = e^{hp}$. Let V denote the set of vertices of $G(\beta)$. The set of vertices of G is $V \times \llbracket 1, p \rrbracket$, and the arrows in G are

- $(v, i) \rightarrow (v, i+1)$ if $v \in V$ and $i \in \llbracket 1, p-1 \rrbracket$,
- $(v, p) \rightarrow (w, 1)$ if $v, w \in V$ and $v \rightarrow w$ is an arrow in $G(\beta)$.

According to the properties of $G(\beta)$, G is strongly connected, of period p , and positive recurrent. Moreover, $h(G) = (1/p)h(G(\beta)) = (1/p)\log \beta = h$.

For G' , we do the same starting with $G'(\beta)$. \square

According to Theorem 7, the graphs of Corollary 11 satisfy that the topological Markov chain on G admits a measure of maximal entropy whereas the topological Markov chain on G' admits no measure of maximal entropy; both are transitive, of Gurevich entropy h , and supported by a graph of period p .

References

- [Dajani and Kraaikamp 2002] K. Dajani and C. Kraaikamp, *Ergodic theory of numbers*, Carus Mathematical Monographs **29**, Mathematical Association of America, Washington, DC, 2002. MR Zbl

- [Denker et al. 1976] M. Denker, C. Grillenberger, and K. Sigmund, *Ergodic theory on compact spaces*, Lecture Notes in Mathematics **527**, Springer, 1976. MR Zbl
- [Gurevich 1969] B. M. Gurevich, “Topological entropy of a countable Markov chain”, *Dokl. Akad. Nauk SSSR* **187**:4 (1969), 715–718. In Russian; translated in *Soviet Math. Dokl.* **10** (1969), 911–915. MR Zbl
- [Gurevich 1970] B. M. Gurevich, “Shift entropy and Markov measures in the space of paths of a countable graph”, *Dokl. Akad. Nauk SSSR* **192**:5 (1970), 963–965. In Russian; translated in *Soviet Math. Dokl.* **11** (1970), 744–747. MR Zbl
- [Ruelle 2003] S. Ruelle, “On the Vere-Jones classification and existence of maximal measures for countable topological Markov chains”, *Pacific J. Math.* **209**:2 (2003), 366–380. MR Zbl
- [Salama 1988] I. A. Salama, “Topological entropy and recurrence of countable chains”, *Pacific J. Math.* **134**:2 (1988), 325–341. Correction in **140**:2 (1989), 397. MR Zbl
- [Salama 1992] I. A. Salama, “On the recurrence of countable topological Markov chains”, pp. 349–360 in *Symbolic dynamics and its applications* (New Haven, CT, 1991), edited by P. Walters, Contemp. Math. **135**, Amer. Math. Soc., Providence, RI, 1992. MR Zbl
- [Vere-Jones 1962] D. Vere-Jones, “Geometric ergodicity in denumerable Markov chains”, *Quart. J. Math. Oxford Ser. (2)* **13** (1962), 7–28. MR Zbl

Received June 26, 2018.

SYLVIE RUETTE
 LABORATOIRE DE MATHÉMATIQUES D’ORSAY, UMR 8628
 UNIVERSITÉ PARIS-SACLAY
 ORSAY
 FRANCE
sylvie.ruette@universite-paris-saclay.fr

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2019 is US \$490/year for the electronic version, and \$665/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 303 No. 1 November 2019

Contrasting various notions of convergence in geometric analysis	1
BRIAN ALLEN and CHRISTINA SORMANI	
Explicit formulae and discrepancy estimates for a -points of the Riemann zeta-function	47
SIEGFRED BALUYOT and STEVEN M. GONEK	
Diffeological vector spaces	73
J. DANIEL CHRISTENSEN and ENXIN WU	
Degree-one, monotone self-maps of the Pontryagin surface are near-homeomorphisms	93
ROBERT J. DAVERMAN and THOMAS L. THICKSTUN	
Denoetherianizing Cohen–Macaulay rings	133
LÁSZLÓ FUCHS and BRUCE OLBERDING	
Ordinary points mod p of $\mathrm{GL}_m(\mathbb{R})$ -locally symmetric spaces	165
MARK GORESKY and YUNG SHENG TAI	
Real structures on polarized Dieudonné modules	217
MARK GORESKY and YUNG SHENG TAI	
Spectrahedral representations of plane hyperbolic curves	243
MARIO KUMMER, SIMONE NALDI and DANIEL PLAUMANN	
Deformations of linear Lie brackets	265
PIER PAOLO LA PASTINA and LUCA VITAGLIANO	
A mod- p Artin–Tate conjecture, and generalizing the Herbrand–Ribet theorem	299
DIPENDRA PRASAD	
Transitive topological Markov chains of given entropy and period with or without measure of maximal entropy	317
SYLVIE RUETTE	
Restricted sum formula for finite and symmetric multiple zeta values	325
HIDEKI MURAHARA and SHINGO SAITO	
Frobenius–Schur indicators for near-group and Haagerup–Izumi fusion categories	337
HENRY TUCKER	
Compactness theorems for 4-dimensional gradient Ricci solitons	361
YONGJIA ZHANG	



0030-8730(201911)303:1;1-J