

*Pacific
Journal of
Mathematics*

**PSEUDOINDEX THEORY AND NEHARI METHOD
FOR A FRACTIONAL CHOQUARD EQUATION**

MIN LIU AND ZHONGWEI TANG

Volume 304 No. 1

January 2020

PSEUDOINDEX THEORY AND NEHARI METHOD FOR A FRACTIONAL CHOQUARD EQUATION

MIN LIU AND ZHONGWEI TANG

We study the following nonlinear fractional Choquard equation:

$$(\star) \quad \varepsilon^{2s}(-\Delta)^s w + V(x)w = \varepsilon^{-\theta} W(x)[I_\theta * (W|w|^p)]|w|^{p-2}w, \quad x \in \mathbb{R}^N,$$

where $\varepsilon > 0$, $s \in (0, 1)$, $N > 2s$, I_θ is the Riesz potential with order $\theta \in (0, N)$, $p \in [2, \frac{N+\theta}{N-2s})$, $\min_{\mathbb{R}^N} V > 0$ and $\inf_{\mathbb{R}^N} W > 0$. By specifying the ranges and interdependence of linear and nonlinear potentials, we achieve the existence, convergence, concentration, and decay estimate of positive groundstates for (\star) . The multiplicity of semiclassical solutions is established via pseudoindex theory. The existence of sign-changing solutions is constructed by minimizing the energy on Nehari nodal set.

1. Introduction and main results

This paper contributes to the multiplicity of semiclassical solutions and the convergence, concentration, decay estimate of positive groundstates for the nonlinear fractional Choquard equation,

$$(1-1) \quad \varepsilon^{2s}(-\Delta)^s w + V(x)w = \varepsilon^{-\theta} W(x)[I_\theta * (W|w|^p)]|w|^{p-2}w, \quad w \in H^s(\mathbb{R}^N),$$

where $\varepsilon > 0$, $s \in (0, 1)$, $N > 2s$, $\theta \in (0, N)$, $p \in [2, \frac{N+\theta}{N-2s})$, V and W are Hölder continuous bounded positive functions, and the Riesz potential I_θ is defined as follows:

$$(1-2) \quad I_\theta(x) := \frac{A_\theta}{|x|^{N-\theta}}, \quad x \in \mathbb{R}^N \setminus \{0\}, \quad \text{where } A_\theta := \frac{\Gamma(\frac{N-\theta}{2})}{2^\theta \pi^{N/2} \Gamma(\frac{\theta}{2})}.$$

The Choquard equation first appeared in Fröhlich and Pekar’s model of polaron [Pekar 1954] and was afterwards introduced by Ph. Choquard in the modeling of a one-component plasma [Lieb 1977]. It can be regarded as Schrödinger–Newton equation in models coupling the Schrödinger equation of quantum physics with

Tang is the corresponding author. He is supported by the National Natural Science Foundation of China (11571040, 11671331).

MSC2010: primary 35J20; secondary 35R11.

Keywords: Fractional Choquard equation, pseudoindex, sign-changing solution, concentration.

nonrelativistic Newtonian gravity and it also associates with the Einstein–Klein–Gordon and Einstein–Dirac system [Moroz and Van Schaftingen 2017].

Solitary wave solution with the type

$$\psi(x, t) = e^{-i\lambda t/\varepsilon} w(x)$$

for the time-dependent Hartree equation with $(t, x) \in (0, \infty) \times \mathbb{R}^N$,

$$i\varepsilon \frac{\partial \psi}{\partial t} = \varepsilon^{2s} (-\Delta)^s \psi + (V(x) + \lambda)\psi - \varepsilon^{-\theta} W(x) [I_\theta * (W|\psi|^p)] |\psi|^{p-2} \psi$$

corresponds to the solution of Equation (1-1). In this sense, Choquard equation is known as stationary Hartree equation.

Concerning the early works on Choquard equation, we can go back to E. H. Lieb [1977] and P. L. Lions [1980], where they both dealt with the following equation

$$(1-3) \quad -\Delta u + au = 2 \left(\frac{1}{|x|} * |u|^2 \right) u, \quad x \in \mathbb{R}^3$$

with $a \in \mathbb{R}$. Moreover, Lieb proved the existence and uniqueness of solutions by using symmetric decreasing rearrangement inequalities and Lions obtained the existence of infinitely many radially symmetric solutions.

Thereafter, the study about Choquard equation has grown steadily. Ma and Zhao [2010] settled the classification of all positive solutions to Equation (1-3) with $a = 1$ and proved that all the positive solutions of this equation must be radially symmetric and monotone decreasing about some fixed point by moving plane method. Moroz and Schaftingen [2013] and Ghimenti and Schaftingen [2016] studied Equation (1-1) with $s = \varepsilon = V(x) = W(x) = 1$ successively, the former proved the existence, regularity, symmetry and decay asymptotics of positive groundstates, and the latter constructed minimal energy odd solutions for $p \in \left(\frac{N+\theta}{N}, \frac{N+\theta}{N-2}\right)$ and minimal energy nodal solutions for $p \in \left(2, \frac{N+\theta}{N-2}\right)$. Alves, Gao, Squassina and Yang [Alves et al. 2017] considered

$$-\varepsilon^2 \Delta u + V(x)u = \varepsilon^{-\theta} W(x) \left(\frac{1}{|x|^{3-\theta}} * (WG(u)) \right) g(u), \quad x \in \mathbb{R}^3,$$

where $\varepsilon > 0$, $\theta \in (0, 3)$, V, W are continuous real functions, G is the primitive of g with a critical growth due to the Hardy–Littlewood–Sobolev inequality, and they established the existence and multiplicity of semiclassical solutions and characterized the concentration behavior by variational methods.

Recently, the discussion about fractional Choquard equation has also appeared gradually. Lenzmann [2009] proved the uniqueness of groundstate for the pseudorelativistic Hartree equation

$$\sqrt{-\Delta + m^2} u + au = \left(\frac{1}{|x|} * |u|^2 \right) u, \quad x \in \mathbb{R}^3$$

in the regime of u with sufficiently small L^2 -mass, where $m > 0$, $a \in \mathbb{R}$. Cingolani and Secchi [2015] studied the semiclassical limit for the pseudorelativistic Hartree equation

$$\sqrt{-\varepsilon^2 \Delta + m^2} u + V(x)u = (I_\theta * |u|^p) |u|^{p-2} u, \quad x \in \mathbb{R}^N,$$

where $\varepsilon > 0$, $m > 0$, $p \in [2, \frac{2N}{N-1})$, $\theta \in ((N-1)p - N, N)$ and V is a continuous and bounded function with $\inf_{\mathbb{R}^N} V > -m$. d'Avenia, Siciliano and Squassina [d'Avenia et al. 2015] investigated

$$(-\Delta)^s u + au = \left(\frac{1}{|x|^{N-\theta}} * |u|^p \right) |u|^{p-2} u, \quad x \in \mathbb{R}^N,$$

where $s \in (0, 1)$, $a > 0$, $p > 1$, $N \geq 3$, $\theta \in (0, N)$, and they obtained the regularity, existence, nonexistence, symmetry as well as decay properties of groundstates. Bhattarai [2017] obtained the existence and stability results of solutions for the following fractional Choquard equation with combined power and Hartree type nonlinearities

$$(-\Delta)^s u + au = b|u|^{r-2} u + \lambda \left(\frac{1}{|x|^{N-\theta}} * |u|^p \right) |u|^{p-2} u, \quad x \in \mathbb{R}^N,$$

where $s \in (0, 1)$, $N \geq 2$, $\theta \in (0, N)$, $r \in (2, 2 + \frac{4s}{N})$, $p \in [2, 1 + \frac{2s+\theta}{N})$, $a \in \mathbb{R}$ and b, λ are nonnegative constants satisfying $b + \lambda \neq 0$.

For more results about Choquard equation, see [Belchior et al. 2017; Moroz and Van Schaftingen 2017; Zhong and Tang 2018]. For classical local case, we mainly refer to [Ding and Wei 2017]. For fractional elliptic problems, we refer to [Ao et al. 2017; Brändle et al. 2013; Dávila et al. 2014; Dipierro et al. 2017; Fall et al. 2015; Felmer et al. 2012; Jin et al. 2014].

Ding and Wei [2017] considered the following Schrödinger equation:

$$-\varepsilon^2 \Delta w + V(x)w = W(x)|w|^{p-2} w, \quad x \in \mathbb{R}^N,$$

where $\varepsilon > 0$, $p \in (2, \frac{2N}{N-2})$ and V, W are continuous bounded positive functions. They studied the existence and concentration of positive groundstates and they also constructed the multiplicity of semiclassical solutions including at least a pair of sign-changing solutions by pseudoindex theory and Nehari method.

With regard to the multiplicity of solutions for fractional elliptic problems, Lyapunov–Schmidt reduction is also an important method. We can take [Ao et al. 2017; Dávila et al. 2014] as examples. Dávila, del Pino and Wei [Dávila et al. 2014] considered the following fractional nonlinear Schrödinger equation

$$\varepsilon^{2s} (-\Delta)^s u + V(x)u = |u|^{p-1} u, \quad x \in \mathbb{R}^N,$$

where $\varepsilon > 0$, $s \in (0, 1)$, $p \in (1, \frac{N+2s}{N-2s})$ and $V \in C^{1,\alpha}(\mathbb{R}^N) \cap L^\infty(\mathbb{R}^N)$ with $\inf_{\mathbb{R}^N} V > 0$. For any positive integer k , the authors proved the existence of k -spike solution in $H^{2s}(\mathbb{R}^N)$ by applying Lyapunov–Schmidt variational reduction together with appropriate assumptions. Ao, Wei and Yang [Ao et al. 2017] later studied the above equation with $\varepsilon = 1$, $N = 2$ and constructed infinitely many nonradial high energy positive solutions for it by using an intermediate Lyapunov–Schmidt reduction and assuming

$$V(x) = V_\infty + \frac{a}{|x|^m} + O\left(\frac{1}{|x|^{m+\sigma}}\right), \quad \text{as } |x| \rightarrow \infty.$$

Motivated by the works cited above, we wish to study the multiplicity and concentration of solutions for the nonlinear fractional Choquard equation (1-1) by using pseudoindex theory and the Nehari method. Referring to [Ding and Wei 2017], we shall consider the ranges and interdependence of linear and nonlinear potentials to be the essential assumptions. According to [Felmer et al. 2012], we will suppose the two potentials to be Hölder continuous in order to lift the regularity of solutions and ensure the representation (2-2) holds. Compared with [Ding and Wei 2017], there appear two nonlocal terms in our equation, which are fractional Laplace operator and convolution type Hartree term, which bring us more difficulties. In order to overcome the nonlocality of the fractional Laplace operator, in some cases we will adopt the extension method proposed in [Caffarelli and Silvestre 2007]. On the Hartree term, the Hardy–Littlewood–Sobolev inequality and the Brézis–Lieb type lemma (see Lemma 3.5) play an important role. We finally obtain the existence of multiple solutions, groundstates and sign-changing solutions for Equation (1-1). In addition, we prove the positive groundstate concentrates in a special set related to the minimum of linear potential and the maximum of nonlinear potential.

Now we state our assumptions and main results.

(A0) $V, W \in C^{0,\lambda}(\mathbb{R}^N, \mathbb{R})$ are bounded with some $\lambda \in (0, 1)$, V achieves a global minimum on \mathbb{R}^N with $\min_{\mathbb{R}^N} V > 0$, and W achieves a global maximum on \mathbb{R}^N with $\inf_{\mathbb{R}^N} W > 0$.

To describe our results and proofs, denote

$$\begin{aligned} \tau &:= \min_{\mathbb{R}^N} V, & \mathcal{V} &:= \{x \in \mathbb{R}^N : V(x) = \tau\}, & \tau_\infty &:= \liminf_{|x| \rightarrow \infty} V(x); \\ k &:= \max_{\mathbb{R}^N} W, & \mathcal{W} &:= \{x \in \mathbb{R}^N : W(x) = k\}, & k_\infty &:= \limsup_{|x| \rightarrow \infty} W(x); \\ x_v &\in \mathcal{V} : k_v := W(x_v) = \max_{\mathcal{V}} W; \\ x_w &\in \mathcal{W} : \tau_w := V(x_w) = \min_{\mathcal{W}} V. \end{aligned}$$

(A1) Either (i) or (ii) holds, where

- (i) $\tau < \tau_\infty$, and there exists $R_v > 0$ such that $W(x) \leq k_v$, for all $|x| \geq R_v$;
- (ii) $k > k_\infty$, and there exists $R_w > 0$ such that $V(x) \geq \tau_w$, for all $|x| \geq R_w$.

If (A1)(i) holds, define $\mathcal{A}_v := \{x \in \mathcal{V} : W(x) = k_v\} \cup \{x \notin \mathcal{V} : W(x) > k_v\}$.

If (A1)(ii) holds, define $\mathcal{A}_w := \{x \in \mathcal{W} : V(x) = \tau_w\} \cup \{x \notin \mathcal{W} : V(x) < \tau_w\}$.

In what follows, \mathcal{A} stands for \mathcal{A}_v in the case (A1)(i) and \mathcal{A}_w in the case (A1)(ii). Clearly, \mathcal{A} is bounded. Moreover, $\mathcal{A} = \mathcal{V} \cap \mathcal{W}$ if $\mathcal{V} \cap \mathcal{W} \neq \emptyset$.

Theorem 1.1. *Assume that (A0) holds and*

$$(1-4) \quad \tau < \tau_\infty, \quad k_v \geq k_\infty.$$

Then for the maximal integer $m \in \mathbb{N}$ with

$$(1-5) \quad m < \left(\frac{\tau_\infty}{\tau}\right)^{\frac{\theta+2s}{2s(p-1)} - \frac{N-2s}{2s}} \cdot \left(\frac{k_v}{k_\infty}\right)^{\frac{2}{p-1}},$$

Equation (1-1) possesses at least m pairs of solutions for small $\varepsilon > 0$. Furthermore, when $m \geq 2$ and $p \in (2, \frac{N+\theta}{N-2s})$, among the solutions, at least one is positive, one is negative and two change sign.

Theorem 1.2. *Assume that (A0) holds and*

$$(1-6) \quad \tau_w \leq \tau_\infty, \quad k > k_\infty.$$

Then for the maximal integer $m \in \mathbb{N}$ with

$$(1-7) \quad m < \left(\frac{\tau_\infty}{\tau_w}\right)^{\frac{\theta+2s}{2s(p-1)} - \frac{N-2s}{2s}} \cdot \left(\frac{k}{k_\infty}\right)^{\frac{2}{p-1}},$$

all the conclusions of Theorem 1.1 remain true.

Theorem 1.3. *Assume that (A0)–(A1) hold. Then for sufficiently small $\varepsilon > 0$, Equation (1-1) has a positive groundstate w_ε . If additionally $V, W \in C^1(\mathbb{R}^N, \mathbb{R})$ and $\nabla V, \nabla W$ are bounded, then w_ε satisfies:*

- (i) (concentration) *There exists a maximum point x_ε of w_ε with*

$$\lim_{\varepsilon \rightarrow 0} \text{dist}(x_\varepsilon, \mathcal{A}) = 0.$$

- (ii) (decay estimate) *For $p \in (2, \frac{N+\theta}{N-2s})$, there exist $C_2 > C_1 > 0$ and sufficiently large $R > 0$ such that*

$$\frac{C_1 \varepsilon^{N+2s}}{|x - x_\varepsilon|^{N+2s}} \leq w_\varepsilon(x) \leq \frac{C_2 \varepsilon^{N+2s}}{|x - x_\varepsilon|^{N+2s}}, \quad \text{for all } |x| \geq R.$$

- (iii) (convergence) *Setting $u_\varepsilon(x) := w_\varepsilon(\varepsilon x + x_\varepsilon)$, for any sequence $x_\varepsilon \rightarrow x_0$ (as $\varepsilon \rightarrow 0$),*

$$u_\varepsilon \rightarrow u \quad \text{in } H^s(\mathbb{R}^N),$$

where u is a groundstate of

$$(1-8) \quad (-\Delta)^s u + V(x_0)u = W^2(x_0)(I_\theta * u^p)u^{p-1}, \quad u > 0.$$

In particular, if $\mathcal{V} \cap \mathcal{W} \neq \emptyset$, then $\lim_{\varepsilon \rightarrow 0} \text{dist}(x_\varepsilon, \mathcal{V} \cap \mathcal{W}) = 0$ and up to a sequence, $u_\varepsilon \rightarrow u$ in $H^s(\mathbb{R}^N)$ as $\varepsilon \rightarrow 0$, with u being a groundstate of

$$(1-9) \quad (-\Delta)^s u + \tau u = k^2(I_\theta * u^p)u^{p-1}, \quad u > 0.$$

Our paper is organized as follows: Section 2 is a review of the fractional Sobolev space and the fractional Laplace operator. Section 3 offers some valuable information about the Riesz potential. Section 4 contains some preliminary results which are established by variational methods and play a key role in the proofs of main theorems. Sections 5 and 6 contribute to the proofs of main results. In Section 5, we prove the multiplicity of semiclassical solutions via Benci pseudoindex theory and show the existence of the groundstates and the sign-changing solutions. In Section 6, we discuss the convergence, concentration and decay estimate of the positive groundstate.

2. Fractional Sobolev space and fractional Laplace operator

In this section, we shall recall some important facts about the fractional Sobolev space and the fractional Laplace operator.

For any $s \in (0, 1)$, the fractional Sobolev space $H^s(\mathbb{R}^N)$ is defined by

$$H^s(\mathbb{R}^N) = \left\{ u \in L^2(\mathbb{R}^N) : \frac{|u(x) - u(y)|}{|x - y|^{(N+2s)/2}} \in L^2(\mathbb{R}^N \times \mathbb{R}^N) \right\}$$

endowed with the natural norm

$$\|u\|_{H^s(\mathbb{R}^N)} = \left(\int_{\mathbb{R}^N} u^2 \, dx + \int_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} \, dx \, dy \right)^{1/2},$$

where the term

$$[u]_{H^s(\mathbb{R}^N)} := \left(\int_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} \, dx \, dy \right)^{1/2}$$

is the Gagliardo seminorm of u .

For any $u \in H^s(\mathbb{R}^N)$, the fractional Laplace operator $(-\Delta)^s$ is defined via Fourier transform by

$$(2-1) \quad \widehat{(-\Delta)^s u} = |\xi|^{2s} \hat{u} \in L^2(\mathbb{R}^N),$$

where $\widehat{\cdot}$ stands for the Fourier transform. When u is assumed additionally smooth, one can obtain the direct representation

$$(2-2) \quad (-\Delta)^s u(x) = -\frac{C_{N,s}}{2} \int_{\mathbb{R}^N} \frac{u(x+y) + u(x-y) - 2u(x)}{|y|^{N+2s}} dy,$$

where $C_{N,s} := (\int_{\mathbb{R}^N} (1 - \cos \zeta_1) / |\zeta|^{N+2s} d\zeta)^{-1}$ is the dimensional constant.

Lemma 2.1 [Di Nezza et al. 2012]. *Let $s \in (0, 1)$ and $u \in H^s(\mathbb{R}^N)$. Then*

$$[u]_{H^s(\mathbb{R}^N)}^2 = 2C_{N,s}^{-1} \int_{\mathbb{R}^N} |\xi|^{2s} |\widehat{u}(\xi)|^2 d\xi = 2C_{N,s}^{-1} \|(-\Delta)^{s/2} u\|_{L^2(\mathbb{R}^N)}^2.$$

Thus one can see the fractional Sobolev space $H^s(\mathbb{R}^N)$ defined as above coincides with

$$\widehat{H}^s(\mathbb{R}^N) = \left\{ u \in L^2(\mathbb{R}^N) : \int_{\mathbb{R}^N} (1 + |\xi|^{2s}) |\widehat{u}(\xi)|^2 d\xi < \infty \right\}.$$

Obviously, the following norms on $H^s(\mathbb{R}^N)$ are all equivalent:

$$\begin{aligned} u &\mapsto \left(\int_{\mathbb{R}^N} u^2 dx + \int_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy \right)^{1/2}, \\ u &\mapsto \left(\int_{\mathbb{R}^N} u^2 dx + \int_{\mathbb{R}^N} |\xi|^{2s} |\widehat{u}(\xi)|^2 d\xi \right)^{1/2}, \\ u &\mapsto \left(\int_{\mathbb{R}^N} u^2 dx + \int_{\mathbb{R}^N} |(-\Delta)^{s/2} u|^2 dx \right)^{1/2}. \end{aligned}$$

The dual space $H^{-s}(\mathbb{R}^N)$ of $H^s(\mathbb{R}^N)$ is defined in the standard way.

Caffarelli and Silvestre [2007] introduced another valuable local representation of $(-\Delta)^s$, which is via the following boundary value problem

$$\begin{cases} \operatorname{div}(y^{1-2s} \nabla \tilde{u}) = 0 & \text{in } \mathbb{R}_+^{N+1} := \{(x, y) : x \in \mathbb{R}^N, y > 0\}, \\ \tilde{u}(x, 0) = u(x) & \text{on } \mathbb{R}^N. \end{cases}$$

There, \tilde{u} is called the s -harmonic extension of u , and \tilde{u} belongs to

$$X^s(\mathbb{R}_+^{N+1}) := \overline{C_0^\infty(\mathbb{R}_+^{N+1})}^{\|\cdot\|_{X^s(\mathbb{R}_+^{N+1})}}$$

with the norm

$$\|\tilde{u}\|_{X^s(\mathbb{R}_+^{N+1})} := \left(C_s \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{u}|^2 dx dy \right)^{1/2}$$

and the inner product

$$(\tilde{u}, \tilde{v})_{X^s(\mathbb{R}_+^{N+1})} := C_s \int_{\mathbb{R}_+^{N+1}} y^{1-2s} \nabla \tilde{u} \cdot \nabla \tilde{v} dx dy,$$

where $C_s := \Gamma(s)/(2^{1-2s}\Gamma(1-s))$, is a normalization constant ensuring

$$(2-3) \quad \|\tilde{u}\|_{X^s(\mathbb{R}_+^{N+1})}^2 = \int_{\mathbb{R}^N} |\xi|^{2s} |\hat{u}(\xi)|^2 d\xi = \|(-\Delta)^{s/2} u\|_{L^2(\mathbb{R}^N)}^2.$$

Moreover, \tilde{u} can be explicitly expressed by

$$(2-4) \quad \tilde{u}(x, y) = \int_{\mathbb{R}^N} P_s(x-z, y) u(z) dz,$$

where

$$P_s(x, y) = \frac{d_{N,s} y^{2s}}{(|x|^2 + y^2)^{(N+2s)/2}}$$

is the s -Poisson kernel. It has been proved in [Caffarelli and Silvestre 2007] that

$$(2-5) \quad (-\Delta)^s u(x) = -C_s \lim_{y \rightarrow 0^+} y^{1-2s} \partial_y \tilde{u}(x, y).$$

The characterizations (2-1), (2-2) and (2-5) are equivalent, for instance, in Schwartz's space of rapidly decaying C^∞ functions on \mathbb{R}^N . For more properties of the fractional Laplace operator, we refer readers to Section 2 of [Silvestre 2007]. In the following, we set $2_s^* := 2N/(N-2s)$.

Lemma 2.2 [Di Nezza et al. 2012]. *Let $s \in (0, 1)$. Then $H^s(\mathbb{R}^N)$ is continuously embedded into $L^q(\mathbb{R}^N)$ for $q \in [2, 2_s^*]$ and compactly embedded into $L_{\text{loc}}^q(\mathbb{R}^N)$ for $q \in [2, 2_s^*)$.*

Lemma 2.3 [d'Avenia et al. 2015]. *Let $s \in (0, 1)$, $\varrho > 0$, $q \in [2, 2_s^*)$. If $\{u_n\}$ is bounded in $H^s(\mathbb{R}^N)$ and*

$$\sup_{y \in \mathbb{R}^N} \int_{B_\varrho(y)} |u_n|^q dx \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

then $u_n \rightarrow 0$ in $L^r(\mathbb{R}^N)$ for any $r \in (2, 2_s^)$.*

Lemma 2.4 [Brändle et al. 2013]. *Let $s \in (0, 1)$ and $U \in X^s(\mathbb{R}_+^{N+1})$. Then*

$$\left(\int_{\mathbb{R}^N} |u|^{2_s^*} dx \right)^{2/2_s^*} \leq k_{N,s} \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla U|^2 dx dy,$$

where

$$u(x) = \text{tr}_{\mathbb{R}^N} U =: U(x, 0) \quad \text{and} \quad k_{N,s} = \frac{\Gamma(s)\Gamma(\frac{N-2s}{2})}{2\pi^s \Gamma(1-s)\Gamma(\frac{N+2s}{2})} \left[\frac{\Gamma(N)}{\Gamma(N/2)} \right]^{2s/N}.$$

Lemma 2.5 [Dipierro et al. 2017]. *Let $r > 0$, $B_r^+ := \{(x, y) \in \mathbb{R}_+^{N+1} : |(x, y)| < r\}$ and \mathcal{T} be a subset of $X^s(\mathbb{R}_+^{N+1})$ such that*

$$\sup_{U \in \mathcal{T}} \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla U|^2 dx dy < \infty.$$

Then \mathcal{T} is precompact in $L^2(B_r^+, y^{1-2s})$. Namely, $X^s(\mathbb{R}_+^{N+1})$ is compactly embedded into $L_{\text{loc}}^2(\mathbb{R}_+^{N+1}, y^{1-2s})$ with

$$\|U\|_{L^2(\mathbb{R}_+^{N+1}, y^{1-2s})} := \left(\int_{\mathbb{R}_+^{N+1}} y^{1-2s} U^2(x, y) \, dx \, dy \right)^{1/2}.$$

Now we define a Hilbert space

$$E := \left\{ U \in X^s(\mathbb{R}_+^{N+1}) : \int_{\mathbb{R}^N} U^2(x, 0) \, dx < \infty \right\}$$

with norm

$$\|U\|_E := \left(C_s \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla U|^2 \, dx \, dy + \int_{\mathbb{R}^N} U^2(x, 0) \, dx \right)^{1/2}$$

and inner product

$$(U_1, U_2)_E := C_s \int_{\mathbb{R}_+^{N+1}} y^{1-2s} \nabla U_1 \cdot \nabla U_2 \, dx \, dy + \int_{\mathbb{R}^N} U_1(x, 0) U_2(x, 0) \, dx.$$

Clearly, $\text{tr}_{\mathbb{R}^N} E = H^s(\mathbb{R}^N)$. By Lemma 2.4, E is continuously embedded into $L^q(\mathbb{R}^N)$ for $q \in [2, 2_s^*]$. Moreover, E is compactly embedded into $L_{\text{loc}}^q(\mathbb{R}^N)$ for $q \in [2, 2_s^*)$.

In the following, we drop the constant C_s for simplicity and without destruction. Moreover, we shall use different patterns of C to denote various positive constants, and $o(1)$ to denote the quantities that tend to 0 as $n \rightarrow \infty$ or $j \rightarrow \infty$. Meanwhile, set

$$\begin{aligned} \|u\|_s &:= \|u\|_{H^s(\mathbb{R}^N)}, & |u|_q &:= \|u\|_{L^q(\mathbb{R}^N)}, \\ \|u\|_X &:= \|u\|_{X^s(\mathbb{R}_+^{N+1})}, & (u, v)_X &:= (u, v)_{X^s(\mathbb{R}_+^{N+1})}, \\ u^+ &:= \max\{0, u\}, & u^- &:= \min\{0, u\}, & \mathbb{R}_+ &:= (0, \infty). \end{aligned}$$

3. Riesz potential

In this section, we shall review and prove some useful results about the Riesz potential.

The Riesz potential with order $\theta \in (0, N)$ of a function $f \in L_{\text{loc}}^1(\mathbb{R}^N)$ is defined by

$$(3-1) \quad (I_\theta * f)(x) := \int_{\mathbb{R}^N} \frac{A_\theta f(y)}{|x-y|^{N-\theta}} \, dy,$$

where A_θ is the same as in (1-2). The integral in (3-1) converges in the classical Lebesgue sense for a.e. $x \in \mathbb{R}^N$ if and only if

$$(3-2) \quad f \in L^1(\mathbb{R}^N, (1+|x|)^{\theta-N}).$$

Moreover, if (3-2) doesn't hold, then (3-1) diverges everywhere on \mathbb{R}^N . In addition, if $f \in C^{0,\nu}(\mathbb{R}^N)$ with $\nu \in (0, 1)$, then either $I_\theta * f \in C^{0,\nu+\theta}(\mathbb{R}^N)$ for $\nu+\theta \in (0, 1)$ or $I_\theta * f \in C^{0,1}(\mathbb{R}^N)$ for $\nu+\theta \in [1, N+1)$ (see Theorem 1 and its proof in [du Plessis 1955]).

The Riesz potential I_θ is well-defined as an operator in $L^q(\mathbb{R}^N)$ if and only if $q \in [1, \frac{N}{\theta})$. Furthermore, if $q \in (1, \frac{N}{\theta})$ and $r := \frac{Nq}{N-\theta q}$, then

$$I_\theta : L^q(\mathbb{R}^N) \rightarrow L^r(\mathbb{R}^N)$$

is a bounded linear operator, which can be disclosed by the Hardy–Littlewood–Sobolev inequality.

Lemma 3.1 [Hardy et al. 1952]. *Let $\theta \in (0, N)$, $q \in (1, \frac{N}{\theta})$. Then for any $f \in L^q(\mathbb{R}^N)$, $I_\theta * f \in L^{Nq/(N-\theta q)}(\mathbb{R}^N)$ and*

$$\left(\int_{\mathbb{R}^N} |I_\theta * f|^{Nq/(N-\theta q)} dx \right)^{\frac{1}{q} - \frac{\theta}{N}} \leq C_{N,\theta,q} \left(\int_{\mathbb{R}^N} |f|^q dx \right)^{\frac{1}{q}}.$$

Applying Lemma 3.1 to the function $f = |u|^p \in L^{2N/(N+\theta)}(\mathbb{R}^N)$, we get the following result.

Lemma 3.2 [Moroz and Van Schaftingen 2017]. *Let $\theta \in (0, N)$. Then for any $u \in L^{2Np/(N+\theta)}(\mathbb{R}^N)$,*

$$\int_{\mathbb{R}^N} (I_\theta * |u|^p) |u|^p dx \leq C_{N,\theta} \left(\int_{\mathbb{R}^N} |u|^{2Np/(N+\theta)} dx \right)^{(N+\theta)/N}.$$

If, in particular, $s \in (0, 1)$, $N > 2s$, $\theta \in (0, N)$, $p \in [\frac{N+\theta}{N}, \frac{N+\theta}{N-2s}]$ and $u \in H^s(\mathbb{R}^N)$, then

$$\int_{\mathbb{R}^N} (I_\theta * |u|^p) |u|^p dx \leq C_{N,\theta,p} \|u\|_s^{2p}.$$

As a matter of fact, $p \in [\frac{N+\theta}{N}, \frac{N+\theta}{N-2s}]$ if and only if $\frac{2Np}{N+\theta} \in [2, 2s^*]$.

It's worthwhile to mention that the Brézis–Lieb type lemma holds for the Riesz potential. To prove this lemma, we need two essential results.

Lemma 3.3. *Let Ω be a domain on \mathbb{R}^N and $q, r \in (1, \infty)$, $\frac{1}{q} + \frac{1}{r} = 1$. If $u_n \rightarrow u$ in $L^q(\Omega)$ and $v_n \rightharpoonup v$ in $L^r(\Omega)$ as $n \rightarrow \infty$, then*

$$\lim_{n \rightarrow \infty} \int_{\Omega} u_n v_n dx = \int_{\Omega} u v dx.$$

Lemma 3.4. *Let Ω be a domain on \mathbb{R}^N , $r \in [1, \infty)$ and $\{u_n\}$ be a bounded sequence in $L^r(\Omega)$. If $u_n \rightarrow u$ a.e. in Ω as $n \rightarrow \infty$, then for any $q \in [1, r]$,*

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\Omega} \left| |u_n|^q - |u_n - u|^q - |u|^q \right|^{r/q} dx &= 0, \\ \lim_{n \rightarrow \infty} \int_{\Omega} \left| |u_n^\pm|^q - |(u_n - u)^\pm|^q - |u^\pm|^q \right|^{r/q} dx &= 0. \end{aligned}$$

We would like to point out that the constraint $p \in [2, \frac{N+\theta}{N-2s})$ is only needed for the second part of the following Brézis–Lieb type lemma, while the first part permits $p \in [\frac{N+\theta}{N}, \frac{N+\theta}{N-2s}]$.

Lemma 3.5. *Let $s \in (0, 1)$, $N > 2s$, $\theta \in (0, N)$, $p \in [2, \frac{N+\theta}{N-2s})$. If $u_n \rightharpoonup u$ in $H^s(\mathbb{R}^N)$ as $n \rightarrow \infty$, then*

- (i) $\mathcal{D}(u_n) - \mathcal{D}(u_n - u) \rightarrow \mathcal{D}(u)$ as $n \rightarrow \infty$;
- (ii) $\mathcal{D}'(u_n) - \mathcal{D}'(u_n - u) \rightarrow \mathcal{D}'(u)$ in $H^{-s}(\mathbb{R}^N)$ as $n \rightarrow \infty$,

where

$$\mathcal{D}(u) := \int_{\mathbb{R}^N} (I_\theta * |u|^p) |u|^p dx.$$

Proof. The proof is similar to that of Lemma 2.5 in [Liu and Tang 2019] with Lemmas 3.3 and 3.4 being used, and so it is omitted. \square

In terms of Lemma 3.5, the following result is absolutely true.

Lemma 3.6. *Let $s \in (0, 1)$, $N > 2s$, $\theta \in (0, N)$, $p \in [2, \frac{N+\theta}{N-2s})$. If $u_n \rightharpoonup u$ in $H^s(\mathbb{R}^N)$ as $n \rightarrow \infty$, then for any $v \in H^s(\mathbb{R}^N)$,*

$$\langle \mathcal{D}'(u_n), v \rangle \rightarrow \langle \mathcal{D}'(u), v \rangle \quad \text{as } n \rightarrow \infty,$$

where $\mathcal{D}(u)$ is defined as in Lemma 3.5.

4. Preliminary results

In this section, we establish some results which are necessary for the arguments of our main results. Let us first consider the following two equations for $s \in (0, 1)$, $N > 2s$, $\theta \in (0, N)$, $p \in [2, \frac{N+\theta}{N-2s})$,

$$(4-1) \quad (-\Delta)^s v + av = b^2(I_\theta * |v|^p) |v|^{p-2}v, \quad v \in H^s(\mathbb{R}^N),$$

where $a > 0$, $b > 0$, and

$$(4-2) \quad (-\Delta)^s v + V_\varepsilon^a(x)v = W_\varepsilon^b(x)[I_\theta * (W_\varepsilon^b |v|^p)] |v|^{p-2}v, \quad v \in H^s(\mathbb{R}^N),$$

where $\tau \leq a \leq \tau_\infty$, $k_\infty \leq b \leq k$ and

$$\begin{aligned} V^a(x) &:= \max\{a, V(x)\}, & V_\varepsilon^a(x) &:= V^a(\varepsilon x), \\ W^b(x) &:= \min\{b, W(x)\}, & W_\varepsilon^b(x) &:= W^b(\varepsilon x). \end{aligned}$$

Definition 4.1. (i) $v \in H^s(\mathbb{R}^N)$ is a weak solution of Equation (4-1) if for any $\varphi \in H^s(\mathbb{R}^N)$,

$$\int_{\mathbb{R}^N} ((-\Delta)^{s/2}v(-\Delta)^{s/2}\varphi + av\varphi) dx = \int_{\mathbb{R}^N} b^2(I_\theta * |v|^p) |v|^{p-2}v\varphi dx.$$

(ii) $v_\varepsilon \in H^s(\mathbb{R}^N)$ is a weak solution of Equation (4-2) if for any $\varphi \in H^s(\mathbb{R}^N)$,

$$\begin{aligned} \int_{\mathbb{R}^N} ((-\Delta)^{s/2} v_\varepsilon (-\Delta)^{s/2} \varphi + V_\varepsilon^a(x) v_\varepsilon \varphi) dx \\ = \int_{\mathbb{R}^N} W_\varepsilon^b(x) [I_\theta * (W_\varepsilon^b |v_\varepsilon|^p)] |v_\varepsilon|^{p-2} v_\varepsilon \varphi dx. \end{aligned}$$

Associated with Equation (4-1) and Equation (4-2) respectively, we define the energy functionals for each $v \in H^s(\mathbb{R}^N)$,

$$\begin{aligned} J^{ab}(v) &:= \frac{1}{2} \int_{\mathbb{R}^N} (|(-\Delta)^{s/2} v|^2 + av^2) dx - \frac{1}{2p} \int_{\mathbb{R}^N} b^2 (I_\theta * |v|^p) |v|^p dx, \\ J_\varepsilon^{ab}(v) &:= \frac{1}{2} \int_{\mathbb{R}^N} (|(-\Delta)^{s/2} v|^2 + V_\varepsilon^a(x) v^2) dx - \frac{1}{2p} \int_{\mathbb{R}^N} W_\varepsilon^b(x) [I_\theta * (W_\varepsilon^b |v|^p)] |v|^p dx; \end{aligned}$$

the Nehari manifolds

$$\begin{aligned} \mathcal{N}^{ab} &:= \{v \in H^s(\mathbb{R}^N) \setminus \{0\} : \langle (J^{ab})'(v), v \rangle = 0\}, \\ \mathcal{N}_\varepsilon^{ab} &:= \{v \in H^s(\mathbb{R}^N) \setminus \{0\} : \langle (J_\varepsilon^{ab})'(v), v \rangle = 0\}; \end{aligned}$$

the least energies

$$\vartheta^{ab} := \inf_{\mathcal{N}^{ab}} J^{ab}, \quad \vartheta_\varepsilon^{ab} := \inf_{\mathcal{N}_\varepsilon^{ab}} J_\varepsilon^{ab};$$

and the sets of least energy solutions

$$\begin{aligned} \mathcal{A}^{ab} &:= \{v \in H^s(\mathbb{R}^N) : J^{ab}(v) = \vartheta^{ab}, (J^{ab})'(v) = 0\}, \\ \mathcal{A}_\varepsilon^{ab} &:= \{v_\varepsilon \in H^s(\mathbb{R}^N) : J_\varepsilon^{ab}(v_\varepsilon) = \vartheta_\varepsilon^{ab}, (J_\varepsilon^{ab})'(v_\varepsilon) = 0\}. \end{aligned}$$

In particular, we set

$$\begin{aligned} J^\infty &:= J^{\tau_\infty k_\infty}, \quad \mathcal{N}^\infty := \mathcal{N}^{\tau_\infty k_\infty}, \quad \vartheta^\infty := \vartheta^{\tau_\infty k_\infty}, \quad V_\varepsilon^\infty := V_\varepsilon^{\tau_\infty}, \\ J_\varepsilon^\infty &:= J_\varepsilon^{\tau_\infty k_\infty}, \quad \mathcal{N}_\varepsilon^\infty := \mathcal{N}_\varepsilon^{\tau_\infty k_\infty}, \quad \vartheta_\varepsilon^\infty := \vartheta_\varepsilon^{\tau_\infty k_\infty}, \quad W_\varepsilon^\infty := W_\varepsilon^{k_\infty}. \end{aligned}$$

Moreover, $J^{ab}, J_\varepsilon^{ab} \in C^1(H^s(\mathbb{R}^N), \mathbb{R})$. The critical points of J^{ab} and J_ε^{ab} correspond to the weak solutions of Equation (4-1) and Equation (4-2), respectively.

Additionally, Equations (4-1) and (4-2) can be reformulated as

$$(4-3) \quad \begin{cases} \operatorname{div}(y^{1-2s} \nabla \tilde{v}) = 0 & \text{in } \mathbb{R}_+^{N+1}, \\ \tilde{v}(x, 0) = v(x) & \text{on } \mathbb{R}^N, \\ \lim_{y \rightarrow 0^+} y^{1-2s} \partial_y \tilde{v} = av - b^2 (I_\theta * |v|^p) |v|^{p-2} v & \text{in } \mathbb{R}_+^{N+1}, \end{cases}$$

and

$$(4-4) \quad \begin{cases} \operatorname{div}(y^{1-2s} \nabla \tilde{v}) = 0 & \text{in } \mathbb{R}_+^{N+1}, \\ \tilde{v}(x, 0) = v(x) & \text{on } \mathbb{R}^N, \\ \lim_{y \rightarrow 0^+} y^{1-2s} \partial_y \tilde{v} = V_\varepsilon^a(x) v - W_\varepsilon^b(x) [I_\theta * (W_\varepsilon^b |v|^p)] |v|^{p-2} v & \text{in } \mathbb{R}_+^{N+1}, \end{cases}$$

respectively, where $\tilde{v} \in E$ is the s -harmonic extension of $v \in H^s(\mathbb{R}^N)$.

Definition 4.2. (i) $\tilde{v} \in E$ is a weak solution of problem (4-3) if for all $\tilde{\varphi} \in E$

$$\int_{\mathbb{R}_+^{N+1}} y^{1-2s} \nabla \tilde{v} \cdot \nabla \tilde{\varphi} \, dx \, dy + \int_{\mathbb{R}^N} a v \varphi \, dx = \int_{\mathbb{R}^N} b^2 (I_\theta * |v|^p) |v|^{p-2} v \varphi \, dx,$$

(ii) $\tilde{v}_\varepsilon \in E$ is a weak solution of problem (4-4) if for all $\tilde{\varphi} \in E$

$$\begin{aligned} \int_{\mathbb{R}_+^{N+1}} y^{1-2s} \nabla \tilde{v}_\varepsilon \cdot \nabla \tilde{\varphi} \, dx \, dy + \int_{\mathbb{R}^N} V_\varepsilon^a(x) v_\varepsilon \varphi \, dx \\ = \int_{\mathbb{R}^N} W_\varepsilon^b(x) [I_\theta * (W_\varepsilon^b |v|^p)] |v_\varepsilon|^{p-2} v_\varepsilon \varphi \, dx, \end{aligned}$$

where $v = \text{tr}_{\mathbb{R}^N} \tilde{v}$, $v_\varepsilon = \text{tr}_{\mathbb{R}^N} \tilde{v}_\varepsilon$, $\varphi = \text{tr}_{\mathbb{R}^N} \tilde{\varphi}$.

Associated with (4-3) and (4-4) respectively, we define the energy functionals for each $\tilde{v} \in E$,

$$\tilde{J}^{ab}(\tilde{v}) := \frac{1}{2} \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{v}|^2 \, dx \, dy + \frac{1}{2} \int_{\mathbb{R}^N} a v^2 \, dx - \frac{1}{2p} \int_{\mathbb{R}^N} b^2 (I_\theta * |v|^p) |v|^p \, dx,$$

$$\begin{aligned} \tilde{J}_\varepsilon^{ab}(\tilde{v}) := \frac{1}{2} \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{v}|^2 \, dx \, dy + \frac{1}{2} \int_{\mathbb{R}^N} V_\varepsilon^a(x) v^2 \, dx \\ - \frac{1}{2p} \int_{\mathbb{R}^N} W_\varepsilon^b(x) [I_\theta * (W_\varepsilon^b |v|^p)] |v|^p \, dx; \end{aligned}$$

and we denote the Nehari manifolds by $\tilde{\mathcal{N}}^{ab}$, $\tilde{\mathcal{N}}_\varepsilon^{ab}$; the least energies by $\tilde{\vartheta}^{ab}$, $\tilde{\vartheta}_\varepsilon^{ab}$; and the sets of least energy solutions by $\tilde{\mathcal{H}}^{ab}$, $\tilde{\mathcal{H}}_\varepsilon^{ab}$.

In addition, \tilde{J}^{ab} , $\tilde{J}_\varepsilon^{ab} \in C^1(E, \mathbb{R})$. The critical points of \tilde{J}^{ab} and $\tilde{J}_\varepsilon^{ab}$ correspond to the weak solutions of (4-3) and (4-4), respectively.

In terms of (2-3), we find that

$$\tilde{J}^{ab}(\tilde{v}) = J^{ab}(v), \quad \tilde{J}_\varepsilon^{ab}(\tilde{v}) = J_\varepsilon^{ab}(v)$$

for any $\tilde{v} \in E$ as the s -harmonic extension of $v \in H^s(\mathbb{R}^N)$. And $v \in \mathcal{N}^{ab}$ if and only if $\tilde{v} \in \tilde{\mathcal{N}}^{ab}$, $v \in \mathcal{N}_\varepsilon^{ab}$ if and only if $\tilde{v} \in \tilde{\mathcal{N}}_\varepsilon^{ab}$. Hence

$$\tilde{\vartheta}^{ab} = \vartheta^{ab}, \quad \tilde{\vartheta}_\varepsilon^{ab} = \vartheta_\varepsilon^{ab}.$$

Indeed, $\tilde{v} \in E$ is a solution of (4-3) if and only if $v = \text{tr}_{\mathbb{R}^N} \tilde{v}$ is a solution of (4-1), and $\tilde{v}_\varepsilon \in E$ is a solution of (4-4) if and only if $v_\varepsilon = \text{tr}_{\mathbb{R}^N} \tilde{v}_\varepsilon$ is a solution of (4-2). Since (4-1), (4-2) and (4-3), (4-4) are equivalent, respectively, we shall take some advantages in different cases.

Equation (4-1). In this subsection, we will derive some results for Equation (4-1), and the corresponding results are also true for (4-3).

Lemma 4.3. *There exist $\rho > 0$ and $\sigma > 0$ such that*

- (i) $J^{ab}(v) > \sigma$, for all $\|v\|_s = \rho$;
- (ii) $\lim_{t \rightarrow +\infty} J^{ab}(tv) = -\infty$ if $v \neq 0$.

Lemma 4.4. $\vartheta^{ab} = \inf_{v \in H^s(\mathbb{R}^N) \setminus \{0\}} \max_{t \geq 0} J^{ab}(tv) = \inf_{\gamma \in \Gamma^{ab}} \max_{t \in [0,1]} J^{ab}(\gamma(t)) > 0$,

where

$$\Gamma^{ab} := \{\gamma \in C([0, 1], H^s(\mathbb{R}^N)) : \gamma(0) = 0, J^{ab}(\gamma(1)) < 0\}.$$

Lemma 4.5. Let $v \in H^s(\mathbb{R}^N) \setminus \{0\}$. Then

$$\max_{t \geq 0} J^{ab}(tv) = \frac{p-1}{2p} (S^{ab}(v))^{p/(p-1)},$$

where

$$S^{ab}(v) := \frac{\int_{\mathbb{R}^N} (|(-\Delta)^{s/2} v|^2 + av^2) dx}{\left(\int_{\mathbb{R}^N} b^2(I_\theta * |v|^p)|v|^p dx\right)^{1/p}}.$$

From the proof of Theorem 1 in [Moroz and Van Schaftingen 2013], the following result is true.

Lemma 4.6. Both ζ^{ab} and ϑ^{ab} are attained in $H^s(\mathbb{R}^N)$, where

$$\begin{aligned} \zeta^{ab} &:= \inf\{S^{ab}(v) : v \in H^s(\mathbb{R}^N) \setminus \{0\}\} \\ &= \inf\left\{S^{ab}(v) : \int_{\mathbb{R}^N} b^2(I_\theta * |v|^p)|v|^p dx = 1, v \in H^s(\mathbb{R}^N)\right\} \end{aligned}$$

with $S^{ab}(v)$ defined as in Lemma 4.5.

In view of Theorem 1.1 in [d'Avenia et al. 2015], we have the following result.

Lemma 4.7. There exists a groundstate $v \in H^s(\mathbb{R}^N)$ for Equation (4-1), where v is positive, radially symmetric and decreasing. Moreover, if $s \in (0, \frac{1}{2}]$, then $v \in L^1(\mathbb{R}^N) \cap C^{0,\mu}(\mathbb{R}^N)$ for $\mu \in (0, 2s)$; if $s \in (\frac{1}{2}, 1)$, then $v \in L^1(\mathbb{R}^N) \cap C^{\bar{1},\mu}(\mathbb{R}^N)$ for $\mu \in (0, 2s - 1)$.

Lemma 4.8. Let $a_i > 0$ and $b_i > 0$ for $i = 1, 2$.

(i) If $\min\{a_2 - a_1, b_1 - b_2\} \geq 0$, then $\vartheta^{a_1 b_1} \leq \vartheta^{a_2 b_2}$.

(ii) If $\min\{a_2 - a_1, b_1 - b_2\} \geq 0$ and $\max\{a_2 - a_1, b_1 - b_2\} > 0$, then $\vartheta^{a_1 b_1} < \vartheta^{a_2 b_2}$.

Lemma 4.9. $\vartheta^{ab} = \left(\frac{a}{\tau_\infty}\right)^{\frac{\theta+2s}{2s(p-1)} - \frac{N-2s}{2s}} \cdot \left(\frac{k_\infty}{b}\right)^{\frac{2}{p-1}} \vartheta^\infty$.

Moreover, (1-5) holds if and only if $m\vartheta^{\tau k_v} < \vartheta^\infty$; (1-7) holds if and only if $m\vartheta^{\tau w^k} < \vartheta^\infty$.

Proof. Setting

$$u(x) := \left(\left(\frac{\tau_\infty}{a}\right)^{\frac{\theta+2s}{4s}} \frac{b}{k_\infty}\right)^{\frac{1}{p-1}} v\left(\left(\frac{\tau_\infty}{a}\right)^{\frac{1}{2s}} x\right),$$

then Equation (4-1) is equivalent to

$$(-\Delta)^s u + \tau_\infty u = k_\infty^2 (I_\theta * |u|^p) |u|^{p-2} u, \quad u \in H^s(\mathbb{R}^N).$$

One can check that $v \in \mathcal{N}^{ab}$ if and only if $u \in \mathcal{N}^\infty$, and for any $v \in \mathcal{N}^{ab}$,

$$J^{ab}(v) = \left(\frac{a}{\tau_\infty} \right)^{\frac{\theta+2s}{2s(p-1)} - \frac{N-2s}{2s}} \cdot \left(\frac{k_\infty}{b} \right)^{\frac{2}{p-1}} J^\infty(u). \quad \square$$

Lemma 4.10. *For any given $r \geq 2$ and $a > 0$, there exists a continuous function $v \in H^s(\mathbb{R}^N)$ satisfying*

$$(-\Delta)^s v + av = 0, \quad |x| \geq r$$

and

$$\frac{C_1}{|x|^{N+2s}} \leq v(x) \leq \frac{C_2}{|x|^{N+2s}}, \quad \text{for all } |x| \geq r,$$

where $C_2 > C_1 > 0$ dependent on s, a, r, N .

Proof. Consider the function $v := \mathcal{K}_a * \chi_{B_{r/2}}$, where \mathcal{K}_a is the fundamental solution of $(-\Delta)^s + a$ with $\widehat{\mathcal{K}_a}(x) = 1/(|\xi|^{2s} + a)$, and $\chi_{B_{r/2}}$ is the characteristic function of the ball $B_{r/2}$. For any $|x| \geq r$,

$$v(x) = \int_{|z| \leq r/2} \mathcal{K}_a(x-z) dz \quad \text{and} \quad (-\Delta)^s v + av = 0.$$

According to Felmer, Quaas and Tan [Felmer et al. 2012], \mathcal{K}_a is positive, radially symmetric and smooth in $\mathbb{R}^N \setminus \{0\}$, moreover,

$$\frac{\bar{C}_1}{|x|^{N+2s}} \leq \mathcal{K}_a(x) \leq \frac{\bar{C}_2}{|x|^{N+2s}}, \quad \text{for all } |x| \geq 1$$

with $\bar{C}_2 > \bar{C}_1 > 0$ dependent on s, c, N . Thus for all $|x| \geq r$,

$$\int_{|z| \leq r/2} \frac{\bar{C}_1}{|x-z|^{N+2s}} dz \leq \int_{|z| \leq r/2} \mathcal{K}_a(x-z) dz \leq \int_{|z| \leq r/2} \frac{\bar{C}_2}{|x-z|^{N+2s}} dz,$$

where

$$\begin{aligned} \int_{|z| \leq r/2} \frac{\bar{C}_1}{|x-z|^{N+2s}} dz &\geq \int_{|z| \leq r/2} \frac{\bar{C}_1}{(|x|+|z|)^{N+2s}} dz \\ &\geq \int_{|z| \leq r/2} \frac{\bar{C}_1}{(|x|+|x|/2)^{N+2s}} dz = \frac{C_1}{|x|^{N+2s}} \end{aligned}$$

and

$$\begin{aligned} \int_{|z| \leq r/2} \frac{\bar{C}_2}{|x-z|^{N+2s}} dz &\leq \int_{|z| \leq r/2} \frac{\bar{C}_2}{(|x|-|z|)^{N+2s}} dz \\ &\leq \int_{|z| \leq r/2} \frac{\bar{C}_2}{(|x|-|x|/2)^{N+2s}} dz = \frac{C_2}{|x|^{N+2s}}. \quad \square \end{aligned}$$

Equation (4-2). In this subsection, we will establish some results for Equation (4-2), and the corresponding results are also true for (4-4).

Lemma 4.11. *There exist $\rho > 0$ and $\sigma > 0$ both independent of ε, a, b and just dependent on N, θ, p, τ, k , such that*

$$(i) J_\varepsilon^{ab}(v) > \sigma, \quad \text{for all } \|v\|_s = \rho; \quad (ii) \lim_{t \rightarrow +\infty} J_\varepsilon^{ab}(tv) = -\infty \quad \text{if } v \neq 0.$$

Lemma 4.12. $\vartheta_\varepsilon^{ab} = \inf_{v \in H^s(\mathbb{R}^N) \setminus \{0\}} \max_{t \geq 0} J_\varepsilon^{ab}(tv) = \inf_{\gamma \in \Gamma_\varepsilon^{ab}} \max_{t \in [0,1]} J_\varepsilon^{ab}(\gamma(t)) > 0$,

where

$$\Gamma_\varepsilon^{ab} := \{\gamma \in C([0, 1], H^s(\mathbb{R}^N)) : \gamma(0) = 0, J_\varepsilon^{ab}(\gamma(1)) < 0\}.$$

Lemma 4.13. *If J_ε^∞ possesses a $(PS)_c$ sequence, then either $c = 0$ or $c \geq \vartheta_\varepsilon^\infty$. Moreover, $\vartheta_\varepsilon^\infty \geq \vartheta^\infty$.*

Proof. Let $\{v_n\} \subset H^s(\mathbb{R}^N)$ and $J_\varepsilon^\infty(v_n) \rightarrow c$, $(J_\varepsilon^\infty)'(v_n) \rightarrow 0$ in $H^{-s}(\mathbb{R}^N)$ as $n \rightarrow \infty$. Assume $c \neq 0$, we will prove $c \geq \vartheta_\varepsilon^\infty$.

Since $\{v_n\}$ is bounded in $H^s(\mathbb{R}^N)$, we may assume $v_n \rightharpoonup v$ in $H^s(\mathbb{R}^N)$ as $n \rightarrow \infty$ along a subsequence. Set $z_n := v_n - v$. By the Brézis–Lieb lemma, we have

$$(4-5) \quad \begin{aligned} \int_{\mathbb{R}^N} (|(-\Delta)^{s/2} v_n|^2 + V_\varepsilon^\infty(x) v_n^2) dx \\ = \int_{\mathbb{R}^N} (|(-\Delta)^{s/2} v|^2 + V_\varepsilon^\infty(x) v^2) dx \\ + \int_{\mathbb{R}^N} (|(-\Delta)^{s/2} z_n|^2 + V_\varepsilon^\infty(x) z_n^2) dx + o(1). \end{aligned}$$

By slight amendment in the proof of Lemma 3.5, we get that

$$(4-6) \quad \begin{aligned} \int_{\mathbb{R}^N} W_\varepsilon^\infty(x) [I_\theta * (W_\varepsilon^\infty |v_n|^p)] |v_n|^p dx \\ = \int_{\mathbb{R}^N} W_\varepsilon^\infty(x) [I_\theta * (W_\varepsilon^\infty |v|^p)] |v|^p dx \\ + \int_{\mathbb{R}^N} W_\varepsilon^\infty(x) [I_\theta * (W_\varepsilon^\infty |z_n|^p)] |z_n|^p dx + o(1), \end{aligned}$$

and for any $\varphi \in H^s(\mathbb{R}^N)$,

$$(4-7) \quad \begin{aligned} \int_{\mathbb{R}^N} W_\varepsilon^\infty(x) [I_\theta * (W_\varepsilon^\infty |v_n|^p)] |v_n|^{p-2} v_n \varphi dx \\ = \int_{\mathbb{R}^N} W_\varepsilon^\infty(x) [I_\theta * (W_\varepsilon^\infty |v|^p)] |v|^{p-2} v \varphi dx \\ + \int_{\mathbb{R}^N} W_\varepsilon^\infty(x) [I_\theta * (W_\varepsilon^\infty |z_n|^p)] |z_n|^{p-2} z_n \varphi dx + o(1) \|\varphi\|_s. \end{aligned}$$

As the proof of Lemma 3.6, we get that for any $\varphi \in H^s(\mathbb{R}^N)$,

$$\begin{aligned} \int_{\mathbb{R}^N} W_\varepsilon^\infty(x)[I_\theta * (W_\varepsilon^\infty |v_n|^p)]|v_n|^{p-2}v_n\varphi \, dx \\ \rightarrow \int_{\mathbb{R}^N} W_\varepsilon^\infty(x)[I_\theta * (W_\varepsilon^\infty |v|^p)]|v|^{p-2}v\varphi \, dx \quad \text{as } n \rightarrow \infty, \end{aligned}$$

which ensures that $(J_\varepsilon^\infty)'(v) = 0$. In virtue of (4-5), (4-6) and (4-7), we obtain that

$$\begin{aligned} J_\varepsilon^\infty(v_n) &= J_\varepsilon^\infty(v) + J_\varepsilon^\infty(z_n) + o(1), \\ (J_\varepsilon^\infty)'(v_n) &= (J_\varepsilon^\infty)'(v) + (J_\varepsilon^\infty)'(z_n) + o(1), \end{aligned}$$

which imply that

$$(4-8) \quad J_\varepsilon^\infty(z_n) \rightarrow c - J_\varepsilon^\infty(v), \quad \text{as } n \rightarrow \infty,$$

$$(4-9) \quad (J_\varepsilon^\infty)'(z_n) \rightarrow 0 \quad \text{in } H^{-s}(\mathbb{R}^N), \quad \text{as } n \rightarrow \infty.$$

If there exists $z_{n_k} \equiv 0$, that is $v_{n_k} \equiv v$, then $J_\varepsilon^\infty(v) = c \neq 0$ and $v \in \mathcal{N}_\varepsilon^\infty$. Thus $c \geq \vartheta_\varepsilon^\infty$.

If $z_n \neq 0$, for all $n \in \mathbb{N}$, then there exists $t_n > 0$ such that $t_n z_n \in \mathcal{N}_\varepsilon^\infty$. Hence

$$(4-10) \quad J_\varepsilon^\infty(t_n z_n) \geq \vartheta_\varepsilon^\infty.$$

It follows from $\langle (J_\varepsilon^\infty)'(t_n z_n), t_n z_n \rangle = 0$ and $\langle (J_\varepsilon^\infty)'(z_n), z_n \rangle = o(1)$ that

$$(4-11) \quad (1 - t_n^{2p-2}) \int_{\mathbb{R}^N} W_\varepsilon^\infty(x)[I_\theta * (W_\varepsilon^\infty |z_n|^p)]|z_n|^p \, dx = o(1).$$

Additionally,

$$\begin{aligned} \|z_n\|_s^2 &\leq C \int_{\mathbb{R}^N} (|(-\Delta)^{\frac{s}{2}} z_n|^2 + V_\varepsilon^\infty(x)z_n^2) \, dx \\ &= C \int_{\mathbb{R}^N} W_\varepsilon^\infty(x)[I_\theta * (W_\varepsilon^\infty |z_n|^p)]|z_n|^p \, dx + o(1). \end{aligned}$$

If $\int_{\mathbb{R}^N} (I_\theta * |z_n|^p)|z_n|^p \, dx \rightarrow 0$ as $n \rightarrow \infty$, then $\|z_n\|_s \rightarrow 0$ as $n \rightarrow \infty$. Thus $v_n \rightarrow v$ in $H^s(\mathbb{R}^N)$ as $n \rightarrow \infty$ and $c = J_\varepsilon^\infty(v) \geq \vartheta_\varepsilon^\infty$. If $\int_{\mathbb{R}^N} (I_\theta * |z_n|^p)|z_n|^p \, dx \geq \delta > 0$, then $t_n \rightarrow 1$ as $n \rightarrow \infty$ by (4-11). Hence $J_\varepsilon^\infty(t_n z_n) \rightarrow c - J_\varepsilon^\infty(v)$ as $n \rightarrow \infty$ by (4-8), which implies $c \geq J_\varepsilon^\infty(v) + \vartheta_\varepsilon^\infty \geq \vartheta_\varepsilon^\infty$ by (4-10).

Finally, it follows from $V_\varepsilon^\infty(x) \geq \tau_\infty$ and $W_\varepsilon^\infty(x) \leq k_\infty$ for any $x \in \mathbb{R}^N$ that

$$J_\varepsilon^\infty(u) \geq J^\infty(u), \quad \text{for all } u \in H^s(\mathbb{R}^N).$$

In virtue of Lemmas 4.4 and 4.12, $\vartheta_\varepsilon^\infty$ and ϑ^∞ are mountain pass levels of J_ε^∞ and J^∞ , respectively. Therefore, $\vartheta_\varepsilon^\infty \geq \vartheta^\infty$. \square

Remark 4.14. Similarly, if J_ε^{ab} has a $(PS)_c$ sequence, then either $c = 0$ or $c \geq \vartheta_\varepsilon^{ab}$.

Lemma 4.15. J_ε^{ab} satisfies the $(PS)_c$ condition for all $c < \vartheta_\varepsilon^\infty$.

Proof. Let $\{v_n\} \subset H^s(\mathbb{R}^N)$ and $J_\varepsilon^{ab}(v_n) \rightarrow c$, $(J_\varepsilon^{ab})'(v_n) \rightarrow 0$ in $H^{-s}(\mathbb{R}^N)$ as $n \rightarrow \infty$.

Since $\{v_n\}$ is bounded in $H^s(\mathbb{R}^N)$, we may assume $v_n \rightharpoonup v$ in $H^s(\mathbb{R}^N)$ as $n \rightarrow \infty$ along a subsequence. Then $(J_\varepsilon^{ab})'(v) = 0$ by Lemma 3.6.

Set $z_n := v_n - v$. Then $z_n \rightharpoonup 0$ in $H^s(\mathbb{R}^N)$ as $n \rightarrow \infty$ and

$$(4-12) \quad z_n \rightarrow 0 \quad \text{in } L_{\text{loc}}^q(\mathbb{R}^N) \quad \text{as } n \rightarrow \infty, \quad \text{for } q \in [2, 2_s^*).$$

Due to the classical Brézis–Lieb lemma and Lemma 3.5, we have

$$(4-13) \quad J_\varepsilon^{ab}(z_n) \rightarrow c - J_\varepsilon^{ab}(v), \quad (J_\varepsilon^{ab})'(z_n) \rightarrow 0 \quad \text{in } H^{-s}(\mathbb{R}^N) \quad \text{as } n \rightarrow \infty.$$

Now we prove

$$J_\varepsilon^\infty(z_n) \rightarrow c - J_\varepsilon^{ab}(v), \quad (J_\varepsilon^\infty)'(z_n) \rightarrow 0 \quad \text{in } H^{-s}(\mathbb{R}^N) \quad \text{as } n \rightarrow \infty.$$

By definition, we can get that

$$\lim_{|x| \rightarrow \infty} (V_\varepsilon^\infty(x) - V_\varepsilon^a(x)) = 0, \quad \lim_{|x| \rightarrow \infty} (W_\varepsilon^\infty(x) - W_\varepsilon^b(x)) = 0,$$

which imply that for all $\delta > 0$, there exists $R > 0$ such that

$$|V_\varepsilon^\infty(x) - V_\varepsilon^a(x)| \leq \delta, \quad |W_\varepsilon^\infty(x) - W_\varepsilon^b(x)| \leq \delta, \quad \text{for all } |x| > R.$$

Therefore, by Lemma 3.2 and the Hölder inequality, we get that

$$|J_\varepsilon^\infty(z_n) - J_\varepsilon^{ab}(z_n)| \leq \left(\frac{1}{2} |z_n|_2^2 + \frac{k}{p} |z_n|_{2Np/(N+\theta)}^{2p} \right) \delta + C(|z_n|_{L^2(B_R)}^2 + |z_n|_{L^{2Np/(N+\theta)}(B_R)}^p),$$

which jointly with (4-12) and (4-13), implies that

$$(4-14) \quad J_\varepsilon^\infty(z_n) \rightarrow c - J_\varepsilon^{ab}(v), \quad \text{as } n \rightarrow \infty.$$

For any $\varphi \in H^s(\mathbb{R}^N)$, by the Hölder inequality and Lemma 3.1, we have

$$\begin{aligned} & \left| \langle (J_\varepsilon^\infty)'(z_n) - (J_\varepsilon^{ab})'(z_n), \varphi \rangle \right| \\ & \leq \int_{\mathbb{R}^N} |V_\varepsilon^\infty(x) - V_\varepsilon^a(x)| |z_n| |\varphi| \, dx \\ & \quad + k \int_{\mathbb{R}^N} |W_\varepsilon^\infty(x) - W_\varepsilon^b(x)| (I_\theta * |z_n|^p) |z_n|^{p-1} |\varphi| \, dx \\ & \quad + k \int_{\mathbb{R}^N} |W_\varepsilon^\infty(x) - W_\varepsilon^b(x)| [I_\theta * (|z_n|^{p-1} |\varphi|)] |z_n|^p \, dx \\ & \leq C_1 \delta (|z_n|_2 + |z_n|_{2Np/(N+\theta)}^{2p-1}) \|\varphi\|_s + C_2 (|z_n|_{L^2(B_R)} + |z_n|_{L^{2Np/(N+\theta)}(B_R)}^{p-1}) \|\varphi\|_s, \end{aligned}$$

which together with (4-12) and (4-13), implies that

$$(4-15) \quad (J_\varepsilon^\infty)'(z_n) \rightarrow 0 \quad \text{in } H^{-s}(\mathbb{R}^N), \quad \text{as } n \rightarrow \infty.$$

It follows from (4-14) and (4-15) that $\{z_n\}$ is a $(\text{PS})_{c-J_\varepsilon^{ab}(v)}$ sequence of J_ε^∞ . By Lemma 4.13, either $c = J_\varepsilon^{ab}(v)$ or $c \geq J_\varepsilon^{ab}(v) + \vartheta_\varepsilon^\infty$. But the latter contradicts the assumption $c < \vartheta_\varepsilon^\infty$ in this lemma. Thus $c = J_\varepsilon^{ab}(v)$ and

$$(4-16) \quad J_\varepsilon^{ab}(v_n) \rightarrow J_\varepsilon^{ab}(v), \quad \text{as } n \rightarrow \infty.$$

Next, since $\langle (J_\varepsilon^{ab})'(v_n), v_n \rangle = o(1)$ and $\langle (J_\varepsilon^{ab})'(v), v \rangle = 0$, we have

$$\begin{aligned} J_\varepsilon^{ab}(v_n) &= \frac{p-1}{2p} \int_{\mathbb{R}^N} (|(-\Delta)^{s/2} v_n|^2 + V_\varepsilon^a(x) v_n^2) dx + o(1), \\ J_\varepsilon^{ab}(v) &= \frac{p-1}{2p} \int_{\mathbb{R}^N} (|(-\Delta)^{s/2} v|^2 + V_\varepsilon^a(x) v^2) dx, \end{aligned}$$

which together with (4-16) imply that $\|v_n\|_s \rightarrow \|v\|_s$ as $n \rightarrow \infty$. Therefore,

$$v_n \rightarrow v \quad \text{in } H^s(\mathbb{R}^N), \quad \text{as } n \rightarrow \infty. \quad \square$$

Lemma 4.16. $\limsup_{\varepsilon \rightarrow 0} \vartheta_\varepsilon^{ab} \leq \vartheta^{\alpha\beta}$, where $\alpha = V^a(0)$, $\beta = W^b(0)$.

Proof. Let $\bar{V}_\varepsilon(x) := V_\varepsilon^a(x) - \alpha$ and $\bar{W}_\varepsilon(x) := \beta - W_\varepsilon^b(x)$. Then

$$(4-17) \quad \bar{V}_\varepsilon(x) \rightarrow 0, \quad \bar{W}_\varepsilon(x) \rightarrow 0, \quad \text{a.e. on } \mathbb{R}^N, \quad \text{as } \varepsilon \rightarrow 0.$$

Meanwhile,

$$(4-18) \quad \begin{aligned} J_\varepsilon^{ab}(v) &= J^{\alpha\beta}(v) + \frac{1}{2} \int_{\mathbb{R}^N} \bar{V}_\varepsilon(x) v^2 dx + \frac{\beta}{p} \int_{\mathbb{R}^N} \bar{W}_\varepsilon(x) (I_\theta * |v|^p) |v|^p dx \\ &\quad - \frac{1}{2p} \int_{\mathbb{R}^N} \bar{W}_\varepsilon(x) [I_\theta * (\bar{W}_\varepsilon |v|^p)] |v|^p dx. \end{aligned}$$

By Lemma 4.6, there exists $e \in \mathcal{R}^{\alpha\beta}$. Let $t_\varepsilon > 0$ satisfy $t_\varepsilon e \in \mathcal{N}_\varepsilon^{ab}$, then

$$(4-19) \quad \max_{t \geq 0} J_\varepsilon^{ab}(te) = J_\varepsilon^{ab}(t_\varepsilon e) \geq \vartheta_\varepsilon^{ab}.$$

Since $J_\varepsilon^{ab}(te) \rightarrow -\infty$ as $t \rightarrow +\infty$, there exists $T_0 > 0$ such that

$$(4-20) \quad J_\varepsilon^{ab}(te) < 0, \quad \text{for all } t > T_0.$$

By (4-19) and (4-20), we get $t_\varepsilon \leq T_0$. Without loss of generality, we can assume $t_\varepsilon \rightarrow t_0$ as $\varepsilon \rightarrow 0$. Noting (4-17), (4-18) and (4-19), it follows from the Lebesgue dominated convergence theorem that

$$\begin{aligned} \vartheta_\varepsilon^{ab} &\leq J_\varepsilon^{ab}(t_\varepsilon e) = J^{\alpha\beta}(t_\varepsilon e) + \frac{t_\varepsilon^2}{2} \int_{\mathbb{R}^N} \bar{V}_\varepsilon(x) e^2 dx + \frac{\beta t_\varepsilon^{2p}}{p} \int_{\mathbb{R}^N} \bar{W}_\varepsilon(x) (I_\theta * |e|^p) |e|^p dx \\ &\quad - \frac{t_\varepsilon^{2p}}{2p} \int_{\mathbb{R}^N} \bar{W}_\varepsilon(x) [I_\theta * (\bar{W}_\varepsilon |e|^p)] |e|^p dx \\ &\rightarrow J^{\alpha\beta}(t_0 e) \leq J^{\alpha\beta}(e) = \vartheta^{\alpha\beta} \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

Thus $\limsup_{\varepsilon \rightarrow 0} \vartheta_\varepsilon^{ab} \leq \vartheta^{\alpha\beta}$. \square

Lemma 4.17. *Let $\tau \leq a \leq \tau_\infty$, $k_\infty \leq b \leq k$. If*

$$(4-21) \quad \left(\frac{V^a(0)}{\tau_\infty} \right)^{\frac{\theta+2s}{4s} - \frac{(N-2s)(p-1)}{4s}} < \frac{W^b(0)}{k_\infty},$$

then there exists $\varepsilon^{ab} > 0$ such that $\vartheta_\varepsilon^{ab}$ is attained at $v_\varepsilon^{ab} > 0$ for all $\varepsilon \leq \varepsilon^{ab}$.

Proof. Noting Lemma 4.9 and (4-21), we have $\vartheta^{\alpha\beta} < \vartheta^\infty$, where $\alpha = V^a(0)$ and $\beta = W^b(0)$. By Lemmas 4.16 and 4.13, there exists $\varepsilon^{ab} > 0$ such that

$$\vartheta_\varepsilon^{ab} < \vartheta^\infty \leq \vartheta_\varepsilon^\infty, \quad \text{for all } \varepsilon \leq \varepsilon^{ab}.$$

By Lemma 4.15, J_ε^{ab} satisfies the $(PS)_{\vartheta_\varepsilon^{ab}}$ condition for all $\varepsilon \leq \varepsilon^{ab}$, which together with Lemmas 4.11 and 4.12 implies that $\vartheta_\varepsilon^{ab}$ is attained at $v_\varepsilon^{ab} \in H^s(\mathbb{R}^N)$.

Next we shall prove $v_\varepsilon^{ab} > 0$. Let $\tilde{v}_\varepsilon^{ab} \in E$ be the s -harmonic extension of v_ε^{ab} , then $\tilde{v}_\varepsilon^{ab}$ is a groundstate of problem (4-4). Since $\tilde{J}_\varepsilon^{ab}(v_\varepsilon^{ab}) = \tilde{J}_\varepsilon^{ab}(|\tilde{v}_\varepsilon^{ab}|)$, we may assume $\tilde{v}_\varepsilon^{ab} \geq 0$. Then $v_\varepsilon^{ab}(x) = \tilde{v}_\varepsilon^{ab}(x, 0) \geq 0$. As in Theorem 3.2 in [d'Avenia et al. 2015], we have $v_\varepsilon^{ab} \in L^1(\mathbb{R}^N) \cap C^{0,\mu}(\mathbb{R}^N)$ with $\mu \in (0, 1)$, which implies $|v_\varepsilon^{ab}(x)| \rightarrow 0$ as $|x| \rightarrow \infty$. Hence $v_\varepsilon^{ab} \in L^\infty(\mathbb{R}^N)$. Then $I_\theta * |v_\varepsilon^{ab}|^p \in C^{0,\mu+\theta}(\mathbb{R}^N)$ for $\mu + \theta \in (0, 1)$ or $I_\theta * |v_\varepsilon^{ab}|^p \in C^{0,1}(\mathbb{R}^N)$ for $\mu + \theta \in [1, N + 1)$. Noting that $V, W \in C^{0,\lambda}(\mathbb{R}^N)$, we can use a similar proof as Theorem 1.4 in [Felmer et al. 2012] to obtain that $v_\varepsilon^{ab} \in C^{0,2s+\nu}(\mathbb{R}^N)$ for $2s + \nu \leq 1$ and $v_\varepsilon^{ab} \in C^{1,2s+\nu-1}(\mathbb{R}^N)$ for $2s + \nu > 1$, where $\nu = \min\{\lambda, \mu\}$. This regularity makes sure (2-2) holds. If $v_\varepsilon^{ab}(x_0) = 0$ for some $x_0 \in \mathbb{R}^N$, then (4-2) implies that $((-\Delta)^s v_\varepsilon^{ab})(x_0) = 0$. Due to (2-2), we get $v_\varepsilon^{ab} \equiv 0$, which is impossible. Hence $v_\varepsilon^{ab} > 0$. \square

In order to prove the existence of sign-changing solutions for Equation (1-1), we turn to problem (4-4) and define the Nehari nodal set

$$\tilde{\mathcal{M}}_\varepsilon^{ab} := \{ \tilde{v} \in E : \tilde{v}^\pm \neq 0, \langle (\tilde{J}_\varepsilon^{ab})'(\tilde{v}), \tilde{v}^\pm \rangle = 0 \}$$

and the least energy nodal value

$$\tilde{\zeta}_\varepsilon^{ab} := \inf_{\tilde{\mathcal{M}}_\varepsilon^{ab}} \tilde{J}_\varepsilon^{ab}.$$

Indeed, $\tilde{\mathcal{N}}_\varepsilon^{ab} \supset \tilde{\mathcal{M}}_\varepsilon^{ab} \neq \emptyset$ and $\tilde{\zeta}_\varepsilon^{ab} \geq \tilde{\vartheta}_\varepsilon^{ab} > 0$.

Lemma 4.18. *For any $\tilde{v} \in E$ with $\tilde{v}^\pm \neq 0$, there exists a unique pair of $(l_{\tilde{v}}, t_{\tilde{v}}) \in \mathbb{R}_+ \times \mathbb{R}_+$ such that*

$$l_{\tilde{v}} \tilde{v}^+ + t_{\tilde{v}} \tilde{v}^- \in \tilde{\mathcal{M}}_\varepsilon^{ab}, \quad \tilde{J}_\varepsilon^{ab}(l_{\tilde{v}} \tilde{v}^+ + t_{\tilde{v}} \tilde{v}^-) = \max_{l, t \geq 0} \tilde{J}_\varepsilon^{ab}(l \tilde{v}^+ + t \tilde{v}^-).$$

Proof. For any $\tilde{v} \in E$ with $\tilde{v}^\pm \neq 0$, consider the mapping

$$F(l, t) := \left(\langle (\tilde{J}_\varepsilon^{ab})'(l \tilde{v}^+ + t \tilde{v}^-), l \tilde{v}^+ \rangle, \langle (\tilde{J}_\varepsilon^{ab})'(l \tilde{v}^+ + t \tilde{v}^-), t \tilde{v}^- \rangle \right)$$

for $(l, t) \in \mathbb{R}_+ \times \mathbb{R}_+$. Since

$$\begin{aligned} \langle (\tilde{J}_\varepsilon^{ab})'(l\tilde{v}^+ + t\tilde{v}^-), l\tilde{v}^+ \rangle &= l^2 \left(\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{v}^+|^2 dx dy + \int_{\mathbb{R}^N} V_\varepsilon^a(x) |v^+|^2 dx \right) \\ &\quad - l^p t^p \int_{\mathbb{R}^N} W_\varepsilon^b(x) [I_\theta * (W_\varepsilon^b |v^-|^p)] |v^+|^p dx \\ &\quad - l^{2p} \int_{\mathbb{R}^N} W_\varepsilon^b(x) [I_\theta * (W_\varepsilon^b |v^+|^p)] |v^+|^p dx, \\ \langle (\tilde{J}_\varepsilon^{ab})'(l\tilde{v}^+ + t\tilde{v}^-), t\tilde{v}^- \rangle &= t^2 \left(\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{v}^-|^2 dx dy + \int_{\mathbb{R}^N} V_\varepsilon^a(x) |v^-|^2 dx \right) \\ &\quad - t^p l^p \int_{\mathbb{R}^N} W_\varepsilon^b(x) [I_\theta * (W_\varepsilon^b |v^+|^p)] |v^-|^p dx \\ &\quad - t^{2p} \int_{\mathbb{R}^N} W_\varepsilon^b(x) [I_\theta * (W_\varepsilon^b |v^-|^p)] |v^-|^p dx, \end{aligned}$$

where $v^\pm = \text{tr}_{\mathbb{R}^N} \tilde{v}^\pm = \tilde{v}^\pm(x, 0)$, there exist $R > r > 0$ such that for any $l, t \in [r, R]$,

$$\begin{aligned} \langle (\tilde{J}_\varepsilon^{ab})'(r\tilde{v}^+ + t\tilde{v}^-), r\tilde{v}^+ \rangle &> 0, & \langle (\tilde{J}_\varepsilon^{ab})'(R\tilde{v}^+ + t\tilde{v}^-), R\tilde{v}^+ \rangle &< 0, \\ \langle (\tilde{J}_\varepsilon^{ab})'(l\tilde{v}^+ + r\tilde{v}^-), r\tilde{v}^- \rangle &> 0, & \langle (\tilde{J}_\varepsilon^{ab})'(l\tilde{v}^+ + R\tilde{v}^-), R\tilde{v}^- \rangle &< 0. \end{aligned}$$

By Miranda's theorem [1940], there exists $(l_{\tilde{v}}, t_{\tilde{v}}) \in (r, R) \times (r, R)$ such that

$$\langle (\tilde{J}_\varepsilon^{ab})'(l_{\tilde{v}}\tilde{v}^+ + t_{\tilde{v}}\tilde{v}^-), l_{\tilde{v}}\tilde{v}^+ \rangle = 0 = \langle (\tilde{J}_\varepsilon^{ab})'(l_{\tilde{v}}\tilde{v}^+ + t_{\tilde{v}}\tilde{v}^-), t_{\tilde{v}}\tilde{v}^- \rangle.$$

That is $l_{\tilde{v}}\tilde{v}^+ + t_{\tilde{v}}\tilde{v}^- \in \tilde{\mathcal{M}}_\varepsilon^{ab}$.

Noting $p \geq 2$ and $\tilde{J}_\varepsilon^{ab}(l^{1/p}\tilde{v}^+ + t^{1/p}\tilde{v}^-)$ is strictly concave for $(l, t) \in \mathbb{R}_+ \times \mathbb{R}_+$, it follows from

$$\begin{aligned} \frac{\partial \tilde{J}_\varepsilon^{ab}(l^{1/p}\tilde{v}^+ + t^{1/p}\tilde{v}^-)}{\partial l} \Big|_{(l_{\tilde{v}}^p, t_{\tilde{v}}^p)} &= \frac{1}{pl_{\tilde{v}}^p} \langle (\tilde{J}_\varepsilon^{ab})'(l_{\tilde{v}}\tilde{v}^+ + t_{\tilde{v}}\tilde{v}^-), l_{\tilde{v}}\tilde{v}^+ \rangle = 0, \\ \frac{\partial \tilde{J}_\varepsilon^{ab}(l^{1/p}\tilde{v}^+ + t^{1/p}\tilde{v}^-)}{\partial t} \Big|_{(l_{\tilde{v}}^p, t_{\tilde{v}}^p)} &= \frac{1}{pt_{\tilde{v}}^p} \langle (\tilde{J}_\varepsilon^{ab})'(l_{\tilde{v}}\tilde{v}^+ + t_{\tilde{v}}\tilde{v}^-), t_{\tilde{v}}\tilde{v}^- \rangle = 0 \end{aligned}$$

that $(l_{\tilde{v}}^p, t_{\tilde{v}}^p)$ is the unique maximum point of $\tilde{J}_\varepsilon^{ab}(l^{1/p}\tilde{v}^+ + t^{1/p}\tilde{v}^-)$ for $(l, t) \in \mathbb{R}_+ \times \mathbb{R}_+$. Hence

$$\tilde{J}_\varepsilon^{ab}(l_{\tilde{v}}\tilde{v}^+ + t_{\tilde{v}}\tilde{v}^-) = \max_{l, t \geq 0} \tilde{J}_\varepsilon^{ab}(l^{1/p}\tilde{v}^+ + t^{1/p}\tilde{v}^-) = \max_{l, t \geq 0} \tilde{J}_\varepsilon^{ab}(l\tilde{v}^+ + t\tilde{v}^-). \quad \square$$

Now we define for any $\tilde{u}, \tilde{v} \in E$ and $u = \text{tr}_{\mathbb{R}^N} \tilde{u}$, $v = \text{tr}_{\mathbb{R}^N} \tilde{v}$,

$$f_\varepsilon^{ab}(\tilde{u}, \tilde{v}) := \begin{cases} \frac{\int_{\mathbb{R}^N} W_\varepsilon^b(x) [I_\theta * (W_\varepsilon^b(|u|^p + |v|^p))] |u|^p dx}{\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{u}|^2 dx dy + \int_{\mathbb{R}^N} V_\varepsilon^a(x) u^2 dx} & \text{if } \tilde{u} \neq 0, \\ 0 & \text{if } \tilde{u} = 0. \end{cases}$$

Obviously, $\tilde{v} \in \tilde{\mathcal{M}}_\varepsilon^{ab}$ if and only if $f_\varepsilon^{ab}(\tilde{v}^+, \tilde{v}^-) = f_\varepsilon^{ab}(\tilde{v}^-, \tilde{v}^+) = 1$.

Set

$$\tilde{U}_\varepsilon^{ab} := \left\{ \tilde{v} \in E : |f_\varepsilon^{ab}(\tilde{v}^+, \tilde{v}^-) - 1| < \frac{1}{2}, |f_\varepsilon^{ab}(\tilde{v}^-, \tilde{v}^+) - 1| < \frac{1}{2} \right\},$$

then $\tilde{\mathcal{M}}_\varepsilon^{ab} \subset \tilde{U}_\varepsilon^{ab} \neq \emptyset$. Moreover, one can verify that for $p \in (2, \frac{N+\theta}{N-2s})$,

$$(4-22) \quad \tilde{v}_n \in \tilde{U}_\varepsilon^{ab} \text{ and } \|\tilde{v}_n\|_E \leq C \implies \|\tilde{v}_n^\pm\|_E \geq \bar{C} > 0.$$

We continue to denote $P := \{\tilde{v} \in E : \tilde{v} \geq 0\}$ and let Q be the set of mappings q such that

$$(i) \quad q \in C(D, E), \quad \text{where } D := [0, 1] \times [0, 1],$$

and for all $t \in [0, 1]$,

$$(ii) \quad q(t, 0) = 0,$$

$$(iii) \quad q(0, t) \in P,$$

$$(iv) \quad q(1, t) \in -P,$$

$$(v) \quad (\tilde{J}_\varepsilon^{ab} \circ q)(t, 1) \leq 0, \quad f_\varepsilon^{ab}(q(t, 1), 0) \geq 2.$$

Then $Q \neq \emptyset$. By a standard proof (see Lemma 3.2 in [Cerami et al. 1986]), we have the following result.

Lemma 4.19.
$$\inf_{q \in Q} \sup_{\tilde{v} \in q(D)} \tilde{J}_\varepsilon^{ab}(\tilde{v}) = \inf_{\tilde{v} \in \tilde{\mathcal{M}}_\varepsilon^{ab}} \tilde{J}_\varepsilon^{ab}(\tilde{v}).$$

5. Proofs of multiplicity of semiclassical solutions

Setting $v(x) := w(\varepsilon x)$, the Equation (1-1) is equivalent to

$$(5-1) \quad (-\Delta)^s v + V(\varepsilon x)v = W(\varepsilon x)[I_\theta * (W(\varepsilon y)|v(y)|^p)]|v|^{p-2}v, \quad v \in H^s(\mathbb{R}^N).$$

If $v_\varepsilon(x)$ is a solution of Equation (5-1), then $w_\varepsilon(x) = v_\varepsilon(x/\varepsilon)$ is a solution of Equation (1-1).

By extension, we can reformulate Equation (5-1) as

$$(5-2) \quad \begin{cases} \operatorname{div}(y^{1-2s}\nabla\tilde{v}) = 0 & \text{in } \mathbb{R}_+^{N+1}, \\ \tilde{v}(x, 0) = v(x) & \text{on } \mathbb{R}^N, \\ \lim_{y \rightarrow 0^+} y^{1-2s}\partial_y\tilde{v} = V(\varepsilon x)v - W(\varepsilon x)[I_\theta * (W(\varepsilon y)|v(y)|^p)]|v|^{p-2}v & \text{in } \mathbb{R}_+^{N+1}. \end{cases}$$

Noting $V(\varepsilon x) = V_\varepsilon^\tau(x)$, $W(\varepsilon x) = W_\varepsilon^k(x)$, we find that Equation (5-1) and problem (5-2) are particular forms of Equation (4-2) and problem (4-4), respectively. For simplicity, we denote

$$\begin{aligned} J_\varepsilon &:= J_\varepsilon^{\tau k}, & \mathcal{N}_\varepsilon &:= \mathcal{N}_\varepsilon^{\tau k}, & \vartheta_\varepsilon &:= \vartheta_\varepsilon^{\tau k}, & \mathcal{R}_\varepsilon &:= \mathcal{R}_\varepsilon^{\tau k}, \\ \tilde{J}_\varepsilon &:= \tilde{J}_\varepsilon^{\tau k}, & \tilde{\mathcal{N}}_\varepsilon &:= \tilde{\mathcal{N}}_\varepsilon^{\tau k}, & \tilde{\vartheta}_\varepsilon &:= \tilde{\vartheta}_\varepsilon^{\tau k}, & \tilde{\mathcal{R}}_\varepsilon &:= \tilde{\mathcal{R}}_\varepsilon^{\tau k}, & V_\varepsilon &:= V_\varepsilon^\tau, \\ f_\varepsilon &:= f_\varepsilon^{\tau k}, & \tilde{\mathcal{M}}_\varepsilon &:= \tilde{\mathcal{M}}_\varepsilon^{\tau k}, & \tilde{\zeta}_\varepsilon &:= \tilde{\zeta}_\varepsilon^{\tau k}, & \tilde{U}_\varepsilon &:= \tilde{U}_\varepsilon^{\tau k}, & W_\varepsilon &:= W_\varepsilon^k. \end{aligned}$$

Proof of Theorem 1.1. Without loss of generality, we may assume $x_v = 0$. Then $V(0) = \tau$, $W(0) = k_v$. By (1-4) and (1-5), we get $m \geq 1$.

Step 1. We shall construct an m -dimensional subspace E_{r_m} of E such that

$$\sup_{\tilde{v} \in E_{r_m}} \tilde{J}_\varepsilon(\tilde{v}) < \vartheta^\infty, \quad \text{for all } r \geq r_m, \varepsilon \leq \varepsilon_m,$$

where r_m and ε_m are existing constants depending on m .

Choose $a = \tau$, $b = k_v$ in Equation (4-1), by Lemmas 4.6 and 4.7, there exist $v \in \mathcal{D}^{\tau k_v}$ and $v(x) = v(|x|) > 0$. Let $\tilde{v} \in E$ be the s -harmonic extension of v , then $\tilde{v} \in \tilde{\mathcal{H}}^{\tau k_v}$ and $\tilde{v}(x, 0) = v(x)$. Let $r > 0$, $\chi_r \in C_0^\infty(\mathbb{R}_+)$ satisfy $\chi_r(t) = 1$ for $t \leq r$ and $\chi_r(t) = 0$ for $t \geq r + 1$ with $|\chi_r'(t)| \leq 2$. Set

$$\tilde{v}_r(x, y) := \chi_r(|(x, y)|)\tilde{v}(x, y), \quad (x, y) \in \mathbb{R}_+^{N+1}.$$

Taking into account Lemmas 2.5, 2.2, 3.1 and 3.3, it follows from

$$\begin{aligned} \|\tilde{v}_r - \tilde{v}\|_E^2 &= \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla(\tilde{v}_r - \tilde{v})|^2 \, dx \, dy + \int_{\mathbb{R}^N} |v_r - v|^2 \, dx \\ &\leq C \left(\int_{\mathbb{R}_+^{N+1} \setminus B_r^+(0)} y^{1-2s} |\nabla \tilde{v}|^2 \, dx \, dy + \int_{B_{r+1}^+(0) \setminus B_r^+(0)} y^{1-2s} \tilde{v}^2 \, dx \, dy \right. \\ &\quad \left. + \int_{|x|>r} v^2 \, dx \right) \\ &\leq \bar{C} \left(\int_{\mathbb{R}_+^{N+1} \setminus B_r^+(0)} y^{1-2s} |\nabla \tilde{v}|^2 \, dx \, dy + \int_{|x|>r} v^2 \, dx \right) \rightarrow 0, \quad \text{as } r \rightarrow \infty \end{aligned}$$

that

$$\begin{aligned} \tilde{v}_r &\rightarrow \tilde{v} \quad \text{in } E, & \text{as } r \rightarrow \infty, \\ v_r &\rightarrow v \quad \text{in } H^s(\mathbb{R}^N), & \text{as } r \rightarrow \infty, \\ v_r &\rightarrow v \quad \text{in } L^{2Np/(N+\theta)}(\mathbb{R}^N), & \text{as } r \rightarrow \infty, \\ \int_{\mathbb{R}^N} (I_\theta * v_r^p) v_r^p \, dx &\rightarrow \int_{\mathbb{R}^N} (I_\theta * v^p) v^p \, dx, & \text{as } r \rightarrow \infty, \end{aligned}$$

where $v_r := \text{tr}_{\mathbb{R}^N} \tilde{v}_r$. Therefore, we obtain that as $r \rightarrow \infty$,

$$\begin{aligned} (5-3) \quad \max_{t \geq 0} \tilde{J}^{\tau k_v}(t \tilde{v}_r) &= \frac{p-1}{2p} \left(\frac{\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{v}_r|^2 \, dx \, dy + \int_{\mathbb{R}^N} \tau v_r^2 \, dx}{\left(\int_{\mathbb{R}^N} k_v^2 (I_\theta * v_r^p) v_r^p \, dx \right)^{1/p}} \right)^{\frac{p}{p-1}} \\ &\rightarrow \frac{p-1}{2p} \left(\frac{\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{v}|^2 \, dx \, dy + \int_{\mathbb{R}^N} \tau v^2 \, dx}{\left(\int_{\mathbb{R}^N} k_v^2 (I_\theta * v^p) v^p \, dx \right)^{1/p}} \right)^{\frac{p}{p-1}} \\ &= \max_{t \geq 0} \tilde{J}^{\tau k_v}(t \tilde{v}) = \tilde{J}^{\tau k_v}(\tilde{v}) = \tilde{\vartheta}^{\tau k_v} = \vartheta^{\tau k_v}. \end{aligned}$$

Additionally,

$$(5-4) \quad V_\varepsilon(x) \rightarrow V(0) = \tau, \quad W_\varepsilon(x) \rightarrow W(0) = k_v$$

as $\varepsilon \rightarrow 0$ uniformly on any bounded set of x . Thus (5-3) and (5-4) imply that

$$(5-5) \quad \begin{aligned} \max_{t \geq 0} \tilde{J}_\varepsilon(t\tilde{v}_r) &= \frac{p-1}{2p} \left(\frac{\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{v}_r|^2 dx dy + \int_{\mathbb{R}^N} V_\varepsilon(x) v_r^2 dx}{\left(\int_{\mathbb{R}^N} W_\varepsilon(x) [I_\theta * (W_\varepsilon v_r^p)] v_r^p dx \right)^{1/p}} \right)^{\frac{p}{p-1}} \\ &\rightarrow \frac{p-1}{2p} \left(\frac{\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{v}_r|^2 dx dy + \int_{\mathbb{R}^N} \tau v_r^2 dx}{\left(\int_{\mathbb{R}^N} k_v^2 (I_\theta * v_r^p) v_r^p dx \right)^{1/p}} \right)^{\frac{p}{p-1}} \\ &= \max_{t \geq 0} \tilde{J}^{\tau k_v}(t\tilde{v}_r) \rightarrow \vartheta^{\tau k_v} \quad \text{as } \varepsilon \rightarrow 0, r \rightarrow \infty, \text{ respectively.} \end{aligned}$$

Define

$$\tilde{\phi}_{rj}(x, y) := \tilde{v}_r(x_1 - 2j(r+1), x_2, \dots, x_N, y), \quad \text{for } j = 0, 1, \dots, m-1$$

and set

$$E_{rm} := \text{span}\{\tilde{\phi}_{rj}(x, y) : j = 0, 1, \dots, m-1\}.$$

One can check that $(\tilde{\phi}_{ri}, \tilde{\phi}_{rj})_E = 0$ if $i \neq j$. Thus $\dim E_{rm} = m$. Similarly as (5-5), for all $j = 1, 2, \dots, m-1$ and $\phi_{rj} = \text{tr}_{\mathbb{R}^N} \tilde{\phi}_{rj}$,

$$(5-6) \quad \begin{aligned} \max_{t \geq 0} \tilde{J}_\varepsilon(t\tilde{\phi}_{rj}) &= \frac{p-1}{2p} \left(\frac{\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{\phi}_{rj}|^2 dx dy + \int_{\mathbb{R}^N} V_\varepsilon(x) \phi_{rj}^2 dx}{\left(\int_{\mathbb{R}^N} W_\varepsilon(x) [I_\theta * (W_\varepsilon \phi_{rj}^p)] \phi_{rj}^p dx \right)^{1/p}} \right)^{\frac{p}{p-1}} \\ &\rightarrow \frac{p-1}{2p} \left(\frac{\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{\phi}_{rj}|^2 dx dy + \int_{\mathbb{R}^N} \tau \phi_{rj}^2 dx}{\left(\int_{\mathbb{R}^N} k_v^2 (I_\theta * \phi_{rj}^p) \phi_{rj}^p dx \right)^{1/p}} \right)^{\frac{p}{p-1}} \\ &= \max_{t \geq 0} \tilde{J}^{\tau k_v}(t\tilde{\phi}_{rj}) = \max_{t \geq 0} \tilde{J}^{\tau k_v}(t\tilde{v}_r) \\ &\rightarrow \vartheta^{\tau k_v} \quad \text{as } \varepsilon \rightarrow 0, r \rightarrow \infty, \text{ respectively.} \end{aligned}$$

Consequently, for all $\delta > 0$, there exist $r_\delta > 0$, $\varepsilon_\delta > 0$ such that

$$(5-6) \quad \max_{t \geq 0} \tilde{J}_\varepsilon(t\tilde{\phi}_{rj}) \leq \vartheta^{\tau k_v} + \delta, \quad \text{for all } r \geq r_\delta, \varepsilon \leq \varepsilon_\delta, j = 0, 1, \dots, m-1.$$

For any $\tilde{v} \in E_{rm}$, we may assume $\tilde{v} = t_0\tilde{\phi}_{r0} + t_1\tilde{\phi}_{r1} + \dots + t_{m-1}\tilde{\phi}_{r(m-1)}$, where $t_j \in \mathbb{R}$ for $j = 0, 1, \dots, m-1$. Then

$$[I_\theta * (W_\varepsilon |\tilde{v}|^p)] |\tilde{v}|^p \geq \sum_{j=0}^{m-1} [I_\theta * (W_\varepsilon |t_j \tilde{\phi}_{rj}|^p)] |t_j \tilde{\phi}_{rj}|^p.$$

Hence by (5-6),

$$\begin{aligned} \tilde{J}_\varepsilon(\tilde{v}) &\leq \tilde{J}_\varepsilon(t_0\tilde{\phi}_{r0}) + \tilde{J}_\varepsilon(t_1\tilde{\phi}_{r1}) + \cdots + \tilde{J}_\varepsilon(t_{m-1}\tilde{\phi}_{r(m-1)}) \\ &\leq m(\vartheta^{\tau k_v} + \delta), \quad \text{for all } r \geq r_\delta, \varepsilon \leq \varepsilon_\delta, \end{aligned}$$

which implies that

$$\sup_{\tilde{v} \in E_{r_m}} \tilde{J}_\varepsilon(\tilde{v}) \leq m(\vartheta^{\tau k_v} + \delta), \quad \text{for all } r \geq r_\delta, \varepsilon \leq \varepsilon_\delta.$$

Noting Lemma 4.9, we can choose $0 < \delta < \frac{\vartheta^\infty}{m} - \vartheta^{\tau k_v}$, then there exist $r_m > 0$, $\varepsilon_m > 0$ such that

$$(5-7) \quad \sup_{\tilde{v} \in E_{r_m}} \tilde{J}_\varepsilon(\tilde{v}) < \vartheta^\infty, \quad \text{for all } r \geq r_m, \varepsilon \leq \varepsilon_m.$$

Step 2. We will define constants c_1, c_2, \dots, c_m and verify that they are critical values of \tilde{J}_ε .

Consider the symmetric group $\mathbb{Z}_2 = \{\text{id}, -\text{id}\}$ and denote

$$\Sigma := \{A \subset E : A \text{ is closed and } A = -A\}.$$

For any $A \in \Sigma$, the Krasnoselskii genus of A is defined by

$$\text{gen}(A) := \inf\{n : \text{there exists } g \in C(A, \mathbb{R}^n \setminus \{0\}) \text{ and } g \text{ is odd}\}.$$

Set

$$\mathcal{H} := \{h \in C(E, E) : h \text{ is an odd homeomorphism}\}$$

and for any $A \in \Sigma$, define

$$i(A) := \min_{h \in \mathcal{H}} \text{gen}(h(A) \cap \partial B_\rho),$$

where $\rho > 0$ is a constant defined in Lemma 4.11. Thus $i(A)$ is a version of Benci pseudoindex of A . Let

$$c_j := \inf_{i(A) \geq j} \sup_{\tilde{v} \in A} \tilde{J}_\varepsilon(\tilde{v}), \quad j = 1, 2, \dots, m.$$

Clearly, $c_1 \leq c_2 \leq \cdots \leq c_m$. Next we will show $c_1 \geq \sigma$ and $c_m \leq \sup_{\tilde{v} \in E_{r_m}} \tilde{J}_\varepsilon(\tilde{v})$, where $\sigma > 0$ was defined in Lemma 4.11.

For any $A \in \Sigma$ and $i(A) \geq 1$, we have $\text{gen}(A \cap \partial B_\rho) \geq 1$, which implies $A \cap \partial B_\rho \neq \emptyset$. It follows from Lemma 4.11(i) that

$$\tilde{J}_\varepsilon(\tilde{v}) = J_\varepsilon(v) > \sigma, \quad \text{for all } \|\tilde{v}\|_E = \|v\|_S = \rho,$$

where $v = \text{tr}_{\mathbb{R}^N} \tilde{v}$. Hence $\sup_{\tilde{v} \in A} \tilde{J}_\varepsilon(\tilde{v}) > \sigma$ and $c_1 \geq \sigma$.

Taking into account that the Krasnoselskii genus satisfies the dimension property (see [Benci 1982]), we get

$$\text{gen}(h(E_{r_m}) \cap \partial B_\rho) = \dim E_{r_m} = m, \quad \text{for all } h \in \mathcal{H},$$

which implies $i(E_{r_m}) = m$. Thus $c_m \leq \sup_{\tilde{v} \in E_{r_m}} \tilde{J}_\varepsilon(\tilde{v})$.

Noting (5-7) and Lemma 4.13, we get that for any $r \geq r_m$, $\varepsilon \leq \varepsilon_m$,

$$(5-8) \quad \sigma \leq c_1 \leq c_2 \leq \cdots \leq c_m \leq \sup_{\tilde{v} \in E_{r_m}} \tilde{J}_\varepsilon(\tilde{v}) < \vartheta^\infty \leq \vartheta_\varepsilon^\infty.$$

Now we prove c_1, c_2, \dots, c_m are critical values of \tilde{J}_ε by applying the Theorem 1.4 in [Benci 1982]. Set

$$c_0 := \sigma, \quad c_\infty := \sup_{\tilde{v} \in E_{r_m}} \tilde{J}_\varepsilon(\tilde{v}),$$

$$(\tilde{J}_\varepsilon)^c := \{\tilde{v} \in E : \tilde{J}_\varepsilon(\tilde{v}) \leq c\}, \quad K_c := \{\tilde{v} \in E : \tilde{J}_\varepsilon(\tilde{v}) = c, (\tilde{J}_\varepsilon)'(\tilde{v}) = 0\}.$$

Since \tilde{J}_ε is an even functional,

$$(5-9) \quad (\tilde{J}_\varepsilon)^c \in \Sigma, \quad K_c \in \Sigma, \quad \text{for all } c \in [c_0, c_\infty].$$

By (5-8) and Lemma 4.15, \tilde{J}_ε satisfies the $(\text{PS})_c$ condition for any $c \in [c_0, c_\infty]$, which implies that

$$(5-10) \quad K_c \text{ is compact in } E, \quad \text{for all } c \in [c_0, c_\infty].$$

For any $c \in [c_0, c_\infty]$, $d > 0$ and $(K_c)_d := \{\tilde{v} \in E : \text{dist}(\tilde{v}, K_c) < d\}$, choose $\delta = \frac{d}{4}$, then by the contradiction method we can get that there exists $\tilde{\varepsilon} > 0$ such that

$$(5-11) \quad \|(\tilde{J}_\varepsilon)'(\tilde{v})\| \geq \frac{8\tilde{\varepsilon}}{\delta}, \quad \text{for all } \tilde{v} \in \tilde{J}_\varepsilon^{-1}([c - 2\tilde{\varepsilon}, c + 2\tilde{\varepsilon}]) \setminus \overline{(K_c)_d/2}.$$

Choose $S := E \setminus (K_c)_d$ and by the deformation lemma (see Lemma 2.3 in [Willem 1996]), there exists $\tilde{\eta} \in C([0, 1] \times E, E)$ such that $\tilde{\eta}(t, \cdot)$ is an odd homeomorphism of E for any $t \in [0, 1]$ and $\tilde{\eta}(1, (\tilde{J}_\varepsilon)^{c+\tilde{\varepsilon}} \cap S) \subset (\tilde{J}_\varepsilon)^{c-\tilde{\varepsilon}}$. Set $\eta(\cdot) := \tilde{\eta}(1, \cdot)$; then η is an odd homeomorphism of E and

$$(5-12) \quad \eta((\tilde{J}_\varepsilon)^{c+\tilde{\varepsilon}} \setminus (K_c)_d) \subset (\tilde{J}_\varepsilon)^{c-\tilde{\varepsilon}}.$$

For any $A \in \Sigma$ and $A \subset (\tilde{J}_\varepsilon)^{c_0}$, it follows from Lemma 4.11(i) that $A \cap \partial B_\rho = \emptyset$. Thus $\text{gen}(A \cap \partial B_\rho) = 0$ and

$$(5-13) \quad i(A) = \min_{h \in \mathcal{H}} \text{gen}(h(A) \cap \partial B_\rho) = 0.$$

Choose $\tilde{A} = E_{r_m}$, then

$$(5-14) \quad \tilde{A} \subset (\tilde{J}_\varepsilon)^{c_\infty} \quad \text{and} \quad i(\tilde{A}) = i(E_{r_m}) = m \geq 1.$$

Uniting (5-9), (5-10), (5-12), (5-13) and (5-14), we obtain that c_1, c_2, \dots, c_m are critical values of \tilde{J}_ε , and $\text{gen}(K_c) \geq r + 1$ if $c := c_k = c_{k+1} = \dots = c_{k+r}$ with $k \geq 1$ and $k + r \leq m$. Since \tilde{J}_ε is even, we conclude that \tilde{J}_ε has at least m pairs of critical points which are also solutions of Equation (5-2). Consequently, J_ε has at least m pairs of critical points being solutions of Equation (5-1).

Step 3. We will prove Equation (5-1) has at least one positive and one negative groundstate.

Noting (1-5) and $m \geq 1$, we get that

$$\left(\frac{\tau}{\tau_\infty} \right)^{\frac{\theta+2s}{4s} - \frac{(N-2s)(p-1)}{4s}} < \frac{k_v}{k_\infty},$$

which together with $V(0) = \tau$, $W(0) = k_v$ implies that

$$\left(\frac{V^\tau(0)}{\tau_\infty} \right)^{\frac{\theta+2s}{4s} - \frac{(N-2s)(p-1)}{4s}} < \frac{W^k(0)}{k_\infty}.$$

By Lemma 4.17, there exists $\varepsilon^{\tau k} > 0$ such that ϑ_ε is attained at $v_\varepsilon > 0$ for all $\varepsilon \leq \varepsilon^{\tau k}$. Therefore, v_ε and $-v_\varepsilon$ are positive and negative groundstates of Equation (5-1).

Step 4. We will prove Equation (5-1) has at least one pair of sign-changing solutions when $m \geq 2$ and $2 < p < \frac{N+\theta}{N-2s}$.

Firstly, let $e \in \mathcal{R}^{\tau k_v}$ with $e > 0$ and $\tilde{e} \in E$ be the s -harmonic extension of e . Then $\tilde{e} \in \tilde{\mathcal{R}}^{\tau k_v}$ with $\tilde{e} > 0$. Let $\chi_r \in C_0^\infty(\mathbb{R}_+)$ satisfy $\chi_r(t) = 1$ for $t \leq r$ and $\chi_r(t) = 0$ for $t \geq r + 1$ with $|\chi_r'(t)| \leq 2$. Choose $r > 0$ and $X_r := (x_r, y_r) \in \mathbb{R}_+^{N+1}$ satisfying $|X_r|$ large and

$$\text{dist}(B_{r+1}^+(0), B_{r+1}^+(X_r)) > 0.$$

Define for $(x, y) \in \mathbb{R}_+^{N+1}$,

$$\begin{aligned} \tilde{e}_r^+(x, y) &:= \chi_r(|(x, y)|) \tilde{e}(x, y) \geq 0, \\ \tilde{e}_r^-(x, y) &:= -\chi_r(|(x - x_r, y - y_r)|) \tilde{e}(x, y) \leq 0. \end{aligned}$$

Then $\text{supp } \tilde{e}_r^+ \cap \text{supp } \tilde{e}_r^- = \emptyset$ and

$$(5-15) \quad \tilde{e}_r^+ \rightarrow e, \quad \tilde{e}_r^- \rightarrow -e \quad \text{in } E, \quad \text{as } r \rightarrow \infty.$$

By Lemma 4.18, there exists $(l_r, t_r) \in \mathbb{R}_+ \times \mathbb{R}_+$ such that

$$(5-16) \quad l_r \tilde{e}_r^+ + t_r \tilde{e}_r^- \in \tilde{\mathcal{M}}_\varepsilon.$$

Secondly, since $\langle (\tilde{J}_\varepsilon)'(l_r \tilde{e}_r^+ + t_r \tilde{e}_r^-), l_r \tilde{e}_r^+ \rangle = \langle (\tilde{J}_\varepsilon)'(l_r \tilde{e}_r^+ + t_r \tilde{e}_r^-), t_r \tilde{e}_r^- \rangle = 0$, we have

$$(5-17) \quad \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{e}_r^+|^2 dx dy + \int_{\mathbb{R}^N} V_\varepsilon(x) |e_r^+|^2 dx \\ = l_r^{p-2} t_r^p \int_{\mathbb{R}^N} W_\varepsilon(x) [I_\theta * (W_\varepsilon |e_r^-|^p)] |e_r^+|^p dx \\ + l_r^{2p-2} \int_{\mathbb{R}^N} W_\varepsilon(x) [I_\theta * (W_\varepsilon |e_r^+|^p)] |e_r^+|^p dx$$

and

$$(5-18) \quad \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{e}_r^-|^2 dx dy + \int_{\mathbb{R}^N} V_\varepsilon(x) |e_r^-|^2 dx \\ = t_r^{p-2} l_r^p \int_{\mathbb{R}^N} W_\varepsilon(x) [I_\theta * (W_\varepsilon |e_r^+|^p)] |e_r^-|^p dx \\ + t_r^{2p-2} \int_{\mathbb{R}^N} W_\varepsilon(x) [I_\theta * (W_\varepsilon |e_r^-|^p)] |e_r^-|^p dx,$$

where $e_r^\pm = \text{tr}_{\mathbb{R}^N} \tilde{e}_r^\pm$, which imply that l_r and t_r are bounded for $r \in (0, \infty)$. Without loss of generality, we may assume that

$$l_r \rightarrow l_0, \quad t_r \rightarrow t_0, \quad \text{as } r \rightarrow \infty.$$

Letting $\varepsilon \rightarrow 0$ in (5-17) and (5-18), we get that

$$(5-19) \quad \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{e}_r^+|^2 dx dy + \int_{\mathbb{R}^N} \tau |e_r^+|^2 dx \\ = l_r^{p-2} t_r^p \int_{\mathbb{R}^N} k_v^2(I_\theta * |e_r^-|^p) |e_r^+|^p dx + l_r^{2p-2} \int_{\mathbb{R}^N} k_v^2(I_\theta * |e_r^+|^p) |e_r^+|^p dx$$

and

$$(5-20) \quad \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{e}_r^-|^2 dx dy + \int_{\mathbb{R}^N} \tau |e_r^-|^2 dx \\ = t_r^{p-2} l_r^p \int_{\mathbb{R}^N} k_v^2(I_\theta * |e_r^+|^p) |e_r^-|^p dx + t_r^{2p-2} \int_{\mathbb{R}^N} k_v^2(I_\theta * |e_r^-|^p) |e_r^-|^p dx.$$

Letting $r \rightarrow \infty$ in (5-19) and (5-20), by (5-15) we obtain

$$\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{e}|^2 dx dy + \int_{\mathbb{R}^N} \tau e^2 dx = (l_0^{p-2} t_0^p + l_0^{2p-2}) \int_{\mathbb{R}^N} k_v^2(I_\theta * e^p) e^p dx, \\ \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{e}|^2 dx dy + \int_{\mathbb{R}^N} \tau e^2 dx = (t_0^{p-2} l_0^p + t_0^{2p-2}) \int_{\mathbb{R}^N} k_v^2(I_\theta * e^p) e^p dx.$$

Since $\tilde{e} \in \mathcal{N}^{\tau k_v}$, we have $l_0 = t_0 = 2^{1/(2-2p)}$.

Thirdly, by (5-16) we have

$$\begin{aligned}
(5-21) \quad \tilde{\zeta}_\varepsilon &\leq \tilde{J}_\varepsilon(l_r \tilde{e}_r^+ + t_r \tilde{e}_r^-) \\
&= \frac{p-1}{2p} \left(\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla(l_r \tilde{e}_r^+ + t_r \tilde{e}_r^-)|^2 dx dy + \int_{\mathbb{R}^N} V_\varepsilon(x) |l_r e_r^+ + t_r e_r^-|^2 dx \right) \\
&\rightarrow \frac{p-1}{2p} \left(\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla(l_r \tilde{e}_r^+ + t_r \tilde{e}_r^-)|^2 dx dy + \int_{\mathbb{R}^N} \tau |l_r e_r^+ + t_r e_r^-|^2 dx \right) \\
&\rightarrow \frac{p-1}{2p} (l_0^2 + t_0^2) \left(\int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{e}|^2 dx dy + \int_{\mathbb{R}^N} \tau e^2 dx \right) \\
&= (l_0^2 + t_0^2) \tilde{\vartheta}^{\tau k_v} < 2\vartheta^{\tau k_v}, \quad \text{as } \varepsilon \rightarrow 0 \quad \text{and} \quad r \rightarrow \infty.
\end{aligned}$$

Noting $m \geq 2$, due to Lemma 4.9 and Lemma 4.13, we get

$$(5-22) \quad 2\vartheta^{\tau k_v} < \vartheta^\infty \leq \vartheta_\varepsilon^\infty.$$

In view of (5-21) and (5-22), we conclude that $\tilde{\zeta}_\varepsilon < \vartheta_\varepsilon^\infty$ for ε small enough, which implies that \tilde{J}_ε satisfies the $(PS)_{\tilde{\zeta}_\varepsilon}$ condition for ε small enough.

Fourthly, consider a minimizing sequence $\{\tilde{v}_n\} \subset \tilde{\mathcal{M}}_\varepsilon$ of $\tilde{\zeta}_\varepsilon$ and denote by $\{q_n\} \subset Q$ a mapping sequence satisfying

$$q_n(D) \subset \{\alpha \tilde{v}_n^+ + \beta \tilde{v}_n^- : \alpha, \beta \in \mathbb{R}_+ \cup \{0\}\}.$$

Then

$$(5-23) \quad \lim_{n \rightarrow \infty} \max_{\tilde{v} \in q_n(D)} \tilde{J}_\varepsilon(\tilde{v}) = \lim_{n \rightarrow \infty} \tilde{J}_\varepsilon(\tilde{v}_n) = \tilde{\zeta}_\varepsilon.$$

By Lemma 4.19, (5-23) and a similar statement in [Cerami et al. 1986], there exists $\{\tilde{u}_n\} \subset E$ such that

$$(5-24) \quad \lim_{n \rightarrow \infty} \text{dist}(\tilde{u}_n, q_n(D)) = 0, \quad \lim_{n \rightarrow \infty} \tilde{J}_\varepsilon(\tilde{u}_n) = \tilde{\zeta}_\varepsilon, \quad \lim_{n \rightarrow \infty} (\tilde{J}_\varepsilon)'(\tilde{u}_n) = 0.$$

Next we will show $\tilde{u}_n \in \tilde{U}_\varepsilon$ for n large enough. Indeed we only need to prove $\tilde{u}_n^\pm \neq 0$ for n large enough, which jointly with $\lim_{n \rightarrow \infty} \langle (\tilde{J}_\varepsilon)'(\tilde{u}_n), \tilde{u}_n^\pm \rangle = 0$ ensures that

$$\lim_{n \rightarrow \infty} f_\varepsilon(\tilde{u}_n^+, \tilde{u}_n^-) = \lim_{n \rightarrow \infty} f_\varepsilon(\tilde{u}_n^-, \tilde{u}_n^+) = 1.$$

By (5-24), there exists $\tilde{z}_n = \alpha_n \tilde{v}_n^+ + \beta_n \tilde{v}_n^- \in q_n(D)$ such that

$$\lim_{n \rightarrow \infty} \|\tilde{u}_n - \tilde{z}_n\|_E = 0.$$

Thus it suffices to show $\alpha_n \tilde{v}_n^+ \neq 0$ and $\beta_n \tilde{v}_n^- \neq 0$ for n large enough. It follows from $\tilde{v}_n \in \tilde{\mathcal{M}}_\varepsilon$ and (5-24) that $0 < C_1 \leq \|\tilde{v}_n^\pm\|_E \leq C_2$. We turn to show $\alpha_n \not\rightarrow 0$ and $\beta_n \not\rightarrow 0$ as $n \rightarrow \infty$. For a contradiction, assume that $\lim_{n \rightarrow \infty} \alpha_n = 0$, then

$$0 < \tilde{\zeta}_\varepsilon = \lim_{n \rightarrow \infty} \tilde{J}_\varepsilon(\tilde{v}_n) = \lim_{n \rightarrow \infty} \tilde{J}_\varepsilon(\beta_n \tilde{v}_n^-).$$

Hence $\{\beta_n\}$ is bounded and $\lim_{n \rightarrow \infty} \beta_n \neq 0$. Noting $p > 2$, we have

$$\begin{aligned} \tilde{\zeta}_\varepsilon &= \lim_{n \rightarrow \infty} \tilde{J}_\varepsilon(\tilde{v}_n) = \lim_{n \rightarrow \infty} \max_{\alpha, \beta \geq 0} \tilde{J}_\varepsilon(\alpha \tilde{v}_n^+ + \beta \tilde{v}_n^-) \geq \lim_{n \rightarrow \infty} \max_{\alpha \geq 0} \tilde{J}_\varepsilon(\alpha \tilde{v}_n^+ + \beta_n \tilde{v}_n^-) \\ &= \lim_{n \rightarrow \infty} \max_{\alpha \geq 0} \left[\frac{\alpha^2}{2} \int_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla \tilde{v}_n^+|^2 \, dx \, dy + \frac{\alpha^2}{2} \int_{\mathbb{R}^N} V_\varepsilon(x) |v_n^+|^2 \, dx \right. \\ &\quad - \frac{\alpha^{2p}}{2p} \int_{\mathbb{R}^N} W_\varepsilon(x) [I_\theta * (W_\varepsilon |v_n^+|^p)] |v_n^+|^p \, dx \\ &\quad \left. - \frac{\alpha^p \beta_n^p}{p} \int_{\mathbb{R}^N} W_\varepsilon(x) [I_\theta * (W_\varepsilon |v_n^+|^p)] |v_n^-|^p \, dx \right] + \lim_{n \rightarrow \infty} \tilde{J}_\varepsilon(\beta_n \tilde{v}_n^-) \\ &\geq \max_{\alpha \geq 0} (C_3 \alpha^2 - C_4 \alpha^p - C_5 \alpha^{2p}) + \tilde{\zeta}_\varepsilon > \tilde{\zeta}_\varepsilon, \end{aligned}$$

which is impossible.

Finally, noting that both of the above sequences $\{\tilde{v}_n\}$ and $\{\tilde{u}_n\}$ are related to ε , going if necessary to a subsequence, for ε small enough we can assume

$$\tilde{u}_n \rightarrow \tilde{u}_\varepsilon \quad \text{in } E, \quad \text{as } n \rightarrow \infty,$$

which together with (4-22) and (5-24) implies that

$$\tilde{u}_\varepsilon \in \tilde{\mathcal{M}}_\varepsilon, \quad \tilde{J}_\varepsilon(\tilde{u}_\varepsilon) = \tilde{\zeta}_\varepsilon, \quad (\tilde{J}_\varepsilon)'(\tilde{u}_\varepsilon) = 0.$$

That is $\pm \tilde{u}_\varepsilon$ are a pair of sign-changing solutions for problem (5-2). Hence we conclude that $\pm u_\varepsilon = \pm \operatorname{tr}_{\mathbb{R}^N} \tilde{u}_\varepsilon$ are a pair of sign-changing solutions for Equation (5-1).

This completes the proof. \square

Proof of Theorem 1.2. We can assume without loss of generality that $x_w = 0$. Then $V(0) = \tau_w$, $W(0) = k$. By (1-6) and (1-7), we get $m \geq 1$. Taking $a = \tau_w$, $b = k$ in Equation (4-1), there exists $v \in \mathcal{R}^{\tau_w k}$. The following arguments are similar to the proof of Theorem 1.1, thus the details are omitted. \square

6. Proofs of convergence, concentration and decay estimate of groundstates

In this section, we shall prove the convergence, concentration and decay estimate of the positive groundstates for Equation (1-1). Namely we offer the proof of Theorem 1.3.

Proof of Theorem 1.3. We shall deal with the case (A1)(i), the other case can be handled similarly. Without loss of generality, we assume $x_v = 0$. Then $V(0) = \tau$, $W(0) = k_v$. Obviously, (A1)(i) implies that (1-4) holds. It follows from Theorem 1.1 that Equation (1-1) has a positive groundstate $w_\varepsilon(x)$. Moreover, Equation (5-1) has a positive groundstate $v_\varepsilon(x) = w_\varepsilon(\varepsilon x)$, and problem (5-2) has a positive groundstate $\tilde{v}_\varepsilon(x, y)$ which is also the s -harmonic extension of v_ε .

Step 1. We shall prove the convergence of v_ε as $\varepsilon \rightarrow 0$ up to a sequence after translations.

Let $\varepsilon_j \rightarrow 0$ (as $j \rightarrow \infty$), $v_j := v_{\varepsilon_j} \in \mathcal{R}_{\varepsilon_j}$ with $v_j > 0$. Since

$$\vartheta_{\varepsilon_j} = J_{\varepsilon_j}(v_j) = \frac{p-1}{2p} \int_{\mathbb{R}^N} (|(-\Delta)^{s/2} v_j|^2 + V_{\varepsilon_j}(x) v_j^2) dx \geq C \|v_j\|_s^2,$$

it follows from Lemma 4.16 that $\{v_j\}$ is bounded in $H^s(\mathbb{R}^N)$. By $v_j \in \mathcal{N}_{\varepsilon_j}$ and Lemma 3.2, we have

$$(6-1) \quad \|v_j\|_s^2 \leq C \int_{\mathbb{R}^N} (I_\theta * v_j^p) v_j^p dx \leq \bar{C} \|v_j\|_s^{2p}.$$

Suppose $\lim_{j \rightarrow \infty} \sup_{y \in \mathbb{R}^N} \int_{B_1(y)} v_j^2 dx = 0$, then by Lemmas 2.3, 3.1 and 3.3, we have

$$\begin{aligned} v_j &\rightarrow 0 \quad \text{in } L^{2Np/(N+\theta)}(\mathbb{R}^N), \quad \text{as } j \rightarrow \infty, \\ \int_{\mathbb{R}^N} (I_\theta * v_j^p) v_j^p dx &\rightarrow 0, \quad \text{as } j \rightarrow \infty, \end{aligned}$$

which contradicts (6-1). Therefore, there exist $\delta > 0$ and $y'_j \in \mathbb{R}^N$ such that

$$(6-2) \quad \int_{B_1(y'_j)} v_j^2 dx \geq \delta.$$

Set $u_j(x) := v_j(x + y'_j)$, $\widehat{V}_{\varepsilon_j}(x) := V_{\varepsilon_j}(x + y'_j)$, $\widehat{W}_{\varepsilon_j}(x) := W_{\varepsilon_j}(x + y'_j)$. Then u_j solves

$$(6-3) \quad (-\Delta)^s u_j + \widehat{V}_{\varepsilon_j}(x) u_j = \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} u_j^p)] u_j^{p-1}, \quad u_j > 0$$

with least energy

$$\begin{aligned} (6-4) \quad \hat{\vartheta}_{\varepsilon_j} = \hat{J}_{\varepsilon_j}(u_j) &:= \frac{1}{2} \int_{\mathbb{R}^N} (|(-\Delta)^{s/2} u_j|^2 + \widehat{V}_{\varepsilon_j}(x) u_j^2) dx \\ &\quad - \frac{1}{2p} \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} u_j^p)] u_j^p dx \\ &= \frac{p-1}{2p} \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} u_j^p)] u_j^p dx. \end{aligned}$$

Moreover, $[I_\theta * (W_{\varepsilon_j} v_j^p)](x + y'_j) = [I_\theta * (\widehat{W}_{\varepsilon_j} u_j^p)](x)$ for any $x \in \mathbb{R}^N$, which ensures that

$$(6-5) \quad \hat{\vartheta}_{\varepsilon_j} = \hat{J}_{\varepsilon_j}(u_j) = J_{\varepsilon_j}(v_j) = \vartheta_{\varepsilon_j}.$$

In view of the boundedness of $\{u_j\}$, we can assume without loss of generality that

$$(6-6) \quad u_j \rightharpoonup u \quad \text{in } H^s(\mathbb{R}^N), \quad \text{as } j \rightarrow \infty,$$

$$(6-7) \quad u_j \rightarrow u \quad \text{in } L_{\text{loc}}^q(\mathbb{R}^N), \quad \text{as } j \rightarrow \infty, \quad \text{for } q \in [2, 2_s^*),$$

which together with (6-2) implies $u \neq 0$.

Since V and W are bounded, going if necessary to a subsequence, we assume

$$(6-8) \quad V_{\varepsilon_j}(y'_j) \rightarrow V_0 \quad \text{and} \quad W_{\varepsilon_j}(y'_j) \rightarrow W_0, \quad \text{as } j \rightarrow \infty.$$

By the boundedness of $\nabla V : |\nabla V(x)| \leq M$, for all $x \in \mathbb{R}^N$, we get that for any given $r > 0$,

$$|\widehat{V}_{\varepsilon_j}(x) - V_{\varepsilon_j}(y'_j)| = \left| \int_0^1 \nabla V(\varepsilon_j y'_j + t\varepsilon_j x) \varepsilon_j x \, dt \right| \leq \varepsilon_j M r, \quad \text{for all } x \in B_r(0).$$

Thus $\widehat{V}_{\varepsilon_j}(x) \rightarrow V_0$ as $j \rightarrow \infty$ uniformly on any bounded set of x . Similarly, $\widehat{W}_{\varepsilon_j}(x) \rightarrow W_0$ as $j \rightarrow \infty$ uniformly on any bounded set of x . As in the proof of Lemma 4.16, we have

$$(6-9) \quad \limsup_{j \rightarrow \infty} \widehat{\vartheta}_{\varepsilon_j} \leq \vartheta^{V_0 W_0}.$$

By (6-3), (6-6), (6-7), and Lemmas 3.1 and 3.3, we obtain that for any $\varphi \in C_0^\infty(\mathbb{R}^N)$,

$$\begin{aligned} 0 &= \lim_{j \rightarrow \infty} \int_{\mathbb{R}^N} [(-\Delta)^{s/2} u_j (-\Delta)^{s/2} \varphi + \widehat{V}_{\varepsilon_j}(x) u_j \varphi - \widehat{W}_{\varepsilon_j}(x) (I_\theta * (\widehat{W}_{\varepsilon_j} u_j^p)) u_j^{p-1} \varphi] \, dx \\ &= \int_{\mathbb{R}^N} [(-\Delta)^{s/2} u (-\Delta)^{s/2} \varphi + V_0 u \varphi - W_0 (I_\theta * (W_0 u^p)) u^{p-1} \varphi] \, dx, \end{aligned}$$

which implies that u solves

$$(6-10) \quad (-\Delta)^s u + V_0 u = W_0^2 (I_\theta * u^p) u^{p-1}, \quad u > 0$$

with energy

$$\begin{aligned} (6-11) \quad J^{V_0 W_0}(u) &:= \frac{1}{2} \int_{\mathbb{R}^N} (|(-\Delta)^{s/2} u|^2 + V_0 u^2) \, dx - \frac{1}{2p} \int_{\mathbb{R}^N} W_0^2 (I_\theta * u^p) u^p \, dx \\ &= \frac{p-1}{2p} \int_{\mathbb{R}^N} W_0^2 (I_\theta * u^p) u^p \, dx \geq \vartheta^{V_0 W_0}. \end{aligned}$$

According to Fatou's Lemma,

$$(6-12) \quad \int_{\mathbb{R}^N} W_0^2 (I_\theta * u^p) u^p \, dx \leq \liminf_{j \rightarrow \infty} \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} u_j^p)] u_j^p \, dx.$$

Uniting (6-4), (6-9), (6-11) and (6-12), we obtain

$$\vartheta^{V_0 W_0} \leq J^{V_0 W_0}(u) \leq \liminf_{j \rightarrow \infty} \widehat{J}_{\varepsilon_j}(u_j) \leq \limsup_{j \rightarrow \infty} \widehat{\vartheta}_{\varepsilon_j} \leq \vartheta^{V_0 W_0}.$$

Therefore,

$$(6-13) \quad \lim_{j \rightarrow \infty} \hat{\vartheta}_{\varepsilon_j} = \vartheta^{V_0 W_0} = J^{V_0 W_0}(u).$$

Let \tilde{u}_j and \tilde{u} be the s -harmonic extension of u_j and u , respectively. Then as $j \rightarrow \infty$,

$$\tilde{u}_j \rightharpoonup \tilde{u}, \quad \text{in } E \text{ and in } X^s(\mathbb{R}_+^{N+1}).$$

Let $\eta \in C_0^\infty(\mathbb{R}_+)$ satisfy $\eta(t) = 1$ for $t \leq 1$ and $\eta(t) = 0$ for $t \geq 2$ with $|\eta'(t)| \leq 2$. Define

$$\tilde{\mu}_j(x, y) := \eta\left(\frac{|(x, y)|}{j}\right) \tilde{u}(x, y) \quad \text{and} \quad \tilde{z}_j(x, y) := \tilde{u}_j(x, y) - \tilde{\mu}_j(x, y)$$

for $(x, y) \in \mathbb{R}_+^{N+1}$ and denote $\mu_j := \text{tr}_{\mathbb{R}^N} \tilde{\mu}_j$, $z_j := \text{tr}_{\mathbb{R}^N} \tilde{z}_j$. Then as $j \rightarrow \infty$,

$$\begin{aligned} \tilde{\mu}_j &\rightarrow \tilde{u} \quad \text{in } E \text{ and in } X^s(\mathbb{R}_+^{N+1}), \\ \mu_j &\rightarrow u \quad \text{in } H^s(\mathbb{R}^N), \\ \mu_j &\rightarrow u \quad \text{in } L^q(\mathbb{R}^N), \quad \text{for } q \in [2, 2_s^*], \\ \mu_j &\rightarrow u \quad \text{a.e. on } \mathbb{R}^N, \end{aligned}$$

and

$$\begin{aligned} \tilde{z}_j &\rightarrow 0 \quad \text{in } E, \\ z_j &\rightarrow 0 \quad \text{in } H^s(\mathbb{R}^N), \\ z_j &\rightarrow 0 \quad \text{in } L_{\text{loc}}^q(\mathbb{R}^N), \quad \text{for } q \in [2, 2_s^*], \\ z_j &\rightarrow 0 \quad \text{a.e. on } \mathbb{R}^N. \end{aligned}$$

Next we will show $\tilde{J}_{\varepsilon_j}(\tilde{z}_j) \rightarrow 0$ and $\langle (\tilde{J}_{\varepsilon_j})'(\tilde{z}_j), \tilde{z}_j \rangle \rightarrow 0$ as $j \rightarrow \infty$, where

$$\begin{aligned} \tilde{J}_{\varepsilon_j}(\tilde{z}_j) &:= \frac{1}{2} \int_{\mathbb{R}_+^{N+1}} y^{1-2s} |\nabla \tilde{z}_j|^2 \, dx \, dy + \frac{1}{2} \int_{\mathbb{R}^N} \widehat{V}_{\varepsilon_j}(x) |z_j|^2 \, dx \\ &\quad - \frac{1}{2p} \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} |z_j|^p)] |z_j|^p \, dx. \end{aligned}$$

Firstly, we claim

$$(6-14) \quad \|\tilde{z}_j\|_X^2 = \|\tilde{u}_j\|_X^2 - \|\tilde{\mu}_j\|_X^2 + o(1).$$

Indeed, $\|\tilde{u}_j\|_X^2 - \|\tilde{\mu}_j\|_X^2 - \|\tilde{z}_j\|_X^2 = 2(\tilde{u}_j, \tilde{\mu}_j)_X - 2\|\tilde{\mu}_j\|_X^2$, where $\|\tilde{\mu}_j\|_X \rightarrow \|\tilde{u}\|_X$ and

$$|(\tilde{u}_j, \tilde{\mu}_j)_X - (\tilde{u}, \tilde{u})_X| \leq |(\tilde{u}_j, \tilde{\mu}_j - \tilde{u})_X| + |(\tilde{u}_j - \tilde{u}, \tilde{u})_X| \rightarrow 0, \quad \text{as } j \rightarrow \infty.$$

Secondly, we claim

$$(6-15) \quad \int_{\mathbb{R}^N} \widehat{V}_{\varepsilon_j}(x) |z_j|^2 dx = \int_{\mathbb{R}^N} \widehat{V}_{\varepsilon_j}(x) |u_j|^2 dx - \int_{\mathbb{R}^N} \widehat{V}_{\varepsilon_j}(x) |\mu_j|^2 dx + o(1).$$

Indeed, $\{z_j\}$ is bounded in $H^s(\mathbb{R}^N)$ and in $L^q(\mathbb{R}^N)$ for any $q \in [2, 2_s^*]$. Moreover, for all $\delta > 0$, there exists $c(\delta) > 0$ such that

$$||u_j|^q - |z_j|^q| = ||z_j + \mu_j|^q - |z_j|^q| \leq \delta |z_j|^q + c(\delta) |\mu_j|^q$$

and

$$g_j^\delta := (||u_j|^q - |z_j|^q - |\mu_j|^q| - \delta |z_j|^q)^+ \leq (1 + c(\delta)) |\mu_j|^q \leq (1 + c(\delta)) |u|^q.$$

By the Lebesgue dominated convergence theorem, $\int_{\mathbb{R}^N} g_j^\delta dx \rightarrow 0$ as $j \rightarrow \infty$. Thus

$$(6-16) \quad \int_{\mathbb{R}^N} ||u_j|^q - |z_j|^q - |\mu_j|^q| dx \leq \int_{\mathbb{R}^N} g_j^\delta dx + \delta \int_{\mathbb{R}^N} |z_j|^q dx = o(1).$$

Choose $q = 2$ in (6-16), we find (6-15) holds.

Thirdly, we claim

$$(6-17) \quad \begin{aligned} \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} |z_j|^p)] |z_j|^p dx \\ = \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} |u_j|^p)] |u_j|^p dx \\ - \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} |\mu_j|^p)] |\mu_j|^p dx + o(1). \end{aligned}$$

Indeed, taking into account that $\{z_j\}$ is bounded in $L^{2Np/(N+\theta)}(\mathbb{R}^N)$, it follows from (6-16) with $q = p$ that

$$\begin{aligned} |u_j|^p - |\mu_j|^p - |z_j|^p &\rightarrow 0 \quad \text{in } L^{2N/(N+\theta)}(\mathbb{R}^N), \quad \text{as } j \rightarrow \infty, \\ I_\theta * (|u_j|^p - |\mu_j|^p - |z_j|^p) &\rightarrow 0 \quad \text{in } L^{2N/(N-\theta)}(\mathbb{R}^N), \quad \text{as } j \rightarrow \infty, \end{aligned}$$

which together with

$$\begin{aligned} |z_j|^p &\rightarrow 0 && \text{in } L^{2N/(N+\theta)}(\mathbb{R}^N), \quad \text{as } j \rightarrow \infty, \\ \mu_j^p &\rightarrow u^p && \text{in } L^{2N/(N+\theta)}(\mathbb{R}^N), \quad \text{as } j \rightarrow \infty, \\ I_\theta * \mu_j^p &\rightarrow I_\theta * u^p && \text{in } L^{2N/(N-\theta)}(\mathbb{R}^N), \quad \text{as } j \rightarrow \infty \end{aligned}$$

and Lemma 3.3 imply that

$$\begin{aligned}
 & \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) \left[(I_\theta * (\widehat{W}_{\varepsilon_j} |u_j|^p)) |u_j|^p - (I_\theta * (\widehat{W}_{\varepsilon_j} |\mu_j|^p)) |\mu_j|^p \right. \\
 & \qquad \qquad \qquad \left. - (I_\theta * (\widehat{W}_{\varepsilon_j} |z_j|^p)) |z_j|^p \right] dx \\
 &= \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} (|u_j|^p - |\mu_j|^p - |z_j|^p))] (|u_j|^p - |\mu_j|^p - |z_j|^p) dx \\
 & \quad + 2 \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} (|u_j|^p - |\mu_j|^p - |z_j|^p))] |\mu_j|^p dx \\
 & \quad \quad + 2 \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} (|u_j|^p - |\mu_j|^p - |z_j|^p))] |z_j|^p dx \\
 & \quad \quad \quad + 2 \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} |\mu_j|^p)] |z_j|^p dx \\
 & \rightarrow 0 \quad \text{as } j \rightarrow \infty.
 \end{aligned}$$

Fourthly, it follows from the Lebesgue dominated convergence theorem that

$$(6-18) \quad \int_{\mathbb{R}^N} \widehat{V}_{\varepsilon_j}(x) \mu_j^2 dx = \int_{\mathbb{R}^N} V_0 u^2 dx + o(1),$$

$$(6-19) \quad \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} \mu_j^p)] \mu_j^p dx = \int_{\mathbb{R}^N} W_0^2 (I_\theta * u^p) u^p dx + o(1).$$

Moreover,

$$(6-20) \quad |(-\Delta)^{s/2} \mu_j|_2^2 = |(-\Delta)^{s/2} u|_2^2 + o(1).$$

Uniting (6-14), (6-15), (6-17), (6-18), (6-19), (6-20), (6-13), (6-3) and (6-10), we obtain

$$(6-21) \quad \tilde{J}_{\varepsilon_j}(\tilde{z}_j) = \tilde{J}_{\varepsilon_j}(\tilde{u}_j) - \tilde{J}_{\varepsilon_j}(\tilde{\mu}_j) + o(1) = \hat{v}_{\varepsilon_j} - J^{V_0 W_0}(u) + o(1) = o(1)$$

and

$$\begin{aligned}
 (6-22) \quad \langle (\tilde{J}_{\varepsilon_j})'(\tilde{z}_j), \tilde{z}_j \rangle &= \langle (\tilde{J}_{\varepsilon_j})'(\tilde{u}_j), \tilde{u}_j \rangle - \langle (\tilde{J}_{\varepsilon_j})'(\tilde{\mu}_j), \tilde{\mu}_j \rangle + o(1) \\
 &= \langle (\hat{J}_{\varepsilon_j})'(u_j), u_j \rangle - \langle (J^{V_0 W_0})'(u), u \rangle + o(1) = o(1).
 \end{aligned}$$

Finally, by (6-21) and (6-22), we obtain that

$$\begin{aligned}
 o(1) &= \tilde{J}_{\varepsilon_j}(\tilde{z}_j) - \frac{1}{2p} \langle (\tilde{J}_{\varepsilon_j})'(\tilde{z}_j), \tilde{z}_j \rangle \\
 &= \frac{p-1}{2p} \left(\int_{\mathbb{R}^{N+1}} y^{1-2s} |\nabla \tilde{z}_j|^2 dx dy + \int_{\mathbb{R}^N} \widehat{V}_{\varepsilon_j}(x) |z_j|^2 dx \right) \geq C \|\tilde{z}_j\|_E^2,
 \end{aligned}$$

which implies $\tilde{z}_j \rightarrow 0$ in E and $z_j \rightarrow 0$ in $H^s(\mathbb{R}^N)$ as $j \rightarrow \infty$. Thus

$$\|u_j - u\|_s \leq \|z_j\|_s + \|\mu_j - u\|_s \rightarrow 0, \quad \text{as } j \rightarrow \infty.$$

That is

$$u_j \rightarrow u \quad \text{in } H^s(\mathbb{R}^N), \quad \text{as } j \rightarrow \infty.$$

Step 2. We claim $u_j(x) \rightarrow 0$ as $|x| \rightarrow \infty$ uniformly in $j \in \mathbb{N}$.

We can use the contradiction method to obtain that

$$(6-23) \quad \sup_j \int_{|x| \geq r} u_j^2 dx \rightarrow 0, \quad \text{as } r \rightarrow \infty.$$

Moreover, noting that $(-\Delta)^s u_j - \widehat{W}_{\varepsilon_j}(x)[I_\theta * (\widehat{W}_{\varepsilon_j} u_j^p)] u_j^{p-1} \leq 0$ on \mathbb{R}^N and

$$[I_\theta * (\widehat{W}_{\varepsilon_j} u_j^p)] u_j^{p-2} \in L_{\text{loc}}^q(\mathbb{R}^N) \quad \text{for some } q > \frac{N}{2s},$$

we deduce from Proposition 2.6 in [Jin et al. 2014] that for any compact set K ,

$$\max_K u_j(x) \leq C \left(\int_K u_j^2 dx \right)^{1/2},$$

which together with (6-23) ensures the claim true.

Step 3. We claim $\{\xi_j y'_j\}_j$ is bounded on \mathbb{R}^N .

Otherwise, there exists $|\xi_j y'_j| \rightarrow \infty$ as $j \rightarrow \infty$ along a subsequence. Hence $V_0 \geq \tau_\infty > \tau$ and $W_0 \leq k_\infty \leq k_v$, which together with Lemma 4.8, imply that $\vartheta^{V_0 W_0} > \vartheta^{\tau k_v}$. However, by (6-5), (6-13) and Lemma 4.16,

$$\vartheta^{V_0 W_0} = \lim_{j \rightarrow \infty} \widehat{\vartheta}_{\varepsilon_j} = \lim_{j \rightarrow \infty} \vartheta_{\varepsilon_j} \leq \limsup_{j \rightarrow \infty} \vartheta_{\varepsilon_j} \leq \vartheta^{\tau k_v}.$$

That is a contradiction.

Therefore, without loss of generality we may assume

$$(6-24) \quad \varepsilon_j y'_j \rightarrow x_0, \quad \text{as } j \rightarrow \infty.$$

Noting (6-8), we have

$$(6-25) \quad V_0 = V(x_0) \quad \text{and} \quad W_0 = W(x_0).$$

By (6-10), we find that u is a groundstate of Equation (1-8).

Step 4. We claim $\{\varepsilon y_\varepsilon\}_\varepsilon$ is bounded, where $y_\varepsilon \in \mathbb{R}^N$ is a maximum point of v_ε .

For a contradiction, assume that there is $\varepsilon_j \rightarrow 0$ with $|\varepsilon_j y_j| \rightarrow \infty$, where $y_j := y_{\varepsilon_j}$ is a maximum point of $v_j := v_{\varepsilon_j}$. Repeating Steps 1, 2, 3, one can get that there exists $y'_j \in \mathbb{R}^N$ such that

$$(6-26) \quad u_j = v_j(\cdot + y'_j) \rightarrow u \neq 0 \quad \text{in } H^s(\mathbb{R}^N), \quad \text{as } j \rightarrow \infty,$$

$$u_j(x) \rightarrow 0, \quad \text{as } |x| \rightarrow \infty, \quad \text{uniformly in } j \in \mathbb{N},$$

$$\{\varepsilon_j y'_j\}_j \text{ is bounded on } \mathbb{R}^N.$$

Thus $|\varepsilon_j y_j - \varepsilon_j y'_j| \geq |\varepsilon_j y_j| - |\varepsilon_j y'_j| \rightarrow \infty$, as $j \rightarrow \infty$, which implies that $|y_j - y'_j| \rightarrow \infty$ as $j \rightarrow \infty$. Then $\max_{\mathbb{R}^N} v_j = v_j(y_j) = u_j(y_j - y'_j) \rightarrow 0$ as $j \rightarrow \infty$. Hence $\max_{\mathbb{R}^N} u_j \rightarrow 0$ as $j \rightarrow \infty$. Noting $u_j > 0$, one has $u_j(x) \rightarrow 0$ as $j \rightarrow \infty$ uniformly in $x \in \mathbb{R}^N$, which contradicts (6-26).

Step 5. We claim $\lim_{\varepsilon \rightarrow 0} \text{dist}(\varepsilon y_\varepsilon, \mathcal{A}_v) = 0$.

By Step 4, there exists $\varepsilon_j \rightarrow 0$ with

$$(6-27) \quad \varepsilon_j y_j \rightarrow y_0, \quad \text{as } j \rightarrow \infty,$$

where $y_j := y_{\varepsilon_j}$ is a maximum point of $v_j := v_{\varepsilon_j}$. It is sufficient to verify that $y_0 \in \mathcal{A}_v$.

By Step 1 and Step 3, there exists $y'_j \in \mathbb{R}^N$ satisfying $u_j(x) = v_j(x + y'_j)$ and (6-24). By Step 2, we can assume $u_j(x'_j) = \max_{\mathbb{R}^N} u_j$ and $\{x'_j\}_j$ is bounded on \mathbb{R}^N . Thus $y_j = x'_j + y'_j$ and $\varepsilon_j y_j - \varepsilon_j y'_j = \varepsilon_j x'_j \rightarrow 0$ as $j \rightarrow \infty$, which together with (6-24), (6-25) and (6-27), imply that

$$(6-28) \quad y_0 = x_0, \quad V(y_0) = V_0, \quad W(y_0) = W_0.$$

For a contradiction, assume that $y_0 \notin \mathcal{A}_v$. Then either $V(y_0) = \tau$, $W(y_0) < k_v$ or $V(y_0) > \tau$, $W(y_0) \leq k_v$. By Lemma 4.8,

$$(6-29) \quad \vartheta^{V(y_0)W(y_0)} > \vartheta^{\tau k_v}.$$

Uniting (6-5), (6-13), (6-28), (6-29) and Lemma 4.16, we obtain

$$\lim_{j \rightarrow \infty} \vartheta_{\varepsilon_j} = \lim_{j \rightarrow \infty} \widehat{\vartheta}_{\varepsilon_j} = \vartheta^{V_0 W_0} = \vartheta^{V(y_0)W(y_0)} > \vartheta^{\tau k_v} \geq \limsup_{j \rightarrow \infty} \vartheta_{\varepsilon_j}.$$

That is impossible.

In particular, if $\mathcal{V} \cap \mathcal{W} \neq \emptyset$, then $x_0 \in \mathcal{A}_v = \mathcal{V} \cap \mathcal{W}$. Hence

$$\lim_{\varepsilon \rightarrow 0} \text{dist}(\varepsilon y_\varepsilon, \mathcal{V} \cap \mathcal{W}) = 0 \quad \text{and} \quad V(x_0) = \tau, \quad W(x_0) = k,$$

which jointly with Equation (1-8) imply that u is a groundstate of Equation (1-9).

Step 6. For $p \in (2, \frac{N+\theta}{N-2s})$, we claim there exist $C_2 > C_1 > 0$ and $r > 2$ large enough such that for all small $\varepsilon > 0$,

$$\frac{C_1}{|x|^{N+2s}} \leq u_\varepsilon(x) \leq \frac{C_2}{|x|^{N+2s}}, \quad \text{for all } |x| \geq r.$$

We verify its correctness for any sequence.

Now choose $a = \sup_{\mathbb{R}^N} \widehat{V}_{\varepsilon_j}(x)$ and $r = 2$ in Lemma 4.10, then there is a continuous function $v_1 \in H^s(\mathbb{R}^N)$ such that

$$(6-30) \quad (-\Delta)^s v_1 + a v_1 = 0 \quad \text{and} \quad v_1(x) \geq \frac{\bar{C}_1}{|x|^{N+2s}}, \quad \text{for all } |x| \geq 2,$$

where \bar{C}_1 depends on a, N, s . By the continuity of u_{ε_j} and v_1 , there is $M_1 > 0$ such that

$$u_{\varepsilon_j} - M_1 v_1 \geq 0, \quad \text{for all } |x| \leq 2.$$

Additionally, by (6-3) and (6-30), we have

$$((-\Delta)^s + a)(u_{\varepsilon_j} - M_1 v_1) \geq \int_{\mathbb{R}^N} \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} u_{\varepsilon_j}^p)] u_{\varepsilon_j}^{p-1} dx \geq 0, \quad \text{for all } |x| \geq 2.$$

Hence $u_{\varepsilon_j} - M_1 v_1 \geq 0$ for any $|x| \geq 2$, which together with (6-30) implies that

$$u_{\varepsilon_j}(x) \geq \frac{C_1}{|x|^{N+2s}}, \quad \text{for all } |x| \geq 2, \quad \text{where } C_1 = M_1 \bar{C}_1.$$

Next, by Step 2 and (6-3), choose $0 < a < \tau$ and $r > 2$ sufficiently large such that

$$(6-31) \quad (-\Delta)^s u_{\varepsilon_j} + a u_{\varepsilon_j} \\ = \widehat{W}_{\varepsilon_j}(x) [I_\theta * (\widehat{W}_{\varepsilon_j} u_{\varepsilon_j}^p)] u_{\varepsilon_j}^{p-1} + (a - \widehat{V}_{\varepsilon_j}(x)) u_{\varepsilon_j} \leq 0, \quad \text{for all } |x| \geq r.$$

By Lemma 4.10, there is a continuous function $v_2 \in H^s(\mathbb{R}^N)$ such that

$$(6-32) \quad (-\Delta)^s v_2 + a v_2 = 0 \quad \text{and} \quad v_2(x) \leq \frac{\bar{C}_2}{|x|^{N+2s}}, \quad \text{for all } |x| \geq r,$$

where \bar{C}_2 depends on a, N, s, r . Moreover, there is $M_2 > 0$ such that

$$u_{\varepsilon_j} - M_2 v_2 \leq 0, \quad \text{for all } |x| \leq r.$$

By (6-31) and (6-32), we have

$$((-\Delta)^s + a)(u_{\varepsilon_j} - M_2 v_2) = (-\Delta)^s u_{\varepsilon_j} + a u_{\varepsilon_j} \leq 0, \quad \text{for all } |x| \geq r.$$

Thus $u_{\varepsilon_j} - M_2 v_2 \leq 0$ for any $|x| \geq r$, which together with (6-32) implies that

$$u_{\varepsilon_j}(x) \leq \frac{C_2}{|x|^{N+2s}}, \quad \text{for all } |x| \geq r, \quad \text{where } C_2 = M_2 \bar{C}_2.$$

Step 7. Set $x_\varepsilon = \varepsilon y_\varepsilon$. Then $w_\varepsilon(x_\varepsilon) = v_\varepsilon(y_\varepsilon)$. By Step 4, x_ε is a maximum point of w_ε and $\{x_\varepsilon\}_\varepsilon$ is bounded on \mathbb{R}^N . By Step 5, $\lim_{\varepsilon \rightarrow 0} \text{dist}(x_\varepsilon, \mathcal{A}_v) = 0$. By Step 1 and Step 2, $u_\varepsilon(x) = v_\varepsilon(x + y'_\varepsilon) = w_\varepsilon(\varepsilon x + x_\varepsilon - \varepsilon x'_\varepsilon)$, where $x'_\varepsilon = y_\varepsilon - y'_\varepsilon$ is a maximum point of u_ε with $\varepsilon x'_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$. By Step 6,

$$\frac{C_1 \varepsilon^{N+2s}}{|x - x_\varepsilon|^{N+2s}} \leq w_\varepsilon(x) \leq \frac{C_2 \varepsilon^{N+2s}}{|x - x_\varepsilon|^{N+2s}}, \quad \text{for all } |x| \geq R,$$

where $R = 2r + \sup_\varepsilon |x_\varepsilon|$ and C_1, C_2 depend on $R, N, s, \tau, \sup_{\mathbb{R}^N} V(x)$.

The proof is completed. \square

References

- [Alves et al. 2017] C. O. Alves, F. Gao, M. Squassina, and M. Yang, “Singularly perturbed critical Choquard equations”, *J. Differential Equations* **263**:7 (2017), 3943–3988. MR Zbl
- [Ao et al. 2017] W. Ao, J. Wei, and W. Yang, “Infinitely many positive solutions of fractional nonlinear Schrödinger equations with non-symmetric potentials”, *Discrete Contin. Dyn. Syst.* **37**:11 (2017), 5561–5601. MR Zbl
- [Belchior et al. 2017] P. Belchior, H. Bueno, O. H. Miyagaki, and G. A. Pereira, “Remarks about a fractional Choquard equation: ground state, regularity and polynomial decay”, *Nonlinear Anal.* **164** (2017), 38–53. MR Zbl
- [Benci 1982] V. Benci, “On critical point theory for indefinite functionals in the presence of symmetries”, *Trans. Amer. Math. Soc.* **274**:2 (1982), 533–572. MR Zbl
- [Bhattarai 2017] S. Bhattarai, “On fractional Schrödinger systems of Choquard type”, *J. Differential Equations* **263**:6 (2017), 3197–3229. MR Zbl
- [Brändle et al. 2013] C. Brändle, E. Colorado, A. de Pablo, and U. Sánchez, “A concave-convex elliptic problem involving the fractional Laplacian”, *Proc. Roy. Soc. Edinburgh Sect. A* **143**:1 (2013), 39–71. MR Zbl
- [Caffarelli and Silvestre 2007] L. Caffarelli and L. Silvestre, “An extension problem related to the fractional Laplacian”, *Comm. Partial Differential Equations* **32**:7-9 (2007), 1245–1260. MR Zbl
- [Cerami et al. 1986] G. Cerami, S. Solimini, and M. Struwe, “Some existence results for superlinear elliptic boundary value problems involving critical exponents”, *J. Funct. Anal.* **69**:3 (1986), 289–306. MR Zbl
- [Cingolani and Secchi 2015] S. Cingolani and S. Secchi, “Semiclassical analysis for pseudo-relativistic Hartree equations”, *J. Differential Equations* **258**:12 (2015), 4156–4179. MR Zbl
- [d’Avenia et al. 2015] P. d’Avenia, G. Siciliano, and M. Squassina, “On fractional Choquard equations”, *Math. Models Methods Appl. Sci.* **25**:8 (2015), 1447–1476. MR Zbl
- [Dávila et al. 2014] J. Dávila, M. del Pino, and J. Wei, “Concentrating standing waves for the fractional nonlinear Schrödinger equation”, *J. Differential Equations* **256**:2 (2014), 858–892. MR Zbl
- [Di Nezza et al. 2012] E. Di Nezza, G. Palatucci, and E. Valdinoci, “Hitchhiker’s guide to the fractional Sobolev spaces”, *Bull. Sci. Math.* **136**:5 (2012), 521–573. MR Zbl
- [Ding and Wei 2017] Y. Ding and J. Wei, “Multiplicity of semiclassical solutions to nonlinear Schrödinger equations”, *J. Fixed Point Theory Appl.* **19**:1 (2017), 987–1010. MR Zbl
- [Dipierro et al. 2017] S. Dipierro, M. Medina, and E. Valdinoci, *Fractional elliptic problems with critical growth in the whole of \mathbb{R}^n* , Appunti. Scuola Normale Superiore di Pisa (Nuova Serie) **15**, Edizioni della Normale, Pisa, 2017. MR Zbl
- [Fall et al. 2015] M. M. Fall, F. Mahmoudi, and E. Valdinoci, “Ground states and concentration phenomena for the fractional Schrödinger equation”, *Nonlinearity* **28**:6 (2015), 1937–1961. MR Zbl
- [Felmer et al. 2012] P. Felmer, A. Quaas, and J. Tan, “Positive solutions of the nonlinear Schrödinger equation with the fractional Laplacian”, *Proc. Roy. Soc. Edinburgh Sect. A* **142**:6 (2012), 1237–1262. MR Zbl
- [Ghimenti and Van Schaftingen 2016] M. Ghimenti and J. Van Schaftingen, “Nodal solutions for the Choquard equation”, *J. Funct. Anal.* **271**:1 (2016), 107–135. MR Zbl
- [Hardy et al. 1952] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, 2nd ed., Cambridge University Press, 1952. MR Zbl
- [Jin et al. 2014] T. Jin, Y. Li, and J. Xiong, “On a fractional Nirenberg problem, part I: blow up analysis and compactness of solutions”, *J. Eur. Math. Soc. (JEMS)* **16**:6 (2014), 1111–1171. MR Zbl

- [Lenzmann 2009] E. Lenzmann, “Uniqueness of ground states for pseudorelativistic Hartree equations”, *Anal. PDE* **2**:1 (2009), 1–27. MR Zbl
- [Lieb 1977] E. H. Lieb, “Existence and uniqueness of the minimizing solution of Choquard’s nonlinear equation”, *Studies in Appl. Math.* **57**:2 (1977), 93–105. MR Zbl
- [Lions 1980] P.-L. Lions, “The Choquard equation and related questions”, *Nonlinear Anal.* **4**:6 (1980), 1063–1072. MR Zbl
- [Liu and Tang 2019] M. Liu and Z. Tang, “Multiplicity and concentration of solutions for Choquard equation via Nehari method and pseudo-index theory”, *Discrete Contin. Dyn. Syst.* **39**:6 (2019), 3365–3398. MR Zbl
- [Ma and Zhao 2010] L. Ma and L. Zhao, “Classification of positive solitary solutions of the nonlinear Choquard equation”, *Arch. Ration. Mech. Anal.* **195**:2 (2010), 455–467. MR Zbl
- [Miranda 1940] C. Miranda, “Un’osservazione su un teorema di Brouwer”, *Boll. Un. Mat. Ital.* (2) **3** (1940), 5–7. MR JFM
- [Moroz and Van Schaftingen 2013] V. Moroz and J. Van Schaftingen, “Groundstates of nonlinear Choquard equations: existence, qualitative properties and decay asymptotics”, *J. Funct. Anal.* **265**:2 (2013), 153–184. MR Zbl
- [Moroz and Van Schaftingen 2017] V. Moroz and J. Van Schaftingen, “A guide to the Choquard equation”, *J. Fixed Point Theory Appl.* **19**:1 (2017), 773–813. MR Zbl
- [Pekar 1954] S. I. Pekar, *Untersuchungen über die Elektronentheorie der Kristalle*, Akademie-Verlag, Berlin, 1954. Zbl
- [du Plessis 1955] N. du Plessis, “Some theorems about the Riesz fractional integral”, *Trans. Amer. Math. Soc.* **80** (1955), 124–134. MR Zbl
- [Silvestre 2007] L. Silvestre, “Regularity of the obstacle problem for a fractional power of the Laplace operator”, *Comm. Pure Appl. Math.* **60**:1 (2007), 67–112. MR Zbl
- [Willem 1996] M. Willem, *Minimax theorems*, Progress in Nonlinear Differential Equations and their Applications **24**, Birkhäuser, Boston, 1996. MR Zbl
- [Zhong and Tang 2018] X.-J. Zhong and C.-L. Tang, “Ground state sign-changing solutions for a class of subcritical Choquard equations with a critical pure power nonlinearity in \mathbb{R}^N ”, *Comput. Math. Appl.* **76**:1 (2018), 23–34. MR Zbl

Received October 9, 2018. Revised August 11, 2019.

MIN LIU
 SCHOOL OF MATHEMATICAL SCIENCES
 BEIJING NORMAL UNIVERSITY
 BEIJING
 CHINA
 minliu@mail.bnu.edu.cn

ZHONGWEI TANG
 SCHOOL OF MATHEMATICAL SCIENCES
 BEIJING NORMAL UNIVERSITY
 LABORATORY OF MATHEMATICS AND COMPLEX SYSTEMS
 MINISTRY OF EDUCATION
 BEIJING
 CHINA
 tangzw@bnu.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 304 No. 1 January 2020

The moduli space of real vector bundles of rank two over a real hyperelliptic curve	1
THOMAS JOHN BAIRD and SHENGDA HU	
Bounds of double zeta-functions and their applications	15
DEBIKA BANERJEE, T. MAKOTO MINAMIDE and YOSHIO TANIGAWA	
Commensurability growth of branch groups	43
KHALID BOU-RABEE, RACHEL SKIPPER and DANIEL STUDENMUND	
Asymptotic orders of vanishing along base loci separate Mori chambers	55
CHIH-WEI CHANG and SHIN-YAO JOW	
Local Langlands correspondence in rigid families	65
CHRISTIAN JOHANSSON, JAMES NEWTON and CLAUS SORENSEN	
PseudoindeX theory and Nehari method for a fractional Choquard equation	103
MIN LIU and ZHONGWEI TANG	
Symmetry and nonexistence of positive solutions for fractional Choquard equations	143
PEI MA, XUDONG SHANG and JIHUI ZHANG	
Decomposability of orthogonal involutions in degree 12	169
ANNE QUÉGUINER-MATHIEU and JEAN-PIERRE TIGNOL	
Zelevinsky operations for multisegments and a partial order on partitions	181
PETER SCHNEIDER and ERNST-WILHELM ZINK	
Langlands parameters, functoriality and Hecke algebras	209
MAARTEN SOLLEVELD	
On the archimedean local gamma factors for adjoint representation of GL_3 , part I	303
FANGYANG TIAN	
An explicit CM type norm formula and effective nonvanishing of class group L-functions for CM fields	347
LIYANG YANG	



0030-8730(2020)304:1;1-S