

*Pacific
Journal of
Mathematics*

GRAPHS ADMITTING ONLY CONSTANT SPLINES

KATIE ANDERS, ALISSA S. CRANS, BRIANA FOSTER-GREENWOOD,
BLAKE MELLOR AND JULIANNA TYMOCZKO

Volume 304 No. 2

February 2020

GRAPHS ADMITTING ONLY CONSTANT SPLINES

KATIE ANDERS, ALISSA S. CRANS, BRIANA FOSTER-GREENWOOD,
BLAKE MELLOR AND JULIANNA TYMOCZKO

We study *generalized graph splines*, introduced by Gilbert, Tymoczko, and Viel (2016). For a large class of rings, we characterize the graphs that only admit constant splines. To do this, we prove that if a graph has a particular type of cutset (e.g., a bridge), then the space of splines naturally decomposes as a certain direct sum of submodules. As an application, we use these results to describe splines on a triangulation studied by Zhou and Lai, but over a different ring than they used.

1. Introduction

This paper studies generalized splines, which are parametrized by a ring, a graph, and a map from the edges of the graph to ideals in the ring. As the name suggests, they generalize the classical splines from analysis and applied mathematics. The main goal of this paper is to describe when the module of generalized splines has rank one over the base ring. We give multiple equivalent conditions for this to be true over a large family of rings. Our main tools are of independent interest: different reductions on the module of splines depending on either algebraic characteristics of the edge labeling or combinatorial characteristics of the graph.

Classically, a spline is a collection of polynomials on the faces of a polyhedral complex that agree to specified degree of smoothness on the intersections of faces. More formally, given a simplicial complex Δ we define the vector space of *splines* $S_d^r(\Delta)$ to be the space of all piecewise polynomial functions on Δ that have degree d and order of smoothness r . Splines are a standard topic in numerical analysis and are used in data interpolation, geometric design, and to approximate solutions to partial differential equations, among other applications. Splines are also studied from a more theoretical perspective by analysts (see [Lai and Schumaker 2007] for a survey). Two fundamental problems in both contexts are to find a basis (often satisfying specified constraints) or to compute the dimension of the vector space of splines over a given polyhedral complex, e.g., [Strang 1973; Alfeld and Schumaker 1987; Alfeld et al. 1987; Chui and Lai 1990; Lai and Schumaker 2007].

MSC2010: 05C25, 05C78, 05E15.

Keywords: splines, generalized graph splines.

Our approach is essentially dual to that of classical splines. Billera and Rose [1991] observed that splines can be viewed as functions on the *dual graph* of the polyhedral complex, reinterpreting the order-of-smoothness condition as a compatibility condition across each edge of the graph. Independently, splines were reinvented in a combinatorial construction of equivariant cohomology, often called *GKM theory* by symplectic geometers and algebraic topologists [Goresky et al. 1998; Tymoczko 2016], and also arise naturally in the context of toric varieties in algebraic geometry [Payne 2006].

The following definition unifies and generalizes previous work, allowing us to define splines on an arbitrary graph (not just those that can be realized in geometric settings) and to consider a much larger class of rings (not just polynomial rings). It first appeared in work of Gilbert, Viel, and the last author of this work [Gilbert et al. 2016], and extends earlier work by Guillemin and Zara [2000] to put GKM theory in a combinatorial context (see also, e.g., [Guillemin and Zara 2001a; 2001b; 2003]).

Definition 1.1. Given a graph $G = (V, E)$ and a commutative ring R with unit, an *edge labeling* of G is a function $\alpha : E \rightarrow I(R)$, where $I(R)$ is the set of ideals of R . A *spline* p on (G, α) is a vertex labeling $p : V \rightarrow R$ such that for each edge uv , the difference $p(u) - p(v)$ lies in the ideal $\alpha(uv)$. Let $S_R(G, \alpha)$ denote the set of all splines on G with labels from R and edge labeling α . When R and α are clear from context, we write $S(G)$.

In geometric applications, we have an underlying complex algebraic variety with a well behaved torus action. The ring R is the collection of polynomials in n variables, where n is the rank of the torus. The graph corresponds to the 1-skeleton of the moment polytope with respect to the torus action and the labeling α records the weight of the torus action on each 1-dimensional torus orbit in the variety.

Note that $S_R(G, \alpha)$ is both a ring (with addition and multiplication of splines defined pointwise) and an R -module [Gilbert et al. 2016]. We focus on the R -module structure of $S_R(G, \alpha)$ in this paper. Any collection of R -module generators of $S_R(G, \alpha)$ generates $S_R(G, \alpha)$ as a ring. However, a minimal set of ring generators can be much smaller, and computing a multiplication table is generally very difficult — in fact, it is a longstanding open problem to compute the multiplication table with respect to the Schubert basis for the ring of splines corresponding to the equivariant cohomology of the full flag variety.

Analogous to the classical problem of finding a basis for the vector space of splines, a core problem in the theory of generalized graph splines is to describe minimal generating sets for $S_R(G, \alpha)$ as an R -module. This is not difficult to do when the graph is a tree [Gilbert et al. 2016] but is significantly harder for graphs containing cycles. Various authors have studied the existence, size and construction

of these generating sets for cycles and other graphs over the integers, the rings \mathbb{Z}_m , and other rings [Altinok and Sarioglan 2019a; 2019b; Bowden et al. 2015; Bowden and Tymoczko 2015; DiPasquale 2017; Guillemin and Zara 2001a; 2003; Handschy et al. 2014; Philbin et al. 2017].

When the ring R is an integral domain, then every generating set for $S_R(G, \alpha)$ has at least n elements, where n is the number of vertices in the graph G [Gilbert et al. 2016]. If R is not an integral domain, however, this is no longer true; in fact, there are nontrivial labeled graphs that admit only constant splines [Bowden and Tymoczko 2015]. Determining whether there are nonconstant splines is thus a fundamental question in the field.

Our primary goal in this paper is to describe the edge labeled graphs which only admit constant splines. In Theorem 3.5 we provide two equivalent combinatorial characterizations of these graphs for a large family of rings, one in terms of cutsets and the other in terms of spanning trees.

Theorem 3.5. *Suppose $R = \bigoplus_{i=1}^k R_i$ and that each R_i is an irreducible commutative ring with identity 1_{R_i} . Let G be a connected graph with edge set $E(G)$ and edge labeling α . Then the following are equivalent.*

- (1) *The edge labeled graph (G, α) has rank one over R .*
- (2) *For any cutset $C \subset E(G)$, the intersection $\bigcap_{e \in C} \alpha(e) = 0$.*
- (3) *The graph G has spanning trees T_1, \dots, T_k such that for each i , all edge labels of T_i are contained in the kernel of the canonical projection $R \rightarrow R_i$.*

Along the way, we study operations that induce a decomposition of the module of splines. We take two approaches, with the following main results:

- In Corollary 2.5 we give an algebraic condition on the labeling α under which the splines $S_R(G, \alpha)$ can be reduced (as a ring and as an R -module) to the splines on a spanning tree of G . This condition is essentially that the edge labels form a set of nested ideals.
- In Theorem 2.10 we give a combinatorial condition on the graph G under which the R -module $S_R(G, \alpha)$ decomposes into a direct sum of specific submodules. Our condition generalizes the graph-theoretic notion of a bridge and applies to any choice of ring R .

As an application, we consider splines on a triangulation studied by Zhou and Lai [2013] but over integers mod m rather than their polynomial rings. The contrast between our results and theirs demonstrates the significant differences that can occur when changing the base ring.

2. Decomposing graphs

In this section we decompose splines based on different combinatorial conditions of the graph or algebraic conditions on the labeling. In what follows, we assume that our graphs G are connected because the ring (and module) of splines naturally decomposes over disconnected components.

The following lemma describes operations on graphs that allow us to assume no edge is labeled either by the ideal (0) or by the ideal (1). It was proven by Gilbert, Tymoczko, and Viel.

Lemma 2.1 [Gilbert et al. 2016]. *If $\alpha(uv)$ is the ideal generated by 0 and G/uv is the graph obtained by contracting edge uv then $S(G) \cong S(G/uv)$. If $\alpha(uv)$ is the whole ring (generated by (1) in R), then $S(G) \cong S(G - uv)$, where $G - uv$ is the graph obtained by deleting the edge uv .*

Contractions can produce loops or multiple edges, so using Lemma 2.1 appears to require splines on *multigraphs* (which can have loops and multiple edges between two vertices). However, the next result is the key step to prove that for each multigraph, there is a simple graph with the same ring of splines. Our definition for splines on a multigraph is that the spline condition must be satisfied for each edge individually.

Lemma 2.2. *Suppose (G, α) is a multigraph, with two edges e_1 and e_2 between vertices u and v . Suppose G' is the graph obtained by replacing e_1 and e_2 with a single edge e . Suppose α' is defined by setting $\alpha'(e) = \alpha(e_1) \cap \alpha(e_2)$ and by setting $\alpha'(e') = \alpha(e')$ for all other edges e' . Then $S(G, \alpha) = S(G', \alpha')$.*

Proof. First suppose $p \in S(G, \alpha)$. Then $p(u) - p(v) \in \alpha(e_1)$ and $p(u) - p(v) \in \alpha(e_2)$. Thus $p(u) - p(v) \in \alpha(e_1) \cap \alpha(e_2) = \alpha'(e)$ so $p \in S(G', \alpha')$.

Conversely, if $p \in S(G', \alpha')$, then $p(u) - p(v) \in \alpha'(e) = \alpha(e_1) \cap \alpha(e_2)$. Thus $p(u) - p(v)$ is in both ideals $\alpha(e_i)$ and so $p \in S(G, \alpha)$. This proves the claim. \square

The next corollary is immediate.

Corollary 2.3. *Suppose (G, α) is a multigraph. Let G' be the graph obtained by erasing every loop (namely edge of the form vv for some vertex v) from G , and by erasing all but one edge in the event of multiple edges between any two vertices uv . Let α' be the edge labeling obtained by assigning to the edge uv in G' the edge label $\alpha(e_1) \cap \alpha(e_2) \cap \cdots \cap \alpha(e_k)$, where e_1, e_2, \dots, e_k are all edges between u and v in G . Then $S(G, \alpha) = S(G', \alpha')$.*

Proof. Loops start and end at the same vertex so the spline condition is trivially satisfied on each loop. Thus loops can be removed without changing the space of splines. Applying Lemma 2.2 repeatedly then gives the result. \square

If R is a field, then its only ideals are (0) and (1) . Lemma 2.1 then says that the ring $S(G)$ is isomorphic to the ring of splines on an isolated set of, say, n vertices, which is simply R^n . For this reason, we generally do not consider splines over fields.

Edges of cycles. We give a new condition under which an edge of a cycle may be deleted without losing any information: that the ideal associated to one edge uv of the cycle contains all of the other edge labels of the cycle. Informally, in this situation the edge uv contains information that is redundant with the rest of the cycle and so may be erased. The next lemma states this formally; in the rest of the section, we elaborate some consequences.

Lemma 2.4. *Let (G, α) be an edge labeled graph. Let uv be an edge and suppose there is a cycle C in G that contains uv such that the ideal $\alpha(uv)$ contains every edge label from C . Then $S(G) = S(G - \{uv\})$.*

Proof. Every spline on G is also a spline on $G - \{uv\}$, so $S(G) \subseteq S(G - \{uv\})$.

Now suppose that $p \in S(G - \{uv\})$. Let $w_0w_1w_2 \cdots w_nw_{n+1}w_0$ be a cycle in G that contains the edge uv , say with $w_0 = u$ and $w_{n+1} = v$. Since

$$p(u) - p(v) = p(u) - p(w_1) + p(w_1) - p(w_2) + \cdots + p(w_n) - p(v)$$

and since $p(w_i) - p(w_{i+1}) \in \alpha(w_iw_{i+1})$ for each i , we know that

$$p(u) - p(v) \in \sum_{i=0}^n \alpha(w_iw_{i+1}).$$

This is contained in $\alpha(uv)$ by the hypothesis that $\alpha(uv)$ contains all the edge labels $\alpha(w_iw_{i+1})$ for $0 \leq i \leq n$. Hence p is also a spline for G and so $S(G - \{uv\}) \subseteq S(G)$. The claim follows. □

Lemma 2.4 has particularly interesting consequences when the set of edge labels is linearly ordered (namely, when any two distinct edge labels α_1 and α_2 satisfy either $\alpha_1 \subset \alpha_2$ or $\alpha_2 \subset \alpha_1$). In this case, we can reduce to considering the R -module of splines on a tree.

Corollary 2.5. *Suppose (G, α) is a connected graph whose set of edge labels can be linearly ordered by inclusion. Then there exists a spanning tree T such that $S(G) = S(T)$.*

Proof. If G is a tree, then we are done. Otherwise, the graph G must contain a cycle. Since the set of edge labels can be linearly ordered, there must be an edge, say uv , of the cycle such that $\alpha(uv)$ contains all other ideals labeling the edges of that cycle. By Lemma 2.4, we have $S(G) = S(G - \{uv\})$. Note that $G - \{uv\}$ is still connected, so either $G - \{uv\}$ is a tree or we can find another cycle and find an

edge in that cycle to delete. Continuing until no cycles remain, we obtain a tree T with $S(G) = S(T)$. \square

In particular Corollary 2.5 applies when working over rings for which the set of *all* ideals is linearly ordered; such rings are known as *uniserial rings*. Examples include \mathbb{Z}_{p^k} for p prime and $F[x]/(x^n)$ where F is a field. Furthermore, commutative uniserial rings that are domains are exactly valuation rings.

Remark 2.6. Corollaries 2.5 and 2.11 can together be used to describe bases of splines over \mathbb{Z}_{p^e} for any prime p (see Remark 2.12 for more detail on how to generate bases of trees). However, our results do not give an explicit construction like that in Philbin–Swift–Tammaro–Williams [Philbin et al. 2017].

Bridges. We now turn to the case of graphs that contain a *bridge*, i.e., an edge whose deletion disconnects the graph. We will show that the module of splines on the graph is (almost) the direct sum of the spline modules for the two components after the bridge is deleted. We find it convenient to define splines *based at a vertex*.

Definition 2.7. Given a vertex v of a graph G , we say that a spline $p \in S_R(G, \alpha)$ is *based at v* if $p(v) = 0$. We denote the submodule of splines based at v by $S_R(G, \alpha; v)$ and call v the *basepoint* of this submodule. We write $S(G; v)$ if R and α are clear.

Our terminology is new but the idea is not, e.g., [Guillemin and Zara 2003; Knutson and Tao 2003; Gilbert et al. 2016].

Note that $S(G; v)$ may equal $S(G; w)$ for distinct vertices v and w . This happens precisely when $p(v) = p(w)$ for every spline $p \in S(G)$.

The next lemma restates an earlier result using our terminology. It says that $S(G; v)$ is almost the same as $S(G)$. We use $\mathbb{1}_G \in S(G)$ to denote the constant spline that takes the value 1 on every vertex.

Lemma 2.8 [Gilbert et al. 2016, Theorem 2.12]. *For any vertex v of G , there is an R -module decomposition*

$$S_R(G) \cong S_R(G; v) \oplus \langle \mathbb{1}_G \rangle \cong S_R(G; v) \oplus R.$$

Remark 2.9. The submodule of splines based at vertex v is closed under multiplication and so is an ideal in $S_R(G)$. However, the decomposition in Lemma 2.8 is not a ring isomorphism.

Extending Lemma 2.8, we next show that if a graph has a bridge, its space of splines is (almost) the direct sum of the modules of splines on the two components created by removing the bridge. In fact, we prove this in an even more general setting, one in which the “bridge” is a subgraph H that is more complicated than an edge. We still require this “generalized bridge” to disconnect the graph and to meet each component G_i of the disconnected graph in a single vertex h_i , as in Figure 1.




Figure 1. A generalized bridge.

Intuitively, our proof decomposes $S(G)$ into a direct sum of splines on H and on each G_i . More formally, we have the following.

Theorem 2.10. *Suppose (G, α) is an edge labeled graph with a subgraph H such that each connected component of the graph $G - E(H)$ contains exactly one vertex of H . Suppose H has vertices h_1, \dots, h_n and G_i is the component of $G - E(H)$ containing h_i . Then $S(G)$ is isomorphic to the direct sum of R -modules:*

$$S(G) \cong \langle \mathbb{1}_G \rangle \oplus S(H; h_1) \oplus S(G_1; h_1) \oplus \dots \oplus S(G_n; h_n).$$

Moreover, for each i , let $\tilde{S}(G_i; h_i)$ denote the subset of splines in $S(G)$ that are zero when restricted to the vertices in $H \cup \bigcup_{j \neq i} G_j$. Let $\tilde{S}(H)$ denote the splines in $S(G)$ such that for each i the restriction to G_i is constant. (If $i \neq j$ then the restrictions to G_i and G_j need not agree.) Then $S(G)$ is the internal direct sum of submodules

$$S(G) = \langle \mathbb{1}_G \rangle \oplus \tilde{S}(H; h_1) \oplus \tilde{S}(G_1; h_1) \oplus \dots \oplus \tilde{S}(G_n; h_n).$$

Proof. Suppose p is a spline in $S(G)$. The spline defined by $p' = p - p(h_1)\mathbb{1}_G$ is in $S(G; h_1)$. Let p'_H be the restriction of p' to H and let \tilde{p}'_H be the extension of p'_H to G defined by setting $\tilde{p}'_H(u) = p'(h_i)$ for all i and vertices u in G_i . Note that $\tilde{p}'_H \in \tilde{S}(H; h_1) \subseteq S(G; h_1)$. Now for each i , let q_i be the restriction of $p' - \tilde{p}'_H$ to G_i , so $q_i \in S(G_i; h_i)$. Extend q_i to the spline $\tilde{q}_i \in S(G)$ by setting $\tilde{q}_i(u) = 0$ for all vertices u off of G_i . Note that $\tilde{q}_i \in \tilde{S}(G_i; h_i)$. Thus by construction

$$p = p(h_1)\mathbb{1}_G + \tilde{p}'_H + \sum_{i=1}^n \tilde{q}_i \in \langle \mathbb{1}_G \rangle \oplus \tilde{S}(H; h_1) \oplus \tilde{S}(G_1; h_1) \oplus \dots \oplus \tilde{S}(G_n; h_n).$$

Now we show that this decomposition of p is unique. Suppose

$$p = r\mathbb{1}_G + q_0 + q_1 + q_2 + \dots + q_n = 0,$$

where $q_0 \in \widetilde{S}(H; h_1)$ and $q_i \in \widetilde{S}(G_i, h_i)$ for $1 \leq i \leq n$. Note that $p(h_i) = q_0(h_i) + r$ for each i and, in particular, $p(h_1) = r$. Since $p = 0$ this means $r = 0$. Then

$$\widetilde{p}_H = q_0 + r\mathbb{1}_G = q_0,$$

since each q_i is zero on H and p is constant on each G_i . But since $p = 0$, we can explicitly compute $p'_H = 0$ and so $q_0 = 0$. An analogous argument shows that for each G_i , the spline

$$\widetilde{p}_{G_i} = q_i + (q_0(h_i) + r)\mathbb{1}_{G_i} = q_i = 0,$$

and so the decomposition is unique.

Finally, note that the restriction map is a natural module isomorphism between the submodule $\widetilde{S}(H; h_1)$ and $S(H; h_1)$, respectively $\widetilde{S}(G_i; h_i)$ and $S(G_i; h_i)$. This proves the claim. \square

We apply this decomposition to the special case of a graph with a bridge that is labeled by a principal ideal; it generalizes to any graph with a bridge if the single generator $\beta\chi_B$ is replaced by the set of generators of the ideal labeling the bridge.

Corollary 2.11. *Let G be a connected graph with bridge ab and edge labeling α such that the bridge is labeled by the principal ideal $\alpha(ab) = (\beta)$. The graph $G - \{ab\}$ has two components; denote the component that contains a by A and the component that contains b by B . Let χ_B represent the function (**not** necessarily a spline) that is identically 1 on the vertices of B and 0 on the vertices of A . Then we have an R -module decomposition*

$$S(G) \cong S(A; a) \oplus S(B; b) \oplus \langle \mathbb{1}_G \rangle \oplus \langle \beta\chi_B \rangle.$$

Proof. This follows directly from the previous theorem once we note that the graph consisting simply of the edge ab is generated as an R -module by the identity spline together with the spline that is β on b and 0 on a and that χ_B is the extension of this latter spline to all of G . \square

Remark 2.12. If G is a tree, we can recursively apply Corollary 2.11 to obtain a minimal generating set for $S(G)$ as an R -module. In this sense, our result generalizes the construction of bases of splines for trees in [Gilbert et al. 2016].

The following example illustrates how Lemma 2.4, and Corollaries 2.5, and 2.11 can be used to find a minimal set of generators for the R -module of splines on a graph whose edge labels can be linearly ordered by inclusion.

Example 2.13. Let $\beta, \gamma, \delta \in R$. Suppose we have the graph G with vertices a, b, c and edge labeling α given in Figure 2. Let $G' = G - \{bc\}$ and $G'' = G' - \{ab\}$.

Since $\alpha(bc) = (\delta)$ contains every other edge label of the cycle, we have $S(G) = S(G')$ by Lemma 2.4. Thus we have reduced to a tree as described by Corollary 2.5.




Figure 2. Graphs of Example 2.13.

To apply Corollary 2.11, we consider the bridge ab and let $G'' = G' - \{ab\}$. Let A be the component of G'' containing a and B be the component of G'' containing b . If the coordinates are ordered (a, b, c) , then the generators of $S(A; a)$ are $\mathcal{B}_a = \{(0, 0, \delta\beta\gamma)\}$ and $S(B; b) = 0$. By Corollary 2.11 we have

$$S(G) = S(G') = S(A; a) \oplus S(B; b) \oplus \langle \mathbb{1}_G \rangle \oplus \langle \delta\beta\chi_B \rangle$$

for which a minimal generating set is $U = \{(0, 0, \delta\beta\gamma), (1, 1, 1), (0, \delta\beta, 0)\}$.

3. Graphs with rank one

In this section, we consider labeled graphs that admit only constant splines, which we call graphs with rank one.

Definition 3.1. Given a graph G with edge labeling α , a *constant spline* on (G, α) takes the same value on every vertex of G . The graph (G, α) has *rank one* over the ring R if it admits only constant splines.

Graphs with rank one are particularly interesting from the point of view of subgraphs. By Theorem 2.10, if the “bridge” graph H has rank one, then its contribution to the space of splines is just zero. More generally, if a subgraph has rank one, it can be contracted to a vertex without changing the space of splines. Our main result is a characterization of rank one graphs over rings which are direct sums of irreducible rings.

Definition 3.2. Recall that an ideal I of a ring R is *irreducible* (sometimes called *meet-irreducible*) if it is not the intersection of two strictly larger ideals. A ring R is *irreducible* if (0) is an irreducible ideal in R . In other words, a ring R is irreducible if the intersection of any two nonzero ideals is nonzero.

Remark 3.3. Examples of irreducible rings include integral domains, uniserial rings such as \mathbb{Z}_{p^k} for p prime, and Artinian Gorenstein rings. An irreducible ring cannot be decomposed as the direct sum of two nontrivial rings, though the converse is not true. For example, the ring $\mathbb{R}[x, y]/(x^2, y^2, xy)$ cannot be decomposed as a direct sum of two of its ideals because one of those two ideals must contain

polynomials with a nonzero constant term, and $ax + by + c$ generates the entire ring whenever $c \neq 0$. However, this quotient is not irreducible since $(x) \cap (y) = (0)$.

We now consider splines over a ring $R = \bigoplus_{i=1}^k R_i$, where each R_i is an irreducible commutative ring with identity 1_{R_i} . Let $\pi_i : R \rightarrow R_i$ be the canonical projection homomorphism and let M_i denote the kernel of π_i .

Lemma 3.4. *Suppose $R = \bigoplus_{i=1}^k R_i$ and that each R_i is an irreducible commutative ring with identity 1_{R_i} . If I and J are ideals of R that are not contained in M_i , then $I \cap J$ is not contained in M_i .*

Proof. Suppose that I and J are ideals of R that are not contained in M_i for some i . It follows that $\pi_i(I) \neq 0$ and $\pi_i(J) \neq 0$. Since R_i is irreducible, the ideal $\pi_i(I) \cap \pi_i(J)$ contains a nonzero element, say x . There must exist elements $r \in I$ and $s \in J$ such that $\pi_i(r) = \pi_i(s) = x$. Multiplying r and s each by $(0, \dots, 0, 1_{R_i}, 0, \dots, 0)$ produces an element $(0, \dots, 0, x, 0, \dots, 0)$ which belongs to both I and J but has nonzero i -th coordinate. Hence $I \cap J$ is not contained in M_i . This proves the claim. \square

We use this lemma to prove our main theorem, which is a complete graph-theoretic characterization of the graphs of rank one over rings R of this form.

Theorem 3.5. *Suppose $R = \bigoplus_{i=1}^k R_i$ and that each R_i is an irreducible commutative ring with identity 1_{R_i} . Let G be a connected graph with edge set $E(G)$ and edge labeling α . Then the following are equivalent.*

- (1) *The edge labeled graph (G, α) has rank one over R .*
- (2) *For any cutset $C \subset E(G)$, the intersection $\bigcap_{e \in C} \alpha(e) = 0$.*
- (3) *The graph G has spanning trees T_1, \dots, T_k such that all edge labels of T_i are contained in M_i for all $1 \leq i \leq k$.*

Remark 3.6. When R itself is irreducible, equivalently $k = 1$, the third condition reduces to G having a spanning tree with edges labeled by the zero ideal. In general, the spanning trees T_i need not be disjoint (or even distinct).

Proof. First we prove (1) \Rightarrow (2) by contrapositive. Let $C \subset E(G)$ be a cutset of G . Without loss of generality, we assume that C is minimal in the sense that no proper subset of C is a cutset. Then there exists a partition of the vertex set V into nonempty subsets V_1 and V_2 such that $C = \{uv \in E(G) \mid u \in V_1, v \in V_2\}$. Suppose there exists a nonzero element x of $\bigcap_{e \in C} \alpha(e)$. Define a vertex labeling $p : V \rightarrow R$ by $p(u) = x$ for all $u \in V_1$ and $p(v) = 0$ for all $v \in V_2$. The restriction of p to V_1 is a spline, as is the restriction of p to V_2 . For any $u \in V_1$ and any $v \in V_2$, we have $p(u) - p(v) = x \in \bigcap_{e \in C} \alpha(e)$. The edge uv is in C , so we conclude $x \in \alpha(uv)$. Thus p is a nonconstant spline on G , so (G, α) does not have rank one.




Figure 3. Graph of Example 3.7.

Next we prove (2) \Rightarrow (3) by contrapositive. For $1 \leq i \leq k$, consider the equivalence relation \equiv_i on $V(G)$ defined by $u \equiv_i v$ if and only if there exists a path from u to v having all edge labels contained in M_i . Now suppose i is an index such that there is no spanning tree T_i such that all of its edge labels are contained in M_i . Then the relation \equiv_i determines more than one equivalence class. Let V_1 be one of the equivalence classes and let V_2 be the union of the rest. Note that if $v_1 v_2$ is an edge with $v_1 \in V_1$ and $v_2 \in V_2$, then $\alpha(v_1 v_2)$ cannot be contained in M_i since $v_2 \not\equiv_i v_1$. Now $C := \{v_1 v_2 \in E(G) \mid v_1 \in V_1, v_2 \in V_2\}$ is a cutset (which is nonempty since G is connected). Lemma 3.4 implies that $\bigcap_{e \in C} \alpha(e) \not\subseteq M_i$. Hence C is a cutset for which the intersection $\bigcap_{e \in C} \alpha(e)$ is nonzero.

Finally, we prove (3) \Rightarrow (1). Suppose that for all $1 \leq i \leq k$ the graph (G, α) has a spanning tree T_i whose edge labels are all in M_i . Suppose p is a spline on G . Since every pair of vertices u and v is connected by a path in T_i , we know $p(u) - p(v) \in M_i$. But then $p(u) - p(v) \in \bigcap_{i=1}^k M_i = 0$. Hence p is a constant spline. \square

The implications (3) \Rightarrow (1) \Rightarrow (2) hold for any direct product $R = \bigoplus_{i=1}^k R_i$, but Examples 3.7 and 3.8 illustrate how the converse implications can fail if the factors are not irreducible.

Example 3.7. Let $R = \mathbb{R}[x, y]/(x^2, xy, y^2)$ and consider the graph shown in Figure 3. Each cutset of a triangle contains at least two edges, so since the pairwise intersections of the edge labels are trivial, Condition (2) of Theorem 3.5 holds. But the vertex labeling above is a nonconstant spline, so Condition (1) is false.

Example 3.8. Consider the labeled graph (K_4, α) over $R = \mathbb{R}[x, y]/(x^2, y^2, xy)$ shown in Figure 4.

Suppose p is a spline on this graph. Inspect the following triangles:

- Triangle 124 shows $p(4) - p(1) \in (x) \cap (y) = 0$.
- Triangle 234 shows $p(4) - p(2) \in (y) \cap (x + y) = 0$.
- Triangle 314 shows $p(4) - p(3) \in (x + y) \cap (x) = 0$.




Figure 4. Graph of Example 3.8.

Thus $p(4) = p(i)$ for $i = 1, 2, 3$, so p is a constant spline.

Hence this graph satisfies Condition (1) in Theorem 3.5. However, since R cannot be decomposed as a direct sum (see Remark 3.6), to satisfy Condition (3) the graph must have a spanning tree with edges labeled (0), which it does not.

As a corollary to Theorem 3.5, we characterize when a labeled tree has rank one over any ring.

Corollary 3.9. *If (G, α) is a tree over a ring R , it is rank one if and only if every edge is labeled (0).*

Proof. If every edge is labeled (0), then the graph can be contracted to a single vertex without changing the space of splines, by Lemma 2.1. Hence the space of splines will be generated by the constant splines, and the graph has rank one.

Conversely, since every edge of a tree is a bridge, if there is an edge whose label is not (0), then by the first part of Theorem 3.5 there is a nonconstant spline, so the graph does not have rank one. (Note that the proof that (1) \Rightarrow (2) in Theorem 3.5 does not impose any restriction on the ring.) \square

Integers mod m . As a particular case of Theorem 3.5, we consider the ring \mathbb{Z}_m of integers modulo m . Splines over \mathbb{Z}_m were previously studied by Bowden and Tymoczko [2015], who showed that the space of splines on a graph of n vertices could have any rank between 1 and n . We characterize when the space of splines has rank one.

Corollary 3.10. *Fix an integer m with prime factorization $m = p_1^{e_1} \cdots p_k^{e_k}$ and let $R = \mathbb{Z}_m$. The graph (G, α) has rank one if and only if G contains spanning trees T_1, \dots, T_k such that all edge labels of T_i are contained in the ideal $(p_i^{e_i})$. In particular, if $m = p^e$ then (G, α) has rank one if and only if G has a spanning tree with all edges labeled (0).*

Proof. If the prime factorization of m is $m = p_1^{e_1} \cdots p_k^{e_k}$, then the Chinese Remainder Theorem implies $\mathbb{Z}_m \cong \mathbb{Z}_{p_1^{e_1}} \oplus \cdots \oplus \mathbb{Z}_{p_k^{e_k}}$. Each ring $\mathbb{Z}_{p_i^{e_i}}$ is uniserial and hence




Figure 5. Graph (K_4, α) of Example 3.11.

irreducible, and $M_i = (p_i^{e_i})$ for each i . The claim thus follows from Theorem 3.5 and Remark 3.6. □

Example 3.11. Let $R = \mathbb{Z}_{pq}$, where p and q are distinct primes. The only ideals in R are (0) , (p) , (q) , (1) . Consider the graph (K_4, α) shown in Figure 5.

The thick and thin edges indicate two spanning trees, one with all edges labeled (p) and the other with all edges labeled (q) . By Corollary 3.10 this graph has rank one. Note this graph has rank one even though *none* of the edges are labeled (0) .

Example 3.12. Instead of simply asking whether a particular labeled graph has rank one, as in the previous example, we could instead ask more generally *which* labelings of a given (unlabeled) graph will yield a labeled graph with rank one.

Motivated by classical splines, we examine the dual graph of a particular triangulation of the plane. We begin with a rectangular grid each of whose squares is divided by a diagonal into two triangles, with the triangles then subdivided by a Clough–Tocher refinement as shown on the left in Figure 6. Starting with an $m \times n$ grid, we denote this graph $G_{m,n}$. The graph dual to the triangulated grid (ignoring the exterior region) is shown on the right in Figure 6; we denote this graph $G_{m,n}^*$.

Splines on these graphs were studied by Zhou and Lai [2013], though we study them over \mathbb{Z}_r rather than the base ring Zhou and Lai used (namely polynomials with two variables). The module of splines in this example differs very dramatically from that of Zhou and Lai, demonstrating the impact of changing the base ring.

Note that the graph $G_{m,n}^*$ always contains at least two bridges. If the graph has rank one, then all of the bridges must be labeled (0) . In the following proposition we use Corollary 3.10 to give a better lower bound on the number of edges labeled (0) than simply by counting bridges.

Proposition 3.13. *If r has at most 3 distinct prime factors and α is a labeling of the edges of $G_{m,n}^*$ over \mathbb{Z}_r such that $(G_{m,n}^*, \alpha)$ has rank one, then a lower bound on the number of edges labeled (0) is*

- $6mn - 1$ if $r = p^a$,




Figure 6. Dual graph for a triangulated rectangular grid.

- $3mn + m + n - 2$ if $r = p_1^{a_1} p_2^{a_2}$, and
- $2m + 2n - 3$ if $r = p_1^{a_1} p_2^{a_2} p_3^{a_3}$.

Proof. Our argument depends on counting the vertices and edges in $G_{m,n}^*$. Observe that $G_{m,n}$ has $6mn$ regions and $9mn + m + n$ edges. The exterior edges do not contribute to the dual graph and there are exactly $9mn - m - n$ interior edges. So $G_{m,n}^*$ has $6mn$ vertices and $9mn - m - n$ edges.

If r has only one prime factor, then by Corollary 3.10 the graph $G_{m,n}^*$ has a spanning tree with all edges labeled (0). Since $G_{m,n}^*$ has $6mn$ vertices, its spanning tree has $6mn - 1$ edges.

Next suppose $\mathbb{Z}_r = \mathbb{Z}_{p^a q^b}$. Also by Corollary 3.10, for every α the labeled graph $(G_{m,n}^*, \alpha)$ has rank one over $\mathbb{Z}_{p^a q^b}$ if and only if $G_{m,n}^*$ contains spanning trees T_1 and T_2 with all edge labels of T_1 in (p^a) and all edge labels of T_2 in (q^b) . These ideals only intersect in (0) so T_1 and T_2 only overlap on edges labeled (0).

Since $G_{m,n}^*$ has $6mn$ vertices, each spanning tree has $6mn - 1$ edges. Since there are only $9mn - m - n$ edges total, the two spanning trees must overlap in at least $2(6mn - 1) - (9mn - m - n) = 3mn + m + n - 2$ edges. Thus for $(G_{m,n}^*, \alpha)$ to have rank one, there must be at least $3mn + m + n - 2$ edges labeled (0).

Finally, suppose $\mathbb{Z}_r = \mathbb{Z}_{p_1^{a_1} p_2^{a_2} p_3^{a_3}}$, where p_1, p_2, p_3 are distinct primes. Then $(G_{m,n}^*, \alpha)$ has rank one if and only if there are three spanning trees T_i such that all labels of T_i are in $(p_i^{a_i})$. The intersection $T_1 \cap T_2$ contains at least $3mn + m + n - 2$ edges, all with labels contained in $(p_1^{a_1} p_2^{a_2})$. Hence $(T_1 \cap T_2) \cap T_3$ contains at least

$$(6mn - 1) + (3mn + m + n - 2) - (9mn - m - n) = 2m + 2n - 3 \geq 1$$

edges, all of which must be labeled (0). □

Acknowledgements

We thank the Institute for Computational and Experimental Research in Mathematics and the American Institute of Mathematics for their generous support of the Research Experience for Undergraduate Faculty (REUF) program, which provided the authors

the opportunity to meet for a week during the summers of 2017 and 2018. We also thank the Simons Foundation for its support (#360097, Alissa Crans) and the National Science Foundation for its support (NSF-DMS 1362855 and 1800773, Julianna Tymoczko). Finally, we are deeply grateful to the referee for their careful reading and helpful suggestions, which greatly improved our work.

References

- [Alfeld and Schumaker 1987] P. Alfeld and L. L. Schumaker, “The dimension of bivariate spline spaces of smoothness r for degree $d \geq 4r + 1$ ”, *Constr. Approx.* **3**:2 (1987), 189–197. MR Zbl
- [Alfeld et al. 1987] P. Alfeld, B. Piper, and L. L. Schumaker, “An explicit basis for C^1 quartic bivariate splines”, *SIAM J. Numer. Anal.* **24**:4 (1987), 891–911. MR Zbl
- [Altinok and Sarioglan 2019a] S. Altinok and S. Sarioglan, “Basis criteria for generalized spline modules via determinant”, preprint, 2019. arXiv
- [Altinok and Sarioglan 2019b] S. Altinok and S. Sarioglan, “Flow-up bases for generalized spline modules on arbitrary graphs”, preprint, 2019. arXiv
- [Billera and Rose 1991] L. J. Billera and L. L. Rose, “A dimension series for multivariate splines”, *Discrete Comput. Geom.* **6**:2 (1991), 107–128. MR Zbl
- [Bowden and Tymoczko 2015] N. Bowden and J. Tymoczko, “Splines mod m ”, preprint, 2015. arXiv
- [Bowden et al. 2015] N. Bowden, S. Hagen, M. King, and S. Reinders, “Bases and structure constants of generalized splines with integer coefficients on cycles”, preprint, 2015. arXiv
- [Chui and Lai 1990] C. K. Chui and M. J. Lai, “Multivariate vertex splines and finite elements”, *J. Approx. Theory* **60**:3 (1990), 245–343. MR Zbl
- [DiPasquale 2017] M. DiPasquale, “Generalized splines and graphic arrangements”, *J. Algebraic Combin.* **45**:1 (2017), 171–189. MR Zbl
- [Gilbert et al. 2016] S. Gilbert, J. Tymoczko, and S. Viel, “Generalized splines on arbitrary graphs”, *Pacific J. Math.* **281**:2 (2016), 333–364. MR Zbl
- [Goresky et al. 1998] M. Goresky, R. Kottwitz, and R. MacPherson, “Equivariant cohomology, Koszul duality, and the localization theorem”, *Invent. Math.* **131**:1 (1998), 25–83. MR Zbl
- [Guillemin and Zara 2000] V. Guillemin and C. Zara, “Equivariant de Rham theory and graphs”, pp. 221–257 in *Surveys in differential geometry*, edited by S.-T. Yau, Surv. Differ. Geom. **7**, International Press, Somerville, MA, 2000. MR Zbl
- [Guillemin and Zara 2001a] V. Guillemin and C. Zara, “1-skeleta, Betti numbers, and equivariant cohomology”, *Duke Math. J.* **107**:2 (2001), 283–349. MR Zbl
- [Guillemin and Zara 2001b] V. Guillemin and C. Zara, “ G -actions on graphs”, *Internat. Math. Res. Notices* **10** (2001), 519–542. MR Zbl
- [Guillemin and Zara 2003] V. Guillemin and C. Zara, “The existence of generating families for the cohomology ring of a graph”, *Adv. Math.* **174**:1 (2003), 115–153. MR Zbl
- [Handschy et al. 2014] M. Handschy, J. Melnick, and S. Reinders, “Integer generalized splines on cycles”, preprint, 2014. arXiv
- [Knutson and Tao 2003] A. Knutson and T. Tao, “Puzzles and (equivariant) cohomology of Grassmannians”, *Duke Math. J.* **119**:2 (2003), 221–260. MR Zbl
- [Lai and Schumaker 2007] M.-J. Lai and L. L. Schumaker, *Spline functions on triangulations*, Encyclopedia of Mathematics and its Applications **110**, Cambridge University Press, 2007. MR Zbl

- [Payne 2006] S. Payne, “Equivariant Chow cohomology of toric varieties”, *Math. Res. Lett.* **13**:1 (2006), 29–41. MR Zbl
- [Philbin et al. 2017] M. Philbin, L. Swift, A. Tamaro, and D. Williams, “Splines over integer quotient rings”, preprint, 2017. arXiv
- [Strang 1973] G. Strang, “Piecewise polynomials and the finite element method”, *Bull. Amer. Math. Soc.* **79** (1973), 1128–1137. MR Zbl
- [Tymoczko 2016] J. Tymoczko, “Splines in geometry and topology”, *Comput. Aided Geom. Design* **45** (2016), 32–47. MR Zbl
- [Zhou and Lai 2013] T. Zhou and M.-J. Lai, “Scattered data interpolation by bivariate splines with higher approximation order”, *J. Comput. Appl. Math.* **242** (2013), 125–140. MR Zbl

Received August 8, 2018. Revised July 25, 2019.

KATIE ANDERS
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF TEXAS AT TYLER
TYLER, TX
UNITED STATES
kanders@uttyler.edu

ALISSA S. CRANS
DEPARTMENT OF MATHEMATICS
LOYOLA MARYMOUNT UNIVERSITY
LOS ANGELES, CA
UNITED STATES
acrans@lmu.edu

BRIANA FOSTER-GREENWOOD
DEPARTMENT OF MATHEMATICS AND STATISTICS
CALIFORNIA STATE POLYTECHNIC UNIVERSITY
POMONA, CA
UNITED STATES
brianaf@cpp.edu

BLAKE MELLOR
DEPARTMENT OF MATHEMATICS
LOYOLA MARYMOUNT UNIVERSITY
LOS ANGELES, CA
UNITED STATES
blake.mellor@lmu.edu

JULIANNA TYMOCZKO
DEPARTMENT OF MATHEMATICS AND STATISTICS
SMITH COLLEGE
NORTHAMPTON, MA
UNITED STATES
jtymoczko@smith.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 304 No. 2 February 2020

Graphs admitting only constant splines	385
KATIE ANDERS, ALISSA S. CRANS, BRIANA FOSTER-GREENWOOD, BLAKE MELLOR and JULIANNA TYMOCZKO	
Centers of disks in Riemannian manifolds	401
IGOR BELEGRADEK and MOHAMMAD GHOMI	
The geometry of the flex locus of a hypersurface	419
LAURENT BUSÉ, CARLOS D'ANDREA, MARTÍN SOMBRA and MARTIN WEIMANN	
Morse inequalities for Fourier components of Kohn–Rossi cohomology of CR covering manifolds with S^1 -action	439
RUNG-TZUNG HUANG and GUOKUAN SHAO	
On Seifert fibered spaces bounding definite manifolds	463
AHMAD ISSA and DUNCAN MCCOY	
Regularity of quotients of Drinfeld modular schemes	481
SATOSHI KONDO and SEIDAI YASUDA	
Sums of algebraic trace functions twisted by arithmetic functions	505
MAXIM KOROLEV and IGOR SHPARLINSKI	
Twisted calculus on affinoid algebras	523
BERNARD LE STUM and ADOLFO QUIRÓS	
Symplectic (-2) -spheres and the symplectomorphism group of small rational 4-manifolds	561
JUN LI and TIAN-JUN LI	
Addendum to the article Contact stationary Legendrian surfaces in S^5	607
YONG LUO	
The Hamiltonian dynamics of magnetic confinement in toroidal domains	613
GABRIEL MARTINS	
Gluing Bartnik extensions, continuity of the Bartnik mass, and the equivalence of definitions	629
STEPHEN MCCORMICK	
Decomposable Specht modules indexed by bihooks	655
LIRON SPEYER and LOUISE SUTTON	
On the global well-posedness of one-dimensional fluid models with nonlocal velocity	713
ZHUAN YE	
Hopf cyclic cohomology for noncompact G -manifolds with boundary	753
XIN ZHANG	



0030-8730(2020)304:2;1-R