

Pacific Journal of Mathematics

CENTERS OF DISKS IN RIEMANNIAN MANIFOLDS

IGOR BELEGRADEK AND MOHAMMAD GHOMI

Volume 304 No. 2

February 2020

CENTERS OF DISKS IN RIEMANNIAN MANIFOLDS

IGOR BELEGRADEK AND MOHAMMAD GHOMI

We prove the existence of a center, or continuous selection of a point, in the relative interior of C^1 embedded k -disks in Riemannian n -manifolds. If $k \leq 3$ the center can be made equivariant with respect to the isometries of the manifold, and under mild assumptions the same holds for $k = 4 = n$. By contrast, for every $n \geq k \geq 6$ there are examples where an equivariant center does not exist. The center can be chosen to agree with any of the classical centers defined on the set of convex compacta in the Euclidean space.

1. Introduction

There are several distinguished points inside a compact convex subset of the Euclidean space; see [Kaiser et al. 1991; Moszyńska 2006, Chapter 12; Schneider 2014, Section 5.4.1]. In this paper we investigate if there is a geometrically meaningful point inside every embedded disk in a Riemannian manifold. Any such point should depend continuously on the disk and be equivariant under isometries.

To set the stage let G be a subgroup of the isometry group of a smooth (i.e., C^∞) connected Riemannian manifold M , and $\mathcal{X}(M)$ be the space of compact connected C^1 embedded submanifolds of M equipped with the C^1 topology (see Section 3). For a subspace X of $\mathcal{X}(M)$, a *center* is a continuous map $\mathfrak{c} : X \rightarrow M$ such that $\mathfrak{c}(D)$ lies in the relative interior of D for every $D \in X$. If in addition X is G -invariant and $\mathfrak{c}$ is G -equivariant, we say that $\mathfrak{c}$ is *G -equivariant*, which simply means that $\mathfrak{c}(gD) = g\mathfrak{c}(D)$ for all $D \in X$, $g \in G$. We investigate the existence of $\mathfrak{c}$ for a given triple (M, G, X) .

The best-known examples of a center are the center of mass and the Steiner point, which we call the *classical centers*. They are $\text{Iso}(\mathbb{R}^n)$ -equivariant centers on the space of convex compacta in $\mathcal{X}(\mathbb{R}^n)$, where $\text{Iso}(\mathbb{R}^n)$ is the group of Euclidean isometries; see [Schneider 2014].

To state our findings, for any $D \in \mathcal{X}(M)$ let G_D be the subgroup of G which maps D to itself, and D^{G_D} be the points of D which are fixed by G_D (see Section 2).

Ghomi's research was supported in part by NSF grants DMS-1308777 and DMS-1711400.

MSC2010: primary 53C40, 54C65; secondary 52A20, 57S25.

Keywords: continuous selection, equivariant, proper actions, actions on disks.

Note that if $\mathfrak{c}$ is G -equivariant, then $\mathfrak{c}(D) \in D^{G_D}$, and in particular, no G -equivariant center exists if $D^{G_D} = \emptyset$ for some $D \in X$. Our main result is as follows:

Theorem 1.1. *The space of submanifolds $D \in \mathcal{X}(M)$ such that D^{G_D} is contractible admits a G -equivariant center. Furthermore, this center can be chosen to agree with any given G -equivariant center defined on a closed subset A of $\mathcal{X}(M)$.*

An important example of A is the set of convex compacta in $\mathbb{R}^n$ equipped with one of the classical centers mentioned above. In this case our results are summarized in Corollary 1.7 below.

We prove Theorem 1.1 by reducing the problem to finding sections of certain bundles with D^{G_D} as fibers, and then fitting the sections together to define a global center. The construction involves some choices, and the resulting center is not canonical.

Let us review some natural conditions under which D^{G_D} is contractible, which yield the following corollaries of Theorem 1.1. It is to be understood that *each of these corollaries has a relative version*, i.e., the corresponding center can be chosen to agree with any given G -equivariant center defined on a closed subset of $\mathcal{X}(M)$. First note that if G is trivial, then $D^{G_D} = D$. So we obtain:

Corollary 1.2. *The space of contractible submanifolds $D \in \mathcal{X}(M)$ admits a center.*

Another condition that ensures contractibility of D^{G_D} is that each $D \in \mathcal{X}(M)$ is homeomorphic to a disk of dimension ≤ 3 . This is due to the fact that any smooth action of a compact Lie group on a disk of dimension ≤ 3 is smoothly equivalent to a linear action, which implies that D^{G_D} is a disk. For actions on 2-disks linearity is a standard consequence of the uniformization theorem (see the beginning of the Appendix), while the 3-dimensional case is established in [Kwasik and Schultz 1992, Theorem B].

Let $\mathfrak{D}^k(M)$ be the subspace of $D \in \mathcal{X}(M)$ such that D is homeomorphic to a disk of dimension $\leq k$. From the previous paragraph we obtain:

Corollary 1.3. *$\mathfrak{D}^3(M)$ admits a G -equivariant center.*

On the other hand, contractibility of D^{G_D} generally fails when D is homeomorphic to a 4-disk, because then D^{G_D} may be an acyclic noncontractible 3-manifold, see Lemma A.4. For this case to occur G_D must be generated by an involution whose action on D reverses the orientation; see Lemmas A.1 and A.3. Avoiding this case, we obtain the existence of centers in dimension 4:

Corollary 1.4. *If $\dim(M) = 4$ and the G -action on M is orientation-preserving, then $\mathfrak{D}^4(M)$ admits a G -equivariant center.*

Another result for 4-manifolds can be obtained by observing that no compact acyclic noncontractible 3-manifold smoothly embeds into S^3 due to the topological Schoenflies theorem [Brown 1960]. In Lemma 4.2 we note that D^{G_D} is a C^1

submanifold of M^{G_D} , the fixed point set of G_D in M . Therefore, a noncontractible D^{G_D} cannot occur if for any involution in G its fixed point set in M smoothly embeds into S^3 . The latter can be forced by the following geometric assumptions.

Corollary 1.5. *If either $M = S^4$ with its standard action of $G = O(5)$, or M is a Hadamard 4-manifold with isometry group G , then $\mathfrak{D}^4(M)$ admits a G -equivariant center.*

Recall that a *Hadamard manifold* is a contractible complete Riemannian manifold of nonpositive sectional curvature, and the fixed point set of any isometric action on a Hadamard manifold is diffeomorphic to a Euclidean space, due to the Cartan–Hadamard theorem.

The above 4-dimensional results depend on establishing that the fixed point set of a smooth compact Lie group action on the 4-disk is either a disk or a compact acyclic 3-manifold. We check this in Lemmas A.1 and A.3. A key ingredient is that any smooth finite group action on the 4-disk fixes a point. This was established in [Buchdahl et al. 1990, Theorem II.1] modulo an announcement of Thurston that any nonfree smooth finite group action on S^3 preserves a round metric. Thurston’s claim was finally proved in [Dinkelbach and Leeb 2009] via a Ricci flow arguments, and his outline was made rigorous in [Boileau et al. 2005, Corollary 1.1] for orientation-preserving actions.

In dimension 5 the situation is unclear, e.g., it does not seem to be known whether every smooth action of a compact Lie group on a 5-disk has a fixed point. The strategy used in Corollary 1.5 for linear actions on $\mathbb{R}^4$ breaks down one dimension up, because there is a linear involution of $\mathbb{R}^5$ whose fixed point set on some embedded disk is not contractible, see Lemma A.5.

Starting from dimension 6 an equivariant center need not exist, namely, there are triples (M, X, G) such that $D^{G_D} = \emptyset$ for some smoothly embedded disk $D \in X$, which rules out the existence of a G -equivariant center, as we had mentioned earlier. Indeed, for each $n \geq 6$ there is a smooth action of the alternating group A_5 on S^n with exactly one fixed point; see, e.g., [Bak and Morimoto 2005], and excising an invariant tubular neighborhood of this point yields a fixed point free smooth action of A_5 on the n -disk. One can then embed the resulting A_5 -action on the n -disk into a fixed point free A_5 -action on a boundaryless manifold, e.g., into $\mathbb{R}^n$ by attaching a collar along the boundary, or into S^n by doubling along the boundary.

The Mostow–Palais theorem [Bredon 1972, Theorem VI.4.1] yields a smooth embedding of the above fixed point free A_5 -action on the 6-disk into an orthogonal A_5 -action on some $\mathbb{R}^r$. (With some effort one can obtain an explicit upper bound for r but we will not attempt it here because we are unable to determine the optimal r .) As we shall explain at the end of Section 4, this discussion easily implies the following:

Proposition 1.6. *For any m, n such that either $m \geq n + r - 6 \geq r$ or $m \geq n \geq r$ there is an orthogonal A_5 -action on $\mathbb{R}^m$ that preserves a smoothly embedded n -disk on which A_5 acts without a fixed point. In particular, $\mathfrak{D}^n(\mathbb{R}^m) \setminus \mathfrak{D}^{n-1}(\mathbb{R}^m)$ does not admit an $O(m)$ -equivariant center.*

At this point it seems worthwhile to summarize our results for the Euclidean space:

Corollary 1.7. *Any $\text{Iso}(\mathbb{R}^n)$ -equivariant center on the set of convex compacta in $\mathbb{R}^n$ extends to a center $\mathfrak{c} : \mathfrak{D}^n(\mathbb{R}^n) \rightarrow \mathbb{R}^n$ that is $\text{Iso}(\mathbb{R}^n)$ -equivariant on $\mathfrak{D}^3(\mathbb{R}^n)$. Moreover, if $n = 4$, then $\mathfrak{c}$ is $\text{Iso}(\mathbb{R}^4)$ -equivariant on $\mathfrak{D}^4(\mathbb{R}^4)$. If k is sufficiently large, then $\mathfrak{D}^k(\mathbb{R}^n)$ does not admit an $\text{Iso}(\mathbb{R}^n)$ -equivariant center.*

The results of this paper belong to the subject of *continuous selections of multivalued mappings*, an established branch of topology which was pioneered by Michael [1956a; 1956b; 1959] and extensively surveyed in [Repovš and Semenov 1998; 2002; 2014]. Indeed, assigning to each $D \in \mathcal{X}(M)$ the relative interior of D^{G_D} in M yields a multivalued map $\mathcal{X}(M) \rightarrow M$ which is lower semicontinuous in the sense of [Repovš and Semenov 1998, Definition 0.43]. What we call a G -equivariant center is a continuous G -equivariant selection of this multivalued map over a subset X of $\mathcal{X}(M)$. The theory of continuous selections implies existence of a nonequivariant center under certain assumptions on $X \subset \mathcal{X}(M)$ that tend to come in two flavors: either every $D \in X$ needs to satisfy a suitable generalized convexity condition, or X is required to be finite-dimensional (see [Repovš and Semenov 1998] for details). By contrast, our focus is on equivariant centers, and there seems to be no prior work analogous to the results of this paper.

The classical centers are continuous in the Hausdorff topology on the set of convex compacta, and hence one might expect that this would be a natural topology for $\mathcal{X}(M)$. However, in Remark 3.4 we shall see that Theorem 1.1 fails when $\mathcal{X}(\mathbb{R}^n)$ is given the Hausdorff topology. On the other hand, for any k the Hausdorff topology on the set of k -dimensional convex compacta in $\mathcal{X}(\mathbb{R}^n)$ coincides with the C^0 topology, and hence we ask:

Question 1.8. Does Theorem 1.1 remain valid when $\mathcal{X}(M)$ is replaced with the space of C^0 submanifolds equipped with the C^0 topology?

2. Background on group actions

Here we review the basic facts on Lie group actions used in this work. Throughout this section G is a Lie group and X is a metrizable space. A G -action on X is a continuous map $a : G \times X \rightarrow G$, written as $gx := a(g, x)$, such that the map $g \rightarrow a(g, \cdot)$ is a homomorphism between G and the homeomorphism group of X . A space with a G -action is a G -space. The orbit space X/G is the set of G -orbits

with the quotient topology. A map $f : X \rightarrow Y$ of G -spaces is a G -map if f is continuous and $f(gx) = gf(x)$ for all $x \in X$, and $g \in G$; we will also refer to f as G -equivariant or just *equivariant* when G is understood. For $S \subseteq X$, we let $gS := \{gx : x \in S\}$ and use the following notations:

$G_S := \{g \in G : gS = S\}$ = the isotropy subgroup of S in G ,

$GS := \{gS : g \in G\}$ = the G -orbit of S ,

$X^G := \{x \in X : gx = x \text{ for each } g \in G\}$ = the fixed point set of G in X .

If $S = \{x\}$, then GS , G_S are denoted by Gx , G_x , respectively. Note that G_x is a closed subgroup of G . If $GS = S$, then S is G -invariant. If $x \in S_x \subseteq X$, then S_x is called a G_x -slice at x provided that GS_x is open in X and there is a G -map $f : GS_x \rightarrow G/G_x$ with $S_x = f^{-1}(G_x)$.

Lemma 2.1. *Let $\alpha : A \rightarrow X$ be a G -map of G -spaces with $\alpha(a) = x$ and $G_a = G_x$. If S_x is a G_x -slice at x , then $\alpha^{-1}(S_x)$ is a G_a -slice at a .*

Proof. Note that

$$\alpha \circ f : A \rightarrow G/G_x = G/G_a$$

is a G -map with $(\alpha \circ f)^{-1}(G_a) = \alpha^{-1}(S_x)$. For any $g \in G$, $z \in X$, we have $g\alpha^{-1}(z) = \alpha^{-1}(gz)$, and hence $G\alpha^{-1}(S_x) = \alpha^{-1}(GS_x)$. Thus openness of GS_x implies openness of $G\alpha^{-1}(S_x)$. $\square$

A G -space X is *Palais-proper* if any $x \in X$ has a neighborhood V_x such that every $y \in Y$ has a neighborhood V_y for which $\{g \in G : gV_x \cap V_y \neq \emptyset\}$ is precompact in G . For example, if G is compact, then any G -space is Palais-proper. If G is an isometry group of a smooth Riemannian manifold, then the G -space is Palais-proper [Kobayashi and Nomizu 1963, Theorem I.4.7], and conversely, any smooth Palais-proper action preserves a smooth Riemannian metric [Palais 1961, Theorem 4.3.1].

Note that if x is a point of a Palais-proper G -space X , then G_x is compact (because it is closed and precompact in G). A key result established in [Palais 1961, Section 2.3] is that every point x in a Palais-proper G -space is contained in a G_x -slice. In [Palais 1961, Section 2.1] one finds the following characterization of slices at points with compact isotropy subgroups:

Lemma 2.2 [Palais 1961]. *Let $x \in S \subseteq X$ and suppose that G_x is compact. Then S is a G_x -slice at x if and only if the following conditions hold:*

- (a) $gS \cap S \neq \emptyset$ if and only if $g \in G_x$.
- (b) S is closed in GS , and GS is open in X .
- (c) There is an open set O such that $S \subset O \subset GS$ and $\{g \in G : gO \cap O \neq \emptyset\}$ is precompact in G . $\square$

The following lemma summarizes what we need to know about slices.

Lemma 2.3. *If S_x is a G_x -slice at x and G_x is compact, then the following statements hold:*

- (i) $G_y \subset G_x$ for every $y \in S_x$.
- (ii) If W is open in a slice S_x , then GW is open in X .
- (iii) The inclusion induced map $S_x/G_x \rightarrow X/G$ is an open embedding.
- (iv) Any open G_x -invariant neighborhood of x in S_x is a G_x -slice at x .
- (v) Any neighborhood of x in S_x contains an open set that is a G_x -slice at x .
- (vi) For any neighborhood U of the identity in G there is a neighborhood V of x in GS such that for each $y \in V$ there is $u \in U$ with $u^{-1}G_y u \subseteq G_x$.

Proof. Statement (i) is immediate by Lemma 2.2(a), while (ii) is proved in [Palais 1961, Corollary on p.306]. The rest of the items are established as follows:

(iii): The map here is one-to-one by Lemma 2.2(a). Further, it is a homeomorphism by (ii) and the defining properties of the quotient topology.

(iv): Let W be an open G_x -invariant neighborhood of x in S_x , and appeal to Lemma 2.2. The conditions (a) and (c) are immediate, and (i) implies that GW is open in X . To see that W is closed in GW take $w_i \rightarrow gw$, where $g \in G$, $w_i, w \in W$, and note that $w, w_i \in S_x$ implies $gw \in S_x$, so $g \in G_x$ and hence $gw \in W$.

(v): Note that S_x has a G_x -invariant metric [Palais 1960, Proposition 1.1.12], so any open metric ball centered at x is G_x -invariant, and hence is a slice by (iii). Any neighborhood of x contains such a ball.

(vi): Choose V inside O of Lemma 2.2(c). Then G_x, G_y lie in a compact subgroup of G , in which case a proof can be found in [Bredon 1972, Corollary II.5.5]. $\square$

3. Space of submanifolds

In this section we give a precise definition of the topology on $\mathcal{X}(M)$. Further we show that this topology is Palais-proper and induced by a G -invariant metric. For any submanifold $D \in \mathcal{X}(M)$ let $\mathcal{X}_D(M) \subset \mathcal{X}(M)$ be the collection of submanifolds which are C^1 diffeomorphic to D . In other words,

$$\mathcal{X}_D(M) := \text{Emb}^1(D, M) / \text{Diff}^1(D),$$

the space of C^1 embeddings of D into M modulo C^1 diffeomorphisms of D . We equip $\mathcal{X}_D(M)$ with its standard C^1 topology, which is induced by $C^1(D, M)$, the space of C^1 mappings $D \rightarrow M$. Thus a pair of submanifolds $A, B \in \mathcal{X}_D(M)$ are close if they admit parametrizations $f, g \in \text{Emb}^1(D, M)$ that are C^1 close. Finally we topologize $\mathcal{X}(M)$ as the disjoint union of $\mathcal{X}_D(M)$ where D ranges over C^1 diffeomorphism classes of submanifolds $D \in \mathcal{X}(M)$. Note that the obvious G -action on $\mathcal{X}(M)$ given by $\alpha(g, D) = gD$ is effective.

Lemma 3.1. *The following statements hold:*

- (1) *The topology on $\mathcal{X}_D(M)$ is induced by a G -invariant metric d_D .*
- (2) *The G -space $\mathcal{X}_D(M)$ is Palais-proper.*
- (3) *The orbit space $\mathcal{X}_D(M)/G$ is metrizable.*

Proof. A convenient way to describe $\mathcal{X}_D(M)$ is to consider a smooth G -equivariant embedding of M into a Hilbert space $\mathcal{H}$ equipped with some orthogonal G -action; see [Kankaanrinta 1994, Theorem 0.1]. In general, one cannot equivariantly embed M into a finite-dimensional linear space; this can be done, e.g., if both G and M are compact [Bredon 1972, Theorem VI.4.1].

To prove (1) we define a G -invariant metric d_D on $\mathcal{X}_D(\mathcal{H})$, and then restrict it to a smoothly embedded G -invariant copy of M in $\mathcal{H}$. Define $d_D(A, B)$ as the infimum of $\|\alpha - \beta\|_{C^1}$ taken over all C^1 embeddings $\alpha, \beta \in C^1(D, \mathcal{H})$ with images A, B , respectively. Here the C^1 norm is computed using the norm on $\mathcal{H}$. The triangle inequality follows from the one for $\mathcal{H}$ and properties of the infimum, and the G -invariance holds because the G -action on $\mathcal{H}$ preserves the norm. For nondegeneracy note that if say $a \in A \setminus B$, then $d_D(A, B)$ is bounded below by the distance from a to B , hence $d_D(A, B) = 0$ implies $A = B$.

To prove (2) fix $r \in (0, 1)$ and let V_x be the r -ball about x in $(\mathcal{X}_D(M), d)$. If $A, B \in \mathcal{X}_D(M)$ with $V_A \cap gV_B \neq \emptyset$, then gB lies in the $3r$ -neighborhood of A , and since the G -action on M is Palais-proper, so is the G -action on $\mathcal{X}(M)$.

Statement (3) is proved in [Palais 1961, Theorem 4.3.4] assuming (1)–(2) with the metric given by $\bar{d}_D(Gx, Gy) = \inf\{d_D(x, gy) : g \in G\}$. $\square$

Remark 3.2. The proof of Lemma 3.1 works for $\mathcal{X}(M)$ in place of $\mathcal{X}_D(M)$ by replacing d_D with $d(A, B) = \min(1, d_D(A, B))$ if $A, B \in \mathcal{X}_D(M)$ for some D , and $d(A, B) = 1$ otherwise.

Remark 3.3. The proof of Lemma 3.1 goes through as written if we give $\mathcal{X}(M)$ the C^k topology, where k is any nonnegative integer.

Remark 3.4. Let G be any group of isometries of $\mathbb{R}^2$ that contains a reflection r in the y -axis, as well as a (nontrivial) rotation about the origin. Let us justify the claim made before Question 1.8 that there is no Hausdorff continuous G -equivariant center defined on the set of 2-disks in $\mathcal{X}(\mathbb{R}^2)$. The unit disk $D^2 \subset \mathbb{R}^2$ is Hausdorff close to a smoothly embedded 2-disk D obtained by removing from D^2 a small r -invariant neighborhood of the segment $\{(0, y) \in D^2 : y \geq -\frac{1}{2}\}$. Since G contains r , any G -equivariant center of D is contained in $\{(0, y) \in D^2 : y < -\frac{1}{2}\}$, while the G -invariant center of D^2 is the origin because G contains a rotation.

4. Proof of Theorem 1.1

In this section M , G are as in Section 1. Thus M^G is a smooth submanifold of M , and if D is a compact subset of M , then G_D is a compact subgroup of G . Also $D^{G_D} = D \cap M^{G_D}$. By *small* we mean close to the identity.

Lemma 4.1. *Any $D \in \mathcal{X}(M)$ has a compatible smooth atlas α such that the given G_D -action on (D, α) can be C^1 approximated by a smooth G_D -action, and moreover, these two actions are conjugate by a C^1 diffeomorphism that is C^1 close to the identity.*

Proof. The existence of a compatible smooth atlas is proved in [Hirsch 1976, Theorem 3.6, Chapter 2]. If D is a closed manifold, the other claims are contained in [Palais 1970, Theorem C]. If D has boundary, we first double the given C^1 actions of G_D along ∂D . To equivariantly smooth the double at ∂D we may have to adjust the action, conjugating it by a small C^1 isotopy near ∂D . Then we apply [Palais 1970, Theorem C] to the double to approximate the C^1 action by a C^1 equivalent smooth action. This action stabilizes a copy of D that is C^1 close to D , so after a conjugation by a small smooth diffeomorphism it can be made to stabilize D inside the double. $\square$

Lemma 4.2. *Let $D \in \mathcal{X}(M)$ and suppose $D^{G_D} \neq \emptyset$. Then D^{G_D} is a properly embedded C^1 submanifold of D that intersects the relative interior of D and is transverse to ∂D .*

Proof. In the compatible atlas of Lemma 4.1 the G_D -action on D is near a smooth action, and the two actions are conjugate by a small C^1 diffeomorphism. Let ϕ denote the smooth G_D -action on the double of D , and fix a Riemannian metric invariant under ϕ . Its fixed point set is a smooth compact boundaryless submanifold which is totally geodesic in the metric. Since ∂D is ϕ -invariant if ϕ fixes $x \in \partial D$, then it also fixes every point in a convex neighborhood of x that lies on the geodesic through x that is orthogonal to ∂D . By assumption G_D fixes a point of D , and hence so does ϕ . It follows that the fixed point of ϕ intersects the relative interior of D and is transverse to ∂D . Hence the same is true for the given G_D action on the double, and the claim follows. $\square$

Remark 4.3. Lemma 4.2 is immediate by transversality if $\dim(D) = \dim(M)$. In general, the intersection of ∂D and D^{G_D} is not transverse, e.g., a round disk D in the xy -plane in $\mathbb{R}^3$ is invariant under rotation by π about the x -axis, which would be disjoint from ∂D if they met transversely.

We partition $\mathcal{X}(M)$ as $\bigcup_{k,l} \mathcal{D}^{k,l}$, where $\mathcal{D}^{k,l}(M)$ is the set of all $D \in \mathcal{X}(M)$ such that D^{G_D} is a k -dimensional manifold with l connected components.

Remark 4.4. Even when D is a disk, the manifold D^{G_D} may have any finite number of components, e.g., if G is connected and nonabelian and F is any finite

CW complex, then there is a smooth G -action on some high-dimensional disk D such that D^{G_D} is homotopy equivalent to F ; see [Oliver 1976, Theorems 3 and 5]. On the other hand, we shall see in the Appendix that if D is a disk of dimension ≤ 4 , then D^{G_D} is acyclic, and in particular, connected.

Henceforth, we need the following notion of a tubular neighborhood of a C^1 submanifold N of a Riemannian manifold M . By [Hirsch 1976, Theorem 3.6, Chapter 2] there is a C^1 diffeomorphism $\phi : M \rightarrow M$ such that $\tilde{N} := \phi(N)$ is a C^∞ submanifold of M . If $\tilde{U}$ is a Riemannian tubular neighborhood of $\tilde{N}$ (produced via the normal exponential map), then we call $U := \phi^{-1}(\tilde{U})$ a *tubular neighborhood* of N , and furthermore, if $\tilde{\pi} : \tilde{U} \rightarrow \tilde{N}$ is the nearest point projection, we call $\pi := \phi^{-1} \circ \tilde{\pi} \circ \phi : U \rightarrow N$ the *normal bundle projection* of U .

Lemma 4.5. *The map $\mathcal{D}^{k,l}(M) \rightarrow \mathcal{X}(M)$ given by $D \rightarrow D^{G_D}$ is continuous.*

Proof. Fix any $J \in \mathcal{D}^{k,l}(M)$ and prove continuity at J . By Lemma 2.3(vi) there is a neighborhood U of J in $\mathcal{D}^{k,l}(M)$ such that G_D can be conjugated into G_J by a small element of G , and since such conjugation results in only small C^1 changes of D and D^{G_D} we may assume that $G_D \leq G_J$. Thus both D and J are invariant under $H = G_D$.

After possibly shrinking U further we can find a C^1 diffeomorphism ψ with $\psi(D) = J$ such that ψ is supported in a small neighborhood of J . (To find ψ we first isotope ∂D to ∂J in the tubular neighborhood ∂J , then extend J to a boundaryless embedded C^1 submanifold, use its tubular neighborhood to isotope D to J relative boundary, and finally extend the isotopy to an ambient one.) Thus $\psi H \psi^{-1}$ and H are two C^1 close H -actions on J . By Lemma 4.1 these actions are C^1 diffeomorphic via a small diffeomorphism.

By Lemma 4.2 the fixed point sets of the actions of $\psi H \psi^{-1}$, H , G_J on J are compact properly embedded k -dimensional C^1 submanifolds of J with l connected components. Any proper embedding of compact k -manifolds with the same number of components is surjective, so since $H \leq G_J$ the fixed point sets of H , G_J in J coincide. Thus J^{G_J} is C^1 close to the fixed point set of $\psi H \psi^{-1}$ which is D^{G_D} . $\square$

Lemma 4.6. *If B is a compact C^1 submanifold of a manifold F , then there are two nested tubular neighborhoods $T \subset T_2$ of ∂B in F and a C^1 self-map q_B of F that is the identity outside T_2 , and that equals the normal bundle projection on T .*

Proof. Fix a tubular neighborhood T_2 of ∂B in F , and identify it with $[-2, 2] \times \partial B$ where ∂B corresponds to $\{0\} \times \partial B$. Let T be the subset of T_2 corresponding under the identification to $[-1, 1] \times \partial B$. Fix a smooth nondecreasing self-map τ of $[-2, 2]$ that vanishes on $[-1, 1]$ and equals the identity near the endpoints $-2, 2$. Then the self-map of $[-2, 2] \times \partial B$ given by $(t, z) \rightarrow (\tau(t), z)$ interpolates between the identity near the boundary and the projection $[-1, 1] \times \partial B \rightarrow \{0\} \times \partial B$. $\square$

Let $\mathcal{E}^{k,l}(M)$ be the subspace of $\mathcal{D}^{k,l}(M) \times M$ consisting of all the pairs (D, u) with $D \in \mathcal{D}^{k,l}(M)$ and $u \in D^{G_D}$. Let $\pi : \mathcal{E}^{k,l}(M) \rightarrow \mathcal{D}^{k,l}(M)$ be the coordinate projection, i.e., $\pi(D, u) = D$.

Lemma 4.7. *The map $\pi : \mathcal{E}^{k,l}(M) \rightarrow \mathcal{D}^{k,l}(M)$ is a locally trivial bundle.*

Proof. Fix $J \in \mathcal{D}^{k,l}(M)$, set $B = J^{G_J}$, and extend B to a k -dimensional boundaryless C^1 submanifold F of M . Let q_F be a projection of a tubular neighborhood of F in M . For such B, F let q_B and T be as in Lemma 4.6. Let $q = q_B \circ q_F$. Using Lemma 4.5 we can assume that D is so close to J that $q_F(D^{G_D}) \subset T$ and q_B restricts to a C^1 diffeomorphism of $q_F(D^{G_D})$ onto B . The map $(D, u) \rightarrow (D, q(u))$ is the desired local trivialization where $u \in D^{G_D}$. $\square$

Remark 4.8. The homeomorphism type of the fiber $\pi^{-1}(D)$ may depend on D .

The group G acts on $\mathcal{E}^{k,l}(M)$ diagonally, i.e., $g(D, u) = (gD, gu)$. In the following commutative diagram

$$(4.9) \quad \begin{array}{ccc} \mathcal{E}^{k,l}(M) & \xrightarrow{\pi} & \mathcal{D}^{k,l}(M) \\ p \downarrow & & \downarrow \bar{p} \\ \mathcal{E}^{k,l}(M)/G & \xrightarrow{\bar{\pi}} & \mathcal{D}^{k,l}(M)/G \end{array}$$

the vertical arrows are the G -orbit maps, and $\bar{\pi}$ sends the G -orbit of (D, u) to the G -orbit of D .

By Remark 3.2 the G -spaces $\mathcal{D}^{k,l}(M)$ and $\mathcal{E}^{k,l}(M)$ are Palais-proper because the property is clearly inherited by invariant subspaces and preimages under equivariant maps. Also $\mathcal{E}^{k,l}(M)$ has a G -invariant metric induced by the G -invariant metrics on the factors of $\mathcal{D}^{k,l}(M) \times M$. The following result is a key observation of this paper, which reduces to Lemma 4.7 when G is trivial.

Lemma 4.10. *$\bar{\pi} : \mathcal{E}^{k,l}(M)/G \rightarrow \mathcal{D}^{k,l}(M)/G$ is a locally trivial fiber bundle.*

Proof. Let us first sketch the proof. The G -action permutes the fibers of π which are of the form $\pi^{-1}(y) = \{(y, u) : u \in y^{G_y}\}$. Moreover, each fiber projects homeomorphically to $\mathcal{E}^{k,l}(M)/G$ because the fiber is compact and the projection p is injective on it: If $g(y, u) = (y, v)$ with $u, v \in y^{G_y}$, then $g \in G_y$ and $gu = v$. Hence $gu = u$ implies $u = v$. To establish local triviality we analyze the structure of the orbit spaces via local slices. Using that $G_y \leq G_x$ for every y in a G_x -slice we show that the local trivialization of Lemma 4.7 can be made G_x -equivariant, so it passes to the G_x -quotients of the slices, yielding a local trivialization of $\bar{\pi}$.

Let us make this sketch rigorous. Set $X = \mathcal{D}^{k,l}(M)$. Fix $x \in X$ and set $H = G_x$. Let S be a H -slice at x . Thus G fixes $\pi^{-1}(x)$ pointwise, and hence, Lemma 2.1 implies that $E = \pi^{-1}(S)$ is an H -slice at any point of $\pi^{-1}(x)$.

Set $\bar{E} = p(E)$ and $\bar{S} = \bar{p}(S)$. Since S is a slice, $\bar{S}$ is a neighborhood of $\bar{p}(x)$, and it suffices to show that $\bar{\pi}$ is a locally trivial bundle over $\bar{S}$.

Let us show that $\bar{E} = \bar{\pi}^{-1}(\bar{S})$. One inclusion follows from a diagram chase: If $z \in p(\pi^{-1}(S))$, then $\bar{\pi}(z) \in (\bar{\pi} \circ p)(\pi^{-1}(S)) = (\bar{p} \circ \pi)(\pi^{-1}(S)) = p(S)$. Conversely, if $z \in \bar{\pi}^{-1}(\bar{S})$, then $\bar{\pi}(z) = Gy$ for some $y \in S$. Also z is the G -orbit of some (a, v) with $a \in X$, $v \in a^{G_a}$, and hence $\bar{\pi}(z)$ is the G -orbit of $\pi(a, v) = a$. Thus $a = gy$ for some $g \in G$. Hence z is the G -orbit of $(y, g^{-1}v)$, and moreover, $g^{-1}v \in y^{G_y}$. Thus $(y, g^{-1}v) \in E$, and so $z \in \bar{E}$.

Since E and S are H -slices, the inclusion induced maps $E/H \rightarrow \bar{E}$, $S/H \rightarrow \bar{S}$ are homeomorphisms, so we can identify the map $\bar{\pi} : \bar{E} \rightarrow \bar{S}$ with the map $E/H \rightarrow S/H$ also induced by π .

Set $B = x^H$ and note that B is contained in M^H , which is a boundaryless properly embedded smooth submanifold of M . We can extend B to a k -dimensional boundaryless C^1 submanifold F of M^H . Fix an H -invariant tubular neighborhood of F in M , and let q_F be an H -equivariant projection of the neighborhood onto M (obtained, e.g., as the nearest point projection of some smooth H -invariant Riemannian metric on M).

For such B , F let q_B and T be as in Lemma 4.6. Now Lemma 4.5 implies that by making S smaller we can assume that any $y \in S$ satisfies $q_F(y^{G_y}) \subset T$ and q_B restricts to a C^1 diffeomorphism of $q_F(y^{G_y})$ onto B . (Lemma 2.3(v) allows us to make S arbitrarily small.) Then for each $y \in S$ the composite $q = q_B \circ q_F$ restricts to a C^1 -diffeomorphism of y^{G_y} onto B .

For every $y \in S$ we have $G_y \leq H$, and hence $M^{G_y} \supseteq M^H \supseteq F$. In particular, if $z \in y^{G_y}$ and $y \in S$, then $q(hz) = q(z)$ because q is H -equivariant and H acts trivially on the image of q .

The map $\phi : E \rightarrow S \times B$ given by $(y, u) \rightarrow (y, q(u))$ is H -equivariant because it sends $h(y, u) = (hy, hu)$ to $(hy, q(hu)) = (hy, q(u)) = (hy, hq(u)) = h(y, q(u))$. Also ϕ is a homeomorphism whose inverse sends (y, v) to $(y, (q|_{y^{G_y}})^{-1}(v))$. Note that H acts trivially on the B -factor. So ϕ descends to a homeomorphism of the H -quotients $E/H \rightarrow S/H \times B$ which gives a desired local trivialization. $\square$

Let $\mathcal{D}^k(M)$ be the subspace of $D \in \mathcal{X}(M)$ such that D^{G_D} is contractible and k -dimensional. Set $\mathcal{E}^k(M) = \{(D, u) : D \in \mathcal{D}^k(M), u \in D^{G_D}\}$. Note that $\mathcal{D}^k(M)$, $\mathcal{E}^k(M)$ are G -invariant subsets of $\mathcal{D}^{k,1}(M)$, $\mathcal{E}^{k,1}(M)$, respectively, and moreover, $\mathcal{E}^k(M)$ is the π -preimage of $\mathcal{D}^k(M)$. Thus $\pi : \mathcal{E}^k(M) \rightarrow \mathcal{D}^k(M)$ and $\bar{\pi} : \mathcal{E}^k(M)/G \rightarrow \mathcal{D}^k(M)/G$ are locally trivial fiber bundles.

Lemma 4.11. *If A is a closed G -invariant subset of $\mathcal{D}^k(M)$, then every continuous G -equivariant section $s_A : A \rightarrow \mathcal{E}^k(M)$ of π over A extends to a continuous G -equivariant section s of π over $\mathcal{D}^k(M)$. Moreover, if s_A takes values in the interiors of the fibers, then one can choose s with the same property.*

Proof. By G -equivariance, s_A descends to a continuous map $\bar{s}_A : A/G \rightarrow \mathcal{E}^{k,l}(M)/G$ that takes the G -orbit of $x \in A$ to the G -orbit of $s(x)$, i.e., $\bar{s}_A(\bar{p}(x)) = p(s_A(x))$. The following section extension property can be found in [Palais 1966, Theorem 9]: Given any locally trivial fiber bundle whose base is metrizable and fiber is an absolute retract, any section of the bundle defined on a closed subset can be extended to the whole base. In our case the fiber is either a compact contractible manifold with boundary, or its interior, which are absolute retracts.

By definition of the quotient topology, A/G is closed in $\mathcal{D}^{k,l}(M)/G$. Thus $\bar{s}_A$ extends to a continuous section $\bar{s}$ of $\bar{\pi}$. If s_A takes values in the interiors of the fibers, then so does $\bar{s}_A$, and hence it can be extended to $\bar{s}$ with the same property. Now define $s : \mathcal{D}^{k,l}(M) \rightarrow \mathcal{E}^{k,l}(M)$ by letting $s(x)$ be the intersection of the fiber $\pi^{-1}(x)$ and the G -orbit $p^{-1}(\bar{s}(\bar{p}(x)))$; see diagram (4.9). Let us check that s has the claimed properties.

The intersection consists of a single point, indeed, if z, gz are in the intersection, then $\pi(z) = x = \pi(gz) = g\pi(z)$ and hence g is in the isotropy subgroup of z , which fixes $\pi^{-1}(\pi(z))$ pointwise so that $gz = z$. If $a \in A$, then $s(a)$ and $s_A(a)$ are points of $\pi^{-1}(a)$ that are mapped by p to $\bar{s}(\bar{p}(a)) = \bar{s}_A(\bar{p}(a))$, and since such a point is unique, s extends s_A .

By construction $\pi(s(x)) \subseteq \pi(\pi^{-1}(x)) = \{x\}$, so s is a section. The map s is G -equivariant because $s(gx)$ is the intersection of $\pi^{-1}(gx) = g\pi^{-1}(x)$ and $p^{-1}(\bar{s}(\bar{p}(gx))) = p^{-1}(\bar{s}(\bar{p}(x))) = g(p^{-1}(\bar{s}(\bar{p}(x))))$, which equals $gs(x)$.

We write $u \approx v$ to indicate that u, v are close to each other. To prove continuity of s write $s(x) = (x, c)$ and $s(y) = (y, d)$, where $c \in x, d \in y$, assume $x \approx y$, and try to show that $c \approx d$. From $x \approx y$ we get closeness of the G -orbits $p^{-1}(\bar{s}(\bar{p}(x)))$, $p^{-1}(\bar{s}(\bar{p}(y)))$ of $s(x), s(y)$, respectively. Hence there is $g \in G$ such that $gs(x) \approx s(y)$, or equivalently, $gx \approx y$ and $gc \approx d$. From $x \approx y$ we get $gx \approx gy$, and hence $x \approx y \approx gx \approx gy$. Then g is close to some elements of the isotropy subgroups of x, y , which fix the fibers $\pi^{-1}(x), \pi^{-1}(y)$ pointwise. Hence $c \approx gc$ and $d \approx gd$ which together with $gc \approx d$ implies $c \approx d$. $\square$

The above proof also yields the following observation:

Lemma 4.12. *If A is a closed G -invariant subset of $\mathcal{D}^k(M)$, every G -equivariant section $\bar{s}_A : A/G \rightarrow \mathcal{E}^{k,l}(M)/G$ extends to a continuous section $\bar{s}$ of $\bar{\pi}$ over $\mathcal{D}^k(M)/G$ which lifts to G -equivariant section of π over $\mathcal{D}^k(M)$. Moreover, if $\bar{s}_A$ takes values in the interiors of the fibers, one can choose s with the same property.*

Set $\mathcal{D}(M) = \bigcup_{k=0}^n \mathcal{D}^k(M)$. We are now ready to prove Theorem 1.1.

Theorem 4.13. *For any G -invariant closed subset Z of $\mathcal{X}(M)$ every G -equivariant center $\mathfrak{z} : Z \rightarrow M$ extends to a G -equivariant center $\mathfrak{c} : \mathcal{D}(M) \rightarrow M$.*

Proof. Given a nonnegative integer m let $\mathcal{S}^m = \bigcup_{i \leq m} \mathcal{D}^i$. Note that $\mathcal{S}^m$ is closed in $\mathcal{D}(M)$ for if $D_i \rightarrow D$, then G_D contains a conjugate of G_{D_i} by Lemma 2.3(vi) and, in particular, if each D_i lies in $\mathcal{S}^m$, then so does D .

We proceed by induction on the dimension of D^{G_D} . Let l be the smallest integer with $\mathcal{S}^l \neq \emptyset$. Use Lemma 4.12 with $A = Z \cap \mathcal{S}^l$ to define a G -equivariant center $\mathfrak{c} : \mathcal{S}^l \rightarrow M$ that extends $\mathfrak{z}$. Inductively, we suppose that a map $\mathfrak{c}$ with claimed properties is defined on $\mathcal{S}^k$, and hence on $\mathcal{Z}^k = \mathcal{S}^k \cup (Z \cap \mathcal{S}^{k+1})$, and then try to extend it to $\mathcal{S}^{k+1}$.

To this end fix a smooth G -equivariant embedding of $\tau : M \rightarrow \mathcal{H}$ where $\mathcal{H}$ is a Hilbert space equipped with the orthogonal G -action [Kankaanrinta 1994, Theorem 0.1]. The latter means that each element of g is continuous, linear, and preserves the inner product. Then the usual Riemannian open tubular neighborhood T_τ of τ is G -invariant and its projection $p_\tau : T_\tau \rightarrow M$ is G -equivariant [Kankaanrinta 2007, Theorem 5.1].

In [Antonyan 2005, Corollary 3.5 and Example 4.1] one finds a general extension result for G -maps from a Palais-proper space X with a G -invariant metric to any locally convex linear G -space, such as $\mathcal{H}$. It implies that for any closed G -invariant subset A of X every G -map $A \rightarrow \mathcal{H}$ can be extended to a G -map $X \rightarrow \mathcal{H}$. For this result the G -action on $\mathcal{H}$ must be linear, continuous, and every compact subgroup of G has to fix a point of $\mathcal{H}$. The latter is true for orthogonal actions of compact groups on $\mathcal{H}$, and in fact, any isometric group action with a bounded orbit on a CAT(0) space fixes the circumcenter of the orbit, and in a CAT(0) space, such as $\mathcal{H}$, every bounded set has a unique circumcenter [Bridson and Haefliger 1999, Proposition II.2.7].

Thus $\mathfrak{c} : \mathcal{Z}^k \rightarrow M \subset \mathcal{H}$ can be extended to a G -map $\tilde{\mathfrak{c}} : \mathcal{S}^{k+1} \rightarrow \mathcal{H}$. Set $O_\tau = \tilde{\mathfrak{c}}^{-1}(T_\tau)$ and $\bar{\mathfrak{c}} = p_\tau \circ \tilde{\mathfrak{c}}|_{O_\tau}$. Thus $\bar{\mathfrak{c}} : O_\tau \rightarrow M$ is a G -map that extends $\mathfrak{c}$.

Let us show that there is a closed neighborhood V of $\mathcal{Z}^k$ in O_τ on which $\bar{\mathfrak{c}}(D) \in \text{Int}(D)$. To this end fix a G -invariant metric d on $\mathcal{D}$ (using the metric of Remark 3.2 in each $\mathcal{D}^k(M)$ and setting the distance between points of different $\mathcal{D}^k(M)$ to be 1). Since $\bar{\mathfrak{c}}$ is continuous for any $D \in \mathcal{D}$ with $\bar{\mathfrak{c}}(D) \in \text{Int}(D)$ there is $\varepsilon_D > 0$ such that $\bar{\mathfrak{c}}(I) \in \text{Int}(I)$ for every I in the ε_D -ball centered at D . We can use the same ε_D for every D in the same G -orbit. The intersection with $\mathcal{S}^{k+1}$ of the union of all such balls is a G -invariant neighborhood U of $\mathcal{Z}^k$ in $\mathcal{S}^{k+1}$, and $\bar{\mathfrak{c}}|_U(D) \in \text{Int}(D)$. Finally, U contains a closed G -invariant neighborhood V of $\mathcal{Z}^k$, e.g., let V be the f -preimage of $[0, \frac{1}{2}]$ where

$$f(x) = \frac{d(x, \mathcal{Z}^k)}{d(x, \mathcal{Z}^k) + d(x, \mathcal{S}^{k+1} \setminus U)}.$$

It remains to enlarge V to $\mathcal{S}^{k+1}$. The restriction of $\bar{\mathfrak{c}}$ to $V_0 = V \cap \mathcal{D}^{k+1}$ is a G -equivariant section of the bundle $\mathcal{E}^{k+1} \rightarrow \mathcal{D}^{k+1}$ over V_0 . Applying Lemma 4.11 to the bundle whose fibers are $\text{Int}(D^{G_D})$, we extend $\bar{\mathfrak{c}}|_{V_0}$ to a G -equivariant section $\mathfrak{c}_0$

over $\mathcal{D}^{k+1}$. Since $\mathcal{D}^{k+1}$, $\text{Int}(V)$ are open in $\mathcal{S}^{k+1}$ and $c_0, \bar{c}|_{\text{Int}(V)}$ agree on $\mathcal{D}^{k+1} \cap \text{Int}(V)$, the maps define a G -map $\mathcal{S}^{k+1} \rightarrow M$ that takes each D to a point in $\text{Int}(D)$. This completes the induction step and proves the theorem. $\square$

Remark 4.14. The above proof reveals that Theorem 1.1 is a formal consequence of Lemma 4.7, and hence if the lemma is true for the C^0 topology on $\mathcal{X}(M)$, then the answer to Question 1.8 is affirmative.

Proof of Proposition 1.6. As discussed before the statement of the corollary, there exists an orthogonal A_5 -action on $\mathbb{R}^r$ that has no fixed points on an A_5 -invariant smoothly embedded 6-disk in $\mathbb{R}^r$. Denote the 6-disk by Δ , consider the product of Δ and the $(n-6)$ -disk with the trivial A_5 -action, and smooth corners. The result is a smoothly embedded A_5 -invariant n -disk in $\mathbb{R}^{n+r-6}$ on which A_5 acts without fixed points. Composing with the standard inclusion $\mathbb{R}^{n+r-6} \subset \mathbb{R}^m$ we can think of the n -disk as sitting in $\mathbb{R}^m$, where we let A_5 act trivially on the orthogonal complement of $\mathbb{R}^{n+r-6}$ in $\mathbb{R}^m$. This gives the desired claim when $m \geq n + r - 6 \geq r$.

For the other case consider the inclusion $\mathbb{R}^r \subset \mathbb{R}^n$, where A_5 acts trivially on the orthogonal complement of $\mathbb{R}^r$ in $\mathbb{R}^n$. Let Δ' be an A_5 -invariant smooth tubular neighborhood of Δ in $\mathbb{R}^n$ on which the A_5 -action is fixed point free. Note that Δ' is an embedded n -disk in $\mathbb{R}^n$, so composing with the inclusion $\mathbb{R}^n \subset \mathbb{R}^m$ gives the claim when $m \geq n \geq r$. $\square$

Appendix: Compact Lie group actions on low dimensional disks

The results of this appendix are surely known to experts but we could not find them in the literature.

It was shown by B. Kerékjártó that any compact topological group action on a 2-disk is equivalent to a linear action; see, e.g., [Kolev 2006]. More precisely, cone off the boundary of the G -action on D^2 , use [Kolev 2006] to conclude that the resulting action on S^2 is topologically equivalent to an action by a subgroup of $O(3)$, which is actually in $O(2)$ since it has a fixed point, and finally restrict the equivalence to D^2 .

For a smooth compact Lie group action, the uniformization gives an alternative route: Find a G -invariant Riemannian metric on the interior, and map it conformally to the standard hyperbolic disk so that the Lie group becomes a compact group of hyperbolic isometries, and hence after conjugation the standard $O(2)$. The same reasoning gives linearity of smooth actions on the closed interval, whose isometry group in any metric is $O(1)$.

It was shown in [Kwasik and Schultz 1992, Theorem B] that any smooth action of a compact Lie group on the 3-disk is smoothly equivalent to a linear action.

Lemma A.1. *If G is a nontrivial finite group that acts smoothly and effectively on the 4-disk D , then either D^G is a disk of dimension ≤ 2 , or G is an order*

two group generated by an orientation-reversing involution and D^G is a compact acyclic 3-manifold.

Proof. Any smooth finite group action on a 4-disk has a fixed point [Buchdahl et al. 1990, Theorem II.2]. Thus D^G is a compact smooth properly embedded submanifold of D . Let S be the double of D along ∂D equipped with the smooth G -action. Thus S is a smooth homotopy 4-sphere and S^G is nonempty.

The fixed point set of any smooth (or, more generally, locally linear) orientation-preserving action of a finite group on a homology 4-sphere is either empty or homeomorphic to a sphere [De Michelis 1989, Theorem 2.1]. Let $H \leq G$ be the subgroup of the orientation-preserving elements. The group G/H has order at most two, because H is the kernel of the G -action on $H_4(D, \partial D; \mathbb{Z}) \cong \mathbb{Z}$.

Thus S^H is a closed smooth submanifold of S that is homeomorphic to a sphere of dimension $l \in [0, 4]$. Since H preserves orientation, we have $l \neq 3$ else H would have to act nontrivially on a one-dimensional fiber of the normal bundle to D^G . If $l \leq 2$, then D^H is homeomorphic to the l -disk, the only manifold whose double is the l -sphere. Since D^G is the fixed point set of the G/H -action on D^H we conclude for $l \leq 2$ that G/H acts linearly on D^G , and hence D^G is a subdisk of D^H as claimed.

It remains to consider the case $l = 4$ where H is trivial and G is generated by an orientation-reversing involution of D . Then D^G is a homology disk [Bredon 1972, Theorem III.5.2] of dimension $l \in [0, 3]$. The G -action in a fiber of the normal bundle to D^G only fixes the origin and reverses the orientation, hence l is odd. If $l = 1$, then D^G is a one-dimensional disk, as claimed, and we are left with the exceptional case $l = 3$ where D^H is acyclic. $\square$

Remark A.2. Any compact acyclic 3-dimensional smooth submanifold of S^3 is homeomorphic to D^3 because its boundary is a homology 2-sphere, and hence is homeomorphic to S^2 , so that the Schoenflies theorem applies [Brown 1960]. In particular, if M and G are as in Corollary 1.5 and D is a C^1 embedded 4-disk in M , then D^{G_D} is homeomorphic to a disk.

Lemma A.3. *If G is an infinite compact Lie group that acts smoothly and effectively on the 4-disk D , then D^G is a disk of dimension ≤ 2 .*

Proof. If the claim is true for the connected component G_0 of G , then it is true for G because if G_0 fixes a disk F of dimension ≤ 2 , then G/G_0 acts on F . Since any compact Lie group action on a disk of dimension ≤ 2 is equivalent to a linear action, G/G_0 fixes a subdisk of F of dimension ≤ 2 , which is then fixed by G .

Thus we may assume that G is connected. By the structure theory of compact Lie groups there is a surjective homomorphism $T \times \bar{G} \rightarrow G$ with finite kernel where T is a torus and $\bar{G}$ is a semisimple connected compact Lie group. The resulting (possibly ineffective) T -action on D^4 has a fixed point set D^T which is an integral homology disk of dimension 0, 2, or 4; see [Bredon 1972, Theorem III.10.3], and in particular

it is nonempty. If T is nontrivial, then D^T is a nowhere dense [Bredon 1972, Theorem III.9.5] smooth submanifold of D , and so $\dim(D^T) < 4$. Thus D^T is diffeomorphic to a disk of dimension ≤ 2 . Since $\bar{G}$ normalizes T , it acts on D^T , and the argument of the previous paragraph applies to show that G fixes a disk of dimension ≤ 2 .

Thus we may assume that G is semisimple. The principal orbit G/H cannot be a point because then G fixes an open subset of D [Bredon 1972, Theorem IV.3.1], and cannot be a circle as, e.g., in the homotopy sequence of the bundle $G \rightarrow G/H$ the group $\pi_1(G/H)$ sits between finite groups $\pi_1(G)$, $\pi_0(H)$. Also G/H is a closed manifold embedded in D^4 , so its dimension is at most 3. Thus G/H has codimension 1 or 2. Then [Bredon 1972, Theorems IV.8.1 and IV.8.5] imply that the G -action on $\text{Int}(D)$ is equivalent to an orthogonal action $\mathbb{R}^4$, so its fixed point set is a linear subspace V , and in particular is nonempty. Since D^G is a smooth submanifold that is transverse to ∂D , we get $\text{Int}(D^G) = V$. Thus

$$0 \leq \dim(D^G) = \dim(V) \leq \dim(\text{Int}(D)/G) = 4 - \dim(G/H) \leq 2$$

and therefore D^G is a disk of dimension ≤ 2 . □

Lemma A.4. *There is a smooth involution on D^4 whose fixed point set is an acyclic nonsimply connected 3-manifold with boundary that is properly embedded in D^4 .*

Proof. In [Mazur 1961; Poenaru 1960] one finds a smooth compact contractible 4-manifold C such that ∂C is not simply connected while the double DC of C along ∂C is diffeomorphic to S^4 . Let ι be the doubling involution. Removing a small ι -invariant open ball B centered at ∂C yields a ι -invariant copy of D^4 that is the double of $C \setminus B$ along $\partial C \setminus B$. Here ι permutes two copies of $C \setminus B$ and fixes $\partial C \setminus B$ pointwise. Finally, since ∂C is a nonsimply connected homology sphere, $\partial C \setminus B$ is a compact acyclic nonsimply connected 3-manifold. □

Here is an example of a 5-disk D in $\mathbb{R}^5$ such that G_D is generated by a linear involution and D^{G_D} is not contractible.

Lemma A.5. *If r is the reflection in the equator of S^5 and $k \in \{4, 5\}$, then S^5 contains a smoothly embedded r -invariant copy of D^k which transversely intersects the equator in an acyclic nonsimply connected $(k-1)$ -manifold with boundary.*

Proof. In the notations of Lemma A.4 consider a proper embedding of C into D^5 (e.g., obtained by isotoping $C \subset S^4 = \partial D^5$ into the interior of D^5 and concatenating the result with the track of isotopy over ∂C). Removing from C a small r -invariant open 4-disk B centered at a point of $\partial C \subset \partial D^5$ and then doubling along ∂D^5 gives an r -invariant copy of $DC \setminus B$ of Lemma A.4 in S^5 that intersects the equator along $\partial C \setminus B$. This covers the case $k = 4$. For the case $k = 5$ set $I = [-1, 1]$ and note that $DC \times I$ embeds as an r -invariant tubular neighborhood of DC in S^5 , and after smoothing corners $(DC \setminus B) \times I$ is an r -invariant 5-disk in S^5 that intersects the equator along $(C \setminus B) \times I$. □

Acknowledgments

Ghomi first learned about the problem of finding a center for planar Jordan domains from Eugenio Calabi in 1995. Both authors thank John Etnyre, Ralph Howard, Dan Margalit and Krzysztof Pawalowski for helpful discussions.

References

- [Antonyan 2005] S. A. Antonyan, “Extending equivariant maps into spaces with convex structure”, *Topology Appl.* **153**:2-3 (2005), 261–275. MR Zbl
- [Bak and Morimoto 2005] A. Bak and M. Morimoto, “The dimension of spheres with smooth one fixed point actions”, *Forum Math.* **17**:2 (2005), 199–216. MR Zbl
- [Boileau et al. 2005] M. Boileau, B. Leeb, and J. Porti, “Geometrization of 3-dimensional orbifolds”, *Ann. of Math. (2)* **162**:1 (2005), 195–290. MR Zbl
- [Bredon 1972] G. E. Bredon, *Introduction to compact transformation groups*, Pure and Applied Mathematics **46**, Academic Press, New York, 1972. MR Zbl
- [Bridson and Haefliger 1999] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften **319**, Springer, 1999. MR Zbl
- [Brown 1960] M. Brown, “A proof of the generalized Schoenflies theorem”, *Bull. Amer. Math. Soc.* **66** (1960), 74–76. MR Zbl
- [Buchdahl et al. 1990] N. P. Buchdahl, S. Kwasik, and R. Schultz, “One fixed point actions on low-dimensional spheres”, *Invent. Math.* **102**:3 (1990), 633–662. MR Zbl
- [De Michelis 1989] S. De Michelis, “The fixed point set of a finite group action on a homology four sphere”, *Enseign. Math. (2)* **35**:1-2 (1989), 107–116. MR Zbl
- [Dinkelbach and Leeb 2009] J. Dinkelbach and B. Leeb, “Equivariant Ricci flow with surgery and applications to finite group actions on geometric 3-manifolds”, *Geom. Topol.* **13**:2 (2009), 1129–1173. MR Zbl
- [Hirsch 1976] M. W. Hirsch, *Differential topology*, Graduate Texts in Mathematics **33**, Springer, New York, 1976. MR Zbl
- [Kaiser et al. 1991] M. J. Kaiser, T. L. Morin, and T. B. Trafalis, “Centers and invariant points of convex bodies”, pp. 367–385 in *Applied geometry and discrete mathematics*, edited by P. Gritzmann and B. Sturmfels, DIMACS Ser. Discrete Math. Theoret. Comput. Sci. **4**, American Mathematical Society, Providence, RI, 1991. MR Zbl
- [Kankaanrinta 1994] M. Kankaanrinta, “On embeddings of proper smooth G -manifolds”, *Math. Scand.* **74**:2 (1994), 208–214. MR Zbl
- [Kankaanrinta 2007] M. Kankaanrinta, “Equivariant collaring, tubular neighbourhood and gluing theorems for proper Lie group actions”, *Algebr. Geom. Topol.* **7** (2007), 1–27. MR Zbl
- [Kobayashi and Nomizu 1963] S. Kobayashi and K. Nomizu, *Foundations of differential geometry*, vol. I, Interscience, New York, 1963. MR Zbl
- [Kolev 2006] B. Kolev, “Sous-groupes compacts d’homéomorphismes de la sphère”, *Enseign. Math. (2)* **52**:3-4 (2006), 193–214. MR Zbl
- [Kwasik and Schultz 1992] S. Kwasik and R. Schultz, “Icosahedral group actions on $\mathbb{R}^3$ ”, *Invent. Math.* **108**:2 (1992), 385–402. MR Zbl
- [Mazur 1961] B. Mazur, “A note on some contractible 4-manifolds”, *Ann. of Math. (2)* **73** (1961), 221–228. MR Zbl

- [Michael 1956a] E. Michael, “Continuous selections, I”, *Ann. of Math. (2)* **63** (1956), 361–382. MR Zbl
- [Michael 1956b] E. Michael, “Continuous selections, II”, *Ann. of Math. (2)* **64** (1956), 562–580. MR Zbl
- [Michael 1959] E. Michael, “Paracompact sets”, *Math. Scand.* **7** (1959), 372–376. MR Zbl
- [Moszyńska 2006] M. Moszyńska, *Selected topics in convex geometry*, Birkhäuser, Boston, 2006. MR Zbl
- [Oliver 1976] R. Oliver, “Smooth compact Lie group actions on disks”, *Math. Z.* **149**:1 (1976), 79–96. MR Zbl
- [Palais 1960] R. S. Palais, *The classification of G-spaces*, Mem. Amer. Math. Soc. **36**, American Mathematical Society, Providence, RI, 1960. MR Zbl
- [Palais 1961] R. S. Palais, “On the existence of slices for actions of non-compact Lie groups”, *Ann. of Math. (2)* **73** (1961), 295–323. MR Zbl
- [Palais 1966] R. S. Palais, “Homotopy theory of infinite dimensional manifolds”, *Topology* **5** (1966), 1–16. MR Zbl
- [Palais 1970] R. S. Palais, “ C^1 actions of compact Lie groups on compact manifolds are C^1 -equivalent to C^∞ actions”, *Amer. J. Math.* **92** (1970), 748–760. MR Zbl
- [Poenaru 1960] V. Poenaru, “Les decompositions de l’hypercube en produit topologique”, *Bull. Soc. Math. France* **88** (1960), 113–129. MR Zbl
- [Repovš and Semenov 1998] D. Repovš and P. V. Semenov, *Continuous selections of multivalued mappings*, Mathematics and its Applications **455**, Kluwer Academic, Dordrecht, 1998. MR Zbl
- [Repovš and Semenov 2002] D. Repovš and P. V. Semenov, “Continuous selections of multivalued mappings”, pp. 423–461 in *Recent progress in general topology, II*, edited by M. Hušek and J. van Mill, North-Holland, Amsterdam, 2002. MR Zbl
- [Repovš and Semenov 2014] D. Repovš and P. V. Semenov, “Continuous selections of multivalued mappings”, pp. 711–749 in *Recent progress in general topology*, vol. III, edited by K. P. Hart et al., Atlantis, Paris, 2014. MR Zbl
- [Schneider 2014] R. Schneider, *Convex bodies: the Brunn–Minkowski theory*, expanded ed., Encyclopedia of Mathematics and its Applications **151**, Cambridge University Press, 2014. MR Zbl

Received May 21, 2018. Revised July 1, 2019.

IGOR BELEGRADEK
SCHOOL OF MATHEMATICS
GEORGIA INSTITUTE OF TECHNOLOGY
ATLANTA, GA
UNITED STATES
ib@math.gatech.edu

MOHAMMAD GHOMI
SCHOOL OF MATHEMATICS
GEORGIA INSTITUTE OF TECHNOLOGY
ATLANTA, GA
UNITED STATES
ghomi@math.gatech.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing


<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 304 No. 2 February 2020

Graphs admitting only constant splines	385
KATIE ANDERS, ALISSA S. CRANS, BRIANA FOSTER-GREENWOOD, BLAKE MELLOR and JULIANNA TYMOCZKO	
Centers of disks in Riemannian manifolds	401
IGOR BELEGRADEK and MOHAMMAD GHOMI	
The geometry of the flex locus of a hypersurface	419
LAURENT BUSÉ, CARLOS D'ANDREA, MARTÍN SOMBRA and MARTIN WEIMANN	
Morse inequalities for Fourier components of Kohn–Rossi cohomology of CR covering manifolds with S^1 -action	439
RUNG-TZUNG HUANG and GUOKUAN SHAO	
On Seifert fibered spaces bounding definite manifolds	463
AHMAD ISSA and DUNCAN MCCOY	
Regularity of quotients of Drinfeld modular schemes	481
SATOSHI KONDO and SEIDAI YASUDA	
Sums of algebraic trace functions twisted by arithmetic functions	505
MAXIM KOROLEV and IGOR SHPARLINSKI	
Twisted calculus on affinoid algebras	523
BERNARD LE STUM and ADOLFO QUIRÓS	
Symplectic (-2) -spheres and the symplectomorphism group of small rational 4-manifolds	561
JUN LI and TIAN-JUN LI	
Addendum to the article Contact stationary Legendrian surfaces in $\mathbb{S}^5$	607
YONG LUO	
The Hamiltonian dynamics of magnetic confinement in toroidal domains	613
GABRIEL MARTINS	
Gluing Bartnik extensions, continuity of the Bartnik mass, and the equivalence of definitions	629
STEPHEN MCCORMICK	
Decomposable Specht modules indexed by bihooks	655
LIRON SPEYER and LOUISE SUTTON	
On the global well-posedness of one-dimensional fluid models with nonlocal velocity	713
ZHUAN YE	
Hopf cyclic cohomology for noncompact G -manifolds with boundary	753
XIN ZHANG	



0030-8730(2020)304:2;1-R