

Pacific Journal of Mathematics

**ON SEIFERT FIBERED SPACES
BOUNDING DEFINITE MANIFOLDS**

AHMAD ISSA AND DUNCAN MCCOY

Volume 304 No. 2

February 2020

ON SEIFERT FIBERED SPACES BOUNDING DEFINITE MANIFOLDS

AHMAD ISSA AND DUNCAN MCCOY

We establish an inequality which gives strong restrictions on when the standard definite plumbing intersection lattice of a Seifert fibered space over S^2 can embed into a standard diagonal lattice, and give two applications. First, we answer a question of Neumann and Zagier on the relationship between Donaldson's theorem and Fintushel–Stern's R -invariant. We also give a short proof of the characterization of Seifert fibered spaces which smoothly bound rational homology $S^1 \times D^3$'s.

1. Introduction

Donaldson's diagonalization theorem [1987] has led to many great successes in understanding several important questions in low dimensional topology, and in knot theory in particular. For example, Donaldson's theorem can often be used to answer questions concerning sliceness, unknotting number, 3-manifolds bounding rational homology balls, and surgery questions. In these cases, one typically uses Donaldson's theorem to obstruct a certain 3-manifold from bounding a certain type of smooth negative definite 4-manifold, with the obstruction taking the form of the existence of a certain map of intersection lattices. However, understanding this obstruction for large families of examples is often highly nontrivial, and can require combinatorial ingenuity.

One appealing application of Donaldson's theorem is to prove the well-known fact that the Poincaré homology sphere $P = S^2(2; 2, \frac{3}{2}, \frac{5}{4})$ does not bound a smooth integral homology 4-ball. This fact can, of course, be proved in many other ways, for example by using Rokhlin's theorem, Fintushel–Stern's R -invariant, or by using the d -invariant coming from Heegaard Floer homology. Assuming that P is oriented to bound the positive E_8 plumbing, the proof by Donaldson's theorem is as follows. If P were the boundary of a smooth integral homology 4-ball W , then we could form a closed positive definite manifold by gluing $-W$ to the positive E_8 plumbing. Donaldson's theorem would then imply that the E_8 intersection form is diagonalizable, which is, of course, untrue. In fact, as the E_8 intersection form

MSC2010: 57M27.

Keywords: Seifert fibered spaces, Donaldson's theorem, lattices, definite 4-manifolds.

does not embed into any positive definite diagonal lattice, this argument shows that P does not bound any smooth negative definite 4-manifold. The purpose of this paper is to generalize this argument to other Seifert fibered spaces. We prove the following theorem.

Theorem 1. *Let $Y = S^2(e; \frac{p_1}{q_1}, \dots, \frac{p_k}{q_k})$ be a Seifert fibered space over S^2 in standard form, that is, with $e \geq 0$, $\frac{p_i}{q_i} > 1$ for all $i \in \{1, 2, \dots, k\}$ and $\varepsilon(Y) \geq 0$. Suppose that Y bounds a smooth 4-manifold W such that $\sigma(W) = -b_2(W)$ and the inclusion induced map $H_1(Y; \mathbb{Q}) \rightarrow H_1(W; \mathbb{Q})$ is injective. Then there is a partition of $\{1, 2, \dots, k\}$ into at most e classes such that for each class C ,*

$$\sum_{i \in C} \frac{q_i}{p_i} \leq 1.$$

We note that the condition that $\varepsilon(Y) := e - \sum_{i=1}^k \frac{q_i}{p_i} \geq 0$ in Theorem 1 guarantees that Y is oriented to bound a positive (semi)definite plumbing 4-manifold. When Y is a rational homology sphere the map $H_1(Y; \mathbb{Q}) \rightarrow H_1(W; \mathbb{Q})$ is automatically injective so in this case we are simply obstructing the existence of a negative definite manifold bounding Y . Although we do not discuss the details in this paper, one can easily obtain analogous results for Seifert fibered spaces over any orientable base surface. In our notation, the Poincaré homology sphere oriented to bound the positive E_8 plumbing is $P = S^2(2; 2, \frac{3}{2}, \frac{5}{4})$, see Figure 2. The reader can easily verify that Theorem 1 obstructs P from bounding a negative definite manifold. When $k < 3$, the Seifert fibered spaces are lens spaces which are well known to bound both positive and negative definite smooth 4-manifolds. Finally, we note that the converse to Theorem 1 is not true. The integer homology sphere $S^2(1; 3, 5, \frac{13}{6})$ passes the obstruction, but does not bound a negative definite manifold as it bounds a positive definite plumbing whose intersection form does not embed in a diagonal lattice.

We give two applications of Theorem 1. First, we prove the following theorem.

Theorem 2. *Let $Y = S^2(e; \frac{p_1}{q_1}, \dots, \frac{p_k}{q_k})$ be a Seifert fibered integral homology sphere in standard form, that is, with $\frac{p_i}{q_i} > 1$ for all $i \in \{1, 2, \dots, k\}$, $e > 0$ and with Y oriented to bound a smooth positive definite plumbing 4-manifold. If Y bounds a smooth negative definite 4-manifold, then $e = 1$.*

In the course of proving Theorem 2, we obtain a positive answer to the following question asked by Neumann and Zagier [1985].

Question. *Let Y be as in Theorem 2. If the intersection form of the plumbing of Y is diagonalizable over \mathbb{Z} , must e be equal to 1?*

The motivation for this question comes from the R -invariant. Fintushel and Stern [1985] used gauge theory to define an invariant $R(Y)$ of Seifert fibered integral

homology spheres with the property that if $R(Y) > 0$ then Y does not bound a smooth negative definite 4-manifold W with $H_1(W)$ having no 2-torsion. Fintushel and Stern originally gave an expression for $R(Y)$ as a trigonometric sum involving the Seifert invariants of Y . Neumann and Zagier [1985] proved that these sums could be simply evaluated in terms of the central weight e of the standard positive definite plumbing bounding Y , showing that $R(Y) = 2e - 3$. Thus, if $e > 1$ then the R -invariant shows that Y does not bound a smooth negative definite 4-manifold W with $H_1(W)$ having no 2-torsion. In this light, the positive answer to the Neumann–Zagier question implies that this result obtained from the R -invariant is also a consequence of Donaldson’s theorem.

We are in fact able to prove a more general version of Theorem 2 which holds for all $|H_1(Y)| \in \{1, 2, 3, 5, 6, 7\}$; see Theorem 8 of Section 5. Some particular cases of Theorem 2 are known. In their original paper, Neumann and Zagier [1985] claimed to have proved the cases when $k = 3$, and when $k = 4$ and $e \neq 3$, but do not provide a proof, remarking that their proof was “clearly not the right proof.” The special case when $e = k - 1$ follows from [Lecuona and Lisca 2011, Lemma 3.3].

Finally, we note that a positive answer to the Neumann–Zagier question is a special case of a more general conjecture made by Neumann [1989], stating that if an integral homology sphere Y is given as the boundary of a positive definite plumbing tree Γ and the intersection lattice of Γ is isomorphic to a diagonal lattice, then some vertex of Γ has weight 1. This general form of Neumann’s conjecture for graph manifolds remains open.

Lidman and Tweedy [2018, Remark 4.3] asked whether a Seifert fibered integral homology sphere with central weight different from 1 must have nonvanishing Heegaard–Floer d -invariant. As a corollary of Theorem 2, we answer their question positively.

Corollary 3. *Let Y be a Seifert fibered integral homology sphere, and let $e \in \mathbb{Z}$ be the central weight in the standard definite plumbing graph for Y . If $|e| \neq 1$, then $d(Y) \neq 0$.*

As a second application, we give a short proof of the following theorem which, in particular, gives a classification of the Seifert fibered spaces bounding rational homology $S^1 \times D^3$ ’s.

Theorem 4. *Let Y be a Seifert fibered space over S^2 with*

$$H_*(Y; \mathbb{Q}) \cong H_*(S^1 \times S^2; \mathbb{Q}).$$

The following are equivalent:

- (1) *Y is of the form $S^2(k; \frac{p_1}{q_1}, \frac{p_1}{p_1 - q_1}, \dots, \frac{p_k}{q_k}, \frac{p_k}{p_k - q_k})$, where $k \geq 0$ and $\frac{p_i}{q_i} > 1$ for all $i \in \{1, \dots, k\}$.*

- (2) $Y = \partial W$, where W is a smooth 4-manifold with $H_*(W; \mathbb{Q}) \cong H_*(S^1 \times D^3; \mathbb{Q})$.
- (3) Y is the boundary of smooth 4-manifolds W_+ and W_- such that $\sigma(W_{\pm}) = \pm b_2(W_{\pm})$ and each of the inclusion-induced maps $H_1(Y; \mathbb{Q}) \rightarrow H_1(W_{\pm}; \mathbb{Q})$ is injective.

Seifert fibered spaces bounding rational homology $S^1 \times D^3$'s naturally arise in two contexts. First, a Seifert fibered space rational homology $S^1 \times S^2$ which embeds in S^4 necessarily bounds a rational homology $S^1 \times D^3$. Indeed, in this context Donald [2015, Proof of Theorem 1.3] proved the implication (2) implies (1) of Theorem 4. Second, a smoothly slice 2-component Montesinos link has double branched cover a Seifert fibered space over S^2 bounding a rational homology $S^1 \times D^3$. Motivated by trying to determine the slice 2-component Montesinos links, Aceto [2015, Theorem 1.2] also classified Seifert fibered spaces bounding rational homology $S^1 \times D^3$'s.

Much like the proofs by Donald and Aceto, our proof also proceeds by means of Donaldson's theorem. However, their proofs rely on the work of Lisca [2007] which gives a detailed analysis on sums of linear lattices embedding in a full-rank lattice. We give a short proof of Theorem 4 circumventing the reliance on Lisca's work. We obtain the additional equivalent condition (3) in Theorem 4, since our method does not require the lattice embeddings to have full-rank.

Finally, we note that Theorem 1 also plays a key role in the paper [Issa and McCoy 2018], where we analyze which Seifert fibered spaces smoothly embed in S^4 , and in particular, completely determine the Seifert fibered spaces

$$Y = S^2(e; r_1, \dots, r_k) \quad \text{with } r_i \in \mathbb{Q}_{>1} \text{ for all } i, \quad \varepsilon(Y) > 0 \quad \text{and} \quad e > k/2$$

which smoothly embed in S^4 .

In Section 2, we recall some standard facts and establish notation and conventions. In Section 3, we prove the key technical theorem used to prove Theorem 1. In Section 4, we analyze when gluing compact 4-manifolds with boundary results in a definite 4-manifold and give a proof of Theorem 1. In Section 5, we prove Theorem 8 answering the Neumann–Zagier question, as well as prove Corollary 3. Finally, in Section 6 we prove Theorem 4 determining the Seifert fibered spaces which bound rational homology $S^1 \times D^3$'s.

2. Preliminaries

In this section we briefly recall some standard facts about Seifert fibered spaces and intersection lattices, and establish notation and conventions. See [Neumann and Raymond 1978] for a more in-depth treatment on Seifert fibered spaces and plumbings.




Figure 1. Weighted linear chain representing $r = [a_1, \dots, a_h]^-$.




Figure 2. Surgery presentation for the Seifert fibered space $S^2(e; \frac{p_1}{q_1}, \dots, \frac{p_k}{q_k})$.

Given $r \in \mathbb{Q}_{>1}$, there is a unique (negative) continued fraction expansion

$$r = [a_1, \dots, a_h]^- := a_1 - \frac{1}{a_2 - \frac{1}{\ddots a_{h-1} - \frac{1}{a_h}}},$$

where $h \geq 1$ and $a_i \geq 2$ are integers for all $i \in \{1, \dots, h\}$. We associate to r the weighted linear graph (or linear chain) given in Figure 1. We call the vertex with weight labeled by a_i the i -th vertex of the linear chain associated to r , so that the vertex labeled with weight a_1 is the first, or starting vertex of the linear chain.

We denote by $Y = S^2(e; \frac{p_1}{q_1}, \dots, \frac{p_k}{q_k})$ the Seifert fibered space over S^2 given in Figure 2, where $e \in \mathbb{Z}$, and $\frac{p_i}{q_i} \in \mathbb{Q}$ is nonzero for all $i \in \{1, \dots, k\}$. The generalized Euler invariant of Y is given by $\varepsilon(Y) = e - \sum_{i=1}^k \frac{q_i}{p_i}$. Every Seifert fibered space Y is (possibly orientation reversing) homeomorphic to one in standard form, i.e., such that $\varepsilon(Y) \geq 0$ and $\frac{p_i}{q_i} > 1$ for all $i \in \{1, \dots, k\}$. We henceforth assume that Y is in standard form. If $\varepsilon(Y) \neq 0$ then Y is a rational homology sphere with $|H_1(Y)| = |p_1 \cdots p_k \varepsilon(Y)|$, and if $\varepsilon(Y) = 0$ then Y is a rational homology $S^1 \times S^2$.

For each $i \in \{1, \dots, k\}$, we have the unique continued fraction expansion $\frac{p_i}{q_i} = [a_1^i, \dots, a_{h_i}^i]^-$ where $h_i \geq 1$ and $a_j^i \geq 2$ are integers for all $j \in \{1, \dots, h_i\}$. We associate to $Y = S^2(e; \frac{p_1}{q_1}, \dots, \frac{p_k}{q_k})$ the weighted star-shaped graph in Figure 3. The i -th leg of the star-shaped graph is the weighted linear subgraph for p_i/q_i generated by the vertices labeled with weights $a_1^i, \dots, a_{h_i}^i$. The degree k vertex labeled with weight e is called the central vertex.

Let Γ be either the weighted star-shaped graph for Y , or a disjoint union of weighted linear graphs. There is an oriented smooth 4-manifold X_Γ given by plumbing D^2 -bundles over S^2 according to the weighted graph Γ . We denote by




Figure 3. The weighted star-shaped plumbing graph Γ .

$|\Gamma|$ the number of vertices in Γ . Let $N = |\Gamma|$ and denote the vertices of Γ by v_1, v_2, \dots, v_N . The zero-sections of the D^2 -bundles over S^2 corresponding to each of v_1, \dots, v_N in the plumbing together form a natural spherical basis for $H_2(X_\Gamma)$. With respect to this basis, which we call the vertex basis, the intersection form of X_Γ is given by the weighted adjacency matrix Q_Γ with entries Q_{ij} , $1 \leq i, j \leq N$ given by

$$Q_{ij} = \begin{cases} w(v_i) & \text{if } i = j, \\ -1 & \text{if } v_i \text{ and } v_j \text{ are connected by an edge,} \\ 0 & \text{otherwise,} \end{cases}$$

where $w(v_i)$ is the weight of vertex v_i . Denoting by Q_{X_Γ} the intersection form of X_Γ , we call $(H_2(X_\Gamma), Q_{X_\Gamma}) \cong (\mathbb{Z}^N, Q_\Gamma)$ the intersection lattice of X_Γ (or of Γ). We denote the intersection pairing of two elements $x, y \in \mathbb{Z}^N$ by $x \cdot y = x^T Q_\Gamma y$. Now assume that Γ is the star-shaped plumbing for Y . If $\varepsilon(Y) > 0$ then X_Γ is a positive definite 4-manifold and Γ is the standard positive definite plumbing graph for Y . If $\varepsilon(Y) = 0$, then X_Γ is a positive semidefinite manifold.

Let $\iota : (\mathbb{Z}^N, Q_\Gamma) \rightarrow (\mathbb{Z}^m, \text{Id})$, $m > 0$, be a map of lattices, i.e., a \mathbb{Z} -linear map preserving pairings, where $(\mathbb{Z}^m, \text{Id})$ is the standard positive diagonal lattice. We call ι a lattice embedding if it is injective. We adopt the following standard abuse of notation. First, for each $i \in \{1, \dots, N\}$, we identify the vertex v_i with the corresponding i -th basis element of (\mathbb{Z}^N, Q_Γ) . Moreover, we shall identify an element $v \in (\mathbb{Z}^N, Q_\Gamma)$ with its image $\iota(v) \in (\mathbb{Z}^m, \text{Id})$.

3. The embedding inequality

In this section, we prove Theorem 6 below, which is the key technical result of this paper. In particular, it will be used in the next section to prove Theorem 1. We begin with some continued fraction identities which we will need.

Lemma 5. Let $\{a_i\}_{i \geq 1}$ be a sequence of integers with $a_i \geq 2$ for all i , and let $p_k/q_k = [a_1, \dots, a_k]^-$ for all $k \geq 1$. Then we have the following identities:

- (a) $q_n p_{n-1} - p_n q_{n-1} = 1$ for all $n \geq 2$.
- (b) $[a_1, \dots, a_n, x]^- = \frac{x p_n - p_{n-1}}{x q_n - q_{n-1}}$, for all $n \geq 2$ and $x \in \mathbb{R}$ such that both sides are well defined.
- (c) $p_n = \det \begin{pmatrix} a_1 & -1 & 0 & 0 \\ -1 & a_2 & -1 & 0 \\ 0 & -1 & \ddots & -1 \\ 0 & 0 & -1 & a_n \end{pmatrix}$ and $q_n = \det \begin{pmatrix} a_2 & -1 & 0 & 0 \\ -1 & a_3 & -1 & 0 \\ 0 & -1 & \ddots & -1 \\ 0 & 0 & -1 & a_n \end{pmatrix}$ for all $n \geq 2$.

Proof. For $a \in \mathbb{R}$, let M_a denote the matrix $M_a = \begin{pmatrix} a & -1 \\ 1 & 0 \end{pmatrix}$. If $q/r = [a_2, \dots, a_n]^-$, then $\frac{p_n}{q_n} = a_1 - \frac{r}{q}$. In particular, we have

$$\begin{pmatrix} p_n \\ q_n \end{pmatrix} = M_{a_1} \begin{pmatrix} q \\ r \end{pmatrix}.$$

Thus, one can inductively show that

$$(1) \quad \begin{pmatrix} p_n \\ q_n \end{pmatrix} = M_{a_1} \cdots M_{a_n} \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

and furthermore that

$$(2) \quad \begin{pmatrix} p_n & -p_{n-1} \\ q_n & -q_{n-1} \end{pmatrix} = M_{a_1} \cdots M_{a_n}.$$

Identity (a) follows by taking determinants of (2) and observing that $\det M_a = 1$ for any a . Identity (b) follows from combining (1) and (2) to get

$$M_{a_1} \cdots M_{a_n} M_x \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} p_n & -p_{n-1} \\ q_n & -q_{n-1} \end{pmatrix} \begin{pmatrix} x \\ 1 \end{pmatrix} = \begin{pmatrix} x p_n - p_{n-1} \\ x q_n - q_{n-1} \end{pmatrix}.$$

The identities in (c) can easily be proven by induction using the observation that

$$\det \begin{pmatrix} a_1 & -1 & 0 \\ -1 & \ddots & -1 \\ 0 & -1 & a_n \end{pmatrix} = a_1 \det \begin{pmatrix} a_2 & -1 & 0 \\ -1 & \ddots & -1 \\ 0 & -1 & a_n \end{pmatrix} - \det \begin{pmatrix} a_3 & -1 & 0 \\ -1 & \ddots & -1 \\ 0 & -1 & a_n \end{pmatrix}. \quad \square$$

The following theorem is the key technical result of this paper.

Theorem 6. Let $\iota : (\mathbb{Z}^{|\Gamma|}, Q_\Gamma) \rightarrow (\mathbb{Z}^m, \text{Id})$ be a map of lattices, where $m > 0$ and Γ is a disjoint union of weighted linear chains representing fractions $\frac{p_1}{q_1}, \dots, \frac{p_k}{q_k} \in \mathbb{Q}_{>1}$. Suppose that there is a unit vector $w \in (\mathbb{Z}^m, \text{Id})$ which pairs nontrivially with (the

image of) the starting vertex of each linear chain. Then

$$\sum_{i=1}^k \frac{q_i}{p_i} \leq 1.$$

Moreover, if we have equality then w has pairing ± 1 with the starting vertex of each linear chain.

Proof. Let $\{e_1, \dots, e_m\}$ denote the orthonormal basis of coordinates vectors of $(\mathbb{Z}^m, \text{Id})$. Since the unit vectors in $(\mathbb{Z}^m, \text{Id})$ are precisely those vectors of the form $\pm e_i$ where $i \in \{1, \dots, m\}$, by a change of basis if necessary, we may assume that $w = e_1 \in (\mathbb{Z}^m, \text{Id})$. Write $\iota: (\mathbb{Z}^{|\Gamma|}, Q_\Gamma) \rightarrow (\mathbb{Z}^m, \text{Id})$ as an integer matrix with respect to the vertex basis of $(\mathbb{Z}^{|\Gamma|}, Q_\Gamma)$, and let M be the transpose of this matrix. Since ι preserves intersection pairings we have, $u^T Q_\Gamma v = \iota(u)^T \iota(v) = (M^T u)^T (M^T v) = u^T M M^T v$ for all $u, v \in (\mathbb{Z}^{|\Gamma|}, Q_\Gamma)$. Thus,

$$M M^T = Q_\Gamma = \begin{pmatrix} A_1 & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & A_k \end{pmatrix},$$

where for each $i \in \{1, \dots, k\}$, A_i on the diagonal represents a block matrix of the form

$$A_i = \begin{pmatrix} a_1^i & -1 & 0 & 0 \\ -1 & a_2^i & -1 & 0 \\ 0 & -1 & \ddots & -1 \\ 0 & 0 & -1 & a_{h_i}^i \end{pmatrix},$$

where $[a_1^i, \dots, a_{h_i}^i]^-$ is the continued fraction expansion for p_i/q_i . If a matrix A can be written as a product $M' M'^T$, then¹

$$(3) \quad \det A \geq 0.$$

We will prove the theorem by applying (3) to a matrix of the form $A = M' M'^T$, where M' is a suitable modification of M .

We may write M in the form

$$M = \begin{pmatrix} M_1 \\ \vdots \\ M_k \end{pmatrix},$$

where for all $i \in \{1, \dots, k\}$, M_i is a matrix such that $M_i M_i^T = A_i$. By the assumption that e_1 pairs nontrivially with each of the starting vertices of the linear chains, we may assume that each matrix M_i is nonzero in its top left entry. For each $i \in \{1, \dots, k\}$,

¹Let $v \neq 0$ be an eigenvector of A with eigenvalue λ . We have $v^T A v = \lambda \|v\|^2 = \|M'^T v\|^2 \geq 0$, thus $\lambda \geq 0$. Since $\det A$ is a product of eigenvalues, this implies that $\det A \geq 0$ as required.

let $p_i/q_i = [a_1^i, \dots, a_{h_i}^i]^-$ be the standard continued fraction expansion and choose M'_i to be the submatrix of M_i obtained by taking the first l_i rows, where l_i is chosen so that the first column w_i of M'_i takes one of the two forms as follows:

(Form 1) If $a_1^i = 2$ and the first column of M_i is of the form $\pm (1 \ -1 \ \dots)^T$ then we may take $w_i = \pm (1 \ -1 \ 0 \ \dots \ 0 \ v)^T$, where $v = 0$ only if $M'_i = M_i$.

(Form 2) Otherwise, we may take w_i to be of the form $w_i = (u \ 0 \ \dots \ 0 \ v)^T$, where $v = 0$ only if $M'_i = M_i$.

Let M' be the matrix

$$M' = \begin{pmatrix} 1 & 0 & \dots & 0 \\ & M'_1 & & \\ & \vdots & & \\ & M'_k & & \end{pmatrix}.$$

Then the product $A = M' M'^T$ takes the form of the block matrix

$$M' M'^T = \begin{pmatrix} 1 & w_1^T & \dots & w_k^T \\ w_1 & A'_1 & & 0 \\ \vdots & & \ddots & \\ w_k & 0 & & A'_k \end{pmatrix}.$$

Claim. We can write $\det A$ in the form

$$\det A = (P_1 \dots P_k) \left(1 - \sum_{i=1}^k \frac{Q_i}{P_i} \right),$$

where $P_i = \det A'_i > 0$ and

$$Q_i = -\det \begin{pmatrix} 0 & w_i^T \\ w_i & A'_i \end{pmatrix}$$

is a quantity depending only on A'_i and w_i .

Proof. Since Γ represents a positive definite lattice, $P_i > 0$ for all i . By the multilinearity of the determinant in both rows and columns we have

$$\det \begin{pmatrix} 1 & w_1^T & \dots & w_k^T \\ w_1 & A'_1 & & 0 \\ \vdots & & \ddots & \\ w_k & 0 & & A'_k \end{pmatrix} = \det \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & A'_1 & & 0 \\ \vdots & & \ddots & \\ 0 & 0 & & A'_k \end{pmatrix} + \sum_{1 \leq i, j \leq k} \det B_{ij},$$

where B_{ij} is the matrix

$$B_{ij} = \begin{pmatrix} 0 & \cdots & w_j^T & \cdots & 0 \\ \vdots & A'_1 & & & \\ w_i & & \ddots & & 0 \\ \vdots & & 0 & \ddots & \\ 0 & & & & A'_k \end{pmatrix}.$$

Since A'_i has full rank, w_i can be expressed as a rational linear combination of the columns of A'_i , and hence $\det B_{ij} = 0$ for all $i \neq j$. For $i \in \{1, \dots, k\}$, by row and column operations, we can put B_{ii} into the form of a diagonal block matrix with diagonal blocks $\begin{pmatrix} 0 & w_i^T \\ w_i & A'_i \end{pmatrix}$, $A'_1, \dots, A'_{i-1}, A'_{i+1}, \dots, A'_k$ without changing the determinant. Hence, $\det B_{ii}$ is the product of the determinants of these blocks, that is, $\det B_{ii} = -(P_1 \cdots P_k) \frac{Q_i}{P_i}$. \square

Since $P_i > 0$ for all i , the previous claim combined with $\det A \geq 0$ (see (3)) shows that

$$(4) \quad \sum_{i=1}^k \frac{Q_i}{P_i} \leq 1.$$

So to prove the inequality in the theorem it suffices to show that $Q_i/P_i \geq q_i/p_i$ for each $i \in \{1, \dots, k\}$. To do this it suffices to consider some fixed $i \in \{1, \dots, k\}$. For convenience, let $P/Q = P_i/Q_i$ and $p/q = p_i/q_i = [a_1, a_2, \dots, a_h]^-$, where $a_j \geq 2$ for all $j \in \{1, \dots, h\}$, and let $l = l_i$ be the number of rows of A'_i .

Consider the following identity obtained by adding the second row to the first row, and the second column to the first column:

$$(5) \quad \det \begin{pmatrix} 0 & -1 & 1 & \cdots & v \\ -1 & 2 & -1 & & 0 \\ 1 & -1 & a_2 & -1 & \\ \vdots & & -1 & \ddots & -1 \\ v & 0 & & -1 & a_l \end{pmatrix} = \det \begin{pmatrix} 0 & 1 & 0 & \cdots & v \\ 1 & 2 & -1 & & 0 \\ 0 & -1 & a_2 & -1 & \\ \vdots & & -1 & \ddots & -1 \\ v & 0 & & -1 & a_l \end{pmatrix}.$$

Recall that w_i takes one of two possible forms. By applying the above identity if w_i takes the form (Form 1)orm 1, we see that regardless of the form that w_i takes, Q is equal to the determinant of a matrix of the following form

$$(6) \quad Q = -\det \begin{pmatrix} 0 & +u & +0 & \cdots & +v \\ u & +a_1 & -1 & & +0 \\ 0 & -1 & +a_2 & -1 & \\ \vdots & & -1 & \ddots & -1 \\ v & +0 & & -1 & +a_l \end{pmatrix},$$

where if $(u, v) = (\pm 1, \mp 1)$ then either $l > 2$ or $a_1 > 2$. If w_i takes the form (Form 1)orm 1, we define $u \in \{\pm 1\}$ via Equation (6) by applying the identity in (5).

For $j \in \{1, \dots, h\}$, let r_j/s_j denote the continued fraction $[a_1, \dots, a_j]^-$. Note that $P = r_l$.

Claim. $Q = u^2 s_l + 2uv + v^2 r_{l-1}$

Proof. Applying cofactor expansion along the first column and first row in (6) gives

$$Q = u^2 C_1 + (-1)^{l+1} uv C_2 + (-1)^{l+1} uv C_3 + v^2 C_4,$$

where

$$C_1 = \det \begin{pmatrix} a_2 & -1 & \cdots & 0 \\ -1 & a_3 & & \\ \vdots & -1 & \ddots & -1 \\ 0 & & & a_l \end{pmatrix}, \quad C_2 = \det \begin{pmatrix} -1 & a_2 & -1 & \cdots & 0 \\ 0 & -1 & a_3 & & \\ \vdots & & -1 & \ddots & -1 \\ & & & -1 & a_{l-1} \\ 0 & & & & -1 \end{pmatrix},$$

$$C_3 = \det \begin{pmatrix} -1 & 0 & & \cdots & 0 \\ a_2 & -1 & & & \\ -1 & a_3 & -1 & & \\ & -1 & \ddots & -1 & \\ 0 & & & a_{l-1} & -1 \end{pmatrix}, \quad C_4 = \det \begin{pmatrix} a_1 & -1 & \cdots & 0 \\ -1 & a_2 & -1 & \\ \vdots & -1 & \ddots & -1 \\ 0 & & -1 & a_{l-1} \end{pmatrix}.$$

Using the continued fraction identities in Lemma 5, we see that $C_1 = s_l$ and $C_4 = r_{l-1}$. Finally, notice that C_2 (resp. C_3) is the determinant of an upper (resp. lower) triangular matrix with $l-1$ diagonal entries all of which are -1 , hence $C_2 = C_3 = (-1)^{l-1}$. \square

Claim. We have $Q/P \geq q/p$ with equality only if $u = \pm 1$ and $v = 0$.

Proof. Recall that if $v = 0$ then $r_l/s_l = p/q$. Thus if $v = 0$, $Q/P = u^2 q/p$. Since $u \neq 0$, we clearly have $Q/P \geq q/p$ with equality only if $u^2 = 1$, as required.

From now on assume that $v \neq 0$. Notice that if $l = 1$, then we necessarily have $v = 0$. Thus we can further assume that $l \geq 2$. This implies that $s_l \geq 2$ and $r_{l-1} \geq 2$. By completing the square, it follows from the preceding claim that Q can be written in the form

$$(7) \quad Q = (s_l - 1)u^2 + (u + v)^2 + (r_{l-1} - 1)v^2.$$

Since u and v are nonzero integers, it follows from (7) that

$$Q \geq s_l + r_{l-1} - 2.$$

If $r_{l-1} = 2$, then this inequality can be strengthened. It follows from the uniqueness of the negative continued fraction expansions that $r_{l-1} = 2$ only if $l = 2$ and $a_1 = 2$.

By the condition stated immediately after (6) this implies that we do not have $u = -v \in \{\pm 1\}$. Thus we have that either $|u| \geq 2$, $|v| \geq 2$ or $u + v \neq 0$. In any of these eventualities it follows from (7) that

$$Q \geq s_l + r_{l-1} - 1.$$

By combining these two inequalities, we see that there is $\varepsilon \in \{1, 2\}$ such that

$$(8) \quad Q \geq s_l + r_{l-1} - \varepsilon \quad \text{and} \quad r_{l-1} - \varepsilon \geq 1.$$

If $l = h$, or equivalently, $p/q = r_l/s_l$, then $P = p$ and (8) implies that $Q > q = s_l$. It follows that $Q/P > q/p$ in this case. For the rest of the proof, we assume that $p/q \neq r_l/s_l$ and, in particular, that $p/q = [a_1, \dots, a_l, x]^-$, where $x = [a_{l+1}, \dots, a_h]^- > 1$. By Lemma 5 we have

$$\frac{p}{q} = \frac{xr_l - r_{l-1}}{xs_l - s_{l-1}}.$$

Using this identity and (8) we obtain

$$(9) \quad \begin{aligned} \frac{Q}{P} - \frac{q}{p} &\geq \frac{s_l + r_{l-1} - \varepsilon}{r_l} - \frac{xs_l - s_{l-1}}{xr_l - r_{l-1}} \\ &= \frac{(r_{l-1} - \varepsilon)(xr_l - r_{l-1}) + r_ls_{l-1} - s_lr_{l-1}}{r_l(xr_l - r_{l-1})} \\ &= \frac{(r_{l-1} - \varepsilon)(xr_l - r_{l-1}) - 1}{r_l(xr_l - r_{l-1})} \\ &\geq \frac{(xr_l - r_{l-1}) - 1}{r_l(xr_l - r_{l-1})} \\ &> 0, \end{aligned}$$

where we used the identity $r_ls_{l-1} - s_lr_{l-1} = -1$ from Lemma 5 to obtain the third line, $r_{l-1} - \varepsilon \geq 1$ to obtain the fourth line, and finally that $xr_l - r_{l-1} > 1$, which follows from the facts that $r_l \geq r_{l-1} + 1$ and $x > 1$. This gives the desired inequality, proving the claim. \square

The claim together with (4) proves that $\sum_{i=1}^k \frac{q_i}{p_i} \leq 1$ with equality only if $w = e_1$ has pairing ± 1 with each starting vertex, which completes the proof. \square

4. Definite 4-manifolds and the Seifert fibered space inequality

Now we consider when gluing two 4-manifolds can result in a closed definite 4-manifold.

Proposition 7. *Let U_1 and U_2 be connected 4-manifolds with $\partial U_1 = -\partial U_2 = Y$. Then the closed 4-manifold $X = U_1 \cup_Y U_2$ is positive definite if and only if*

- (a) *the inclusion-induced map $(i_1)_* \oplus (i_2)_* : H_1(Y; \mathbb{Q}) \rightarrow H_1(U_1; \mathbb{Q}) \oplus H_1(U_2; \mathbb{Q})$ is injective, and*
- (b) *for $i = 1, 2$, U_i is positive semidefinite, or equivalently has signature*

$$\sigma(U_i) = b_2(U_i) + b_1(U_i) - b_3(U_i) - b_2(Y).$$

Proof. In this proof all homology groups will be taken with rational coefficients. First, for $i = 1, 2$, consider the following segment of the long exact sequence in homology of the pair (U_i, Y) :

$$(10) \quad 0 \rightarrow H_3(U_i) \rightarrow H_3(U_i, Y) \rightarrow H_2(Y) \rightarrow H_2(U_i) \xrightarrow{Q} H_2(U_i, Y),$$

where Q can be represented by the intersection form matrix with respect to suitable bases. Hence, the null space of Q is precisely the image of $H_2(Y) \rightarrow H_2(U_i)$. By exactness and Lefschetz duality, the rank of the map $H_2(Y) \rightarrow H_2(U_i)$ is $b_2(Y) - b_1(U_i) + b_3(U_i)$. This gives an upper bound on the signature of U_i :

$$(11) \quad \sigma(U_i) \leq b_2(U_i) + b_1(U_i) - b_3(U_i) - b_2(Y),$$

with equality if and only if U_i is positive semidefinite. Now consider the segment of the Mayer–Vietoris sequence

$$(12) \quad 0 \rightarrow H_3(U_1) \oplus H_3(U_2) \rightarrow H_3(X) \rightarrow H_2(Y) \rightarrow H_2(U_1) \oplus H_2(U_2) \\ \rightarrow H_2(X) \rightarrow H_1(Y) \rightarrow H_1(U_1) \oplus H_1(U_2) \rightarrow H_1(X) \rightarrow 0.$$

The last three terms in this sequence show that

$$(13) \quad b_1(U_1) + b_1(U_2) \leq b_1(Y) + b_1(X),$$

with equality if and only if the map induced by the inclusions

$$(i_1)_* \oplus (i_2)_* : H_1(Y) \rightarrow H_1(U_1) \oplus H_1(U_2)$$

is injective.

Since the Euler characteristic of an exact sequence is zero, (12) shows that

$$(14) \quad b_2(X) = 2b_1(X) + \sum_{i=1}^2 (b_2(U_i) - b_1(U_i) - b_3(U_i)),$$

where we also used that $b_1(Y) = b_2(Y)$ and $b_1(X) = b_3(X)$.

By Novikov additivity, we have that $\sigma(X) = \sigma(U_1) + \sigma(U_2)$. So by summing the inequalities in (11) for $i = 1, 2$ and comparing with (14) we obtain

$$(15) \quad b_2(X) \geq 2(b_1(X) + b_2(Y) - b_1(U_1) - b_1(U_2)) + \sigma(X),$$

with equality if and only if we have equality in (11) for both $i = 1, 2$. Hence, X can be positive definite if and only if

$$(16) \quad b_1(U_1) + b_1(U_2) = b_2(Y) + b_1(X)$$

and we have equality in (11) for $i = 1, 2$. However we have already seen that equality occurs in (13) if and only if $(i_1)_* \oplus (i_2)_*$ is injective. \square

This allows us to prove the main theorem.

Theorem 1. *Let $Y = S^2(e; \frac{p_1}{q_1}, \dots, \frac{p_k}{q_k})$ be a Seifert fibered space over S^2 in standard form, that is, with $e \geq 0$, $\frac{p_i}{q_i} > 1$ for all $i \in \{1, 2, \dots, k\}$ and $\varepsilon(Y) \geq 0$. Suppose that Y bounds a smooth 4-manifold W such that $\sigma(W) = -b_2(W)$ and the inclusion induced map $H_1(Y; \mathbb{Q}) \rightarrow H_1(W; \mathbb{Q})$ is injective. Then there is a partition of $\{1, 2, \dots, k\}$ into at most e classes such that for each class C ,*

$$\sum_{i \in C} \frac{q_i}{p_i} \leq 1.$$

Proof. Let X be the standard positive (semi)definite plumbing 4-manifold with $\partial X = Y$, and let $Z = X \cup_Y -W$. It follows from Proposition 7 that Z is positive definite. Condition (b) of Proposition 7 holds, since $\sigma(W) = -b_2(W)$ implies that $-W$ is positive definite. Thus, Z is a smooth positive definite 4-manifold, so by Donaldson's theorem Z has standard positive diagonal intersection form. The inclusion $X \subset Z$ induces a map $H_2(X) \rightarrow H_2(Z)$ which preserves the intersection pairing. Thus, there is a map of lattices $(H_2(X), Q_X) \rightarrow (\mathbb{Z}^m, \text{Id})$ for some $m > 0$.

We construct a partition of $\{1, 2, \dots, k\}$ into at most e classes as follows. Denote the orthonormal basis of coordinate vectors of $(\mathbb{Z}^m, \text{Id})$ by $\{e_1, \dots, e_m\}$. For $v \in (H_2(X), Q_X)$, we call $\{e_i : 1 \leq i \leq m, e_i \cdot v \neq 0\}$ the support of v . Without loss of generality, we may assume that the central vertex has support $\{e_1, e_2, \dots, e_n\}$, where $n \leq e$. Let v_1, v_2, \dots, v_k be the vertices of the plumbing adjacent to the central vertex, so that v_i is a vertex belonging to the i -th leg of the plumbing graph (with fraction p_i/q_i). For $i \in \{1, \dots, n\}$, let $B_i = \{1 \leq j \leq k \mid v_j \cdot e_i \neq 0\}$ and define $B_0 = \emptyset$. Let $C_i = B_i \setminus \bigcup_{j < i} B_j$ for $i \in \{1, \dots, n\}$. Then C_1, \dots, C_n are disjoint and $\bigcup_i C_i = \{1, \dots, k\}$. Thus the nonempty classes $\{C_i : C_i \neq \emptyset\}$ form a partition of $\{1, 2, \dots, k\}$ into at most e classes. By definition for each $i \in \{1, 2, \dots, n\}$, the starting vertices of the linear chains indexed by C_i all have support containing the common unit vector e_i . Hence, by Theorem 6, we have that $\sum_{j \in C_i} \frac{q_j}{p_j} \leq 1$. \square

5. The Neumann–Zagier question

We prove Theorem 8 below which, when combined with Donaldson's theorem, immediately implies Theorem 2. Note that the following theorem also positively answers the Neumann–Zagier question stated in the introduction.

Theorem 8. *Let $Y = S^2(e; \frac{p_1}{q_1}, \dots, \frac{p_k}{q_k})$, $k \geq 3$, be in standard form, that is, with $\frac{p_i}{q_i} > 1$ for all $i \in \{1, 2, \dots, k\}$, $e > 0$ and with Y bounding a smooth positive definite plumbing X . Suppose that $|H_1(Y)| \in \{1, 2, 3, 5, 6, 7\}$ and the intersection lattice $(H_2(X), Q_X)$ embeds into a positive standard diagonal lattice. Then $e = 1$.*

Proof. For sake of contradiction, assume that $e > 1$. We may apply Theorem 1, noting that the existence of W in the hypothesis of Theorem 1 is only required to ensure that there is a map of lattices of $(H_2(X), Q_X)$ into a positive standard diagonal lattice. Hence, there is a partition $\{C_1, \dots, C_n\}$ of $\{1, \dots, k\}$ into $n \leq e$ classes. Moreover, for each class C , $1 - \sum_{i \in C} \frac{q_i}{p_i} \geq 0$, and we call C *complementary* if equality occurs, and *noncomplementary* otherwise.

We have

$$\begin{aligned}
 (17) \quad |H_1(Y)| &= p_1 \cdots p_k \cdot \varepsilon(Y) = p_1 \cdots p_k \left(e - \sum_{i=1}^k \frac{q_i}{p_i} \right) \\
 &= p_1 \cdots p_k \left((e - n) + \sum_{i=1}^n \left(1 - \sum_{j \in C_i} \frac{q_j}{p_j} \right) \right) \\
 &= p_1 \cdots p_k (e - n) + \sum_{i=1}^n a_i \prod_{\substack{1 \leq l \leq k \\ l \notin C_i}} p_l,
 \end{aligned}$$

where $a_i = (\prod_{j \in C_i} p_j) \cdot (1 - \sum_{l \in C_i} \frac{q_l}{p_l})$ is an integer for all $i \in \{1, \dots, n\}$. Notice that all terms in (17) are nonnegative integers. Since we are assuming that $|H_1(Y)| \in \{1, 2, 3, 5, 6, 7\}$, we must have $n = e$, otherwise $|H_1(Y)| \geq p_1 \cdots p_k \geq 2 \cdot 2 \cdot 2 = 8$ since $k \geq 3$.

We claim that $|C_i| \leq k - 2$ for some $i \in \{1, \dots, e\}$ with C_i noncomplementary. To see this, we argue as follows. There are $n = e \geq 2$ classes in the partition, and at least one noncomplementary class since $|H_1(Y)| > 0$. If there are two noncomplementary classes then at least one has size at most $k - 2$ since $k \geq 3$. If there is only one noncomplementary class, then there is a complementary class which necessarily has size at least 2, and hence the noncomplementary class satisfies the claim.

Combining the above claim with (17), we see that $|H_1(Y)|$ is a sum of integers greater than 1, and at least one of these integers is not prime. For $|H_1(Y)| \in \{1, 2, 3, 5, 6, 7\}$, this is only possible for $|H_1(Y)| = 7$ with decomposition $7 = 3 + 2 \cdot 2$, and for $|H_1(Y)| = 6$ with the two decompositions $|H_1(Y)| = 2 + 2 \cdot 2 = 2 \cdot 3$. We address these cases in turn. For $|H_1(Y)| = 7 = 3 + 2 \cdot 2$, comparing this decomposition with (17), we see that there must exist some noncomplementary C_i with $|C_i| = 2$ and $p_j = 2$ for all $j \in C_i$. However, such a C_i must be complementary since $1 - \frac{1}{2} - \frac{1}{2} = 0$, a contradiction. A similar argument rules out the decomposition $|H_1(Y)| = 2 + 2 \cdot 2$. Finally, in the case $|H_1(Y)| = 2 \cdot 3$, the decomposition implies

that there exists a complementary class $C_i = \{a, b\}$ with $p_a = 2$ and $p_b = 3$, which is impossible. \square

We obtain the following corollary, answering a question of Lidman and Tweedy [2018, Remark 4.3].

Corollary 3. *Let Y be a Seifert fibered integral homology sphere, and let $e \in \mathbb{Z}$ be the central weight in the standard definite plumbing graph for Y . If $|e| \neq 1$, then $d(Y) \neq 0$.*

Proof. We prove the contrapositive. Assume that $d(Y) = 0$. Note that reversing the orientation of Y simply changes the sign of the weight of the central vertex in the definite plumbing bounding Y . Thus, by reversing the orientation of Y if necessary we assume that Y bounds a smooth negative definite plumbing X^4 . Let

$$C = \{ \xi \in H_2(X; \mathbb{Z}) \mid \xi \cdot v = v \cdot v \pmod{2} \text{ for all } v \in H_2(X; \mathbb{Z}) \}$$

be the set of characteristic vectors, and let $n = \text{rk}(H_2(X))$. Elkies [1995] proved that $0 \leq n + \max_{\xi \in C} \xi \cdot \xi$, with equality if and only if Q_X is diagonalizable over \mathbb{Z} . However, it follows from [Ozsváth and Szabó 2003, Theorem 9.6] that $n + \max_{\xi \in C} \xi \cdot \xi \leq 4d(Y) = 0$. Therefore Q_X is diagonalizable over \mathbb{Z} , in particular $(H_2(-X), Q_{-X})$ embeds into a positive standard diagonal lattice. Hence, Theorem 8 implies that $|e| = 1$. \square

6. Seifert fibered spaces bounding rational homology $S^1 \times D^3$'s

In this section we prove Theorem 4, which in particular gives a classification of the Seifert fibered spaces which smoothly bound rational homology $S^1 \times D^3$'s. We note that the implication (2) implies (1) was proved by Donald [2015, Proof of Theorem 1.3], and the equivalence of (1) and (2) was shown by Aceto [2015, Theorem 1.2].

Theorem 4. *Let Y be a Seifert fibered space over S^2 with*

$$H_*(Y; \mathbb{Q}) \cong H_*(S^1 \times S^2; \mathbb{Q}).$$

The following are equivalent:

- (1) *Y is of the form $S^2(k; \frac{p_1}{q_1}, \frac{p_1}{p_1 - q_1}, \dots, \frac{p_k}{q_k}, \frac{p_k}{p_k - q_k})$, where $k \geq 0$ and $\frac{p_i}{q_i} > 1$ for all $i \in \{1, \dots, k\}$.*
- (2) *$Y = \partial W$, where W is a smooth 4-manifold with $H_*(W; \mathbb{Q}) \cong H_*(S^1 \times D^3; \mathbb{Q})$.*
- (3) *Y is the boundary of smooth 4-manifolds W_+ and W_- such that $\sigma(W_{\pm}) = \pm b_2(W_{\pm})$ and each of the inclusion-induced maps $H_1(Y; \mathbb{Q}) \rightarrow H_1(W_{\pm}; \mathbb{Q})$ is injective.*

Proof. First suppose that (1) holds, that is, $Y = S^2(k; \frac{p_1}{q_1}, \frac{p_1}{p_1 - q_1}, \dots, \frac{p_k}{q_k}, \frac{p_k}{p_k - q_k})$, where $k \geq 0$ and $\frac{p_i}{q_i} \in \mathbb{Q}_{>1}$ for all $i \in \{1, \dots, k\}$. By Rolfsen twisting, Y can be put into the form $S^2(0; \frac{p_1}{q_1}, -\frac{p_1}{q_1}, \dots, \frac{p_k}{q_k}, -\frac{p_k}{q_k})$. Let $M = S^2(0; \frac{p_1}{q_1}, \frac{p_2}{q_2}, \dots, \frac{p_k}{q_k})$, let M° be the 3-manifold with torus boundary given by removing a tubular neighborhood of a regular fiber of M and let $W = M \times [0, 1]$. Then $\partial W = M^\circ \cup_\partial -M^\circ$ is the double of M° , which is precisely Y . Finally, notice that

$$H_*(W; \mathbb{Q}) = H_*(M^\circ; \mathbb{Q}) = H_*(S^1 \times D^3; \mathbb{Q}),$$

where the last equality follows from the fact that M is a rational homology S^3 and M° is obtained by removing a neighborhood of a simple closed curve from M . This proves (2).

The implication (2) implies (3) holds by taking $W_\pm = W$ and noting that $H_1(Y; \mathbb{Q}) \rightarrow H_1(W; \mathbb{Q})$ is injective by the long exact sequence of the pair (Y, W) .

Finally assume that (3) holds. Hence, Y is the boundary of smooth 4-manifolds W_+ and W_- satisfying $\sigma(W_\pm) = \pm b_2(W_\pm)$ and such that the inclusion induced maps $H_1(Y) \rightarrow H_1(W_\pm)$ are injective. Write Y as $S^2(e; \frac{p_1}{q_1}, \dots, \frac{p_k}{q_k})$ with $k \geq 3$ and $\frac{p_i}{q_i} \in \mathbb{Q}_{>1}$ for all $i \in \{1, \dots, k\}$. Notice that $-Y = S^2(k - e; \frac{p_1}{p_1 - q_1}, \dots, \frac{p_k}{p_k - q_k})$, and Y is of the form given in (1) if and only if $-Y$ is of this form. Thus, by reversing the orientations of both Y and W_\pm if necessary, we may assume that $e \geq \frac{k}{2}$.

By Theorem 1, there is a partition $\{C_1, \dots, C_n\}$ of $\{1, \dots, k\}$ into $n \leq e$ classes such that for each class C , $1 - \sum_{i \in C} \frac{q_i}{p_i} \geq 0$. Since Y is a rational homology $S^1 \times S^2$, we thus have

$$\begin{aligned} 0 &= p_1 \cdots p_k \cdot \varepsilon(Y) = p_1 \cdots p_k \left(e - \sum_{i=1}^k \frac{q_i}{p_i} \right) \\ &= p_1 \cdots p_k \left((e - n) + \sum_{i=1}^n \left(1 - \sum_{j \in C_i} \frac{q_j}{p_j} \right) \right), \end{aligned}$$

where all terms in the sum are nonnegative. Hence, we must have $n = e$ and $1 - \sum_{i \in C} \frac{q_i}{p_i} = 0$, for all $i \in \{1, \dots, n\}$. This implies that $|C_i| \geq 2$ for all $i \in \{1, \dots, n\}$. Thus, there are at least $2n = 2e$ legs, so $e \leq \frac{k}{2}$. However, by assumption $e \geq \frac{k}{2}$ so $e = \frac{k}{2}$ and $|C_i| = 2$ for all $i \in \{1, \dots, n\}$. Thus, C_1, \dots, C_n partition $\{1, \dots, k\}$ into pairs of indices indexing pairs of fractions of the form $\frac{p}{q}, \frac{p}{p-q} \in \mathbb{Q}_{>1}$, and thus (1) holds. \square

Acknowledgements

Issa would like to thank Cameron Gordon for his support and encouragement, and Josh Greene for a helpful conversation on the Neumann–Zagier question. Both authors would like to thank James Davis for his comment on an earlier version of this paper and an anonymous referee for their feedback.

References

- [Aceto 2015] P. Aceto, “Rational homology cobordisms of plumbed 3-manifolds”, preprint, 2015. arXiv
- [Donald 2015] A. Donald, “Embedding Seifert manifolds in S^4 ”, *Trans. Amer. Math. Soc.* **367**:1 (2015), 559–595. MR Zbl
- [Donaldson 1987] S. K. Donaldson, “The orientation of Yang–Mills moduli spaces and 4-manifold topology”, *J. Differential Geom.* **26**:3 (1987), 397–428. MR Zbl
- [Elkies 1995] N. D. Elkies, “A characterization of the \mathbb{Z}^n lattice”, *Math. Res. Lett.* **2**:3 (1995), 321–326. MR Zbl
- [Fintushel and Stern 1985] R. Fintushel and R. J. Stern, “Pseudofree orbifolds”, *Ann. of Math.* (2) **122**:2 (1985), 335–364. MR Zbl
- [Issa and McCoy 2018] A. Issa and D. McCoy, “Smoothly embedding Seifert fibered spaces in S^4 ”, preprint, 2018. arXiv
- [Lecuona and Lisca 2011] A. G. Lecuona and P. Lisca, “Stein fillable Seifert fibered 3-manifolds”, *Algebr. Geom. Topol.* **11**:2 (2011), 625–642. MR Zbl
- [Lidman and Tweedy 2018] T. Lidman and E. Tweedy, “A note on concordance properties of fibers in Seifert homology spheres”, *Canad. Math. Bull.* **61**:4 (2018), 754–767. MR Zbl
- [Lisca 2007] P. Lisca, “Sums of lens spaces bounding rational balls”, *Algebr. Geom. Topol.* **7** (2007), 2141–2164. MR Zbl
- [Neumann 1989] W. D. Neumann, “On bilinear forms represented by trees”, *Bull. Austral. Math. Soc.* **40**:2 (1989), 303–321. MR Zbl
- [Neumann and Raymond 1978] W. D. Neumann and F. Raymond, “Seifert manifolds, plumbing, μ -invariant and orientation reversing maps”, pp. 163–196 in *Algebraic and geometric topology* (Santa Barbara, CA, 1977), edited by K. C. Millett, Lecture Notes in Math. **664**, Springer, Berlin, 1978. MR Zbl
- [Neumann and Zagier 1985] W. D. Neumann and D. Zagier, “A note on an invariant of Fintushel and Stern”, pp. 241–244 in *Geometry and topology* (College Park, MD, 1983/84), edited by J. Alexander and J. Harer, Lecture Notes in Math. **1167**, Springer, Berlin, 1985. MR Zbl
- [Ozsváth and Szabó 2003] P. Ozsváth and Z. Szabó, “Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary”, *Adv. Math.* **173**:2 (2003), 179–261. MR Zbl

Received August 29, 2018. Revised August 20, 2019.

AHMAD ISSA
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF BRITISH COLUMBIA
 VANCOUVER, BC
 CANADA
 aissa@math.ubc.ca

DUNCAN MCCOY
 DÉPARTEMENT DE MATHÉMATIQUES
 UNIVERSITÉ DU QUÉBEC À MONTRÉAL
 MONTRÉAL, QC
 CANADA
 duncan.mccoy@cirget.ca

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 304 No. 2 February 2020

Graphs admitting only constant splines	385
KATIE ANDERS, ALISSA S. CRANS, BRIANA FOSTER-GREENWOOD, BLAKE MELLOR and JULIANNA TYMOCZKO	
Centers of disks in Riemannian manifolds	401
IGOR BELEGRADEK and MOHAMMAD GHOMI	
The geometry of the flex locus of a hypersurface	419
LAURENT BUSÉ, CARLOS D'ANDREA, MARTÍN SOMBRA and MARTIN WEIMANN	
Morse inequalities for Fourier components of Kohn–Rossi cohomology of CR covering manifolds with S^1 -action	439
RUNG-TZUNG HUANG and GUOKUAN SHAO	
On Seifert fibered spaces bounding definite manifolds	463
AHMAD ISSA and DUNCAN MCCOY	
Regularity of quotients of Drinfeld modular schemes	481
SATOSHI KONDO and SEIDAI YASUDA	
Sums of algebraic trace functions twisted by arithmetic functions	505
MAXIM KOROLEV and IGOR SHPARLINSKI	
Twisted calculus on affinoid algebras	523
BERNARD LE STUM and ADOLFO QUIRÓS	
Symplectic (-2) -spheres and the symplectomorphism group of small rational 4-manifolds	561
JUN LI and TIAN-JUN LI	
Addendum to the article Contact stationary Legendrian surfaces in \mathbb{S}^5	607
YONG LUO	
The Hamiltonian dynamics of magnetic confinement in toroidal domains	613
GABRIEL MARTINS	
Gluing Bartnik extensions, continuity of the Bartnik mass, and the equivalence of definitions	629
STEPHEN MCCORMICK	
Decomposable Specht modules indexed by bihooks	655
LIRON SPEYER and LOUISE SUTTON	
On the global well-posedness of one-dimensional fluid models with nonlocal velocity	713
ZHUAN YE	
Hopf cyclic cohomology for noncompact G -manifolds with boundary	753
XIN ZHANG	



0030-8730(2020)304:2;1-R