

*Pacific
Journal of
Mathematics*

**THE POINCARÉ HOMOLOGY SPHERE,
LENS SPACE SURGERIES, AND SOME KNOTS WITH
TUNNEL NUMBER TWO**

KENNETH L. BAKER

APPENDIX WRITTEN JOINTLY WITH NEIL R. HOFFMAN

THE POINCARÉ HOMOLOGY SPHERE, LENS SPACE SURGERIES, AND SOME KNOTS WITH TUNNEL NUMBER TWO

KENNETH L. BAKER

APPENDIX WRITTEN JOINTLY WITH NEIL R. HOFFMAN

We exhibit an infinite family of knots in the Poincaré homology sphere with tunnel number 2 that have a lens space surgery. Notably, these knots are not doubly primitive and provide counterexamples to a few conjectures. Additionally, we update (and correct) our earlier work on Hedden’s almost-simple knots. In the appendix, it is shown that a hyperbolic knot in the Poincaré homology sphere with a lens space surgery has either no symmetries or just a single strong involution.

1. A family of knots in the Poincaré homology sphere with lens space surgeries

Figure 1 shows a one-parameter family of arcs κ_n , $n \in \mathbb{Z}$, on the pretzel knot $P(-2, 3, 5) = P(3, 5, -2)$ for which a -1 -framed banding produces the two bridge knot $B(3n^2 + n + 1, -3n + 2)$. Indeed,

$$[n, -1, -n, 3] = -\frac{3n^2 + n + 1}{-3n + 2}.$$

Through double branched coverings and the Montesinos trick, the arc κ_n lifts to a knot K_n in the Poincaré homology sphere on which an integral surgery yields the lens space $L(3n^2 + n + 1, -3n + 2)$.

As presented, the arc κ_n lies in a bridge sphere that presents $P(-2, 3, 5)$ as a 3-bridge link. Thus the knot K_n lies in a genus 2 Heegaard surface for the Poincaré homology sphere \mathcal{P} . Nevertheless, as we shall see, generically K_n does not have tunnel number one.

Theorem 1. *For each integer n , the knot K_n in the Poincaré homology sphere \mathcal{P} has an integral surgery to the lens space $L(3n^2 + n + 1, -3n + 2)$. If $|n| \geq 4$, then K_n has tunnel number 2.*

MSC2010: 57M27.

Keywords: Poincaré homology sphere, Dehn surgery, lens space, tunnel number.




Figure 1. Top left shows a family of arcs κ_n with end points on the pretzel knot $P(-2, 3, 5)$. A numbered vertical box signifies a stack of half-twists where the sign informs the handedness. We perform a banding on κ_n (framed relative to the horizontal bridge sphere) followed by a sequence of isotopies to recognize the result as the two-bridge knot $B(3n^2 + n + 1, -3n + 2)$. The arc κ_n^* dual to the banding is also carried along.

These knots arose from an application of the “longitudinally jointly primitive” ideas presented in [Baker et al. 2019]. After observing that the negative Whitehead link is a two-component genus 1 fibered link for which -1 surgery on both components produces \mathcal{P} , we found a family of jointly primitive knots in the twice-punctured torus bundle that is the exterior of this Whitehead link. Our knots K_n are the images of these jointly primitive knots in the \mathcal{P} filling. This is discussed further in [Baker et al. 2019, §2.3.3].

Previously, the only known knots in \mathcal{P} admitting a surgery to a lens space are surgery duals to Tange knots [2009a] and certain Hedden knots [Hedden 2011;

[Rasmussen 2007; Baker 2014b]. (See Section 5 for a discussion of the Hedden knots.) These knots in \mathcal{P} are all *doubly primitive*, they may be presented as curves in a genus 2 Heegaard splitting that represent a generator in π_1 of each handlebody bounded by the Heegaard surface. We see this as Tange knots and Hedden knots all admit descriptions by doubly pointed genus 1 Heegaard diagrams. (Tange knots are all simple knots in lens spaces and Hedden knots are “almost simple”.) Equivalently, they are all 1-bridge knots in their lens spaces. Any integral surgery on a 1-bridge knot in a lens space naturally produces a 3-manifold with a genus 2 Heegaard surface in which the dual knot sits as a doubly primitive knot.

The Berge conjecture posits that any knot in S^3 with a lens space surgery is a doubly primitive knot [Berge 1990]; see [Gordon 1991, Question 5.5]. It is also proposed that any knot in $S^1 \times S^2$ with a lens space surgery is a doubly primitive knot [Baker et al. 2016, Conjecture 1.1; Greene 2013, Conjecture 1.9]. By analogy and due to a sense of simplicity¹, one may suspect that any knot in \mathcal{P} with a lens space surgery should also be doubly primitive. However this is not the case. Since our knots K_n have tunnel number 2 for $|n| \geq 4$, they cannot be doubly primitive.

Corollary 2. *There are knots in the Poincaré homology sphere with lens space surgeries that are not doubly primitive. That is, the analogy to the Berge conjecture fails for the Poincaré homology sphere.*

Corollary 3. *[Greene 2013, Conjecture 1.10] is false. For example, the lens space $L(5, 1)$ contains a knot K_1^* with a surgery to \mathcal{P} but does not contain a Tange knot or a Hedden knot with a \mathcal{P} surgery.*

Proof. For $n = 1$, surgery on K_1 produces $L(5, -1)$ as shown in Figure 1. A quick check of [Tange 2009a, Table 2] shows that $L(5, -1)$ contains no Tange knots. The Hedden knots with surgery to $\pm\mathcal{P}$ are homologous to Berge lens space knots of type VII [Rasmussen 2007; Greene 2013]: one further observes that $L(5, -1)$ does not contain any of these knots since $k^2 + k + 1 = 0 \pmod{5}$ has no integral solution. \square

Remark 4. The knot K_1 is actually the $(2, 3)$ -cable of the knot J that is surgery dual to the trefoil in S^3 .

As we shall see in Theorem 7(3), it is the $(2g(K_n) - 1)$ -surgery on our knots K_n that produces a lens space; $g(K_n)$ is the Seifert genus of the knot K_n . Hence they also provide counterexamples to [Hedden 2011, Conjecture 1.7] (see [Baker et al. 2008, Conjecture 1.6]) and [Greene 2013, Conjecture 1.10]. It seems plausible that it is the largeness of the knot genus with respect to the lens space surgery

¹The manifolds S^3 , \mathcal{P} , and its mirror $-\mathcal{P}$ are the only homology 3-spheres with finite fundamental group (by Perelman) and the only known irreducible L-space homology 3-spheres, e.g., [Eftekhary 2009; Eftekhary 2018].

slope that enables the failure of our knots to be doubly primitive. Hedden and Rasmussen also observe a distinction at this slope for lens space surgeries on knots in L-space homology spheres [Hedden 2011; Rasmussen 2007]. Indeed, from the view of Heegaard Floer homology for integral slopes on knots in homology spheres, this slope is right at the threshold at which a knot could have an L-space surgery [Ozsváth and Szabó 2011, Proposition 9.6] (see also [Hedden 2009, Lemma 2.13])² and just below what implies it has simple knot Floer homology [Eftekhary 2011]³. With this in mind, we adjust and update [Greene 2013, Conjecture 1.10].

Conjecture 5. *Suppose that p -surgery on a knot K in the Poincaré homology sphere produces a lens space. If $p > 2g(K) - 1$ is an integer, then K is a doubly primitive knot. Furthermore it is surgery dual to one of the Tange knots.*

Conjecture 6. *If a knot in the Poincaré homology sphere is doubly primitive, then it is surgery dual to one of the Tange knots or one of the Hedden knots.*

As mentioned in the paragraph following its statement, [Hedden 2011, Theorem 3.1] (presented here in Theorem 14) applies to any L-space homology sphere, not just S^3 . Thus, since doubly primitive knots are surgery dual to one-bridge knots in lens spaces, [Hedden 2011, Theorem 3.1] implies that such surgery duals are either simple knots or one of the Hedden knots T_L or T_R . In Section 5 we clarify and correct our work in [Baker 2014b] and further classify when Hedden's knots are dual to integral surgeries on knots homology spheres, \mathcal{P} in particular. Tange has produced a list of simple knots in lens spaces that are surgery dual to knots in \mathcal{P} [Tange 2009a], and it is expected that this list is complete. Verifying its completeness will affirm Conjecture 6.

Let us collect various properties of our knots.

Theorem 7. *Let $p = 3n^2 + n + 1$ and $q = -3n + 2$. Then the following hold for our family of knots K_n in \mathcal{P} :*

- (1) *Positive p -surgery on K_n produces a lens space $L(p, q)$.*
- (2) *K_n is fibered and supports the tight contact structure on \mathcal{P} .*
- (3) $g(K_n) = \frac{1}{2}(p + 1)$
- (4) $\text{rk HFK}(L(p, q), K_n^*) = p + 2$
- (5) *Let T_n be the $(3n + 1, n)$ -torus knot in S^3 , and note that p -surgery on T_n also produces $L(p, q)$. The surgery dual to K_n is homologous to the surgery dual to T_n .*
- (6) $\Delta_{K_n}(t) = \Delta_{T_n}(t) - (t^{(p-1)/2} + t^{-(p-1)/2}) + (t^{(p+1)/2} + t^{-(p+1)/2})$.

²This threshold may be lower for knots in homology spheres with $\tau(K) < g(K)$.

³A knot K^* in a rational homology sphere Y has *simple knot Floer homology* if $\text{rk HFK}(Y, K^*) = \text{rk HF}(Y)$. This definition does not require Y to be an L-space itself.

Theorem 7(5) is given as **Lemma 12**. The remainder of **Theorem 7** follows from assembling the works [**Hedden 2011**; **Rasmussen 2007**; **Tange 2011, 2009b**]. In fact, appealing to Greene’s proof of the lens space realization problem [**Greene 2013**], we tease out the following general theorem from which **Theorem 7(3)** and (6) follow.

Theorem 8. *Suppose that p -surgery on knot K in an L -space homology sphere Y with $d(Y) = 2$ produces the lens space $L(p, q)$. Then $p = 2g(K) - 1$ if and only if p -surgery on some Berge knot B in S^3 also produces the lens space $L(p, q)$ in which the surgery duals to K and B are homologous.*

When this holds, $\Delta_K(t) = \Delta_B(t) - (t^{(p-1)/2} + t^{-(p-1)/2}) + (t^{(p+1)/2} + t^{-(p+1)/2})$.

2. Questions about knots in the Poincaré homology sphere with lens space surgeries.

2.1. Homology classes of knots.

Question 9. (1) For which Berge knots B that have a positive, odd p -surgery producing the lens space $L(p, q)$, is there a knot K in \mathcal{P} such that $2g(K) - 1 = p$, p -surgery produces the lens space $L(p, q)$, and the surgery duals B^* and K^* are homologous?

(2) For a Berge knot B as above, can there be two distinct knots K_1 and K_2 in \mathcal{P} with p -surgery to $L(p, q)$ such that the surgery duals B^* , K_1^* , and K_2^* are all homologous?

Presently the only Berge knots known to answer **Question 9(1)** are the Berge knots of type VII (those that embed in the fiber of the trefoil) due to [**Hedden 2011**; **Baker 2014b**] and the $(3n + 1, n)$ -torus knots with $p = 3n^2 + n + 1$ due to our knots introduced here. Indeed, what about the case of the $(3n + 1, n)$ -torus knots when $p = 3n^2 + n - 1$?

To clarify the word “distinct” in **Question 9(2)**, note that the mapping class group of the Poincaré homology sphere is trivial [**Boileau and Otal 1991**, **Théorém 3** and **Corollaire 4**]. Thus knots in \mathcal{P} that are related by a diffeomorphism of \mathcal{P} are also isotopic.

2.2. Symmetries. One may observe that, thus far, all the known examples of knots in \mathcal{P} with a lens space surgery are strongly invertible. Hence there is an involution of \mathcal{P} taking each of these knots with some orientation to its reverse. Wang and Zhou [**1992**] have shown that if a knot in S^3 other than a torus knots has a nontrivial lens space surgery, then its only possible symmetry is a strong involution.

Question 10. (1) Is every knot in \mathcal{P} with a lens space surgery strongly invertible?

(2) What are the possible symmetries of a knot in \mathcal{P} with a lens space surgery?

In the appendix to this article, the author and Hoffman provide [Theorem A.1](#) which addresses [Question 10\(2\)](#): A hyperbolic knot in \mathcal{P} with a lens space surgery either has no symmetries or just a single strong involution.

2.3. Hopf plumbings. Let us inquire about another potential analogy with the Berge knots in S^3 . The Giroux correspondence says that any two fibered knots supporting the same contact structure are related by a sequence of plumbings and de-plumbings of positive Hopf bands [[Giroux 2002](#)]. Since the Berge knots (with positive surgeries to lens spaces) can all be expressed as closures of positive braids⁴, each Berge knot can be obtained from the unknot by a sequence of plumbings of Hopf bands. No de-plumbings are necessary.

In \mathcal{P} the unknot is not fibered, but the knot J that is -1 -surgery dual to the (negative) trefoil is. As we shall see in [Lemma 13](#), J is a reasonable surrogate for the unknot in \mathcal{P} : it is the unique genus one fibered knot in \mathcal{P} , and it supports the tight contact structure on \mathcal{P} . Since a knot in \mathcal{P} with a lens space surgery is fibered and supports the tight contact structure (by [Theorem 7\(2\)](#)), the Giroux correspondence says that it may be obtained from J by a sequence of plumbings and de-plumbings of positive Hopf bands.

Question 11. If a knot in \mathcal{P} has a lens space surgery, then can it be obtained from J by a sequence of plumbings of Hopf bands?

We have yet to systematically check this question for the knots produced here or for the duals to the Tange and Hedden knots. As noted in [Remark 4](#), the knot K_1 is the $(2, 3)$ -cable of J . One may use this to show that K_1 is indeed obtained by plumbing two Hopf bands onto J .

3. Notation and conventions

Let K be a knot in an oriented 3-manifold M . Choose an orientation on K and let μ be a meridian of K in the torus $\partial\mathcal{N}(K)$ that positively links K . Let λ be an oriented curve in $\partial\mathcal{N}(K)$ that is isotopic to K in $\mathcal{N}(K)$; if K is null-homologous in M , we choose λ so that it is null-homologous in $M - \mathcal{N}(K)$. If γ is an essential simple closed curve in $\partial\mathcal{N}(K)$, then when it is oriented $[\gamma] = p[\mu] + q[\lambda] \in H_1(\partial\mathcal{N}(K))$ where p, q are coprime integers; changing the orientation of γ changes the signs of both p and q . We refer to both the unoriented isotopy class of γ in $\partial\mathcal{N}(K)$ and the number $p/q \in \mathbb{Q} \cup \{\infty\}$ as a *slope* and (when λ is null-homologous) say it is *positive* if $0 < p/q < \infty$. If the slope is integral, we also say it is a *longitude* or a *framing* of K .

The result of γ -Dehn surgery on the knot $K \subset M$ is the manifold $M_K(\gamma)$ obtained by attaching $S^1 \times D^2$ to $M - \mathcal{N}(K)$ along the torus $\partial\mathcal{N}(K)$ so that $\text{pt} \times \partial D^2$ is

⁴This is remarked preceding the conjecture of Goda and Teragaito [[2000](#)], for example.

identified with the slope γ . The image of the curve $S^1 \times \text{pt}$ in $M_K(\gamma)$ is a knot called the *surgery dual* to K and denoted K^* . Observe that γ is the meridian of K^* and μ -surgery on K^* returns M with surgery dual $K = (K^*)^*$.

If M is the double branched cover of S^3 over a link L , an arc κ such that $\kappa \cap L = \partial\kappa$ lifts to a knot K in the cover M . Via the Montesinos trick [1975], each integral surgery on K corresponds to a *banding* along κ . With $I = [-1, 1]$, if $R = I \times I$ is a disk embedded in S^3 such that $R \cap L = \partial I \times I$ and $\{0\} \times I = \kappa$, then the link $L' = (L - \partial I \times I) \cup I \times \partial I$ is the result of a banding along κ . The arc $\kappa^* = I \times \{0\}$ is the dual arc of the banding. In the double branched cover over L' , κ^* lifts to the dual knot K^* of the corresponding integral surgery on K .

The lens space $L(p, q)$ is defined to be the manifold that results from $-p/q$ -surgery on the unknot in S^3 . The two-bridge link $B(p, q)$ is the link in S^3 whose double branched cover is $L(p, q)$ [Hodgson and Rubinstein 1985]. Using the continued fraction $[x_1, x_2, \dots, x_n] = x_1 - 1/(x_2 - 1/(\dots - 1/x_n))$ to express $-p/q$ describes the two-bridge link $B(p, q)$ geometrically in plat presentation as in the lower left of Figure 1.

A nontrivial knot in a lens space $L(p, q)$ is *simple* (or *grid number one*) if, in the standard genus one Heegaard diagram of the lens space, it may be represented by two of the p intersection points. The simple knot is then the union of the arcs connecting those points in the two meridional disks whose boundaries are described by the diagram. Equivalently, a simple knot (including the trivial knot) is a knot represented by a doubly pointed genus 1 Heegaard diagram of $L(p, q)$ that has p intersection points. There is one for each homology class $k \in H_1(L(p, q))$ in $L(p, q)$, denoted $K(p, q, k)$. See for example [Rasmussen 2007; Hedden 2011; Baker et al. 2008].

4. Proofs

Proof of Theorem 1. Figure 1 exhibits arcs κ_n on the pretzel knot $P(-2, 3, 5)$ that have a banding to a two-bridge link. By passing to the double branched covers (and using the Montesinos trick), this describes knots K_n in \mathcal{P} that have an integral surgery to a lens space.

Since K_n lies in a genus 2 Heegaard surface, its tunnel number is at most 2. If K_n were to have tunnel number one, then every surgery on K_n would have Heegaard genus at most 2. However we will see that surgery along the framing induced by the Heegaard surface produces a toroidal manifold which, according to Kobayashi's classification [1984], does not have Heegaard genus 2. (The same scheme was recently employed in [Eudave-Muñoz et al. 2015] to demonstrate a family of strongly invertible knots in S^3 with a Seifert fibered surgery and tunnel number 2.)




Figure 2. Left: The bridge sphere S for $P(-2, 3, 5)$ splits $\kappa_n(0)$ into two tangles, T_n^α above and T_n^β below. Middle: Filling T_n^α with a rational tangle to get $T(2, n+2)\#T(2, 3)$ makes T_n^β into $P(1-n, -2, 2)$. Right: Filling T_n^β with a rational tangle to get $T(2, 1-n)\#T(2, 2)$ makes T_n^α into $P(n+3, 3, -2)$.

Kobayashi shows that if $M = M^\alpha \cup_T M^\beta$ is a genus 2 manifold decomposed along a torus T into two atoroidal manifolds, then one of M^α or M^β admits a Seifert fibering over the disk with 2 or 3 exceptional fibers or over the Mobius band with up to 2 exceptional fibers such that filling the other along the slope induced by a regular fiber in T produces a lens space. (This is more general than Kobayashi's classification but suitable for our needs.) The slope of a regular fiber may be identified since filling one of these Seifert fibered spaces produces a reducible manifold if and only if the filling is done along the slope of a regular fiber.

Figure 2 (left) shows that the result of a 0-framed banding along κ_n has a sphere S dividing it into two 2-string tangles. Up to homeomorphism, these two tangles are the tangle sums $T_n^\alpha = 1/(n+2) + \frac{1}{3}$ and $T_n^\beta = 1/(1-n) - \frac{1}{2}$. Their corresponding double branched covers, M_n^α and M_n^β , are in general each Seifert fiber spaces over the disk with exactly two exceptional (and nondegenerate) fibers: M_n^α has type $D^2(|n+2|, 3)$ and M_n^β has type $D^2(|1-n|, 2)$. This fails for M_n^α when $n = -1, -2, -3$ and for M_n^β when $n = 0, 1, 2$; in each, the middle value yields a degenerate Seifert fibration while the other values yield solid tori. In these cases the torus T that is the double branched cover of S is compressible and so Kobayashi's classification does not apply; moreover, one may observe that K_n has tunnel number one in these cases. Hence we assume $n \notin \{-3, -2, -1, 0, 1, 2\}$.

When $n = -1$ or 3 , M_n^β has type $D^2(2, 2)$ and thus is a twisted I -bundle over the Klein bottle. Therefore it has an alternative Seifert fibration over the Mobius band with no exceptional fibers. We already assume $n \neq -1$. We shall assume $n \neq 3$ as well.




Figure 3. Left: An isotopy of the arc κ_n^* on the two bridge link $B(3n^2 + n + 1, -3n + 2)$ from the lower left of Figure 1 has two clasp changes performed. The resulting arc τ_n^* is then isotoped to lie in a bridge sphere. Right: A clasp change of an arc around a segment of a branch locus lifts to a crossing change about a segment of the fixed set in the double branched cover.

Figure 2 (middle) shows the results of filling T_n^α and T_n^β with the rational tangle ρ^α defined by a pair of arcs in S so that $T_n^\alpha(\rho^\alpha)$ is composite. We then observe that the filling $T_n^\beta(\rho^\alpha)$ is the pretzel link $P(1 - n, 2, 2)$, and this is a two-bridge link if and only if $n = 0, 2$. We have already omitted these values of n .

Figure 2 (right) shows the results of filling T_n^α and T_n^β with the rational tangle ρ^β defined by a pair of arcs in S so that $T_n^\beta(\rho^\beta)$ is composite. We then observe that the filling $T_n^\alpha(\rho^\beta)$ is the pretzel link $P(n + 2, 3, -2)$, and this is a two-bridge link if and only if $n = -1, -3$. We have already omitted these values of n too.

Therefore, to conclude that the knot K_n has tunnel number 2, it is sufficient to require that $n \notin \{-3, -2, -1, 0, 1, 2, 3\}$. □

Lemma 12. *The knot K_n^* in $L(3n^2 + n + 1, -3n + 2)$ that is surgery dual to K_n is homologous to the knot T_n^* that is surgery dual to the $(3n + 1, n)$ -torus knot in S^3 .*

Proof. Starting from the lower left of Figure 1 where the dual arc κ_n^* is presented on a standard form of the two bridge link $B(3n^2 + n + 1, -3n + 2)$, Figure 3 (left) shows how two clasp changes (and isotopy) transforms κ_n^* into an arc τ_n^* in a bridge sphere. Since τ_n^* lies in the bridge sphere, its lift to the double branched cover is a knot T_n^* in the Heegaard torus of the lens space. Lifting the transformation of Figure 3 (left) to the double branched cover thus shows that K_n^* and T_n^* are related by two crossing changes. (The clasp changes lift to crossing changes as indicated in Figure 3 (right).) Hence these knots are homotopic, and therefore homologous, in the lens space.




Figure 4. Left: A banding of $B(3n^2 + n + 1, -3n + 2)$ along the arc τ_n^* produces the unknot. Right: A different banding of $B(3n^2 + n + 1, -3n + 2)$ along the arc τ_n^* produces the connected sum of $B(n, -1)$ and $B(3n + 1, 3)$.

Finally, [Figure 3](#) (left) shows a banding of $B(3n^2 + n + 1, -3n + 2)$ along the arc τ_n^* to the unknot. Thus we may identify T_n^* as surgery dual to a torus knot T_n in S^3 . Because [Figure 3](#) (left) shows a banding of $B(3n^2 + n + 1, -3n + 2)$ along the arc τ_n^* to the connected sum $B(n, -1)\#B(3n + 1, 3)$, the torus knot T_n has a reducible surgery to $L(n, -1)\#L(3n + 1, 3) \cong L(-n, 3n + 1)\#L(3n + 1, -n)$ which is the mirror of $L(n, 3n + 1)\#L(3n + 1, n)$. Thus T_n must be the $(3n + 1, n)$ -torus knot (for example, see [\[Moser 1971\]](#)). \square

Proof of Theorem 7. (1) [Theorem 1](#) shows that either $\pm p$ -surgery on K_n is a lens space. Tange shows that any integral lens space surgery on a knot in \mathcal{P} must be positive [\[Tange 2011, Theorem 1.1\]](#).

(2) In [\[Tange 2011, Theorem 3.1\]](#), the paragraph following, and its proof, Tange further shows that if an L-space homology sphere Y contains a knot K with irreducible exterior for which a positive integer surgery produces an L-space, then K is a fibered knot supporting a tight contact structure. (Tange notes that the fiberedness of K follows from the proof of [\[Ozsváth and Szabó 2005, Theorem 1.2\]](#) and [\[Ni 2007\]](#). Tange then demonstrates that the Heegaard Floer contact invariant of the contact structure supported by K is nonzero.)

(5) If T_n is the $(3n + 1, n)$ -torus knot in S^3 , then p -surgery on T_n produces the lens space $L(p, q)$. By [Lemma 12](#), the surgery dual knot T_n^* is homologous to the knot K_n^* that is surgery dual to K_n .

(3) and (6) Since the dual knots K_n^* and T_n^* are homologous, [Theorem 8](#) implies $2g(K^n) = p - 1$ and the stated relationship of the Alexander polynomials of K_n and T_n .




Figure 5. Left: A regular neighborhood of an oriented knot K in a homology sphere Y has meridian μ linking K positively and longitude λ oriented as the boundary of a Seifert surface. The surgery curve μ^* of positive slope $5/1$ shown here is oriented as $[\mu^*] = [\lambda] + 5[\mu]$ in $H_1(\partial\mathcal{N}(K))$. Right: Performing μ^* -surgery on K produces the rational homology sphere Z and surgery dual knot K^* . Orienting K^* to be linked positively by its meridian μ^* , the boundary of the (rational) Seifert surface λ is homologous to $5[K^*]$ in $\mathcal{N}(K^*)$ while the longitude μ is homologous to $-[K^*]$.

(4) Knowing that $g(K_n) = (p+1)/2$ implies $\text{rk } \widehat{\text{HFK}}(L(p, q), K_n^*) = p+2$ [Hedden 2011, Theorem 1.4] (see also [Rasmussen 2007, Proposition 4.5]). \square

Proof of Theorem 8. Let K^* be the surgery dual to K in $L(p, q)$ and (for some choice of orientations) let J^* be the simple knot in $L(p, q)$ homologous to K^* . Then J^* has an integral surgery to an L-space homology sphere Y_J [Rasmussen 2007, Theorem 2].

Since *positive* p -surgery on K gives $L(p, q)$, then the self-linking number of K^* is $-1/p \pmod{1}$; see [Rasmussen 2007, §2]. This is demonstrated in Figure 5: If Σ is an oriented Seifert surface for K giving the oriented longitude $\lambda = \partial\Sigma$, then the positively linking meridian μ of K is oriented as shown. In order for λ to run positively along K^* , we need $\mu^* \cdot \lambda > 0$. Thus we must orient μ^* so that $[\mu^*] = [\lambda] + p[\mu]$. This forces μ to be an anti-parallel longitude of K^* . Hence we may use $-\mu$ to calculate the self-linking number of K^* as $(-\mu \cdot \lambda)/p \pmod{1}$.

With this set-up, μ -surgery on K^* may be regarded as -1 -surgery on K^* . This means $Y = K_{-1}^*$ in the notation of [Rasmussen 2007], though not explicitly stated. Similarly we also have $Y_J = J_{-1}^*$.

If $2g(K) - 1 \neq p$, then the knots J^* and K^* have isomorphic knot Floer homology [Rasmussen 2007, Theorem 2] (see also [Hedden 2011, Theorem 1.4]). Having isomorphic (and simple) knot Floer homology would imply that the L-space homology spheres Y_J and Y have the same d -invariants. Hence $d(Y_J) = 2$ too. But now Y_J cannot be S^3 since $d(S^3) = 0$.

If $2g(K) - 1 = p$ then width $\widehat{\text{HF}}(L(p, q), K^*) = 2p$ [Ni 2009]; see [Rasmussen 2007, Theorem 4.3]. Thus $d(Y) = d(Y_J) + 2$ by [Rasmussen 2007, Proposition 5.4] and so $d(Y_J) = 0$. Greene's solution [2013] to the lens space realization problem proceeds by first identifying the pairs (lens space, homology class) that contain knots for which some integral surgery produces an L-space homology sphere with $d = 0$, and then observing that each of these pairs contains the surgery dual to a Berge knot. Since the lens space surgery duals to Berge knots are simple knots, we have that in fact J^* is surgery dual to a Berge knot B and $Y_J = S^3$.

For the statement about Alexander polynomials, let us first summarize work of Tange [2009b] on Alexander polynomials of knots in L-space homology spheres for which a positive integral surgery yields a lens space. The symmetrized Alexander polynomial of a knot K in an L-space homology sphere with a nontrivial L-space surgery can be expressed as

$$\Delta_K(t) = \sum_{i \in \mathbb{Z}} a_i(K) t^i = (-1)^k + \sum_{j=1}^k (-1)^{k-j} (t^{n_j} + t^{-n_j}) \in \mathbb{Z}[t^{\pm 1}]$$

following the arguments of [Ozsváth and Szabó 2005]. For a given positive integer p , pass to the quotient $\mathbb{Z}[t^{\pm 1}]/(t^p - 1)$ to obtain a polynomial $\tilde{\Delta}_K(t) = \sum_{i \in \mathbb{Z}/p\mathbb{Z}} \tilde{a}_i(K) t^i$ with coefficients $\tilde{a}_i(K) = \sum_{j=i \bmod p} a_j(K)$. Assuming that p -surgery gives a lens space Z , then $2g(K) - 1 \leq p$ [Kronheimer et al. 2007]. Note that K is fibered⁵ and hence the degree of $\Delta_K(t)$ equals $2g(K)$. Because of this and that the coefficients of $\Delta_K(t)$ are $a_i(K) = 0$ or ± 1 (and the nonzero ones alternate in sign), it follows that $\tilde{a}_i(K) = 0, \pm 1$, or 2 where $\tilde{a}_i(K) = 2$ implies that both p is even and $i = \frac{1}{2}p$ [Tange 2009b, Corollary 2]. Furthermore, one may determine that the coefficients $\tilde{a}_i(K)$ and the polynomial $\tilde{\Delta}_K(t)$ only depend on the lens space Z and the homology class of the surgery dual knot K^* [Brody 1960] (see [Rasmussen 2007, §5]). In particular, if $p = 2g(K) - 1$ then

$$\Delta_K(t) = \tilde{\Delta}_K(t) - (t^{(p-1)/2} + t^{-(p-1)/2}) + (t^{(p+1)/2} + t^{-(p+1)/2})$$

where the indices for the coefficients of $\tilde{\Delta}_K(t)$ are taken to be the representatives of $\mathbb{Z}/p\mathbb{Z}$ from $-\frac{1}{2}(p-1)$ and $\frac{1}{2}(p-1)$. See also the discussion preceding [Tange 2009a, Proposition 3.3].

⁵Since lens spaces are irreducible, a knot in a homology sphere with a lens space surgery must have irreducible exterior. Thus K has irreducible exterior and so it is fibered by [Ozsváth and Szabó 2005, Theorem 1.2] and [Ni 2007]; see Theorem 7(2).

Now in our present situation, since the Berge knot B in S^3 has p -surgery to $L(p, q)$ in which the surgery dual is the simple knot J^* , we therefore have $2g(B) < p$ because p is odd [Hedden 2011, Theorem 3.1]⁶. Hence $\tilde{a}_i(B) = a_i(B)$ for all $i \in \mathbb{Z}/p\mathbb{Z}$ and $\Delta_B(t) = \tilde{\Delta}_B(t)$. Because the surgery dual K^* is homologous to the simple knot B^* in $L(p, q)$, $\tilde{\Delta}_K(t) = \tilde{\Delta}_B(t)$. Therefore we have

$$\Delta_K(t) = \Delta_B(t) - (t^{(p-1)/2} + t^{-(p-1)/2}) + (t^{(p+1)/2} + t^{-(p+1)/2})$$

as claimed. □

Lemma 13. *There is a unique genus one fibered knot J in \mathcal{P} . As an open book, it supports the tight contact structure on \mathcal{P} . Furthermore, it is surgery dual to the negative trefoil knot in S^3 .*

Proof. This can be seen in the spirit of [Baker 2014a] which relates genus one fibered knots to axes of closed 3-braids as follows: Noting that since \mathcal{P} has a unique genus 2 Heegaard splitting [Boileau and Otal 1991], $P(-2, 3, 5)$ is the only 3-bridge link whose double branched cover gives \mathcal{P} . (In fact \mathcal{P} is the double branched cover of no other link.) Since $P(-2, 3, 5)$ is isotopic to the $(3, 5)$ -torus knot, its 3-braid axis A lifts to a genus one fibered knot $J \subset \mathcal{P}$. By [Birman and Menasco 1993, classification theorem] and [Baker 2014a, Lemma 3.8], for instance, A is the only 3-braid axis for $P(-2, 3, 5)$ up to isotopy of unoriented links. Hence J is the only genus one fibered knot up to homeomorphism in \mathcal{P} . Since the mapping class group of \mathcal{P} is trivial [Boileau and Otal 1991], J is the only genus one fibered knot in \mathcal{P} up to isotopy.

In the double branched cover, a braid presentation of the branch locus lifts to a Dehn twist presentation of the monodromy of the lift of the axis. Since the axis A presents $P(-2, 3, 5)$ as a positive braid (indeed, as the $(3, 5)$ -torus knot), we obtain a presentation of the monodromy of J as a product of positive Dehn twists. Hence it supports the tight contact structure on \mathcal{P} [Giroux 2002].

One may view J as surgery dual to the negative trefoil in several ways. Taking the route through branched covers, this is demonstrated in [Baker 2014b, Proof 2], though for the mirrored situation. □

5. Hedden’s almost-simple knots.

In each lens space $L(p, q)$ with coprime $p > q > 0$, Hedden diagrammatically describes two unoriented 1-bridge knots T_L and T_R (*the Hedden knots*) via the doubly pointed genus 1 Heegaard diagrams of $L(p, q)$ with $p + 2$ intersection points [Hedden 2011, Figure 3]. An alternative presentation of local pictures of these diagrams near the two marked points are shown in the top row of Figure 6.

⁶See also [Greene 2013, Theorem 1.4].




Figure 6. Local portions of the doubly pointed diagrams for the almost-simple knots T_L and T_R and their associated simple knots $K(p, q, q+1)$ and $K(p, q, q-1)$; see [Baker 2014b, Figure 1; Hedden 2011, Figure 3]. In the bottom two pictures, the intersection points of the red α -curve and blue β -curve are numbered in order mod p along the α -curve.

Since the *simple knots* in $L(p, q)$ are those that can be described via doubly pointed genus 1 Heegaard diagrams of $L(p, q)$ with only p intersection points, we like to regard the Hedden knots as *almost-simple*. The bottom row of Figure 6 shows the same portion of the diagram for related simple knots. These knots are notable because of the following theorem.

Theorem 14 [Hedden 2011, Theorem 3.1]. *If K^* is a 1-bridge knot in $L(p, q)$ with an integral surgery to an L-space homology sphere, then either*

- $p > 2g - 1$ and K^* is a simple knot, or
- $p = 2g - 1$ and K^* is either T_L or T_R ,

where g is the Seifert genus of the surgery dual knot.

Though the statement of this theorem in [Hedden 2011] is explicitly only for surgeries to S^3 , Hedden notes that his proof applies for surgeries to any L-space homology sphere.

Here we would like to clarify a few points about these almost-simple knots, correct a couple of misstatements in [Baker 2014b], and discuss the homology spheres that may be obtained by integral surgery on them following the arguments of [Baker 2014b]. Afterwards, we summarize these results in Proposition 15.

The knots T_L and T_R are not homologous in general, contrary to what was stated in [Baker 2014b]. One can observe this directly from the descriptions in Figure 6 (or [Hedden 2011, Figure 3]), noting that the magnitude of their algebraic intersection numbers with the blue β -curve differ by 2. One may further observe that the mirror of the knot T_L in $L(p, q)$ is the knot T_R in $L(p, -q)$.

Furthermore, the proof in [Baker 2014b] that T_L differs from (and thus is homologous to) $K(p, q, q + 1)$ by a crossing change also shows that T_R differs from $K(p, q, q - 1)$ by a crossing change. Unfortunately, [Baker 2014b, Figure 2] should be mirrored through the Heegaard torus so that the red α -disks are below, the blue β -disks are above, and the twist in T_L is right-handed in order to be consistent with the orientation conventions. In particular, the crossing change from $K(p, q, q + 1)$ to T_L is achieved by a -1 -surgery on a loop C_L in the Heegaard that encircles the two base-points in the diagram for $K(p, q, q + 1)$ of Figure 6 (bottom left). The crossing change from $K(p, q, q - 1)$ to T_R may be similarly achieved by a $+1$ -surgery on a loop C_R .

A knot homologous to $K(p, q, k)$ has an integral surgery to a homology sphere if and only if its self linking is $\pm 1/p \pmod{1}$, or equivalently if and only if $k^2 = \pm q \pmod{p}$ [Rasmussen 2007, Lemma 2.6]; see [Fintushel and Stern 1980]. Here the choice of sign of \pm is consistent and agrees with whether this integral surgery is a ± 1 -surgery; see Figure 5 and the discussion in the proof of Theorem 8.

If T_L in $L(p, q)$ is dual to positive p -surgery on a knot K in a homology sphere Y , then so is $K(p, q, q + 1)$ and hence $(q + 1)^2 = -q \pmod{p}$. Making the substitution $-k = q + 1$ gives the equation $k^2 + k + 1 = 0 \pmod{p}$. Thus $K(p, q, q + 1)$ is dual to a type VII Berge knot B , one that lies in the fiber of a trefoil knot; see [Rasmussen 2007, §6.2; Greene 2013, §1.2]. Since its fiber positively frames B , this trefoil knot is the *negative* trefoil (not the positive trefoil as stated in [Baker 2014b]), and it can be identified with C_L under the S^3 surgery on $K(p, q, q + 1)$. Performing -1 -surgery on the negative trefoil C_L in S^3 produces \mathcal{P} and takes $B \subset S^3$ to $K \subset \mathcal{P}$. Since the linking of B and C_L is 0, the positive p -surgery on B becomes the positive p -surgery on K . Consequently, if -1 -surgery on T_L is a homology sphere, it is \mathcal{P} .

Similarly, if T_R in $L(p, q)$ is dual to positive p -surgery on a knot K' in a homology sphere, then $(q - 1)^2 = -q \pmod{p}$. Making the substitution $k = q - 1$ gives the equation $k^2 - k - 1 = 0 \pmod{p}$. Thus $K(p, q, q - 1)$ is dual to a type VIII Berge knot B' , one that lies in the fiber of the figure eight knot. This figure eight knot may be identified with C_R . Performing $+1$ -surgery on C_R in S^3 produces

the Brieskorn sphere $\Sigma(2, 3, 7)$ taking the knot B' to the knot K' . Consequently, if -1 -surgery on T_R is a homology sphere, it is $\Sigma(2, 3, 7)$.

This coincides with the difference between the τ -invariants of these knots as noted by Rasmussen in the last two paragraphs of [Rasmussen 2007, §5]: $\tau(T_L, \mathfrak{s}_0) = -1$ and so $\tau(T_R, \mathfrak{s}_0) = +1$. Rasmussen further shows that if integral surgery on T_L or T_R produces a homology sphere, then it is an L-space homology sphere if and only if the surgery is a -1 -surgery on T_L or a $+1$ -surgery on T_R [Rasmussen 2007, Proposition 4.5].

In summary, the above discussion shows:

- Proposition 15.** (1) *In $L(p, q)$, T_L is homologous to the simple knot $K(p, q, q + 1)$ and T_R is homologous to the simple knot $K(p, q, q - 1)$.*
- (2) *The mirror of $(L(p, q), T_L)$ is $(L(p, -q), T_R)$.*
- (3) *If -1 -surgery on T_L is a homology sphere, then it is $\mathcal{P} = \Sigma(2, 3, 5)$ and $K(p, q, q + 1)$ is positive surgery dual to a type VII Berge knot.*
- (4) *If -1 -surgery on T_R is a homology sphere, then it is $\Sigma(2, 3, 7)$ and $K(p, q, q - 1)$ is positive surgery dual to a type VIII Berge knot.*

Appendix: Symmetries of knots with lens space and Poincaré homology sphere surgeries

Kenneth L. Baker and Neil R. Hoffman

The aim of this appendix is to compute the symmetry group of a hyperbolic knot in a lens space with a surgery to the Poincaré homology sphere \mathcal{P} . Specifically, it will provide a proof of the following theorem:

Theorem A.1. *Let X be a one-cusped hyperbolic manifold admitting both a Poincaré homology sphere filling and a lens space filling. Then the symmetry group of X is trivial or generated by a single strong involution.*

Our method blends the classification of spherical orbifolds with results involving exceptional Dehn fillings. We refer the reader to [Boileau et al. 2012] for the most closely related paper on this topic and Thurston’s notes [1979, Chapter 13] for a more general introduction to orbifolds. In preparation, we first extend the notation of Section 3 to address orbifolds and orbifold fillings and then discuss symmetries and quotients of hyperbolic manifolds. Thereafter, the argument in this appendix will gradually “whittle-down” the symmetry group of the exterior X of a hyperbolic knot K in a lens space with a Dehn surgery to \mathcal{P} . First, we give an argument which eliminates orientation reversing symmetries so that $\text{Sym}(X) = \text{Sym}^+(X)$. Then we proceed to analyze a subgroup $Z(X)$ of $\text{Sym}^+(X)$, showing

- (a) that it has index at most 2,

- (b) if the index is 2 then $\text{Sym}(X)$ contains a strong involution, and ultimately
- (c) that $Z(X)$ is trivial.

This is pulled together in the proof of [Theorem A.1](#) at the end of this appendix.

Background on orbifolds and symmetries. Let Q be a 3-dimensional manifold or orbifold with *singular set* $\Sigma(Q)$, the set of points fixed by some nontrivial element of $\pi_1^{\text{orb}}(Q)$. (So Q is a manifold if $\Sigma(Q) = \emptyset$.) If K is a knot in Q , then its complement is $Q - K$ while its exterior is $X = Q - \mathcal{N}(K)$. We further assume that K is either a component of $\Sigma(Q)$ or disjoint from $\Sigma(Q)$.

The torus $\partial\mathcal{N}(K)$ represents the *cusps* of $Q - K$ corresponding to K . In such a torus we consider essential closed curves up to free homotopy. A *primitive curve* is homotopic to an essential simple closed curve, and hence is a slope. Any nonprimitive essential curve is a multiple of a primitive curve. Given two closed curves α and β in the torus we say the *distance* between α and β , $\Delta(\alpha, \beta)$ is the minimal (unoriented) geometric intersection number between the two curves. If α is a curve in $\partial\mathcal{N}(K)$ that is an n -fold multiple of a slope γ , then the result of α -Dehn surgery on $K \subset Q$ is the orbifold $X(\alpha)$ with underlying manifold $|X(\alpha)|$ obtained as the γ -Dehn surgery on $K \subset |Q|$ and a new singular set $\Sigma(X(\alpha))$ consisting of $\Sigma(X)$ and the core of the attached $S^1 \times D^2$ with order n (unless $n = 1$ in which case the singular set remains only $\Sigma(X)$). The orbifold attached to X is a *orbi-solid torus* of order n . An orbi-solid torus of order 1 is just a solid torus.

Following [[Boileau et al. 2012](#), §3], an *orbi-lens space* is the quotient of S^3 by a finite cyclic subgroup of $SO(4)$ and is consequentially a union of two orbi-solid tori with coprime orders along their common boundary. Its singular set is the union of the cores of these orbi-solid tori that have order at least 2.

A knot K in a closed 3-manifold M is *hyperbolic* if its complement $M - K$ is a hyperbolic 3-manifold with a single torus cusp. We also say the exterior $X = M - \mathcal{N}(K)$ of such a knot is *hyperbolic* if the interior of X is hyperbolic. For a hyperbolic knot K with exterior X , the symmetry group $\text{Sym}(X)$ of homeomorphisms modulo isotopy is identified with the isometry group of the interior of X . The orientation preserving symmetry group is denoted $\text{Sym}^+(X)$.

Within $\text{Sym}^+(X)$ is the subgroup $Z(X)$, the maximal subgroup that acts freely on ∂X . In particular, $Z(X)$ is the maximal subgroup of $\text{Sym}^+(X)$ for which the quotient orbifold (or manifold) $X/Z(X)$ has a torus cusp.

Since X is hyperbolic, the quotient $X/\text{Sym}^+(X)$ is an orientable cusped hyperbolic 3-orbifold with $\text{vol}(X/\text{Sym}^+(X)) = \text{vol}(X)/|\text{Sym}^+(X)|$. Since the minimal volume of an orientable cusped hyperbolic 3-orbifold is positive (e.g., [[Meyerhoff 1986](#)]) and X has finite volume, it follows that $\text{Sym}^+(X)$ is a finite group.

In a knot exterior X , the *homological longitude* is the slope λ in ∂X so that $[\lambda] = 0$ in $H_1(X; \mathbb{Q})$. This slope is unique up to sign as it is a generator of the

kernel of the map $i_*: H_1(\partial X; \mathbb{Q}) \rightarrow H_1(X; \mathbb{Q})$ which has rank 1 by the half-lives, half-dies lemma. Since it is unique, elements of $\text{Sym}(X)$ must preserve the slope λ up to sign. Elements of $\text{Sym}^+(X)$ of order 2 that reverse the orientation of the homological longitude λ of the manifold X are called *strong involutions*.

If a group G acts on a hyperbolic knot exterior X , the cusp of the quotient orbifold X/G records important information about the action of G on the cusp of X . If the quotient X/G has a torus cusp, then G acts freely on ∂X . Furthermore, each element of G restricts to a translation on ∂X . If X/G has an $S^2(2, 2, 2, 2)$ cusp, then G contains an order 2 element which fixes a discrete set of points on ∂X . In particular, if $X/\text{Sym}^+(X)$ has an $S^2(2, 2, 2, 2)$ boundary, then such an element of $\text{Sym}^+(X)$ must preserve isotopy classes of slopes in ∂X but reverse their orientations. Hence $\text{Sym}^+(X)$ contains a strong involution. Furthermore, this means that there is a degree 2 cover with a torus cusp, and hence $Z(X)$ has index 2 in $\text{Sym}^+(X)$.

Whittling down the symmetry group.

Proposition A.2. *Let X be the exterior of a knot in a lens space $L(p, q)$ with $p \geq 2$ admitting a Dehn surgery to an integral homology sphere. Then either X is Seifert fibered or $\text{Sym}(X) = \text{Sym}^+(X)$.*

Proof. Assume X admits an orientation reversing symmetry τ , and let λ be the homological longitude of X . Then $\tau(\lambda)$ is isotopic to $\pm\lambda$ in ∂X . Since τ restricts to a homeomorphism on the boundary torus ∂X , there must be a slope δ with $\Delta(\delta, \lambda) = 1$ such that $\tau(\delta) = \pm\delta$. Because this restriction of τ to ∂X must also be orientation reversing, τ reverses the orientation on exactly one of δ and λ .

Let μ be the slope in ∂X such that $X(\mu) = L(p, q)$. Since X has a filling producing an integer homology sphere, X must be a homology solid torus. Thus, $\mu = p\delta + q'\lambda$ in $H_1(\partial X)$ for some integer q' coprime to p and $p \geq 2$. Since $\tau(\mu) = \pm(p\delta - q'\lambda)$ and $q' \neq 0$, we have that $\Delta(\mu, \tau(\mu)) = |2pq'| > 2$.

The symmetry τ implies $X(\mu) \cong X(\tau(\mu))$ so that $\pi_1(X(\mu))$ and $\pi_1(X(\tau(\mu)))$ are both cyclic. Thus, since $\Delta(\mu, \tau(\mu)) > 1$, X must be Seifert fibered by the cyclic surgery theorem [Culler et al. 1987]. \square

We now further classify $Z(X)$.

Lemma A.3. *Let X be the exterior of a hyperbolic knot K in a lens space $L(p, q)$ with $p \geq 2$ admitting a Dehn surgery to an integral homology sphere. Then the following hold:*

- (1) $Z(X)$ is cyclic.
- (2) $[\text{Sym}^+(X) : Z(X)] \leq 2$.
- (3) If $[\text{Sym}^+(X) : Z(X)] = 2$, then X has a symmetry which is a strong involution.

- (4) In the quotient orbifold $\mathcal{Z} = X/Z(X)$, the meridian μ of K maps to a primitive curve $\bar{\mu}$ in $\partial\mathcal{Z}$.
- (5) The Dehn filling $\mathcal{Z}(\bar{\mu})$ is an orbi-lens space in which the core \bar{K} of the filling is disjoint from the singular set of $\mathcal{Z}(\bar{\mu})$.

Proof. We first utilize the fact that X is the exterior of a hyperbolic knot in an integral homology sphere to show that the boundary of $X/\text{Sym}^+(X)$ is either a torus or $S^2(2, 2, 2, 2)$.

Since X admits an integer homology sphere filling, X is a homology solid torus. Furthermore, once oriented, the slope γ in ∂X of this filling generates $H_1(X)$. Thus together with a choice of orientation on the homological longitude λ it forms a basis (γ, λ) for $H_1(\partial X)$.

Now consider the action of $\text{Sym}^+(X)$ on ∂X . Since the homological longitude is the unique slope in ∂X that is null homologous in $H_1(X; \mathbb{Q})$, any element of $\text{Sym}^+(X)$ must send λ to $\pm\lambda$. Since $\text{Sym}^+(X)$ is finite because X is hyperbolic, consideration of the action on $H_1(\partial X; \mathbb{Q})$ shows that any element of $\text{Sym}^+(X)$ must send γ to $\pm\gamma$. Then, because $\text{Sym}^+(X)$ is orientation preserving on ∂X too, any element of $\text{Sym}^+(X)$ must send the basis (γ, λ) to $\pm(\gamma, \lambda)$. Hence all slopes in ∂X are preserved by the action of $\text{Sym}^+(X)$. Consequently, the restriction to ∂X of a symmetry of X either is trivial, has no fixed point (and hence is a translation), or has order 2. Therefore, by consideration of the Euclidean orbifolds which are orientable quotients of the torus, the boundary of $X/\text{Sym}^+(X)$ is either a torus or $S^2(2, 2, 2, 2)$.

We now further refine this analysis by incorporating the lens space filling. Since its exterior X is a homology solid torus, the knot K in $L(p, q)$ represents a generator of $H_1(L(p, q), \mathbb{Z})$. Hence it lifts to a knot $\tilde{K} \subset S^3$ in the p -fold cyclic cover $S^3 \rightarrow L(p, q)$. This cover restricts to a p -fold cyclic cover of knot exteriors $\tilde{X} \rightarrow X$ in which the meridian $\mu \subset \partial X$ of K lifts to p disjoint copies of the meridian of \tilde{K} . (Here \tilde{X} is the exterior of \tilde{K} .) The deck transformation group H of this cover is therefore isomorphic to the cyclic group $\mathbb{Z}/p\mathbb{Z}$ and extends to an action on S^3 that is free on $\partial\tilde{X}$. Hence H is a subgroup of $Z(\tilde{X})$.

Consider the composition $\alpha: \tilde{X} \rightarrow X/\text{Sym}^+(X)$ of the covers $\tilde{X} \rightarrow X$ and $X \rightarrow X/\text{Sym}^+(X)$ with deck transformation groups H and $\text{Sym}^+(X)$ respectively. Let G be the deck transformation group of α . If α is regular, then G is a subgroup of $\text{Sym}^+(\tilde{X})$ and H must be a normal subgroup of G with $\text{Sym}^+(X) = G/H$. As the boundary of $X/\text{Sym}^+(X)$ is either a torus or $S^2(2, 2, 2, 2)$, we claim that α is indeed a regular cover. If this boundary is a torus, then this follows directly from [Reid 1991, Lemma 4]. However if this boundary is $S^2(2, 2, 2, 2)$, then the argument of [Reid 1991, Lemma 4] extends to also show that the covering $\alpha: \tilde{X} \rightarrow X/\text{Sym}^+(X)$ is a regular cover as noted in the proof of [Neumann and

Reid 1992, Proposition 9.1]. (Ultimately, this is because conjugation by an order 2 element of the peripheral subgroup $\pi_1(S^2(2, 2, 2, 2))$ inverts elements of the maximal abelian subgroup. This maximal abelian subgroup contains the image of the peripheral subgroup of \tilde{X} .)

Because \tilde{X} is the exterior of a hyperbolic knot in S^3 , we know that $\text{Sym}^+(\tilde{X})$ must be cyclic or dihedral [Boileau et al. 2012, p. 627] (see also [Neumann and Reid 1992, Proposition 9.1]). More specifically $Z(\tilde{X})$ is cyclic, $[\text{Sym}^+(\tilde{X}) : Z(\tilde{X})] \leq 2$, and $[\text{Sym}^+(\tilde{X}) : Z(\tilde{X})] = 2$ only if $\text{Sym}^+(\tilde{X})$ contains a strong involution.

Since G is a subgroup of $\text{Sym}^+(\tilde{X})$, either it is a subgroup of $Z(\tilde{X})$ and hence cyclic and acting freely on $\partial\tilde{X}$ itself, or it contains a strong involution. In the former case, $\text{Sym}^+(X) = G/H$ is cyclic as it is the quotient of a cyclic group G and $X/\text{Sym}^+(X)$ has torus boundary so that $\text{Sym}^+(X) = Z(X)$. In the latter case, $\text{Sym}^+(X) = G/H$ must also contain a strong involution since H is a subgroup of $Z(\tilde{X})$. Furthermore, because $H \cong \mathbb{Z}/p\mathbb{Z}$ is a nontrivial subgroup of $Z(\tilde{X})$ (since $p \geq 2$), G cannot be generated by a single strong involution. Thus G must be a dihedral group (of order ≥ 4), and so $\text{Sym}^+(X) = G/H$ is a dihedral group too (though possibly of order 2). Since $X/\text{Sym}^+(X)$ has $S^2(2, 2, 2, 2)$ boundary, $[\text{Sym}^+(X) : Z(X)] = 2$ as noted in the introduction to this appendix. Since $Z(X)$ cannot contain a strong involution, is a cyclic group. Consequently, in both cases claims (1), (2), and (3) of the lemma hold.

Finally we examine the quotient orbifold $\mathcal{Z} = X/Z(X)$ and its filling $\mathcal{Z}(\bar{\mu})$. Since \tilde{X} is the exterior of the knot $\tilde{K} \subset S^3$, $\text{Sym}^+(\tilde{X})$ naturally identifies with $\text{Sym}^+(S^3, \tilde{K})$. In particular, $Z(\tilde{X})$ acts freely on \tilde{K} . Therefore, if $\tilde{\mu}$ is the lift of the slope μ to \tilde{X} and $\bar{\mu}$ is its image in the quotient orbifold $\mathcal{Z} = \tilde{X}/Z(\tilde{X}) = X/Z(X)$, then $\bar{\mu}$ must be a primitive slope and (4) holds.

Filling \mathcal{Z} along the slope $\bar{\mu}$ produces

$$\mathcal{Z}(\bar{\mu}) = X(\mu)/Z(X),$$

which we recognize as an orbi-lens space since it is a cyclic quotient of $S^3 = \tilde{X}(\tilde{\mu})$ (see [Boileau et al. 2012, §3] for context). Because $\bar{\mu}$ is a primitive slope, the core of the solid torus in the filling $\mathcal{Z}(\bar{\mu})$ is a knot \bar{K} that is disjoint from the singular set $\Sigma(\mathcal{Z}(\bar{\mu}))$ of the orbi-lens space. In particular, \mathcal{Z} is the exterior of a knot in an orbi-lens space that is disjoint from the singular set. Thus (5) holds, which completes the proof. \square

In light of the above lemma which employs a classification of cyclic quotients of S^3 , it will also prove useful to have a list of manifolds and orbifolds that are cyclic quotients of \mathcal{P} . This is accomplished by Lemma A.4 below which essentially follows from combining the classification of elliptic manifolds with the corresponding classification of elliptic orbifolds in Dunbar [1988].




Figure 7. The possible singular sets of orbifolds cyclically covered by \mathcal{P} with underlying space S^3 . These are the relevant cases from a more general classification in [Dunbar 1988]. Following the notation of that paper, a link component labeled by n indicates a cone angle of $2\pi/n$ along that component in the singular set of the corresponding orbifold.

Lemma A.4. *If the finite cyclic quotient of \mathcal{P} by symmetries is an orbifold Q , then Q is homeomorphic to one of the following:*

- (1) *a manifold M with $\pi_1(M) \cong \pi_1(\mathcal{P}) \times \mathbb{Z}/n\mathbb{Z}$ for some integer n coprime to 30,*
- (2) *an orbifold that fibers over $S^2(2, 3, 5)$ where the singular set of Q has 1, 2, or 3 components, or*
- (3) *an orbifold with base space S^3 and singular set a knot or link as pictured in Figure 7.*

Moreover, in the case that Q is an orbifold, the components of its singular set are fixed locally by finite cyclic groups of relatively prime orders.

Proof. By the orbifold theorem (see [Boileau et al. 2003; Cooper et al. 2000, Chapter 7]), we know that Q is a spherical 3-orbifold.

Case 1 follows directly from the classification of elliptic manifolds (see for example [Thurston 1997, Theorem 4.4.14]), while Cases 2 and 3 follow from a careful reading of [Dunbar 1988]. Dunbar [1988] also classifies orbifolds covered by elliptic manifolds where the singular sets are trivalent graphs. However, no orbifold of this type can be cyclically covered by a manifold because the group of isometries that fixes a vertex in the trivalent graph is never cyclic. \square

A simple case analysis shows that drilling the singular sets of orbifolds from the previous lemma results in several interesting manifolds.

Lemma A.5. *In Cases 2 and 3 of Lemma A.4:*

- If $|\Sigma(Q)| = 1$, then the complement of $\Sigma(Q)$ is a Seifert fibered space over the disk with two exceptional fibers.
- If $|\Sigma(Q)| = 2$, then the complement of $\Sigma(Q)$ is a Seifert fibered space over the annulus with one exceptional fiber.
- If $|\Sigma(Q)| = 3$, then the complement of $\Sigma(Q)$ is $F \times S^1$, where F is a pair of pants.

Lemma A.6. *Let X be the exterior of a hyperbolic knot K in a lens space $L(p, q)$ with $p \geq 2$ admitting a Dehn surgery to \mathcal{P} . Then every nontrivial subgroup of $Z(X)$ acts nonfreely on X .*

Proof. Assume $Z_f \subset Z(X)$ is a nontrivial subgroup of elements that act freely on X . Then the quotient X/Z_f is a hyperbolic manifold with a torus boundary. Since Z_f is nontrivial, $|Z_f| \geq 2$. Since $Z(X)$ is cyclic by Lemma A.3, the subgroup Z_f must be cyclic.

Assume μ is the meridian of the knot K in the lens space, and γ is the slope of is \mathcal{P} surgery. Let $\bar{\mu}$ and $\bar{\gamma}$ be the images of μ and γ in the quotient X/Z_f . Then the orbifold fillings of X/Z_f are quotients of the Dehn fillings of X . In particular, $(X/Z_f)(\bar{\mu}) = X(\mu)/Z_f$ and $(X/Z_f)(\bar{\gamma}) = X(\gamma)/Z_f$ where the singular sets, if nonempty, are connected, consisting solely of the cores of the filling orbifold tori. Note that these quotients are manifolds exactly when their curves $\bar{\mu}$ and $\bar{\gamma}$ are primitive.

First we observe Lemma A.3 implies that $\bar{\mu}$ must be a primitive curve because X/Z_f is a quotient of X that covers the orbifold $\mathcal{Z} = X/Z(X)$. Thus we have two cases depending on whether $\bar{\gamma}$ is a primitive curve.

Case 1: $\bar{\gamma}$ are primitive. Observe that $\Delta(\bar{\mu}, \bar{\gamma}) = |Z_f| \Delta(\mu, \gamma)$, where $\Delta(\bar{\mu}, \bar{\gamma})$ is the distance in the boundary of the quotient and $\Delta(\mu, \gamma)$ is measured in ∂X .

Since μ and γ have primitive images in the quotient manifold X/Z_f , Z_f acts freely on the fillings $X(\mu)$ and $X(\gamma)$. Hence $(X/Z_f)(\bar{\mu}) = X(\mu)/Z_f$ is a manifold quotient of the lens space $X(\mu)$ and $(X/Z_f)(\bar{\gamma}) = X(\gamma)/Z_f$ is a manifold quotient of $\mathcal{P} = X(\gamma)$. Thus they are both covered by S^3 , and therefore they are both finite manifolds (i.e., have finite π_1). Since $X(\gamma)/Z_f$ is covered by \mathcal{P} , $|Z_f|$ must be relatively prime to $|\pi_1(\mathcal{P})| = 120$ by Lemma A.4(1). Thus we must have $|Z_f| > 6$. Therefore $\Delta(\bar{\mu}, \bar{\gamma}) > 6$ which contradicts the bound on distance between the slopes of two finite surgeries on a hyperbolic knot provided by Boyer and Zhang [1996]

Case 2: $\bar{\gamma}$ is not primitive. In this case $(X/Z_f)(\bar{\gamma})$ is an orbifold that is a cyclic quotient of \mathcal{P} whose singular set is the core of the filling of X/Z_f . However Lemma A.5 shows that the exterior of any such core would be the exterior of a torus knot in S^3 and hence Seifert fibered, contradicting that X/Z_f must be hyperbolic. \square

The final lemma needed to establish the theorem can be proved in a similar manner to the lemma above.

Lemma A.7. *Let X be the exterior of a hyperbolic knot K in a lens space $L(p, q)$ with $p \geq 2$ admitting a Dehn surgery to \mathcal{P} . Then $|Z(X)| = 1$.*

Proof. Set $Z = Z(X)$ and assume $|Z| > 1$. Let μ be the meridian of K and let γ be the slope of the surgery to \mathcal{P} . Then let $\bar{\mu}$ and $\bar{\gamma}$ be the images of μ and γ in the quotient $\mathcal{Z} = X/Z$.

By Lemma A.6, \mathcal{Z} is an orbifold (with nonempty singular set). Since \mathcal{P} is a homology sphere, \mathcal{Z} is the exterior of a knot \bar{K} in an orbi-lens space $\mathcal{Z}(\bar{\mu})$ that is disjoint from the singular set by Lemma A.3. Because \mathcal{Z} is a suborbifold of $\mathcal{Z}(\bar{\mu})$, the singular set $\Sigma(\mathcal{Z})$ may be identified with the singular set $\Sigma(\mathcal{Z}(\bar{\mu}))$. Thus $\Sigma(\mathcal{Z})$ has at most two components. Since $\mathcal{P} = X(\gamma)$, the quotient orbifold $\mathcal{P}/Z = \mathcal{Z}(\bar{\gamma})$ has a singular set $\Sigma(\mathcal{Z}(\bar{\gamma}))$ consisting of one, two, or three embedded circles according to whether the singular set $\Sigma(\mathcal{Z})$ has one or two components and whether or not the slope $\bar{\gamma}$ is a primitive curve.

Case 1: $\bar{\gamma}$ is not primitive. Then \mathcal{Z} is the exterior of one of the components of $\Sigma(\mathcal{Z}(\bar{\gamma}))$. Hence by Lemmas A.4 and A.5, \mathcal{Z} must be Seifert fibered. Thus X must be Seifert fibered contrary to K being a hyperbolic knot.

Case 2: $\bar{\gamma}$ is primitive. Let \bar{K}_γ be the core of the filling solid torus in $\mathcal{Z}(\bar{\gamma})$ so that the exterior of \bar{K}_γ is \mathcal{Z} . Since $\bar{\gamma}$ is primitive, the singular set $\Sigma = \Sigma(\mathcal{Z}(\bar{\gamma}))$ of $\mathcal{Z}(\bar{\gamma})$ is contained in the knot exterior \mathcal{Z} and thus has either one or two components. Since \mathcal{Z} is a hyperbolic orbifold in which Σ is a geodesic link of one or two components, the exterior of Σ is a hyperbolic manifold $X' = \mathcal{Z}_\Sigma = \mathcal{Z} - \mathcal{N}(\Sigma)$ [Kojima 1988; Sakai 1991].

Let us write $X'(\alpha, \beta)$ (or $X'(\alpha, \beta_1, \beta_2)$) to denote the Dehn filling of X' along the α and β (or α and β_1, β_2) slopes in the components of $\partial X'$ corresponding to $\partial\mathcal{N}(\bar{K}_\gamma) = \partial\mathcal{Z}$ and $\partial\mathcal{N}(\Sigma)$ respectively. We will use $-$ to indicate that the boundary component is left unfilled.

Case 2a: Σ has just one component. By Lemma A.4, $X'(\bar{\gamma}, -)$ is a Seifert fibered space over the disk with two exceptional fibers. Thus $X'(\bar{\gamma}, -)$ has a one-parameter family of slopes σ_i giving lens space fillings $X'(\bar{\gamma}, \sigma_i)$. Since $X'(\bar{\mu}, -)$ is a solid torus (as it is the complement of the core of an orbi-solid torus in an orbifold lens space with connected singular set), $X'(\bar{\mu}, \beta_i)$ is also a lens space for each β_i . Hence, by Thurston’s hyperbolic dehn filling theorem [1979, Theorem 5.8.2], we can choose β_i so that $X'(-, \beta_i)$ is hyperbolic. However, $\Delta(\bar{\mu}, \bar{\gamma}) = |Z|\Delta(\mu, \gamma) \geq 2$, which contradicts the cyclic surgery theorem [Culler et al. 1987].

Case 2b: Σ has two components. This case is nearly identical to the previous.

By [Lemma A.5](#), $X'(\bar{\gamma}, -, -)$ is a Seifert fibered space over the annulus with one exceptional fiber. Then there are infinitely many pairs of slopes $\{(\sigma_n, \sigma'_{n,m})\}_{(n,m) \in \mathbb{Z}^2}$ such that $X'(\bar{\gamma}, \sigma_n, \sigma'_{n,m})$ is a lens space. Since $X'(\bar{\mu}, -, -)$ is a thickened torus (being the complement of the two cores of the orbi-solid tori in the orbi-lens space), $X(\bar{\mu}, \sigma_n, \sigma'_{n,m})$ is a lens space for all these pairs of slopes.

Using Thurston’s hyperbolic Dehn filling theorem twice, there are infinitely many of these pairs of slopes such that $X(-, \sigma_n, \sigma'_{n,m})$ is hyperbolic. Again, since $\Delta(\bar{\mu}, \bar{\gamma}) = |Z| \Delta(\mu, \gamma) \geq 2$, we obtain a contradiction to the cyclic surgery theorem [\[Culler et al. 1987\]](#) for these pairs. \square

It may be tempting to try to use $|Z| \cdot \Delta(\mu, \gamma)$ in the previous lemma place try and improve the bound in [\[Boyer and Zhang 1996\]](#) that $\Delta(\mu, \gamma) \leq 2$. However, this bound is known to be sharp. Also, when $Z(M)$ is trivial, we lose the ability to “match up” the cores of the solid tori coming from the Heegaard splitting of the lens space with the exceptional fibers in \mathcal{P} , and so the arguments in this appendix will not apply.

Proof of [Theorem A.1](#). By [Proposition A.2](#),

$$\text{Sym}(X) = \text{Sym}^+(X).$$

From [Lemma A.3\(2\)](#), $Z(X)$ has index at most 2 in $\text{Sym}(X)$. Then since $Z(X) = 1$ by [Lemma A.7](#), we see that $\text{Sym}(X)$ is either trivial or $\mathbb{Z}/2\mathbb{Z}$. According to [Lemma A.3\(3\)](#), the nontrivial element in this latter case must be a strong involution. \square

Acknowledgments.

The main article benefited from Baker’s discussions with Joshua Greene and Tye Lidman. Parts of this work and writing were done during his visits to the University of Texas and Boston College in the winter of 2015; he thanks them for their hospitality.

Baker was partially supported by a grant from the Simons Foundation (#209184). Hoffman was supported by the Australian Research Council Discovery Grant DP130103694 and by a grant from the Simons Foundation (#524123). The appendix benefited from the thoughtful suggestions and corrections of the referee.

References

- [Baker 2014a] K. L. Baker, “Counting genus one fibered knots in lens spaces”, *Michigan Math. J.* **63**:3 (2014), 553–569. [MR](#) [Zbl](#)
- [Baker 2014b] K. L. Baker, “The Poincaré homology sphere and almost-simple knots in lens spaces”, *Proc. Amer. Math. Soc.* **142**:3 (2014), 1071–1074. [MR](#) [Zbl](#)

- [Baker et al. 2008] K. L. Baker, J. E. Grigsby, and M. Hedden, “Grid diagrams for lens spaces and combinatorial knot Floer homology”, *Int. Math. Res. Not.* **2008**:10 (2008), art. id. rnm024, 39 pp. [MR](#) [Zbl](#)
- [Baker et al. 2016] K. L. Baker, D. Buck, and A. G. Lecuona, “Some knots in $S^1 \times S^2$ with lens space surgeries”, *Comm. Anal. Geom.* **24**:3 (2016), 431–470. [MR](#) [Zbl](#)
- [Baker et al. 2019] K. L. Baker, N. R. Hoffman, and J. E. Licata, “Jointly primitive knots and surgeries between lens spaces”, preprint, 2019. [arXiv](#)
- [Berge 1990] J. Berge, “Some knots with surgeries yielding lens spaces”, preprint, 1990. [arXiv](#)
- [Birman and Menasco 1993] J. S. Birman and W. W. Menasco, “Studying links via closed braids, III: Classifying links which are closed 3-braids”, *Pacific J. Math.* **161**:1 (1993), 25–113. [MR](#) [Zbl](#)
- [Boileau and Otal 1991] M. Boileau and J.-P. Otal, “Scindements de Heegaard et groupe des homéotopies des petites variétés de Seifert”, *Invent. Math.* **106**:1 (1991), 85–107. [MR](#) [Zbl](#)
- [Boileau et al. 2003] M. Boileau, S. Maillot, and J. Porti, *Three-dimensional orbifolds and their geometric structures*, Panoramas et Synthèses **15**, Société Mathématique de France, Paris, 2003. [MR](#) [Zbl](#)
- [Boileau et al. 2012] M. Boileau, S. Boyer, R. Cebanu, and G. S. Walsh, “Knot commensurability and the Berge conjecture”, *Geom. Topol.* **16**:2 (2012), 625–664. [MR](#) [Zbl](#)
- [Boyer and Zhang 1996] S. Boyer and X. Zhang, “Finite Dehn surgery on knots”, *J. Amer. Math. Soc.* **9**:4 (1996), 1005–1050. [MR](#) [Zbl](#)
- [Brody 1960] E. J. Brody, “The topological classification of the lens spaces”, *Ann. of Math. (2)* **71** (1960), 163–184. [MR](#) [Zbl](#)
- [Cooper et al. 2000] D. Cooper, C. D. Hodgson, and S. P. Kerckhoff, *Three-dimensional orbifolds and cone-manifolds*, MSJ Memoirs **5**, Math. Soc. Japan, Tokyo, 2000. [MR](#) [Zbl](#)
- [Culler et al. 1987] M. Culler, C. M. Gordon, J. Luecke, and P. B. Shalen, “Dehn surgery on knots”, *Ann. of Math. (2)* **125**:2 (1987), 237–300. [MR](#) [Zbl](#)
- [Dunbar 1988] W. D. Dunbar, “Geometric orbifolds”, *Rev. Mat. Univ. Complut. Madrid* **1**:1-3 (1988), 67–99. [MR](#) [Zbl](#)
- [Eftekhary 2009] E. Eftekhary, “Seifert fibered homology spheres with trivial Heegaard Floer homology”, preprint, 2009. [arXiv](#)
- [Eftekhary 2011] E. Eftekhary, “Knots which admit a surgery with simple knot Floer homology groups”, *Algebr. Geom. Topol.* **11**:3 (2011), 1243–1256. [MR](#) [Zbl](#)
- [Eftekhary 2018] E. Eftekhary, “Bordered Floer homology and existence of incompressible tori in homology spheres”, *Compos. Math.* **154**:6 (2018), 1222–1268. [MR](#) [Zbl](#)
- [Eudave-Muñoz et al. 2015] M. Eudave-Muñoz, E. Jasso, K. Miyazaki, and K. Motegi, “Seifert fibered surgeries on strongly invertible knots without primitive/Seifert positions”, *Topology Appl.* **196**:B (2015), 729–753. [MR](#) [Zbl](#)
- [Fintushel and Stern 1980] R. Fintushel and R. J. Stern, “Constructing lens spaces by surgery on knots”, *Math. Z.* **175**:1 (1980), 33–51. [MR](#) [Zbl](#)
- [Giroux 2002] E. Giroux, “Géométrie de contact: de la dimension trois vers les dimensions supérieures”, pp. 405–414 in *Proceedings of the International Congress of Mathematicians* (Beijing, 2002), vol. II, edited by T. Li, Higher Ed. Press, Beijing, 2002. [MR](#) [Zbl](#)
- [Goda and Teragaito 2000] H. Goda and M. Teragaito, “Dehn surgeries on knots which yield lens spaces and genera of knots”, *Math. Proc. Cambridge Philos. Soc.* **129**:3 (2000), 501–515. [MR](#) [Zbl](#)

- [Gordon 1991] C. M. Gordon, “Dehn surgery on knots”, pp. 631–642 in *Proceedings of the International Congress of Mathematicians* (Kyoto, 1990), vol. I, Math. Soc. Japan, Tokyo, 1991. [MR](#) [Zbl](#)
- [Greene 2013] J. E. Greene, “The lens space realization problem”, *Ann. of Math.* (2) **177**:2 (2013), 449–511. [MR](#) [Zbl](#)
- [Hedden 2009] M. Hedden, “On knot Floer homology and cabling, II”, *Int. Math. Res. Not.* **2009**:12 (2009), 2248–2274. [MR](#) [Zbl](#)
- [Hedden 2011] M. Hedden, “On Floer homology and the Berge conjecture on knots admitting lens space surgeries”, *Trans. Amer. Math. Soc.* **363**:2 (2011), 949–968. [MR](#) [Zbl](#)
- [Hodgson and Rubinstein 1985] C. Hodgson and J. H. Rubinstein, “Involutions and isotopies of lens spaces”, pp. 60–96 in *Knot theory and manifolds* (Vancouver, 1983), edited by D. Rolfsen, Lecture Notes in Math. **1144**, Springer, 1985. [MR](#) [Zbl](#)
- [Kobayashi 1984] T. Kobayashi, “Structures of the Haken manifolds with Heegaard splittings of genus two”, *Osaka J. Math.* **21**:2 (1984), 437–455. [MR](#) [Zbl](#)
- [Kojima 1988] S. Kojima, “Isometry transformations of hyperbolic 3-manifolds”, *Topology Appl.* **29**:3 (1988), 297–307. [MR](#) [Zbl](#)
- [Kronheimer et al. 2007] P. Kronheimer, T. Mrowka, P. Ozsváth, and Z. Szabó, “Monopoles and lens space surgeries”, *Ann. of Math.* (2) **165**:2 (2007), 457–546. [MR](#) [Zbl](#)
- [Meyerhoff 1986] R. Meyerhoff, “Sphere-packing and volume in hyperbolic 3-space”, *Comment. Math. Helv.* **61**:2 (1986), 271–278. [MR](#) [Zbl](#)
- [Montesinos 1975] J. M. Montesinos, “Surgery on links and double branched covers of S^3 ”, pp. 227–259 in *Knots, groups, and 3-manifolds*, edited by L. P. Neuwirth, Ann. of Math. Studies **84**, 1975. [MR](#) [Zbl](#)
- [Moser 1971] L. Moser, “Elementary surgery along a torus knot”, *Pacific J. Math.* **38** (1971), 737–745. [MR](#) [Zbl](#)
- [Neumann and Reid 1992] W. D. Neumann and A. W. Reid, “Arithmetic of hyperbolic manifolds”, pp. 273–310 in *Topology '90* (Columbus, OH, 1990), edited by B. Apanasov et al., Ohio State Univ. Math. Res. Inst. Publ. **1**, de Gruyter, Berlin, 1992. [MR](#) [Zbl](#)
- [Ni 2007] Y. Ni, “Knot Floer homology detects fibred knots”, *Invent. Math.* **170**:3 (2007), 577–608. [MR](#) [Zbl](#)
- [Ni 2009] Y. Ni, “Link Floer homology detects the Thurston norm”, *Geom. Topol.* **13**:5 (2009), 2991–3019. [MR](#) [Zbl](#)
- [Ozsváth and Szabó 2005] P. Ozsváth and Z. Szabó, “On knot Floer homology and lens space surgeries”, *Topology* **44**:6 (2005), 1281–1300. [MR](#) [Zbl](#)
- [Ozsváth and Szabó 2011] P. S. Ozsváth and Z. Szabó, “Knot Floer homology and rational surgeries”, *Algebr. Geom. Topol.* **11**:1 (2011), 1–68. [MR](#) [Zbl](#)
- [Rasmussen 2007] J. Rasmussen, “Lens space surgeries and L-space homology spheres”, preprint, 2007. [arXiv](#)
- [Reid 1991] A. W. Reid, “Arithmeticity of knot complements”, *J. London Math. Soc.* (2) **43**:1 (1991), 171–184. [MR](#) [Zbl](#)
- [Sakai 1991] T. Sakai, “Geodesic knots in a hyperbolic 3-manifold”, *Kobe J. Math.* **8**:1 (1991), 81–87. [MR](#) [Zbl](#)
- [Tange 2009a] M. Tange, “Lens spaces given from L-space homology 3-spheres”, *Experiment. Math.* **18**:3 (2009), 285–301. [MR](#) [Zbl](#)

- [Tange 2009b] M. Tange, “Ozsváth Szabó’s correction term and lens surgery”, *Math. Proc. Cambridge Philos. Soc.* **146**:1 (2009), 119–134. [MR](#) [Zbl](#)
- [Tange 2011] M. Tange, “On the non-existence of L-space surgery structure”, *Osaka J. Math.* **48**:2 (2011), 541–547. [MR](#) [Zbl](#)
- [Thurston 1979] W. P. Thurston, “The geometry and topology of three-manifolds”, lecture notes, Princeton University, 1979, Available at <http://library.msri.org/books/gt3m/>.
- [Thurston 1997] W. P. Thurston, *Three-dimensional geometry and topology*, vol. 1, edited by S. Levy, Princeton Mathematical Series **35**, Princeton University Press, 1997. [MR](#) [Zbl](#)
- [Wang and Zhou 1992] S. C. Wang and Q. Zhou, “Symmetry of knots and cyclic surgery”, *Trans. Amer. Math. Soc.* **330**:2 (1992), 665–676. [MR](#) [Zbl](#)

Received May 16, 2017. Revised June 27, 2019.

KENNETH L. BAKER
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MIAMI
CORAL GABLES, FL
UNITED STATES
k.baker@math.miami.edu

NEIL R. HOFFMAN
DEPARTMENT OF MATHEMATICS
OKLAHOMA STATE UNIVERSITY
STILLWATER, OK
UNITED STATES
neil.r.hoffman@okstate.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 305 No. 1 March 2020

The Poincaré homology sphere, lens space surgeries, and some knots with tunnel number two	1
KENNETH L. BAKER	
Fusion systems of blocks of finite groups over arbitrary fields	29
ROBERT BOLTJE, ÇISIL KARAGÜZEL and DENİZ YILMAZ	
Torsion points and Galois representations on CM elliptic curves	43
ABBEY BOURDON and PETE L. CLARK	
Stability of the positive mass theorem for axisymmetric manifolds	89
EDWARD T. BRYDEN	
Index estimates for free boundary constant mean curvature surfaces	153
MARCOS P. CAVALCANTE and DARLAN F. DE OLIVEIRA	
A criterion for modules over Gorenstein local rings to have rational Poincaré series	165
ANJAN GUPTA	
Generalized Cartan matrices arising from new derivation Lie algebras of isolated hypersurface singularities	189
NAVEED HUSSAIN, STEPHEN S.-T. YAU and HUAIQING ZUO	
On the commutativity of coset pressure	219
BING LI and WEN-CHIAO CHENG	
Signature invariants related to the unknotting number	229
CHARLES LIVINGSTON	
The global well-posedness and scattering for the 5-dimensional defocusing conformal invariant NLW with radial initial data in a critical Besov space	251
CHANGXING MIAO, JIANWEI YANG and TENGFEI ZHAO	
Liouville-type theorems for weighted p-harmonic 1-forms and weighted p-harmonic maps	291
KEOMKYO SEO and GABJIN YUN	
Remarks on the Hölder-continuity of solutions to parabolic equations with conic singularities	311
YUANQI WANG	
Deformation of Milnor algebras	329
ZHENJIAN WANG	
Preservation of log-Sobolev inequalities under some Hamiltonian flows	339
BO XIA	
Ground state solutions of polyharmonic equations with potentials of positive low bound	353
CAIFENG ZHANG, JUNGANG LI and LU CHEN	