

Pacific Journal of Mathematics

**INDEX ESTIMATES FOR FREE BOUNDARY
CONSTANT MEAN CURVATURE SURFACES**

MARCOS P. CAVALCANTE AND DARLAN F. DE OLIVEIRA

INDEX ESTIMATES FOR FREE BOUNDARY CONSTANT MEAN CURVATURE SURFACES

MARCOS P. CAVALCANTE AND DARLAN F. DE OLIVEIRA

We consider compact constant mean curvature surfaces with boundary immersed in a mean convex region of the Euclidean space or in the unit sphere. We prove that the weak Morse index is bounded from below by a linear function of the genus and number of boundary components.

1. Introduction

Let W be a Riemannian manifold with nonempty boundary such that its boundary ∂W is a union of smooth hypersurfaces. Let $M \subset W$ be a compact constant mean curvature hypersurface such that M intersects the regular part of ∂W along its boundary in a right angle. It is well known that such hypersurfaces are critical points of the area functional for variations of M that preserve the enclosed volume and keep the boundary freely on ∂W . We recall that the variations allowed of M are variations $\phi : (-\varepsilon, \varepsilon) \times M \rightarrow W$ whose immersions $\phi_t : M \rightarrow W$ satisfy $\phi_t(\text{int } M) \subset \text{int } W$ and $\phi_t(\partial M) \subset \text{int } \partial W$ for all $t \in (-\varepsilon, \varepsilon)$; see [Ros and Vergasta 1995, Section 1]. These hypersurfaces arise in many geometrical and physical problems and are referred as *free boundary CMC hypersurfaces* (FBCMC hypersurfaces, for short). They have been studied since the 19th century and still form a very active topic in differential geometry. We refer the reader to the books of Finn [1986] and López [2013] for a nice introduction to this subject.

An important problem about FBCMC hypersurfaces is to classify those ones that are *stable*, that is, whose second variation of the area is nonnegative for volume preserving variations. For instance, in the case that W is a geodesic ball in a space form, a well-known conjecture asserts that the totally geodesic ball and the spherical caps are the only solutions. It was confirmed by Ros and Vergasta [1995] and Nunes [2017] in dimension two, and more recently by Wang and Xia [2019] in any dimension and also for capillary hypersurfaces. Other results on stable FBCMC hypersurfaces can be found for instance in [Ainouz and Souam 2016; Athanassenas

Cavalcante was partially supported by CNPq-Brazil (309543/2015-0), CAPES-Brazil (897/18) and FAPEAL-Brazil (14/2016).

MSC2010: 35P15, 49Q10, 53A10.

Keywords: constant mean curvature surfaces, Morse index, free boundary surfaces.

1987; Barbosa 2018; Choe and Koiso 2016; Li and Xiong 2017; 2018; López 2014; Ros 2008; Ros and Souam 1997].

When M is a stable FBCMC surface immersed in a mean convex region $W \subset \mathbb{R}^3$, Ros [2008, Theorem 9] showed that there are just a few possibilities for the genus and the number of boundary components of M (see Corollary 1.2 below).

When M is not stable, there is a Schrödinger operator L associated to the second variation of the area which has nonzero *weak Morse index* (see Section 2 for precise definitions). Geometrically, the weak Morse index of M is the number of directions whose volume preserving variations decrease area. It will be denoted by $\text{Ind}_w(M)$.

The case of free boundary minimal surfaces is of special interest and the works of Fraser and Schoen [2011; 2013; 2016] have motivated much research in this case. For free boundary minimal hypersurfaces all variations keeping the boundary freely in the boundary are allowed, not only volume preserving variations. In this setting Sargent [2017] proved that if M is a free boundary minimal surface immersed in a convex region $W \subset \mathbb{R}^3$ then the index is bounded from below by $(2g + k - 1)/3$, where g is the genus of M and k is the number of boundary components. In [Ambrozio et al. 2018], Ambrozio, Carlotto and Sharp proved that it is valid in weakly mean convex domains in $\mathbb{R}^3$. In fact, their results apply more generally to domains in Riemannian manifolds with boundary satisfying suitable curvature conditions. In higher dimensions, they also obtained lower bounds for the index in terms of the dimension of the first relative homology group with real coefficients. The technique presented in these results uses the coordinates of harmonic forms as test functions and is inspired by previous works on eigenvalue estimates and index estimates for minimal hypersurfaces without boundary; see [Ros 2006; Savo 2010; Ambrozio et al. 2018].

Following these lines, in this paper, we obtain lower bounds for the weak Morse index of FBCMC surfaces in weakly mean convex regions of the Euclidean space or the unit sphere. More precisely, our results are the following.

Theorem 1.1. *Let W be a region of $\mathbb{R}^3$ such that its boundary is a union of smooth weakly mean convex surfaces, and let M^2 be a compact, orientable, FBCMC surface immersed in the mean convex side of W and whose boundary intersects the regular part of ∂W . If M has genus g and k boundary components, then*

$$\text{Ind}_w(M) \geq \frac{2g+k-4}{6}.$$

As an immediate consequence we obtain the result of Ros cited above:

Corollary 1.2 [Ros 2008, Theorem 9]. *Under the conditions of Theorem 1.1, if M is stable, then the only possibilities for g and k are*

- (1) $g = 0$ and $k \leq 4$;
- (2) $g = 1$ and $k = 1$ or 2 .

In the case of FBCMC surfaces immersed in weakly mean convex domains of $\mathbb{S}^3$, the result reads as follows:

Theorem 1.3. *Let W be a region in the unit sphere $\mathbb{S}^3$ such that its boundary is a union of smooth weakly mean convex surfaces, and let M^2 be a compact, orientable, FBCMC surface immersed in the mean convex side of W and whose boundary intersects the regular part of ∂W . If M has genus g and k boundary components,*

$$\text{Ind}_w(M) \geq \frac{2g+k-5}{8}.$$

Corollary 1.4. *Under the conditions of [Theorem 1.3](#), if M is stable, then the only possibilities for g and k are*

- (1) $g = 0$ and $k \leq 5$;
- (2) $g = 1$ and $k \leq 3$;
- (3) $g = 2$ and $k = 1$.

This paper is organized as follows. In [Section 2](#), we present some definitions and basic results to be used in the proofs. [Section 3](#) is devoted to computing the Jacobi operator of the test functions given by the coordinates of harmonic forms. In [Section 4](#), we present the proof of [Theorem 1.1](#). The proof of [Theorem 1.3](#) is analogous and it is sketched in [Section 5](#).

2. Preliminaries

Let us denote by W a connected domain of the Euclidean space $\mathbb{R}^3$ which is not necessarily compact. For simplicity, let us assume that W has smooth boundary and fix a unit normal vector field ν along each component of ∂W . We recall that the second fundamental form and the mean curvature of ∂W with respect to ν are defined respectively by

$$II^{\partial W}(X, Y) = \langle -D_X \nu, Y \rangle, \quad \text{for } X, Y \in T\partial W,$$

and

$$H^{\partial W} = \frac{1}{2} \text{tr } II^{\partial W},$$

where D is the Levi-Civita connection in the Euclidean space. Since the boundary ∂W is orientable, at a point on ∂W we have two choices for ν , one pointing inward from W and the other one pointing outward from W . From now on, we fix the vector field ν pointing outward from W . In this case, we say that W is *weakly convex* if $II^{\partial W}$ is nonpositive defined. If $H^{\partial W} \leq 0$, W is said to be *weakly mean convex*.

Let $x : M \rightarrow W$ be a compact oriented surfaced with boundary which is properly immersed, that is, $x(M) \cap W = x(\partial M)$. Fixing a unit normal vector field N along x ,

we denote by A the shape operator associated to the second fundamental form II^M of M with respect to N , namely

$$\begin{aligned} AX &= -D_X N, & \text{for } X \in TM, \\ II^M(X, Y) &= \langle AX, Y \rangle, & \text{for } X, Y \in TM. \end{aligned}$$

We say that M is free boundary if $x(\partial M)$ meets ∂W orthogonally.

From now on, let us assume that W is a mean convex domain of $\mathbb{R}^3$ and M is a *free boundary constant mean curvature surface* properly immersed in W . Such surfaces are critical points of the area functional for normal variations whose variational vector field is given by $X = uN$, where $u \in \mathcal{F}$ and

$$\mathcal{F} = \left\{ u : M \rightarrow \mathbb{R} : u \text{ is smooth up to the boundary and } \int_M u \, dM = 0 \right\}.$$

The second variation of area functional is given by the quadratic form $Q : C^\infty(M) \times C^\infty(M) \rightarrow \mathbb{R}$ (see [Ros and Vergasta 1995; Ros 2008]),

$$Q(u, u) = \int_M (u \Delta u - \|A\|^2 u^2) \, dM + \int_{\partial M} (u \eta(u) + II^{\partial W}(N, N) u^2) \, ds.$$

Here η is the outward unit conormal vector field on ∂M , that is, the unique unit vector field on ∂M that is tangent to M , normal to ∂M and pointing outward from M . Note that, under our notations, the free boundary condition means that $\eta = \nu$ along ∂M . We point out that in this paper we are using the geometric definition of the Laplacian operator, that is, $\Delta u = \operatorname{div} \nabla u$, where $\operatorname{div} X = -\operatorname{tr} \nabla X$.

The *weak Morse index* of M , denoted by $\operatorname{Ind}_w(M)$, is defined as the maximal dimension of a subspace of $\mathcal{F}$ on which $Q|_{\mathcal{F}}$ is negative definite. Geometrically, the index indicates the number of directions whose variations decrease area. In particular, we say that M is *stable* if the weak Morse index is zero.

We say that $u \in \mathcal{F}$ is an eigenfunction of $Q|_{\mathcal{F}}$ associated to the eigenvalue $\lambda \in \mathbb{R}$ if and only if $Q|_{\mathcal{F}}(u, v) = \lambda \int_M uv \, dM$ for all $v \in \mathcal{F}$. This is equivalent to saying that u solves the following eigenvalue problem:

$$(2-1) \quad \begin{cases} Lu = \lambda u & \text{in } M, \\ \frac{\partial u}{\partial \eta} = -II^{\partial W}(N, N)u & \text{on } \partial M, \end{cases}$$

where $L : \mathcal{F} \rightarrow \mathcal{F}$ is given by

$$(2-2) \quad Lu = Ju - \frac{1}{\operatorname{vol}(M)} \int_M Ju \, dM$$

and $J = \Delta - \|A\|^2$ is the Jacobi operator. We conclude that the weak Morse index coincides with the number of negative eigenvalues of the boundary problem (2-1).

Also, it is well known that such eigenvalues are given in a nondecreasing sequence $\lambda_1^L \leq \lambda_2^L \leq \dots \leq \lambda_k^L \leq \dots \nearrow \infty$ associated to a $L^2(M)$ -orthonormal basis,

$$\{\phi_1, \phi_2, \dots, \phi_k, \dots\} \quad \text{of} \quad L^2(M) \cap \mathcal{F}$$

of solutions of the eigenvalue problem (2-1), satisfying the min-max characterization

$$\lambda_k^L = \min_{u \in \mathcal{J}^{k-1} \setminus \{0\}} \frac{Q|_{\mathcal{F}}(u, u)}{\int_M u^2 dM},$$

where $\mathcal{J}^{k-1} = \langle \phi_1, \dots, \phi_{k-1} \rangle^\perp$.

In order to give lower bounds for the weak Morse index of M in terms of its topological invariants we will construct admissible eigenfunctions in $\mathcal{F}$ using harmonic vector fields, or equivalently harmonic 1-forms. Let us denote by $i : \partial M \rightarrow M$ the inclusion map. We also set $\mathcal{H}_T^1(M)$ the space of closed and coclosed 1-forms that are tangential at ∂M , that is,

$$\mathcal{H}_T^1(M) := \{w \in \Omega^1(M), dw = 0, \delta w = 0 \text{ and } i_\eta w = 0 \text{ along } \partial M\}.$$

Here d is the exterior derivative operator and δ is the interior derivative operator defined by $\delta = -\star d\star$, where $\star : \Omega^1(M) \rightarrow \Omega^1(M)$ is the Hodge star operator.

It is important to note that $\mathcal{H}_T^1(M)$ coincides with the space of harmonic 1-forms satisfying the *absolute boundary conditions*, that is

$$\mathcal{H}_T^1(M) = \{w \in \Omega^1(M), \Delta w = 0, i_\eta w = 0 \text{ and } i_\eta dw = 0 \text{ along } \partial M\}.$$

This space is closely related to the topology of the underline manifold. In fact we have the following result; see [Ambrozio et al. 2018] or [Sargent 2017].

Lemma 2.1. *Let M^2 be a compact, orientable surface with nonempty boundary ∂M . If M has genus g and $k \geq 1$ boundary components, then*

$$\dim \mathcal{H}_T^1(M) = 2g + k - 1.$$

3. Test functions and harmonic vector fields

Denoting by $\mathcal{E} = \{\bar{E}_1, \bar{E}_2, \bar{E}_3\}$ the canonical basis in $\mathbb{R}^3$ we will consider $E_i := \bar{E}_i - \langle \bar{E}_i, N \rangle N$, the orthogonal projection of $\bar{E}_i$ on TM . We also consider the smooth support functions $g_i : M \rightarrow \mathbb{R}$, $g_i := \langle \bar{E}_i, N \rangle$, for $1 \leq i \leq 3$.

Given a 1-form ω on M we denote by ξ its dual vector field, that is, $\xi^\flat = \omega$. Abusing notation slightly, we denote by $\star \xi$ the vector field dual of $\star \omega$. In the following, we will use the coordinates of ξ and $\star \xi$ as test functions. Namely, for each $1 \leq i \leq 3$, we define $w_i, \bar{w}_i : M \rightarrow \mathbb{R}$ as

$$w_i := \omega(E_i) = \langle E_i, \xi \rangle \quad \text{and} \quad \bar{w}_i := \star \omega(E_i) = \langle E_i, \star \xi \rangle.$$

To compute the Jacobi operator of w_i and $\bar{w}_i$ we need the following lemma of local nature proved in [Cavalcante and de Oliveira 2020]; see also [Ros 2007].

Lemma 3.1. *Let M^2 be an orientable CMC surface in $\mathbb{R}^3$. Then, using the above notation we have*

$$\begin{aligned}\Delta w_i &= (\|A\|^2 - 4H^2)w_i + 2H\langle AE_i, \xi \rangle - 2g_i\langle A, \nabla \xi \rangle + \langle E_i, \Delta \xi \rangle, \\ \Delta \bar{w}_i &= (\|A\|^2 - 4H^2)\bar{w}_i + 2H\langle AE_i, \star \xi \rangle - 2g_i\langle A, \nabla \star \xi \rangle + \langle E_i, \Delta \star \xi \rangle,\end{aligned}$$

for $1 \leq i \leq 3$.

Now we note that when the vector field ξ is harmonic and tangential along ∂M its coordinates are admissible functions to compute the weak Morse index of CMC surfaces. More precisely we have:

Lemma 3.2. *If $\xi \in TM$ is a harmonic vector field which is tangential in ∂M , then $w_i \in \mathcal{F}$, that is,*

$$\int_M w_i dM = 0,$$

for $1 \leq i \leq 3$.

Proof. Note that $E_i = \nabla x_i$, $1 \leq i \leq 3$, where $x = (x_1, x_2, x_3) : M \rightarrow W$ is the immersion map. Then we have

$$\begin{aligned}\int_M w_i dM &= \int_M \langle \nabla x_i, \xi \rangle dM \\ &= \int_M x_i \operatorname{div} \xi dM + \int_{\partial M} x_i \langle \xi, \eta \rangle ds = 0.\end{aligned}$$

In fact, $\operatorname{div} \xi = 0$ since ξ is harmonic, and $\langle \xi, \eta \rangle = 0$ since ξ tangential to ∂M . $\square$

Remark 3.3. In general the functions $\bar{w}_i$, $1 \leq i \leq 3$, do not have mean value zero. However, we will see in [Section 4](#) that if $\dim \mathcal{H}_T^1(M)$ is large enough then we can choose ξ such that

$$\int_M \bar{w}_i dM = 0, \quad \text{for } 1 \leq i \leq 3.$$

We conclude this section by computing the boundary term of the quadratic form Q on w_i and $\bar{w}_i$.

Lemma 3.4. *If $\xi \in TM$ is a vector field such that its dual 1-form satisfies the absolute boundary condition, then*

$$(3-1) \quad \sum_i \int_{\partial M} (w_i \eta(w_i) + II^{\partial W}(N, N) w_i^2) ds = 2 \int_{\partial M} H^{\partial W} \|\xi\|^2 ds,$$

$$(3-2) \quad \sum_i \int_{\partial M} (\bar{w}_i \eta(\bar{w}_i) + II^{\partial W}(N, N) \bar{w}_i^2) ds = 2 \int_{\partial M} H^{\partial W} \|\xi\|^2 ds.$$

Proof. We first note that for any vector field $X \in TM$ we have

$$\begin{aligned}\langle \nabla_\eta E_i, X \rangle &= \eta \langle \bar{E}_i, X \rangle - \langle \bar{E}_i, \nabla_\eta X \rangle \\ &= \langle \bar{E}_i, D_\eta X - \nabla_\eta X \rangle \\ &= \langle \bar{E}_i, N \rangle \langle X, A\eta \rangle.\end{aligned}$$

Let ω be the dual 1-form of the vector field ξ . Since $i_\eta d\omega = 0$ we have

$$0 = d\omega(\eta, \xi) = \langle \nabla_\eta \xi, \xi \rangle - \langle \nabla_\xi \xi, \eta \rangle.$$

Thus,

$$\begin{aligned}\sum_i \int_{\partial M} w_i \eta(w_i) ds &= \sum_i \int_{\partial M} w_i (\langle \nabla_\eta \xi, E_i \rangle + \langle \bar{E}_i, N \rangle \langle \xi, A\eta \rangle) ds \\ &= \int_{\partial M} \langle \nabla_\eta \xi, \xi \rangle ds = \int_{\partial M} \langle \nabla_\xi \xi, \eta \rangle ds \\ &= - \int_{\partial M} \langle \nabla_\xi \eta, \xi \rangle ds = \int_{\partial M} II^{\partial W}(\xi, \xi) ds.\end{aligned}$$

Since ξ and N form an orthogonal basis of the tangent space of ∂W along ∂M we conclude the proof by noting that

$$II^{\partial W}(\xi, \xi) + II^{\partial W}(N, N) \|\xi\|^2 = 2H^{\partial W} \|\xi\|^2.$$

The proof of assertion (3-2) follows the same steps as above, noting additionally that the Levi-Civita connection ∇ commutes with the Hodge star operator $\star$. $\square$

4. Proof of Theorem 1.1

Proof. The proofs follow the same spirit as our proofs in [Cavalcante and de Oliveira 2020] but take into account the boundary term. Let $\xi_1, \xi_2, \dots, \xi_m$, be the first m eigenfunctions of the Hodge Laplacian Δ on M , which satisfy the absolute boundary condition [Gilkey et al. 1999, Theorem 1.5.4]. Set $\mathcal{L}_m^\Delta = \text{span}\{\xi_1, \dots, \xi_m\}$ the vector space generated by these functions. By Lemma 2.1, we know that $\dim \mathcal{H}_T^1(M) = 2g + k - 1$. Let us assume that $m \geq 2g - k - 1$, and so $\mathcal{H}_T^1(M)$ is a subspace of $\mathcal{L}_m^\Delta$.

Next, we choose an orthonormal basis of $L^2(M)$ given by eigenfunctions of the operator L defined in (2-2), say $\{\phi_1, \phi_2, \dots, \phi_k, \dots\}$. We denote by $\mathcal{J}^n := \langle \phi_1, \dots, \phi_n \rangle^\perp$ the linear subspace of $\mathcal{F}$ orthogonal to the first n eigenfunctions of L .

Initially, we look for harmonic forms $\xi \in \mathcal{L}_m^\Delta$ such that the functions $w_i, \bar{w}_i \in \mathcal{J}^{\alpha-1}$, for some $\alpha \in \mathbb{N}$ and $i \in \{1, 2, 3\}$. It is equivalent to find a solution to the following system with $6(\alpha - 1)$ homogenous linear equations in the variable ξ :

$$(4-1) \quad \int_M w_i \phi_k dM = \int_M \bar{w}_i \phi_k dM = 0,$$

where $1 \leq i \leq 3$ and $1 \leq k \leq \alpha - 1$. In particular, if $m(\alpha) := \dim \mathcal{L}_m^\Delta > 6(\alpha - 1)$, then

the system (4-1) has at least one nontrivial solution $\xi \in \mathcal{L}_m^\Delta$ such that $w_i, \bar{w}_i \in \mathcal{J}^{\alpha-1}$ for all $1 \leq i \leq 3$. By min-max characterization we have

$$\lambda_\alpha^J \int_M w_i^2 dM \leq Q(w_i, w_i) \quad \text{and} \quad \lambda_\alpha^J \int_M \bar{w}_i^2 dM \leq Q(\bar{w}_i, \bar{w}_i).$$

Now, using Lemma 3.1 we get

$$\begin{aligned} \lambda_\alpha^J \int_M w_i^2 dM &\leq -4H^2 \int_M w_i^2 dM + 2H \int_M \langle E_i, A\xi \rangle w_i dM \\ &\quad + \int_M \langle E_i, \Delta\xi \rangle w_i dM - 2 \int_M g_i \langle A, \nabla\xi \rangle w_i dM \\ &\quad + \int_{\partial M} (w_i \eta(w_i) + H^{\partial W(N,N)} w_i^2) ds. \end{aligned}$$

Summing up $i = 1, 2, 3$ and using Lemma 3.4 we obtain

$$\begin{aligned} \lambda_\alpha^J \int_M \|\xi\|^2 dM &\leq -4H^2 \int_M \|\xi\|^2 dM + 2H \int_M \langle A\xi, \xi \rangle dM \\ &\quad + \int_M \langle \Delta\xi, \xi \rangle dM + 2 \int_{\partial M} H^{\partial W} \|\xi\|^2 dM. \end{aligned}$$

Applying the same arguments to the test functions $\bar{w}_i$ we get

$$\begin{aligned} \lambda_\alpha^J \int_M \|\xi\|^2 dM &\leq -4H^2 \int_M \|\xi\|^2 dM + 2H \int_M \langle A\star\xi, \star\xi \rangle dM \\ &\quad + \int_M \langle \Delta\star\xi, \star\xi \rangle dM + 2 \int_{\partial M} H^{\partial W} \|\xi\|^2 ds. \end{aligned}$$

Then, summing these last two inequalities and noting that $\langle A\xi, \xi \rangle + \langle A\star\xi, \star\xi \rangle = 2H\|\xi\|^2$, we have

$$\begin{aligned} (4-2) \quad \lambda_\alpha^J \int_M \|\xi\|^2 dM &\leq +2 \int_{\partial M} H^{\partial W} \|\xi\|^2 dM - 2H^2 \int_M \|\xi\|^2 dM \\ &\quad + \frac{1}{2} \int_M (\langle \Delta\xi, \xi \rangle + \langle \Delta\star\xi, \star\xi \rangle) dM. \end{aligned}$$

Finally, if $\xi \in \mathcal{L}_m^\Delta$ we get $\xi = \sum_i \alpha_i \xi_i$ and therefore

$$(4-3) \quad \int_M \langle \Delta\star\xi, \star\xi \rangle dM = \int_M \langle \Delta\xi, \xi \rangle dM = \lambda_{m(\alpha)} \int_M \|\xi\|^2 dM.$$

Substituting (4-3) into (4-2) and using the fact that $H^{\partial W} \leq 0$ we obtain

$$\lambda_\alpha^J \leq -2H^2 + \lambda_{m(\alpha)}^\Delta,$$

where $m(\alpha) > 6(\alpha - 1)$. This concludes the first part of Theorem 1.1.

In order to get the lower bound for the weak Morse index of M we take $\mathcal{J}^{\alpha-1} := \langle \phi_1, \dots, \phi_{\alpha-1} \rangle$, where $\phi_1, \dots, \phi_{\alpha-1}$ are the first eigenfunctions of the eigenvalue equation (2-1). From Lemma 3.2 we know that if $\xi \in \mathcal{H}_T^1(M)$, then the test functions w_1, w_2 and w_3 , belong to $\mathcal{F}$.

We look for vector fields $\xi \in \mathcal{H}_T^1(M)$ such that for $1 \leq i \leq 3$, the test functions $w_i, \bar{w}_i$ are in $\mathcal{J}^{\alpha-1}$, for some $\alpha \in \mathbb{N}$, and $\bar{w}_i$ is in $\mathcal{F}$. In this case, we have the following system with $6\alpha - 3$ homogeneous linear equations in the variable ξ :

$$(4-4) \quad \int_M \bar{w}_i = \int_M w_i \phi_k = \int_M \bar{w}_i \phi_k = 0,$$

where $1 \leq i \leq 3$ and $1 \leq k \leq \alpha - 1$.

If $\dim \mathcal{H}_T^1(M) = 2g + k - 1 > 6\alpha - 3$, then the system (4-4) has at least one nontrivial solution $\xi \in \mathcal{H}_T^1(M)$. Following the same steps as above we get

$$\lambda_\alpha^L \int_M \|\xi\|^2 \leq -2H^2 \int_M \|\xi\|^2.$$

This implies that $\lambda_\alpha^L < 0$ and then $\text{Ind}_w(M) \geq \alpha$. Since α can be chosen as the largest integer such that $2g + k - 1 > 6\alpha - 3$ we get

$$\text{Ind}_w(M) \geq \frac{2g + k - 4}{6}.$$

□

5. Proof of Theorem 1.3

Proof. Composing the immersion $x : M \rightarrow W \subset \mathbb{S}^3$ with the canonical immersion of the unit sphere into the Euclidean space, we may consider $x : M \rightarrow \mathbb{R}^4$. Let $\mathcal{E} = \{\bar{E}_1, \bar{E}_2, \bar{E}_3, \bar{E}_4\}$ be the canonical basis in $\mathbb{R}^4$ and $E_i := \bar{E}_i - \langle \bar{E}_i, N \rangle N - \langle \bar{E}_i, x \rangle x$, the orthogonal projections of $\bar{E}_i$ on TM . Choosing $\nu = -x$ as an orientation of $\mathbb{S}^3$ we have

$$D_Y X - \nabla_Y X = \langle AX, Y \rangle N + \langle X, Y \rangle \nu, \quad X, Y \in TM,$$

and also

$$\begin{aligned} \langle \nabla_X E_i, Y \rangle &= X \langle \bar{E}_i, Y \rangle - \langle \bar{E}_i, \nabla_X Y \rangle = \langle \bar{E}_i, D_Y X - \nabla_Y X \rangle \\ &= \langle AX, Y \rangle \langle \bar{E}_i, N \rangle + \langle X, Y \rangle \langle \bar{E}_i, \nu \rangle. \end{aligned}$$

Using that $\langle \xi, N \rangle = \langle \star \xi, N \rangle = \langle \xi, \nu \rangle = \langle \star \xi, \nu \rangle = 0$ we get

$$\begin{aligned} \sum_i \int_{\partial M} w_i \eta(w_i) ds &= \sum_i \int_{\partial M} w_i (\langle \nabla_\eta \xi, E_i \rangle) ds \\ &\quad + \sum_i \int_{\partial M} w_i (\langle \bar{E}_i, N \rangle \langle \xi, A\eta \rangle + \langle \eta, \xi \rangle \langle \bar{E}_i, \nu \rangle) ds \\ &= \int_{\partial M} \langle \nabla_\eta \xi, \xi \rangle ds + \int_{\partial M} \langle \xi, N \rangle \langle \xi, A\eta \rangle + \int_{\partial M} \langle \eta, \xi \rangle \langle \xi, \nu \rangle \\ &= \int_{\partial M} \langle \nabla_\eta \xi, \xi \rangle ds, \end{aligned}$$

and analogously

$$\sum_i \int_{\partial M} \bar{w}_i \eta(\bar{w}_i) ds = \int_{\partial M} \langle \nabla_\eta \star \xi, \star \xi \rangle ds = \int_{\partial M} \langle \nabla_\eta \xi, \xi \rangle ds.$$

So, [Lemma 3.4](#) holds in the spherical case. The Laplacian of the test functions w_i and $\bar{w}_i$ are given by (see [\[Cavalcante and de Oliveira 2020\]](#))

$$\begin{aligned} \Delta w_i &= (\|A\|^2 - 4H^2)w_i + 2H\langle AE_i, \xi \rangle - 2g_i\langle A, \nabla \xi \rangle + \langle E_i, \Delta \xi \rangle - 2\langle x, \bar{E}_i \rangle \operatorname{div} \xi, \\ \Delta \bar{w}_i &= (\|A\|^2 - 4H^2)\bar{w}_i + 2H\langle AE_i, \bar{\xi} \rangle - 2g_i\langle A, \nabla \bar{\xi} \rangle + \langle E_i, \Delta \bar{\xi} \rangle - 2\langle x, \bar{E}_i \rangle \operatorname{div} \bar{\xi}. \end{aligned}$$

Under these considerations, and taking into account that the Jacobi operator for immersions into $\mathbb{S}^3$ is given by $J = \Delta - (\|A\|^2 + 2)$ the proof follows as in the proof of [Theorem 1.1](#). $\square$

Acknowledgments

The authors would like to thank Lucas Ambrozio and Ivaldo Nunes for useful comments, and Fernando Codá Marques for illuminating conversations about this work.

References

- [Ainouz and Souam 2016] A. Ainouz and R. Souam, “Stable capillary hypersurfaces in a half-space or a slab”, *Indiana Univ. Math. J.* **65**:3 (2016), 813–831. [MR](#) [Zbl](#)
- [Ambrozio et al. 2018] L. Ambrozio, A. Carlotto, and B. Sharp, “Index estimates for free boundary minimal hypersurfaces”, *Math. Ann.* **370**:3-4 (2018), 1063–1078. [MR](#) [Zbl](#)
- [Athanasenas 1987] M. Athanassenas, “A variational problem for constant mean curvature surfaces with free boundary”, *J. Reine Angew. Math.* **377** (1987), 97–107. [MR](#) [Zbl](#)
- [Barbosa 2018] E. Barbosa, “On CMC free-boundary stable hypersurfaces in a Euclidean ball”, *Math. Ann.* **372**:1-2 (2018), 179–187. [MR](#) [Zbl](#)
- [Cavalcante and de Oliveira 2020] M. P. Cavalcante and D. F. de Oliveira, “Lower bounds for the index of compact constant mean curvature surfaces in $\mathbb{R}^3$ and $\mathbb{S}^3$ ”, *Rev. Mat. Iberoam.* **36**:1 (2020), 195–206. [MR](#)
- [Choe and Koiso 2016] J. Choe and M. Koiso, “Stable capillary hypersurfaces in a wedge”, *Pacific J. Math.* **280**:1 (2016), 1–15. [MR](#) [Zbl](#)
- [Finn 1986] R. Finn, *Equilibrium capillary surfaces*, Grundlehren der Mathematischen Wissenschaften **284**, Springer, 1986. [MR](#) [Zbl](#)
- [Fraser and Schoen 2011] A. Fraser and R. Schoen, “The first Steklov eigenvalue, conformal geometry, and minimal surfaces”, *Adv. Math.* **226**:5 (2011), 4011–4030. [MR](#) [Zbl](#)
- [Fraser and Schoen 2013] A. Fraser and R. Schoen, “Minimal surfaces and eigenvalue problems”, pp. 105–121 in *Geometric analysis, mathematical relativity, and nonlinear partial differential equations*, edited by M. Ghomi et al., Contemp. Math. **599**, Amer. Math. Soc., Providence, RI, 2013. [MR](#) [Zbl](#)
- [Fraser and Schoen 2016] A. Fraser and R. Schoen, “Sharp eigenvalue bounds and minimal surfaces in the ball”, *Invent. Math.* **203**:3 (2016), 823–890. [MR](#) [Zbl](#)
- [Gilkey et al. 1999] P. B. Gilkey, J. V. Leahy, and J. Park, *Spectral geometry, Riemannian submersions, and the Gromov–Lawson conjecture*, Chapman & Hall, Boca Raton, FL, 1999. [MR](#) [Zbl](#)

- [Li and Xiong 2017] H. Li and C. Xiong, “Stability of capillary hypersurfaces with planar boundaries”, *J. Geom. Anal.* **27**:1 (2017), 79–94. [MR](#) [Zbl](#)
- [Li and Xiong 2018] H. Li and C. Xiong, “Stability of capillary hypersurfaces in a Euclidean ball”, *Pacific J. Math.* **297**:1 (2018), 131–146. [MR](#) [Zbl](#)
- [López 2013] R. López, *Constant mean curvature surfaces with boundary*, Springer, 2013. [MR](#) [Zbl](#)
- [López 2014] R. López, “Capillary surfaces with free boundary in a wedge”, *Adv. Math.* **262** (2014), 476–483. [MR](#) [Zbl](#)
- [Nunes 2017] I. Nunes, “On stable constant mean curvature surfaces with free boundary”, *Math. Z.* **287**:1-2 (2017), 473–479. [MR](#) [Zbl](#)
- [Ros 2006] A. Ros, “One-sided complete stable minimal surfaces”, *J. Differential Geom.* **74**:1 (2006), 69–92. [MR](#) [Zbl](#)
- [Ros 2007] A. Ros, “Stable periodic constant mean curvature surfaces and mesoscopic phase separation”, *Interfaces Free Bound.* **9**:3 (2007), 355–365. [MR](#) [Zbl](#)
- [Ros 2008] A. Ros, “Stability of minimal and constant mean curvature surfaces with free boundary”, *Mat. Contemp.* **35** (2008), 221–240. [MR](#) [Zbl](#)
- [Ros and Souam 1997] A. Ros and R. Souam, “On stability of capillary surfaces in a ball”, *Pacific J. Math.* **178**:2 (1997), 345–361. [MR](#) [Zbl](#)
- [Ros and Vergasta 1995] A. Ros and E. Vergasta, “Stability for hypersurfaces of constant mean curvature with free boundary”, *Geom. Dedicata* **56**:1 (1995), 19–33. [MR](#) [Zbl](#)
- [Sargent 2017] P. Sargent, “Index bounds for free boundary minimal surfaces of convex bodies”, *Proc. Amer. Math. Soc.* **145**:6 (2017), 2467–2480. [MR](#) [Zbl](#)
- [Savo 2010] A. Savo, “Index bounds for minimal hypersurfaces of the sphere”, *Indiana Univ. Math. J.* **59**:3 (2010), 823–837. [MR](#) [Zbl](#)
- [Wang and Xia 2019] G. Wang and C. Xia, “Uniqueness of stable capillary hypersurfaces in a ball”, *Math. Ann.* **374**:3-4 (2019), 1845–1882. [MR](#) [Zbl](#)

Received March 25, 2018. Revised March 25, 2019.

MARCOS P. CAVALCANTE
UNIVERSIDADE FEDERAL DE ALAGOAS
MACEIÓ, AL
BRAZIL
marcos@pos.mat.ufal.br

DARLAN F. DE OLIVEIRA
UNIVERSIDADE ESTADUAL DE FEIRA DE SANTANA
FEIRA DE SANTANA, BA
BRAZIL
darlanfdeoliveira@gmail.com

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic.
Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY



mathematical sciences publishers

nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 305 No. 1 March 2020

The Poincaré homology sphere, lens space surgeries, and some knots with tunnel number two	1
KENNETH L. BAKER	
Fusion systems of blocks of finite groups over arbitrary fields	29
ROBERT BOLTJE, ÇISIL KARAGÜZEL and DENİZ YILMAZ	
Torsion points and Galois representations on CM elliptic curves	43
ABBEY BOURDON and PETE L. CLARK	
Stability of the positive mass theorem for axisymmetric manifolds	89
EDWARD T. BRYDEN	
Index estimates for free boundary constant mean curvature surfaces	153
MARCOS P. CAVALCANTE and DARLAN F. DE OLIVEIRA	
A criterion for modules over Gorenstein local rings to have rational Poincaré series	165
ANJAN GUPTA	
Generalized Cartan matrices arising from new derivation Lie algebras of isolated hypersurface singularities	189
NAVEED HUSSAIN, STEPHEN S.-T. YAU and HUAIQING ZUO	
On the commutativity of coset pressure	219
BING LI and WEN-CHIAO CHENG	
Signature invariants related to the unknotting number	229
CHARLES LIVINGSTON	
The global well-posedness and scattering for the 5-dimensional defocusing conformal invariant NLW with radial initial data in a critical Besov space	251
CHANGXING MIAO, JIANWEI YANG and TENGFEI ZHAO	
Liouville-type theorems for weighted p -harmonic 1-forms and weighted p -harmonic maps	291
KEOMKYO SEO and GABJIN YUN	
Remarks on the Hölder-continuity of solutions to parabolic equations with conic singularities	311
YUANQI WANG	
Deformation of Milnor algebras	329
ZHENJIAN WANG	
Preservation of log-Sobolev inequalities under some Hamiltonian flows	339
BO XIA	
Ground state solutions of polyharmonic equations with potentials of positive low bound	353
CAIFENG ZHANG, JUNGANG LI and LU CHEN	