

*Pacific  
Journal of  
Mathematics*

THE GLOBAL WELL-POSEDNESS AND SCATTERING FOR  
THE 5-DIMENSIONAL DEFOCUSING CONFORMAL  
INVARIANT NLW WITH RADIAL INITIAL DATA IN  
A CRITICAL BESOV SPACE

CHANGXING MIAO, JIANWEI YANG AND TENGFEI ZHAO



**THE GLOBAL WELL-POSEDNESS AND SCATTERING FOR  
THE 5-DIMENSIONAL DEFOCUSING CONFORMAL  
INVARIANT NLW WITH RADIAL INITIAL DATA IN  
A CRITICAL BESOV SPACE**

CHANGXING MIAO, JIANWEI YANG AND TENGFEI ZHAO

We obtain the global well-posedness and scattering for the radial solution to the defocusing conformal invariant nonlinear wave equation with initial data in the critical Besov space  $\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)$ . This is the 5-dimensional analogue of Dodson's result (2019), which was the first on the global well-posedness and scattering of the energy subcritical nonlinear wave equation without the uniform boundedness assumption on the critical Sobolev norms employed as a substitute of the missing conservation law with respect to the scaling invariance of the equation. The proof is based on exploiting the structure of the radial solution, developing the Strichartz-type estimates and incorporation of Dodson's strategy (2019), where we also avoid a logarithm-type loss by employing the inhomogeneous Strichartz estimates.

## 1. Introduction

We consider the solutions  $u$  to

$$(1-1) \quad \begin{cases} \partial_{tt}u - \Delta u + \mu|u|^p u = 0, & (t, x) \in \mathbb{R} \times \mathbb{R}^d, \\ (u(0), \partial_t u(0)) = (u_0, u_1), & x \in \mathbb{R}^d, \end{cases}$$

where  $\mu = \pm 1$ ,  $d \geq 1$ , and  $p > 0$ . If  $\mu = 1$ , (1-1) is described as defocusing, otherwise focusing. There is a natural scaling symmetry for (1-1), i.e., if we let  $u_\lambda(t, x) = \lambda^{2/p}u(\lambda t, \lambda x)$  for  $\lambda > 0$ , then  $u_\lambda$  is also a solution to (1-1) with initial data  $(\lambda^{2/p}u_0(\lambda x), \lambda^{(2/p)+1}u_1(\lambda x))$  preserving the  $\dot{H}^{s_p} \times \dot{H}^{s_p-1}(\mathbb{R}^d)$  norm of the initial data, where we define the critical regularity as  $s_p = \frac{d}{2} - \frac{2}{p}$ . At least, the solutions to (1-1) formally conserve the energy

$$(1-2) \quad E(u(t), \partial_t u(t)) := \frac{1}{2} \int_{\mathbb{R}^5} |\nabla_x u(t)|^2 dx + \frac{1}{2} \int_{\mathbb{R}^5} |\partial_t u(t)|^2 dx + \frac{\mu}{p+2} \int_{\mathbb{R}^5} |u(t)|^{p+2} dx,$$

*MSC2010:* primary 35B40, 35L05; secondary 35Q40.

*Keywords:* nonlinear wave equation, Strichartz estimates, scattering, hyperbolic coordinates, Morawetz estimates.

which is also invariant under the scaling if  $s_p = 1$ . In view of this, we say the Cauchy problem (1-1) is energy critical when  $s_p = 1$ , subcritical for  $s_p < 1$  and supercritical when  $s_p > 1$ .

Lindblad and Sogge [1995] proved the local theory of the Cauchy problem (1-1) in the minimal regularity spaces. In fact, if  $d \geq 2$  and  $p \geq (d+3)/(d-1)$ , the Cauchy problem (1-1) with initial data in the critical spaces  $\dot{H}^{s_p} \times \dot{H}^{s_p-1}(\mathbb{R}^d)$  is locally well-posed. The global theory for the Cauchy problem (1-1) with  $\mu = 1$  and  $s_p \leq 1$  has been studied extensively. While for the focusing case, even the solution with smooth initial data may blow up at finite time. For more related results see [Sogge 1995].

We will consider global existence and scattering of the solutions to (1-1). In general, a solution  $u$  is said to be scattering if it is a global solution and approaches a linear solution as  $t \rightarrow \pm\infty$ . In the cases of  $d \geq 2$  and  $p \geq (d+3)/(d-1)$ , the solution to (1-1) with small initial datum in the critical Sobolev spaces is globally well-posed and scattering; see [Lindblad and Sogge 1995].

For the defocusing energy critical wave equation (1-1), Grillakis [1990] first established the global existence theory for classical solution when  $d = 3$ . The results for other dimensions are proved in [Grillakis 1992; Shatah and Struwe 1993]. Scattering results for large energy data are proved in [Bahouri and Gérard 1999; Bahouri and Shatah 1998; Nakanishi 1999] by establishing variants of the Morawetz estimates [1968]

$$(1-3) \quad \iint_{\mathbb{R}^{1+d}} \frac{|u|^{\frac{2d}{d-2}}}{|x|} dx dt \leq C_d E(u_0, u_1),$$

where  $C_d$  is a constant depending on  $d$ . While in focusing energy critical cases, the Morawetz estimates (1-3) fails. The scattering results do not hold in general, since (1-1) has a ground state

$$W(x) = \left(1 + \frac{|x|^2}{d(d-2)}\right)^{-\frac{d-2}{2}}.$$

In the cases of  $3 \leq d \leq 5$ , Kenig and Merle [2008] proved the scattering result for solution with initial data such that  $E(u_0, u_1) < E(W, 0)$  and  $\|u_0\|_{\dot{H}^1(\mathbb{R}^d)} < \|W\|_{\dot{H}^1(\mathbb{R}^d)}$ . In their proofs, the main ingredient is the concentration compactness/rigidity theorem method introduced by [Kenig and Merle 2006]. This method is powerful and plays an important role in study of many other nonlinear dispersive equations. We refer to [Killip and Vişan 2013; Koch et al. 2014; Kenig 2015].

For the defocusing subcritical equation (1-1), the global existence has been proved for solution with initial data in the energy space  $\dot{H}^1 \times L^2(\mathbb{R}^d)$  by Ginibre and Velo [1985; 1989]. However, there are no scattering results even for solutions with initial datum in  $(\dot{H}^1 \cap \dot{H}^{s_p}) \times (L^2 \cap \dot{H}^{s_p-1})(\mathbb{R}^d)$ .

Recently, Dodson [2019] proved scattering results for the defocusing cubic wave equation with the initial datum belonging to the space  $\dot{B}_{1,1}^2 \times \dot{B}_{1,1}^1(\mathbb{R}^3)$ , which is a subspace of  $\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^3)$ . We remark that this is the first work that gives scattering results for large data in the critical Sobolev space without any a priori bound on the critical norm of the solution. Dodson's strategy consists of three steps:

- (1) By establishing some new Strichartz-type estimates, one can show that the solution is in the energy space  $\dot{H}^1 \times L^2(\mathbb{R}^3)$  up to some free evolutions. Then this decomposition enables one to prove the global well-posedness of the solution.
- (2) To obtain the scattering result, a conformal transformation is applied to show that the energy part of the solution has finite energy in hyperbolic coordinates. Then from the conformal invariance of the equation and a Morawetz-type inequality, one can deduce that  $\|u\|_{L_{t,x}^4(\mathbb{R} \times \mathbb{R}^5)} \leq C(\|(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)}, \delta_1)$ , where the parameter  $\delta_1$  relies on the scaling and spatial profiles of the initial data.
- (3) Finally, one can remove the dependence of  $\delta_1$  by employing the profile decomposition, which completes the proof.

Let  $S(t)(f, g)$  be the solution of Cauchy problem to the free wave equation

$$(1-4) \quad \begin{cases} \partial_{tt} v - \Delta v = 0, & (t, x) \in \mathbb{R} \times \mathbb{R}^5, \\ (v, \partial_t v)|_{t=0} = (f(x), g(x)), & x \in \mathbb{R}^5. \end{cases}$$

For the sake of statement, we introduce the following notation as

$$\dot{S}(t)(f, g) \triangleq \partial_t S(t)(f, g), \quad \text{and} \quad \vec{S}(t)(f, g) \triangleq (S(t)(f, g), \dot{S}(t)(f, g)).$$

We consider the Cauchy problem of nonlinear wave equation

$$(1-5) \quad \begin{cases} \partial_{tt} u - \Delta u + |u|u = 0, & (t, x) \in \mathbb{R} \times \mathbb{R}^d, \\ (u(0), \partial_t u(0)) = (u_0, u_1), & x \in \mathbb{R}^d, \end{cases}$$

Our main result can be stated as:

**Theorem 1.1.** *For any radial initial data  $(u_0, u_1) \in \dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)$ , the solution  $u$  to (1-5) is globally well-posed and scattering, i.e., there exists  $(u_0^\pm, u_1^\pm) \in \dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)$  such that*

$$(1-6) \quad \lim_{t \rightarrow \pm\infty} \|(u(t), \partial_t u(t)) - \vec{S}(t)(u_0^\pm, u_1^\pm)\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)} \rightarrow 0.$$

Furthermore, there is a function  $A : [0, \infty) \rightarrow [0, \infty)$ , such that

$$(1-7) \quad \|u\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} \leq A(\|(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)}).$$

**Remark 1.2.** (1) This theorem extends the results of [Dodson 2019] to the 5-dimension case. The proof will utilize the strategy given in [Dodson 2019], but it is highly nontrivial.

(2) Unlike the 3-dimensional case, the dispersive estimate (see (2-20)) gives a decay in time of order  $-2$ , which may cause a logarithmic failure when one estimates

$$\|u\|_{L_t^{4/3} L_x^4(J \times \mathbb{R}^5)} + \sup_{t \in J} (t^{3/4} \|u\|_{L_x^4(\mathbb{R}^5)}),$$

where  $0 \in J$  is a local time interval. We circumvent this difficulty by using the inhomogeneous Strichartz estimates in [Taggart 2010] and prove the global well-posedness of  $u$ .

(3) For the scattering result, by reductions, we need to bound the  $L_{t,x}^3$  of  $w$  on the light cone  $\{|x| \leq t + \frac{1}{2}\}$ . We will define the hyperbolic energy by rewriting (1-5) as the form

$$\partial_{tt}(r^2 w) - \partial_{rr}(r^2 w) = -2w - r^2 |w|w.$$

Observing that the additional term  $2w$  and the nonlinear term  $r^2 |w|w$  enjoy the same sign, we can bound the  $L_{t,x}^3$  norm of  $w$  by applying a Morawetz-type inequality, if we assume the hyperbolic energy of  $w$  is bounded.

(4) To certify the above assumption, we will make full use of (2-19) for radial solution and the sharp Hardy inequality. In contrast to the 3-dimensional case, some terms in (2-19) seem more difficult to dealt with. However, the integration domains of these terms are symmetric about the radius  $r$ , which is also consistent with the Huygens principle. This fact allows us apply the Hardy–Littlewood maximal functions to verify the assumption.

Now, we give the outline of the proof. By the Strichartz estimates and a standard fix point argument, for initial data  $(u_0, u_1)$ , there exists a maximal time interval  $I \subset \mathbb{R}$  such that there exists a unique solution  $u$  (see Definition 2.9 in Section 2) to (1-5) on  $I \times \mathbb{R}^5$ . We consider the global well-posedness by developing some Strichartz-type estimates (3-30). Utilizing the standard blowup criterion, we can show the global well-posedness of  $u$ .

Next, we claim the following proposition:

**Proposition 1.3.** *For every radial initial data  $(u_0, u_1) \in \dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)$ , let  $u$  be the corresponding solution to (1-5). Then there exists a parameter  $\delta_1$  depending the initial data  $(u_0, u_1)$  and a function  $A : [0, \infty)^2 \rightarrow [0, \infty)$  such that*

$$(1-8) \quad \|u\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} \leq A(\|(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)}, \delta_1).$$

We prove Theorem 1.1 by employing this and establishing Proposition 4.2, where the proof provides an alternate proof of Lemma 6.2 in [Dodson 2019].

Finally, we need to prove Proposition 1.3. From the partition  $u = v + w$ , it suffices to show the boundedness of  $L_{t,x}^3$  norm of  $w$ . We prove the hyperbolic energy of  $w$  is uniformly bounded. Then, a Morawetz-type inequality yields that the  $L_{t,x}^3$  norm of  $w$  is bounded in the cone, which finishes the proof.

This paper is organized as follows: Section 2 gives some tools from harmonic analysis and basic properties for the wave equation. In Section 3 we give the decomposition of  $u$  and prove its global well-posedness. The existence of the function  $A$  in (1-7) is shown in Section 4 based on the Proposition 1.3. Finally, in Section 5, we complete the proof by showing Proposition 1.3.

We end the introduction with some notations used throughout this paper. We use  $\mathcal{S}(\mathbb{R}^d)$  to denote the space of Schwartz functions on  $\mathbb{R}^d$ . For  $1 \leq p \leq \infty$ , we define  $L^p(\mathbb{R}^d)$  by the spaces of Lebesgue measurable functions with finite  $L^p(\mathbb{R}^d)$ -norm, which is defined by

$$\|f\|_{L^p(\mathbb{R}^d)} = \left( \int_{\mathbb{R}^d} |f(x)|^p dx \right)^{1/p}, \quad \text{for } 1 \leq p < \infty,$$

and  $\|f\|_{L^\infty(\mathbb{R}^d)} = \text{ess sup}_{x \in \mathbb{R}^d} |f(x)|$ . We let  $\ell^p$  be the spaces of complex number sequences  $\{a_n\}_{n \in \mathbb{Z}}$  such that  $\{a_n\}_{n \in \mathbb{Z}} \in \ell^p$  if and only if

$$\|\{a_n\}\|_{\ell_n^p(\mathbb{Z})} \triangleq \left( \sum_n |a_n|^p \right)^{1/p} < \infty, \quad \text{for } 1 \leq p < \infty$$

and  $\|\{a_n\}\|_{\ell_n^\infty(\mathbb{Z})} := \sup_n |a_n| < \infty$ . We use  $X \lesssim Y$  to mean that there exists a constant  $C > 1$  such that  $X \leq CY$ , where the dependence of  $C$  on the parameters will be clear from the context. We use  $X \sim Y$  to denote  $X \lesssim Y$  and  $Y \lesssim X$ .  $A \ll B$  denotes there is a sufficiently large number  $C$  such that  $A \leq C^{-1}B$ .

## 2. Basic tools and some elementary properties for the wave equation

In this section, we recall some tools from harmonic analysis and useful results for the wave equation.

**2A. Some tools from harmonic analysis.** Recall the Fourier transform of  $f \in \mathcal{S}(\mathbb{R}^d)$  is defined by

$$\widehat{f}(\xi) = (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(x) e^{-ix\xi} dx,$$

which can be extended to Schwartz distributions naturally. We will make frequent use of the Littlewood–Paley projection operators. Specifically, we let  $\varphi$  be a radial smooth function supported on the ball  $|\xi| \leq 2$  and equal to 1 on the ball  $|\xi| \leq 1$ . For  $j \in \mathbb{Z}$ , we define the Littlewood–Paley projection operators by

$$\begin{aligned} \widehat{P_{\leq j} f}(\xi) &:= \varphi(\xi/2^j) \widehat{f}(\xi), \\ \widehat{P_{>j} f}(\xi) &:= (1 - \varphi(\xi/2^j)) \widehat{f}(\xi), \\ \widehat{P_j f}(\xi) &:= (\varphi(\xi/2^j) - \varphi(\xi/2^{j-1})) \widehat{f}(\xi). \end{aligned}$$

The Littlewood–Paley operators commute with derivative operators and are bounded on the general Sobolev spaces. These operators also obey the following standard Bernstein estimates:

**Lemma 2.1** (Bernstein estimates). *For  $1 \leq r \leq q \leq \infty$  and  $s \geq 0$ ,*

$$\begin{aligned} \|\nabla^{\pm s} P_j f\|_{L_x^r(\mathbb{R}^d)} &\sim 2^{\pm js} \|P_j f\|_{L_x^r(\mathbb{R}^d)}, \\ \|\nabla^s P_{\leq j} f\|_{L_x^r(\mathbb{R}^d)} &\lesssim 2^{js} \|P_{\leq j} f\|_{L_x^r(\mathbb{R}^d)}, \\ \|P_{>j} f\|_{L_x^r(\mathbb{R}^d)} &\lesssim 2^{-js} \|\nabla^s P_{>j} f\|_{L_x^r(\mathbb{R}^d)}, \\ \|P_{\leq j} f\|_{L^q(\mathbb{R}^d)} &\lesssim 2^{(d/r-d/q)j} \|P_{\leq j} f\|_{L_x^r(\mathbb{R}^d)}, \end{aligned}$$

where the fractional derivative operator  $|\nabla|^\sigma$  is defined by  $\widehat{|\nabla|^\sigma f}(\xi) = |\xi|^\sigma \widehat{f}(\xi)$ , for  $\sigma \in \mathbb{R}$ .

**Definition 2.2** (homogeneous Besov spaces). Let  $s$  be a real number and let  $1 \leq p, r \leq \infty$ . We denote the homogeneous Besov norm by

$$(2-1) \quad \|f\|_{\dot{B}_{p,r}^s(\mathbb{R}^d)} := \left\| \{2^{js} \|P_j f\|_{L^p(\mathbb{R}^d)} \} \right\|_{\ell_j^r(\mathbb{Z})},$$

for  $f \in \mathcal{S}(\mathbb{R}^d)$ . Then the Besov space  $\dot{B}_{p,r}^s(\mathbb{R}^d)$  is the completion of the Schwartz function under this norm.

We shall give the following radial Sobolev-type inequalities, which are analogous to the 3-dimensional cases established in [Dodson 2019]. We denote radial derivative by  $\partial_r f(x) = \left(\frac{x}{|x|} \cdot \nabla\right) f(x)$  for any function  $f$  defined on  $\mathbb{R}^5$ .

**Lemma 2.3** (radial Sobolev-type inequalities in Besov spaces). *For any radial function  $f \in \mathcal{S}(\mathbb{R}^5)$ , we have*

$$(2-2) \quad \||x|^2 f\|_{L^\infty(\mathbb{R}^5)} \lesssim \|f\|_{\dot{B}_{2,1}^{1/2}(\mathbb{R}^5)}.$$

Let  $(u_0, u_1) \in \dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)$  be a radial function; then we have

$$\begin{aligned} (2-3) \quad & \left\| \frac{1}{|x|^2} \partial_r u_0(x) \right\|_{L_x^1(\mathbb{R}^5)} + \left\| \frac{1}{|x|} \partial_{rr} u_0(x) \right\|_{L_x^1(\mathbb{R}^5)} \\ & + \left\| \frac{1}{|x|^3} u_0(x) \right\|_{L_x^1(\mathbb{R}^5)} + \||x|^3 \partial_r u_0\|_{L_x^\infty(\mathbb{R}^5)} \lesssim \|u_0\|_{\dot{B}_{1,1}^3(\mathbb{R}^5)}, \end{aligned}$$

$$(2-4) \quad \left\| \frac{1}{|x|} \partial_r u_1(x) \right\|_{L_x^1(\mathbb{R}^5)} + \left\| \frac{1}{|x|^2} u_1(x) \right\|_{L_x^1(\mathbb{R}^5)} + \||x|^3 u_1(x)\|_{L_x^\infty(\mathbb{R}^5)} \lesssim \|u_1\|_{\dot{B}_{1,1}^2(\mathbb{R}^5)}.$$

*Proof.* We first consider (2-2). Since  $f$  is radial, using polar coordinates, we have

$$(2-5) \quad \begin{aligned} P_j f(|x|) &= P_j f(x) = \int_{\mathbb{R}^5} \widehat{P_j f}(\xi) e^{ix\xi} d\xi \\ &= \int_0^\infty \int_{\mathbb{S}^4} \widehat{P_j f}(r) r^4 e^{irx\omega} d\sigma(\omega) dr. \end{aligned}$$

Recall the decay estimates of Fourier transform of the surface measure on the sphere

$$d\widehat{\sigma}_{\mathbb{S}^4}(\xi) \leq C(1 + |\xi|)^{-2},$$

which, with Hölder's inequality, yields

$$(2-6) \quad |P_j f(|x|)| \lesssim \int_0^\infty |\widehat{P_j f}(r)| r^2 |x|^{-2} dr \lesssim |x|^{-2} 2^{\frac{1}{2}j} \|P_j f\|_{L^2}.$$

Then the inequality (2-2) follows from (2-6) and the definition of the Besov space.

Next, we consider (2-3) and (2-4). By the density of Schwartz functions in  $\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)$ , we may assume that  $u_0, u_1 \in \mathcal{S}(\mathbb{R}^5)$ . We claim it suffices to show

$$(2-7) \quad \begin{aligned} \left\| \frac{1}{|x|^2} \partial_r u_0(x) \right\|_{L_x^1(\mathbb{R}^5)} + \left\| \frac{1}{|x|} |\Delta u_0(x)| \right\|_{L_x^1(\mathbb{R}^5)} + \left\| \frac{1}{|x|^3} u_0(x) \right\|_{L_x^1(\mathbb{R}^5)} \\ \lesssim \|u_0\|_{\dot{B}_{1,1}^3(\mathbb{R}^5)}, \end{aligned}$$

$$(2-8) \quad \left\| \frac{1}{|x|} \partial_r u_1(x) \right\|_{L_x^1(\mathbb{R}^5)} + \left\| \frac{1}{|x|^2} u_1(x) \right\|_{L_x^1(\mathbb{R}^5)} \lesssim \|u_1\|_{\dot{B}_{1,1}^2(\mathbb{R}^5)}.$$

To see this, by using the fact  $\Delta f = \partial_{rr} f + \frac{4}{r} \partial_r f$  for radial function  $f(x)$  on  $\mathbb{R}^5$ , we have

$$(2-9) \quad \begin{aligned} \left\| \frac{1}{|x|} \partial_{rr} u_0(x) \right\|_{L_x^1(\mathbb{R}^5)} &\lesssim \left\| \frac{1}{|x|^2} \partial_r u_0(x) \right\|_{L_x^1(\mathbb{R}^5)} + \left\| \frac{1}{|x|} |\Delta u_0(x)| \right\|_{L_x^1(\mathbb{R}^5)} \\ &\lesssim \|u_0\|_{\dot{B}_{1,1}^3(\mathbb{R}^5)} \end{aligned}$$

From the fundamental theorem of calculus and polar coordinates, for  $y \in \mathbb{R}^5 \setminus \{0\}$ ,

$$(2-10) \quad \begin{aligned} |y|^3 |\partial_r u_0(y)| &\lesssim \int_{|y|}^\infty \int_{\mathbb{S}^4} r^3 |\partial_{rr} u_0(r)| d\sigma(\omega) dr \\ &\lesssim \left\| \frac{1}{|x|} \partial_{rr} u_0(x) \right\|_{L_x^1(\mathbb{R}^5)} \lesssim \|u_0\|_{\dot{B}_{1,1}^3(\mathbb{R}^5)} \end{aligned}$$

and

$$(2-11) \quad \begin{aligned} |y|^3 |u_1(y)| &\lesssim \int_{|y|}^\infty \int_{\mathbb{S}^4} r^3 |\partial_{rr} u_1(r)| d\sigma(\omega) dr \\ &\lesssim \left\| \frac{1}{|x|} \partial_r u_1(x) \right\|_{L_x^1(\mathbb{R}^5)} \lesssim \|u_1\|_{\dot{B}_{1,1}^2(\mathbb{R}^5)}. \end{aligned}$$

Hence, we are reduced to proving (2-7) and (2-8). We just give the estimate for the first term on the left-hand side of (2-7), since others can be handled similarly. For  $j \in \mathbb{Z}$ , utilizing Bernstein's estimates and polar coordinates, we obtain

$$\begin{aligned}
(2-12) \quad & \left\| \frac{1}{|x|^2} \partial_r P_j u_0(x) \right\|_{L_x^1(\mathbb{R}^5)} \lesssim \int_0^\infty \int_{\mathbb{S}^4} r^2 \partial_r P_j u_0(r) d\sigma(\omega) dr \\
& \lesssim \int_0^{2^{-j}} \frac{1}{r^2} |\partial_r(P_j u_0)| r^4 dr + \int_{2^{-j}}^\infty \frac{1}{r^2} |\partial_r(P_j u_0)| r^4 dr \\
& \lesssim 2^{-3j} \|\partial_r(P_j u_0)\|_{L_r^\infty(\mathbb{R}_+)} + 2^{2j} \|\partial_r(P_j u_0)\|_{L_x^1(\mathbb{R}^5)} \\
& \lesssim 2^{3j} \|P_j u_0\|_{L_x^1(\mathbb{R}^5)}.
\end{aligned}$$

Thus, we have  $\|(1/|x|^2) \partial_r u_0(x)\|_{L_x^1(\mathbb{R}^5)} \lesssim \|u_0\|_{\dot{B}_{1,1}^3(\mathbb{R}^5)}$ .  $\square$

As a direct consequence of Lemma 2.3, we have

$$\begin{aligned}
(2-13) \quad & \| |x|^{1/2} \partial_r u_0(x) \|_{L_x^2(\mathbb{R}^5)} + \| |x|^{-1/2} u_0(x) \|_{L_x^2(\mathbb{R}^5)} + \| |x|^{1/2} u_1(x) \|_{L_x^2(\mathbb{R}^5)} \\
& \lesssim \| (u_0, u_1) \|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)}.
\end{aligned}$$

**Lemma 2.4.** Suppose  $\chi(x) \in C_c^\infty(\mathbb{R}^5)$ . Let  $R = 2^k$  be a dyadic number for  $k \in \mathbb{Z}$  and denote  $\chi_R(x) = \chi(\frac{x}{R})$ . Then we have

$$(2-14) \quad \|\chi_R(x) f\|_{\dot{B}_{2,1}^{1/2}(\mathbb{R}^5)} \lesssim \|f\|_{\dot{B}_{2,1}^{1/2}(\mathbb{R}^5)},$$

$$(2-15) \quad \|\chi_R(x) g\|_{\dot{B}_{2,1}^{-1/2}(\mathbb{R}^5)} \lesssim \|g\|_{\dot{B}_{2,1}^{-1/2}(\mathbb{R}^5)},$$

where the bound is independent of  $R$ . Furthermore, if  $\chi(x) = 1$  on  $|x| \leq 1$ , then for  $(f, g) \in \dot{B}_{2,1}^{1/2} \times \dot{B}_{2,1}^{-1/2}(\mathbb{R}^5)$ , we have

$$(2-16) \quad \lim_{R \rightarrow \infty} \|(1 - \chi_R(x)) f\|_{\dot{B}_{2,1}^{1/2}(\mathbb{R}^5)} + \|(1 - \chi_R(x)) g\|_{\dot{B}_{2,1}^{-1/2}(\mathbb{R}^5)} = 0.$$

*Proof.* By scaling, to prove the inequalities (2-14) and (2-15), it suffices to prove the cases for  $R = 1$ , which follows from a similar proof of Lemma 2.2 in [Dodson 2019]. On the other hand, (2-16) follows from (2-14), (2-15), and the fact that  $C_c^\infty \times C_c^\infty(\mathbb{R}^5)$  is dense in  $\dot{B}_{2,1}^{1/2} \times \dot{B}_{2,1}^{-1/2}(\mathbb{R}^5)$ .  $\square$

Finally, we need the following chain rule estimates for later use.

**Lemma 2.5** ( $C^1$ -fractional chain rule [Christ and Weinstein 1991]). Suppose  $G \in C^1(\mathbb{C})$ ,  $s \in (0, 1]$ , and  $1 < q, q_1, q_2 < \infty$  satisfying  $\frac{1}{q} = \frac{1}{q_1} + \frac{1}{q_2}$ . Then

$$(2-17) \quad \|\nabla^s G(u)\|_{L^q(\mathbb{R}^d)} \lesssim \|G'(u)\|_{L^{q_1}(\mathbb{R}^d)} \|\nabla^s u\|_{L^{q_2}(\mathbb{R}^d)}.$$

**2B. Fundamental properties of the wave equations.** Throughout the paper, by abuse of notations, we often write  $u(t) = u(t, x)$  for simplicity and  $u(t, r) = u(t, x)$  when  $u(t, \cdot)$  is radially symmetric.

Recall the explicit formula for solution to the linear wave equation in 5 dimensions,

$$(2-18) \quad \begin{aligned} S(t)(f, g)(x) &= \cos(t|\nabla|)f(x) + \frac{\sin(t|\nabla|)}{|\nabla|}g(x) \\ &= \frac{1}{3\omega_5} \partial_t \left[ \frac{1}{t} \partial_t \right] \left( t^3 \int_{|y|=1} f(x+ty) d\sigma(y) \right) \\ &\quad + \frac{1}{3\omega_5} \frac{1}{t} \partial_t \left( t^3 \int_{|y|=1} g(x+ty) d\sigma(y) \right), \end{aligned}$$

where  $\omega_5$  is the surface area of the unit sphere in  $\mathbb{R}^5$ . When  $(f, g)$  is radially symmetric, for  $t > 0$ , (2-18) can be rewritten as

$$(2-19) \quad \begin{aligned} S(t)(f, g)(r) &= \frac{1}{2r^2} [(r-t)^2 f(r-t) + (r+t)^2 f(r+t)] \\ &\quad - \frac{t}{2r^3} \int_{|r-t|}^{r+t} s f(s) ds + \frac{1}{4r^3} \int_{|r-t|}^{r+t} s(s^2 + r^2 - t^2) g(s) ds. \end{aligned}$$

See also [Rammaha 1987; Lindblad and Sogge 1996; Colzani et al. 2002] for the radial solutions to general dimensions linear wave equation. From the explicit formula (2-18), we can obtain the following dispersive estimate.

**Proposition 2.6** (dispersive estimate).

$$(2-20) \quad \|S(t)(u_0, u_1)\|_{L^\infty(\mathbb{R}^5)} \lesssim \frac{1}{t^2} [\|\nabla^3 u_0\|_{L^1(\mathbb{R}^5)} + \|\nabla^2 u_1\|_{L^1(\mathbb{R}^5)}].$$

*Proof.* We give the proof for completeness. A similar proof for the 3-dimensional case can be found in [Killip and Visan 2011]. By (2-18), the free solution  $S(t)(u_0, u_1)$  can be rewritten as

$$(2-21) \quad \begin{aligned} &\frac{1}{\omega_5} \int_{|y|=1} u_0(x+ty) d\sigma(y) + \frac{5t}{3\omega_5} \int_{|y|=1} y(\nabla u_0)(x+ty) d\sigma(y) \\ &+ \frac{t^2}{3\omega_5} \int_{|y|=1} y(\nabla^2 u_0)(x+ty) y d\sigma(y) + \frac{t}{\omega_5} \int_{|y|=1} u_1(x+ty) d\sigma(y) \\ &+ \frac{t^2}{3\omega_5} \int_{|y|=1} y(\nabla u_1)(x+ty) d\sigma(y), \end{aligned}$$

which, with the fundamental theorem of calculus, yields (2-20). For instance, using

polar coordinates, we can estimate the first term of (2-21) as

$$\begin{aligned}
(2-22) \quad & \left| \frac{1}{\omega_5} \int_{|y|=1} u_0(x+ty) d\sigma(y) \right| \\
&= \left| \frac{1}{\omega_5} \int_t^\infty \int_s^\infty \int_\tau^\infty \int_{|y|=1} \frac{d^3}{d\rho^3} [u_0(x+\rho y)] d\sigma(y) d\rho d\tau ds \right| \\
&\lesssim \int_t^\infty \int_s^\infty \int_\tau^\infty \int_{|y|=1} |\nabla^3 u_0|(x+\rho y) d\sigma(y) d\rho d\tau ds \\
&\lesssim \int_t^\infty \int_s^\infty \frac{1}{\tau^4} d\tau ds \|\nabla^3 u_0\|_{L^1(\mathbb{R}^5)} \\
&\lesssim \frac{1}{t^2} \|\nabla^3 u_0\|_{L^1(\mathbb{R}^5)}.
\end{aligned}$$

The other terms can be dealt with similarly.  $\square$

We recall the Strichartz estimates of the wave equation in  $\mathbb{R}^5$ . Let  $I \subset \mathbb{R}$  be an interval. We denote the spacetime norm  $L_t^q W_x^{s,r}(I \times \mathbb{R}^5)$  of a function  $u(t, x)$  on  $I \times \mathbb{R}^5$  by

$$\|u\|_{L_t^q W_x^{s,r}(I \times \mathbb{R}^5)} := \left\| \|u(t, x)\|_{W_x^{s,r}(\mathbb{R}^5)} \right\|_{L_t^q(I)},$$

for  $s \in \mathbb{R}$ ,  $1 \leq q, r \leq \infty$ . We denote that a pair  $(q, r)$  of exponents is admissible, if

$$(2-23) \quad 2 \leq q \leq \infty, \quad 2 \leq r < \infty, \quad \text{and} \quad \frac{1}{q} + \frac{2}{r} \leq 1.$$

Moreover, we say  $(q, r)$  is wave acceptable, provided

$$(2-24) \quad 1 \leq q < \infty, \quad 2 \leq r \leq \infty, \quad \frac{1}{q} < 4 \left( \frac{1}{2} - \frac{1}{r} \right),$$

or  $(q, r) = (\infty, 2)$ .

**Proposition 2.7** (Strichartz estimates [Lindblad and Sogge 1995; Ginibre and Velo 1995; Keel and Tao 1998]). *Let  $(u_0, u_1) \in \dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)$  and  $(q, r), (\tilde{q}, \tilde{r})$  be two admissible pairs. If  $u$  is a weak solution to the wave equation  $\partial_{tt}u - \Delta u = F(t, x)$  with initial data  $(u_0, u_1)$ , then we have*

$$\begin{aligned}
(2-25) \quad & \left\| |\nabla|^\rho u \right\|_{L_t^q L_x^r(I \times \mathbb{R}^5)} + \sup_{t \in I} \|(u, u_t)\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)} \\
&\lesssim \|(u_0, u_1)\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)} + \left\| |\nabla|^{-\mu} F \right\|_{L_t^{\tilde{q}'} L_x^{\tilde{r}'}(I \times \mathbb{R}^5)},
\end{aligned}$$

provided that

$$(2-26) \quad \rho = \frac{1}{q} + \frac{5}{r} - 2 \quad \text{and} \quad \mu = \frac{1}{\tilde{q}} + \frac{5}{\tilde{r}} - 2.$$

**Proposition 2.8** (inhomogeneous Strichartz estimates [Taggart 2010]). *Suppose that the exponent pairs  $(q_1, r_1)$  and  $(\tilde{q}_1, \tilde{r}_1)$  are wave acceptable, and satisfy the scaling condition*

$$\frac{1}{q} + \frac{1}{\tilde{q}} = 2 - 2\left(\frac{1}{r_1} + \frac{1}{\tilde{r}_1}\right)$$

and the conditions

$$\frac{1}{q} + \frac{1}{\tilde{q}} < 1, \quad \frac{1}{2} \leq \frac{\tilde{r}_1}{r_1} \leq 2.$$

Let  $r \geq r_1$ ,  $\tilde{r} \geq \tilde{r}_1$ ,  $\rho \in \mathbb{R}$  be such that

$$\rho + 5\left(\frac{1}{2} - \frac{1}{r}\right) - \frac{1}{q} = 1 - \left(\tilde{\rho} + 5\left(\frac{1}{2} - \frac{1}{\tilde{r}}\right) - \frac{1}{\tilde{q}}\right).$$

If  $F(t, x)$  is in  $L_t^{\tilde{q}'}(\mathbb{R}; \dot{B}_{\tilde{r}', 2}^{-\tilde{\rho}}(\mathbb{R}^5))$  and  $u$  is a weak solution to the inhomogeneous wave equation

$$(2-27) \quad -\partial_t^2 u + \Delta u = F(t, x), \quad u(0) = u_t(0) = 0,$$

then

$$(2-28) \quad \|u\|_{L_t^q(\mathbb{R}; \dot{B}_{r, 2}^\rho(\mathbb{R}^5))} \lesssim \|F(t, x)\|_{L_t^{\tilde{q}'}(\mathbb{R}; \dot{B}_{\tilde{r}', 2}^{-\tilde{\rho}}(\mathbb{R}^5))}.$$

Next, we recall the well-posedness theory and the perturbation theory of the Cauchy problem (1-5).

**Definition 2.9** (solution). Let  $I$  be a time interval such that  $0 \in I$ . We say function  $u : I \times \mathbb{R}^5 \rightarrow \mathbb{R}$  is a (strong) solution to the Cauchy problem (1-5) in  $I$  if it satisfies  $(u, u_t)(0) = (u_0, u_1)$ ,

$$(u, u_t) \in C(I; \dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)) \cap L_{t,x}^3(I \times \mathbb{R}^5),$$

and the integral equation

$$(2-29) \quad u(t) = S(t)(u_0, u_1) - \int_0^t S(t-\tau)(0, |u|u(\tau)) d\tau$$

for all  $t \in I$ .

**Theorem 2.10** (local well-posedness [Lindblad and Sogge 1995; Rodriguez 2017]). *Let  $(u_0, u_1) \in \dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)$  with*

$$\|(u_0, u_1)\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)} \leq A.$$

*There exists  $\delta = \delta(A) > 0$  such that, if*

$$(2-30) \quad \|S(t)(u_0, u_1)\|_{L_{t,x}^3([0, T] \times \mathbb{R}^5)} \leq \delta, \quad \text{for some } T > 0,$$

*then there exists a unique solution  $u$  to (1-5) in  $[0, T] \times \mathbb{R}^5$ , such that*

$$(2-31) \quad \sup_{0 \leq t \leq T} \|(u, u_t)\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)} + \|u\|_{L_{t,x}^3([0, T] \times \mathbb{R}^5)} \leq C(A).$$

In addition, if  $A > 0$  is small enough, we can take  $T = \infty$ .

We define  $T_+(u_0, u_1) := \sup I$ , where  $I$  is the maximal interval of existence of the solution  $u$ .

**Lemma 2.11** (standard blowup criterion). *Suppose  $u$  is the solution to the Cauchy problem (1-5) with initial data  $(u_0, u_1) \in \dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)$  and  $T_+(u_0, u_1) < \infty$ . Then we have*

$$(2-32) \quad \|u\|_{L_{t,x}^3([0, T_+(u_0, u_1)) \times \mathbb{R}^5)} = \infty.$$

The proof is standard and similar to the energy critical case in [Kenig 2015].

We end this section by recalling the stability lemma for the Cauchy problem (1-5), which plays an important role in the Theorem 4.1.

**Theorem 2.12** (perturbation theory [Rodriguez 2017]). *Let  $I \subset \mathbb{R}$  be a time interval with  $0 \in I$ . Let  $(u_0, u_1) \in \dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)$  and some constants  $M, A, A' > 0$  be given. Let  $\tilde{u}$  be defined on  $I \times \mathbb{R}^5$  and satisfy*

$$(2-33) \quad \sup_{t \in I} \|(\tilde{u}, \partial_t \tilde{u})\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)} \leq A,$$

$$(2-34) \quad \|\tilde{u}\|_{L_{t,x}^3(I \times \mathbb{R}^5)} \leq M.$$

Assume that  $\partial_{tt} \tilde{u} - \Delta \tilde{u} = -|\tilde{u}| \tilde{u} + e$  on  $I \times \mathbb{R}^5$ ,

$$(2-35) \quad \|(u_0 - \tilde{u}(0), u_1 - \partial_t \tilde{u}(0))\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)} \leq A'$$

and that

$$(2-36) \quad \|e\|_{L_{t,x}^{3/2}(I \times \mathbb{R}^5)} + \|S(t)[(\tilde{u}(0), \partial_t \tilde{u}(0)) - (u_0, u_1)]\|_{L_{t,x}^3(I \times \mathbb{R}^5)} < \varepsilon.$$

Then, there exist  $\beta > 0$  and  $\varepsilon_0 = \varepsilon_0(M, A, A') > 0$  such that, for  $0 < \varepsilon < \varepsilon_0$ , there exists a solution  $u$  to (1-5) in  $I$  such that  $(u(0), \partial_t u(0)) = (u_0, u_1)$ , with

$$(2-37) \quad \|u\|_{L_{t,x}^3(I \times \mathbb{R}^5)} \leq C(M, A, A'),$$

$$(2-38) \quad \sup_{t \in I} \|(\partial_t \tilde{u}(t), \tilde{u}(t)) - (u, \partial_t u(t))\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)} \leq C(M, A, A')(A' + \varepsilon^\beta).$$

### 3. Decomposition of the solution and global well-posedness

In this section, we will prove that for any given initial data  $(u_0, u_1) \in \dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)$ , the corresponding solution  $u$  to (1-5) is globally well-posed. To prove this, we first show the solution  $u$  belongs to some suitable Strichartz-type spaces on a local time interval. Then, we split it into two parts:  $u = v + w$ . Based on the inhomogeneous Strichartz estimates (2-28), we will derive a decay property for  $v$  and prove that  $w$  is in the energy space  $\dot{H}^1 \times L^2(\mathbb{R}^5)$ . We remark that the constants in “ $\lesssim$ ” in this section depend upon  $\|(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)}$ .

For the sake of simplicity, we denote  $F(u) = |u|u$ . Recall that  $u_\lambda$  is also a solution to the Cauchy problem (1-5) with initial data  $(\lambda^2 u_0(\lambda x), \lambda^3 u_1(\lambda x))$ , where

$$(3-1) \quad u_\lambda(t, x) = \lambda^2 u(\lambda t, \lambda x).$$

Given  $(u_0, u_1) \in \dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)$ , for any  $\eta > 0$ , there exists  $j_0 = j_0(u_0, u_1, \eta) < \infty$  such that

$$(3-2) \quad \sum_{j \geq j_0} 2^{2j} \|P_j u_0\|_{L^1(\mathbb{R}^5)} < \eta.$$

Replace  $u$  by  $u_\lambda$  for  $\lambda = 2^{-j_0}$ , then we have

$$(3-3) \quad \sum_{j \geq 0} 2^{3j} \|P_j u_0\|_{L^1(\mathbb{R}^5)} + \sum_{j \geq 0} 2^{2j} \|P_j u_1\|_{L^1(\mathbb{R}^5)} < \eta.$$

**Lemma 3.1.** *Let  $\epsilon_0 > 0$  be a small constant and  $\eta \ll \epsilon_0$ . If the initial data  $(u_0, u_1) \in \dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)$  satisfies (3-3) and  $u$  is the solution to (1-5) with initial data  $(u_0, u_1)$  given by Theorem 2.10, then there exists*

$$\delta = \delta(\epsilon_0, \|(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2}) > 0$$

such that

$$(3-4) \quad \|u\|_{L_{t,x}^3([-\delta, \delta] \times \mathbb{R}^5)} \leq \sum_{j \in \mathbb{Z}} \|P_j u\|_{L_{t,x}^3([-\delta, \delta] \times \mathbb{R}^5)} \lesssim \epsilon_0,$$

$$(3-5) \quad \|u\|_{L_t^\infty([-\delta, \delta]; \dot{B}_{2,1}^{1/2}(\mathbb{R}^5))} \lesssim \sum_{j \in \mathbb{Z}} \|P_j u\|_{L_t^\infty \dot{H}^{1/2}([-\delta, \delta] \times \mathbb{R}^5)} \lesssim 1.$$

*Proof.* By Strichartz's estimates in Proposition 2.7 and (3-3), we obtain

$$(3-6) \quad \|S(t) P_{\geq 0}(u_0, u_1)\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} \leq \frac{1}{2} \epsilon_0.$$

On the other hand, for every  $t \in \mathbb{R}$ , by Bernstein, we have

$$(3-7) \quad \|S(t) P_{\leq 0}(u_0, u_1)\|_{L_x^3(\mathbb{R}^5)} \lesssim 1.$$

Hence, taking  $\delta$  small enough, we have,

$$(3-8) \quad \|S(t)(u_0, u_1)\|_{L_{t,x}^3([-\delta, \delta] \times \mathbb{R}^5)} \leq \epsilon_0.$$

Then, by the Strichartz estimates, we have

$$(3-9) \quad \|u\|_{L_{t,x}^3([-\delta, \delta] \times \mathbb{R}^5)} \lesssim \|S(t)(u_0, u_1)\|_{L_{t,x}^3([-\delta, \delta] \times \mathbb{R}^5)} + \|u\|_{L_{t,x}^3([-\delta, \delta] \times \mathbb{R}^5)}^2,$$

from which, by a standard continuity argument, we deduce that

$$(3-10) \quad \|u\|_{L_{t,x}^3([-\delta, \delta] \times \mathbb{R}^5)} \lesssim \epsilon_0.$$

Let

$$(3-11) \quad a_k = \|P_k u\|_{L_{t,x}^3([-\delta, \delta] \times \mathbb{R}^5)} + 2^{\frac{1}{2}k} \|P_k u\|_{L_t^\infty L_x^2([-\delta, \delta] \times \mathbb{R}^5)} + 2^{\frac{1}{4}k} \|P_k u\|_{L_t^6 L_x^{12/5}([-\delta, \delta] \times \mathbb{R}^5)},$$

$$(3-12) \quad b_k = 2^{\frac{1}{2}k} \|P_k u_0\|_{L_x^2} + 2^{-\frac{1}{2}k} \|P_k u_1\|_{L_x^2}.$$

By Young's inequality, it suffices to show there is a recurrence relation

$$(3-13) \quad a_k \lesssim b_k + \epsilon_0 \sum_j 2^{-\frac{1}{4}|j-k|} a_j.$$

To prove this, making use of the Strichartz estimates, we have

$$(3-14) \quad a_k \lesssim b_k + 2^{\frac{1}{4}k} \|P_k F(u)\|_{L_t^2 L_x^{4/3}([0, \delta] \times \mathbb{R}^5)}.$$

First, we consider the low frequency part of the second term in the right-hand side of (3-14). By Lemma 2.5 and Hölder, we have

$$(3-15) \quad \begin{aligned} & \|P_k F(u_{\leq k})\|_{L_t^2 L_x^{4/3}([-\delta, \delta] \times \mathbb{R}^5)} \\ & \lesssim 2^{-\frac{1}{2}k} \|P_k |\nabla_x|^{\frac{1}{2}} F(u_{\leq k})\|_{L_t^2 L_x^{4/3}([-\delta, \delta] \times \mathbb{R}^5)} \\ & \lesssim 2^{-\frac{1}{2}k} \|u\|_{L_{t,x}^3([-\delta, \delta] \times \mathbb{R}^5)} \||\nabla|^{1/2} (P_{\leq k} u)\|_{L_t^6 L_x^{12/5}([-\delta, \delta] \times \mathbb{R}^5)} \\ & \lesssim \|u\|_{L_{t,x}^3([-\delta, \delta] \times \mathbb{R}^5)} \sum_{j \leq k} 2^{-\frac{1}{2}(k-j)} \|P_j u\|_{L_t^6 L_x^{12/5}([-\delta, \delta] \times \mathbb{R}^5)}, \end{aligned}$$

from which it follows that

$$(3-16) \quad \begin{aligned} 2^{\frac{k}{4}} \|P_k F(u_{\leq k})\|_{L_t^2 L_x^{4/3}([-\delta, \delta] \times \mathbb{R}^5)} & \lesssim \epsilon_0 \sum_{j \leq k} 2^{-\frac{1}{4}(k-j)} 2^{\frac{1}{4}j} \|P_j u\|_{L_t^6 L_x^{12/5}([-\delta, \delta] \times \mathbb{R}^5)} \\ & \lesssim \epsilon_0 \sum_{j \leq k} 2^{-\frac{1}{4}(k-j)} a_j. \end{aligned}$$

On the other hand, by Hölder's inequality,

$$(3-17) \quad \begin{aligned} & 2^{\frac{1}{4}k} \|P_k (F(u) - F(u_{\leq k}))\|_{L_t^2 L_x^{4/3}([-\delta, \delta] \times \mathbb{R}^5)} \\ & \lesssim \|u\|_{L_{t,x}^3([-\delta, \delta] \times \mathbb{R}^5)} 2^{\frac{1}{4}k} \|P_{\geq k+1} u\|_{L_t^6 L_x^{12/5}([-\delta, \delta] \times \mathbb{R}^5)} \\ & \lesssim \epsilon_0 \sum_{j \geq k+1} 2^{-\frac{1}{4}(j-k)} 2^{\frac{1}{4}j} \|P_j u\|_{L_t^6 L_x^{12/5}([-\delta, \delta] \times \mathbb{R}^5)} \lesssim \epsilon_0 \sum_{j \geq k+1} 2^{-\frac{1}{4}(j-k)} a_j. \end{aligned}$$

Then the recurrence relation (3-13) follows from (3-16) and (3-17).  $\square$

Note that by the inequality (3-5), the inequalities (3-16) and (3-17) yield that

$$(3-18) \quad \sum_{k \in \mathbb{Z}} 2^{\frac{1}{4}k} \|P_k (F(u))\|_{L_t^2 L_x^{4/3}([0, \delta] \times \mathbb{R}^5)} \lesssim \epsilon_0.$$

As an application of Lemma 3.1 and the radial Sobolev inequality (2-2), we will see that the solution  $u$  possesses some space decay property in the region  $\{|x| \geq |t| + C\}$  for some large constant  $C > 0$ . Let  $\chi(x)$  be a smooth cutoff function such that  $\chi(x) = 1$  for  $|x| \leq \frac{1}{2}$  and  $\chi(x) = 0$  for  $|x| \geq 1$ . By Lemma 2.4, there exists a dyadic integer  $R = R(u_0, u_1, \epsilon_0)$  such that  $\|(1 - \chi(\frac{x}{R}))(u_0, u_1)\|_{\dot{B}_{2,1}^{1/2} \times \dot{B}_{2,1}^{-1/2}(\mathbb{R}^5)} \leq \epsilon_0$ . Then by scaling, we have

$$(3-19) \quad \|(1 - \chi(2x))((2R)^2 u_0(2Rx), (2R)^3 u_1(2Rx))\|_{\dot{B}_{2,1}^{1/2} \times \dot{B}_{2,1}^{-1/2}(\mathbb{R}^5)} \leq \epsilon_0.$$

By abuse of notations, we will still use  $(u_0, u_1)$  to represent the initial data  $((2R)^2 u_0(2Rx), (2R)^3 u_1(2Rx))$ . Then we have,

$$(3-20) \quad \|(1 - \chi(2x))(u_0, u_1)\|_{\dot{B}_{2,1}^{1/2} \times \dot{B}_{2,1}^{-1/2}(\mathbb{R}^5)} \leq \epsilon_0.$$

In addition, by Lemma 3.1, we have

$$(3-21) \quad \|u\|_{L_{t,x}^3([-\delta/(2R), \delta/(2R)] \times \mathbb{R}^5)} \lesssim \epsilon_0,$$

$$(3-22) \quad \|u\|_{L_t^\infty([-\delta/(2R), \delta/(2R)]; \dot{B}_{2,1}^{1/2}(\mathbb{R}^5))} \lesssim 1.$$

**Lemma 3.2.** *Let  $J \subset \mathbb{R}$  be an interval such that  $u$  is a solution to (1-5) on  $J$ . Then*

$$(3-23) \quad \|u\|_{L_{t,x}^3(\{(t,x) \in J \times \mathbb{R}^5 : |x| \geq |t| + \frac{1}{2}\})} + \sup_{t \in J} \| |x|^2 u(t, x) \|_{L_x^\infty(\{x \in \mathbb{R}^5 : |x| \geq |t| + \frac{1}{2}\})} \lesssim \epsilon_0.$$

*Proof.* Let  $U(t, x)$  be the solution to (1-5) with initial data  $(1 - \chi(2x))(u_0(x), u_1(x))$ . Employing Theorem 2.10 and arguing by similar arguments in Lemma 3.1, one can deduce (3-14) when  $u$  is replaced by  $U$  and  $[-\delta, \delta]$  is replaced by  $\mathbb{R}$ . Thus

$$(3-24) \quad \|U\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} + \|U\|_{L_t^\infty \dot{B}_{2,1}^{1/2}(\mathbb{R} \times \mathbb{R}^5)} \lesssim \epsilon_0.$$

Due to the finite propagation speed property of the wave equation (1-5), we have  $u(t, x) = U(t, x)$  when  $|x| \geq |t| + \frac{1}{2}$ . Then (3-23) follows from (3-24) and the radial Sobolev inequality (2-2).  $\square$

Next, we want to show the following local properties, which will play an important role in Section 3B. Unlike the case of three dimensions in [Dodson 2019], we will make use of the inhomogeneous Strichartz estimates (2-28) to conquer the difficulties caused by the higher order decay of time.

**Lemma 3.3.** *If  $\epsilon_0$  is sufficiently small and  $\delta$  is as given in Lemma 3.1, then, for  $3 < r < 4$ , we have*

$$(3-25) \quad \sup_{-\frac{2\delta}{2R} < t < \frac{2\delta}{2R}} t^{(2r-5)/r} \|u\|_{L_x^r(\mathbb{R}^5)} + \|u\|_{L_t^{5/4} L_x^{25/6}([0, \delta/(2R)] \times \mathbb{R}^5)} \lesssim 1.$$

We remark that for  $3 < r < \infty$ , the space  $L_t^{r/(2r-5)} L_x^r(\mathbb{R} \times \mathbb{R}^5)$  is  $\dot{H}^{1/2}$ -critical but not admissible.

*Proof.* First, we consider the estimates for the linear part. Utilizing dispersive estimate (2-20), we have

$$(3-26) \quad \|S(t)P_j(u_0, u_1)\|_{L_x^\infty(\mathbb{R}^5)} \lesssim \frac{1}{t^2} [2^{3j} \|P_j u_0\|_{L^1(\mathbb{R}^5)} + 2^{2j} \|P_j u_1\|_{L^1(\mathbb{R}^5)}].$$

By Bernstein, we have

$$(3-27) \quad \begin{aligned} \|S(t)P_j(u_0, u_1)\|_{L_x^2(\mathbb{R}^5)} &\lesssim \|P_j u_0\|_{L_x^2(\mathbb{R}^5)} + 2^{-j} \|P_j u_1\|_{L_x^2(\mathbb{R}^5)} \\ &\lesssim 2^{\frac{5}{2}j} \|P_j u_0\|_{L^1(\mathbb{R}^5)} + 2^{\frac{3}{2}j} \|P_j u_1\|_{L^1(\mathbb{R}^5)}. \end{aligned}$$

Interpolating this inequality with the estimate (3-26) yields that,

$$(3-28) \quad \|S(t)P_j(u_0, u_1)\|_{L_x^r(\mathbb{R}^5)} \lesssim t^{-2(1-\frac{2}{r})} 2^{-\frac{j}{r}} [2^{3j} \|P_j u_0\|_{L^1(\mathbb{R}^5)} + 2^{2j} \|P_j u_1\|_{L^1(\mathbb{R}^5)}].$$

On the other hand, for  $r \geq \frac{5}{2}$ , employing Bernstein's estimates, we have

$$(3-29) \quad \begin{aligned} \|S(t)P_j(u_0, u_1)\|_{L_x^r(\mathbb{R}^5)} &\lesssim 2^{5(\frac{1}{2}-\frac{1}{r})j} \|S(t)P_j(u_0, u_1)\|_{L_x^2(\mathbb{R}^5)} \\ &\lesssim 2^{(2-\frac{5}{r})j} [2^{3j} \|P_j u_0\|_{L^1(\mathbb{R}^5)} + 2^{2j} \|P_j u_1\|_{L^1(\mathbb{R}^5)}]. \end{aligned}$$

This estimate and (3-28) yield that, for  $r \geq \frac{5}{2}$ ,

$$(3-30) \quad \begin{aligned} \sup_{t \in \mathbb{R}} t^{(2r-5)/r} \|S(t)(u_0, u_1)\|_{L_x^r(\mathbb{R}^5)} + \|S(t)(u_0, u_1)\|_{L_t^{r/(2r-5)} L_x^r(\mathbb{R} \times \mathbb{R}^5)} \\ \lesssim \|(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)}. \end{aligned}$$

By the reversal property of the wave equation, it suffices to prove (3-25) for  $t \geq 0$ . Using the inhomogeneous Strichartz estimates (2-28), Lemmas 2.5, 3.1, and Hölder, we have

$$(3-31) \quad \begin{aligned} \left\| \int_0^t S(t-\tau)(0, F(u(\tau))) d\tau \right\|_{L_t^{5/4} L_x^{25/6}([0, \frac{\delta}{2R}] \times \mathbb{R}^5)} \\ \lesssim \|\nabla|^{1/4} F(u)\|_{L_t^{30/29} L_x^{300/197}([0, \frac{\delta}{2R}] \times \mathbb{R}^5)} \\ \lesssim \|\nabla|^{1/2} u\|_{L_t^\infty L_x^2([0, \frac{\delta}{2R}] \times \mathbb{R}^5)}^{1/2} \|u\|_{L_{t,x}^3([0, \frac{\delta}{2R}] \times \mathbb{R}^5)}^{1/2} \|u\|_{L_t^{5/4} L_x^{25/6}([0, \frac{\delta}{2R}] \times \mathbb{R}^5)}. \end{aligned}$$

This estimates together with the estimate (3-30) yields

$$(3-32) \quad \|u\|_{L_t^{5/4} L_x^{25/6}([0, \frac{\delta}{2R}] \times \mathbb{R}^5)} \lesssim 1,$$

provided  $0 < \epsilon_0 \ll \min(1, \|(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)}^{-1})$ .

Let  $c \in (0, 1)$  to be chosen later. First, employing the dispersive estimate (2-20), Lemma 2.5, and interpolation, for  $r \in (3, 4)$ , we have

$$\begin{aligned}
(3-33) \quad & \sup_{t \in [0, \frac{\delta}{2R}]} t^{\frac{2r-5}{r}} \left\| \int_{(1-c)t}^t S(t-\tau)(0, F(u(\tau))) d\tau \right\|_{L^r(\mathbb{R}^5)} \\
& \lesssim \sup_{t \in [0, \frac{\delta}{2R}]} t^{\frac{2r-5}{r}} \int_{(1-c)t}^t \frac{1}{(t-\tau)^{2-\frac{4}{r}}} \left\| |\nabla|^{2-\frac{6}{r}} F(u(\tau)) \right\|_{L^{r'}(\mathbb{R}^5)} d\tau \\
& \lesssim \sup_{t \in [0, \frac{\delta}{2R}]} t^{\frac{2r-5}{r}} \int_{(1-c)t}^t \frac{1}{(t-\tau)^{2-\frac{4}{r}}} \|u(\tau)\|_{L^r(\mathbb{R}^5)}^{\frac{r-1}{2r-5}} \\
& \quad \times \|u(\tau)\|_{\dot{H}_x^{1/2}(\mathbb{R}^5)}^{4-\frac{12}{r}} \|u(\tau)\|_{L_x^{5/2}(\mathbb{R}^5)}^{(\frac{12}{r}-2-\frac{r-1}{2r-5})} d\tau \\
& \lesssim \left( \sup_{t \in [0, \frac{\delta}{2R}]} t^{\frac{2r-5}{r}} \|u(t)\|_{L^r(\mathbb{R}^5)} \right)^{\frac{r-1}{2r-5}} \|u\|_{L_t^\infty \dot{H}_x^{1/2}([0, \frac{\delta}{2R}] \times \mathbb{R}^5)}^{4-\frac{12}{r}} \\
& \quad \times \|u\|_{L_t^\infty L_x^{5/2}([0, \frac{\delta}{2R}] \times \mathbb{R}^5)}^{(\frac{12}{r}-2-\frac{r-1}{2r-5})} \cdot \int_{(1-c)t}^t \frac{1}{(t-\tau)^{2-\frac{4}{r}}} t^{1-\frac{4}{r}} d\tau \\
& \lesssim c^{4/r-1} \left( \sup_{t \in [0, \frac{\delta}{2R}]} t^{\frac{2r-5}{r}} \|u(t)\|_{L^r(\mathbb{R}^5)} \right)^{\frac{r-1}{2r-5}}.
\end{aligned}$$

For the remainder part, we utilize the dispersive estimate (2-20), Lemma 2.5, the Hölder inequality, and interpolation to obtain

$$\begin{aligned}
(3-34) \quad & t^{\frac{2r-5}{r}} \left\| \int_0^{(1-c)t} S(t-\tau)(0, F(u(\tau))) d\tau \right\|_{L^r(\mathbb{R}^5)} \\
& \lesssim t^{\frac{2r-5}{r}} \int_0^{(1-c)t} \frac{1}{(t-\tau)^{2-\frac{4}{r}}} \left\| |\nabla|^{2-\frac{6}{r}} F(u(\tau)) \right\|_{L_x^{r'}(\mathbb{R}^5)} d\tau \\
& \lesssim c^{\frac{4}{r}-2} \left\| |\nabla|^{2-\frac{6}{r}} F(u) \right\|_{L_{t,x}^{r'}(\mathbb{R} \times \mathbb{R}^5)} \\
& \lesssim c^{\frac{4}{r}-2} \|u\|_{L_t^{r/(2r-5)} L_x^r(\mathbb{R} \times \mathbb{R}^5)} \|u\|_{L_t^\infty \dot{H}_x^{1/2}(\mathbb{R} \times \mathbb{R}^5)}^{4-\frac{12}{r}} \|u\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)}^{\frac{12}{r}-3} \\
& \lesssim c^{\frac{4}{r}-2} \epsilon_0^{\frac{159}{7r}-\frac{39}{7}},
\end{aligned}$$

where in the last step we used the fact that

$$\|u\|_{L_t^{r/(2r-5)} L_x^r([0, \frac{\delta}{2R}] \times \mathbb{R}^5)} \lesssim \|u\|_{L_{t,x}^3([0, \frac{\delta}{2R}] \times \mathbb{R}^5)}^{\frac{75}{7}(\frac{1}{r}-\frac{6}{25})} \|u\|_{L_t^{5/4} L_x^{25/6}([0, \frac{\delta}{2R}] \times \mathbb{R}^5)}^{\frac{25}{7}(1-\frac{3}{r})} \lesssim \epsilon_0^{\frac{75}{7}(\frac{1}{r}-\frac{6}{25})}.$$

Hence, by (3-30) and (3-32)–(3-34), taking  $c > 0$  small enough and using a continuity method, there exists  $\epsilon_0 = \epsilon_0(\|(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2}) > 0$  such that

$$(3-35) \quad \sup_{t \in [0, \frac{\delta}{2R}]} t^{\frac{2r-5}{r}} \|u\|_{L_x^r} \lesssim 1, \quad \text{for } 3 < r < 4. \quad \square$$

We denote  $\delta_1 = \frac{\delta}{2R}$  for simplicity. Let  $\psi \in C_0^\infty(\mathbb{R}^5)$  be supported in  $|x| \leq \frac{\delta_1}{10}$  and  $\psi(x) = 1$  when  $|x| \leq \frac{\delta_1}{20}$ . Then we can assume that  $|\nabla \psi(x)| \lesssim \frac{1}{\delta_1}$ . For  $t \geq \delta_1$ , we split  $u(t, x) = v(t, x) + w(t, x)$ , where

$$(3-36) \quad v(t) = S(t)(\psi u_0, \psi u_1) - \int_0^{\delta_1/10} S(t-\tau)(0, \psi F(u(\tau))) d\tau.$$

We will prove that  $v$  has a decay property and  $w$  has finite energy.

### 3A. The decay part of the solution $u$ .

**Lemma 3.4.** *For  $t \geq \delta_1$ , we have*

$$(3-37) \quad \|v(t)\|_{L_x^\infty(\mathbb{R}^5)} \lesssim \delta_1^{-1/2} t^{-2}.$$

*In addition, we have*

$$(3-38) \quad \|v\|_{L_t^\infty(\mathbb{R}; \dot{B}_{2,1}^{1/2}(\mathbb{R}^5))} + \|v\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} \lesssim \delta_1^{-1/2}.$$

*Proof.* We first estimate the linear part of  $v$ . By the Huygens principle, the radial Sobolev inequality (2-2) and Lemma 2.4, we have, for  $t \geq \delta_1$

$$(3-39) \quad \|S(t)(\psi u_0, \psi u_1)\|_{L_x^\infty(\mathbb{R}^5)} \lesssim \frac{1}{t^2} \|(u_0, u_1)\|_{\dot{B}_{2,1}^{1/2} \times \dot{B}_{2,1}^{-1/2}} \lesssim \frac{1}{t^2}.$$

For the second part of  $v$  and  $t \geq \delta_1$ , using the Radial Sobolev inequality (2-2), the Huygens principle and the Strichartz estimates, we obtain

$$(3-40) \quad \begin{aligned} & \left\| \int_0^{\delta_1/10} S(t-\tau)(0, \psi F(u(\tau))) d\tau \right\|_{L_x^\infty(\mathbb{R}^5)} \\ & \lesssim \frac{1}{t^2} \left\| \int_0^{\delta_1/10} S(t-\tau)(0, \tilde{\chi} F(u(\tau))) d\tau \right\|_{\dot{B}_{2,1}^{1/2}(\mathbb{R}^5)} \\ & \lesssim \frac{1}{t^2} \left\| \int_0^{\delta_1/10} \frac{\sin(\tau|\nabla|)}{|\nabla|} (\psi F(u(\tau))) d\tau \right\|_{\dot{B}_{2,1}^{1/2}} \\ & \quad + \frac{1}{t^2} \left\| \int_0^{\delta_1/10} \frac{\cos(\tau|\nabla|)}{|\nabla|} (\psi F(u(\tau))) d\tau \right\|_{\dot{B}_{2,1}^{1/2}} \end{aligned}$$

$$(3-41) \quad \lesssim \frac{1}{t^2} \sum_{j < 0} 2^{-\frac{1}{2}j} \|P_j[\psi F(u(\tau))]\|_{L_t^1 L_x^2([0, \delta_1/10] \times \mathbb{R}^5)}$$

$$(3-42) \quad + \frac{1}{t^2} \sum_{j \geq 0} 2^{\frac{1}{4}j} \|P_j[\psi F(u(\tau))]\|_{L_t^2 L_x^{4/3}([0, \delta_1/10] \times \mathbb{R}^5)}$$

For the low frequency part (3-41), by Bernstein's estimates and Hölder's inequality, we have

$$(3-43) \quad t^2 \times (3-41) \lesssim \sum_{j < 0} 2^{\frac{j}{3}} \|F(u)\|_{L_t^1 L_x^{3/2}([0, \delta_1/10] \times \mathbb{R}^5)} \lesssim \delta_1^{\frac{1}{3}} \|u\|_{L_{t,x}^3([0, \delta_1/10] \times \mathbb{R}^5)}^2 \lesssim 1.$$

For (3-42), it suffices to estimate

$$(3-44) \quad \sum_{j \geq 0} 2^{\frac{1}{4}j} \| [P_j, \psi] F(u(\tau)) \|_{L_t^2 L_x^{4/3}([0, \delta_1/10] \times \mathbb{R}^5)}$$

$$(3-45) \quad + \sum_{j \geq 0} 2^{\frac{1}{4}j} \| \psi P_j F(u(\tau)) \|_{L_t^2 L_x^{4/3}([0, \delta_1/10] \times \mathbb{R}^5)}.$$

For (3-44), by commutator estimates, Young's inequality, the Sobolev embedding and Lemma 3.1, we have

$$(3-46) \quad \begin{aligned} (3-44) &\lesssim \sum_{j \geq 0} 2^{-\frac{3}{4}j} \delta_1^{-1} \| Q_j F(u(\tau)) \|_{L_t^2 L_x^{4/3}([0, \delta_1/10] \times \mathbb{R}^5)} \\ &\lesssim \delta_1^{-1} \sum_{j \geq 0} 2^{-\frac{3}{4}j} 2^{\frac{1}{4}j} \| F(u(\tau)) \|_{L_t^2 L_x^{5/4}([0, \delta_1/10] \times \mathbb{R}^5)} \\ &\lesssim \delta_1^{-\frac{1}{2}} \| u \|_{L_t^\infty L_x^{5/2}([0, \delta_1/10] \times \mathbb{R}^5)}^2 \lesssim \delta_1^{-\frac{1}{2}}, \end{aligned}$$

where

$$Q_j f(x) = 2^{6j} \int_{\mathbb{R}^5} |y| \phi(2^j y) |f|(x-y) dy$$

and in the first inequality we used the mean value theorem. For (3-45), by the estimate (3-18), we have

$$(3-47) \quad (3-45) \lesssim \sum_{j \geq 0} 2^{\frac{1}{4}j} \| P_j F(u) \|_{L_t^2 L_x^{4/3}} \lesssim \epsilon_0.$$

Hence, by (3-39)–(3-47), we have  $\| v(t) \|_{L_x^\infty(\mathbb{R}^5)} \lesssim \delta_1^{-\frac{1}{2}} t^{-2}$ .

Now we consider (3-38). For simplicity, we write

$$\| v \|_{S(\mathbb{R})} := \| v \|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} + \| v \|_{L_t^\infty(\mathbb{R}; \dot{B}_{2,1}^{1/2}(\mathbb{R}^5))}.$$

For the linear part, by the Strichartz estimates and Lemma 2.4, we have

$$(3-48) \quad \| S(t)(\psi u_0, \psi u_1) \|_{S(\mathbb{R})} \lesssim \| (u_0, u_1) \|_{\dot{B}_{2,1}^{1/2} \times \dot{B}_{2,1}^{-1/2}(\mathbb{R}^5)} \lesssim 1.$$

By the Strichartz estimates and repeating the arguments that deal with (3-40),

$$\begin{aligned} &\left\| \int_0^{\delta_1/10} S(t-\tau)(0, \psi F(u(\tau))) d\tau \right\|_{S(\mathbb{R})} \\ &\lesssim \left\| \int_0^{\delta_1/10} \frac{\sin(\tau|\nabla|)}{|\nabla|} (\psi F(u(\tau))) d\tau \right\|_{\dot{B}_{2,1}^{1/2}(\mathbb{R}^5)} \\ &\quad + \left\| \int_0^{\delta_1/10} \frac{\cos(\tau|\nabla|)}{|\nabla|} (\psi F(u(\tau))) d\tau \right\|_{\dot{B}_{2,1}^{1/2}(\mathbb{R}^5)} \lesssim \delta_1^{-\frac{1}{2}}. \end{aligned}$$

This completes the proof.  $\square$

### 3B. The energy part of the solution $u$ .

**Lemma 3.5.** *We have*

$$(3-49) \quad \|\vec{w}(\delta_1)\|_{\dot{H}^1 \times L^2(\mathbb{R}^5)} \lesssim \delta_1^{-1/2}.$$

*Proof.* By the definition of  $w$ , for  $t \geq \delta_1$ , we have

$$(3-50) \quad w(t) = S(t)((1-\psi)u_0, (1-\psi)u_1) - \int_0^{\frac{\delta_1}{10}} S(t-\tau)(0, (1-\psi)F(u(\tau)))d\tau - \int_{\frac{\delta_1}{10}}^t S(t-\tau)(0, F(u(\tau)))d\tau.$$

First, we consider the contribution of the third term of (3-50). Taking  $r = \frac{50}{13}$  in (3-25), by interpolation, we have

$$(3-51) \quad \begin{aligned} \|u\|_{L_t^2 L_x^4([\delta_1/10, \delta_1] \times \mathbb{R}^5)}^2 &= \int_{\delta_1/10}^{\delta_1} \|u\|_{L_x^{50/13}(\mathbb{R}^5)} \|u\|_{L_x^{25/6}(\mathbb{R}^5)} dt \\ &\lesssim \delta_1^{-\frac{1}{2}} \sup_{t \in [\frac{\delta_1}{10}, \delta_1]} [t^{\frac{7}{10}} \|u(t)\|_{L_x^{50/13}(\mathbb{R}^5)}] \|u\|_{L_t^{5/4} L_x^{25/6}([\frac{\delta_1}{10}, \delta_1] \times \mathbb{R}^5)} \lesssim \delta_1^{-\frac{1}{2}}. \end{aligned}$$

From this inequality and Strichartz, we have

$$(3-52) \quad \begin{aligned} \left\| \int_{\delta_1/10}^{\delta_1} S(\delta_1 - \tau)(0, F(u(\tau)))d\tau \right\|_{\dot{H}_x^1(\mathbb{R}^5)} &+ \left\| \partial_t \left[ \int_{\delta_1/10}^t S(t - \tau)(0, F(u(\tau)))d\tau \right]_{t=\delta_1} \right\|_{L_x^2(\mathbb{R}^5)} \\ &\lesssim \|F(u)\|_{L_t^1 L_x^2([\frac{\delta_1}{10}, \delta_1] \times \mathbb{R}^5)} \lesssim \|u\|_{L_t^2 L_x^4([\frac{\delta_1}{10}, \delta_1] \times \mathbb{R}^5)}^2 \lesssim \delta_1^{-\frac{1}{2}}. \end{aligned}$$

By Strichartz, radial Sobolev inequality (2-2) and Hölder, the second term of (3-50) can be estimated as

$$(3-53) \quad \begin{aligned} \left\| \int_0^{\delta_1/10} \vec{S}(t - \tau)(0, (1 - \psi)F(u(\tau)))d\tau \right\|_{t=\delta_1} &\lesssim \|(1 - \psi)F(u(\tau))\|_{L_t^1 L_x^2([0, \frac{\delta_1}{10}] \times \mathbb{R}^5)} \\ &\lesssim \delta_1^{\frac{1}{2}} \|(1 - \psi)u\|_{L_{t,x}^\infty([0, \frac{\delta_1}{10}] \times \mathbb{R}^5)}^{\frac{1}{2}} \|u\|_{L_{t,x}^3([0, \frac{\delta_1}{10}] \times \mathbb{R}^5)}^{\frac{3}{2}} \\ &\lesssim \delta_1^{-\frac{1}{2}} \|u\|_{L_t^\infty \dot{B}_{2,1}^{1/2}}^{\frac{1}{2}} \lesssim \delta_1^{-\frac{1}{2}}. \end{aligned}$$

Hence, it remains to estimate

$$(3-54) \quad \|(1 - \psi)(u_0, u_1)\|_{\dot{H}^1 \times L^2(\mathbb{R}^5)}.$$

For  $u_0$ , by radial Sobolev inequality (2-2) and polar coordinates,

$$(3-55) \quad \|(\nabla \psi)u_0\|_{L^2(\mathbb{R}^5)}^2 \lesssim \delta_1^{-1} \int_{\delta_1/100}^{2\delta_1} \int_{\mathbb{S}^4} u_0^2(r) d\sigma(\omega) r^4 dr \lesssim \|u_0\|_{\dot{B}_{2,1}^{1/2}(\mathbb{R}^5)}^2 \lesssim 1.$$

By the inequality (2-13), we have

$$(3-56) \quad \|(1 - \psi)\partial_r u_0\|_{L_x^2(\mathbb{R}^5)} \lesssim \delta_1^{-\frac{1}{2}}.$$

For  $u_1$ , by the inequality (2-8) and polar coordinates, one can deduce that

$$(3-57) \quad \|(1 - \psi)u_1\|_{L_x^2(\mathbb{R}^5)}^2 \lesssim \int_{\frac{\delta_1}{10}}^{\infty} \int_{\mathbb{S}^4} |u_1(r)|^2 r^4 d\sigma(\omega) dr \lesssim \int_{\frac{\delta_1}{10}}^{\infty} r^{-2} dr \lesssim \delta_1^{-1}.$$

This inequality together with (3-55) and (3-56) implies that

$$(3-58) \quad \|(1 - \psi)(u_0, u_1)\|_{\dot{H}^1 \times L^2(\mathbb{R}^5)} \lesssim \delta_1^{-\frac{1}{2}}.$$

This completes the proof.  $\square$

**3C. Global well-posedness.** In this subsection, we prove that the solution  $u$  is globally well-posed. We emphasize that the constants in “ $\lesssim$ ” in this subsection depend only upon  $\delta_1$  and  $\|(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)}$ .

**Theorem 3.6.** *Let  $u$  be the solution to (1-5) with initial data  $(u_0, u_1) \in \dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2$ . Then  $u$  is globally well-posed and such that for any compact interval  $J \subset \mathbb{R}$ ,*

$$(3-59) \quad \|u\|_{L_{t,x}^3(J \times \mathbb{R}^5)} < C(J, \|(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2}, \delta_1).$$

*Proof.* By Lemma 2.11, it suffices to show (3-59). By the time reversibility of the wave equation, we just need consider the part of  $t \geq 0$ .

For  $t \geq \delta_1$ , by  $u(t) = w(t) + v(t)$  and the formula (3-36) of  $v(t)$ , we have

$$(3-60) \quad w_{tt} - \Delta w = -|u|u.$$

We define the energy of  $w$  as (1-2) by

$$(3-61) \quad E(w(t)) = \frac{1}{2} \int_{\mathbb{R}^5} |\nabla_t w(t)|^2 dx + \frac{1}{2} \int_{\mathbb{R}^5} |\nabla_x w(t)|^2 dx + \frac{1}{3} \int_{\mathbb{R}^5} |w(t)|^3 dx.$$

By the estimates (3-22), (3-38), Lemma 3.5 and interpolation, we have

$$(3-62) \quad E(w(\delta_1)) \lesssim 1.$$

Now, we consider

$$\begin{aligned}
 (3-63) \quad & \left| \frac{d}{dt} E(w(t)) \right| \\
 &= \left| \int_{\mathbb{R}^5} (|w|w - |u|u) w_t \, dx \right| \\
 &\lesssim \|v\|_{L_x^6(\mathbb{R}^5)} \|w_t\|_{L_x^2(\mathbb{R}^5)} \|w\|_{L_x^3(\mathbb{R}^5)} + \|w_t\|_{L_x^2(\mathbb{R}^5)} \|v\|_{L_x^6(\mathbb{R}^5)} \|v\|_{L_x^3(\mathbb{R}^5)}.
 \end{aligned}$$

By interpolation and the dispersive estimate (3-37) of  $v$ , we have

$$\begin{aligned}
 (3-64) \quad & \|v\|_{L_x^6(\mathbb{R}^5)} \|w_t\|_{L_x^2(\mathbb{R}^5)} \|w\|_{L_x^3(\mathbb{R}^5)} \lesssim \frac{1}{t} \|v\|_{L_x^3(\mathbb{R}^5)}^{\frac{1}{2}} \|w_t\|_{L_x^2(\mathbb{R}^5)} \|w\|_{L_x^3(\mathbb{R}^5)} \\
 &\lesssim \frac{1}{t} E(w(t))^{\frac{5}{6}} \|v\|_{L_x^3(\mathbb{R}^5)}^{\frac{1}{2}} \\
 &\lesssim \frac{1}{t} [E(w)(t) + \|v\|_{L_x^3(\mathbb{R}^5)}^3],
 \end{aligned}$$

$$\begin{aligned}
 (3-65) \quad & \|w_t\|_{L_x^2(\mathbb{R}^5)} \|v\|_{L_x^6(\mathbb{R}^5)} \|v\|_{L_x^3(\mathbb{R}^5)} \lesssim \frac{1}{t} E(w(t))^{\frac{1}{2}} \|v\|_{L_x^3(\mathbb{R}^5)}^{\frac{3}{2}} \\
 &\lesssim \frac{1}{t} [E(w)(t) + \|v\|_{L_x^3(\mathbb{R}^5)}^3].
 \end{aligned}$$

Substituting (3-64) and (3-65) into (3-63), we obtain

$$(3-66) \quad \frac{d}{dt} E(w(t)) \leq C \frac{1}{t} (E(w(t)) + \|v\|_{L_x^3(\mathbb{R}^5)}^3).$$

Hence,

$$(3-67) \quad \frac{d}{dt} [t^{-C} (E(w(t)))] \leq t^{-C-1} \|v\|_{L_x^3(\mathbb{R}^5)}^3.$$

This estimate and the inequality (3-38) imply that

$$(3-68) \quad E(w(t)) \leq C_1 (1 + |t|)^{C_2}.$$

Thus, for any compact interval  $J \subset \mathbb{R}$ , we have

$$(3-69) \quad \|u\|_{L_{t,x}^3(J \times \mathbb{R}^5)} \leq \|v\|_{L_{t,x}^3(J \times \mathbb{R}^5)} + \|w\|_{L_{t,x}^3(J \times \mathbb{R}^5)} < \infty. \quad \square$$

#### 4. Scattering

In this section we prove Theorem 1.1 by assuming Proposition 1.3, that is, removing the dependence of  $\delta_1$  in (1-8). From the arguments in Section 3, we have  $\delta_1 = \delta/(2R)$ , where  $\delta$  and  $R$  depending the scaling and spatial profile of the initial data, respectively. We give the heuristic idea of the proof by analyzing the effect of the parameters  $\delta$  and  $R$  on the critical norm  $L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)$ .

Note that the critical norm  $L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)$  of the solution to the nonlinear equation (1-5) is invariant under the scaling transform. Hence the parameter  $\delta$  may not

be the main difficulty in proving Theorem 1.1. On the other hand, the latter parameter  $R$  relies on the spatial profile of the initial data. For example, let  $R$  be the parameter corresponding to the initial data  $(u_0, u_1)$  with compact support. The linear evolution  $\vec{S}(t)$  for  $t \in \mathbb{R}$  does not change the critical norm  $\dot{H}^{1/2} \times \dot{H}^{-1/2}$  by the Plancherel theorem, but owing to the Huygens principle for the odd-dimension linear wave equations, it does change the spatial support of the initial datum. Thus, for the initial data  $\vec{S}(t_0)(u_0, u_1)$ , the spatial parameter (may be chosen as  $R + t_0$ ) is likely very huge, when  $t_0$  is large enough. However, the  $\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2$  norm may become huge under the evolution of  $\vec{S}(t)$ . Indeed, if  $\|(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)} = 1$ , then  $\|\vec{S}(t_0)(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)} \rightarrow \infty$  as  $t_0 \rightarrow \infty$ .<sup>1</sup> Hence, if

$$\|\vec{S}(t_0)(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)} \lesssim 1,$$

then  $t_0$  remains bounded. Taking account of this fact, one may conquer the difficulties caused by the parameter  $R$ .

To finish the proof of Theorem 1.1, we need the following theorem of profile decomposition.

**4A. Profile decomposition.** Now, we recall the linear profile decomposition from [Ramos 2012] in the radial case. We refer to [Bahouri and Gérard 1999] for the profile decompositions in the energy critical spaces.

**Theorem 4.1** (profile decomposition). *Let  $C > 0$  be a fixed number and let  $(u_0^n, u_1^n)_n$  be a sequence in  $\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)$ , with*

$$(4-1) \quad \|(u_0^n, u_1^n)\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)} \leq C.$$

*Then there exist a subsequence of  $(u_0^n, u_1^n)$  (still denoted by  $(u_0^n, u_1^n)$ ), a sequence*

$$(\phi_0^j, \phi_1^j)_{j \in \mathbb{N}} \subset \dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5),$$

*a sequence*

$$(R_{0,n}^N, R_{1,n}^N)_{N \geq 1} \subset \dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)$$

*and a sequence of parameters  $(t_j^n, \lambda_j^n) \subset \mathbb{R} \times (0, \infty)$  such that for each  $N \geq 1$ ,*

$$(4-2) \quad S(t)(u_0^n, u_1^n) = \sum_{j=1}^N (\lambda_j^n)^2 [S(\lambda_j^n(t - t_j^n))(\phi_0^j, \phi_1^j)](\lambda_j^n x) + S(t)(R_{0,n}^N, R_{1,n}^N)$$

*with*

$$(4-3) \quad \lim_{N \rightarrow \infty} \limsup_{n \rightarrow \infty} \|S(t)(R_{0,n}^N, R_{1,n}^N)\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} = 0.$$

---

<sup>1</sup>By interpolation and density, it suffices to show that, for  $f \in \mathcal{S}(\mathbb{R}^5)$ ,  $\lim_{t \rightarrow \infty} \|e^{it|\nabla|} f\|_{\dot{B}_{\infty,\infty}^{-2}(\mathbb{R}^5)} = 0$ , which follows from the dispersive estimate (2-20) and Bernstein's estimates.

In addition, the parameters  $(t_j^n, \lambda_j^n)$  satisfy the orthogonality property: for any  $j \neq k$ ,

$$(4-4) \quad \lim_{n \rightarrow \infty} \left( \frac{\lambda_j^n}{\lambda_k^n} + \frac{\lambda_k^n}{\lambda_j^n} + (\lambda_k^n)^{\frac{1}{2}} (\lambda_j^n)^{\frac{1}{2}} |t_j^n - t_k^n| \right) = \infty.$$

Furthermore, for every  $N \geq 1$ ,

$$(4-5) \quad \begin{aligned} & \| (u_0^n, u_1^n) \|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)}^2 \\ &= \sum_{j=1}^N \| (\phi_0^j, \phi_1^j) \|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)}^2 + \| (R_{0,n}^N, R_{1,n}^N) \|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)}^2 + o_n(1). \end{aligned}$$

**4B. End of the proof of the main theorem.** Now, we apply the strategy in [Dodson 2019] to finish the proof of Theorem 1.1, that is, remove the  $\delta_1$  in Proposition 1.3.

We prove Theorem 1.1 by contradiction. We assume  $u$  is the solution to (1-5) with the initial data such that  $(u_0, u_1) \in \dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)$ . For  $M \geq 0$ , let

$$(4-6) \quad f(M) = \sup \{ \|u\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} : \| (u_0, u_1) \|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)} \leq M \}.$$

Then by Proposition 1.3,  $f$  is well defined. Also, by Bernstein and Theorem 2.10,  $f(M) < \infty$  when  $M$  is small enough. From the definition,  $f(M)$  is nondecreasing as  $M$  increases.

If Theorem 1.1 fails, then there exist  $M_0 < \infty$  and a sequence  $\{(u_0^n, u_1^n)\}_{n \in \mathbb{N}} \subset \dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)$ , such that

$$(4-7) \quad \| (u_0^n, u_1^n) \|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)} \leq M_0,$$

and the solution  $u^n$  to (1-5) with  $(u^n(0), (\partial_t u^n)(0)) = (u_0^n, u_1^n)$  satisfying

$$(4-8) \quad \|u^n\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} \rightarrow \infty,$$

as  $n \rightarrow \infty$ . By Theorem 4.1, we have

$$(4-9) \quad S(t)(u_0^n, u_1^n) = \sum_{j=1}^N (\lambda_j^n)^2 [S(\lambda_j^n(t - t_j^n))(\phi_0^j, \phi_1^j)] (\lambda_j^n x) + S(t)(R_{0,n}^N, R_{1,n}^N).$$

In the proof of Theorem 4.1, Ramos [2012] actually proved that

$$(4-10) \quad \vec{S}(t + t_j^n \lambda_j^n) \left[ (\lambda_j^n)^{-2} u_0^n \left( \frac{\cdot}{\lambda_j^n} \right), (\lambda_j^n)^{-3} u_1^n \left( \frac{\cdot}{\lambda_j^n} \right) \right] \Big|_{t=0} \rightharpoonup (\phi_0^j, \phi_1^j),$$

weakly in  $\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)$  as  $n \rightarrow \infty$ . From this we can prove the following:

**Proposition 4.2.** *For fixed  $j \in \mathbb{N}$ , if  $(\phi_0^j, \phi_1^j) \neq 0$ , then  $|t_j^n \lambda_j^n|$  is bounded as  $n \rightarrow \infty$ .*

*Proof.* First, if  $t_j^n \lambda_j^n$  is unbounded for  $n \in \mathbb{N}$ , then by taking a subsequence of  $n$  (still denoted by  $n$ ), we assume that  $|t_j^n \lambda_j^n| \rightarrow \infty$ , as  $n \rightarrow \infty$ . In light of the heuristic analysis at the beginning of this section, we have

$$(4-11) \quad \vec{S}(t_j^n \lambda_j^n) \left[ (\lambda_j^n)^{-2} u_0^n \left( \frac{\cdot}{\lambda_j^n} \right), (\lambda_j^n)^{-3} u_1^n \left( \frac{\cdot}{\lambda_j^n} \right) \right] \rightarrow (0, 0).$$

in  $L_x^3 \times W_x^{-1,3}(\mathbb{R}^5)$ . In fact, using (4-7) and the estimate (3-30) in Section 3, we have

$$\begin{aligned} (4-12) \quad & \left\| S(t_j^n \lambda_j^n) \left[ (\lambda_j^n)^{-2} u_0^n \left( \frac{\cdot}{\lambda_j^n} \right), (\lambda_j^n)^{-3} u_1^n \left( \frac{\cdot}{\lambda_j^n} \right) \right] \right\|_{L_x^3(\mathbb{R}^5)} \\ & \lesssim |t_j^n \lambda_j^n|^{-\frac{1}{3}} \left\| \left( (\lambda_j^n)^{-2} u_0^n \left( \frac{\cdot}{\lambda_j^n} \right), (\lambda_j^n)^{-3} u_1^n \left( \frac{\cdot}{\lambda_j^n} \right) \right) \right\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)} \\ & \lesssim |t_j^n \lambda_j^n|^{-\frac{1}{3}} \|(u_0^n, u_1^n)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)} \rightarrow 0, \end{aligned}$$

as  $n \rightarrow \infty$ . Similarly, by the dispersive estimate (2-20), Bernstein and interpolation, we have

$$\begin{aligned} (4-13) \quad & \left\| \dot{S}(t_j^n \lambda_j^n) \left[ (\lambda_j^n)^{-2} u_0^n \left( \frac{\cdot}{\lambda_j^n} \right), (\lambda_j^n)^{-3} u_1^n \left( \frac{\cdot}{\lambda_j^n} \right) \right] \right\|_{\dot{W}_x^{-1,3}(\mathbb{R}^5)} \\ & \lesssim |t_j^n \lambda_j^n|^{-\frac{1}{3}} \|(u_0^n, u_1^n)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)} \rightarrow 0, \end{aligned}$$

as  $n \rightarrow \infty$ . Hence, from the weak convergence relation (4-10), (4-12) and (4-13) imply that  $(\phi_0^j, \phi_1^j) = (0, 0)$ .  $\square$

For simplicity, we assume that every  $(\phi_0^j, \phi_1^j)$  in (4-9) is nontrivial. By Proposition 4.2,  $t_j^n \lambda_j^n$  is bounded for each fixed  $j$ , and therefore after taking a suitable subsequence of  $n$  (still denoted by  $n$ ), we can assume  $t_j^n \lambda_j^n \rightarrow t_j \in \mathbb{R}$  as  $n \rightarrow \infty$ . Hence, if we denote  $(\varphi_0^j, \varphi_1^j) = \vec{S}(-t_j)(\phi_0^j, \phi_1^j)$ , then

$$(4-14) \quad \vec{S}(-\lambda_j^n t_j^n)(\phi_0^j, \phi_1^j) - (\varphi_0^j, \varphi_1^j) \rightarrow 0,$$

in  $\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)$  as  $n \rightarrow \infty$ . Let

$$(4-15) \quad \begin{cases} \tilde{R}_{0,n}^N = R_{0,n}^N + \sum_{j=1}^N (\lambda_j^n)^2 [S(-\lambda_j^n t_j^n)(\phi_0^j, \phi_1^j)](\lambda_j^n x) - (\lambda_j^n)^2 (\varphi_0^j, \varphi_1^j)(\lambda_j^n x), \\ \tilde{R}_{1,n}^N = R_{1,n}^N + \sum_{j=1}^N (\lambda_j^n)^3 [\dot{S}(-\lambda_j^n t_j^n)(\phi_0^j, \phi_1^j)](\lambda_j^n x) - (\lambda_j^n)^3 [(\varphi_0^j, \varphi_1^j)](\lambda_j^n x), \end{cases}$$

then

$$(4-16) \quad \lim_{N \rightarrow \infty} \limsup_{n \rightarrow \infty} \|S(t)(\tilde{R}_{0,n}^N, \tilde{R}_{1,n}^N)\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} = 0.$$

Taking  $t = 0$  in (4-9), by (4-14) and (4-15), we have

$$(4-17) \quad (u_0^n, u_1^n) = \sum_{j=1}^N ((\lambda_j^n)^2 (\varphi_0^j(\lambda_j^n x), (\lambda_j^n)^3 \varphi_1^j(\lambda_j^n x)) + (\tilde{R}_{0,n}^N, \tilde{R}_{1,n}^N)).$$

In addition, by the orthogonality (4-4) and Proposition 4.2, we have for each  $j \neq k$

$$(4-18) \quad \lim_{n \rightarrow \infty} \frac{\lambda_j^n}{\lambda_k^n} + \frac{\lambda_k^n}{\lambda_j^n} = \infty,$$

as  $n \rightarrow \infty$ . Thus, for fixed  $j \in \mathbb{N}$ , we have

$$(4-19) \quad \left( (\lambda_j^n)^{-2} u_0^n \left( \frac{x}{\lambda_j^n} \right), (\lambda_j^n)^{-3} u_1^n \left( \frac{x}{\lambda_j^n} \right) \right) \rightharpoonup (\varphi_0^j, \varphi_1^j)$$

weakly in  $\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)$  as  $n \rightarrow \infty$ . By Fatou's lemma, this fact and the inequality (4-7) imply

$$(4-20) \quad \|(\varphi_0^j, \varphi_1^j)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)} \leq M_0.$$

On the other hand, (4-5) and (4-14) yield that

$$(4-21) \quad \sum_{j \geq 1} \|(\varphi_0^j, \varphi_1^j)\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)}^2 \lesssim \sup_{n \geq 1} \|(u_0^n, u_1^n)\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)}^2 \lesssim C_0^2.$$

Hence, for fixed  $\epsilon > 0$ , there exists a finite integer  $N_0$  such that

$$(4-22) \quad \sum_{j \geq N_0 + 1} \|(\varphi_0^j, \varphi_1^j)\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)}^2 \leq \epsilon.$$

By the local well-posedness theory, if  $\epsilon > 0$  is small enough, then the solution  $v^j$  to (1-5) with the initial data  $(\varphi_0^j, \varphi_1^j)$  is globally well-posed and

$$(4-23) \quad \|v^j\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} \lesssim \|(\varphi_0^j, \varphi_1^j)\|_{\dot{H}^{1/2} \times \dot{H}^{-1/2}(\mathbb{R}^5)}, \quad \text{for every } j \geq N_0 + 1.$$

For  $1 \leq j \leq N_0$ , as a consequence of Proposition 1.3, the solution to (1-5) with the initial data  $(\varphi_0^j, \varphi_1^j)$  is globally well-posed and such that

$$(4-24) \quad \|v^j\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} \lesssim_{M_0, j} 1.$$

By the orthogonality property (4-18), for any  $j \neq k$ ,

$$(4-25) \quad \lim_{n \rightarrow \infty} \iint_{\mathbb{R} \times \mathbb{R}^5} |(\lambda_j^n)^2 v^j(\lambda_j^n t, \lambda_j^n x)|^2 |(\lambda_k^n)^2 v^k(\lambda_k^n t, \lambda_k^n x)| dx dt = 0.$$

This together with the estimates (4-22)–(4-24) implies

$$(4-26) \quad \sup_{N \geq N_0 + 1} \lim_{n \rightarrow \infty} \left\| \sum_{1 \leq j \leq N} (\lambda_j^n)^2 v^j(\lambda_j^n t, \lambda_j^n x) \right\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)}$$

is bounded. Similarly, as a consequence of the trivial estimate

$$\begin{aligned} & \left| F\left(\sum_{j=1}^N (\lambda_j^n)^2 v^j(\lambda_j^n t, \lambda_j^n x)\right) - \sum_{j=1}^N F((\lambda_j^n)^2 v^j(\lambda_j^n t, \lambda_j^n x)) \right| \\ & \lesssim \sum_{1 \leq j, k \leq N, j \neq k} |(\lambda_j^n)^2 v^j(\lambda_j^n t, \lambda_j^n x)| |(\lambda_k^n)^2 v^k(\lambda_k^n t, \lambda_k^n x)| \end{aligned}$$

and the orthogonality property (4-18), we have

$$(4-27) \quad \lim_{n \rightarrow \infty} \left\| F\left(\sum_{j=1}^N (\lambda_j^n)^2 v^j(\lambda_j^n t, \lambda_j^n x)\right) - \sum_{j=1}^N F((\lambda_j^n)^2 v^j(\lambda_j^n t, \lambda_j^n x)) \right\|_{L_{t,x}^{3/2}(\mathbb{R} \times \mathbb{R}^5)} = 0.$$

Let  $u_N^n$  be an approximate solution to (1-5) defined by

$$u_N^n = \sum_{j=1}^N (\lambda_j^n)^2 v^j(\lambda_j^n x) + S(t)(\tilde{R}_{0,n}^N, \tilde{R}_{1,n}^N).$$

Then, recall the property (4-16) for  $(\tilde{R}_{0,n}^N, \tilde{R}_{1,n}^N)$  and the fact that (4-26) is uniformly bounded for  $N \geq N_0 + 1$ , we obtain

$$(4-28) \quad \limsup_{N \rightarrow \infty} \lim_{n \rightarrow \infty} \|u_N^n\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)} \lesssim 1.$$

Moreover, combining (4-27), the property (4-16) for  $(\tilde{R}_{0,n}^N, \tilde{R}_{1,n}^N)$ , and Hölder's inequality, we have

$$(4-29) \quad \limsup_{N \rightarrow \infty} \lim_{n \rightarrow \infty} \left\| F(u_N^n) - \sum_{j=1}^N F((\lambda_j^n)^2 v^j(\lambda_j^n t, \lambda_j^n x)) \right\|_{L_{t,x}^{3/2}(\mathbb{R} \times \mathbb{R}^5)} = 0.$$

Utilizing Theorem 2.12, by (4-17), (4-28) and (4-29), we have that for  $n$  large enough, the solution  $u^n$  to (1-5) with initial data  $(u_0^n, u_1^n)$  is global and such that

$$(4-30) \quad \lim_{n \rightarrow \infty} \|u^n\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)}$$

is bounded, which contradicts the hypothesis (4-8) of  $u^n$ . Thus, we have proved Theorem 1.1.

## 5. Hyperbolic coordinates and spacetime estimates

In this section, we will finish the proof of Proposition 1.3. We first reduce Proposition 1.3 to estimating the  $L_{t,x}^3$  norm of  $w$  on the region  $\Omega_2$ , which will be defined below. Without loss of generality, we assume that  $\delta_1 < \frac{1}{4}$ . As in Theorem 3.6, we also note the constants in “ $\lesssim$ ” in this section may be different in each step and are dependent on  $\delta_1$  and  $\|(u_0, u_1)\|_{\dot{B}_{1,1}^3 \times \dot{B}_{1,1}^2(\mathbb{R}^5)}$ .

**5A. Reduction of the proof of Proposition 1.3.** Now we consider the  $L_{t,x}^3$  norm of  $u$  on  $\mathbb{R}_+ \times \mathbb{R}^5$ . First, we split time-spatial region  $\mathbb{R}_+ \times \mathbb{R}^5$  as the union

$$(5-1) \quad \mathbb{R}_+ \times \mathbb{R}^5 = \Omega_1 \cup \Omega_2 \cup \Omega_3,$$

where

$$\begin{aligned} \Omega_1 &= \{(t, x) \in \mathbb{R}_+ \times \mathbb{R}^5 : |x| \geq t + \frac{1}{2}\}, \\ \Omega_2 &= \{(t, x) \in \mathbb{R}_+ \times \mathbb{R}^5 : (t + (1 - \delta_1))^2 - |x|^2 \geq 1\}. \end{aligned}$$

Since  $\delta_1 < \frac{1}{4}$ , there exists a large constant  $C > 0$ , such that

$$\Omega_3 \subset \{(t, x) \in \mathbb{R}_+ \times \mathbb{R}^5 : t + |x| \leq C\}.$$

Recalling the estimate (3-23) in Section 3, we obtain  $\|u\|_{L_{t,x}^3(\Omega_1)} \lesssim 1$ . For the bounded region  $\Omega_3$ , Theorem 3.6 yields  $\|u\|_{L_{t,x}^3(\Omega_3)} \lesssim 1$ . Hence, we just need to consider the  $L_{t,x}^3$  norm of  $u$  on the region  $\Omega_2$ . By the estimate (3-38) for  $v$ , we are reduced to showing  $\|w\|_{L_{t,x}^3(\Omega_2)} \lesssim 1$ .

**5B. Hyperbolic coordinates.** For the radial solution  $u(t, x)$  to (1-5), if we denote  $u(t, r) = u(t, x)$  for  $r = |x|$ , then

$$(5-2) \quad \partial_{tt}(r^2 u) - \partial_{rr}(r^2 u) = -2u - r^2|u|u.$$

Denote  $\bar{u}(t, r) = u(t - (1 - \delta_1), r)$  and denote  $\bar{v}, \bar{w}$  similarly. Let  $(t, r) = (e^\tau \cosh s, e^\tau \sinh s)$ ; then  $dr dt = e^{2\tau} d\tau ds$ . We denote the hyperbolic transforms by

$$(5-3) \quad \tilde{u} = \frac{e^{2\tau} \sinh^2 s}{s^2} \bar{u}(e^\tau \cosh s, e^\tau \sinh s),$$

$$(5-4) \quad \tilde{v} = \frac{e^{2\tau} \sinh^2 s}{s^2} \bar{v}(e^\tau \cosh s, e^\tau \sinh s),$$

$$(5-5) \quad \tilde{w} = \frac{e^{2\tau} \sinh^2 s}{s^2} \bar{w}(e^\tau \cosh s, e^\tau \sinh s).$$

Hence, we have

$$(5-6) \quad \partial_{\tau\tau}(s^2 \tilde{u}) - \partial_{ss}(s^2 \tilde{u}) = -\frac{2s^2}{\sinh^2 s} \tilde{u} - \frac{s^4}{\sinh^2 s} |\tilde{u}| \tilde{u},$$

$$(5-7) \quad \partial_{\tau\tau}(s^2 \tilde{v}) - \partial_{ss}(s^2 \tilde{v}) = -\frac{2s^2}{\sinh^2 s} \tilde{v},$$

$$(5-8) \quad \partial_{\tau\tau}(s^2 \tilde{w}) - \partial_{ss}(s^2 \tilde{w}) = -\frac{2s^2}{\sinh^2 s} \tilde{w} - \frac{s^4}{\sinh^2 s} |\tilde{u}| \tilde{u}.$$

Define the hyperbolic energy of  $\tilde{w}$  by

$$(5-9) \quad E_h(\tilde{w})(\tau) = \int_0^\infty \left[ \frac{1}{2} |(s^2 \tilde{w})_\tau|^2 + \frac{1}{2} |(s^2 \tilde{w})_s|^2 + \frac{|s^2 \tilde{w}|^2}{\sinh^2 s} + \frac{1}{3} \frac{|s^2 \tilde{w}|^3}{\sinh^2 s} \right] ds.$$

**5C. The hyperbolic energy for some  $\tau_0 \geq 0$ .** First, we want to prove  $E_h(\tilde{w})(\tau)$  is bounded for some  $\tau_0 \geq 0$ . We claim that it suffices to show the boundedness of

$$(5-10) \quad \int_0^\infty \left[ |(s^2 \tilde{w})_\tau(\tau_0, s)|^2 + |(s^2 \tilde{w})_s(\tau_0, s)|^2 \right] ds$$

for some  $\tau_0 > 0$ .

To prove this claim, we need the sharp Hardy inequality,

$$(5-11) \quad \left( \frac{d-2}{2} \right)^2 \int_{\mathbb{R}^d} \frac{|f(x)|^2}{|x|^2} dx \leq \int_{\mathbb{R}^d} |\nabla f|^2(x) dx.$$

By polar coordinates, we rewrite this inequality for radial functions,

$$(5-12) \quad \left( \frac{d-2}{2} \right)^2 \int_0^\infty |f(r)|^2 r^{d-3} dr \leq \int_0^\infty |\partial_r f(r)|^2(r) r^{d-1} dr.$$

Then, this inequality and integration by parts imply that

$$(5-13) \quad \begin{aligned} & \int_0^\infty |(s^2 \tilde{w}(\tau_0))_s|^2 ds \\ &= \int_0^\infty s^4 \tilde{w}_s^2(\tau_0) ds + 4 \int_0^\infty s^2 \tilde{w}(\tau_0) s \tilde{w}_s(\tau_0) ds + 4 \int_0^\infty s^2 \tilde{w}^2(\tau_0) ds \\ &= \int_0^\infty s^4 \tilde{w}_s^2(\tau_0) ds - 2 \int_0^\infty s^2 \tilde{w}(\tau_0)^2 ds \geq \frac{1}{9} \int_0^\infty s^4 \tilde{w}_s^2(\tau_0) ds. \end{aligned}$$

In addition, by Hölder and Sobolev in polar coordinates, we have

$$(5-14) \quad \begin{aligned} \int_0^\infty \frac{|s^2 \tilde{w}(\tau_0)|^3}{\sinh^2 s} ds &= \int_0^\infty \frac{s^2}{\sinh^2 s} |\tilde{w}(\tau_0)|^3 s^4 ds \\ &\lesssim \left( \int_0^\infty |\tilde{w}(\tau_0)|^{\frac{10}{3}} s^4 ds \right)^{\frac{9}{10}} \lesssim \left( \int_0^\infty |\tilde{w}_s(\tau_0)|^2 s^4 ds \right)^{\frac{3}{2}}. \end{aligned}$$

By Hardy's inequality, we have

$$(5-15) \quad \int_0^\infty \frac{|s^2 \tilde{w}(\tau_0)|^2}{\sinh^2 s} ds \lesssim \int_0^\infty \frac{1}{s^2} |\tilde{w}(\tau_0)|^2 s^4 ds \lesssim \int_0^\infty |\tilde{w}_s(\tau_0)|^2 s^4 ds.$$

Hence the claim follows.

**5C1. The hyperbolic energy for  $s > s_0 > 0$ .** For  $\tau \in [0, 1]$  and sufficiently large  $s_0 > 0$ , we can assume that  $e^{\tau-s_0} < \frac{1}{2} - \delta_1$ . By the finite speed of propagation,  $v(t, r)$  are supported in the region  $\{(t, r) \in \mathbb{R} \times \mathbb{R}_+ : r - t \lesssim \delta_1/5\}$ . Then, for  $\tau \in [0, 1]$  and  $s \geq s_0$ , we have

$$e^\tau \sinh s - [e^\tau \cosh s - (1 - \delta_1)] = 1 - \delta_1 - e^{\tau-s} > \frac{1}{2} > \frac{\delta_1}{5},$$

which leads to  $v(e^\tau \cosh s - (1 - \delta_1), e^\tau \sinh s) = 0$ . Hence, for  $\tau \in [0, 1]$ , we have

$$(5-16) \quad \begin{aligned} \int_{s_0}^{\infty} \frac{1}{2} |(s^2 \tilde{w})_\tau(\tau, s)|^2 + \frac{1}{2} |(s^2 \tilde{w})_s(\tau, s)|^2 ds \\ = \int_{s_0}^{\infty} \frac{1}{2} |(s^2 \tilde{u})_\tau(\tau, s)|^2 + \frac{1}{2} |(s^2 \tilde{u})_s(\tau, s)|^2 ds. \end{aligned}$$

Since  $u$  is a radial solution to (1-5), we have, by (2-19),

$$(5-17) \quad \begin{aligned} r^2 u(t, r) = \frac{1}{2} [(r-t)^2 u_0(r-t) + (r+t)^2 u_0(r+t)] \\ - \frac{1}{2} t r^{-1} \int_{r-t}^{r+t} s u_0(s) ds + \frac{1}{4r} \int_{r-t}^{r+t} s (s^2 + r^2 - t^2) u_1(s) ds \\ + \frac{1}{4r} \int_0^t \int_{r-t+s}^{r+t-s} \rho (\rho^2 + r^2 - (t-s)^2) |u| u(s, \rho) d\rho ds, \end{aligned}$$

for  $r \geq t \geq 0$ . Hence, by the hyperbolic transform (5-3), we have

$$(5-18) \quad s^2 \tilde{u}(\tau, s) = \frac{1}{2} [(1 - \delta_1 - e^{\tau-s})^2 u_0(1 - \delta_1 - e^{\tau-s}) + (e^{\tau+s} - (1 - \delta_1))^2 u_0(e^{\tau+s} - (1 - \delta_1))]$$

$$(5-19) \quad - \frac{1}{2} (e^\tau \cosh s - (1 - \delta_1)) (e^\tau \sinh s)^{-1} \int_{1 - \delta_1 - e^{\tau-s}}^{e^{\tau+s} - (1 - \delta_1)} \rho u_0(\rho) d\rho$$

$$(5-20) \quad + \frac{1}{4} \int_{1 - \delta_1 - e^{\tau-s}}^{e^{\tau+s} - (1 - \delta_1)} \rho \frac{\rho^2 + (e^{\tau+s} - (1 - \delta_1))(1 - \delta_1 - e^{\tau-s})}{e^\tau \sinh s} u_1(\rho) d\rho$$

$$(5-21) \quad + \frac{1}{4} \int_{1 - \delta_1}^{e^\tau \cosh s} \int_{t - e^{\tau-s}}^{e^{\tau+s} - t} \rho \frac{\rho^2 + (e^{\tau+s} - t)(t - e^{\tau-s})}{e^\tau \sinh s} |\bar{u}| \bar{u}(t, \rho) d\rho dt.$$

For (5-18), by a direct calculation, we obtain

$$(5-22) \quad (\partial_\tau + \partial_s)(5-18) = 2(e^{\tau+s} - (1 - \delta_1)) e^{\tau+s} u_0(e^{\tau+s} - (1 - \delta_1)) \\ + (e^{\tau+s} - (1 - \delta_1))^2 u'_0(e^{\tau+s} - (1 - \delta_1)) e^{\tau+s},$$

$$(5-23) \quad (\partial_\tau - \partial_s)(5-18) = 2(1 - \delta_1 - e^{\tau-s}) e^{\tau-s} u_0(1 - \delta_1 - e^{\tau-s}) \\ + ((1 - \delta_1 - e^{\tau-s})^2 u'_0(1 - \delta_1 - e^{\tau-s}) e^{\tau-s}).$$

Using the estimate (2-13) in Section 2 and polar coordinates, we deduce that

$$(5-24) \quad \int_{s_0}^{\infty} |(e^{\tau+s} - (1-\delta_1))e^{\tau+s} u_0(e^{\tau+s} - (1-\delta_1))|^2 ds \lesssim \int_0^{\infty} |u_0(r)|^2 r^3 dr \lesssim 1,$$

$$(5-25) \quad \int_{s_0}^{\infty} |(e^{\tau+s} - (1-\delta_1))^2 e^{\tau+s} u'_0(e^{\tau+s} - (1-\delta_1))|^2 ds \lesssim \int_0^{\infty} |\partial_r u_0(r)|^2 r^5 dr \lesssim 1.$$

By the radial Sobolev inequality (2-2), we have  $|u_0(r)| \lesssim r^{-2}$ . This estimate and the inequality (2-3) imply

$$(5-26) \quad \begin{aligned} & \int_{s_0}^{\infty} |(1 - \delta_1 - e^{\tau-s})e^{\tau-s} u_0(1 - \delta_1 - e^{\tau-s})|^2 ds \\ & \quad + \int_{s_0}^{\infty} |((1 - \delta_1 - e^{\tau-s})^2 u'_0(1 - \delta_1 - e^{\tau-s})e^{\tau-s})|^2 ds \\ & \lesssim \int_{s_0}^{\infty} e^{-2s} ds \lesssim 1. \end{aligned}$$

We now take the derivatives of (5-19) with respect to  $\tau$  and  $s$ ,

$$(5-27) \quad \partial_{\tau}(5-19) = \frac{1 - \delta_1}{2e^{\tau} \sinh s} \int_{1 - \delta_1 - e^{\tau-s}}^{e^{\tau+s} - (1 - \delta_1)} \rho u_0(\rho) d\rho + I_1 + I_2$$

$$(5-28) \quad \partial_s(5-19) = \partial_s \left( \frac{e^{\tau} \cosh s - (1 - \delta_1)}{2e^{\tau} \sinh s} \right) \int_{1 - \delta_1 - e^{\tau-s}}^{e^{\tau+s} - (1 - \delta_1)} \rho u_0(\rho) d\rho + I_1 - I_2,$$

where

$$(5-29) \quad I_1 = \frac{e^{\tau} \cosh s - (1 - \delta_1)}{2e^{\tau} \sinh s} e^{\tau+s} (e^{\tau+s} - (1 - \delta_1)) u_0(e^{\tau+s} - (1 - \delta_1)),$$

$$(5-30) \quad I_2 = \frac{e^{\tau} \cosh s - (1 - \delta_1)}{2e^{\tau} \sinh s} e^{\tau-s} (1 - \delta_1 - e^{\tau-s}) u_0(1 - \delta_1 - e^{\tau-s}).$$

For the first term in the right-hand side of (5-27), by the inequality (2-3), we have

$$(5-31) \quad \int_{s_0}^{\infty} \left| e^{-s} \int_{1 - \delta_1 - e^{\tau-s}}^{e^{\tau+s} - (1 - \delta_1)} \rho u_0(\rho) d\rho \right|^2 ds \lesssim \int_{s_0}^{\infty} e^{-2s} ds \lesssim 1.$$

By similar estimates, one can find that the contribution of the first term in the right-hand side of (5-28) to (5-16) is finite. For  $I_1$ , a change of variables and the inequality (2-13) yield

$$(5-32) \quad \int_{s_0}^{\infty} |I_1|^2 ds \lesssim \int_{s_0}^{\infty} |e^{2s} u_0(e^{\tau+s} - (1 - \delta_1))|^2 ds \lesssim \int_{\frac{1}{2}e^{s_0}}^{\infty} \rho^3 |u_0(\rho)|^2 d\rho \lesssim 1.$$

For  $I_2$ , by (2-2), we obtain

$$(5-33) \quad \int_{s_0}^{\infty} |I_2| ds \lesssim \int_{s_0}^{\infty} |e^{-s}|^2 ds \lesssim 1.$$

Next, we consider the contribution of (5-20) to (5-16). For simplicity, we consider

$$(5-34) \quad \frac{1}{2}(\partial_{\tau} - \partial_s)(5-20) \\ = e^{\tau-s}(1 - \delta_1 - e^{\tau-s})^2 u_1(1 - \delta_1 - e^{\tau-s})$$

$$(5-35) \quad + \frac{e^{\tau-s}(e^{\tau+s} - (1 - \delta_1))^2}{(e^{\tau+s} - e^{\tau-s})^2} \int_{1-\delta_1-e^{\tau-s}}^{e^{\tau+s}-(1-\delta_1)} \rho u_1(\rho) d\rho$$

$$(5-36) \quad + \frac{e^{\tau-s}}{(e^{\tau+s} - e^{\tau-s})^2} \int_{1-\delta_1-e^{\tau-s}}^{e^{\tau+s}-(1-\delta_1)} \rho^3 u_1(\rho) d\rho,$$

and

$$(5-37) \quad \frac{1}{2}(\partial_{\tau} + \partial_s)(5-20) \\ = e^{\tau+s}(e^{\tau+s} - (1 - \delta_1))^2 u_1(e^{\tau+s} - (1 - \delta_1))$$

$$(5-38) \quad + \frac{e^{\tau+s}(1 - \delta_1 - e^{\tau-s})^2}{(e^{\tau+s} - e^{\tau-s})^2} \int_{1-\delta_1-e^{\tau-s}}^{e^{\tau+s}-(1-\delta_1)} \rho u_1(\rho) d\rho$$

$$(5-39) \quad + \frac{e^{\tau+s}}{(e^{\tau+s} - e^{\tau-s})^2} \int_{1-\delta_1-e^{\tau-s}}^{e^{\tau+s}-(1-\delta_1)} \rho^3 u_1(\rho) d\rho.$$

Using the estimates (2-4) and (2-13), we can easily estimate the contributions of (5-34)–(5-38) to (5-16). Let  $\mathbb{I}_J(y)$  be the characteristic function of an interval  $J \subset \mathbb{R}$ . For (5-39), by the inequality (2-13) and a change of variables, we see that

$$(5-40) \quad \begin{aligned} \int_{s_0}^{\infty} \left| e^{-s} \int_{1-\delta_1-e^{\tau-s}}^{e^{\tau+s}-(1-\delta_1)} \rho^3 u_1(\rho) d\rho \right|^2 ds &\lesssim \int_{s_0}^{\infty} \left| e^{-s} \int_0^{2e^s} \rho^3 |u_1(\rho)| d\rho \right|^2 ds \\ &\lesssim \int_0^{\infty} \left| \frac{1}{\eta} \int_0^{2\eta} \rho^3 |u_1(\rho)| d\rho \right|^2 \frac{1}{\eta} d\eta \\ &\lesssim \int_0^{\infty} \left| \frac{1}{\eta} \int_0^{2\eta} \rho^{\frac{5}{2}} |u_1(\rho)| d\rho \right|^2 d\eta \\ &\lesssim \int_0^{\infty} \left| \mathcal{M}(\mathbb{I}_{[0,\infty)}(\rho) \rho^{\frac{5}{2}} u_1(\rho)) \right|^2(\eta) d\eta \\ &\lesssim \int_0^{\infty} |u_1(\rho)|^2 \rho^5 d\rho \lesssim 1, \end{aligned}$$

where  $\mathcal{M}$  is the Hardy–Littlewood maximal function and we used the fact that  $\mathcal{M}$  is bounded in  $L^2$ .

Next, we consider the contribution of (5-21) to the energy (5-16). Also, for simplicity, we consider

$$(\partial_\tau + \partial_s)(5-21)$$

$$(5-41) \quad = e^{\tau+s} \int_{1-\delta_1}^{e^\tau \cosh s} (e^{\tau+s} - t)^2 (|\bar{u}| \bar{u})(t, e^{\tau+s} - t) dt$$

$$(5-42) \quad + \frac{e^{\tau+s}}{(e^{\tau+s} - e^{\tau-s})^2} \int_{1-\delta_1}^{e^\tau \cosh s} \int_{t-e^{\tau-s}}^{e^{\tau+s}-t} \rho (t - e^{\tau-s})^2 (|\bar{u}| \bar{u})(t, \rho) d\rho dt$$

$$(5-43) \quad - \frac{e^{\tau+s}}{(e^{\tau+s} - e^{\tau-s})^2} \int_{1-\delta_1}^{e^\tau \cosh s} \int_{t-e^{\tau-s}}^{e^{\tau+s}-t} \rho^3 (|\bar{u}| \bar{u})(t, \rho) d\rho dt,$$

and

$$(\partial_\tau - \partial_s)(5-21)$$

$$(5-44) \quad = e^{\tau-s} \int_{1-\delta_1}^{e^\tau \cosh s} (t - e^{\tau-s})^2 |\bar{u}| \bar{u}(t, t - e^{\tau-s}) dt$$

$$(5-45) \quad - \frac{e^{\tau-s}}{(e^{\tau+s} - e^{\tau-s})^2} \int_{1-\delta_1}^{e^\tau \cosh s} \int_{t-e^{\tau-s}}^{e^{\tau+s}-t} \rho (e^{\tau+s} - t)^2 |\bar{u}| \bar{u}(t, \rho) d\rho dt$$

$$(5-46) \quad + \frac{e^{\tau-s}}{(e^{\tau+s} - e^{\tau-s})^2} \int_{1-\delta_1}^{e^\tau \cosh s} \int_{t-e^{\tau-s}}^{e^{\tau+s}-t} \rho^3 |\bar{u}| \bar{u}(t, \rho) d\rho dt.$$

By the definition of  $\bar{u}$ , the inequality (3-23) and Hölder, the contribution of (5-41) can be estimated as

$$(5-47) \quad \begin{aligned} & \int_{s_0}^{\infty} \left| e^{\tau+s} \int_{1-\delta_1}^{e^\tau \cosh s} (e^{\tau+s} - t)^2 |\bar{u}|^2 (t, e^{\tau+s} - t) dt \right|^2 ds \\ & \lesssim \int_{s_0}^{\infty} \int_{1-\delta_1}^{e^\tau \cosh s} |\bar{u}|^4 (t, e^{\tau+s} - t) e^{7s} dt ds \\ & \lesssim \int_{\frac{1}{4}e^{s_0}}^{\infty} \int_{1-\delta_1}^{\rho} |\bar{u}|^4 (t, \rho) \rho^6 dt d\rho \lesssim \int_0^{\infty} \int_{\rho > t + \frac{1}{2}} |\bar{u}|^4 (t, \rho) \rho^6 d\rho dt \\ & \lesssim \|u\|_{L^3(\{|x| > |t| + \frac{1}{2}\})}^3 \sup_{t \geq 0} \| |x|^2 u(t, x) \|_{L_x^{\infty}(\{|x| > t + \frac{1}{2}\})} \lesssim 1. \end{aligned}$$

Now, we consider (5-42) and (5-43). By Hölder, a change of variables, and the inequality (3-23), we have

$$(5-48) \quad \begin{aligned} & \int_{s_0}^{\infty} \left| \int_{1-\delta_1}^{e^\tau \cosh s} e^{-s} \int_{t-e^{\tau-s}}^{e^{\tau+s}-t} (t^2 \rho + \rho^3) |\bar{u}|^2 (t, \rho) d\rho dt \right|^2 ds \\ & \lesssim \int_{s_0}^{\infty} \left| \int_{1-\delta_1}^{e^\tau \cosh s} \mathcal{M}(\mathbb{I}_{[t-\frac{1}{2}+\delta_1, \infty)}(\rho) \rho^3 |\bar{u}|^2 (t, \rho)) (e^\tau \sinh s) dt \right|^2 ds \end{aligned}$$

$$\begin{aligned}
&\lesssim \int_{s_0}^{\infty} \int_{1-\delta_1}^{e^{\tau} \cosh s} [\mathcal{M}(\mathbb{I}_{[t-\frac{1}{2}+\delta_1, \infty)}(\rho) \rho^3 |\bar{u}|^2(t, \rho))(e^{\tau} \sinh s)]^2 e^{\tau} \cosh s dt ds \\
&\lesssim \int_{1-\delta_1}^{\infty} \int_{e^{s_0}}^{\infty} [\mathcal{M}(\mathbb{I}_{[t-\frac{1}{2}+\delta_1, \infty)}(\rho) \rho^3 |\bar{u}|^2(t, \rho))(r)]^2 dr dt \\
&\lesssim \int_{1-\delta_1}^{\infty} \int_{r > t - \frac{1}{2} + \delta_1} r^6 |\bar{u}|^4(t, r) dr dt \\
&\lesssim \int_0^{\infty} \int_{\rho > t + \frac{1}{2}} |u|^4(t, \rho) \rho^6 d\rho dt \\
&\lesssim \|u\|_{L^3(\{|x| > |t| + \frac{1}{2}\})}^3 \sup_{t \geq 0} \| |x|^2 u(t, x) \|_{L_x^{\infty}(\{|x| > t + \frac{1}{2}\})} \\
&\lesssim 1.
\end{aligned}$$

Thus, the contribution of (5-42) and (5-43) to (5-16) is finite.

For (5-44), by the fact that  $e^{\tau-s_0} < \frac{1}{2} - \delta_1$ , the definition of  $\bar{u}$ , and the inequality (3-23), we have

$$\begin{aligned}
(5-49) \quad &\int_{s_0}^{\infty} e^{-2s} \left| \int_{1-\delta_1}^{e^{\tau} \cosh s} (t - e^{\tau-s})^2 |\bar{u}|^2(t, t - e^{\tau-s}) dt \right|^2 ds \\
&\lesssim \int_{s_0}^{\infty} e^{-2s} \left| \int_{1-\delta_1}^{e^{\tau} \cosh s} t^{-2} dt \right|^2 \lesssim 1.
\end{aligned}$$

Similarly, for (5-45) and (5-46), by the fact that  $e^{\tau-s_0} < \frac{1}{2} - \delta_1$  and the inequality (3-23), we can obtain that

$$\begin{aligned}
(5-50) \quad &\int_{s_0}^{\infty} e^{-6s} \left| \int_{1-\delta_1}^{e^{\tau} \cosh s} \int_{t-e^{\tau-s}}^{e^{\tau+s}-t} [\rho^3 + \rho(e^{\tau+s} - t)^2] |\bar{u}|^2(t, \rho) d\rho dt \right|^2 ds \\
&\lesssim \int_{s_0}^{\infty} e^{-2s} \left| \int_{1-\delta_1}^{e^{\tau} \cosh s} \int_{t-e^{\tau-s}}^{e^{\tau+s}-t} \rho |\bar{u}|^2(t, \rho) d\rho dt \right|^2 ds \\
&\lesssim \int_{s_0}^{\infty} e^{-2s} \left| \int_{1-\delta_1}^{e^{\tau} \cosh s} \int_{t-e^{\tau-s}}^{e^{\tau+s}-t} \rho^{-3} d\rho dt \right|^2 ds \\
&\lesssim \int_{s_0}^{\infty} e^{-2s} \left| \int_{1-\delta_1}^{e^{\tau} \cosh s} t^{-2} dt \right|^2 ds \\
&\lesssim 1.
\end{aligned}$$

Hence, combining (5-24)–(5-50), we have

$$(5-51) \quad \int_{s_0}^{\infty} \frac{1}{2} |(s^2 \tilde{w})_{\tau}|^2 + \frac{1}{2} |(s^2 \tilde{w})_s|^2 ds \lesssim 1.$$

**5C2. The hyperbolic energy for  $0 \leq s \leq s_0$ .** By the hyperbolic transform (5-5), we have

$$(5-52) \quad \begin{aligned} (s^2 \tilde{w})_\tau(\tau, s) = & 2e^{2\tau} \sinh^2 s \bar{w}(e^\tau \cosh s, e^\tau \sinh s) \\ & + e^{3\tau} \sinh^2 s \cosh s \bar{w}_t(e^\tau \cosh s, e^\tau \sinh s) \\ & + e^{3\tau} \sinh^3 s \bar{w}_r(e^\tau \cosh s, e^\tau \sinh s), \end{aligned}$$

$$(5-53) \quad \begin{aligned} (s^2 \tilde{w})_s(\tau, s) = & 2e^{2\tau} \sinh s \cosh s \bar{w}(e^\tau \cosh s, e^\tau \sinh s) \\ & + e^{3\tau} \sinh^3 s \bar{w}_t(e^\tau \cosh s, e^\tau \sinh s) \\ & + e^{3\tau} \sinh^2 s \cosh s \bar{w}_r(e^\tau \cosh s, e^\tau \sinh s). \end{aligned}$$

Hence

$$(5-54) \quad \begin{aligned} & \int_0^1 \int_0^{s_0} |(s^2 \tilde{w})_\tau|^2 + |(s^2 \tilde{w})_s|^2 ds d\tau \\ & \lesssim \int_0^1 \int_0^{s_0} e^{4\tau} \sinh^2 s (\sinh^2 s + \cosh^2 s) |\bar{w}|^2(e^\tau \cosh s, e^\tau \sinh s) ds d\tau \end{aligned}$$

$$(5-55) \quad + \int_0^1 \int_0^{s_0} e^{6\tau} \sinh^4 s (\sinh^2 s + \cosh^2 s) [\bar{w}_t^2 + \bar{w}_r^2](e^\tau \cosh s, e^\tau \sinh s) ds d\tau.$$

Taking  $C_0 = e^{1+s_0}$ , by a change of variables, the Hardy inequality and the inequality (3-68), we obtain

$$(5-56) \quad (5-54) \lesssim \iint_{|x|+|t| \leq C_0} \frac{1}{|x|^2} |\bar{w}|^2(t, x) dx dt \lesssim \sup_{0 < t < C_0} \|\nabla_x \bar{w}\|_{L_x^2(\mathbb{R}^5)} \lesssim 1.$$

Similarly, for (5-55), by a change of variables, we have

$$(5-57) \quad (5-55) \lesssim \iint_{|x|+|t| \leq C_0} |\nabla_{t,x} \bar{w}|^2(t, x) dx dt \lesssim \sup_{0 < t < C_0} \|\nabla_{t,x} \bar{w}\|_{L_x^2(\mathbb{R}^5)} \lesssim 1.$$

Then, by the mean value theorem, there exists  $\tau_0 \in [0, 1]$ , such that

$$(5-58) \quad \int_0^{s_0} |(s^2 \tilde{w})_\tau|^2(\tau_0, s) + |(s^2 \tilde{w})_s|^2(\tau_0, s) ds \lesssim 1.$$

This estimate along with (5-51) implies

$$(5-59) \quad \int_0^\infty |(s^2 \tilde{w})_\tau|^2(\tau_0, s) + |(s^2 \tilde{w})_s|^2(\tau_0, s) ds \lesssim 1.$$

**5D. Uniform boundedness of the hyperbolic energy of  $\tilde{w}$ .** We are now going to show that  $E_h(\tilde{w})(\tau)$  is uniformly bounded for  $\tau \in \mathbb{R}_+$ .

A simple calculation gives

$$(5-60) \quad \frac{d}{d\tau} E_h(\tilde{w}(\tau)) = \int \frac{|s^2 \tilde{w}| s^2 \tilde{w} - |s^2 \tilde{u}| s^2 \tilde{u}}{\sinh^2 s} s^2 \tilde{w}_\tau ds.$$

Utilizing the decay property (3-37) of  $v$ , we have, for  $\tau, s \geq 0$ ,

$$(5-61) \quad (e^\tau \cosh s - (1 - \delta_1))^2 v(e^\tau \cosh s - (1 - \delta_1), e^\tau \sinh s) \lesssim \delta_1^{-\frac{1}{2}}.$$

The Huygens principle implies that  $v(e^\tau \cosh s - (1 - \delta_1), e^\tau \sinh s) = 0$  unless  $1 - \frac{6}{5}\delta_1 \leq e^{\tau-s} \leq 1 - \frac{4}{5}\delta_1$ . Thus, for  $\tau, s \geq 0$ , we have

$$(5-62) \quad \begin{aligned} \frac{s^2 |\tilde{v}(\tau, s)|}{\sinh^2 s} &= e^{2\tau} |v(e^\tau \cosh s - (1 - \delta_1), e^\tau \sinh s)| \\ &\lesssim \frac{e^{2\tau} \mathbb{I}_{\{s \geq 0: e^{\tau-s} \leq 1 - \frac{4}{5}\delta_1\}}}{(e^\tau \cosh s - (1 - \delta_1))^2} \lesssim e^{-\tau}. \end{aligned}$$

Hence, by Hölder and interpolation, we have

$$(5-63) \quad \begin{aligned} &\int_0^\infty \frac{1}{\sinh^2 s} |s^2 \tilde{w}_\tau| |s^2 \tilde{v}|^2 ds \\ &\lesssim \left( \int_0^\infty |s^2 \tilde{w}_\tau|^2 ds \right)^{\frac{1}{2}} \left( \int_0^\infty \frac{1}{\sinh^2 s} |s^2 \tilde{v}|^3 ds \right)^{\frac{1}{2}} \left\| \frac{s^2}{\sinh^2 s} \tilde{v}(\tau, s) \right\|_{L_s^\infty}^{\frac{1}{2}} \\ &\lesssim e^{-\tau/2} E_h(\tilde{w}(\tau))^{\frac{1}{2}} \left( \int_0^\infty \frac{1}{\sinh^2 s} |s^2 \tilde{v}|^3 ds \right)^{\frac{1}{2}}, \end{aligned}$$

$$(5-64) \quad \begin{aligned} &\int_0^\infty \frac{1}{\sinh^2 s} |s^2 \tilde{w}_\tau| |s^2 \tilde{v}| |s^2 \tilde{w}| ds \\ &\lesssim \left( \int_0^\infty |s^2 \tilde{w}_\tau|^2 ds \right)^{\frac{1}{2}} \left( \int_0^\infty \frac{1}{\sinh^2 s} |s^2 \tilde{w}|^3 ds \right)^{\frac{1}{3}} \\ &\quad \times \left( \int_0^\infty |s^2 \tilde{v}(\tau, s)|^6 \frac{1}{\sinh^8 s} ds \right)^{\frac{1}{6}} \\ &\lesssim e^{-\tau/2} E_h(\tilde{w}(\tau))^{\frac{5}{6}} \left( \int_0^\infty \frac{1}{\sinh^2 s} |s^2 \tilde{v}|^3 ds \right)^{\frac{1}{6}}. \end{aligned}$$

Combining (5-63) with (5-64) and employing Hölder again, we have

$$(5-65) \quad \frac{d}{d\tau} E_h(\tilde{w}(\tau)) \lesssim e^{-\tau/2} \left[ E_h(\tilde{w}(\tau)) + \int_0^\infty \frac{1}{\sinh^2 s} |s^2 \tilde{v}|^3 ds \right].$$

On the other hand, by a change of variables, we have

$$(5-66) \quad \int_0^\infty \int_0^\infty \frac{1}{\sinh^2 s} |s^2 \tilde{v}|^3 ds d\tau \leq \int_{\delta_1}^\infty \int_0^\infty |v(t, r)|^3 r^4 dr dt \\ \leq \|v\|_{L_{t,x}^3([\delta_1, \infty) \times \mathbb{R}^5)}^3 \lesssim 1.$$

Hence, by Gronwall's inequality, (5-65) and (5-66) yield that  $E_h(\tilde{w})(\tau)$  is uniformly bounded in  $\mathbb{R}_+$ .

**5E. Conclusion of the proof of Proposition 1.3.** We complete the proof by studying the Morawetz action in hyperbolic coordinates.

**Proposition 5.1.** *Let  $\tilde{w}$  be defined in (5-5), then*

$$(5-67) \quad \iint_{\Omega_2} |w(t, r)|^3 r^4 dt dr = \int_0^\infty \int_0^\infty \frac{|s^2 \tilde{w}|^3}{\sinh^2 s} ds d\tau \lesssim 1.$$

*Proof.* Define the Morawetz action by

$$(5-68) \quad M(\tau) = \int_0^\infty (s^2 \tilde{w})_\tau (s^2 \tilde{w})_s ds = \int_0^\infty \tilde{w}_\tau \left( \tilde{w}_s + \frac{2}{s} \tilde{w} \right) s^4 ds.$$

One can easily find that  $|M(\tau)| \leq E_h(\tilde{w})(\tau)$ . By (5-8), we have

$$(5-69) \quad \begin{aligned} \frac{d}{d\tau} M(\tau) &= - \int_0^\infty \frac{2s^2 \tilde{w}}{\sinh^2 s} (s^2 \tilde{w})_s ds - \int_0^\infty \frac{s^4}{\sinh^2 s} |\tilde{u}| \tilde{u} s^2 \tilde{w}_s ds \\ &= - 2 \int_0^\infty \frac{|s^2 \tilde{w}|^2 \cosh s}{\sinh^2 s \sinh s} ds - \frac{2}{3} \int_0^\infty \frac{|s^2 \tilde{w}|^3 \cosh s}{\sinh^2 s \sinh s} ds \\ &\quad + \int_0^\infty \frac{s^4}{\sinh^2 s} (|\tilde{w}| \tilde{w} - |\tilde{u}| \tilde{u}) s^2 \tilde{w}_s ds. \end{aligned}$$

By Hölder, the estimate (5-62), and the fact that  $E_h(\tilde{w}(\tau))$  is uniformly bounded for  $\tau \geq 0$ , we have

$$(5-70) \quad \begin{aligned} &\left| \int_0^\infty \int_0^\infty \frac{s^4}{\sinh^2 s} (|\tilde{w}| \tilde{w} - |\tilde{u}| \tilde{u}) s^2 \tilde{w}_s ds d\tau \right| \\ &\lesssim \int_0^\infty \int_0^\infty \frac{1}{\sinh^2 s} |s^2 \tilde{w}_\tau| |s^2 \tilde{v}|^2 ds d\tau \\ &\quad + \int_0^\infty \int_0^\infty \frac{1}{\sinh^2 s} |s^2 \tilde{w}_\tau| |s^2 \tilde{v}| |s^2 \tilde{w}| ds d\tau \end{aligned}$$

$$\begin{aligned}
&\lesssim \int_0^\infty e^{-\tau} E_h(\tilde{w}(\tau))^{\frac{1}{2}} \left( \int_0^\infty \frac{1}{\sinh^2 s} |s^2 \tilde{v}|^3 ds \right)^{\frac{1}{2}} \tau \\
&\quad + \int_0^\infty e^{-\tau} E_h(\tilde{w}(\tau))^{\frac{5}{6}} \left( \int_0^\infty \frac{1}{\sinh^2 s} |s^2 \tilde{v}|^3 ds \right)^{\frac{1}{6}} d\tau \\
&\lesssim \|v\|_{L_{t,x}^3(\mathbb{R} \times \mathbb{R}^5)}^3 \lesssim 1.
\end{aligned}$$

This together with the equality (5-69) and the fact  $M(\tau)$  is uniformly bounded for  $\tau \geq 0$ , implies that

$$(5-71) \quad \int_0^\infty \int_0^\infty \frac{|s^2 \tilde{w}|^3}{\sinh^2 s} \frac{\cosh s}{\sinh s} ds d\tau \lesssim 1.$$

Thus, we have

$$(5-72) \quad \int_0^\infty \int_0^\infty \frac{|s^2 \tilde{w}|^3}{\sinh^2 s} ds d\tau \lesssim 1.$$

This yields (5-67) by the definition of  $\tilde{w}$ .  $\square$

### Acknowledgments

Miao is supported in part by the National Natural Science Foundation of China under grant No. 11831004, No. 11926303 and No. 11671047. Yang is supported by the NNSF of China under grant No. 11901032 and the Labex MME-DII. Zhao is supported in part by the Chinese Postdoc Foundation Grant 2019M650457 and National Natural Science Foundation of China (NSAF-U1930402).

### References

- [Bahouri and Gérard 1999] H. Bahouri and P. Gérard, “High frequency approximation of solutions to critical nonlinear wave equations”, *Amer. J. Math.* **121**:1 (1999), 131–175. MR Zbl
- [Bahouri and Shatah 1998] H. Bahouri and J. Shatah, “Decay estimates for the critical semilinear wave equation”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **15**:6 (1998), 783–789. MR Zbl
- [Christ and Weinstein 1991] F. M. Christ and M. I. Weinstein, “Dispersion of small amplitude solutions of the generalized Korteweg–de Vries equation”, *J. Funct. Anal.* **100**:1 (1991), 87–109. MR Zbl
- [Colzani et al. 2002] L. Colzani, A. Cominardi, and K. Stempak, “Radial solutions to the wave equation”, *Ann. Mat. Pura Appl.* (4) **181**:1 (2002), 25–54. MR Zbl
- [Dodson 2019] B. Dodson, “Global well-posedness and scattering for the radial, defocusing, cubic wave equation with initial data in a critical Besov space”, *Anal. PDE* **12**:4 (2019), 1023–1048. MR Zbl
- [Ginibre and Velo 1985] J. Ginibre and G. Velo, “The global Cauchy problem for the nonlinear Klein–Gordon equation”, *Math. Z.* **189**:4 (1985), 487–505. MR Zbl
- [Ginibre and Velo 1989] J. Ginibre and G. Velo, “The global Cauchy problem for the nonlinear Klein–Gordon equation, II”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **6**:1 (1989), 15–35. MR Zbl

[Ginibre and Velo 1995] J. Ginibre and G. Velo, “Generalized Strichartz inequalities for the wave equation”, pp. 153–160 in *Partial differential operators and mathematical physics* (Holzhau, Germany, 1994), edited by M. Demuth and B.-W. Schulze, Oper. Theory Adv. Appl. **78**, Birkhäuser, Basel, 1995. MR Zbl

[Grillakis 1990] M. G. Grillakis, “Regularity and asymptotic behaviour of the wave equation with a critical nonlinearity”, *Ann. of Math.* (2) **132**:3 (1990), 485–509. MR Zbl

[Grillakis 1992] M. G. Grillakis, “Regularity for the wave equation with a critical nonlinearity”, *Comm. Pure Appl. Math.* **45**:6 (1992), 749–774. MR Zbl

[Keel and Tao 1998] M. Keel and T. Tao, “Endpoint Strichartz estimates”, *Amer. J. Math.* **120**:5 (1998), 955–980. MR Zbl

[Kenig 2015] C. E. Kenig, *Lectures on the energy critical nonlinear wave equation*, CBMS Regional Conference Series in Mathematics **122**, American Mathematical Society, Providence, RI, 2015. MR Zbl

[Kenig and Merle 2006] C. E. Kenig and F. Merle, “Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case”, *Invent. Math.* **166**:3 (2006), 645–675. MR Zbl

[Kenig and Merle 2008] C. E. Kenig and F. Merle, “Global well-posedness, scattering and blow-up for the energy-critical focusing non-linear wave equation”, *Acta Math.* **201**:2 (2008), 147–212. MR Zbl

[Killip and Vişan 2013] R. Killip and M. Vişan, “Nonlinear Schrödinger equations at critical regularity”, pp. 325–437 in *Evolution equations*, edited by D. Ellwood et al., Clay Math. Proc. **17**, American Mathematical Society, Providence, RI, 2013. MR Zbl

[Killip and Visan 2011] R. Killip and M. Visan, “The defocusing energy-supercritical nonlinear wave equation in three space dimensions”, *Trans. Amer. Math. Soc.* **363**:7 (2011), 3893–3934. MR Zbl

[Koch et al. 2014] H. Koch, D. Tataru, and M. Vişan, *Dispersive equations and nonlinear waves: generalized Korteweg–de Vries, nonlinear Schrödinger, wave and Schrödinger maps*, Oberwolfach Seminars **45**, Springer, 2014. MR Zbl

[Lindblad and Sogge 1995] H. Lindblad and C. D. Sogge, “On existence and scattering with minimal regularity for semilinear wave equations”, *J. Funct. Anal.* **130**:2 (1995), 357–426. MR Zbl

[Lindblad and Sogge 1996] H. Lindblad and C. D. Sogge, “Long-time existence for small amplitude semilinear wave equations”, *Amer. J. Math.* **118**:5 (1996), 1047–1135. MR Zbl

[Morawetz 1968] C. S. Morawetz, “Time decay for the nonlinear Klein–Gordon equations”, *Proc. Roy. Soc. London Ser. A* **306** (1968), 291–296. MR Zbl

[Nakanishi 1999] K. Nakanishi, “Unique global existence and asymptotic behaviour of solutions for wave equations with non-coercive critical nonlinearity”, *Comm. Partial Differential Equations* **24**:1–2 (1999), 185–221. MR Zbl

[Rammaha 1987] M. A. Rammaha, “Finite-time blow-up for nonlinear wave equations in high dimensions”, *Comm. Partial Differential Equations* **12**:6 (1987), 677–700. MR Zbl

[Ramos 2012] J. Ramos, “A refinement of the Strichartz inequality for the wave equation with applications”, *Adv. Math.* **230**:2 (2012), 649–698. MR Zbl

[Rodriguez 2017] C. Rodriguez, “Scattering for radial energy-subcritical wave equations in dimensions 4 and 5”, *Comm. Partial Differential Equations* **42**:6 (2017), 852–894. MR Zbl

[Shatah and Struwe 1993] J. Shatah and M. Struwe, “Regularity results for nonlinear wave equations”, *Ann. of Math.* (2) **138**:3 (1993), 503–518. MR Zbl

[Sogge 1995] C. D. Sogge, *Lectures on nonlinear wave equations*, Monographs in Analysis **II**, International Press, Boston, 1995. MR Zbl

[Taggart 2010] R. J. Taggart, “Inhomogeneous Strichartz estimates”, *Forum Math.* **22**:5 (2010), 825–853. MR Zbl

Received June 25, 2018.

CHANGXING MIAO

INSTITUTE OF APPLIED PHYSICS AND COMPUTATIONAL MATHEMATICS

BEIJING

CHINA

miao\_changxing@iapcm.ac.cn

JIANWEI YANG

BEIJING INSTITUTE OF TECHNOLOGY

BEIJING

CHINA

jw-urbain.yang@bit.edu.cn

TENGFEI ZHAO

BEIJING COMPUTATIONAL SCIENCE RESEARCH CENTER

BEIJING

CHINA

zhao\_tengfei@csrc.ac.cn

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
blasius@math.ucla.edu

Matthias Aschenbrenner  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
matthias@math.ucla.edu

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
cooper@math.ucsb.edu

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
jhu@maths.hku.hk

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
balmer@math.ucla.edu

Wee Teck Gan  
Mathematics Department  
National University of Singapore  
Singapore 119076  
matgwt@nus.edu.sg

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
popa@math.ucla.edu

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
yang@math.princeton.edu

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
chari@math.ucr.edu

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
liu@math.ucla.edu

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
qing@cats.ucsc.edu

PRODUCTION  
Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

---

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 305 No. 1 March 2020

---

The Poincaré homology sphere, lens space surgeries, and some knots with tunnel number two	1
KENNETH L. BAKER	
Fusion systems of blocks of finite groups over arbitrary fields	29
ROBERT BOLTJE, ÇİSİL KARAGÜZEL and DENİZ YILMAZ	
Torsion points and Galois representations on CM elliptic curves	43
ABBEY BOURDON and PETE L. CLARK	
Stability of the positive mass theorem for axisymmetric manifolds	89
EDWARD T. BRYDEN	
Index estimates for free boundary constant mean curvature surfaces	153
MARCOS P. CAVALCANTE and DARLAN F. DE OLIVEIRA	
A criterion for modules over Gorenstein local rings to have rational Poincaré series	165
ANJAN GUPTA	
Generalized Cartan matrices arising from new derivation Lie algebras of isolated hypersurface singularities	189
NAVEED HUSSAIN, STEPHEN S.-T. YAU and HUAIQING ZUO	
On the commutativity of coset pressure	219
BING LI and WEN-CHIAO CHENG	
Signature invariants related to the unknotting number	229
CHARLES LIVINGSTON	
The global well-posedness and scattering for the 5-dimensional defocusing conformal invariant NLW with radial initial data in a critical Besov space	251
CHANGXING MIAO, JIANWEI YANG and TENGFEI ZHAO	
Liouville-type theorems for weighted $p$ -harmonic 1-forms and weighted $p$ -harmonic maps	291
KEOMKYO SEO and GABJIN YUN	
Remarks on the Hölder-continuity of solutions to parabolic equations with conic singularities	311
YUANQI WANG	
Deformation of Milnor algebras	329
ZHENJIAN WANG	
Preservation of log-Sobolev inequalities under some Hamiltonian flows	339
BO XIA	
Ground state solutions of polyharmonic equations with potentials of positive low bound	353
CAIFENG ZHANG, JUNGANG LI and LU CHEN	



0030-8730(2020)305:1;1-R