

Pacific Journal of Mathematics

DEFORMATION OF MILNOR ALGEBRAS

ZHENJIAN WANG

DEFORMATION OF MILNOR ALGEBRAS

ZHENJIAN WANG

We investigate deformations of Milnor algebras of smooth homogeneous polynomials, and prove in particular that any smooth degree d homogeneous polynomial in $n + 1$ variables that is not of Sebastiani–Thom type is determined by the degree k homogeneous component of its Jacobian ideal for any $d - 1 \leq k \leq (n + 1)(d - 2)$. Our results generalize the previous result on the reconstruction of a homogeneous polynomial from its Jacobian ideal.

1. Introduction

The classical theory of variation of Hodge structures for smooth hypersurfaces in a complex projective space gives a variation of Milnor algebras of homogeneous polynomials. The celebrated generic Torelli theorem for hypersurfaces is almost reduced to the study of injectivity of some mappings concerning the deformation of Milnor algebras, see [Voisin 2003, Subsection 6.3.2, p. 179; Donagi 1983]. Nevertheless, the homogeneous components of the Milnor algebra involved there are of specific degrees, namely degrees of the form $pd - n - 1$. In this note, we will investigate homogeneous components of all degrees of the Milnor algebra.

To fix notation, let $S = \mathbb{C}[x_0, \dots, x_n]$ be the homogeneous coordinate ring of the complex projective space $\mathbb{P}^n$,

$$S = \bigoplus_{d=0}^{\infty} S_d,$$

where S_d is the vector space of homogeneous polynomials of degree d . Given a homogeneous polynomial $f \in S_d$, denote by $J(f)$ the Jacobian ideal of f :

$$J(f) = (\partial f / \partial x_0, \partial f / \partial x_1, \dots, \partial f / \partial x_n).$$

Set $M(f) = S/J(f)$, known as the Milnor algebra of f . The algebra $M(f)$ has the natural grading

$$M(f) = \bigoplus_{k=0}^{\infty} M(f)_k,$$

MSC2010: primary 14A25; secondary 14C34, 14J70.

Keywords: Homogeneous polynomials, Milnor algebras, Macaulay inverse system.

where $M(f)_k = S_k/(J(f) \cap S_k)$.

We say that $f \in \mathbb{P}(S_d)$ is a *smooth* polynomial if the hypersurface $V_f : f = 0$ in $\mathbb{P}^n$ is a smooth hypersurface. The discriminant defines a divisor $\Delta \subset \mathbb{P}(S_d)$ such that the complement $\mathbb{P}(S_d)_\Delta$ parametrizes smooth homogeneous polynomials of degree d .

We say that a polynomial $f \in S_d$ is of *Sebastiani–Thom type* (ST type) or a *direct sum* if f can be represented as

$$(1) \quad f(x_0, \dots, x_n) = f_1(x_0, \dots, x_\ell) + f_2(x_{\ell+1}, \dots, x_n)$$

for a choice of homogeneous coordinates $\{x_i\}_{i=0}^n$ of $\mathbb{P}^n$ and some $0 \leq \ell < n$; see [Ueda and Yoshinaga 2009; Wang 2015; Buczyńska et al. 2015]. For various characterizations of polynomials of ST type, we refer to [Fedorchuk 2020]. Denote by $\mathcal{U} \subset \mathbb{P}(S_d)_\Delta$ the set of all smooth homogeneous polynomials that are *not* of ST type.

It is well-known that $\dim M(f)_k$ does not depend on the concrete equation of f for smooth f (see for instance [Dimca 1987, Proposition 7.22, p. 108]); we denote this dimension by $a_{n,d}(k)$. Let $\text{Grass}(S_k, a_{n,d}(k))$ be the Grassmannian parametrizing all $a_{n,d}(k)$ dimensional *quotient spaces* of S_k , then we have the following map

$$(2) \quad \varphi_k : \mathbb{P}(S_d)_\Delta \rightarrow \text{Grass}(S_k, a_{n,d}(k)),$$

defined by $f \mapsto M(f)_k$.

More generally, denote $\text{Grass}(n+1, S_{d-1})$ the Grassmannian of *linear subspaces* of dimension $n+1$ of the space of degree $d-1$ homogeneous polynomials S_{d-1} . Following [Fedorchuk 2017, Subsection 1.2], given $W \in \text{Grass}(n+1, S_{d-1})$, we form the ideal $I_W := (W)$ and the quotient algebra $M(W) = S/I_W$. Let $g_0, \dots, g_n$ be a basis of W , then the sequence $g_0, \dots, g_n$ is a regular sequence if and only if I_W is a complete intersection ideal if and only if $M(W)$ is a standard local Artinian Gorenstein algebra of socle degree $T := (n+1)(d-2)$ if and only if the resultant of $g_0, \dots, g_n$ is nonzero; we refer to [Gelfand et al. 1994, Chapter 13] for the definition and basic properties of the resultant. Therefore, there exists a divisor $\text{Res} \subset \text{Grass}(n+1, S_{d-1})$ parametrizing all W such that I_W is not a complete intersection ideal. We denote by $\text{Grass}(n+1, S_{d-1})_{\text{Res}}$ the affine complement of Res . For more discussions about ideals of the form I_W , see [Fedorchuk 2017, Subsection 1.2].

For $W \in \text{Grass}(n+1, S_{d-1})_{\text{Res}}$, we have $\dim M(W)_k = a_{n,d}(k)$ by [Dimca 1987, Proposition 7.22, p. 108]. Hence the assignment $W \mapsto M(W)_k$ defines a map

$$(3) \quad \Phi_k : \text{Grass}(n+1, S_{d-1})_{\text{Res}} \rightarrow \text{Grass}(S_k, a_{n,d}(k)).$$

Our first result is the following theorem.

Theorem 1.1. *For any $d - 1 \leq k \leq T = (n + 1)(d - 2)$, the map Φ_k is an immersion, that is, it is injective and the differential $d\Phi_k$ is also injective at any point of $\text{Grass}(n + 1, S_{d-1})_{\text{Res}}$.*

Using our previous result on determination of a polynomial by its Jacobian ideal (see [Wang 2015, Theorem 1.1; Ueda and Yoshinaga 2009, Lemma 3]), we further prove the following result.

Theorem 1.2. *For $d - 1 \leq k \leq T = (n + 1)(d - 2)$, the restriction of the map φ_k (defined in (2)) to $\mathcal{U}$,*

$$\varphi_k : \mathcal{U} \rightarrow \text{Grass}(S_k, a_{n,d}(k)),$$

is an immersion.

In particular, we have that a smooth homogeneous polynomial $f \in \mathcal{U}$ can be reconstructed from the degree k homogeneous component of its Jacobian ideal $J(f)$ for any k satisfying $d - 1 \leq k \leq T = (n + 1)(d - 2)$. This gives a generalization of the previous results, in the case of smooth polynomials, in [Wang 2015; Ueda and Yoshinaga 2009].

We will also investigate the map φ_k defined in (2) and discuss its fibers over $\varphi_k(f)$ for homogeneous polynomials f that are of ST type, see Section 4.

Our results are related to the problem of characterizing the hypersurface singularity $\widehat{V}_f = \{x \in \mathbb{C}^{n+1} \mid f(x) = 0\}$ at the origin 0 of $\mathbb{C}^{n+1}$ using the Milnor algebra $M(f)$. In fact, the characterization problem of a singularity by its algebraic data can be proposed and solved in a much more general setting, see [Gaffney and Hauser 1985]. As a general philosophy in singularity theory, the Milnor algebra $M(f)$ is closely connected to the topology and geometry of the hypersurface singularity $(\widehat{V}_f, 0) \subset (\mathbb{C}^{n+1}, 0)$. Instead of giving characterizations of a singularity by *algebras* or *modules* derived from it, as in [Gaffney and Hauser 1985], here using Theorem 1.2, we can give a characterization of the *isolated* hypersurface singularity $(\widehat{V}_f, 0)$ just by a single homogeneous component $M(f)_k$ of the Milnor algebra $M(f)$ for any $d - 1 \leq k \leq (n + 1)(d - 2)$ with $d = \deg f$. This conclusion can obviously be extended to an isolated complete intersection singularity by using Theorem 1.1.

Of course, our results concern only the case when the hypersurface $(\widehat{V}_f, 0)$ is an isolated singularity. It is natural to extend these results to the case where the singularities of the hypersurface $\widehat{V}_f$ have positive dimension. However, the tools used in this note cannot be directly applied in the extended case because they depend heavily on the condition that $M(f)$ is a local Artinian Gorenstein algebra which holds only when 0 is an isolated singularity of $\widehat{V}_f$. In addition, heuristically, the results in [Gaffney and Hauser 1985] also show that any possible extension must

be more complicated and more technical than our results above; see also the results in [Wang 2015] concerning the nonsmooth homogeneous polynomials.

We hope the results in this note can be applied to the study of Lefschetz properties for Milnor algebras. In fact, this is an important impetus to our present work. As is well-known, the strong Lefschetz property holds for $M(f)$ for a generic f . Our naïve idea is to investigate the Lefschetz properties by deforming the Milnor algebras. For an excellent exposition for the Lefschetz properties, we refer to [Harima et al. 2003; Migliore and Nagel 2013; Harima et al. 2013]. In addition, the strong Lefschetz property for $M(f)$ where f is of ST type can be reduced to that where f is not of ST type (see [Harima and Watanabe 2007, Theorem 3.10; Harima et al. 2013, Proposition 3.77, p. 137]), since $M(f)$ is the tensor product of $M(f_1)$ and $M(f_2)$ when f is represented as in (1). This is an important reason why we specifically investigate the set $\mathcal{U}$ in this paper; another reason is about the determination of a homogeneous polynomial by its Jacobian ideal, see the proof of Corollary 3.2 below.

2. Polar paring and Macaulay inverse systems

2.1. Polar pairing. Let $S = \mathbb{C}[x_0, \dots, x_n]$ and $R = \mathbb{C}[z_0, \dots, z_n]$ be two polynomial rings. There is a natural action of S on R by the “polar paring”

$$S \times R \rightarrow R$$

defined by

$$(f(x_0, \dots, x_n), Q(z_0, \dots, z_n)) \mapsto f \cdot Q := f(\partial/\partial z_0, \dots, \partial/\partial z_n)Q(z_0, \dots, z_n).$$

It induces perfect parings $S_\rho \times R_\rho \rightarrow \mathbb{C}$ for every $\rho \in \mathbb{N}$. In particular, for $f \in S_\rho$ written as

$$f(x_1, \dots, x_n) = \sum_{|\alpha|=\rho} a_\alpha x^\alpha$$

and $Q \in R_\rho$ written as

$$Q(z_0, \dots, z_n) = \sum_{|\alpha|=\rho} b_\alpha z^\alpha$$

we have

$$f \cdot Q = \sum_{|\alpha|=\rho} \alpha! a_\alpha b_\alpha.$$

Define the polynomial $q = \sum_{|\alpha|=\rho} b_\alpha x^\alpha$, or equivalently

$$q(x_0, \dots, x_n) = Q(x_0, \dots, x_n),$$

and define the inner product of f and q by

$$(4) \quad \langle f, q \rangle = \sum_{|\alpha|=\rho} \alpha! a_\alpha b_\alpha = f \cdot Q.$$

For any linear space $E \subset S_\rho$, with respect to the above inner product $\langle \cdot, \cdot \rangle$, we have its orthogonal complement, denoted by $E^\perp$.

2.2. Macaulay inverse system. Let $I \subset S$ be a Gorenstein ideal and ν the socle degree of the algebra $\mathcal{A} = S/I$. Recall that a (homogeneous) Macaulay inverse system of $\mathcal{A}$ is an element $Q_{\mathcal{A}} \in \mathbb{P}(R_\nu)$ such that I is equal to the apolar ideal $Q_{\mathcal{A}}^\perp$, namely,

$$I = \{f \in S \mid f \cdot Q_{\mathcal{A}} = 0\}$$

(see [Iarrobino and Kanev 1999, Lemma 2.12] or [Eisenbud 1995, Exercise 2.17]).

Let $W = \text{span}\langle g_0, \dots, g_n \rangle$ such that $W \in \text{Grass}(n+1, S_{d-1})_{\text{Res}}$, the associated form $A_W := A(g_0, \dots, g_n) \in \mathbb{P}(R_T)$ (recall that $T = (n+1)(d-2)$) gives the Macaulay inverse system for $S_W = S/I_W$; see [Alper and Isaev 2018, Proposition 2.1]. We write

$$A_W = \sum_{|\alpha|=T} c_\alpha z^\alpha.$$

In this case, define $B_W \in \mathbb{P}(S_T)$ by

$$B_W = \sum_{|\alpha|=T} c_\alpha x^\alpha.$$

The polynomial B_W , by definition, determines and is determined by A_W . Moreover, by the definition of Macaulay inverse systems, we have that $(I_W)_T^\perp = \mathbb{C}B_W$, namely, the line $\mathbb{C}B_W$ is exactly the orthogonal complement of $(I_W)_T$ with respect to the inner product $\langle \cdot, \cdot \rangle$ on S_T . Therefore, $A_W \in \mathbb{P}(R_T)$ is uniquely determined by $(I_W)_T$.

Lemma 2.3. *For two points $U, W \in \text{Grass}(n+1, S_{d-1})_{\text{Res}}$, the following statements are equivalent:*

- (1) $U = W$.
- (2) $I_U = I_W$.
- (3) For any k satisfying $d-1 \leq k \leq T = (n+1)(d-2)$, we have $(I_U)_k = (I_W)_k$.
- (4) For some k satisfying $d-1 \leq k \leq T$, we have $(I_U)_k = (I_W)_k$.
- (5) $(I_U)_T = (I_W)_T$.

Proof. It is obvious that (1), (2), (3) are all equivalent and (3) implies (4).

(4) $\Rightarrow$ (5): Since I_U is generated by polynomials all of which have degree $d - 1$, we have that $(I_U)_T$ is the image of $S_{T-k} \times (I_U)_k$ under the multiplication map $S_{T-k} \times S_k \rightarrow S_T$. Hence $(I_U)_T = (I_W)_T$ whenever $(I_U)_k = (I_V)_k$ for $d - 1 \leq k \leq T$.

(5) $\Rightarrow$ (1): This is clear once we note that A_U can be uniquely determined by $(I_U)_T$, and I_U is the apolar ideal $A_U^\perp$. $\square$

Recall that as it is shown in the introduction, $a_{n,d}(k) = \dim M(f)_k$ for any $f \in \mathbb{P}(S_d)_\Delta$, which is also the dimension of $S_k/(I_W)_k$ for any $W \in \text{Grass}(n+1, S_{d-1})_{\text{Res}}$. Set $b_{n,d}(k) = \dim S_k - a_{n,d}(k)$ and let $\text{Grass}(b_{n,d}(k), S_k)$ be the Grassmannian parametrizing all $b_{n,d}(k)$ dimensional linear *subspaces* of S_k . For a subspace $E \subset S_k$ of dimension $b_{n,d}(k)$, we obtain the quotient space S_k/E of dimension $a_{n,d}(k)$; and the mapping $S_k \supset E \mapsto S_k/E$ clearly defines an isomorphism between the Grassmannians $\text{Grass}(b_{n,d}(k), S_k)$ and $\text{Grass}(S_k, a_{n,d}(k))$. Then to prove [Theorem 1.1](#), it suffices to prove the following theorem.

Theorem 2.4. *For any $d - 1 \leq k \leq T$, the assignment $W \mapsto (I_W)_k$ defines an immersion*

$$(5) \quad \Psi_k : \text{Grass}(n+1, S_{d-1})_{\text{Res}} \rightarrow \text{Grass}(b_{n,d}(k), S_k),$$

that is Ψ_k is injective and the differential $d\Psi_k$ is also injective at any point of $\text{Grass}(n+1, S_{d-1})_{\text{Res}}$.

Proof. The injectivity of Ψ_k follows from the equivalence (1) $\Leftrightarrow$ (4) in [Lemma 2.3](#).

Given $W \in \text{Grass}(n+1, S_{d-1})_{\text{Res}}$ such that $W = \text{span}\langle g_0, \dots, g_n \rangle$. For any $h \in T_W(\text{Grass}(n+1, S_{d-1})_{\text{Res}}) \simeq \text{Hom}(W, S_{d-1}/W)$, choose $h_i \in S_{d-1}$ such that $h(g_i) = h_i \pmod{W}$ for $i = 0, \dots, n$. Then if $h \in \text{Ker}(d\Psi_k)_W$, we have $(d\Psi_k)_W(h) = 0$ as an element in $\text{Hom}((I_W)_k, S_k/(I_W)_k)$. A direct computation gives that

$$(d\Psi_k)_W(h)((I_W)_k) = (I_H)_k + (I_W)_k \pmod{(I_W)_k},$$

where $H = \text{span}\langle h_0, \dots, h_n \rangle$. It follows from $(d\Psi_k)_W(h) = 0$ that $(I_H)_k \subset (I_W)_k$.

For $t \in \mathbb{C}^*$ and $|t|$ sufficiently small, we have that

$$W_t := \text{span}\langle g_0 + th_0, \dots, g_n + h_n \rangle$$

satisfies $W_t \in \text{Grass}(n+1, S_{d-1})_{\text{Res}}$. It then follows from $(I_H)_k \subset (I_W)_k$ that $(I_{W_t})_k \subset (I_W)_k$, hence $(I_{W_t})_k = (I_W)_k$ because $\dim(I_{W_t})_k = b_{n,d}(k) = \dim(I_W)_k$. Therefore $W_t = W$ by (4) $\Rightarrow$ (1) in [Lemma 2.3](#). It follows that $h_i \in W$ for $i = 0, \dots, n$ and thus $h = 0$ as an element of $T_W(\text{Grass}(n+1, S_{d-1})_{\text{Res}})$.

Since h can be arbitrarily chosen, $(d\Psi_k)_W$ is injective. We are done. $\square$

3. Variation of Milnor algebras

3.1. Polynomials not of ST type. Recall that $\mathcal{U} \subset \mathbb{P}(S_d)$ denotes the space of smooth homogeneous polynomials of degree d that are not of ST type, or equivalently, the space of smooth hypersurfaces whose defining equations are not of ST type. From the proof of [Wang 2015, Corollary 6.1], we have that $\mathcal{U}$ is a Zariski open subset of $\mathbb{P}(S_d)_\Delta$.

For $f \in \mathcal{U}$, recall that $J(f)$ denotes the Jacobian ideal of f and $M(f) = S/J(f)$ the Milnor algebra. For $k \geq d - 1$, we denote by $E_k(f) = J(f) \cap S_k$. Then $\dim E_k(f) = b_{n,d}(k) = \dim S_k - a_{n,d}(k)$ is independent of $f \in \mathcal{U}$. Moreover, since $\partial f / \partial x_0, \dots, \partial f / \partial x_n$ form a regular sequence and $J(f) = I_{E_{d-1}(f)}$, from Lemma 2.3, we immediately get the following corollary.

Corollary 3.2. *Given $f, g \in \mathbb{P}(S_d)$ and $f \in \mathcal{U}$, the following conditions are equivalent:*

- (1) $E_{d-1}(f) = E_{d-1}(g)$.
- (2) $J(f) = J(g)$.
- (3) For any k satisfying $d - 1 \leq k \leq T = (n + 1)(d - 2)$, we have $E_k(f) = E_k(g)$.
- (4) For some k satisfying $d - 1 \leq k \leq T$, we have $E_k(f) = E_k(g)$.
- (5) $E_T(f) = E_T(g)$.
- (6) $f = g$.

Proof. The equivalences among the first five statements follow from Lemma 2.3; we here just note that any one of these conditions implies that $E_{T+1}(g) = S_{T+1}$, hence g is also smooth and thus $J(g)$ is a complete intersection ideal.

The equivalence (1) $\Leftrightarrow$ (6) follows from [Wang 2015, Theorem 1.1] or [Ueda and Yoshinaga 2009, Lemma 3]. $\square$

Now we are ready to prove Theorem 1.2. Similar to the proof of Theorem 1.1, it is sufficient to prove the following theorem.

Theorem 3.3. *For any $d - 1 \leq k \leq T$, the assignment $f \mapsto E_k(f)$ defines an immersion*

$$\psi_k : \mathcal{U} \rightarrow \text{Grass}(b_{n,d}(k), S_k),$$

Namely, ψ_k is injective and its differential $d\psi_k$ is also injective at any point $f \in \mathcal{U}$.

Proof. By the equivalence of (4) and (6) in Corollary 3.2, we have that ψ_k is injective.

We will not distinguish an element $f \in \mathbb{P}(S_d)$ and its lifting in S_d . For $f \in \mathcal{U}$, we have $T_f \mathcal{U} = T_f \mathbb{P}(S_d) \simeq \text{Hom}(\mathbb{C}f, S_d/\mathbb{C}f)$. The mapping $\text{Hom}(\mathbb{C}f, S_d/\mathbb{C}f) \ni$

$\eta \mapsto \eta(f) \in S_d/\mathbb{C}f$ then gives an identification $T_f\mathcal{U} \simeq S_d/\mathbb{C}f$. With the help of this identification, the differential of ψ_k at $f \in \mathcal{U}$ is given by

$$(d\psi_k)_f : T_f\mathcal{U} \simeq S_d/\mathbb{C}f \rightarrow \text{Hom}(E_k(f), S_k/E_k(f)).$$

Therefore, we have $(d\psi_k)_f(h) = 0$ as an element of $\text{Hom}(E_k(f), S_k/E_k(f))$ for any $h \in \text{Ker}(d\psi_k)_f$. Represent h by an element in $S_d/\mathbb{C}f$, and lift it to an element in S_d which is still denoted by h . A direct computation gives that

$$(d\psi_k)_f(h)(E_k(f)) = E_k(h) + E_k(f) \pmod{E_k(f)}.$$

Hence it follows from $(d\psi_k)_f(h) = 0$ that $E_k(h) \subset E_k(f)$.

From the semicontinuity of the dimension of $E_k(f)$ with respect to $f \in S_d$, we obtain that for a small positive number $\epsilon > 0$ and for any $t \in \mathbb{C}$ such that $|t| < \epsilon$, the following hold:

$$(i) \dim E_k(f + th) = b_{n,d}(k) = \dim E_k(f);$$

$$(ii) E_k(f + th) \subset E_k(f).$$

Hence $E_k(f + th) = E_k(f)$ for any $|t| < \epsilon$. In particular, choosing $t_0 \neq 0$ satisfying $|t_0| < \epsilon$, we have $E_k(f + t_0h) = E_k(f)$. Using (4) $\Leftrightarrow$ (6) in [Corollary 3.2](#) again, we deduce that $f + t_0h = f$ in $\mathbb{P}(S_d)$, hence $h = f$ in $\mathbb{P}(S_d)$ which implies that the chosen tangent vector $h \in \text{Ker}(d\psi_k)_f$ is equal to zero. Therefore $(d\psi_k)_f$ is also injective. $\square$

The above proof also gives the following corollary, which is interesting in its own right; compare with [Corollary 3.2](#).

Corollary 3.4. *Given $f \in \mathcal{U}$ and $h \in \mathbb{P}(S_d)$. Suppose $E_k(h) \subset E_k(f)$ for some $d-1 \leq k \leq T$, then $h = f$.*

4. Polynomials of Sebastiani–Thom type

In this section, we give a brief discussion about the fibers of the map φ_k in (2) over $\varphi_k(f)$ for a polynomial f of ST type.

By [[Fedorchuk 2020](#), Proposition 4.8 or Corollary 3.15], a smooth homogeneous polynomial $f \in S_d$ admits a unique maximally fine “direct sum decomposition”

$$(6) \quad f(x_0, \dots, x_n) = f_1(x_0, \dots, x_{n_1-1}) + f_2(x_{n_1}, \dots, x_{n_2}) + \dots + f_s(x_{n_{s-1}}, \dots, x_n),$$

for a choice of linear coordinates $\{x_i\}_{i=0}^n$, where $0 \leq n_1 \leq n_2 \leq \dots \leq n_{s-1} \leq n$ and none of the f_j are of ST type. In addition, if $g \in S_d$ satisfies $E_{d-1}(g) \subset E_{d-1}(f)$, then necessarily, g is of the following form

$$(7) \quad g = \lambda_1 f_1 + \dots + \lambda_s f_s, \quad \lambda_i \in \mathbb{C},$$

see [Fedorchuk 2020, Corollary 3.12]. In particular, if g is also smooth, then all the λ_j in (7) are nonzero. With these results at hand, we prove the following theorem.

Theorem 4.1. *For any $d - 1 \leq k \leq T = (n + 1)(d - 1)$ and any $f \in \mathbb{P}(S_d)_\Delta$, the fiber over $\varphi_k(f)$ of φ_k defined in (2), namely,*

$$\varphi_k : \mathbb{P}(S_d)_\Delta \rightarrow \text{Grass}(S_k, a_{n,d}(k)),$$

is

$$\varphi_k^{-1}(\varphi_k(f)) = \{\lambda_1 f_1 + \cdots + \lambda_s f_s \mid \lambda_i \in \mathbb{C}^*, i = 1, \dots, s\}.$$

Proof. It is obvious that for the λ_j nonzero, the polynomial $\lambda_1 f_1 + \cdots + \lambda_s f_s$ is smooth and is mapped under φ_k to $\varphi_k(f)$.

Conversely, if $g \in \mathbb{P}(S_d)_\Delta$ satisfies $\varphi_k(g) = \varphi_k(f)$, then we have $E_k(g) = E_k(f)$. It follows by (4) $\Rightarrow$ (1) in Lemma 2.3 that $E_{d-1}(g) = E_{d-1}(f)$. Hence by [Fedorchuk 2020, Corollary 3.12], we have that g is of the form $\lambda_1 f_1 + \cdots + \lambda_s f_s$ for nonzero λ_j 's. $\square$

In conclusion, for the map φ_k , we can explicitly and completely determine all the fibers.

Acknowledgements

We would like to thank Professor Herwig Hauser for the reference [Gaffney and Hauser 1985]. We thank an anonymous referee for useful remarks and suggestions. We also thank the Yau Mathematical Sciences Center for financial support and stimulating working atmosphere.

References

- [Alper and Isaev 2018] J. Alper and A. Isaev, “Associated forms and hypersurface singularities: the binary case”, *J. Reine Angew. Math.* **745** (2018), 83–104. [MR](#) [Zbl](#)
- [Buczyńska et al. 2015] W. Buczyńska, J. Buczyński, J. Kleppe, and Z. Teitler, “Apolarity and direct sum decomposability of polynomials”, *Michigan Math. J.* **64**:4 (2015), 675–719. [MR](#) [Zbl](#)
- [Dimca 1987] A. Dimca, *Topics on real and complex singularities: an introduction*, Vieweg, Braunschweig, 1987. [MR](#) [Zbl](#)
- [Donagi 1983] R. Donagi, “Generic Torelli for projective hypersurfaces”, *Compositio Math.* **50**:2-3 (1983), 325–353. [MR](#) [Zbl](#)
- [Eisenbud 1995] D. Eisenbud, *Commutative algebra: with a view toward algebraic geometry*, Graduate Texts in Mathematics **150**, Springer, 1995. [MR](#) [Zbl](#)
- [Fedorchuk 2017] M. Fedorchuk, “GIT semistability of Hilbert points of Milnor algebras”, *Math. Ann.* **367**:1-2 (2017), 441–460. [MR](#) [Zbl](#)
- [Fedorchuk 2020] M. Fedorchuk, “Direct sum decomposability of polynomials and factorization of associated forms”, *Proc. Lond. Math. Soc.* (3) **120**:3 (2020), 305–327. [MR](#)
- [Gaffney and Hauser 1985] T. Gaffney and H. Hauser, “Characterizing singularities of varieties and of mappings”, *Invent. Math.* **81**:3 (1985), 427–447. [MR](#) [Zbl](#)

- [Gelfand et al. 1994] I. M. Gelfand, M. M. Kapranov, and A. V. Zelevinsky, *Discriminants, resultants, and multidimensional determinants*, Birkhäuser, Boston, 1994. [MR](#) [Zbl](#)
- [Harima and Watanabe 2007] T. Harima and J. Watanabe, “The strong Lefschetz property for Artinian algebras with non-standard grading”, *J. Algebra* **311**:2 (2007), 511–537. [MR](#) [Zbl](#)
- [Harima et al. 2003] T. Harima, J. C. Migliore, U. Nagel, and J. Watanabe, “The weak and strong Lefschetz properties for Artinian K -algebras”, *J. Algebra* **262**:1 (2003), 99–126. [MR](#) [Zbl](#)
- [Harima et al. 2013] T. Harima, T. Maeno, H. Morita, Y. Numata, A. Wachi, and J. Watanabe, *The Lefschetz properties*, Lecture Notes in Mathematics **2080**, Springer, 2013. [MR](#) [Zbl](#)
- [Iarrobino and Kanev 1999] A. Iarrobino and V. Kanev, *Power sums, Gorenstein algebras, and determinantal loci*, Lecture Notes in Mathematics **1721**, Springer, 1999. [MR](#) [Zbl](#)
- [Migliore and Nagel 2013] J. Migliore and U. Nagel, “Survey article: a tour of the weak and strong Lefschetz properties”, *J. Commut. Algebra* **5**:3 (2013), 329–358. [MR](#) [Zbl](#)
- [Ueda and Yoshinaga 2009] K. Ueda and M. Yoshinaga, “Logarithmic vector fields along smooth divisors in projective spaces”, *Hokkaido Math. J.* **38**:3 (2009), 409–415. [MR](#) [Zbl](#)
- [Voisin 2003] C. Voisin, *Hodge theory and complex algebraic geometry*, vol. II, Cambridge Studies in Advanced Mathematics **77**, Cambridge University Press, 2003. [MR](#) [Zbl](#)
- [Wang 2015] Z. Wang, “On homogeneous polynomials determined by their Jacobian ideal”, *Manuscripta Math.* **146**:3-4 (2015), 559–574. [MR](#) [Zbl](#)

Received July 17, 2019. Revised October 4, 2019.

ZHENJIAN WANG
YAU MATHEMATICAL SCIENCES CENTER
TSINGHUA UNIVERSITY
BEIJING
CHINA
wzhj@mail.tsinghua.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 305 No. 1 March 2020

The Poincaré homology sphere, lens space surgeries, and some knots with tunnel number two	1
KENNETH L. BAKER	
Fusion systems of blocks of finite groups over arbitrary fields	29
ROBERT BOLTJE, ÇISIL KARAGÜZEL and DENİZ YILMAZ	
Torsion points and Galois representations on CM elliptic curves	43
ABBEY BOURDON and PETE L. CLARK	
Stability of the positive mass theorem for axisymmetric manifolds	89
EDWARD T. BRYDEN	
Index estimates for free boundary constant mean curvature surfaces	153
MARCOS P. CAVALCANTE and DARLAN F. DE OLIVEIRA	
A criterion for modules over Gorenstein local rings to have rational Poincaré series	165
ANJAN GUPTA	
Generalized Cartan matrices arising from new derivation Lie algebras of isolated hypersurface singularities	189
NAVEED HUSSAIN, STEPHEN S.-T. YAU and HUAIQING ZUO	
On the commutativity of coset pressure	219
BING LI and WEN-CHIAO CHENG	
Signature invariants related to the unknotting number	229
CHARLES LIVINGSTON	
The global well-posedness and scattering for the 5-dimensional defocusing conformal invariant NLW with radial initial data in a critical Besov space	251
CHANGXING MIAO, JIANWEI YANG and TENGFEI ZHAO	
Liouville-type theorems for weighted p -harmonic 1-forms and weighted p -harmonic maps	291
KEOMKYO SEO and GABJIN YUN	
Remarks on the Hölder-continuity of solutions to parabolic equations with conic singularities	311
YUANQI WANG	
Deformation of Milnor algebras	329
ZHENJIAN WANG	
Preservation of log-Sobolev inequalities under some Hamiltonian flows	339
BO XIA	
Ground state solutions of polyharmonic equations with potentials of positive low bound	353
CAIFENG ZHANG, JUNGANG LI and LU CHEN	