

*Pacific
Journal of
Mathematics*

**ON THE NONEXISTENCE OF
 S^6 TYPE COMPLEX THREEFOLDS
IN ANY COMPACT HOMOGENEOUS COMPLEX MANIFOLDS
WITH THE COMPACT LIE GROUP G_2
AS THE BASE MANIFOLD**

DANIEL GUAN

**ON THE NONEXISTENCE OF
 S^6 TYPE COMPLEX THREEFOLDS
IN ANY COMPACT HOMOGENEOUS COMPLEX MANIFOLDS
WITH THE COMPACT LIE GROUP G_2
AS THE BASE MANIFOLD**

DANIEL GUAN

In a recent preprint Professor Etesi asked a question: could one find a complex three dimensional submanifold S in a compact complex seven dimensional homogeneous space with the compact real 14 dimensional Lie group G_2 as the base manifold, such that S is diffeomorphic to the six dimensional sphere S^6 ? We apply a result of Tits on compact complex homogeneous space, or of H. C. Wang and Hano–Kobayashi on the classification of compact complex homogeneous manifolds with a compact reductive Lie group to give an answer to his question. In particular, we show that one could not obtain a complex structure of S^6 in his way.

1. Introduction

Let M be a complex manifold, h be an Hermitian metric. For a compact complex manifold, there is always some Hermitian metric h by the partition of the unity argument. If h is an Hermitian metric and G is a compact Lie group acting on M biholomorphically, then by taking average on G , we can always assume that h is invariant under G .

A compact complex homogeneous space with an invariant Hermitian structure was classified by H. C. Wang [1954], see also [Hano and Kobayashi 1960]. In fact, they classified the compact complex homogeneous spaces with compact Lie groups. In particular, an Hermitian manifold is a Riemannian manifold. The identity component of the Riemannian isometric group for a compact Riemannian manifold is a compact Lie group. So is the identity component of the Hermitian isometric group for a compact Hermitian manifold.

Therefore, we have:

MSC2010: primary 53C15; secondary 57S25, 53C30, 22E99, 15A75.

Keywords: complex structure, six dimensional sphere, cohomology, invariant structure, complex torus bundles, Hermitian manifolds.

Lemma 1. *If $M = G/H$ is a compact homogeneous Riemannian manifold with G connected, then G is a subgroup of a compact Lie group. In particular, both G and H are reductive with compact semisimple parts.*

We then have (see [Hano and Kobayashi 1960, Theorem B]):

Lemma 2. *Any compact Hermitian homogeneous manifold is a complex torus bundle over a rational (therefore simply connected) projective homogeneous space.*

One could also see [Guan 1994, page 66, Remark] for a detailed understanding of this fibration.

There is also a similar fibration [Tits 1971] closely related to Lemma 2 for any general compact complex homogeneous space:

Proposition 3. *Let $M = G/H$ be a compact complex homogeneous space such that G is a complex Lie group, with H a complex Lie subgroup and $M = G/H$ is the complex quotient. Then there is a complex fibration $G/H \rightarrow G/N$ such that $N = \text{Norm}_G(H^0)$ and G/N is a rational projective homogeneous space.*

Here, H^0 is the identity component of H and

$$\text{Norm}_G(H^0) = \{g \in G \mid gH^0g^{-1} \subset H^0\}$$

is the normalizer of H^0 in G .

Let $G = G_2$ be the compact real 14 dimensional Lie group of type G_2 , $G^{\mathbb{C}}$ be the complex simple Lie group such that the Lie algebra of $G^{\mathbb{C}}$ is the complexification of the Lie algebra of G . Let B be the Borel subgroup such that B is connected and the Lie algebra of B contains all the negative root vector spaces and the given Cartan subalgebra. Then $G^{\mathbb{C}}/B$ has a complex dimension 6. We notice that the Cartan subalgebra has a complex dimension 2. So, let U be the maximal nilpotent subgroup of B , which is generated by the all the negative root spaces; then B/U is a complex two dimensional algebraic torus $(\mathbb{C}^*)^2$. Let $\pi : B \rightarrow B/U$ be the quotient map. Given any complex one dimensional line subspace $l = \mathbb{C}$ in B/U , $S_l = \pi^{-1}(l)$ is a complex codimension one normal subgroup of B . Then $G^{\mathbb{C}}/S_l$ is diffeomorphic to G_2 . Therefore, it gives a complex structure on G_2 .

On the other hand, if $G = G_2$ is a complex homogeneous space itself, then its complexification $G^{\mathbb{C}}$ acts on G also. And $G = G^{\mathbb{C}}/H$. By Lemma 2, the Hano-Kobayashi fiber bundle is $G^{\mathbb{C}}/H \rightarrow G^{\mathbb{C}}/B$. The torus is just the Cartan subgroup of G . By earlier works, e.g., [Guan 2002], we know that the Tits fibration and the HK fibration are the same and therefore, $B = \text{Norm}_{G^{\mathbb{C}}}(H^0)$. But $\dim_{\mathbb{C}} B/H = 1$ and H^0 is normal in B . We see that H contains all the subgroups generated by the negative root spaces. That is, $U \subset H$.

Proposition 4. *Let G be a complex manifold with the base manifold being G_2 such that $G = G_2$ acts on itself biholomorphically. Then there is a holomorphic*

fibration from G into a projective manifold $F : G \rightarrow G^{\mathbb{C}}/B$ such that each fiber is a complex one dimensional complex torus. If S is a complex three-dimensional compact submanifold of G , then either $F(S)$ is complex three dimensional or $F(S)$ is complex two dimensional and $S = F^{-1}(F(S))$.

Therefore, if S is diffeomorphic to S^6 , by the result in [Campana et al. 1998; Campana et al. 2020], the algebraic dimension of a complex structure on S is zero. This implies that S can not be a complex submanifold of G .

However, in this paper, we would like to give an argument which is mildly independent of the result from Campana et al.

We shall prove:

Main Theorem. *The conjugate orbit described in both [Etesi 2015] and [Chaves and Rigas 1991] can not be a complex submanifold of G_2 .*

Recall that $S^6 = G_2/SU(3)$. The Cartan subgroup of $SU(3)$ is also a Cartan subgroup of G_2 . The Cartan subgroup of $SU(3)$ has elements $\text{diag}(a, b, c)$ with $abc = 1$. The center of $SU(3)$ has elements $aI = \text{diag}(a, a, a)$ with $a^3 = 1$. This implies the center of $SU(3)$ has the structure $C = \mathbb{Z}/(3\mathbb{Z}) = \mathbb{Z}_3$. Let A be one of the generator of $C = \mathbb{Z}_3$. Then $S = \{gAg^{-1} \mid g \in G_2\}$. Since A is in the center of $SU(3)$, it is not difficult to see that $SU(3)$ fixes A .

A description of the Lie algebra of G_2 can be also found in the Section 4 of [Guan 2006], in which the Lie algebra of $SU(3)$ is simply generated by the root spaces with the short roots $e_i - e_j$ with $\{i, j\} \in \{1, 2, 3\}$.

2. Proofs

Proof of Proposition 4. If $F(S)$ in Proposition 4 has two complex dimensions, let $p \in F(S)$ be a regular point for F ; then $F^{-1}(p) \cap S$ is a complex one-dimensional manifold. As a closed complex one-dimensional submanifold of $F^{-1}(p)$, it can only be the whole complex torus. Since the fiber bundle is locally free and S a complex submanifold, the regular points for F are a dense set, by continuity we have the last part of Proposition 4 \square

Proof of Main Theorem. Assume that S is a complex submanifold, we shall get a contradiction.

If $F(S)$ is complex three dimensional, by $F(G)$ being projective, the pullback of the Kähler form ω is a nonzero H^2 class since it introduces a nonzero measure ω^3 on S . A contradiction.

If $F(S)$ is of complex dimension two, the conjugate orbit of Λ in [Etesi 2015] or [Chaves and Rigas 1991] is S . Here, Λ is one of the two nonidentity elements in the center of $SU(3)$ and therefore is in the given Cartan subgroup of $SU(3)$, which is exactly the Cartan subgroup T of G_2 . But $S = F^{-1}(F(S))$, we have that

the whole Cartan subgroup of G_2 is $F^{-1}(F(\Lambda))$. Therefore, S could be described as $S_t = \{gtg^{-1} \mid g \in G_2\}$ for any $t \in T$. But $S_t = G/G_t$ with G_t the centralizer $\{g \in G \mid gt = tg\}$. However, in general, $G_t = T$. That is S_t is real 12 dimensional. A contradiction. \square

Acknowledgments

I would like to take this chance to thank Professor S. Feng and the School of Mathematics and Statistics at Henan University for their support and hospitality while I was collaborating on this topic. Thanks also go to Professor Dorfmeister for showing me [Tits 1971; Wang 1954; Hano and Kobayashi 1960] when I was still a graduate and Professor Salamon for support for [Guan 2006].

References

- [Campana et al. 1998] F. Campana, J.-P. Demailly, and T. Peternell, “The algebraic dimension of compact complex threefolds with vanishing second Betti number”, *Compositio Math.* **112**:1 (1998), 77–91. [MR](#) [Zbl](#)
- [Campana et al. 2020] F. Campana, J.-P. Demailly, and T. Peternell, “The algebraic dimension of compact complex threefolds with vanishing second Betti number”, *Compos. Math.* **156**:4 (2020), 679–696. [MR](#) [Zbl](#)
- [Chaves and Rigas 1991] L. M. Chaves and A. Rigas, “On a conjugate orbit of G_2 ”, *Math. J. Okayama Univ.* **33** (1991), 155–161. [MR](#) [Zbl](#)
- [Etesi 2015] G. Etesi, “Explicit construction of the complex structure on the six dimensional sphere”, preprint, 2015. [arXiv](#)
- [Guan 1994] D. Guan, “Examples of compact holomorphic symplectic manifolds which admit no Kähler structure”, pp. 63–74 in *Geometry and analysis on complex manifolds*, edited by T. Mabuchi et al., World Sci. Publ., River Edge, NJ, 1994. [MR](#) [Zbl](#)
- [Guan 2002] D. Guan, “Classification of compact complex homogeneous spaces with invariant volumes”, *Trans. Amer. Math. Soc.* **354**:11 (2002), 4493–4504. [MR](#) [Zbl](#)
- [Guan 2006] D. Guan, “Existence of extremal metrics on almost homogeneous manifolds of cohomogeneity one, IV”, *Ann. Global Anal. Geom.* **30**:2 (2006), 139–167. [MR](#) [Zbl](#)
- [Hano and Kobayashi 1960] J.-i. Hano and S. Kobayashi, “A fibering of a class of homogeneous complex manifolds”, *Trans. Amer. Math. Soc.* **94** (1960), 233–243. [MR](#) [Zbl](#)
- [Tits 1971] J. Tits, “Représentations linéaires irréductibles d’un groupe réductif sur un corps quelconque”, *J. Reine Angew. Math.* **247** (1971), 196–220. [MR](#) [Zbl](#)
- [Wang 1954] H.-C. Wang, “Closed manifolds with homogeneous complex structure”, *Amer. J. Math.* **76** (1954), 1–32. [MR](#) [Zbl](#)

Received May 31, 2019. Revised August 28, 2019.

DANIEL GUAN
 SCHOOL OF MATHEMATICS AND STATISTICS
 HENAN UNIVERSITY
 KAIFENG
 CHINA
guan@henu.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 305 No. 2 April 2020

| | |
|---|-----|
| Singular periodic solutions to a critical equation in the Heisenberg group CLAUDIO AFELTRA | 385 |
| On a theorem of Hegyvári and Hennecart DAO NGUYEN VAN ANH, LE QUANG HAM, DOOWON KOH, THANG PHAM and LE ANH VINH | 407 |
| On the Ekeland–Hofer symplectic capacities of the real bidisc LUCA BARACCO, MARTINO FASSINA and STEFANO PINTON | 423 |
| Freeness characterizations on free chaos spaces SOLESNE BOURGUIN and IVAN NOURDIN | 447 |
| Dominance order and monoidal categorification of cluster algebras ELIE CASBI | 473 |
| On the fine expansion of the unipotent contribution of the Guo–Jacquet trace formula PIERRE-HENRI CHAUDOUARD | 539 |
| Strongly algebraic realization of dihedral group actions KARL HEINZ DOVERMANN | 563 |
| On commuting billiards in higher-dimensional spaces of constant curvature ALEXEY GLUTSYUK | 577 |
| On the arithmetic of a family of twisted constant elliptic curves RICHARD GRIFFON and DOUGLAS ULMER | 597 |
| On the nonexistence of S^6 type complex threefolds in any compact homogeneous complex manifolds with the compact lie group G_2 as the base manifold DANIEL GUAN | 641 |
| On $SU(3)$ Toda system with multiple singular sources ALI HYDER, CHANGSHOU LIN and JUNCHENG WEI | 645 |
| Convergence of mean curvature flow in hyper-Kähler manifolds KEITA KUNIKAWA and RYOSUKE TAKAHASHI | 667 |
| The two-dimensional analogue of the Lorentzian catenary and the Dirichlet problem RAFAEL LÓPEZ | 693 |
| Schwarz D-surfaces in $\mathbb{N}\mathbb{I}_3$ HEYONG SHIN, YOUNG WOOK KIM, SUNG-EUN KOH, HYUNG YONG LEE and SEONG-DEOG YANG | 721 |
| Compactness of constant mean curvature surfaces in a three-manifold with positive Ricci curvature AO SUN | 735 |
| The rational cohomology Hopf algebra of a generic Kac–Moody group ZHAO XU-AN and GAO HONGZHU | 757 |