

Pacific Journal of Mathematics

**SOME CLASSIFICATIONS OF
BIHARMONIC HYPERSURFACES WITH
CONSTANT SCALAR CURVATURE**

SHUN MAETA AND YE-LIN OU

SOME CLASSIFICATIONS OF BIHARMONIC HYPERSURFACES WITH CONSTANT SCALAR CURVATURE

SHUN MAETA AND YE-LIN OU

We give some classifications of biharmonic hypersurfaces with constant scalar curvature. These include biharmonic Einstein hypersurfaces in space forms, compact biharmonic hypersurfaces with constant scalar curvature in a sphere, and some complete biharmonic hypersurfaces of constant scalar curvature in space forms and in a nonpositively curved Einstein space. Our results provide additional cases (Theorem 2.3 and Proposition 2.8) that support the conjecture that a biharmonic submanifold in S^{m+1} has constant mean curvature, and two more cases that support Chen's conjecture on biharmonic hypersurfaces (Corollaries 2.2 and 2.7).

1. Introduction

Biharmonic maps, as a generalization of harmonic maps, are maps between Riemannian manifolds which are critical points of the bienergy functional. Biharmonic submanifolds are the images of biharmonic isometric immersions and they include minimal submanifolds as special cases. As in the study of minimal submanifolds, a fundamental problem in the study of biharmonic submanifolds is to classify nonminimal biharmonic submanifolds (called proper biharmonic submanifolds) in a model space. For example, when the ambient space is a space form, we have the following conjectures which are still far beyond our reach.

Conjecture (Chen's conjecture on biharmonic submanifolds of Euclidean space [1991; 2015]). *Any biharmonic submanifold in a Euclidean space is minimal.*

Maeta is supported partially by the Grant-in-Aid for Young Scientists(B), No.15K17542, Japan Society for the Promotion of Science, and partially by JSPS Overseas Research Fellowships 2017-2019 No. 70. The work was done while he was visiting the Department of Mathematics of Texas A & M University–Commerce as a Visiting Scholar and he is grateful to the department and the university for the hospitality he received during the visit. Ou is supported by a grant from the Simons Foundation (#427231, Ye-Lin Ou).

MSC2010: primary 58E20; secondary 53C12.

Keywords: biharmonic hypersurfaces, Einstein manifolds, constant scalar curvature, constant mean curvature.

Conjecture (Balmuş, Montaldo and Oniciuc's conjectures on biharmonic submanifolds of spheres [2008; 2012]). (1) *A proper biharmonic submanifold of a sphere has constant mean curvature.*

(2) *The only proper biharmonic hypersurface of S^{m+1} is a part of $S^m(1/\sqrt{2})$ or $S^p(1/\sqrt{2}) \times S^q(1/\sqrt{2})$ with $p + q = m$ and $p \neq q$.*

A lot of work related to these conjectures has been done since 2000; some recent developments, partial results and open problems in the topics can be found in the recent survey [Ou 2016].

For a hypersurface $\varphi : (M^m, g) \rightarrow (N^{m+1}, h)$ in a Riemannian manifold, i.e., an isometric immersion of codimension 1, we choose a local unit normal vector field ξ with respect to which, we have the second fundamental form $B(X, Y) = b(X, Y)\xi$, where $b : TM \times TM \rightarrow C^\infty(M)$ is the function-valued second fundamental form. The Gauss and the Weingarten formulae read, respectively,

$$(1) \quad \tilde{\nabla}_X Y = (\tilde{\nabla}_X Y)^\top + (\tilde{\nabla}_X Y)^\perp = \nabla_X Y + b(X, Y)\xi,$$

and

$$(2) \quad \tilde{\nabla}_X \xi = (\tilde{\nabla}_X \xi)^\top + (\tilde{\nabla}_X \xi)^\perp = -AX + \nabla_X^\perp \xi = -AX,$$

where A is the shape operator of the hypersurface with respect to the unit normal vector ξ .

We use R^M , Ric^M , and Scal^M (respectively, R^N , Ric^N , and Scal^N) to denote the Riemannian, Ricci and scalar curvatures of (M^m, g) (respectively, of (N, h)) with the following conventions:

$$\begin{aligned} R^M(X, Y, Z, W) &= \langle R^M(Z, W)Y, X \rangle, \\ R^M(Z, W)Y &= \nabla_Z \nabla_W Y - \nabla_W \nabla_Z Y - \nabla_{[Z, W]} Y, \\ \text{Ric}^M(X, Y) &= \sum_{i=1}^m R^M(X, e_i, Y, e_i) \quad \text{for an orthonormal base } \{e_i\}. \end{aligned}$$

Using these, together with (1) and (2), we have the Gauss equation

$$(3) \quad R^N(X, Y, Z, W) = R^M(X, Y, Z, W) + b(X, W)b(Y, Z) - b(X, Z)b(Y, W),$$

where

$$(4) \quad b(X, Y) = \langle AX, Y \rangle = \langle B(X, Y), \xi \rangle,$$

for any $X, Y, Z, W \in TM$.

From (3) we have the relationship between the Ricci curvatures of the hypersurface and the ambient space

$$(5) \quad \text{Ric}^N(X, Y) = \text{Ric}^M(X, Y) + \langle AX, AY \rangle - mH \langle AX, Y \rangle + R^N(X, \xi, Y, \xi),$$

and the relationship between the scalar curvatures is

$$(6) \quad \text{Scal}^N = \text{Scal}^M + |A|^2 - m^2 H^2 + 2 \text{Ric}^N(\xi, \xi).$$

It was proved in [Ou 2010] that a hypersurface $\varphi : M^m \rightarrow N^{m+1}$ with mean curvature vector $\eta = H\xi$ is biharmonic if and only if

$$(7) \quad \begin{cases} \Delta H - H|A|^2 + H \text{Ric}^N(\xi, \xi) = 0, \\ 2A(\text{grad } H) + \frac{m}{2} \text{grad } H^2 - 2H(\text{Ric}^N(\xi))^\top = 0, \end{cases}$$

where $\text{Ric}^N : T_q N \rightarrow T_q N$ denotes the Ricci operator of the ambient space defined by $\langle \text{Ric}^N(Z), W \rangle = \text{Ric}^N(Z, W)$.

In particular, for biharmonic hypersurfaces in an Einstein space, we have:

Corollary 1.1 [Ou 2010]. *A hypersurface $\varphi : M^m \rightarrow (N^{m+1}, h)$ of an Einstein manifold with $\text{Ric}^N = \lambda h$ is biharmonic if and only if its mean curvature function H solves the equations*

$$(8) \quad \begin{cases} \Delta H - H(|A|^2 - \lambda) = 0, \\ A(\text{grad } H) + \frac{m}{2} H \text{grad } H = 0. \end{cases}$$

In this note, we give some classifications of biharmonic hypersurfaces with constant scalar curvature. These include biharmonic Einstein hypersurfaces in space forms, compact biharmonic hypersurfaces with constant scalar curvature in a sphere, and some complete biharmonic hypersurfaces with constant scalar curvature in space forms and in a nonpositively curved Einstein space. Our results provide a further case (Theorem 2.3) that supports Balmuş, Montaldo and Oniciuc's conjecture that a biharmonic submanifold in S^{m+1} has constant mean curvature, and two more cases that support Chen's conjecture on biharmonic hypersurfaces (Corollaries 2.2 and 2.7).

2. Biharmonic hypersurfaces with constant scalar curvature in Einstein spaces

Recall that a Riemannian manifold (M^m, g) is called an Einstein space if its Ricci curvature is proportional to the metric, i.e., $\text{Ric}^M = \lambda g$. It is well known that every 2-dimensional manifold is an Einstein space; any space form is of Einstein, and any 3-dimensional Einstein space has to be a space form. One can also check that for $m \geq 3$, if (M^m, g) is an Einstein space with $\text{Ric}^M = \lambda g$, then λ has to be a constant and hence an Einstein space (M^m, g) of dimension $m \geq 3$ has constant scalar curvature since $\text{Scal}^M = m\lambda$.

Our first result is the following theorem which gives a classification of biharmonic Einstein hypersurfaces in a space form.

Theorem 2.1. *An Einstein hypersurface $M^m \hookrightarrow (N^{m+1}(C), h)$ ($m \geq 3$) in a space form is biharmonic if and only if it is minimal or $|A|^2 = mC$. In the latter case, the hypersurface has positive scalar curvature, i.e., $\text{Scal}^M = m(m-2)C + m^2H^2 > 0$.*

Proof. Suppose that the mean curvature function H is not constant; then there exists an open neighborhood U of M on which $\nabla H \neq 0$. Substituting $X = Y = \nabla H$ into (5) gives

$$(9) \quad \text{Ric}^M(\nabla H, \nabla H) = (m-1)C|\nabla H|^2 + mHg(A(\nabla H), \nabla H) - g(A(\nabla H), A(\nabla H)).$$

Substituting the second equation $A(\nabla H) = -\frac{m}{2}H\nabla H$ of the biharmonic equation for a hypersurface into the above equation we have

$$(10) \quad \text{Ric}^M(\nabla H, \nabla H) = ((m-1)C - \frac{3}{4}m^2H^2)|\nabla H|^2.$$

If (M^m, g) is an Einstein hypersurface with $\text{Ric}^M = \mu g$ for some constant μ , then (10) becomes

$$(11) \quad \mu|\nabla H|^2 = ((m-1)C - \frac{3}{4}m^2H^2)|\nabla H|^2.$$

Since $\nabla H \neq 0$ on $U \subset M$, we have $\mu = (m-1)C - \frac{3}{4}m^2H^2$, which implies H is a constant on $U \subset M$ since μ is a constant. This contradicts the assumption that $\nabla H \neq 0$ on $U \subset M$. The contradiction shows that the mean curvature function H has to be a constant. It follows from the first equation $\Delta H - H(|A|^2 - mC) = 0$ of (8) that either $H = 0$ and the hypersurface is minimal, or $|A|^2 = mC$, and in this case, $C > 0$. Using the scalar curvature $\text{Scal}^N = (m+1)mC > 0$ of $(N^{m+1}(C), h)$, $|A|^2 = mC$ and (6) we have $\text{Scal}^M = m(m-2)C + m^2H^2 > 0$.

Thus, we obtain the theorem. $\square$

As an immediate consequence of Theorem 2.1, we have:

Corollary 2.2. *A biharmonic Einstein hypersurface of Euclidean space $\mathbb{R}^{m+1}$ or hyperbolic space H^{m+1} is minimal. A proper biharmonic Einstein hypersurface in S^{m+1} has constant mean curvature and $|A|^2 = m$.*

It was proved in [Chen 1993] (see also [Balmuş et al. 2012]) that for $m \geq 2$, if a compact biharmonic hypersurface in sphere $S^{m+1}(1)$ with the squared norm of the second fundamental form satisfies $|A|^2 \leq m$, then $|A|^2 = 0$, or $|A|^2 = m$ and it has constant mean curvature. Also, Fu [2015] proved that a biharmonic hypersurface with constant scalar curvature in 5-dimensional space forms $M^5(C)$ has constant mean curvature, and in [Fu and Hong 2018], it was proved that a biharmonic hypersurface with constant scalar curvature and at most six distinct principal curvatures in space forms $M^{m+1}(C)$ has constant mean curvature. Our next theorem shows that the same result as in [Fu and Hong 2018] holds when we replace the principal curvature assumption by the compactness of the hypersurface.

Theorem 2.3. *A compact hypersurface with constant scalar curvature $(M^m, g) \hookrightarrow S^{m+1}$ in a sphere is biharmonic if and only if it is minimal, or it has nonzero constant mean curvature, and $|A|^2 = m$.*

Proof. If the scalar curvature Scal^M of the hypersurface is constant, then, by (6), we have $|A|^2 = m^2 H^2 + \text{constant}$, from which we have

$$(12) \quad \nabla |A|^2 = 2m^2 H \nabla H.$$

Using this, together with the first equation of (8) with $\lambda = m$, we have

$$(13) \quad \nabla \Delta H = \nabla[(|A|^2 - m)H] = (|A|^2 - m + 2m^2 H^2) \nabla H.$$

On the other hand, we have the following estimate of the squared norm of the Hessian of H :

$$(14) \quad |\nabla dH|^2 = \sum_{i,j=1}^m [\nabla dH(e_i, e_j)]^2 \geq \sum_{i=1}^m [\nabla dH(e_i, e_i)]^2 \geq \frac{1}{m} \left(\sum_{i=1}^m \nabla dH(e_i, e_i) \right)^2 \\ = \frac{1}{m} (\Delta H)^2.$$

Substituting (10), (13) and (14) into the Bochner formula for $|\nabla H|^2$, we have

$$(15) \quad \frac{1}{2} \Delta |\nabla H|^2 = |\nabla dH|^2 + \text{Ric}^M(\nabla H, \nabla H) + \langle \nabla H, \nabla \Delta H \rangle \\ \geq \frac{1}{m} (\Delta H)^2 + (|A|^2 - 1 + \frac{5}{4} m^2 H^2) |\nabla H|^2.$$

Since M is compact, we integrate both sides of (15) to get

$$(16) \quad \int_M \left[\frac{1}{m} (\Delta H)^2 + (|A|^2 - 1 + \frac{5}{4} m^2 H^2) |\nabla H|^2 \right] dv_g \leq \int_M \frac{1}{2} \Delta |\nabla H|^2 dv_g = 0.$$

Using compactness of M , (13), and the divergence theorem we have

$$(17) \quad \frac{1}{m} \int_M \Delta H \Delta H dv_g = \frac{1}{m} \int_M \Delta H \text{div}(\nabla H) dv_g = -\frac{1}{m} \int_M \langle \nabla \Delta H, \nabla H \rangle dv_g \\ = -\frac{1}{m} \int_M (|A|^2 - m + 2m^2 H^2) |\nabla H|^2 dv_g,$$

Substituting this into (16) we have

$$(18) \quad 0 \leq \int_M \left(\left(1 - \frac{1}{m}\right) |A|^2 + \frac{m}{4} (5m - 8) H^2 \right) |\nabla H|^2 dv_g \leq 0.$$

It follows that $\left[\left(1 - \frac{1}{m}\right) |A|^2 + \frac{m}{4} (5m - 8) H^2 \right] |\nabla H|^2 = 0$ from which, together with Newton's inequality $|A|^2 \geq m H^2$, we have

$$(19) \quad 0 = \left[\left(1 - \frac{1}{m}\right) |A|^2 + \frac{m}{4} (5m - 8) H^2 \right] |\nabla H|^2 \geq \frac{1}{16} (5m^2 - 4m - 4) |\nabla H^2|^2.$$

From this, we conclude that $H = \text{constant}$. In the case where $H = \text{constant} \neq 0$, we use the first equation of (8) to have $|A|^2 = m$. Thus, we obtain the theorem. $\square$

- Remark 1.** (i) Notice that [Theorem 2.3](#), together with the results in [\[Chen 1993\]](#) and [\[Balmuş et al. 2012\]](#), implies that for a compact biharmonic hypersurface in a sphere, if one of the data Scal^M , H and $|A|^2$ is constant, then so are the other two.
- (ii) We would like to point out that according to a classical result of Chern, do Carmo and Kobayashi [\[Chern et al. 1970\]](#), any compact minimal hypersurface in S^{m+1} with $|A|^2 = m$ is locally the Clifford tori $S^p \times S^{m-p}$. On the other hand, a hypersurface of S^{m+1} with constant mean curvature and $|A|^2 = m$ is biharmonic. So it would be very interesting to classify hypersurfaces of S^{m+1} with constant mean curvature and $|A|^2 = m$. This would be an important case to solve the second of Balmuş, Montaldo and Oniciuc's conjectures.
- (iii) Finally, we remark that there are infinitely many compact hypersurfaces of constant scalar curvatures in a sphere. In fact, it was proved in [\[Ejiri 1982\]](#) that there exist countably many isometric immersions of $S^1 \times S^{m-1}$ into a sphere S^{m+1} so that $S^1 \times S^{m-1}$ is a warped product of constant scalar curvature $m(m-1)$ with respect to the induced metric.

Theorem 2.4. *A compact Einstein hypersurface $M^m \hookrightarrow (N^{m+1}, h)$ in an Einstein manifold with $\text{Ric}^N = \lambda h$ is biharmonic if and only if it is minimal or $|A|^2 = \lambda$. Furthermore, in the latter case, the hypersurface has positive scalar curvature, i.e.,*

$$\text{Scal}^M = (m-2)\lambda + m^2 H^2 > 0.$$

Proof. Suppose the hypersurface M is Einstein with $\text{Ric}^M = \mu g$, then, by (6),

$$(m+1)\lambda = m\mu + |A|^2 - m^2 H^2 + 2\lambda.$$

Hence,

$$(20) \quad \mu = \left(1 - \frac{1}{m}\right)\lambda - \frac{1}{m}|A|^2 + mH^2.$$

It follows that

$$(21) \quad \begin{aligned} \langle \nabla H, \nabla \Delta H \rangle &= \langle \nabla H, \nabla (H|A|^2 - \lambda H) \rangle \\ &= |A|^2 |\nabla H|^2 + H \langle \nabla H, \nabla |A|^2 \rangle - \lambda |\nabla H|^2 \\ &= (|A|^2 - \lambda + 2m^2 H^2) |\nabla H|^2, \end{aligned}$$

where the first equality was obtained by using the first equation of (8), and the third equality follows from (6) and the fact that the scalar curvature of an Einstein hypersurface is constant.

Using these and a similar computation used in obtaining (15) we have,

$$\begin{aligned}
 (22) \quad \frac{1}{2} \Delta |\nabla H|^2 &= |\nabla dH|^2 + \text{Ric}^M(\nabla H, \nabla H) + \langle \nabla H, \nabla \Delta H \rangle \\
 &\geq \frac{1}{m} (\Delta H)^2 + \mu |\nabla H|^2 + (|A|^2 - \lambda + 2m^2 H^2) |\nabla H|^2 \\
 &= \frac{1}{m} (\Delta H)^2 + \left(-\frac{1}{m} \lambda + \left(1 - \frac{1}{m} \right) |A|^2 + m(2m+1) H^2 \right) |\nabla H|^2.
 \end{aligned}$$

Similar to (17), we have

$$\begin{aligned}
 (23) \quad \frac{1}{m} \int_M \Delta H \Delta H dv_g &= \frac{1}{m} \int_M \Delta H \text{div}(\nabla H) dv_g = -\frac{1}{m} \int_M \langle \nabla \Delta H, \nabla H \rangle dv_g \\
 &= -\frac{1}{m} \int_M (|A|^2 - \lambda + 2m^2 H^2) |\nabla H|^2 dv_g.
 \end{aligned}$$

Integrating both sides of (22) and using the compactness of M and (23) we obtain

$$\begin{aligned}
 (24) \quad 0 &\geq \int_M \left(\frac{1}{m} (m-2) |A|^2 + m(2m-1) H^2 \right) |\nabla H|^2 dv_g \\
 &\geq \frac{1}{4} m(2m-1) \int_M |\nabla H|^2 dv_g.
 \end{aligned}$$

It follows that H is constant. If $H = 0$, then M is minimal. If $H \neq 0$, by the first equation of (8), we have $|A|^2 = \lambda$. Thus, $\text{Scal}^M = m\mu = (m-2)\lambda + m^2 H^2 > 0$. $\square$

For the classification of complete biharmonic hypersurfaces, we notice that it was proved in [Nakauchi and Urakawa 2011], [Maeta 2014a] and [Luo 2015] that a complete hypersurface $M^m \hookrightarrow (N^{m+1}, h)$ in a manifold of nonpositive Ricci curvature with mean curvature function $H \in L^p(M)$ for some $0 < p < \infty$ is minimal. We will give a classification of biharmonic hypersurfaces of constant scalar curvatures in an Einstein manifold using a condition on the maximum rate of change of the mean curvature function. For that purpose, we will need the following maximum principles:

Theorem 2.5 [Yau 1976; Karp and Li 1982]. *Let M be a complete Riemannian manifold and f be a smooth function on M . If one of the following conditions is satisfied, then f is constant.*

- (i) M has Ricci curvature bounded from below and $\Delta f \geq \varepsilon f$ (for some $\varepsilon > 0$) for an upper bounded function $f \geq 0$.
- (ii) $\Delta f \geq 0$ for $f \geq 0$ and $f \in L^p(M)$ for some $1 < p < \infty$.
- (iii) $f \in L^1(M)$, $\Delta f \geq 0$, $f \geq 0$ and the Ricci curvature is bounded from below by $-c\{1 + r^2(x)\}$, where $r(x)$ is the distance function on M .

Now we are ready to give some classifications of biharmonic hypersurfaces with constant scalar curvature in a nonpositively curved Einstein manifold.

Proposition 2.6. *A complete biharmonic hypersurface $M^m \hookrightarrow (N^{m+1}, h)$ of constant scalar curvature in a nonpositively curved Einstein manifold (N^{m+1}, h) is minimal if one of the following occurs:*

- (a) $|\nabla H| \in L^p$ for some $2 < p < \infty$.
- (b) $|\nabla H| \in L^2$ and the Ricci curvature is bounded from below by $-c\{1 + r^2(x)\}$, where $r(x)$ is the distance function on M .

Proof. If the ambient space (N^{m+1}, h) is a nonpositively curved Einstein space with $\widetilde{\text{Ric}} = \lambda h$, then from (5) we have

$$(25) \quad \begin{aligned} \text{Ric}(\nabla H, \nabla H) &\geq \lambda |\nabla H|^2 + mH \langle A(\nabla H), \nabla H \rangle - \langle A(\nabla H), A(\nabla H) \rangle \\ &= \left(\lambda - \frac{3}{4}m^2 H^2\right) |\nabla H|^2, \end{aligned}$$

where the equality was obtained by using the second equation of the biharmonic hypersurface equation (8).

Substituting (25) and (21) into the Bochner formula we have

$$(26) \quad \begin{aligned} \Delta |\nabla H|^2 &= 2\{|\nabla dH|^2 + \text{Ric}(\nabla H, \nabla H) + \langle \nabla H, \nabla \Delta H \rangle\} \\ &\geq 2\{\text{Ric}(\nabla H, \nabla H) + \langle \nabla H, \nabla \Delta H \rangle\} \\ &= 2\left[\left(\lambda - \frac{3}{4}m^2 H^2\right) + (|A|^2 - \lambda + 2m^2 H^2)\right] |\nabla H|^2 \\ &= 2\left(\frac{5}{4}m^2 H^2 + |A|^2\right) |\nabla H|^2 \geq 2\left(\frac{5}{4}m^2 H^2 + mH^2\right) |\nabla H|^2 \\ &= \frac{1}{8}m(5m + 4) |\nabla H^2|^2 \geq 0. \end{aligned}$$

Using maximum principles (ii) and (iii) in [Theorem 2.5](#) we have that $|\nabla H|$ is constant. Using this and (26) again we conclude that $|\nabla H^2| = 0$. It follows that H is constant. If $H = \text{constant} \neq 0$, then, by the first equation of (8), we have $|A|^2 - \lambda = 0$. It follows that $|A|^2 = \lambda \leq 0$ since N is nonpositively curved. From this, we have $|A|^2 = 0$, which means that M is totally geodesic and hence minimal, which is a contradiction. So we are left with the only conclusion that $H = 0$, that is, the hypersurface is minimal. Thus, we complete the proof of the proposition. $\square$

Remark 2. (i) For a classification of complete biharmonic submanifolds with Ricci curvature bounded from below in a nonpositively curved manifold see [\[Maeta 2014b\]](#).

(ii) Our [Theorem 2.4](#) and [Proposition 2.6](#) give some classifications of biharmonic hypersurfaces in an Einstein space. We refer to [\[Inoguchi and Sasahara 2016; 2017\]](#) for some examples and classifications of constant mean curvature proper biharmonic hypersurfaces in a special class of Einstein spaces—the compact Riemannian symmetric space with a G -invariant metric. Also, for some classifications of f -biharmonic hypersurfaces in an Einstein space, see [\[Ou 2017\]](#).

As a corollary of [Proposition 2.6](#), we have an affirmative partial answer to Chen's conjecture:

Corollary 2.7. *Any complete biharmonic hypersurface with constant scalar curvature and $|\nabla H| \in L^p(M)$ for some $2 < p < \infty$ in a Euclidean space is minimal.*

Proposition 2.8. *Let $M^m \hookrightarrow S^{m+1}$ be a complete biharmonic hypersurface with constant scalar curvature in a sphere with $H^2 \geq (2\varepsilon + 4)/(m(5m + 4))$ (for some $\varepsilon > 0$). If one of the following is satisfied, then the mean curvature is constant.*

- (A) *The Ricci curvature of M is bounded from below, and $|\nabla H|$ from above.*
- (B) *$|\nabla H| \in L^p$ for $2 < p < \infty$.*
- (C) *$|\nabla H| \in L^2$ and the Ricci curvature is bounded from below by $-c\{1 + r^2(x)\}$, where $r(x)$ is the distance function on M .*

Proof. It follows from [\(15\)](#) that

$$\Delta|\nabla H|^2 \geq 2\left(-1 + \frac{5}{4}m^2H^2 + |A|^2\right)|\nabla H|^2.$$

Using Newton's formula $|A|^2 \geq mH^2$ for a hypersurface we have

$$\Delta|\nabla H|^2 \geq 2\left(-1 + \frac{m}{4}(5m + 4)H^2\right)|\nabla H|^2.$$

From this, together with the assumption that $H^2 \geq (2\varepsilon + 4)/(m(5m + 4))$, we have

$$(27) \quad \Delta|\nabla H|^2 \geq \varepsilon|\nabla H|^2.$$

Using the maximum principles (i), (ii), and (iii) in [Theorem 2.5](#) with $f = |\nabla H|^2$ we have obtain the proposition. $\square$

Acknowledgments

We would like to thank C. Oniciuc and the referee for some very useful comments that helped to improve the manuscript.

References

- [Balmuş et al. 2008] A. Balmuş, S. Montaldo, and C. Oniciuc, “Classification results for biharmonic submanifolds in spheres”, *Israel J. Math.* **168** (2008), 201–220. [MR](#) [Zbl](#)
- [Balmuş et al. 2012] A. Balmuş, S. Montaldo, and C. Oniciuc, “New results toward the classification of biharmonic submanifolds in S^n ”, *An. Ştiinţ. Univ. “Ovidius” Constanţa Ser. Mat.* **20:2** (2012), 89–114. [MR](#) [Zbl](#)
- [Chen 1991] B.-Y. Chen, “Some open problems and conjectures on submanifolds of finite type”, *Soochow J. Math.* **17:2** (1991), 169–188. [MR](#) [Zbl](#)
- [Chen 1993] J. H. Chen, “Compact 2-harmonic hypersurfaces in $S^{n+1}(1)$ ”, *Acta Math. Sinica* **36:3** (1993), 341–347. In Chinese. [MR](#) [Zbl](#)
- [Chen 2015] B.-Y. Chen, *Total mean curvature and submanifolds of finite type*, 2nd ed., Series in Pure Mathematics **27**, World Scientific, Hackensack, NJ, 2015. [MR](#) [Zbl](#)

- [Chern et al. 1970] S. S. Chern, M. do Carmo, and S. Kobayashi, “Minimal submanifolds of a sphere with second fundamental form of constant length”, pp. 59–75 in *Functional analysis and related fields* (Chicago, IL, 1968), edited by F. E. Browder, Springer, 1970. [MR](#) [Zbl](#)
- [Ejiri 1982] N. Ejiri, “Some compact hypersurfaces of constant scalar curvature in a sphere”, *J. Geom.* **19**:2 (1982), 197–199. [MR](#) [Zbl](#)
- [Fu 2015] Y. Fu, “On biharmonic hypersurfaces with constant scalar curvatures in $\mathbb{S}^5$ ”, *Proc. Amer. Math. Soc.* **143**:12 (2015), 5399–5409. [MR](#) [Zbl](#)
- [Fu and Hong 2018] Y. Fu and M.-C. Hong, “Biharmonic hypersurfaces with constant scalar curvature in space forms”, *Pacific J. Math.* **294**:2 (2018), 329–350. [MR](#) [Zbl](#)
- [Inoguchi and Sasahara 2016] J.-i. Inoguchi and T. Sasahara, “Biharmonic hypersurfaces in Riemannian symmetric spaces I”, *Hiroshima Math. J.* **46**:1 (2016), 97–121. [MR](#) [Zbl](#)
- [Inoguchi and Sasahara 2017] J.-i. Inoguchi and T. Sasahara, “Biharmonic hypersurfaces in Riemannian symmetric spaces II”, *Hiroshima Math. J.* **47**:3 (2017), 349–378. [MR](#) [Zbl](#)
- [Karp and Li 1982] L. Karp and P. Li, “The heat equation on complete Riemannian manifolds”, preprint, 1982.
- [Luo 2015] Y. Luo, “On biharmonic submanifolds in non-positively curved manifolds”, *J. Geom. Phys.* **88** (2015), 76–87. [MR](#) [Zbl](#)
- [Maeta 2014a] S. Maeta, “Biharmonic maps from a complete Riemannian manifold into a non-positively curved manifold”, *Ann. Global Anal. Geom.* **46**:1 (2014), 75–85. [MR](#) [Zbl](#)
- [Maeta 2014b] S. Maeta, “Properly immersed submanifolds in complete Riemannian manifolds”, *Adv. Math.* **253** (2014), 139–151. [MR](#) [Zbl](#)
- [Nakauchi and Urakawa 2011] N. Nakauchi and H. Urakawa, “Biharmonic hypersurfaces in a Riemannian manifold with non-positive Ricci curvature”, *Ann. Global Anal. Geom.* **40**:2 (2011), 125–131. [MR](#) [Zbl](#)
- [Ou 2010] Y.-L. Ou, “Biharmonic hypersurfaces in Riemannian manifolds”, *Pacific J. Math.* **248**:1 (2010), 217–232. [MR](#) [Zbl](#)
- [Ou 2016] Y.-L. Ou, “Some recent progress of biharmonic submanifolds”, pp. 127–139 in *Recent advances in the geometry of submanifolds*, edited by B. D. Suceavă et al., Contemp. Math. **674**, Amer. Math. Soc., Providence, RI, 2016. [MR](#) [Zbl](#)
- [Ou 2017] Y.-L. Ou, “ f -biharmonic maps and f -biharmonic submanifolds II”, *J. Math. Anal. Appl.* **455**:2 (2017), 1285–1296. [MR](#) [Zbl](#)
- [Yau 1976] S. T. Yau, “Some function-theoretic properties of complete Riemannian manifold and their applications to geometry”, *Indiana Univ. Math. J.* **25**:7 (1976), 659–670. [MR](#) [Zbl](#)

Received November 4, 2018. Revised September 8, 2019.

SHUN MAETA
DEPARTMENT OF MATHEMATICS
SHIMANE UNIVERSITY
MATSUE
JAPAN
maeta@riko.shimane-u.ac.jp

YE-LIN OU
DEPARTMENT OF MATHEMATICS
TEXAS A & M UNIVERSITY
COMMERCE, TX
UNITED STATES
yelin_ou@tamu-commerce.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY



mathematical sciences publishers

nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 306 No. 1 May 2020

Tensor structure for Nori motives	1
LUCA BARBIERI-VIALE, ANNETTE HUBER and MIKE PREST	
On the Garden of Eden theorem for endomorphisms of symbolic algebraic varieties	31
TULLIO CECCHERINI-SILBERSTEIN, MICHEL COORNAERT and XUAN KIEN PHUNG	
Bergman kernels of elementary Reinhardt domains	67
DEBRAJ CHAKRABARTI, AUSTIN KONKEL, MEERA MAINKAR and EVAN MILLER	
Central splitting of manifolds with no conjugate points	95
JAMES DIBBLE	
On the global Gan–Gross–Prasad conjecture for general spin groups	115
MELISSA EMORY	
Schur algebras for the alternating group and Koszul duality	153
THANGAVELU GEETHA, AMRITANSHU PRASAD and SHRADDHA SRIVASTAVA	
A positive mass theorem for manifolds with boundary	185
SVEN HIRSCH and PENGZI MIAO	
Circle patterns on surfaces of finite topological type revisited	203
YUE-PING JIANG, QIANGHUA LUO and ZE ZHOU	
On some conjectures of Heywood	221
DONG LI	
Knapp–Stein dimension theorem for finite central covering groups	265
CAIHUA LUO	
Some classifications of biharmonic hypersurfaces with constant scalar curvature	281
SHUN MAETA and YE-LIN OU	
Surface diffusion flow of arbitrary codimension in space forms	291
DONG PU and HONGWEI XU	
Nonvanishing square-integrable automorphic cohomology classes: the case $GL(2)$ over a central division algebra	321
JOACHIM SCHWERMER	
Invariant Banach limits and applications to noncommutative geometry	357
EVGENII SEMENOV, FEDOR SUKOCHEV, ALEXANDR USACHEV and DMITRIY ZANIN	
A combinatorial identity for the Jacobian of t -shifted invariants	375
OKSANA YAKIMOVA	