

*Pacific
Journal of
Mathematics*

**SURFACE DIFFUSION FLOW OF
ARBITRARY CODIMENSION IN SPACE FORMS**

DONG PU AND HONGWEI XU

Volume 306 No. 1

May 2020

SURFACE DIFFUSION FLOW OF ARBITRARY CODIMENSION IN SPACE FORMS

DONG PU AND HONGWEI XU

Dedicated to Professor Duanzhuang Qian on his 110th anniversary.

We first prove that a properly immersed static n -dimensional submanifold ($\Delta H = 0$) with restricted growth of the curvature at infinity in $\mathbb{F}^{n+p}(c)$ ($c \geq 0$) is totally umbilical if the Willmore functional is pinched by a positive constant depending only on n . Secondly, we obtain a global rigidity theorem for Willmore surfaces in the sphere. Thirdly, we give a lower bound on the lifespan of the surface diffusion flow in $\mathbb{F}^{2+p}(c)$. Finally, we get the longtime existence and convergence of the surface diffusion flow in $\mathbb{F}^{2+p}(c)$ ($c \geq 0$) under the small initial Willmore energy condition.

1. Introduction

An important problem in global differential geometry is the study of curvature and topology of Riemannian manifolds and submanifolds. As we know, curvature flows are powerful tools in the study of sphere theorems as in [Andrews and Baker 2010; Brendle and Schoen 2009; Hamilton 1982; Huisken 1984; Gu et al. 2017; Liu et al. 2018; Wang 2008; Gu and Xu 2012; Xu and Gu 2013], etc. For instance, Brendle and Schoen [2009] proved the remarkable differentiable $\frac{1}{4}$ -pinching sphere theorem via the Ricci flow, which had been open for half a century.

Let $\mathbb{F}^{n+p}(c)$ be the $(n + p)$ -dimensional complete and simply connected space form with constant curvature c . In this paper, we study the motion of an immersed submanifold with the normal velocity $-\Delta H$ in the space form $\mathbb{F}^{n+p}(c)$. More precisely, let $f : M^n \rightarrow \mathbb{F}^{n+p}(c)$ be a compact immersed submanifold. The diffusion flow for submanifolds is a fourth order flow

$$(1-1) \quad \frac{\partial f}{\partial t} = -\Delta H,$$

where $\nabla_X \phi = (D_X \phi)^\perp$ for a tangent vector field X and a normal vector field ϕ

Research supported by the National Natural Science Foundation of China, Grant Nos. 11531012, 11371315.

MSC2010: 53C24, 53C44.

Keywords: surface diffusion, Willmore surface, gap theorem, convergence theorem.

on M , $\Delta = \nabla_i \nabla_i$ and H is the mean curvature vector of M^n . Denote by A the second fundamental form and by $A^\circ = A - \frac{1}{n}g \otimes H$ the trace free part of the second fundamental form. When $\nabla H = 0$, we call M a submanifold with parallel mean curvature. In particular, M is minimal if $H = 0$. When $\Delta H = 0$, we call M a static submanifold for the diffusion flow which is a generalization of the submanifold with parallel mean curvature.

For $n = 2$, surface diffusion flow (1-1) in \mathbb{R}^3 was proposed by Mullins [1957] to describe thermal grooving in material sciences. From the view of geometric analysis it appears naturally as the gradient flow of the area functional with respect to the inner product of H^{-1} ; see [Mayer 2001; Taylor and Cahn 1994]. Escher, Mayer and Simonett [Escher et al. 1998] showed that solutions that start out close to spheres with respect to the $C^{2+\beta}$ -topology for hypersurfaces exist globally and converge exponentially fast to a sphere. Wheeler [2012, Chapter 3] obtained the following convergence theorem.

Theorem A. *Let $f : \Sigma \rightarrow \mathbb{R}^3$ be a compact surface. There exists an absolute constant $C_1 > 0$ such that if*

$$\int_{\Sigma} |A^\circ|^2 d\mu < C_1,$$

then the surface diffusion flow with initial data f exists smoothly for all time and converges exponentially to a round sphere as $t \rightarrow \infty$.

Willmore flow is another important fourth order flow that is closely related to the surface diffusion flow. Letting M be an n -dimensional compact submanifold in the space form, the Willmore functional is

$$(1-2) \quad \mathcal{W}(f) = \int_M |A^\circ|^n d\mu.$$

It is invariant under the conformal (or Moebius) transformations of the ambient space. When $n = 2$, the Willmore functional is also called Willmore energy. For a compact immersed surface $f : \Sigma \rightarrow \mathbb{R}^{2+p}$, the associated Euler–Lagrange operator is

$$(1-3) \quad W(f) = \Delta H + Q(A^\circ)H, \quad Q(A^\circ)H = A^\circ(e_i, e_j)\langle A^\circ(e_i, e_j), H \rangle.$$

When $W(f) = 0$, we call it the Willmore surface. The L_2 -gradient flow of $\mathcal{W}(f)$ for surfaces, briefly called Willmore flow, is a quasilinear geometric evolution equation

$$(1-4) \quad \frac{\partial f}{\partial t} = -W(f).$$

Kuwert and Schätzle [2002] gave a lower bound on the lifespan of a smooth solution, which depends only on how much the curvature of the initial surface is concentrated, and in [Kuwert and Schätzle 2001] they proved the following theorem:

Theorem B. *Let $f : \Sigma \rightarrow \mathbb{R}^{2+p}$ be a compact surface. There exists a constant $C_2(p) > 0$ such that if*

$$\int_{\Sigma} |A^\circ|^2 d\mu < C_2(p),$$

then the Willmore flow with initial data f exists smoothly for all time and converges exponentially to a round sphere as $t \rightarrow \infty$.

Kuwert and Schätzle [2004] investigated point singularities of Willmore surfaces and obtained that the Willmore flow of spheres in \mathbb{R}^3 with energy less than 8π exists at all times and converges to a round sphere.

For other fourth order flows, Bernard, Wheeler and Wheeler [Bernard et al. 2019] introduced the so-called Chen’s flow and investigated the lifespan theorem and finite-time singularities for such a flow. Moreover, one class of fourth order flows derived from the critical point of some constructed functionals in different background manifolds. Metzger, Wheeler and Wheeler [2013] investigated the gradient flow of $\int_{\Sigma} |H|^2 d\mu$ and proved the lifespan theorem in Riemannian 3-manifold. Link [2013] also studied this flow in bounded Riemannian manifolds. Magni [2015] studied the gradient flow of $\int_{\Sigma} |A|^2 d\mu$ and obtained a smooth convergence theorem in three-dimensional Riemannian manifolds.

The local existence of these fourth order curvature flows is standard as we can see in [Escher et al. 1998], for example. In this paper, we first obtain a gap theorem for an n -dimensional properly immersed static submanifold M^n with restricted growth of the curvature at infinity in $\mathbb{F}^{n+p}(c) (c \geq 0)$ under the pinching condition for the Willmore functional, which generalizes the gap lemma in [Wheeler 2012] to any dimension and codimension.

Theorem 1.1. *Let $f : M^n \rightarrow \mathbb{F}^{n+p}(c) (c \geq 0)$ be a properly immersed submanifold with $\Delta H = 0$. There exists a positive constant $\sigma(n)$ depending only on n such that if*

$$\int_M |A^\circ|^n d\mu < \sigma(n),$$

$$\liminf_{\rho \rightarrow \infty} \frac{1}{\rho^4} \int_{f^{-1}(B_\rho(x))} |A|^2 d\mu = 0 \quad \text{for } x \in \mathbb{F}^{n+p}(c),$$

then M is totally umbilical.

Notice that the properness of the immersion implies the completeness of M . For a complete immersion, if we choose the cutoff function on intrinsic balls, the properness of the immersion isn’t essential. For higher dimensions, there are few results about the convergence of fourth order flow, and the above gap theorem provides a feasible method to handle this problem. Xu and Gu [2007] proved a gap theorem for complete surfaces with parallel mean curvature in a space form under the pinching condition $\int_M |A^\circ|^4 d\mu < D(|H|, c)$, where $c + |H|^2/4 > 0$. Moreover,

they proposed an open problem asking if there is global rigidity for surfaces with parallel mean curvature under the pinching condition for the Willmore functional. As a consequence of Theorem 1.1, we have solved this open problem for compact surfaces in $\mathbb{F}^{2+p}(c)$ ($c \geq 0$).

For compact Willmore surfaces in a sphere, Xu and Yang [2016] proved a global rigidity theorem under the pinching condition for $\int_M |A^\circ|^4 d\mu$. Motivated by the convergence result in Theorem B, we obtain the following global rigidity theorem for compact Willmore surfaces in a sphere under the pinching condition for the Willmore functional.

Theorem 1.2. *Let M be a compact Willmore surface in the unit sphere \mathbb{S}^{2+p} . There exists an absolute positive constant σ_1 ($= 6.23 \times 10^{-8}$) such that if*

$$\int_M |A^\circ|^2 d\mu < \sigma_1,$$

then M is totally umbilical.

Next we obtain a prior estimate on the lifespan of the surface diffusion flow in space forms in terms of the concentration of curvature at the initial time. Our proof is standard for the analysis of Willmore flow in Euclidean spaces and later in bounded Riemannian manifolds [Link 2013; Metzger et al. 2013].

Theorem 1.3. *Let $f : \Sigma \rightarrow \mathbb{F}^{2+p}(c)$ be a compact surface. There exist constants $\epsilon_1 > 0$, $C < \infty$ depending only on p , such that if $\rho > 0$ is chosen with*

$$\int_{f^{-1}(B_\rho(x))} |A|^2 d\mu \leq \epsilon \leq \epsilon_1(p) \quad \text{for any } x \in \mathbb{F}^{2+p}(c),$$

then the maximal time T for the surface diffusion flow with initial data f satisfies

$$T \geq \frac{1}{C} \rho^4,$$

and one has the estimate

$$\int_{f^{-1}(B_\rho(x))} |A|^2 d\mu \leq C\epsilon \quad \text{for } 0 \leq t \leq \frac{1}{C} \rho^4.$$

Then we perform a blowup at an assumed singularity and construct a static surface as a limit. Putting the above results together we get the following convergence theorem.

Theorem 1.4 (main theorem). *Let $f : \Sigma \rightarrow \mathbb{F}^{2+p}(c)$ ($c \geq 0$) be a compact surface. There exists a positive constant $\epsilon_0(p)$ such that if*

$$(1-5) \quad \int_\Sigma |A^\circ|^2 d\mu < \epsilon_0(p),$$

then the surface diffusion flow with initial data f exists smoothly for all time and converges exponentially to a round sphere as $t \rightarrow \infty$.

In particular, the convergence theorem for the surface diffusion flow implies the differentiable sphere theorem and the method can be also applied to the Willmore flow in a sphere. The key ingredient of the proof is to establish regularity and stability results in a space form. As we know, the equation of the diffusion flow is a fourth order equation so that tools related to the maximum principle are not available. Here we use the Sobolev inequality due to Michael and Simon [1973] and the Hölder inequality to classify the curvature terms especially for $\int_M |A^\circ|^n d\mu$, then we get the gap theorem for static submanifolds under the pinching condition for the Willmore functional. However, it contradicts the blowup limit at a finite time curvature singularity and thus we obtain the convergence theorem for the surface diffusion flow in the space form $\mathbb{F}^{2+p}(c)(c \geq 0)$.

2. Preparation

Let $(\mathbb{F}^{n+p}(c), \bar{g})$ be the space form with constant curvature c . For an immersion $f : M^n \rightarrow \mathbb{F}^{n+p}(c)$, the basic geometric data associated to f is the induced metric $g = f^*\bar{g}$ and $g(X, Y) = \langle Df \cdot X, Df \cdot Y \rangle$ with corresponding Levi-Civita connection ∇ . Let $\{e_i\}$ be a set of locally defined orthogonal bases and the summation over repeated indices is used. The second fundamental form is given by $A(X, Y) = D_{X,Y}^2 f = D_X(D_Y f) - Df \cdot \nabla_X Y$ with mean curvature vector given by the trace $H = A(e_i, e_i)$ and the trace free part $A^\circ(X, Y) = A(X, Y) - \frac{1}{n}g(X, Y)H$. We have the normal connection $\nabla_X \phi = (D_X \phi)^\perp$ which acts on the normal vector field ϕ along f . In (1-1) the Laplace operator $\Delta \phi = -\nabla^* \nabla \phi$ is understood with respect to the normal connection, where ∇^* denotes the formal adjoint of ∇ .

When computing tensor identities we freely use vector fields with first derivative vanishing at a given point. We define the curvature by $R^\perp(X, Y)\phi = \nabla_{X,Y}^2 \phi - \nabla_{Y,X}^2 \phi$ and the equations of Codazzi, Gauss and Ricci are

$$\begin{aligned}
 (\nabla_X A)(Y, Z) &= (\nabla_Y A)(X, Z), \\
 \nabla H &= -\nabla^* A = -\frac{n}{n-1} \nabla^* A^\circ, \\
 R(X, Y, Z, W) &= c\langle X, Z \rangle \langle Y, W \rangle - c\langle X, W \rangle \langle Y, Z \rangle \\
 &\quad + A(X, Z)A(Y, W) - A(X, W)A(Y, Z), \\
 R^\perp(X, Y)\phi &= A^\circ(e_i, X)\langle A^\circ(e_i, Y), \phi \rangle - A^\circ(e_i, Y)\langle A^\circ(e_i, X), \phi \rangle.
 \end{aligned}$$

The Codazzi equation implies that ∇A and $\nabla^2 A$ can be expressed by ∇A° and $\nabla^2 A^\circ$, respectively. In particular one has inequalities

$$(2-1) \quad |\nabla A| \leq C|\nabla A^\circ|, \quad |\nabla^2 A| \leq C|\nabla^2 A^\circ|.$$

If ϕ, ψ are normal forms, we denote by $\phi * \psi$ any normal-valued, multilinear form depending on ϕ, ψ in a universal, bilinear way. In particular, we have the properties $|\phi * \psi| \leq C|\phi| |\psi|$ and $\nabla(\phi * \psi) = \nabla\phi * \psi + \phi * \nabla\psi$. We use the notation P_r^m for any term of the type

$$(2-2) \quad P_r^m = \sum_{i_1 + \dots + i_r = m} \nabla^{i_1} A * \nabla^{i_2} A * \dots * \nabla^{i_r} A.$$

Now we derive a Sobolev type inequality in \mathbb{S}^{n+p} .

Lemma 2.1 [Michael and Simon 1973]. *Let M^n ($n \geq 2$) be a compact submanifold with or without boundary in the Euclidean space \mathbb{R}^{n+p} with $p \geq 1$. For a nonnegative function $g \in C^1(M)$ such that $g|_{\partial M} = 0$ if $\partial M \neq \emptyset$, we have*

$$\left[\int_M g^{\frac{n}{n-1}} d\mu \right]^{\frac{n-1}{n}} \leq D(n) \int_M (|\nabla g| + |H|g) d\mu,$$

where $D(n) = 4^{n+1} \sigma_n^{-1/n}$, and σ_n is the volume of the unit ball in \mathbb{R}^n .

Let M^n be a compact submanifold in \mathbb{S}^{n+p} . We consider the composition of isometric immersions $M^n \hookrightarrow \mathbb{S}^{n+p} \subset \mathbb{R}^{n+p+1}$ and denote by \bar{H} the mean curvature vector of M^n as a submanifold in \mathbb{R}^{n+p+1} . Then $|\bar{H}|^2 = |H|^2 + n^2$. Hence we have for any nonnegative function $f \in C^1(M)$,

$$\begin{aligned} \left[\int_M g^{\frac{n}{n-1}} d\mu \right]^{\frac{n-1}{n}} &\leq D(n) \int_M (|\nabla g| + |\bar{H}|g) d\mu \\ &\leq D(n) \int_M (|\nabla g| + (n + |H|)g) d\mu. \end{aligned}$$

3. Gap theorem

In this section, using the Sobolev inequality we get some integral estimates, and obtain the gap theorem. First, we need the following lemma.

Lemma 3.1 [Kuwert and Schätzle 2002]. *Let $f : M^n \rightarrow \mathbb{F}^{n+p}(c)$ be a complete submanifold, then for any l -linear normal form ϕ we have*

$$\begin{aligned} ((\nabla\nabla^* - \nabla^*\nabla)\phi)(X_1, X_2 \dots X_l) &= -(\nabla^*T)(X_1, X_2 \dots X_l) + (R^\perp(e_i, X_l)\phi)(e_i, X_2 \dots X_l) \\ &\quad - \sum_{k=1}^l \phi(X_i, \dots, R(e_i, X_l)X_k, \dots, X_l), \end{aligned}$$

where $T(X_0, X_1 \dots X_l) = (\nabla_{X_0}\phi)(X_1, X_2 \dots X_l) - (\nabla_{X_1}\phi)(X_0, X_2 \dots X_l)$.

We have the following Simons' identity in space form: $\mathbb{F}^{n+p}(c)$.

$$\Delta h_{ij} = \nabla^2 H + H \cdot h_{ip} h_{pj} - h_{ij} \cdot h_{pq} h_{pq} + h_{jq} \cdot h_{ip} h_{pq} - h_{jq} \cdot h_{pq} h_{pi} + cnA^\circ + R^\perp * A.$$

Therefore, we can write

$$(3-1) \quad \Delta A^\circ = \nabla^2 H + \frac{H^2}{n} A^\circ + A^\circ * A^\circ * A + cnA^\circ - \frac{\Delta H}{n} g.$$

Here we need to estimate the terms of $A^\circ * A^\circ * A$ and cnA° , which are different from the equation of surfaces in the Euclidean space. Taking $\phi = \nabla H$ in Lemma 3.1, and using the Gauss equation, we have

$$((\nabla \nabla^* - \nabla^* \nabla) \nabla H)(e_1) = A^\circ * A^\circ * \nabla A^\circ - \nabla H(R(e_i, e_1)e_i).$$

Since

$$\begin{aligned} -\langle R(e_i, e_1)e_i, e_k \rangle \nabla_k H &= c(n-1) \nabla_1 H + H \cdot h_{1k} \nabla_k H - h_{1i} \cdot h_{ik} \nabla_k H \\ &= c(n-1) \nabla H(e_1) + \frac{n-1}{n^2} H^2 \nabla H(e_1) + A * A^\circ * \nabla A^\circ, \end{aligned}$$

we have

$$(3-2) \quad \nabla^*(\nabla^2 H) + c(n-1) \nabla H = \nabla(\nabla^* \nabla H) - \frac{n-1}{n^2} H^2 \nabla H + A * A^\circ * \nabla A^\circ.$$

Taking $\phi = \nabla A^\circ$ in Lemma 3.1, we have

$$(3-3) \quad \nabla^*(\nabla^2 A^\circ) = \nabla(\nabla^* \nabla A^\circ) + A * A * \nabla A^\circ + cC \nabla A^\circ.$$

Replacing ϕ by $\nabla \phi$, we have

$$(3-4) \quad \Delta(\nabla \phi) - \nabla(\Delta \phi) = A * A * \nabla \phi + A * \nabla A * \phi + cC \nabla \phi.$$

Since the last terms of (3-2)–(3-4) are new compared with the equations in the Euclidean space, the next lemma is important in our proof.

Lemma 3.2. *If $f : M^n \rightarrow \mathbb{F}^{n+p}(c)$ is a complete submanifold with $\Delta H = F$ and $r \in C_c^1(M^n)$ satisfies $|\nabla r| \leq \Lambda$, then*

$$(3-5) \quad \int |\nabla A^\circ|^2 r^2 d\mu + c \int |A^\circ|^2 r^2 d\mu \leq \frac{C}{\Lambda^2} \int |F|^2 r^4 d\mu + \frac{C}{\Lambda^2} \int |A^\circ|^6 r^4 d\mu + C \Lambda^2 \int_{[r>0]} |A|^2 d\mu,$$

$$(3-6) \quad \int |\nabla A^\circ|^2 r^4 d\mu + c \int |A^\circ|^2 r^4 d\mu \leq C \int |\nabla H|^2 r^4 d\mu + C \int |A^\circ|^4 |A|^2 r^4 d\mu + C \Lambda^4 \int_{[r>0]} |A|^2 d\mu.$$

Proof. Multiplying (3-1) by $r^2 A^\circ$, we have

$$\begin{aligned} & \int \left(|\nabla A^\circ|^2 + cn|A^\circ|^2 + \frac{1}{n} H^2 |A^\circ|^2 \right) r^2 d\mu \\ &= \frac{n-1}{n} \int |\nabla H|^2 r^2 d\mu + C \int |A^\circ|^3 |A| r^2 d\mu + \int A^\circ * \nabla A^\circ * r * \nabla r d\mu \\ &\leq -\frac{n-1}{n} \int \langle H, \Delta H \rangle r^2 d\mu + \frac{C}{\Lambda^2} \int |A^\circ|^6 r^4 d\mu + \Lambda^2 \int_{[r>0]} |A|^2 d\mu + \frac{1}{2} \int |\nabla A^\circ|^2 r^2 d\mu \\ &\leq \frac{C}{\Lambda^2} \int |F|^2 r^4 d\mu + \frac{C}{\Lambda^2} \int |A^\circ|^6 r^4 d\mu + \Lambda^2 \int_{[r>0]} |A|^2 d\mu + \frac{1}{2} \int |\nabla A^\circ|^2 r^2 d\mu. \end{aligned}$$

Multiplying (3-1) by $r^4 A^\circ$, we get another inequality. □

Lemma 3.3. Under the assumption of Lemma 3.2 we have for $\eta = r^4$,

$$\begin{aligned} & \int (|\nabla^2 H|^2 + |A|^2 |\nabla A|^2 + |A|^4 |A^\circ|^2) \eta d\mu + c \int |\nabla H|^2 \eta d\mu \\ & \leq C \int (|A^\circ|^2 |\nabla A^\circ|^2 + |A^\circ|^6) \eta d\mu \\ & \quad + C \int |F|^2 \eta d\mu + \Lambda^4 \int_{[r>0]} |A|^2 d\mu. \end{aligned}$$

Proof. Multiplying (3-2) by $\nabla H \eta$, we have

$$\begin{aligned} (3-7) \quad & \int (|\nabla^2 H|^2 + c(n-1)|\nabla H|^2) \eta d\mu \\ & \leq \int (|\Delta H|^2 - \frac{n-1}{n^2} H^2 |\nabla H|^2) \eta d\mu \\ & \quad + C \int |A| |A^\circ| |\nabla A^\circ|^2 \eta d\mu + \int r^3 * \nabla r * \nabla H \nabla^2 H d\mu, \end{aligned}$$

and the proof is similar to Lemma 2.3 in [Kuwert and Schätzle 2002]. □

Proposition 3.4. Let $f : M^n \rightarrow \mathbb{F}^{n+p}(c)$ ($c \geq 0$) be a complete submanifold with $F = \Delta H$ and $r \in C_c^1(M)$ satisfying $s \geq 4$, and let $\Lambda = \|\nabla r\|_\infty$, then

$$\begin{aligned} & \int (|\nabla^2 A|^2 + |A|^2 |\nabla A|^2 + |A|^4 |A^\circ|^2) r^s d\mu + c \int (|A^\circ|^2 + |\nabla H|^2) r^s d\mu \\ & \leq C \int |F|^2 r^s d\mu + C \Lambda^4 \int_{[r>0]} |A|^2 d\mu \\ & \quad + C \int (|A^\circ|^2 |\nabla A^\circ|^2 + |A^\circ|^6) r^s d\mu. \end{aligned}$$

Proof. Multiplying (3-3) by $\nabla A^\circ \eta$, $\eta = r^4$, using (3-1) we have

$$\begin{aligned} & \int |\nabla^2 A^\circ|^2 \eta \, d\mu \\ &= \int (|\Delta A^\circ|^2 + |A|^2 |\nabla A^\circ|^2 + cC |\nabla A^\circ|^2 + r^{-1} * \nabla r * \nabla^2 A^\circ * \nabla A^\circ) \eta \, d\mu \\ &\leq \int (|\nabla^2 H|^2 + |A|^4 |A^\circ|^2 + |A|^2 |\nabla A^\circ|^2 + |F|^2) \eta \, d\mu \\ &\quad + \frac{1}{2} \int |\nabla^2 A^\circ|^2 \eta \, d\mu + \Lambda^2 \int |\nabla A^\circ|^2 r^2 \, d\mu + cC \int (|\nabla A^\circ|^2 + |A^\circ|^2) \eta \, d\mu. \end{aligned}$$

Using Lemma 3.2, we have

$$\begin{aligned} & \int |\nabla^2 A^\circ|^2 \eta \, d\mu + c \int |A^\circ|^2 \eta \, d\mu \\ &\leq \int (|\nabla^2 H|^2 + |A|^2 |\nabla A^\circ|^2 + |A|^4 |A^\circ|^2 + c|\nabla H|^2) \eta \, d\mu \\ &\quad + \int |F|^2 \eta \, d\mu + \int |A^\circ|^6 \eta \, d\mu + \Lambda^4 \int_{[r>0]} |A|^2 \, d\mu. \end{aligned}$$

The assertion follows from Lemma 3.3. □

Now we can prove Theorem 1.1.

Proof of Theorem 1.1. It follows from Proposition 3.4 that

$$\begin{aligned} & \int (|\nabla^2 A|^2 + |A|^2 |\nabla A|^2 + |A|^4 |A^\circ|^2) r^s \, d\mu + c \int (|A^\circ|^2 + |\nabla H|^2) r^s \, d\mu \\ &\leq C \int |F|^2 r^s \, d\mu + C\Lambda^4 \int_{[r>0]} |A|^2 \, d\mu + C \int (|A^\circ|^2 |\nabla A^\circ|^2 + |A^\circ|^6) r^s \, d\mu. \end{aligned}$$

Then we have

$$\int |A^\circ|^2 |\nabla A^\circ|^2 r^s \, d\mu \leq C \int |A^\circ|^6 r^s \, d\mu + C \int |\nabla A^\circ|^3 r^s \, d\mu,$$

and

$$\begin{aligned} (3-8) \quad & \int |\nabla A^\circ|^3 r^s \, d\mu \leq \int |A^\circ| |\nabla^2 A^\circ| |\nabla A^\circ| r^s \, d\mu + \Lambda \int |A^\circ| |\nabla A^\circ|^2 r^{s-1} \, d\mu \\ &\leq \int (\delta |\nabla^2 A^\circ|^2 + C_\delta |A^\circ|^2 |\nabla A^\circ|^2) r^s \, d\mu \\ &\quad + \Lambda \int |A^\circ|^{\frac{1}{2}} |\nabla A^\circ|^{\frac{1}{2}} |A^\circ|^{\frac{1}{2}} |\nabla A^\circ|^{\frac{3}{2}} r^{s-1} \, d\mu \\ &\leq \int (\delta |\nabla^2 A^\circ|^2 + C_\delta |A^\circ|^6 + \delta |\nabla A^\circ|^3) r^s \, d\mu \\ &\quad + \Lambda^{\frac{4}{3}} \int |A^\circ|^{\frac{2}{3}} |\nabla A^\circ|^2 r^{s-\frac{4}{3}} \, d\mu \\ &\leq \delta \int |\nabla^2 A^\circ|^2 r^s \, d\mu + C_\delta \int |A^\circ|^6 r^s \, d\mu + \Lambda^4 \int_{[r>0]} |A|^2 \, d\mu. \end{aligned}$$

Therefore,

$$\int |A^\circ|^2 |\nabla A^\circ|^2 r^s d\mu \leq \delta \int |\nabla^2 A^\circ|^2 r^s d\mu + C_\delta \int |A^\circ|^6 r^s d\mu + \Lambda^4 \int_{[r>0]} |A|^2 d\mu.$$

When $c = 0$, the Sobolev inequality is

$$\left[\int g^{\frac{n}{n-1}} d\mu \right]^{\frac{n-1}{n}} \leq C(n) \int (|\nabla g| + |H|g) d\mu.$$

If we take $g = |A^\circ|^{6(n-1)/n} r^{(n-1)/(n)s}$, then

$$\begin{aligned} (3-9) \quad & \left(\int |A^\circ|^6 r^s d\mu \right)^{\frac{n-1}{n}} \\ &= \left[\int (|A^\circ|^{\frac{6(n-1)}{n}} r^{\frac{n-1}{n}s})^{\frac{n}{n-1}} d\mu \right]^{\frac{n-1}{n}} \\ &\leq C(n) \int \left[\nabla (|A^\circ|^{\frac{6(n-1)}{n}} r^{\frac{n-1}{n}s}) + |H| |A^\circ|^{\frac{6(n-1)}{n}} r^{\frac{n-1}{n}s} \right] d\mu \\ &\leq \int |\nabla A^\circ| |A^\circ|^{\frac{5n-6}{n}} r^{\frac{n-1}{n}s} d\mu + \Lambda \int |A^\circ|^{\frac{6(n-1)}{n}} r^{\frac{n-1}{n}s-1} d\mu \\ &\quad + \int |A| |A^\circ|^{\frac{6(n-1)}{n}} r^{\frac{n-1}{n}s} d\mu = a + b + c. \end{aligned}$$

Here

$$\begin{aligned} a &= \int |\nabla A^\circ| |A^\circ|^{\frac{n}{3(n-1)}} r^{\frac{1}{3}s} |A^\circ|^{\frac{(2n-3)(7n-6)}{3n(n-1)}} r^{\frac{2n-3}{3n}s} d\mu \\ &\leq \int (|\nabla A^\circ|^{\frac{3(n-1)}{n}} |A^\circ| + |A^\circ|^{\frac{7n-6}{n}}) r^{\frac{n-1}{n}s} d\mu \\ &\leq \left(\int_{[r>0]} |A^\circ|^n d\mu \right)^{\frac{1}{n}} \left[\left(\int |\nabla A^\circ|^3 r^s d\mu \right)^{\frac{n-1}{n}} + \left(\int |A^\circ|^6 r^s d\mu \right)^{\frac{n-1}{n}} \right], \\ b &= \Lambda \int |A| |A^\circ|^{\frac{5n-6}{n}} r^{\frac{n-1}{n}s-1} d\mu \\ &\leq \left(\int_{[r>0]} |A^\circ|^n d\mu \right)^{\frac{1}{n}} \left(\Lambda^{\frac{n}{n-1}} \int |A^\circ|^{\frac{5n-6}{n-1}} r^{s-\frac{n}{n-1}} d\mu \right)^{\frac{n-1}{n}} \\ &= \left(\int_{[r>0]} |A^\circ|^n d\mu \right)^{\frac{1}{n}} \left(\Lambda^{\frac{n}{n-1}} \int |A^\circ|^{\frac{n}{2(n-1)}} |A^\circ|^{\frac{3(3n-4)}{2(n-1)}} r^{s-\frac{n}{n-1}} d\mu \right)^{\frac{n-1}{n}} \\ &\leq \left(\int_{[r>0]} |A^\circ|^n d\mu \right)^{\frac{1}{n}} \left[\Lambda^4 \left(\int_{[r>0]} |A^\circ|^2 d\mu \right) + \left(\int |A^\circ|^6 r^s d\mu \right) \right]^{\frac{n-1}{n}}, \end{aligned}$$

$$\begin{aligned} c &= \int |A| |A^\circ|^{\frac{6(n-1)}{n}} r^{\frac{n-1}{n}s} d\mu \\ &\leq \left(\int_{[r>0]} |A^\circ|^n d\mu \right)^{\frac{1}{n}} \left[\int (|A| |A^\circ|^{\frac{5n-6}{n}})^{\frac{n}{n-1}} r^s d\mu \right]^{\frac{n-1}{n}} \\ &\leq \left(\int_{[r>0]} |A^\circ|^n d\mu \right)^{\frac{1}{n}} \left(\int |A|^2 |A^\circ|^4 r^s d\mu \right)^{\frac{n-1}{n}}. \end{aligned}$$

When $c = 1$, the Sobolev inequality is

$$\left[\int g^{\frac{n}{n-1}} d\mu \right]^{\frac{n-1}{n}} \leq C(n) \int (|\nabla g| + |H|g + g) d\mu.$$

If we take $g = |A^\circ|^{6(n-1)/n} r^{(n-1)/(n)s}$, then the third term in the Sobolev inequality is

$$\begin{aligned} \int |A^\circ|^{\frac{6(n-1)}{n}} r^{\frac{n-1}{n}s} d\mu &= \int |A|^\circ |A^\circ|^{\frac{5n-6}{n}} r^{\frac{n-1}{n}s} d\mu \\ &\leq \left(\int_{[r>0]} |A^\circ|^n d\mu \right)^{\frac{1}{n}} \left(\int |A^\circ|^{\frac{5n-6}{n-1}} r^s d\mu \right)^{\frac{n-1}{n}} \\ &\leq \left(\int_{[r>0]} |A^\circ|^n d\mu \right)^{\frac{1}{n}} \left[\left(\int |A^\circ|^2 r^s d\mu \right) + \left(\int |A^\circ|^6 r^s d\mu \right) \right]^{\frac{n-1}{n}}. \end{aligned}$$

Combining the above estimates, we have

$$\begin{aligned} (3-10) \quad &\int (|A^\circ|^2 |\nabla A^\circ|^2 + |A^\circ|^6) r^s d\mu \\ &\leq \Lambda^4 (1 + \|A^\circ\|_{n,[r>0]}^{\frac{n}{n-1}}) \int_{[r>0]} |A|^2 d\mu + \delta \int |\nabla^2 A^\circ|^2 r^s d\mu \\ &\quad + \|A^\circ\|_{n,[r>0]}^{\frac{n}{n-1}} \int_M (|\nabla^2 A^\circ|^2 + |A^\circ|^6 + |A|^2 |A^\circ|^4 + c |A^\circ|^2) r^s d\mu. \end{aligned}$$

Now take the cutoff function $r(q) = \varphi(\gamma_x(f(q))/\rho)$, where γ_x is the distance function with respect to the fixed point $x \in \mathbb{F}^{2+p}$, $\varphi \in C^1(\mathbb{R})$ and $\chi_{B_{1/2}(x)} \leq \varphi \leq \chi_{B_1(x)}$; then $\Lambda = \frac{C}{\rho}$. From Proposition 3.4, $\|A^\circ\|_{n,[r>0]}^n < \sigma(n)$, and $s \geq 4$, we have

$$\begin{aligned} (3-11) \quad &\int (|\nabla^2 A|^2 + |A|^2 |\nabla A|^2 + |A|^4 |A^\circ|^2) r^s d\mu + c \int |A^\circ|^2 r^s d\mu \\ &\leq \int |F|^2 r^s d\mu + \left(\frac{C}{\rho}\right)^4 \int_{[r>0]} |A|^2 d\mu. \end{aligned}$$

Since f is a proper immersion, taking $\rho \rightarrow \infty$ and using a similar argument as in Theorem 8 of [Wheeler 2012], we finish the proof. \square

A surface in the sphere is a Willmore surface if and only if it satisfies

$$W(f) = \Delta H + Q(A^\circ)H = 0,$$

or else we get (4-1). The estimate of $W(f)$ is easy to handle as in [Kuwert and Schätzle 2002], so we also have the gap theorem for Willmore surfaces in a sphere.

Proof of Theorem 1.2. For $n = 2$ and $c = 1$, the gradient of the second fundamental form can be controlled (see [Andrews and Baker 2010]) by

$$(3-12) \quad \frac{1}{4}|\nabla H|^2 \leq \frac{1}{3}|\nabla A|^2 \leq |\nabla A^\circ|^2.$$

The precise forms of (3-1), (3-2) and (3-3) are

$$(3-13) \quad \nabla^* \nabla A^\circ + 2KA^\circ = -\nabla^2 H - R^\perp(e_k, \cdot)A(e_k, \cdot) + \frac{\Delta H}{2}g(\cdot, \cdot),$$

$$(3-14) \quad \nabla^*(\nabla^2 H) + K\nabla H = \nabla(\nabla^* \nabla H) - R^\perp(e_k, \cdot)\nabla H(e_k),$$

$$(3-15) \quad \langle \nabla^*(\nabla^2 A^\circ), \nabla A^\circ \rangle \leq \langle \nabla(\nabla^* \nabla A^\circ), \nabla A^\circ \rangle - 5|\nabla A^\circ|^2 \\ + 2|\nabla H|^2 + 12|A|^2|\nabla A|^2,$$

where the section curvature $K = 1 + |H|^2/4 - |A^\circ|^2/2$.

Multiplying (3-13) by $r^2 A^\circ$ with $r \in C_c^1(M^2)$ satisfies $\|\nabla r\|_\infty = \Lambda$. We have

$$\begin{aligned} & \int (|\nabla A^\circ|^2 + 2|A^\circ|^2 + \frac{1}{2}H^2|A^\circ|^2)r^2 d\mu \\ & \leq \frac{1}{2} \int |\nabla H|^2 r^2 d\mu + 2 \int |A^\circ|^4 r^2 d\mu + 6\Lambda \int |A^\circ| |\nabla A^\circ| r d\mu \\ & \leq -\frac{1}{2} \int \langle H, W \rangle r^2 d\mu + \frac{1}{2} \int \langle H, Q(A^\circ)H \rangle r^2 d\mu \\ & \quad + \frac{1}{128\Lambda^2} \int |A^\circ|^6 r^4 d\mu + C\Lambda^2 \int_{[r>0]} |A|^2 d\mu + \frac{1}{2} \int |\nabla A^\circ|^2 r^2 d\mu \\ & \leq \frac{C}{\Lambda^2} \int |W|^2 r^4 d\mu + \frac{1}{2} \int H^2 |A^\circ|^2 r^2 d\mu \\ & \quad + \frac{1}{128\Lambda^2} \int |A^\circ|^6 r^4 d\mu + C\Lambda^2 \int_{[r>0]} |A|^2 d\mu + \frac{1}{2} \int |\nabla A^\circ|^2 r^2 d\mu, \end{aligned}$$

and

$$(3-16) \quad \int (|\nabla A^\circ|^2 + 4|A^\circ|^2)r^2 d\mu \\ \leq \frac{C}{\Lambda^2} \int |W|^2 r^4 d\mu + \frac{1}{64\Lambda^2} \int |A^\circ|^6 r^4 d\mu + C\Lambda^2 \int_{[r>0]} |A|^2 d\mu.$$

Multiplying (3-13) by $r^4 A^\circ$, we have

$$(3-17) \quad \int (|\nabla A^\circ|^2 + |A^\circ|^2) r^4 d\mu \leq 2 \int |\nabla H|^2 r^4 d\mu + 2 \int |A^\circ|^6 r^4 d\mu + C \Lambda^4 \int_{[r>0]} |A|^2 d\mu.$$

Similarly, from the proof of Proposition 3.4 we have

$$(3-18) \quad \int \left(\frac{3}{4} |\nabla^2 A|^2 + |A^\circ|^2 + 20 |\nabla A^\circ|^2 + 40 |A|^2 |\nabla A|^2 + |A|^4 |A^\circ|^2 \right) r^4 d\mu \leq \int (3546 |A^\circ|^2 |\nabla A^\circ|^2 + 5230 |A^\circ|^6) r^4 d\mu + C \int |W|^2 r^4 d\mu + C \Lambda^4 \int_{[r>0]} |A|^2 d\mu.$$

Taking $g = |A^\circ|^3 r^2$ in the Sobolev inequality, we get

$$(3-19) \quad \frac{1}{D^2(2)} \int |A^\circ|^6 r^4 d\mu \leq \left[\int (3 |A^\circ|^2 |\nabla A^\circ|^2 r^2 + 2 \Lambda |A^\circ|^3 r + |H| |A^\circ|^3 r^2 + 2 |A^\circ|^3 r^2) d\mu \right]^2 \leq 5 \int_{[r>0]} |A^\circ|^2 d\mu \int (9 |A^\circ|^2 |\nabla A^\circ|^2 + 2 |A^\circ|^6 + 2 |A|^2 |A^\circ|^4 + |A^\circ|^2) r^4 d\mu + C \Lambda^4 \left(\int_{[r>0]} |A^\circ|^2 d\mu \right)^2.$$

Thus

$$(3-20) \quad \left(\frac{1}{5D^2(2)} - 2\sigma_1 \right) \int |A^\circ|^6 r^4 d\mu \leq \int_{[r>0]} |A^\circ|^2 d\mu \int (9 |A^\circ|^2 |\nabla A^\circ|^2 + 2 |A|^2 |A^\circ|^4 + |A^\circ|^2) r^4 d\mu + C \Lambda^4 \left(\int_{[r>0]} |A^\circ|^2 d\mu \right)^2.$$

Taking $g = |A^\circ| |\nabla A^\circ| r^2$ in the Sobolev inequality, we also obtain

$$(3-21) \quad \frac{1}{5D^2(2)} \int |A^\circ|^2 |\nabla A^\circ|^2 r^4 d\mu \leq \int_{[r>0]} |A^\circ|^2 d\mu \int \left(\frac{5}{2} |\nabla^2 A^\circ|^2 + 2 |A|^2 |\nabla A^\circ|^2 + 4 |\nabla A^\circ|^2 \right) r^4 d\mu + C \Lambda^4 \left(\int_{[r>0]} |A^\circ|^2 d\mu \right)^2.$$

It follows from (3-18), (3-20) and (3-21) and the assumption $\|A^\circ\|_2^2 < 6.23 \times 10^{-8}$ that

$$\int (|\nabla^2 A|^2 + |A|^2 |\nabla A|^2 + |A|^4 |A^\circ|^2) r^s d\mu + \int |A^\circ|^2 r^s d\mu \leq C \int |W|^2 r^s d\mu + C \Lambda^4 \int_{[r>0]} |A|^2 d\mu.$$

We take the cutoff function $r(q) = \varphi(\gamma_x(f(q))/(\rho))$, where γ_x is a distance function with respect to the fixed point $x \in \mathbb{S}^{2+p}$, $\varphi \in C^1(\mathbb{R})$ and $\chi_{B_{1/2}(x)} \leq \varphi \leq \chi_{B_1(x)}$; then $\Lambda = C/\rho$. Taking $\rho \rightarrow \infty$, we finish the proof. \square

4. Lifespan theorem

In this section, we give a lower bound on the lifespan of a smooth solution of the surface diffusion flow, which depends only on how much the curvature of the initial surface is concentrated. First, we recall these equations from [Kuwert and Schätzle 2002], and we can see the proof in [Link 2013].

Lemma 4.1. *Let $f : M^n \times [0, T) \rightarrow \mathbb{F}^{n+p}(c)$ be a smooth variation with normal velocity $\partial_t f = -V$, then*

$$\begin{aligned} \partial_t g(X Y) &= 2\langle A(X Y), V \rangle, \\ \partial_t (d\mu) &= \langle H, V \rangle d\mu, \\ \partial_t^\perp (\nabla_X \phi) &= \nabla_X \partial_t^\perp \phi - A(X e_i) \langle \nabla_{e_i} V, \phi \rangle - \nabla_{e_i} V \langle A(X e_i), \phi \rangle, \\ \partial_t^\perp (\nabla_X Y) &= \langle (\nabla_{e_i} A)(X, Y), V \rangle e_i - \langle A(X Y), \nabla_{e_i} V \rangle e_i \\ &\quad + \langle A(X e_i), \nabla_Y V \rangle e_i + \langle A(Y e_i), \nabla_X V \rangle e_i, \\ \partial_t^\perp A(X Y) &= -\nabla_{X,Y}^2 V + A(e_i, X) \langle A(e_i, Y), V \rangle - c g(X Y) V, \\ \partial_t^\perp H &= -\Delta V - A(e_i, e_j) \langle A(e_i, e_j), V \rangle - ncV. \end{aligned}$$

Using Gauss equation $2R_{1212} = 2c + H^2 - |A|^2$ and Lemma 4.1, the first variation formula for the Willmore functional in space forms in the normal direction $-\phi$ is

$$\begin{aligned} (4-1) \quad \frac{d}{d\epsilon} \mathcal{W}(f - \epsilon\phi)|_{\epsilon=0} &= \frac{d}{d\epsilon} \int_{\Sigma} |A^\circ|^2 d\mu|_{\epsilon=0} = \frac{d}{d\epsilon} \int_{\Sigma} \left(\frac{|H|^2}{2} + 2c \right) d\mu|_{\epsilon=0} \\ &= \int_{\Sigma} \langle H, -\Delta\phi - A(e_i, e_j) \langle A(e_i, e_j), \phi \rangle - 2c\phi + 2c\phi \rangle d\mu + \frac{1}{2} \int_{\Sigma} \langle H^3, \phi \rangle d\mu \\ &= - \int_{\Sigma} \langle \phi, \Delta H + Q(A^\circ)H \rangle d\mu, \end{aligned}$$

which proves $\text{grad}_{L^2} \mathcal{W}(f) = \Delta H + Q(A^\circ)H$.

The following lemmas will be needed for computing the evolution of derivatives of the curvature.

Lemma 4.2. *Let ϕ be an $(l - 1)$ -form with normal values along a variation $f : M^n \times I \rightarrow \mathbb{F}^{n+p}(c)$ with normal velocity $\partial_t f = -V$. If $\partial_t^\perp \phi + \Delta^2 \phi = Y$, then $\psi = \nabla \phi$ satisfies an equation*

$$(4-2) \quad \partial_t^\perp \psi + \Delta^2 \psi = \nabla Y + \sum_{i+j+k=3} \nabla^i A * \nabla^j A * \nabla^k \phi + A * \nabla V * \phi + \nabla A * V * \phi + C(\Delta(\nabla \phi) + \nabla(\Delta \phi)).$$

Proof. Let X_1, \dots, X_l be independent of t and such that $\nabla X_k = 0$ at a given point and a given time. Using Lemma 4.1 we have

$$\begin{aligned} (\partial_t^\perp \psi)(X_1, \dots, X_l) &= \partial_t^\perp((\nabla_{X_1} \phi)(X_2, \dots, X_l)) \\ &= \partial_t^\perp(\nabla_{X_1} \phi(X_2, \dots, X_l)) - \partial_t^\perp \sum_{k=2}^l \phi(X_2, \dots, \nabla_{X_1} X_k, \dots, X_l) \\ &= (\nabla(\partial_t^\perp \phi))(X_1, \dots, X_l) + A * \nabla V * \phi + \nabla A * V * \phi. \end{aligned}$$

So,

$$(4-3) \quad \partial_t^\perp \psi + \Delta^2 \psi - \nabla Y = \Delta^2(\nabla \phi) - \nabla(\Delta^2 \phi) + A * \nabla V * \phi + \nabla A * V * \phi.$$

Using (3-4), we have

$$\begin{aligned} \Delta^2(\nabla \phi) - \nabla(\Delta^2 \phi) &= \Delta(\Delta(\nabla \phi) - \nabla(\Delta \phi)) + \Delta(\nabla(\Delta \phi)) - \nabla(\Delta(\Delta \phi)) \\ &= \Delta(A * A * \nabla \phi + A * \nabla A * \phi + C \nabla \phi) + A * A * \nabla(\Delta \phi) \\ &\quad + A * \nabla A * (\Delta \phi) + C \nabla(\Delta \phi) \\ &= \sum_{i+j+k=3} \nabla^i A * \nabla^j A * \nabla^k \phi + C(\Delta(\nabla \phi) + \nabla(\Delta \phi)). \quad \square \end{aligned}$$

Lemma 4.3. *Let $f : \Sigma \times [0, T) \rightarrow \mathbb{F}^{2+p}(c)$ be a compact surface diffusion flow, then*

$$\partial_t^\perp(\nabla^m A) + \Delta^2(\nabla^m A) = P_3^{m+2}(A) + P_1^{m+2}(A),$$

for any $m \in \mathbb{N}_0$.

Proof. We proceed by induction on m , starting with $m = 0$,

$$\begin{aligned} \Delta(\nabla^2 H) - \nabla^2(\Delta H) &= (\nabla \nabla^* - \nabla^* \nabla) \nabla(\nabla H) + \nabla(\nabla \nabla^* - \nabla^* \nabla) \nabla H \\ &= A * A * \nabla^2 H + A * \nabla A * \nabla H + \nabla(A * A * \nabla H + A * \nabla A * \nabla H) + C \nabla^2 H, \end{aligned}$$

and

$$(4-4) \quad \begin{aligned} \partial_t^\perp A &= -\nabla^2(\Delta H) + A * A * \Delta H - c \Delta H g = -\Delta(\nabla^2 H) + P_3^2(A) + P_1^2(A) \\ &= -\Delta^2(A) + P_3^2(A) + P_1^2(A). \end{aligned}$$

Now let $m \geq 1$ and conclude from (4-2), that we have

$$\begin{aligned} \partial_t^\perp(\nabla^m A) + \Delta^2(\nabla^m A) &= \nabla(P_3^{m+1}(A) + P_1^{m+1}(A)) + \sum_{i+j+k=3} \nabla^i A * \nabla^j A * \nabla^k(\nabla^{m-1} A) \\ &\quad + A * \nabla(\Delta H) * \nabla^{m-1} A + \nabla A * \Delta H * \nabla^{m-1} A + \nabla^3(\nabla^{m-1} A), \end{aligned}$$

which yields the result. □

In the following we assume $r = \tilde{r} \circ f$, where $\tilde{r} \in C_c^1(B_\rho)$ and $\|\tilde{r}\|_{C^2} \leq C < \infty$. This implies

$$(4-5) \quad |\nabla r| \leq C, \quad |\nabla^2 r| \leq C(1 + |A|),$$

and for a ball $B_\rho = B_\rho(x_0) \subset \mathbb{F}^{2+p}(c)$ we use the notion

$$(4-6) \quad \Sigma_\rho(x_0) = f^{-1}(B_\rho(x_0)).$$

We denote by $\epsilon_1(p)$ the upper bound of the concentration of curvature for the initial surface, which is a positive constant depending only on the codimension p .

The next lemma can be proved much as in [Kuwert and Schätzle 2002].

Lemma 4.4. *Let $f : \Sigma \times [0, T) \rightarrow \mathbb{F}^{2+p}(c)$ be a compact surface diffusion flow, then for $\phi = \nabla^m A$ with $m \in \mathbb{N}$ and $s \geq 2m + 4$, we have*

$$\begin{aligned} \frac{d}{dt} \int |\phi|^2 r^s d\mu + \frac{3}{4} \int |\nabla^2 \phi|^2 r^s d\mu &\leq \int (P_3^{m+2}(A) + P_5^m(A) + P_1^{m+2}(A)) * \phi r^s d\mu + C \int |A|^2 r^{s-4-2m} d\mu. \end{aligned}$$

Remark 4.5. Suppose $\chi_{B_\rho(x)} \leq \tilde{r} \leq \chi_{B_{2\rho}(x)}$ and $\|D^j \tilde{r}\|_{L^\infty} \leq C' \rho^{-j}$ for $j = 1, 2$. Then we have $C = C' / (\rho^{4+2m})$ in Lemma 4.4.

Proposition 4.6. *Let $f : \Sigma \times [0, T) \rightarrow \mathbb{F}^{2+p}(c)$ be a compact surface diffusion flow, then for $\phi = \nabla^m A$ with $m \in \mathbb{N}$ and $s \geq 2m + 4$, we have*

$$(4-7) \quad \begin{aligned} \frac{d}{dt} \int |\phi|^2 r^s d\mu + \frac{1}{2} \int |\nabla^2 \phi|^2 r^s d\mu &\leq C(\|A\|_{\infty [r>0]}^4 + 1) \|A\|_{2 [r>0]}^2 + C \|A\|_{\infty [r>0]}^4 \int |\phi|^2 r^s d\mu. \end{aligned}$$

Proof. According to Lemma 4.4, we only need to show

$$\int (P_3^{m+2}(A) + P_5^m(A) + P_1^{m+2}(A)) * \phi r^s d\mu + C \int_{[r>0]} |A|^2 d\mu \leq \frac{1}{4} \int |\nabla^2 \phi|^2 r^s d\mu + C(\|A\|_\infty^4_{[r>0]} + 1) \|A\|_2^2_{[r>0]} + C \|A\|_\infty^4_{[r>0]} \int |\phi|^2 r^s d\mu,$$

and the proof is similar to Proposition 4.5 in [Kuwert and Schätzle 2002] with

$$(4-8) \quad \int P_1^{m+2}(A) * \phi r^s d\mu \leq \tau \int |\nabla^2 \phi|^2 r^s d\mu + C(\tau) \|A\|_2^2_{[r>0]}.$$

It follows from Corollaries 5.3 and 5.5 in [Kuwert and Schätzle 2002]. □

Since the Sobolev inequality is similar when in $\mathbb{F}^{2+p}(c)$ for $c \leq 0$, in the next lemma we just check the situation $c > 0$ (see [Kuwert and Schätzle 2002]).

Lemma 4.7. *For any normal 1-form ϕ on Σ and r as in (4-5),*

$$(4-9) \quad \|\phi\|_\infty^4_{[r=1]} \leq C \|\phi\|_2^2_{[r>0]} (\|\nabla^2 \phi\|_2^2_{[r>0]} + \|\phi^2 |A|^4\|_1_{[r>0]} + \|\phi\|_2^2_{[r>0]}).$$

Moreover, if $\phi = A$ and $\|A\|_2^2_{[r>0]} \leq \epsilon_1(p)$ for some ϵ_1 small enough depending on the constants in (4-5), then

$$(4-10) \quad \|A\|_\infty^4_{[r=1]} \leq C \|A\|_2^2_{[r>0]} (\|\nabla^2 A\|_2^2_{[r>0]} + \|A\|_2^2_{[r>0]}).$$

Proposition 4.8. *Let $f : \Sigma \times [0, T] \rightarrow \mathbb{F}^{2+p}(c)$ be a compact surface diffusion flow, then there exists an $\epsilon_1(p) > 0$ such that if*

$$\epsilon = \sup_{[0, T]} \int_{[r>0]} |A|^2 d\mu \leq \epsilon_1,$$

then for any $t \in [0, T]$ we have

$$(4-11) \quad \int_{[r=1]} |A|^2 d\mu + \frac{1}{4} \int_0^t \int_{[r=1]} (|\nabla^2 A|^2 + |A|^2 |\nabla A|^2 + |A|^6) d\mu dt \leq \int_{[r_0>0]} |A_0|^2 d\mu_0 + C\epsilon t.$$

Proof. We know from (3-10) that

$$\int (|A|^2 |\nabla A|^2 + |A|^6) r^4 d\mu \leq C \int_{[r>0]} |A|^2 d\mu + \frac{1}{4} \int_M |\nabla^2 A|^2 r^4 d\mu + \|A\|_{2, [r>0]}^2 \int_M (|\nabla^2 A|^2 + |A|^2 |\nabla A|^2 + |A|^6 + |A|^2) r^4 d\mu.$$

Choosing $m = 0$ in Lemma 4.4, we get

$$\begin{aligned} \frac{d}{dt} \int |A|^2 r^4 d\mu + \frac{3}{4} \int |\nabla^2 A|^2 r^4 d\mu \\ \leq \int (P_3^2(A) + P_5^0(A) + P_1^2(A)) * A r^4 d\mu + C \int_{[r>0]} |A|^2 d\mu. \end{aligned}$$

Combining these two inequalities, we have

$$\begin{aligned} \frac{d}{dt} \int |A|^2 r^4 d\mu + \frac{3}{4} \int (|\nabla^2 A|^2 + |A|^2 |\nabla A|^2 + |A|^6) r^4 d\mu \\ \leq C \int (|A|^3 |\nabla^2 A| + |A|^2 |\nabla A|^2 + |A|^6 + |\nabla^2 A| |A|) r^4 d\mu + C \int_{[r>0]} |A|^2 d\mu \\ \leq \|A\|_{2,[r>0]}^2 \int (|\nabla^2 A|^2 + |A|^2 |\nabla A|^2 + |A|^6) r^4 d\mu + \frac{1}{2} \int |\nabla^2 A|^2 d\mu \\ + C \int_{[r>0]} |A|^2 d\mu + C \left(\int_{[r>0]} |A|^2 d\mu \right)^2. \end{aligned}$$

Since $\int_{[r>0]} |A|^2 d\mu \leq \epsilon$, it yields

$$(4-12) \quad \frac{d}{dt} \int |A|^2 r^4 d\mu + \frac{1}{4} \int (|\nabla^2 A|^2 + |A|^2 |\nabla A|^2 + |A|^6) r^4 d\mu \leq C\epsilon.$$

The proposition follows by integrating over $[0, t]$. □

This proposition is the local estimate for the flow, from which we can see the variation of curvature is locally small. Furthermore, we have the following estimate as Proposition 3.4 in [Kuwert and Schätzle 2001] for the surface diffusion flow in $\mathbb{F}^{2+p}(c)$ ($c \geq 0$) with small initial Willmore energy.

Proposition 4.9. *Let $f : \Sigma \times [0, T] \rightarrow \mathbb{F}^{2+p}(c)$ ($c \geq 0$) be a compact surface diffusion flow with $\int_{\Sigma} |A|^2 d\mu \leq \aleph$, then there exist constants $\epsilon_0(p) \geq 0$ and $c_1 = C(p)/\aleph > 0$, such that if $\rho > 0$ is chosen with*

$$\int_{\Sigma_{\rho}} |A^{\circ}|^2 d\mu \leq \epsilon < \epsilon_0(p) \text{ at time } t = 0 \text{ for all } \Sigma_{\rho} \subset \mathbb{F}^{2+p},$$

then for any time $0 \leq t < t_1 = \min\{c_1 \rho^4, T\}$, we have

$$\int_{\Sigma_{\rho}} |A^{\circ}|^2 d\mu + \int_0^t \int_{\Sigma_{\rho}} (|\nabla^2 A|^2 + |A|^2 |\nabla A|^2 + |A|^6 + c |\nabla H|^2) d\mu dt \leq C(\epsilon + \aleph \rho^{-4} t).$$

Applying Proposition 4.6, Lemma 4.7 and Proposition 4.8, we get the higher order derivative estimate and the proof is similar to Proposition 4.6 in [Kuwert and Schätzle 2002].

Proposition 4.10. *Let $f : \Sigma \times [0, T] \rightarrow \mathbb{F}^{2+p}(c)$ be a compact surface diffusion flow. If*

$$\sup_{[0, T]} \int_{[r>0]} |A|^2 d\mu \leq \epsilon_1(p),$$

where $\epsilon_1(p)$ is small enough, then

$$\|\nabla^m A\|_{\infty [r=1]} \leq C(m, T, \alpha_0(m + 2)),$$

where $\alpha_0(m) = \sum_{j=0}^m \|\nabla^j A_0\|_{2 [r_0>0]}$.

Furthermore, we get the following proposition:

Proposition 4.11 (interior estimates). *Let $f : \Sigma \times (0, T] \rightarrow \mathbb{F}^{2+p}(c)$ be a compact surface diffusion flow satisfying the condition*

$$\sup_{(0, T]} \int_{\Sigma_\rho(x)} |A|^2 d\mu \leq \epsilon < \epsilon_1(p) \quad \text{for } x \in \mathbb{F}^{2+p}(c),$$

where $T \leq C\rho^4$, then for any $k \in \mathbb{N}_0$, at time $t \in (0, T]$, we have the estimates

$$\begin{aligned} \|\nabla^k A\|_{L^\infty(\Sigma_{\rho/2}(x))} &\leq c(k)\sqrt{\epsilon} t^{-\frac{k+1}{4}}, \\ \|\nabla^k A\|_{L^2(\Sigma_{\rho/2}(x))} &\leq c(k)\sqrt{\epsilon} t^{-\frac{k}{4}}. \end{aligned}$$

These are the higher order derivative estimate, which are localized in time. The proof follows similar to Theorem 3.5 in [Kuwert and Schätzle 2001].

Now we prove the lifespan theorem for the surface diffusion flow.

Proof of Theorem 1.3. We may assume that $\rho = 1$ by rescaling. Put

$$\epsilon(t) = \sup_{x \in \mathbb{F}^{2+p}} \int_{\Sigma_1(x)} |A|^2 d\mu.$$

By a trivial covering argument, we get

$$\epsilon(t) \leq \Gamma \cdot \sup_{x \in \mathbb{F}^{2+p}} \int_{\Sigma_{\frac{1}{2}}(x)} |A|^2 d\mu,$$

for some $\Gamma = \Gamma(p)$. The function $\epsilon : [0, T] \rightarrow \mathbb{R}$ is continuous by the compactness of $f(\Sigma \times [0, t])$ for $t < T$. Now let $\lambda > 0$ be a parameter, and define

$$t_0 : \sup\{0 \leq t \leq \min(T, \lambda) : \epsilon(\tau) \leq 3\Gamma\epsilon \text{ for } 0 \leq \tau < t\}.$$

The continuity of $\epsilon(t)$ implies $t_0 \geq 0$ and

$$(4-13) \quad \epsilon(t_0) = 3\Gamma\epsilon \text{ if } t_0 < \min(T, \lambda).$$

Fix a cutoff function $\tilde{r} \in C^2(\mathbb{F}^{2+p})$ with $\|\tilde{r}\|_{C^2(\mathbb{F}^{2+p})} \leq C$ and $\chi_{B_{1/2}(x)} \leq \tilde{r} \leq \chi_{B_1(x)}$, then $r = \tilde{r} \circ f$ satisfies condition (4-5). Thus, it follows from Proposition 4.8 that

$$\int_{\Sigma_{\frac{1}{2}}(x)} |A|^2(t) d\mu \leq \int_{\Sigma_1(x)} |A|^2(0) d\mu_0 + C\Gamma\epsilon t \leq 2\epsilon \quad \text{for } 0 \leq t \leq t_0.$$

Taking $\lambda = (C\Gamma)^{-1}$, we conclude that

$$\epsilon(t) \leq 2\Gamma\epsilon, \quad 0 \leq t \leq t_0,$$

and (4-13) implies $t_0 = \min(T, (C\Gamma)^{-1})$. Now if $t_0 = (C\Gamma)^{-1}$, we prove the proposition with a contraction for

$$(4-14) \quad t_0 = T.$$

We can apply Proposition 4.10 to obtain

$$\|\nabla^m A\|_\infty \leq C(p, m, f_0).$$

With the same argument as in [Magni 2015], we can get

$$\|\partial^m f\|_\infty, \|\partial^m \partial_t f\|_\infty \leq C(p, m, f_0).$$

Then $f(t)$ converges in $C^m(\Sigma)$ as $t \rightarrow T$ to a smooth function $f(T)$. By short time existence, we can extend the flow f to an interval $[0, T + \delta)$, which is contrary to the maximality of T . Hence it contradicts with (4-14). This proves Theorem 1.3. \square

5. Blowup analysis

In this section, we rescale the surface diffusion flow in $\mathbb{F}^{2+p}(c)$ ($c \geq 0$) at an assumed singularity, thereby constructing a static surface as a limit. First we need the following local area bound due to L. Simon [1993] in the Euclidean space, which has been generalized by F. Link [2013] in Riemannian manifolds.

Lemma 5.1. *Let $f : \Sigma \rightarrow \mathbb{F}^{2+p}(c)$ be a compact surface. Then for $0 < \rho < \infty$ and $\Sigma_\rho = \Sigma_\rho(x_0)$, one has*

$$(5-1) \quad \frac{\mu(\Sigma_\rho)}{\rho^2} \leq C \left(\int_\Sigma |A^\circ|^2 d\mu + 4\pi \chi(\Sigma) \right).$$

Moreover, the global area estimate is given as follows:

Proposition 5.2 (area estimate). *Let $f : \Sigma \times [0, T] \rightarrow \mathbb{F}^{2+p}(c)$ be a compact surface diffusion flow with $\int_\Sigma |A^\circ|^2 d\mu \leq \epsilon \leq \epsilon_0(p)$, where $\epsilon_0(p)$ is as in Proposition 4.9, then*

$$(5-2) \quad (1 - C\epsilon)\mu_0(\Sigma) \leq \mu(\Sigma) \leq \mu_0(\Sigma).$$

Proof. Using Lemma 4.1, we have

$$\frac{d}{dt} \int d\mu = - \int |\nabla H|^2 d\mu < 0,$$

so, $\mu(\Sigma) \leq \mu_0(\Sigma)$, and the area is decreasing. On the other hand, by the Sobolev inequalities we have

$$\int |\nabla H|^2 d\mu \leq C(n)\mu(\Sigma) \int |\nabla^2 H|^2 + |H|^2 |\nabla H|^2 + c|\nabla H|^2 d\mu.$$

We obtain

$$\frac{d}{dt} \mu(\Sigma) \geq -C\mu(\Sigma) \int |\nabla^2 H|^2 + |H|^2 |\nabla H|^2 + c|\nabla H|^2 d\mu.$$

Using Proposition 4.9 with $\rho = \infty$ implies

$$\mu(\Sigma) \geq \mu_0(\Sigma)e^{-C \int_0^t \int_{\Sigma} |\nabla^2 H|^2 + |H|^2 |\nabla H|^2 + c|\nabla H|^2 d\mu dt} \geq \mu_0(\Sigma)(1 - C\epsilon). \quad \square$$

Now we state the required compactness theorem, which was originally proved by Langer [1985] for surfaces. Recently it has been generalized by Breuning [2015] to any dimension and codimension in the Euclidean space and even to Riemannian setting, where the ambient space can be isometrically embedded into some Euclidean space with bounded second fundamental form. Here we use a simplified version from [Cooper 2011; Magni 2015].

Theorem 5.3 [Cooper 2011, Theorem 1.2]. *Given a compact surface Σ , a sequence of complete Riemannian manifolds $\{(N^{2+p}, \bar{g}_j)\}_{j \in \mathbb{N}}$ with uniformly bounded geometry and two sequences of points $\{q_j\}_{j \in \mathbb{N}} \subset \Sigma$ and $\{x_j\}_{j \in \mathbb{N}} \subset N^{2+p}$, let $f_j : (\Sigma, g_j) \rightarrow (N^{2+p}, \bar{g}_j)$ be a sequence of isometric proper immersions such that $f_j(q_j) = x_j$. Suppose that*

$$(5-3) \quad \mu_j(\Sigma_R(x_j)) \leq C(R) \quad \text{for any } R > 0,$$

$$(5-4) \quad \|\nabla^k A_j\|_{L^\infty(\Sigma_R)} \leq C_k(R) \quad \text{for any } R > 0 \text{ and any } k \in \mathbb{N}_0.$$

Then there exist a surface $\hat{\Sigma}$, a complete Riemannian manifold (M^{2+p}, \bar{g}) and two points $q \in \hat{\Sigma}$ and $x \in M^{2+p}$ such that:

- *There exists an increasing exhaustion $\{U_j\}_{j \in \mathbb{N}}$ of $\hat{\Sigma}$ made of open relatively compact sets, and there are diffeomorphisms $\phi_j : U_j \rightarrow \Sigma$ with $\phi_j(q) = q_j$, such that for any $R > 0$ we have $\Sigma_R^{\bar{g}_j}(q_j) \subset \phi_j(U_j)$, for all $j \geq j_0(R)$.*
- *There is an increasing exhaustion $\{V_j\}_{j \in \mathbb{N}}$ of M^{2+p} made of open relatively compact sets, and diffeomorphisms $\psi_j : V_j \rightarrow N^{2+p}$ with $\psi_j(x) = x_j$, such that for any $R > 0$ we have $B_R^{\bar{g}_j}(x_j) \subset \psi_j(V_j)$, for all $j \geq j_0(R)$.*
- *$\phi_j(U_j) \subset \psi_j(V_j)$ and $\psi_j^* \bar{g}_j \rightarrow \bar{g}$ smoothly.*

- *There exists a proper immersion $\hat{f} : \hat{\Sigma} \rightarrow M^n$ such that $\psi_j^{-1} \circ f_j \circ \phi_j \rightarrow \hat{f}$ smoothly with respect to a global isometric embedding of (M^{2+p}, \bar{g}) into a suitable Euclidean space \mathbb{R}^K . The immersion \hat{f} also satisfies (5-3) and (5-4) with respect to x .*

Here, the sequence of proper immersions f_j converging as in Theorem 5.3 to a proper immersion $\hat{f} : \hat{\Sigma} \rightarrow M^{2+p}$, will be denoted by $f_j \rightarrow \hat{f}$. Let $f : \Sigma \times [0, T) \rightarrow \mathbb{F}^{2+p}(c)$ be a surface diffusion flow defined on a compact surface Σ , where $0 < T \leq \infty$. Define

$$\mathfrak{N}(r, t) = \sup_{x \in \mathbb{F}^{2+p}} \int_{\Sigma_r(x)} |A(t)|^2 d\mu_t, \quad \Sigma_r(x) = f^{-1}(B_r(x)).$$

Choose an arbitrary sequence $r_j \searrow 0$ and assume concentration in the sense that for all j ,

$$(5-5) \quad t_j = \inf\{t \geq 0 : \mathfrak{N}(r_j, t) > \epsilon_2\} < T,$$

where $\epsilon_2 = \epsilon_1/C$, and ϵ_1 and C are the constants from the lifespan theorem. Clearly,

$$\int_{\Sigma_{r_j}(x)} |A(t_j)|^2 d\mu_{t_j} \leq \epsilon_2 \quad \text{for any } x \in \mathbb{F}^{2+p}(c).$$

On the other hand, choosing an appropriate sequence of balls at times $\tau \searrow t_j$, we find a point $x_j \in \mathbb{F}^{2+p}$ satisfying

$$\int_{f^{-1}(\overline{B_{r_j}(x_j)})} |A(t_j)|^2 d\mu_{t_j} \geq \epsilon_2.$$

Now we rescale by considering

$$f_j : (\Sigma, g_j) \times [-r_j^{-4}t_j, r_j^{-4}(T - t_j)] \rightarrow (\mathbb{F}^{2+p}, \bar{g}_j),$$

$$f_j(p, t) = f(p, t_j + r_j^4 t),$$

where $\bar{g}_j = r_j^{-2} \bar{g}$ and $g_j = f_j(\cdot, t)^* \bar{g}_j$. From Theorem 5.2 in [Magni 2015], we know that the rescaled flows converge locally smoothly on $\hat{\Sigma} \times \mathbb{R}$ to a static surface diffusion flow represented by a static properly immersed surface $\hat{f} : \hat{\Sigma} \rightarrow \mathbb{R}^{2+p}$, open sets with the property $\mathfrak{N}_j(1, t) \leq \epsilon_2$, for all $t \leq 0$ and

$$(5-6) \quad \int_{\hat{f}^{-1}(\overline{B_1(0)})} |\hat{A}|^2 d\hat{\mu} \geq \epsilon_2.$$

More precisely, the lifespan theorem yields $r_j^{-4}(T - t_j) \geq c_0$ and in fact

$$\mathfrak{N}_j(1, t) \leq \epsilon_1 \quad \text{for } 0 < t \leq c_0.$$

We may now apply Proposition 4.11 on parabolic cylinders $B_1(x) \times (t - 1, t]$ to obtain

$$\|\nabla^k A\|_{L^\infty} \leq c(k) \quad \text{for } r_j^{-4} + 1 \leq t \leq c_0.$$

Then we can apply the compactness theorem to the sequence

$$f_j = f_j(\cdot, 0) : \Sigma \rightarrow \mathbb{F}^{2+p},$$

thus obtaining a limit immersion $\hat{f}_0 : \hat{\Sigma} \rightarrow \mathbb{R}^{2+p}$. Then by reparametrization $f_j(\phi_j, \cdot)$ is a surface diffusion flow and has initial data converging locally in C^k to the immersion \hat{f}_0 . By standard estimates for geometric evolution equations in [Kuwert and Schätzle 2002], we deduce the locally smooth convergence $f_j \rightarrow \hat{f}$, where

$$\hat{f} : \hat{\Sigma} \times [0, c_0] \rightarrow \mathbb{R}^{2+p}$$

is a surface diffusion flow with initial data \hat{f}_0 .

In order to get the static blowup limit, we need the monotonicity of the Willmore functional.

Lemma 5.4. *Let $f : \Sigma \times (0, T] \rightarrow \mathbb{F}^{2+p}(c)$ be a compact surface diffusion flow with small initial Willmore energy satisfying Proposition 5.2; we have*

$$\begin{aligned} \frac{d}{dt} \int |A^\circ|^2 d\mu &\leq \frac{1}{2} \frac{d}{dt} \int |A|^2 d\mu \\ &\leq \frac{1}{2} \frac{d}{dt} \int |H|^2 d\mu \leq -\frac{1}{2} \int |\Delta H|^2 d\mu. \end{aligned}$$

Proof. The first two inequalities are obvious. From (3-7) and (3-8) with $r = 1$, we know

$$\begin{aligned} c \int |\nabla H|^2 d\mu &\leq C \int |A| |A^\circ| |\nabla A^\circ|^2 d\mu \leq C \int |A|^3 |A^\circ|^3 + |\nabla A^\circ|^3 d\mu \\ &\leq C \int |A|^4 |A^\circ|^2 d\mu + \delta \int |\nabla^2 A^\circ|^2 d\mu + C_\delta \int |A^\circ|^6 d\mu. \end{aligned}$$

From (3-11) we know

$$\int (|\nabla^2 A|^2 + |A|^2 |\nabla A|^2 + |A|^4 |A^\circ|^2) d\mu + c \int |A^\circ|^2 d\mu \leq \int |\Delta H|^2 d\mu,$$

and (3-10) implies

$$\begin{aligned} \int |A|^2 |A^\circ|^4 d\mu + c \int |\nabla H|^2 d\mu &\leq \int (\delta |A|^4 |A^\circ|^2 + C_\delta |A^\circ|^6 + c |\nabla H|^2) d\mu \\ &\leq (\delta + c\epsilon + \epsilon C_\delta) \int |\Delta H|^2 d\mu. \end{aligned}$$

So,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int |H|^2 d\mu &= - \int (\Delta H + H|A^\circ|^2 + 2cH)(\Delta H) d\mu \\ &\leq -\frac{3}{4} \int |\Delta H|^2 d\mu + 2 \int |A|^2 |A^\circ|^4 d\mu + 2c \int |\nabla H|^2 d\mu \\ &\leq -\frac{1}{2} \int |\Delta H|^2 d\mu. \end{aligned} \quad \square$$

Proposition 5.5. *Let $f : \Sigma \times (0, T] \rightarrow \mathbb{F}^{2+p}(c)$ be a compact surface diffusion flow with small initial Willmore energy, then the blow up \hat{f} which is constructed above is static.*

Proof. Let $U \Subset \hat{\Sigma}$ be an open set and ϕ_j be the diffeomorphisms in Theorem 5.3, then from Lemma 5.4 and the scale invariance of $\|A\|_2^2$,

$$\begin{aligned} \int_0^{c_0} \int_U |\Delta H(f_j(\phi_j, t))|^2 d\mu_{f_j(\phi_j, \cdot)} dt &= \int_0^{c_0} \int_{\phi_j(U)} |\Delta H_j|^2 d\mu_j dt \\ &\leq \int_\Sigma |A_j(0)|^2 d\mu_j - \int_\Sigma |A_j(c_0)|^2 d\mu_j = \int_\Sigma |A(t_j)|^2 d\mu - \int_\Sigma |A(t_j + r_j^4 c_0)|^2 d\mu, \end{aligned}$$

and it converges to zero as $j \rightarrow \infty$, Therefore $\Delta H(\hat{f}) \equiv 0$ and the blow up \hat{f} is static, which means that $\hat{f}(\cdot, t) = \hat{f}_0$. Furthermore (5-6) implies

$$\int_{\hat{f}^{-1}(B_1(0))} |\hat{A}|^2 d\hat{\mu} \geq \epsilon_2 > 0.$$

Thus \hat{f} is not totally geodesic. □

Lemma 5.6. *Let $\hat{f} : \hat{\Sigma} \rightarrow \mathbb{R}^{2+p}$ be the blowup constructed above. If $\hat{\Sigma}$ contains a compact component C , then in fact $\hat{\Sigma} = C$ and Σ is diffeomorphic to C .*

Proposition 5.7 (nontriviality of the blowup). *Let $\hat{f} : \hat{\Sigma} \rightarrow \mathbb{R}^{2+p}$ be the blowup of a compact surface diffusion flow as constructed above. Then none of the components of \hat{f} is compact. In particular, the blowup has a component which is a noncompact nonumbilical surface with $\Delta H \equiv 0$.*

Proof. Assume that there is a compact umbilical component of \hat{f} , then Lemma 5.6 implies \hat{f} has no further components. It follows that, up to diffeomorphism $\phi_i : \hat{\Sigma} \rightarrow \Sigma$, we have

$$\begin{aligned} \int_\Sigma |A^\circ(t_j)|^2 d\mu &= \int_\Sigma |A_j^\circ(0)|^2 d\mu_j \rightarrow 0, \\ \mu(t_j)(\Sigma) &= r_j \mu_j(0)(\Sigma) \rightarrow 0. \end{aligned}$$

This contradicts the area estimate. □

Now we prove the longtime existence. Combining the previous theorems, we can finally rule out concentration of curvature in finite time.

Proposition 5.8. *Let $f : \Sigma \times [0, T) \rightarrow \mathbb{F}^{2+p}(c)$ be a compact surface diffusion flow, then there exists a constant $\epsilon_0(p)$ such that if*

$$(5-7) \quad \int_{\Sigma} |A^\circ|^2 d\mu|_{t=0} < \epsilon_0 \leq \sigma(2),$$

then $T = \infty$ and there exists a radius $r_0 > 0$ such that

$$(5-8) \quad \int_{\Sigma_{r_0}(x)} |A|^2 d\mu < \epsilon_2,$$

for all $x \in \mathbb{F}^{2+p}$, $t \in [0, +\infty)$, where $\epsilon_2 > 0$ as in (5-5).

Proof. First, if curvature concentrates at time T , then we perform a blowup as above at T and get a static surface \hat{f} with small tracefree curvature. Now the gap theorem implies \hat{f} must be a plane or sphere, which contradicts the nontriviality of the blowup. There does not exist a finite time when curvature concentrates, so $T = \infty$.

Then we claim there exists a radius $r_0 > 0$, such that

$$\int_{\Sigma_{r_0}(x)} |A|^2 d\mu < \epsilon_2 \quad \text{for all } x \in \mathbb{F}^{2+p}, \quad t \in [0, \infty),$$

where $\epsilon_2 > 0$ is as in(5-5).

If not, in the same way we can perform a blowup as above, and get a static surface $\hat{f} : \hat{\Sigma} \rightarrow \mathbb{R}^{2+p}$ with

$$\int_{\hat{f}^{-1}(B_1(0))} |\hat{A}|^2 d\hat{\mu} \geq \epsilon_2,$$

whereas

$$\int_{\hat{\Sigma}} |\hat{A}^\circ|^2 d\hat{\mu} \leq \epsilon_0,$$

due to the scale invariance of $\|A^\circ\|_2^2$. Now the gap theorem implies \hat{f} must be totally umbilical, which contradicts the nontriviality of the blowup. □

Using a similar argument as in Proposition 5.5, we get the smooth convergence from Propositions 4.11 and 5.8.

Proof of Theorem 1.4. For any sequence $t_j \rightarrow \infty$, there exist $\phi_j \in \text{Diff}(\Sigma)$ such that, after passing to a subsequence, $f(\phi_j, t_j)$ converges smoothly to a static surface. The gap theorem implies it is a union of planes and spheres. After excluding several components as in [Kuwert and Schätzle 2001] and investigating the asymptotic behavior as in [Wheeler 2012], we get the global existence and exponential convergence for the surface diffusion flow with small initial Willmore energy. □

In particular, the convergence theorem implies a differentiable sphere theorem.

Corollary 5.9. *Let $f : \Sigma \rightarrow \mathbb{F}^{2+p}(c)$ ($c \geq 0$) be a compact surface. If there exists a positive constant $\epsilon_0(p)$ such that*

$$\int_{\Sigma} |A^\circ|^2 d\mu < \epsilon_0(p),$$

then Σ is diffeomorphic to the unit sphere.

The proof of Theorem 1.4 also applies to the Willmore flow in a sphere.

6. Open problems

In this section, we propose several problems for the convergence of the curvature flow. Applying the Morse theory of submanifolds, Shiohama and Xu [2000] obtained a topological sphere theorem for n -dimensional compact submanifolds in $\mathbb{F}^{n+p}(c)$ ($c \geq 0$) under the pinching condition for the Willmore functional. Thus, a natural problem is whether or not M^n is diffeomorphic to \mathbb{S}^n under the pinching condition for the Willmore functional. More precisely, we propose the following problem.

Conjecture 6.1. *Let $f : M^n \rightarrow \mathbb{F}^{n+p}(c)$ be a compact submanifold. There exists a positive constant $C_3(n)$ such that if*

$$(6-1) \quad \int_M |A^\circ|^n d\mu < C_3(n),$$

then the diffusion flow for submanifolds with initial data f exists smoothly for all time and converges to a round sphere as $t \rightarrow \infty$. In particular, M is diffeomorphic to \mathbb{S}^n .

Liu, Xu, Ye and Zhao [2018] investigated the convergence of the mean curvature flow of compact n -dimensional submanifolds in \mathbb{R}^{n+p} . They proved if the initial submanifold satisfies some suitable integral curvature conditions, then along the mean curvature flow it will shrink to a round point in finite time.

T. J. Willmore [1968] proved the Willmore inequality about the total mean curvature $\int |H|^2 d\mu$ for compact surfaces in \mathbb{R}^3 , and then B. Y. Chen [1971] obtained a general version of the Willmore inequality for compact submanifolds in \mathbb{R}^{n+p} , as follows:

Theorem 6.2. *If M is an n -dimensional compact submanifold in \mathbb{R}^{n+p} , then*

$$n^n \text{Vol}(\mathbb{S}^n) \leq \int_M |H|^n d\mu,$$

where the equality holds if and only if $M^n = \mathbb{S}^n(r)$.

Therefore, the total mean curvature for compact submanifolds in \mathbb{R}^{n+p} has a natural lower bound. Applying the Morse theory of submanifolds, Xu [2007] obtained a topological sphere theorem under the total mean curvature pinching condition. For other topological sphere theorems we can see [Shiohama and Xu 1994; 2000]. Similarly, we have the following problem for the diffusion flow under the total mean curvature pinching condition.

Conjecture 6.3. *Let $f : M^n \rightarrow \mathbb{R}^{n+p}$ be a compact submanifold. There exists a positive constant $C_4(n)$ such that if*

$$\int_M |H|^n d\mu < n^n \text{Vol}(\mathbb{S}^n) + C_4(n),$$

then the diffusion flow for submanifolds with initial data f exists smoothly for all time and converges to a round sphere as $t \rightarrow \infty$. In particular, M is diffeomorphic to \mathbb{S}^n .

For surfaces in \mathbb{R}^3 , the small Willmore energy assumption means a pinching condition for $\int |H|^2 d\mu$ and the convergence of the surface diffusion flow has answered these two problems. The Willmore conjecture verified by Marques and Neves [2014] says the integral of the square of the mean curvature of a torus in \mathbb{R}^3 is at least $8\pi^2$. In fact, they proved the integral inequality of the square of the mean curvature for any compact surfaces with genus greater than or equal to one. Thus, a strong version of the problem related to the Marques and Neves's theorem is: what is the best pinching constant in Conjecture 6.3 for compact surfaces in \mathbb{R}^3 ?

Conjecture 6.4. *Let $f : \Sigma \rightarrow \mathbb{R}^3$ be a compact surface satisfying*

$$\int_{\Sigma} |H|^2 d\mu < 8\pi^2,$$

then the surface diffusion flow with initial data f exists smoothly for all time and converges to a round sphere as $t \rightarrow \infty$. In particular, Σ is diffeomorphic to \mathbb{S}^2 .

Conjecture 6.5. *Let $f : \Sigma \rightarrow \mathbb{R}^3$ be a compact surface satisfying*

$$\int_{\Sigma} |H|^2 d\mu < 8\pi^2,$$

then the Willmore flow with initial data f exists smoothly for all time and converges to a round sphere as $t \rightarrow \infty$. In particular, Σ is diffeomorphic to \mathbb{S}^2 .

For higher dimensions, there are seldom results about the diffusion flow. However it provides a feasible method to handle the above problems so we can study the gradient flow of some constructed functionals. Li [2002] investigated the rigidity of the Willmore submanifold, which is the critical point of the Willmore functional for n -dimensional compact submanifolds in a sphere. As the Willmore flow for

surfaces, we can also study the higher-dimensional Willmore flow, which is the gradient flow of the Willmore functional for n -dimensional compact submanifolds in space forms. More precisely, we will study the following higher-dimensional Willmore flow

$$(6-2) \quad \frac{\partial f}{\partial t} = -\operatorname{grad} \mathcal{W}(f),$$

where $\operatorname{grad} \mathcal{W}(f)$ is the gradient vector field of the Willmore functional (1-2). We propose the following problems.

Conjecture 6.6. *Let $f : M^n \rightarrow \mathbb{F}^{n+p}(c)$ be a compact submanifold. There exists a positive constant $C_5(n)$ such that if*

$$(6-3) \quad \int_M |A^\circ|^n d\mu < C_5(n),$$

then the Willmore flow for submanifolds with initial data f exists smoothly for all time and converges to a round sphere as $t \rightarrow \infty$. In particular, M is diffeomorphic to \mathbb{S}^n .

Conjecture 6.7. *Let $f : M^n \rightarrow \mathbb{R}^{n+p}$ be a compact submanifold. There exists a positive constant $C_6(n)$ such that if*

$$\int_M |H|^n d\mu < n^n \operatorname{Vol}(\mathbb{S}^n) + C_6(n),$$

then the Willmore flow for submanifolds with initial data f exists smoothly for all time and converges to a round sphere as $t \rightarrow \infty$. In particular, M is diffeomorphic to \mathbb{S}^n .

Acknowledgment. The authors would like to thank the referee for valuable suggestions.

References

- [Andrews and Baker 2010] B. Andrews and C. Baker, “Mean curvature flow of pinched submanifolds to spheres”, *J. Differential Geom.* **85**:3 (2010), 357–395. MR Zbl
- [Bernard et al. 2019] Y. Bernard, G. Wheeler, and V.-M. Wheeler, “Concentration-compactness and finite-time singularities for Chen’s flow”, *J. Math. Sci. Univ. Tokyo* **26**:1 (2019), 55–139. MR Zbl
- [Brendle and Schoen 2009] S. Brendle and R. Schoen, “Manifolds with 1/4-pinched curvature are space forms”, *J. Amer. Math. Soc.* **22**:1 (2009), 287–307. MR Zbl
- [Breuning 2015] P. Breuning, “Immersions with bounded second fundamental form”, *J. Geom. Anal.* **25**:2 (2015), 1344–1386. MR Zbl
- [Chen 1971] B.-y. Chen, “On a theorem of Fenchel–Borsuk–Willmore–Chern–Lashof”, *Math. Ann.* **194** (1971), 19–26. MR Zbl
- [Cooper 2011] A. A. Cooper, “A compactness theorem for the second fundamental form”, preprint, 2011. arXiv

- [Escher et al. 1998] J. Escher, U. F. Mayer, and G. Simonett, “The surface diffusion flow for immersed hypersurfaces”, *SIAM J. Math. Anal.* **29**:6 (1998), 1419–1433. MR Zbl
- [Gu and Xu 2012] J.-R. Gu and H.-W. Xu, “The sphere theorems for manifolds with positive scalar curvature”, *J. Differential Geom.* **92**:3 (2012), 507–545. MR Zbl
- [Gu et al. 2017] J.-R. Gu, H.-W. Xu, Z.-Y. Xu, and E.-T. Zhao, “A survey on rigidity problems in geometry and topology of submanifolds”, pp. 79–99 in *Proceedings of the Sixth International Congress of Chinese Mathematicians*, vol. II, edited by C.-S. Lin et al., Adv. Lect. Math. **37**, International Press, Somerville, MA, 2017. MR Zbl
- [Hamilton 1982] R. S. Hamilton, “Three-manifolds with positive Ricci curvature”, *J. Differential Geometry* **17**:2 (1982), 255–306. MR Zbl
- [Huisken 1984] G. Huisken, “Flow by mean curvature of convex surfaces into spheres”, *J. Differential Geom.* **20**:1 (1984), 237–266. MR Zbl
- [Kuwert and Schätzle 2001] E. Kuwert and R. Schätzle, “The Willmore flow with small initial energy”, *J. Differential Geom.* **57**:3 (2001), 409–441. MR Zbl
- [Kuwert and Schätzle 2002] E. Kuwert and R. Schätzle, “Gradient flow for the Willmore functional”, *Comm. Anal. Geom.* **10**:2 (2002), 307–339. MR Zbl
- [Kuwert and Schätzle 2004] E. Kuwert and R. Schätzle, “Removability of point singularities of Willmore surfaces”, *Ann. of Math. (2)* **160**:1 (2004), 315–357. MR Zbl
- [Langer 1985] J. Langer, “A compactness theorem for surfaces with L_p -bounded second fundamental form”, *Math. Ann.* **270**:2 (1985), 223–234. MR Zbl
- [Li 2002] H. Li, “Willmore submanifolds in a sphere”, *Math. Res. Lett.* **9**:5-6 (2002), 771–790. MR Zbl
- [Link 2013] F. Link, *Gradient flow for the Willmore functional in Riemannian manifolds of bounded geometry*, Ph.D. thesis, Albert-Ludwigs-Universität Freiburg, 2013. Zbl arXiv
- [Liu et al. 2018] K. Liu, H. Xu, F. Ye, and E. Zhao, “The extension and convergence of mean curvature flow in higher codimension”, *Trans. Amer. Math. Soc.* **370**:3 (2018), 2231–2262. MR Zbl
- [Magni 2015] A. Magni, “A convergence result for the gradient flow of $\int |A|^2$ in Riemannian manifolds”, *Geom. Flows* **1**:1 (2015), 1–10. MR Zbl
- [Marques and Neves 2014] F. C. Marques and A. Neves, “Min-max theory and the Willmore conjecture”, *Ann. of Math. (2)* **179**:2 (2014), 683–782. MR Zbl
- [Mayer 2001] U. F. Mayer, “Numerical solutions for the surface diffusion flow in three space dimensions”, *Comput. Appl. Math.* **20**:3 (2001), 361–379. MR Zbl
- [Metzger et al. 2013] J. Metzger, G. Wheeler, and V.-M. Wheeler, “Willmore flow of surfaces in Riemannian spaces I: Concentration-compactness”, preprint, 2013. arXiv
- [Michael and Simon 1973] J. H. Michael and L. M. Simon, “Sobolev and mean-value inequalities on generalized submanifolds of R^n ”, *Comm. Pure Appl. Math.* **26** (1973), 361–379. MR Zbl
- [Mullins 1957] W. W. Mullins, “Theory of thermal grooving”, *J. Appl. Phys.* **28**:3 (1957), 333–339.
- [Shiohama and Xu 1994] K. Shiohama and H. W. Xu, “Rigidity and sphere theorems for submanifolds”, *Kyushu J. Math.* **48**:2 (1994), 291–306. MR Zbl
- [Shiohama and Xu 2000] K. Shiohama and H. W. Xu, “Rigidity and sphere theorems for submanifolds, II”, *Kyushu J. Math.* **54**:1 (2000), 103–109. MR Zbl
- [Simon 1993] L. Simon, “Existence of surfaces minimizing the Willmore functional”, *Comm. Anal. Geom.* **1**:2 (1993), 281–326. MR Zbl
- [Taylor and Cahn 1994] J. E. Taylor and J. W. Cahn, “Linking anisotropic sharp and diffuse surface motion laws via gradient flows”, *J. Statist. Phys.* **77**:1-2 (1994), 183–197. MR Zbl

- [Wang 2008] M.-T. Wang, “Lectures on mean curvature flows in higher codimensions”, pp. 525–543 in *Handbook of geometric analysis*, vol. 1, edited by L. Ji et al., Adv. Lect. Math. 7, International Press, Somerville, MA, 2008. MR Zbl
- [Wheeler 2012] G. Wheeler, “Surface diffusion flow near spheres”, *Calc. Var. Partial Differential Equations* **44**:1-2 (2012), 131–151. MR Zbl
- [Willmore 1968] T. J. Willmore, “Mean curvature of immersed surfaces”, *An. Şti. Univ. “Al. I. Cuza” Iaşi Sect. I a Mat. (N.S.)* **14** (1968), 99–103. MR Zbl
- [Xu 2007] H. W. Xu, “Mean value theorem for critical points and sphere theorems”, pp. 203–217 in *Proceedings of the 4th International Congress of Chinese Mathematicians*, vol. 2, 2007.
- [Xu and Gu 2007] H.-W. Xu and J.-R. Gu, “ L^2 -isolation phenomenon for complete surfaces arising from Yang–Mills theory”, *Lett. Math. Phys.* **80**:2 (2007), 115–126. MR Zbl
- [Xu and Gu 2013] H.-W. Xu and J.-R. Gu, “Geometric, topological and differentiable rigidity of submanifolds in space forms”, *Geom. Funct. Anal.* **23**:5 (2013), 1684–1703. MR Zbl
- [Xu and Yang 2016] H. Xu and D. Yang, “Rigidity theorem for Willmore surfaces in a sphere”, *Proc. Indian Acad. Sci. Math. Sci.* **126**:2 (2016), 253–260. MR Zbl

Received April 10, 2019. Revised December 11, 2019.

DONG PU
CENTER OF MATHEMATICAL SCIENCES
ZHEJIANG UNIVERSITY
HANGZHOU
CHINA
pudong@zju.edu.cn

HONGWEI XU
CENTER OF MATHEMATICAL SCIENCES
ZHEJIANG UNIVERSITY
HANGZHOU
CHINA
xuhw@zju.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 306 No. 1 May 2020

Tensor structure for Nori motives	1
LUCA BARBIERI-VIALE, ANNETTE HUBER and MIKE PREST	
On the Garden of Eden theorem for endomorphisms of symbolic algebraic varieties	31
TULLIO CECCHERINI-SILBERSTEIN, MICHEL COORNAERT and XUAN KIEN PHUNG	
Bergman kernels of elementary Reinhardt domains	67
DEBRAJ CHAKRABARTI, AUSTIN KONKEL, MEERA MAINKAR and EVAN MILLER	
Central splitting of manifolds with no conjugate points	95
JAMES DIBBLE	
On the global Gan–Gross–Prasad conjecture for general spin groups	115
MELISSA EMORY	
Schur algebras for the alternating group and Koszul duality	153
THANGAVELU GEETHA, AMRITANSHU PRASAD and SHRADDHA SRIVASTAVA	
A positive mass theorem for manifolds with boundary	185
SVEN HIRSCH and PENGZI MIAO	
Circle patterns on surfaces of finite topological type revisited	203
YUE-PING JIANG, QIANGHUA LUO and ZE ZHOU	
On some conjectures of Heywood	221
DONG LI	
Knapp–Stein dimension theorem for finite central covering groups	265
CAIHUA LUO	
Some classifications of biharmonic hypersurfaces with constant scalar curvature	281
SHUN MAETA and YE-LIN OU	
Surface diffusion flow of arbitrary codimension in space forms	291
DONG PU and HONGWEI XU	
Nonvanishing square-integrable automorphic cohomology classes: the case $GL(2)$ over a central division algebra	321
JOACHIM SCHWERMER	
Invariant Banach limits and applications to noncommutative geometry	357
EVGENII SEMENOV, FEDOR SUKOCHEV, ALEXANDR USACHEV and DMITRIY ZANIN	
A combinatorial identity for the Jacobian of t -shifted invariants	375
OKSANA YAKIMOVA	



0030-8730(202005)306:1;1-L