

*Pacific
Journal of
Mathematics*

TWO APPLICATIONS OF THE INTEGRAL REGULATOR

MATT KERR AND MUXI LI

Volume 306 No. 2

June 2020

TWO APPLICATIONS OF THE INTEGRAL REGULATOR

MATT KERR AND MUXI LI

We review Li’s refinement of the KLM regulator map, and use it to detect torsion phenomena in higher Chow groups of number fields and families of elliptic curves.

1. Introduction

Let X be a smooth projective variety defined over a subfield of \mathbb{C} . The KLM formula is a morphism of complexes inducing the Bloch–Beilinson regulator map

$$\mathrm{Gr}_\gamma^p K_n^{\mathrm{alg}}(X)_{\mathbb{Q}} \cong H_{\mathcal{M}}^{2p-n}(X, \mathbb{Q}(p)) \rightarrow H_{\mathcal{D}}^{2p-n}(X_{\mathbb{C}}^{\mathrm{an}}, \mathbb{Q}(p))$$

with rational coefficients, developed by the first author together with J. Lewis and S. Müller-Stach [Kerr 2003b; Kerr et al. 2006; Kerr and Lewis 2007] (see Section 3). It works by representing motivic cohomology by cycles in Bloch’s higher Chow complex (Section 2), and associating explicit currents on $X_{\mathbb{C}}^{\mathrm{an}}$ to them. The second author’s refinement now enables the direct computation of the *integral* regulator

$$H_{\mathcal{M}}^{2p-n}(X, \mathbb{Z}(p)) \rightarrow H_{\mathcal{D}}^{2p-n}(X, \mathbb{Z}(p))$$

on the level of higher Chow complexes [Li 2018].

The aim of this note is to offer a brief review of Li’s construction (Section 4) and give some first examples of how it may be used: First, we demonstrate how to find explicit torsion generators in higher Chow groups of number fields (Section 5); the results are summarized in Theorem 5.3. We also apply the formula (in Section 6) to *integrally* calculate a branch of the higher normal function arising from the mirror of local \mathbb{P}^2 (Proposition 6.4), along the way computing some integral regulator periods for curves of any genus (Theorem 6.2).

2. Higher Chow groups

Invented by Spencer Bloch [1986a; 1994] in the mid-1980s to geometrize Quillen’s higher algebraic K -theory, these generalize the usual Chow groups of cycles modulo

MSC2010: 14C25, 14C30, 19E15.

Keywords: higher Chow group, regulator map, normal function, integral regulator.

rational equivalence (the $n = 0$ case). In particular, for X smooth quasiprojective over a field k , they satisfy¹

$$\mathrm{CH}^p(X, n)_{\mathbb{Q}} \cong \mathrm{Gr}_{\mathcal{Y}}^p K_n^{\mathrm{alg}}(X)_{\mathbb{Q}}.$$

For such X (and no restriction on k), Voevodsky [2002] proved that they are *integrally* isomorphic to his motivic cohomology groups:

$$\mathrm{CH}^p(X, n) \cong H_{\mathcal{M}}^{2p-n}(X, \mathbb{Z}(p)).$$

Beyond their role in arithmetic geometry (e.g., Beilinson’s conjectures [1984]), they have recently shown up in several branches of physics (e.g., quantum field theory [Bloch et al. 2015] and topological string theory [del Ángel R. et al. 2019]) and mirror symmetry [Doran and Kerr 2014; Bloch et al. 2017]. We focus on the cubical presentation of $\mathrm{CH}^p(X, n)$ as the n -th homology of a complex of *higher Chow precycles* [Levine 1994]

$$\cdots \rightarrow Z^p(X, n+1) \xrightarrow{\partial} Z^p(X, n) \xrightarrow{\partial} Z^p(X, n-1) \rightarrow \cdots$$

or its (integrally quasi-isomorphic) subcomplex of *normalized precycles* [Bloch 2004]

$$\cdots \rightarrow N^p(X, n+1) \xrightarrow{\partial} N^p(X, n) \xrightarrow{\partial} N^p(X, n-1) \rightarrow \cdots .$$

A *higher Chow cycle* is an element of $\ker(\partial)$. Roughly speaking, these are relative codimension- p cycles on

$$(X \times \mathbb{A}^n, X \times \cup \mathbb{A}^{n-1}),$$

where the \mathbb{A}^{n-1} ’s are inserted into \mathbb{A}^n as a “cubical” configuration of hyperplanes. More precisely, writing $\{z_1, \dots, z_n\}$ for coordinates, $(z_i) = (z_i)_0 - (z_i)_\infty$ for their divisors and $|\cdot|$ for support, the *algebraic n -cube* is defined by

$$\square^n := (\mathbb{P}^1 \setminus \{1\})^n \supset \partial \square^n := \bigcup_i |(z_i)|.$$

Any component of $X \times \partial \square^n$ is called a *facet* of $X \times \square^n$, and a *face* is any intersection of facets. Now set²

$$\begin{aligned} c^p(X, n) &:= \{ \text{cycles in } Z^p(X \times \square^n) \text{ meeting faces of } X \times \partial \square^n \text{ properly,} \\ &\quad \text{i.e., in the expected codimension} \}, \\ d^p(X, n) &:= \sum_i \{ \text{cycles in } c^p(X, n) \text{ which are constant in } z_i \}, \end{aligned}$$

¹The subscript \mathbb{Q} denotes $\otimes \mathbb{Q}$.

²Normalized precycles may be represented (in $Z^p(X, n)$) by Z satisfying $Z \cdot \{z_i = 0\} = 0$ ($\forall i$) and $Z \cdot \{z_i = \infty\} = 0$ ($i < n$) simply by adding an element of $d^p(X, n)$.

$$Z^P(X, n) := c^P(X, n)/d^P(X, n),$$

$$N^P(X, n) := \{Z \mid Z \cdot (z_i)_0 = 0 \ (\forall i) \text{ and } Z \cdot (z_i)_\infty = 0 \ (\forall i < n)\},$$

and for $Z \in Z^P(X, n)$ or $N^P(X, n)$,

$$\partial Z := \sum_{i=1}^n (-1)^i (Z \cdot (z_i)_\infty - Z \cdot (z_i)_0).$$

If $X = \text{Spec}(k)$, write $Z^P(k, n)$ etc. for short.

Example 2.1. For any positive integer ℓ , let ζ_ℓ denote a primitive ℓ -th root of unity. Parametrize a cycle in $N^2(\mathbb{Q}(\zeta_\ell), 3)$ by $t \in \mathbb{P}^1$:

$$Z_\ell^2 := \left(1 - \frac{\zeta_\ell}{t}, 1 - t, t^{-\ell}\right).$$

Intersections with facets $\{z_i = 0, \infty\}$ are given by $t = 0, 1, \zeta_\ell, \infty$. But all these intersections have some $z_j = 1$, so are trivial (as $1 \notin \square$). We also record the cycle

$$\mathcal{Z}_5^2 := Z_1^2 + \left(1 - \frac{\zeta_5}{t}, 1 - t, t^{-5}\right) + \left(1 - \frac{\bar{\zeta}_5}{t}, 1 - t, t^{-5}\right)$$

in $N^2(\mathbb{Q}(\sqrt{5}), 3)$ for later reference.

3. Abel–Jacobi maps

These simultaneously generalize two classical invariants:³

- (1) Griffiths’ AJ map [1969]

$$\text{CH}^p(X, 0) \rightarrow H_{\mathcal{D}}^{2p}(X_{\mathbb{C}}^{\text{an}}, \mathbb{Z}(p))$$

for X smooth projective over \mathbb{C} ; and

- (2) Borel’s regulator map [Borel 1977; Burgos Gil 2002]

$$\text{CH}^p(k, 2p - 1) \rightarrow \mathbb{C}/\mathbb{Z}(p)$$

for $k \subset \mathbb{C}$ a number field.

Defined abstractly by Bloch [1986b], they map higher Chow groups to Deligne cohomology:

$$\text{CH}^p(X, n) \xrightarrow{\text{AJ}_X^{p,n}} H_{\mathcal{D}}^{2p-n}(X_{\mathbb{C}}^{\text{an}}, \mathbb{Z}(p)).$$

³The versions given here are, of course, not the original ones: (1) assembles Griffiths’ map (to the intermediate Jacobian $J^p(X)$) and the fundamental class map (to the Hodge classes $\text{Hg}^p(X)$) into a single invariant; while (2) is an integral lift of the composition $\text{CH}^p(k, 2p - 1) \rightarrow \text{CH}^p(k, 2p - 1)_{\mathbb{Q}} \cong \text{Gr}_\gamma^p K_{2p-1}(k)_{\mathbb{Q}} \xrightarrow{r_{\text{Bo}}} \mathbb{R} \cong \mathbb{C}/\mathbb{R}(p)$.

Here we are taking X to be smooth projective, and defined over a subfield of \mathbb{C} . (In the smooth quasiprojective setting, the correct target is the absolute Hodge cohomology $H_{\mathcal{H}}^{2p-n}(X_{\mathbb{C}}^{\text{an}}, \mathbb{Z}(p))$ [Kerr and Lewis 2007, §2].)

Kerr, Lewis, and Müller-Stach [Kerr et al. 2006] constructed a morphism of complexes, from a subcomplex of $Z^p(X, -\bullet)$ to a complex defining Deligne cohomology, which induces $\text{AJ}_X^{p,n}$. Writing C_{sing}^k for singular chains of real codimension k , D^k for currents of degree k , and F^\bullet for the Hodge filtration, their morphism takes the form

$$\begin{aligned} \widetilde{\text{AJ}}_{\text{KLM}}^{p,-\bullet} : Z_{\mathbb{R}}^p(X, -\bullet) &\rightarrow C_{\mathcal{D}}^{2p+\bullet}(X, \mathbb{Z}(p)) \\ &:= C_{\text{sing}}^{2p+\bullet}(X; \mathbb{Z}(p)) \oplus F^p D^{2p+\bullet}(X) \oplus D^{2p-1+\bullet}(X), \end{aligned}$$

with differential $D_{\mathcal{D}}(\alpha, \beta, \gamma) = (-\partial\alpha, -d\beta, d\gamma - \beta + \alpha)$ on the right. For $Z \in Z_{\mathbb{R}}^p(X, n)$ with projections π_1 (to \square^n) and π_2 (to X), they set

$$\begin{aligned} \widetilde{\text{AJ}}_{\text{KLM}}^{p,n}(Z) &:= (2\pi i)^{p-n}((2\pi i)^n T_Z, \Omega_Z, R_Z) \\ &:= (2\pi i)^{p-n}(\pi_2)_*(\pi_1)^*((2\pi i)^n T_n, \Omega_n, R_n), \end{aligned}$$

where $T_n := \bigcap_{i=1}^n T_{z_i} = \mathbb{R}_{<0}^{\times n}$, $\Omega_n := dz_1/z_1 \wedge \cdots \wedge dz_n/z_n$, and R_n is defined inductively by

$$R_n(z_1, \dots, z_n) := \log(z_1) \frac{dz_2}{z_2} \wedge \cdots \wedge \frac{dz_n}{z_n} - (-1)^n (2\pi i) R_{n-1}(z_2, \dots, z_n) \cdot \delta_{T_{z_1}}.$$

Here $\log(z_i)$ has a branch cut along $T_{z_i} = \{z_i \in \mathbb{R}_{<0}\}$, which is regarded as a 1-cochain on \square^n , oriented so that $\partial T_{z_i} = (z_i)_0 - (z_i)_\infty$; and $\delta_{T_{z_i}}$ is the current of integration along it. (Note that $\widetilde{\text{AJ}}_{\text{KLM}}^{p,n}$ vanishes identically on $d^p(X, n)$.)

The subcomplex $Z_{\mathbb{R}}^p(X, -\bullet) \subset Z^p(X, -\bullet)$ consists of cycles Z for which $Z_{\mathbb{C}}^{\text{an}}$ properly intersects certain combinations⁴ of $\{T_{z_i}\}$ and $\{|(z_j)|\}$; this is necessary in order for T_Z and R_Z to be well-defined. We call such precycles \mathbb{R} -proper. Kerr and Lewis [2007] proved the inclusion is a *rational* quasi-isomorphism, by appealing to Kleiman transversality in K -theory. Unfortunately, the claimed integral moving lemma in [Kerr et al. 2006] (which would have made this quasi-isomorphism integral) was incorrect, and [Kerr and Lewis 2007] was only written after a prolonged and unsuccessful effort to repair the integral version.

Now suppose we have a cycle $Z \in \ker(\partial) \subset Z_{\mathbb{R}}^p(X, n)$ with

$$[\widetilde{\text{AJ}}_{\text{KLM}}^{p,n}(Z)] \in H_{\mathcal{D}}^{2p-n}(X, \mathbb{Z}(p))$$

torsion of order exactly M . This implies $[Z] \in H_n\{Z_{\mathbb{R}}^p(X, \bullet)\}$ is at least of this order. But for $[Z] \in H_n\{Z^p(X, \bullet)\} = \text{CH}^p(X, n)$, it implies no such thing: there could be

⁴Namely, for each $J \subset \{1, \dots, n\}$ and $k \in \{0, \dots, n\}$, set $I := \{1, \dots, k\} \cap (\{1, \dots, n\} \setminus J)$; then we require that $Z_{\mathbb{C}}^{\text{an}}$ meet $(\bigcap_{i \in I} \{T_{z_i}\}) \cap (\bigcap_{j \in J} |(z_j)|)$ properly.

a $W \in Z^p(X, n + 1) \setminus Z_{\mathbb{R}}^p(X, n + 1)$ with $\partial W = Z$. So the KLM map only induces a homomorphism

$$AJ_{\mathbb{Q}}^{p,n} : CH^p(X, n) \rightarrow H_{\mathcal{D}}^{2p-n}(X, \mathbb{Q}(p))$$

consistent with Bloch’s $AJ^{p,n}$. This is frustrating, as the KLM formulas appear to be well-adapted to detecting torsion!

For $X = \text{Spec}(k)$ and $(p, n) = (2, 3)$, consider the portion

$$\begin{array}{ccccccc} \cdots & \rightarrow & Z_{\mathbb{R}}^2(k, 4) & \rightarrow & Z_{\mathbb{R}}^2(k, 3) & \rightarrow & Z_{\mathbb{R}}^2(k, 2) \rightarrow \cdots \\ & & \downarrow (2\pi i)^2 T_4 \cap (\cdot) & & \downarrow \frac{1}{2\pi i} \int_{(\cdot)} R_{2p-1} & & \downarrow 0 \\ \cdots & \rightarrow & \mathbb{Z}(2) & \hookrightarrow & \mathbb{C} & \longrightarrow & 0 \longrightarrow \cdots \end{array}$$

of the KLM map of complexes. We want to use the middle map to detect torsion. Denote its image on a cycle Z by $\mathcal{R}(Z) \in \mathbb{C}/\mathbb{Z}(2)$.

Example 3.1 [Petras 2009]. We calculate

$$\begin{aligned} \mathcal{R}(Z_{\ell}^2) &= \frac{1}{2\pi i} \int_{Z_{\ell}^2} R_3 \\ &= \frac{1}{2\pi i} \int_{Z_{\ell}^2} \left(\log(z_1) dz_2/z_2 \wedge dz_3/z_3 + (2\pi i) \log(z_2) dz_3/z_3 \cdot \delta_{T_{z_1}} \right. \\ &\qquad \qquad \left. + (2\pi i)^2 \log(z_3) \delta_{T_{z_1} \cap T_{z_2}} \right) \\ &= \int_{Z_{\ell} \cap T_{z_1}} \log(z_2) \frac{dz_3}{z_3} = - \int_{T_{1-\zeta_{\ell}/t}} \log(1-t) \frac{dt}{t} \\ &= - \int_0^{\zeta_{\ell}} \log(1-t) \frac{dt}{t} = \text{Li}_2(\zeta_{\ell}). \end{aligned}$$

For $\ell = 1$, this is $\pi^2/6 \in \mathbb{C}/\mathbb{Z}(2)$, which is 24-torsion, while (for the second cycle of Example 2.1) $\mathcal{R}(\mathcal{X}_5^2) = \text{Li}_2(1) + \text{Li}_2(\zeta_5) + \text{Li}_2(\bar{\zeta}_5) = 7\pi^2/30$ is 120-torsion. To deduce that these orders of torsion exist in $\text{CH}^2(\mathbb{Q}, 3)$ and $\text{CH}^2(\mathbb{Q}(\sqrt{5}), 3)$, rather than just in $H_3\{Z_{\mathbb{R}}^2(k, \bullet)\}$, we need an improvement in technology.

4. The integral regulator

Turning to potential strategies for effecting this improvement, one can probably rule out:

- (1) proving an integral moving lemma $(Z_{\mathbb{R}}^p(X, \bullet) \xrightarrow{\cong} Z^p(X, \bullet))$,
- and

- (2) extending KLM to a map of complexes on $Z^p(X, \bullet)$,

as too naive (in view of the history recounted in Section 3); and while

- (3) extending KLM to an infinite family of homotopic maps on nested subcomplexes with union $Z^P(X, \bullet)$,

seemed promising, what ultimately worked was

- (4) extending KLM to an infinite family of homotopic maps on nested subcomplexes with union $N^P(X, \bullet)$.

The heuristic idea of (3) was to perturb the branch cuts $T_{z_i} = \{z_i \in \mathbb{R}_{<0}\}$ in $\log(z_i)$ to $T_{z_i}^\epsilon = \{z_i/e^{i\epsilon} \in \mathbb{R}_{<0}\}$ (for small $\epsilon \in \mathbb{R}_{>0}$) and take a limit as $\epsilon \rightarrow 0$, an approach that had been successfully applied in [Kerr 2003a, §9]. (We write $\log^\epsilon(z_i)$ for the branch of logarithm along $T_{z_i}^\epsilon$.) Unfortunately, there are cycles in $Z^2(\mathbb{C}, 3)$ whose intersection with $T_{z_1}^\epsilon \cap T_{z_2}^\epsilon \cap T_{z_3}^\epsilon$ is improper for every real ϵ near 0 [Li 2018, §3]. So we need to deform the branches by distinct $\{\epsilon_i\}$ (viz., $T_{z_1}^{\epsilon_1} \cap T_{z_2}^{\epsilon_2} \cap T_{z_3}^{\epsilon_3}$); but then we cannot expect a morphism of complexes (or “limit” thereof) on $Z^P(X, \bullet)$. This forces us into strategy (4), and working with normalized subcomplexes.

Let \mathcal{B}_ϵ denote the set of infinite sequences $\underline{\epsilon} = \{\epsilon_i\}_{i>0}$, with

$$0 < \epsilon_1 < \epsilon, \quad 0 < \epsilon_2 < e^{-1/\epsilon_1}, \quad 0 < \epsilon_3 < e^{-1/\epsilon_2}, \quad \text{etc.,}$$

so that when $\epsilon \rightarrow 0$ its projection to any $(S^1)^n$ via $(e^{i\epsilon_1}, \dots, e^{i\epsilon_n})$ eventually avoids any given analytic subvariety. Let $N_\epsilon^P(X, \bullet) \subset N^P(X, \bullet)$ denote the subcomplexes of cycles Z with Z^{an} properly intersecting (for each $\underline{\epsilon} \in \mathcal{B}_\epsilon$) certain⁵ combinations of $\{T_{z_i}^{\epsilon_i}\}$ and $\{(z_j)_0, (z_j)_\infty\}$. Since for $\epsilon' < \epsilon$ we have $\mathcal{B}_{\epsilon'} \subset \mathcal{B}_\epsilon$, we also have $N_{\epsilon'}^P(X, \bullet) \supset N_\epsilon^P(X, \bullet)$.

Lemma 4.1 [Li 2018, Theorems 4.2 and 7.2]. *We have*

$$\bigcup_{\epsilon>0} N_\epsilon^P(X, n) = N^P(X, n) \quad \text{for all } n$$

and

$$\lim_{\epsilon \rightarrow 0} H_n(N_\epsilon^P(X, \bullet)) \cong H_n(N^P(X, \bullet)) \cong \text{CH}^P(X, n).$$

For any $\underline{\epsilon} \in \mathcal{B}_\epsilon$, replacing T_{z_i} by $T_{z_i}^{\epsilon_i}$ and $\log(z_i)$ by $\log^{\epsilon_i}(z_i)$ everywhere in the KLM formula yields a morphism of complexes

$$\widetilde{\text{AJ}}_{\epsilon, \underline{\epsilon}}^{P, -\bullet} : N_\epsilon^P(X, -\bullet) \rightarrow C_{\mathcal{D}}^{2p+\bullet}(X, \mathbb{Z}(p)).$$

Lemma 4.2 [Li 2018, Theorem 6.1]. *Given $\underline{\epsilon}, \underline{\epsilon}' \in \mathcal{B}_\epsilon$, $\widetilde{\text{AJ}}_{\epsilon, \underline{\epsilon}}^P$ and $\widetilde{\text{AJ}}_{\epsilon, \underline{\epsilon}'}^P$ are (\mathbb{Z}) -homotopic.*

Sketch of proof. Write $\bar{\square} := \mathbb{P}^1$ and fix an integer $N \gg 0$. For each multi-index $I \subset \{1, \dots, N\}$ and function $f : I \rightarrow \{0, \infty\}$, define $\partial_f^I \bar{\square}^N := \bigcap_{i \in I} \{z_i = f(i)\} \subset \bar{\square}^N$;

⁵as in the last footnote

and let $\rho_{i,I}^{f(i)} : \partial_f^I \bar{\square}^N \hookrightarrow \partial_f^{I \setminus i} \bar{\square}^N$ be the inclusions. Then we can define a double complex (zero outside $0 \leq a \leq N$)

$$E^{a,b} := \bigoplus_{|I|=N-a} \bigoplus_{f: I \rightarrow \{0, \infty\}} C_{\mathcal{D}}^{2a+b}(\partial_f^I \bar{\square}^N, \mathbb{Z}(a))$$

with $D_{\mathcal{D}}$ the vertical differential, $\delta := \sum_{I,f,i} (-1)^{\text{sgn}_I(i) + \text{sgn}(f(i))} (\rho_{i,I}^{f(i)})_*$ the horizontal differential, and $\mathbb{D} := D_{\mathcal{D}} + (-1)^b \delta$ the total differential.

Put $\underline{\epsilon}_I := \{\epsilon_j\}_{j \notin I}$ and

$$\mathcal{R}_{I,f}^{\underline{\epsilon}} := ((2\pi i)^a T_a^{\underline{\epsilon}}, \Omega_a, R_a^{\underline{\epsilon}}) \in C_{\mathcal{D}}^a(\partial_f^I \bar{\square}^N, \mathbb{Z}(a)) \quad (\cong C_{\mathcal{D}}^a((\mathbb{P}^1)^a, \mathbb{Z}(a))),$$

so that $\mathcal{R}_a^{\underline{\epsilon}} := \{\mathcal{R}_{I,f}^{\underline{\epsilon}}\}_{I,f}^{|I|=N-a}$ belongs to $E^{a,-a}$. Adapting [Kerr et al. 2006, (5.2)–(5.4)] to these perturbed currents, [Li 2018] deduces that $\mathcal{R}_{\square}^{\underline{\epsilon}} := \{\mathcal{R}_a^{\underline{\epsilon}}\}_{0 \leq a \leq N}$ is a 0-cocycle in $E^{\bullet,\bullet}$. The key point of [op. cit.] is then to construct a (-1) -cochain $S_{\square}^{\underline{\epsilon}, \underline{\epsilon}'}$ in $E^{\bullet,\bullet}$ with $\mathbb{D} S_{\square}^{\underline{\epsilon}, \underline{\epsilon}'} = \mathcal{R}_{\square}^{\underline{\epsilon}} - \mathcal{R}_{\square}^{\underline{\epsilon}'}$, and with respect to which the precycles in $N_{\varepsilon}^p(X, \bullet)$ remain proper. This yields at once a homotopy formula in $C_{\mathcal{D}}^{\bullet}(X, \mathbb{Z}(\bullet))$ of the form

$$D_{\mathcal{D}} S_Z^{\underline{\epsilon}, \underline{\epsilon}'} - S_{\partial Z}^{\underline{\epsilon}, \underline{\epsilon}'} = \widetilde{\text{AJ}}_{\varepsilon, \underline{\epsilon}}^p(Z) - \widetilde{\text{AJ}}_{\varepsilon, \underline{\epsilon}'}^p(Z).$$

The construction of $S_{\square}^{\underline{\epsilon}, \underline{\epsilon}'}$ proceeds by replacing each KLM triple

$$((2\pi i)^a T_a^{\underline{\epsilon}}, \Omega_a, R_a^{\underline{\epsilon}}) = \left(2\pi i T_{z_1}^{\varepsilon_1}, \frac{dz_1}{z_1}, \log^{\varepsilon_1}(z_1)\right) \cup \dots \cup \left(2\pi i T_{z_a}^{\varepsilon_a}, \frac{dz_a}{z_a}, \log^{\varepsilon_a}(z_a)\right)$$

by an alternating sum of cup-products in which $(-2\pi i \delta_{\{\arg(z_i) \in (\varepsilon_i, \varepsilon'_i)\}}, 0, 0)$ replaces the i -th term. See [op. cit.] for full details. \square

We therefore have well-defined, compatible maps

$$\text{AJ}_{\varepsilon}^{p,n} : H_n(N_{\varepsilon}^p(X, \bullet)) \rightarrow H_{\mathcal{D}}^{2p-n}(X, \mathbb{Z}(p))$$

for each $\varepsilon > 0$, essentially given by $\lim_{\varepsilon \rightarrow 0} \widetilde{\text{AJ}}_{\varepsilon}^p$ (where the limit is taken so that $\varepsilon > \varepsilon_1 \gg \varepsilon_2 \gg \varepsilon_3 \gg \dots > 0$), and recovering $\widetilde{\text{AJ}}_{\text{KLM}}^p$ on

$$N_{\mathbb{R}}^p(X, \bullet) := Z_{\mathbb{R}}^p(X, \bullet) \cap N^p(X, \bullet).$$

More precisely:

Theorem 4.3 [Li 2018, §7]. *The $\{\widetilde{\text{AJ}}_{\varepsilon, \underline{\epsilon}}^{p, -\bullet}\}$ induce a homomorphism*

$$\text{AJ}_{\mathbb{Z}}^{p,n} : \text{CH}^p(X, n) \cong \lim_{\varepsilon \rightarrow 0} H_n(N_{\varepsilon}^p(X, \bullet)) \rightarrow H_{\mathcal{D}}^{2p-n}(X, \mathbb{Z}(p))$$

factoring $\text{AJ}_{\mathbb{Q}}^{p,n}$.

Corollary 4.4. *If a class $\xi \in \text{CH}^p(X, n)$ is represented by*

$$Z \in \ker(\partial) \subset N_{\mathbb{R}}^p(X, n) \left(\subset \bigcap_{\varepsilon > 0} N_{\varepsilon}^p(X, n) \right),$$

then

$$\text{AJ}_{\mathbb{Z}}^{p,n}(\xi) = \lim_{\epsilon \rightarrow 0} \widetilde{\text{AJ}}_{\epsilon}^{p,n}(Z) = \widetilde{\text{AJ}}_{\text{KLM}}^{p,n}(Z).$$

So the KLM formula holds verbatim on normalized, \mathbb{R} -proper representatives, validating the deductions at the end of Example 3.1. It is this statement that we (primarily) use in the applications that follow.

5. Torsion generators

Let $\mu_{\infty} = \bigcup_{m \in \mathbb{N}} \mu_m \subset \mathbb{C}^*$ denote the roots of unity, and $w_r(k) := |(\mu_{\infty}^{\otimes r})^{\text{Gal}(\bar{\mathbb{Q}}/k)}|$ for any number field $k \subset \mathbb{C}$. By the universal coefficient sequence for motivic cohomology

$$H_{\mathcal{M}}^0(k, \mathbb{Z}(r)) \rightarrow H_{\mathcal{M}}^0(k, \mathbb{Z}/m\mathbb{Z}(r)) \rightarrow H_{\mathcal{M}}^1(k, \mathbb{Z}(r)) \xrightarrow{m} H_{\mathcal{M}}^1(k, \mathbb{Z}(r))$$

and vanishing of $H_{\mathcal{M}}^0(k, \mathbb{Z}(r))$,⁶ we have

$$\text{CH}^r(k, 2r - 1)[m] \cong H_{\mathcal{M}}^1(k, \mathbb{Z}(r))[m] \cong H_{\mathcal{M}}^0(k, \mathbb{Z}/m\mathbb{Z}(r)).$$

Since the norm residue map

$$H_{\mathcal{M}}^0(k, \mathbb{Z}/m\mathbb{Z}(r)) \rightarrow H_{\text{ét}}^0(k, \mu_m^{\otimes r}) \cong (\mu_m^{\otimes r})^{\text{Gal}(\bar{\mathbb{Q}}/k)}$$

is an isomorphism by a celebrated theorem of Rost-Voevodsky (see [Haesemeyer and Weibel 2019]), we conclude that $\text{CH}^r(k, 2r - 1)[m] \cong \mathbb{Z}/(m, w_r(k))\mathbb{Z}$, hence

$$\text{CH}^r(k, 2r - 1)_{\text{tors}} \cong \mathbb{Z}/w_r(k)\mathbb{Z}.$$

Example 5.1. If $k = \mathbb{Q}$, one has⁷ $w_{2n}(k) = \text{denominator of } (|B_{2n}|)/4n$ (written in lowest terms) and $w_{2n+1} = 2$ for $n \geq 1$; so

$$\text{CH}^2(\mathbb{Q}, 3) \cong \mathbb{Z}/24\mathbb{Z}, \quad \text{CH}^3(\mathbb{Q}, 5)_{\text{tors}} \cong \mathbb{Z}/2\mathbb{Z}, \quad \text{CH}^4(\mathbb{Q}, 7) \cong \mathbb{Z}/240\mathbb{Z}.$$

For real quadratic fields $k = \mathbb{Q}(\sqrt{d})$, the situation is more complicated (see [Weibel 2013, §VI.2]); one computes for instance

d	2	3	5	7
$w_2(\mathbb{Q}(\sqrt{d}))$	48	24	120	24
$w_4(\mathbb{Q}(\sqrt{d}))$	480	240	240	240

⁶See [Weibel 2013, Exercise VI.4.6]: since $H_{\mathcal{M}}^{-1}(k, \mathbb{Z}/m\mathbb{Z}(r)) \cong H_{\text{ét}}^{-1}(k, \mu_m^{\otimes r}) = \{0\}$, $\cdot m$ is injective on $H_{\mathcal{M}}^0(k, \mathbb{Z}(r))$ (universal coefficient sequence), which is thus torsion-free; it has rank 0 since $H_{\mathcal{M}}^0(k, \mathbb{Q}(r)) \cong \text{Gr}_r^{\vee} K_{2r}(k) \otimes \mathbb{Q} = \{0\}$ by Borel’s theorem [1974].

⁷Bernoulli numbers: $|B_{2n}| = \frac{1}{6}, \frac{1}{30}, \frac{1}{42}, \dots$ for $n = 1, 2, 3, \dots$

so that only $\text{CH}^2(\mathbb{Q}(\sqrt{2}), 3)$, $\text{CH}^2(\mathbb{Q}(\sqrt{5}), 3)$ and $\text{CH}^4(\mathbb{Q}(\sqrt{2}), 7)$ are different from the $k = \mathbb{Q}$ case. Finally, for cyclotomic $k = \mathbb{Q}(\zeta_p)$ ($p \geq 5$ prime) one can show that $w_3(\mathbb{Q}(\zeta_p)) = 2p$, while $w_3(\mathbb{Q}(\zeta_3)) = 18$.

For computing torsion orders of images under

$$\text{AJ}_{\mathbb{Z}}^{r, 2r-1} : H_{2r-1}(N_{\mathbb{R}}^r(k, \bullet)) \rightarrow \mathbb{C}/(2\pi i)^r \mathbb{Z}, \quad Z \mapsto \frac{1}{(2\pi i)^{r-1}} \int_Z R_{2r-1} =: \mathcal{R}(Z)$$

we use the following basic calculation:

Proposition 5.2. *Suppose that for a given $r \in \mathbb{N}$ there exists a collection of closed precycles $Z_{\ell, a}^r \in N_{\mathbb{R}}^r(\mathbb{Q}(\zeta_{\ell}), 2r - 1)$ with*

$$(5-1) \quad \mathcal{R}(Z_{\ell, a}^r) = (r - 1)! \ell^{r-1} \text{Li}_r(\zeta_{\ell}^a).$$

Then for $Z := \sum_{a=0}^{\ell-1} f(a) Z_{\ell, a}^r$ with $f(-a) = (-1)^r f(a)$, $\text{AJ}(Z)$ is torsion of order given by the denominator of

$$(5-2) \quad \tau(Z) := |(2\pi i)^{-r} \mathcal{R}(Z)| = \pm \frac{\ell^{r-1}}{2^r} \sum_{a=0}^{\ell-1} f(a) B_r(a/\ell),$$

where $B_r(\cdot)$ are the Bernoulli polynomials.⁸

Proof. Writing $\hat{f}(k) := \sum_{a=0}^{\ell-1} f(a) \zeta_{\ell}^{-ka}$ for the finite Fourier transform (for functions on $\mathbb{Z}/\ell\mathbb{Z}$), we have

$$\begin{aligned} \sum_{a=0}^{\ell-1} f(a) \text{Li}_r(\zeta_{\ell}^a) &= \frac{1}{2} \sum_{a=0}^{\ell-1} f(a) \sum_{k \in \mathbb{Z} \setminus \{0\}} \frac{\zeta_{\ell}^{ka}}{k^r} \\ &= \frac{(-1)^r}{2} \sum_{k \in \mathbb{Z} \setminus \{0\}} \frac{\hat{f}(k)}{k^r} =: \frac{(-1)^r}{2} \tilde{L}(\hat{f}, r) \\ &= \frac{(2\pi i)^r}{2 \cdot r!} \sum_{a=0}^{\ell-1} f(a) B_r(a/\ell), \end{aligned}$$

where the last step is [Kerr 2018, Theorem 3.9]. □

In practice, the $\{Z_{\ell, a}^r\}$ will be obtained from a single cycle $Z_{\ell}^r = Z_{\ell, 1}^r$ by Galois conjugation. For $r = 2$, we already have this from Examples 2.1 and 3.1: namely,

$$(5-3) \quad Z_{\ell, a}^2 = \left(1 - \frac{\zeta_{\ell}^a}{t}, 1 - t, t^{-\ell}\right).$$

Now $\text{CH}^r(k, 2r - 1) = \text{CH}^r(k, 2r - 1)_{\text{tors}} \Leftrightarrow r$ is even and k is totally real. In particular, assuming the hypothesis of Proposition 5.2, we obtain generators of

⁸ $B_2(x) = x^2 - x + \frac{1}{6}$, $B_3(x) = x^3 - \frac{3}{2}x^2 + \frac{1}{2}x$, $B_4(x) = x^4 - 2x^3 + x^2 - \frac{1}{30}$, etc.

$\text{CH}^{2n}(k, 4n - 1)$ as follows:

$$\begin{aligned} k = \mathbb{Q}, \text{ any } n : \quad & \tau(Z_1^{2n}) = \frac{|B_{2n}|}{4n} (= \frac{1}{24}, \frac{1}{240}, \dots); \\ k = \mathbb{Q}(\sqrt{2}), n = 1, 2 : \quad & \tau(Z_{8,1}^{2n} + Z_{8,7}^{2n}) = \frac{11}{48}, \frac{1313}{480}; \\ k = \mathbb{Q}(\sqrt{5}), n = 1 : \quad & \tau(Z_1^2 + Z_{5,1}^2 + Z_{5,4}^2) = \frac{7}{120}. \end{aligned}$$

For $\text{CH}^3(k, 5)_{\text{tors}}$ one computes, for example,

$$\begin{aligned} k = \mathbb{Q}(\zeta_3) : \quad & \tau(Z_{3,1}^3 - Z_{3,2}^3) = \frac{1}{9}; \quad \text{and} \\ k = \mathbb{Q}(\zeta_5) : \quad & \tau(Z_{5,1}^3 - Z_{5,4}^3) = \frac{2}{5}, \end{aligned}$$

which miss only the 2-torsion element from $\text{CH}^3(\mathbb{Q}, 5) \hookrightarrow \text{CH}^3(k, 5)$. (So far we have no $N_{\mathbb{R}}^3(\mathbb{Q}, 5)$ representative for this element.)

It remains to construct the cycles of the Proposition for $r = 3, 4$. From [Kerr and Yang 2018, §4.2], for $r = 3$ we have⁹

$$(5-4) \quad Z_{\ell}^3 := -2 \left(\frac{t_1}{t_1 - 1}, \frac{t_2}{t_2 - 1}, 1 - \zeta_{\ell} t_1 t_2, t_1^{\ell}, t_2^{\ell} \right) - \left(\frac{t}{t - 1}, \frac{1}{1 - \zeta_{\ell} t}, \frac{(u - t^{\ell})(u - t^{-\ell})}{(u - 1)^2}, t^{\ell} u, \frac{u}{t^{\ell}} \right)$$

which is normalized since all ‘‘boundaries’’ occur in the third coordinate. We have $\mathcal{R}(Z_{\ell}^3) = 2\text{Li}_3(\zeta_{\ell})$ by [op. cit., Theorem 3.6], with only the first term contributing. (This gives in particular $\mathcal{R}(Z_1^3) = 2\text{Li}_3(1) = 2\zeta(3)$.)

For $r = 4$, the first construction in [Kerr and Yang 2018, §4.3] would be in $N_{\mathbb{R}}^4(\mathbb{Q}(\zeta_{\ell}), 7)$, but there is an error in the computation of the boundary of the last component \mathcal{W}_2 : in fact, it is degenerate,¹⁰ and the cycle \mathcal{Z} is therefore not closed. A correct application of the strategy in [op. cit., §3.1], yields

$$(5-5) \quad \begin{aligned} Z_{\ell}^4 := & 6 \left(\frac{t_1}{t_1 - 1}, \frac{t_2}{t_2 - 1}, \frac{t_3}{t_3 - 1}, 1 - \zeta_{\ell} t_1 t_2 t_3, t_1^{\ell}, t_2^{\ell}, t_3^{\ell} \right) \\ & + \left(\frac{t_1}{t_1 - 1}, \frac{t_2}{t_2 - 1}, \frac{1}{1 - \zeta_{\ell} t_1 t_2}, \frac{(u - t_1^{\ell})(u - t_2^{\ell})(u - t_1^{-\ell} t_2^{-\ell})}{(u - 1)^3}, \frac{t_1^{\ell}}{u}, \frac{t_2^{\ell}}{u}, \frac{1}{u t_1^{\ell} t_2^{\ell}} \right) \\ & + \left(\frac{t_1}{t_1 - 1}, \frac{t_2}{t_2 - 1}, \frac{1}{1 - \zeta_{\ell} t_1 t_2}, \frac{(u - t_1^{\ell})(u - t_2^{\ell})}{(u - t_1^{\ell} t_2^{\ell})(u - 1)}, \frac{t_1^{\ell}}{u}, \frac{t_2^{\ell}}{u}, \frac{u}{t_1^{\ell} t_2^{\ell}} \right) \\ & + \left(\frac{t_1}{t_1 - 1}, \frac{t_2}{t_2 - 1}, \frac{1}{1 - \zeta_{\ell} t_1 t_2}, \frac{(u - t_1^{\ell})(u - t_1^{-\ell} t_2^{-\ell})}{(u - t_2^{-\ell})(u - 1)}, \frac{t_1^{\ell}}{u}, t_2^{\ell} u, \frac{1}{u t_1^{\ell} t_2^{\ell}} \right) \end{aligned}$$

⁹The components are parametrized by (t_1, t_2) and (t, u) respectively.

¹⁰I.e., it belongs to $d^4(\mathbb{Q}(\zeta_{\ell}), 7)$, as can be seen by substituting $v = uw$.

$$\begin{aligned}
 & + \left(\frac{t_1}{t_1 - 1}, \frac{t_2}{t_2 - 1}, \frac{1}{1 - \zeta_\ell t_1 t_2}, \frac{(u - t_2^\ell)(u - t_1^{-\ell} t_2^{-\ell})}{(u - t_1^{-\ell})(u - 1)}, t_1^\ell u, \frac{t_2^\ell}{u}, \frac{1}{t_1^\ell t_2^\ell u} \right) \\
 & + \left(\frac{(v - u)(v - u^{-1})}{(v - 1)^2}, \frac{t}{t - 1}, \frac{1}{1 - \zeta t}, \frac{(u - t^\ell)(u - t^{-\ell})}{(u - 1)^2}, v, \frac{t^\ell}{u}, \frac{1}{ut^\ell} \right) \\
 & + \left(\frac{t}{t - 1}, \frac{(v - u)(v - u^{-1})}{(v - 1)^2}, \frac{1}{1 - \zeta t}, \frac{(u - t^\ell)(u - t^{-\ell})}{(u - 1)^2}, \frac{t^\ell}{u}, v, \frac{1}{ut^\ell} \right) \\
 & + \left(\frac{t}{t - 1}, \frac{1}{1 - \zeta t}, \frac{(v - u)(v - u^{-1})}{(v - 1)^2}, \frac{(u - t^\ell)(u - t^{-\ell})}{(u - 1)^2}, \frac{t^\ell}{u}, \frac{1}{ut^\ell}, v \right),
 \end{aligned}$$

which belongs to $\ker(\partial) \cap N_{\mathbb{R}}^4(\mathbb{Q}(\zeta_\ell), 7)$. Only the first term contributes to $\mathcal{R}(Z_\ell^4) = -6\ell^3 \int_{[0,1]^3} \log(1 - \zeta_\ell t_1 t_2 t_3) dt_1/t_1 \wedge dt_2/t_2 \wedge dt_3/t_3 = 6\ell^3 \text{Li}_4(\zeta_\ell)$, see [op. cit., §3.2]. Again, cycles $Z_{\ell,a}^3$ and $Z_{\ell,a}^4$ are obtained by replacing ζ_ℓ by ζ_ℓ^a in (5-4) and (5-5), respectively.

Summarizing, we have

Theorem 5.3. *The cycles given in (5-3), (5-4), (5-5) generate the (cyclic, torsion) groups as shown in Table 1.*

Remark 5.4. An example of a cycle for which the log-branch perturbations are required for the integral regulator computation is

$$Z := Z_- - Z_+ := \left(\left(\frac{z - i}{z + i} \right)^{-2}, \left(\frac{z - 1}{z + 1} \right)^{-2}, z^{-2} \right) - \left(\left(\frac{z - i}{z + i} \right)^2, \left(\frac{z - 1}{z + 1} \right)^2, z^2 \right)$$

(parametrized by $z \in \mathbb{P}^1$) in $N^2(\mathbb{Q}(i), 3)$. Indeed, T_{z^2} has support on $i\mathbb{R}$ in the complex z -plane, and $T_{((z-1)/(z+1))^2}$ and $T_{((z-i)/(z+i))^2}$ have support on the unit circle S^1 in the complex z -plane, so that the triple intersection (essentially $i\mathbb{R} \cap S^1 \cap S^1$) is nonempty. Though this cycle is *non-torsion*, we briefly describe the computation. After making the deformation, $T_{((z-i)/(z+i))^2}$ and $T_{((z-1)/(z+1))^2}$ intersect twice with

Group	Cycle	Order
$\text{CH}^2(\mathbb{Q}, 3)$	Z_1^2	24
$\text{CH}^4(\mathbb{Q}, 7)$	Z_1^4	240
$\text{CH}^2(\mathbb{Q}(\sqrt{2}), 3)$	$Z_{8,1}^2 + Z_{8,7}^2$	48
$\text{CH}^4(\mathbb{Q}(\sqrt{2}), 7)$	$Z_{8,1}^4 + Z_{8,7}^4$	480
$\text{CH}^2(\mathbb{Q}(\sqrt{5}), 3)$	$Z_1^2 + Z_{5,1}^2 + Z_{5,4}^2$	120
$\frac{\text{CH}^3(\mathbb{Q}(\zeta_3), 5)_{\text{tors}}}{\text{CH}^3(\mathbb{Q}, 5)_{\text{tors}}}$	$Z_{3,1}^3 - Z_{3,2}^3$	9
$\frac{\text{CH}^3(\mathbb{Q}(\zeta_5), 5)_{\text{tors}}}{\text{CH}^3(\mathbb{Q}, 5)_{\text{tors}}}$	$Z_{5,1}^3 - Z_{5,4}^3$	5

Table 1. Some Generators (see Theorem 5.3).

opposite orientations, at points near i and $-i$ with phase just greater than $\pi/2$ and $3\pi/2$, respectively. Since $\epsilon_3 \rightarrow 0$ much faster than ϵ_1 and ϵ_2 , in the limit the

$$2\pi i \int_{Z_+} \log^{\epsilon_3}(z^2) \delta_{T^{\epsilon_1}_{((z-i)/(z+i))^2} \cap T^{\epsilon_2}_{((z-1)/(z+1))^2}}$$

term of $\frac{1}{2\pi i} \int_{Z_+} R_3$ contributes $2\pi i(\pi i - \pi i) = 0$. The remaining term yields

$$(5-6) \quad 2 \int_{Z_+} \log^{\epsilon_2} \left(\left(\frac{1-z}{1+z} \right)^2 \right) \frac{dz}{z} \delta_{T^{\epsilon_1}_{((z-i)/(z+i))^2}},$$

where $T^{\epsilon_1}_{((z-i)/(z+i))^2}$ consists of two paths from $-i$ to i , along which one checks that (in the limit)

$$\log^{\epsilon_2} \left(\left(\frac{1-z}{1+z} \right)^2 \right) = 2 \log^{\epsilon_2}(1-z) - 2 \log^{\epsilon_2}(1+z);$$

and so (5-6) becomes $8 \int_{-i}^i \log(1-z) \frac{dz}{z} - 8 \int_{-i}^i \log(1+z) \frac{dz}{z}$. Combining this with the portion from Z_- , we obtain

$$\mathcal{R}(Z) = 32\text{Li}_2(i) - 32\text{Li}_2(-i) = 64iL(\chi_4, 2) \in \mathbb{C}/\mathbb{Z}(2).$$

6. Local \mathbb{P}^2 revisited

For a reflexive polytope $\Delta \subset \mathbb{R}^2$ with polar polytope Δ° , the mirror of $K_{\mathbb{P}_{\Delta^\circ}}$ (“local $\mathbb{P}_{\Delta^\circ}$ ”) can be identified with a family of $\text{CH}^2(\cdot, 2)$ -elements on a family of anticanonical (elliptic) curves in \mathbb{P}_Δ [Doran and Kerr 2011, §5]. As a second application of the integral regulator, we show how to apply it to compute the correct “torsion term” in the higher normal function associated to one of these families. That is, if $\mathcal{E} \xrightarrow{\pi} \mathbb{P}^1_t$ is smooth away from $\Sigma = \{0\} \cup \Sigma^*$, with fibers $E_t = \pi^{-1}(t)$, and $\Xi \in \text{CH}^2(\mathcal{E} \setminus E_0, 2)$ has fiberwise restrictions $\xi_t \in \text{CH}^2(E_t, 2)$ ($t \notin \Sigma$), we shall compute

$$\begin{aligned} \mathcal{R}_t &:= \text{AJ}^{2,2}(\xi_t) \in H^2_{\mathcal{D}}(E_t, \mathbb{C}/\mathbb{Z}(2)) \cong H^1(E_t, \mathbb{C}/\mathbb{Z}(2)) \\ &\cong \text{Hom}(H_1(E_t, \mathbb{Z}), \mathbb{C}/(2\pi i)^2\mathbb{Z}) \end{aligned}$$

in a neighborhood of $t = 0$. Writing $\{\omega_t\}$ for a section of $\omega_{\mathcal{E}/\mathbb{P}^1}$ vanishing at ∞ , the constant term in (a branch of) the resulting truncated higher normal function

$$(6-1) \quad v(t) := \frac{1}{2\pi i} \langle \omega_t, \mathcal{R}_t \rangle$$

will play a role in forthcoming work of the first author with C. Doran on quantum curves.

To begin in a somewhat more general scenario, let $\Delta \subset \mathbb{R}^2$ be any convex polytope with integer vertices $\{p_i = (a_i, b_i)\}_{i=1}^N$ and interior integer points $\{(v_j, w_j)\}_{j=1}^g$.




Figure 1. For the proof of Lemma 6.1.

Define a multiparameter family

$$\rho : \mathcal{C} \rightarrow \mathbb{C}^g, \quad C_{\underline{\lambda}} := \rho^{-1}(\underline{\lambda})$$

of (where smooth) genus g curves by taking the Zariski closure of

$$\mathcal{C}^* := \{(x, y; \lambda_1, \dots, \lambda_g) \mid 0 = \Phi_{\underline{\lambda}}(x, y) := \phi(x, y) - \sum_{j=1}^g \lambda_j x^{v_j} y^{w_j}\} \subset (\mathbb{C}^*)^2 \times \mathbb{C}^g$$

in $\mathbb{P}^1 \times \mathbb{C}^g$, where $\phi(x, y) := \sum_{(a,b) \in \partial \Delta \cap \mathbb{Z}^2} m_{a,b} x^a y^b$ is uniquely determined by requiring its edge polynomials to be powers of $(t+1)$. (If the edges of Δ have no interior points, then $\phi(x, y) = \sum_{i=1}^N x^{a_i} y^{b_i}$.) The symbol $\{-x, -y\}$ represents a closed precycle $\Xi^* \in Z^2(\mathcal{C}^*, 2)$ parametrized (in $\mathcal{C}^* \times \square^2$) by $(x, y, \underline{\lambda}, -x, -y)_{(x,y,\underline{\lambda}) \in \mathcal{C}^*}$.

Lemma 6.1. *The class of Ξ^* in $\text{CH}^2(\mathcal{C}^*, 2)$ is the restriction of a class $\Xi \in \text{CH}^2(\mathcal{C}, 2)$.*

Proof. We need only check that the Tame symbol of $\{-x, -y\}|_{C_{\underline{\lambda}}^*} \in K_2^M(\mathbb{C}(C_{\underline{\lambda}}^*))$ is zero for general $\underline{\lambda}$. The symbol $\{-x, -y\}$ is invariant under unimodular change of toric coordinates,¹¹ so we may assume that (after shifting Δ by $(-a_m, -b_m)$ for some m) we have a picture (Figure 1) where the bottom edge corresponds to the toric divisor at whose intersection with $C_{\underline{\lambda}}$ we wish to compute $\text{Tame}(\{-x, -y\}|_{C_{\underline{\lambda}}^*}) \in \mathbb{C}^*$. Since the edge polynomial is $(1+x)^c$, this intersection occurs at $(-1, 0)$, so the Tame symbol is 1. □

Now set $\mathcal{R}_{\underline{\lambda}} := \text{AJ}^{2,2}(\Xi|_{C_{\underline{\lambda}}}) \in \text{Hom}(H_1(C_{\underline{\lambda}}), \mathbb{Z}, \mathbb{C}/\mathbb{Z}(2))$. Picking any vertex p_m of Δ , we can (via unimodular coordinate change) put it in the position indicated in Figure 1. In the new coordinates (still denoted (x, y)), $C_{\underline{\lambda}}$ is cut out by an equation of the form

$$\begin{aligned} 0 &= \tilde{\Phi}_{\underline{\lambda}} := x^{-a_m} y^{-b_m} \Phi_{\underline{\lambda}}(x, y) \\ &= (1+x)^{k_m} + y \left\{ \Psi_m(x, y) - \sum_{j=1}^g \lambda_j x^{v_j - a_m} y^{w_j - b_m - 1} \right\}, \end{aligned}$$

¹¹That is, replacing x, y by $x^a y^b, x^c y^d$ with $ad - bc = 1$; the a_i, b_i, v_i, w_i are changed accordingly.

and acquires a node at $(0, 0)$ as $\lambda_\ell \rightarrow \infty$. (Note that ℓ is determined by m .) The corresponding vanishing cycle α_m has image $|x| = |y| = \epsilon$ under

$$H_1(C_{\underline{\lambda}}, \mathbb{Z}) \xrightarrow{\text{Tube}} H_2(\mathbb{P}_\Delta \setminus C_{\underline{\lambda}})$$

for large $|\lambda_\ell|$.

Theorem 6.2. *For $i\lambda_\ell \in \mathfrak{H}$ and $|\lambda_\ell| \gg 0$, and $\lambda_{j \neq \ell}$ sufficiently small,¹² we have*

$$\mathcal{R}_{\underline{\lambda}}(\alpha_m) = 2\pi i \left(-\log(\lambda_\ell) + \sum_{k \geq 1} \frac{1}{k} [\Psi_{\underline{\lambda}, \ell}^k]_0 \right) \in \mathbb{C}/\mathbb{Z}(2),$$

where $\Psi_{\underline{\lambda}, \ell} := (-1/\lambda_\ell)(x^{-v_\ell} y^{-w_\ell} \Phi_{\underline{\lambda}} + \lambda_\ell)$ and $[\cdot]_0$ takes the constant term in a Laurent polynomial.

Proof. We use the notation

$$\begin{aligned} R\{f_1, f_2\} &= \log(f_1) \frac{df_1}{f_1} - 2\pi i \log(f_2) \delta_{T_{f_1}}, \quad \text{and} \\ R\{f_1, f_2, f_3\} &= \log(f_1) \frac{df_2}{f_2} \wedge \frac{df_3}{f_3} + 2\pi i \log(f_2) \frac{df_3}{f_3} \delta_{T_{f_1}} + (2\pi i)^2 \log(f_3) \delta_{T_{f_1} \cap T_{f_2}} \end{aligned}$$

for R_2 and R_3 with f_i replacing z_i . Writing D for the bottom-edge divisor in Figure 1, we have

$$\text{Tame}_D\{\tilde{\Phi}_{\underline{\lambda}}, -x, -y\} = \{\tilde{\Phi}_{\underline{\lambda}}(x, 0), -x\} = \{(1+x)^c, -x\} (= 1).$$

So writing $\Gamma = \{|x| = \epsilon \geq |y|\}$ ($\implies \alpha_m = \Gamma \cap C_{\underline{\lambda}}$) gives

$$\begin{aligned} \mathcal{R}_t(\alpha_m) &= \int_{\alpha_m} R\{-x, -y\} = \int_{\Gamma} R\{-x, -y\} \cdot \delta_{C_{\underline{\lambda}}} \\ &= \frac{-1}{2\pi i} \int_{\Gamma} d[R\{\tilde{\Phi}_{\underline{\lambda}}, -x, -y\}] - \int_{\Gamma} R\{(1+x)^c, -x\} \cdot \delta_D \\ &= \frac{-1}{2\pi i} \int_{\partial\Gamma} R\{\tilde{\Phi}_{\underline{\lambda}}, -x, -y\} - \int_{|x|=\epsilon} R\{(1+x)^c, -x\} \\ &= \frac{-1}{2\pi i} \int_{|x|=|y|=\epsilon} R\{x^{-v_\ell} y^{-w_\ell} \Phi_{\underline{\lambda}}, -x, -y\} \\ &= \frac{-1}{2\pi i} \int_{|x|=|y|=\epsilon} R\{\lambda(1 - \Psi_{\underline{\lambda}, \ell}), -x, -y\} \\ &= \frac{-1}{2\pi i} \int_{|x|=|y|=\epsilon} \{\log(\lambda) + \log(1 - \Psi_{\underline{\lambda}, \ell})\} \frac{dx}{x} \wedge \frac{dy}{y} \\ &= -2\pi i \log(\lambda) + 2\pi i \sum_{k \geq 1} \int_{|x|=|y|=\epsilon} \Psi_{\underline{\lambda}, \ell}^k \frac{dx}{x} \wedge \frac{dy}{y} \end{aligned}$$

¹²E.g., if $c := |\Delta \cap \mathbb{Z}^2|$, then $|\lambda_\ell| > c\epsilon^{-2}$ and $|\lambda_{j \neq \ell}| < 1/(c\epsilon^3)$ will do.

modulo $\mathbb{Z}(2)$. Here only the first term of R_3 enters since $T_{\lambda(1-\psi)} \cap |x| = |y| = \epsilon$ is empty under the given assumptions. \square

Returning to the more specific scenario at the beginning of this section, if $g = 1$ and $\lambda_1 =: \lambda =: \frac{1}{t}$, then $\Phi_\lambda = \phi(x, y) - \lambda$ and $\Psi_{\lambda,1} = t\phi(x, y)$, so that (writing \mathcal{R}_t instead of \mathcal{R}_λ), Theorem 6.2 yields:

Corollary 6.3. *If Δ is reflexive, then the α_m are all homologous ($=: \alpha$), and*

$$\mathcal{R}_t(\alpha) \equiv_{\mathbb{Z}(2)} 2\pi i \left(\log(t) + \sum_{k \geq 1} \frac{[\phi^k]_0}{k} \right)$$

for t small in the right-half-plane.

It remains to compute $\mathcal{R}_t(\beta)$ for a cycle β complementary to α (so that $\mathbb{Z}\langle \alpha, \beta \rangle = H_1(E_t, \mathbb{Z})$), which we shall do for the local \mathbb{P}^2 setting only: Δ the convex hull of $\{(1, 0), (0, 1), (-1, -1)\}$, and $\phi = x + y + x^{-1}y^{-1}$. (So $\mathbb{P}_{\Delta^\circ} \cong \mathbb{P}^2$, while the family of elliptic curves E_t lives in \mathbb{P}_Δ .) Taking $t > 0$ small, write $(0 <) x_0(t) < x_-(t) < x_+(t) < \infty$ for the branch points of

$$E_{t(=\lambda^{-1})} : y^2 + (x - \lambda)y + x^{-1} = 0$$

over \mathbb{P}_x^1 , and $y^\pm(x) = \frac{1}{2} \{ (\lambda - x) \pm \sqrt{(x - \lambda)^2 - 4x^{-1}} \}$. Then β (resp. α) is given by the difference of paths (on the two branches) between $x_0(t)$ and $x_-(t)$ (resp. $x_-(t)$ and $x_+(t)$).

Now $T_{-x} = \mathbb{R}_{>0} \subset \mathbb{P}_x^1$, so taking the y^+ -branch of β to run from x_0 to x_- in \mathfrak{H} , and the y^- -branch of β to run from x_- to x_0 in $-\mathfrak{H}$, we have $\beta \cap T_{-x} = (x_0, y_0) \cup (x_-, y_-)$; moreover, $\log(-x) = \log(x) \mp i\pi$ on the y^\pm -branch of β . The upshot is that

$$\begin{aligned} \int_\beta R\{-x, -y\}|_{E_t} &= \int_\beta \log(-x) \frac{dy}{y} - 2\pi i \sum_{\beta \cap T_{-x}} \log(y) \\ &= - \int_{x_0(t)}^{x_-(t)} \log(x) d\log\left(\frac{y^+(x)}{y^-(x)}\right) = \int_{x_0(t)}^{x_-(t)} \log\left(\frac{y^+(x)}{y^-(x)}\right) \frac{dx}{x} \\ &= \int_{x_0(t)}^{x_-(t)} \log\left(\frac{1 + \sqrt{1 - \xi}}{1 - \sqrt{1 - \xi}}\right) \frac{dx}{x}, \end{aligned}$$

where $\xi = \frac{4t^2}{x(1-xt)^2}$. Writing for $\xi \in (0, 1)$

$$\log\left(\frac{1 + \sqrt{1 - \xi}}{1 - \sqrt{1 - \xi}}\right) + \log\left(\frac{\xi}{4}\right) =: - \sum_{m \geq 1} \alpha_m \xi^m,$$

the above integral decomposes into

$$-2 \log(t) \int_{x_0}^{x_-} \frac{dx}{x} + \int_{x_0}^{x_-} \log(x) \frac{dx}{x} + 2 \int_{x_0}^{x_-} \log(1 - xt) \frac{dx}{x} - \sum_{m \geq 1} \alpha_m \int_{x_0}^{x_-} \xi^m \frac{dx}{x}.$$

Using the approximations $x_0 \simeq 4t^2(1+8t^3)$ and $x_- \simeq t^{-1}(1-2t^{3/2}-2t^3)$, a lengthy direct computation gives that

$$\mathcal{R}_t(\beta) = \frac{9}{2} \log^2(t) - \frac{\pi^2}{2} + \mathcal{O}(t \log(t)).$$

Let $\delta_t := t \frac{d}{dt}$. By the general result [Doran and Kerr 2011, Corollary 4.1], one knows that $\nabla_{\delta_t} \mathcal{R}_t = [\omega_t]$, where

$$\omega_t := \text{Res}_{E_t} \left(\frac{\frac{dx}{x} \wedge \frac{dy}{y}}{1-t\phi(x,y)} \right)$$

has its periods $\omega_t(\gamma) := \int_\gamma \omega_t$ annihilated by the Picard–Fuchs operator

$$\mathcal{L} = \delta_t^2 - 27t^3(\delta_t + 1)(\delta_t + 2).$$

The regulator periods $\mathcal{R}_t(\gamma)$ are therefore killed by $\mathcal{L} \circ \delta_t$. Since $\mathcal{L}(\cdot) = 0$ is known to have the basis of solutions

$$\begin{aligned} \pi_1 &= \sum_{n \geq 0} a_n t^{3n}, \\ \pi_2 &= 3 \log(t) \pi_1 + \sum_{n \geq 1} a_n b_n t^{3n}, \end{aligned}$$

with

$$a_n = \frac{(3n)!}{(n!)^3} \quad \text{and} \quad b_n = \sum_{k=0}^{n-1} \left(\frac{3}{3k+1} + \frac{3}{3k+2} - \frac{2}{k+1} \right),$$

it now follows that (writing $B_n = b_n - 1/n$)

$$\begin{aligned} \mathcal{R}_t(\alpha) &\stackrel{\mathbb{Z}(2)}{\equiv} 2\pi i \left(\log(t) + \sum_{n \geq 1} \frac{a_n}{3n} t^{3n} \right), \\ \mathcal{R}_t(\beta) &\stackrel{\mathbb{Z}(2)}{\equiv} \frac{9}{2} \log^2(t) + 3 \log(t) \sum_{n \geq 1} \frac{a_n}{n} t^{3n} + \sum_{n \geq 1} \frac{a_n B_n}{n} t^{3n} - \frac{\pi^2}{2}, \\ \omega_t(\alpha) &= 2\pi i \sum_{n \geq 0} a_n t^{3n}, \\ \omega_t(\beta) &= 9 \log(t) \sum_{n \geq 0} a_n t^{3n} + 3 \sum_{n \geq 1} a_n b_n t^{3n}, \end{aligned}$$

for $0 < |t| < \frac{1}{3}$. For (6-1), this yields

Proposition 6.4. *The truncated normal function mirror to local \mathbb{P}^2 is*

$$\begin{aligned} v(t) &= \left\langle \frac{\omega_t}{2\pi i}, \mathcal{R}_t \right\rangle = \frac{1}{2\pi i} (\mathcal{R}_t(\alpha) \omega_t(\beta) - \mathcal{R}_t(\beta) \omega_t(\alpha)) \\ &\equiv \frac{9}{2} \log^2(t)(1+6t^3) + 3 \log(t)(9t^3) + \frac{\pi^2}{2} + (3\pi^2 - 9)t^3 + \mathcal{O}(t^6 \log^2 t), \end{aligned}$$

modulo the period lattice $\mathbb{Z}(2)\langle\omega_t(\alpha), \omega_t(\beta)\rangle$.

Remark 6.5. This is closely related to computations in [Hosono 2006; Mohri et al. 2001]; the main difference — and the salient result here — is the identification of $\pi^2/2$ as the correct torsion offset for our motivically defined ν .

Acknowledgments

We thank the National Science Foundation for support under the aegis of FRG Grant DMS-1361147, as well as C. Doran, J. Lewis, and C. Weibel for helpful correspondence, and the referee for a careful reading and constructive comments.

References

- [del Ángel R. et al. 2019] P. L. del Ángel R., C. Doran, M. Kerr, J. Lewis, J. Iyer, S. Müller-Stach, and D. Patel, “Specialization of cycles and the K -theory elevator”, *Commun. Number Theory Phys.* **13**:2 (2019), 299–349. MR Zbl
- [Beilinson 1984] A. A. Beilinson, “Higher regulators and values of L -functions”, pp. 181–238 in *Current problems in mathematics*, vol. 24, edited by R. V. Gamkrelidze, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1984. In Russian; translated in *J. Soviet Math.* **20**:2 (1985), 2036–2070. MR Zbl
- [Bloch 1986a] S. Bloch, “Algebraic cycles and higher K -theory”, *Adv. in Math.* **61**:3 (1986), 267–304. MR Zbl
- [Bloch 1986b] S. Bloch, “Algebraic cycles and the Beilinson conjectures”, pp. 65–79 in *The Lefschetz centennial conference* (Mexico City, 1984), vol. I, edited by D. Sundararaman, Contemp. Math. **58**, Amer. Math. Soc., Providence, RI, 1986. MR Zbl
- [Bloch 1994] S. Bloch, “The moving lemma for higher Chow groups”, *J. Algebraic Geom.* **3**:3 (1994), 537–568. MR Zbl
- [Bloch 2004] S. Bloch, “Some notes on elementary properties of higher Chow groups, including functoriality properties and cubical Chow groups”, preprint, 2004, http://www.math.uchicago.edu/~bloch/cubical_chow.pdf.
- [Bloch et al. 2015] S. Bloch, M. Kerr, and P. Vanhove, “A Feynman integral via higher normal functions”, *Compos. Math.* **151**:12 (2015), 2329–2375. MR Zbl
- [Bloch et al. 2017] S. Bloch, M. Kerr, and P. Vanhove, “Local mirror symmetry and the sunset Feynman integral”, *Adv. Theor. Math. Phys.* **21**:6 (2017), 1373–1454. MR Zbl
- [Borel 1974] A. Borel, “Stable real cohomology of arithmetic groups”, *Ann. Sci. École Norm. Sup.* (4) **7** (1974), 235–272. MR Zbl
- [Borel 1977] A. Borel, “Cohomologie de SL_n et valeurs de fonctions zeta aux points entiers”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **4**:4 (1977), 613–636. MR Zbl
- [Burgos Gil 2002] J. I. Burgos Gil, *The regulators of Beilinson and Borel*, CRM Monograph Series **15**, Amer. Math. Soc., Providence, RI, 2002. MR Zbl
- [Doran and Kerr 2011] C. F. Doran and M. Kerr, “Algebraic K -theory of toric hypersurfaces”, *Commun. Number Theory Phys.* **5**:2 (2011), 397–600. MR Zbl
- [Doran and Kerr 2014] C. F. Doran and M. Kerr, “Algebraic cycles and local quantum cohomology”, *Commun. Number Theory Phys.* **8**:4 (2014), 703–727. MR Zbl
- [Griffiths 1969] P. A. Griffiths, “On the periods of certain rational integrals, II”, *Ann. of Math.* (2) **90** (1969), 496–541. MR Zbl

- [Haesemeyer and Weibel 2019] C. Haesemeyer and C. A. Weibel, *The norm residue theorem in motivic cohomology*, Annals of Mathematics Studies **200**, Princeton University Press, 2019. MR Zbl
- [Hosono 2006] S. Hosono, “Central charges, symplectic forms, and hypergeometric series in local mirror symmetry”, pp. 405–439 in *Mirror symmetry*, vol. V, edited by N. Yui et al., AMS/IP Stud. Adv. Math. **38**, Amer. Math. Soc., Providence, RI, 2006. MR Zbl
- [Kerr 2003a] M. Kerr, “A regulator formula for Milnor K -groups”, *K-Theory* **29**:3 (2003), 175–210. MR Zbl
- [Kerr 2003b] M. D. Kerr, *Geometric construction of regulator currents with applications to algebraic cycles*, Ph.D. thesis, Princeton University, 2003, <https://search.proquest.com/docview/305317544>. MR
- [Kerr 2018] M. Kerr, “Counting, sums, and series”, lecture notes, Alberta Summer Math. Inst., 2018, <http://www.math.wustl.edu/~matkerr/ASMI.pdf>.
- [Kerr and Lewis 2007] M. Kerr and J. D. Lewis, “The Abel–Jacobi map for higher Chow groups, II”, *Invent. Math.* **170**:2 (2007), 355–420. MR Zbl
- [Kerr and Yang 2018] M. Kerr and Y. Yang, “An explicit basis for the rational higher Chow groups of abelian number fields”, *Ann. K-Theory* **3**:2 (2018), 173–191. MR Zbl
- [Kerr et al. 2006] M. Kerr, J. D. Lewis, and S. Müller-Stach, “The Abel–Jacobi map for higher Chow groups”, *Compos. Math.* **142**:2 (2006), 374–396. MR Zbl
- [Levine 1994] M. Levine, “Bloch’s higher Chow groups revisited”, pp. 235–320 in *K-theory* (Strasbourg, 1992), Astérisque **226**, Société Mathématique de France, Paris, 1994. MR Zbl
- [Li 2018] M. Li, “Integral regulators for higher Chow complexes”, *SIGMA Symmetry Integrability Geom. Methods Appl.* **14** (2018), art. no. 118, 12 pp. MR Zbl
- [Mohri et al. 2001] K. Mohri, Y. Onjo, and S.-K. Yang, “Closed sub-monodromy problems, local mirror symmetry and branes on orbifolds”, *Rev. Math. Phys.* **13**:6 (2001), 675–715. MR Zbl
- [Petras 2009] O. Petras, “Functional equations of the dilogarithm in motivic cohomology”, *J. Number Theory* **129**:10 (2009), 2346–2368. MR Zbl
- [Voevodsky 2002] V. Voevodsky, “Motivic cohomology groups are isomorphic to higher Chow groups in any characteristic”, *Int. Math. Res. Not.* **2002**:7 (2002), 351–355. MR Zbl
- [Weibel 2013] C. A. Weibel, *The K-book: an introduction to algebraic K-theory*, Graduate Studies in Mathematics **145**, Amer. Math. Soc., Providence, RI, 2013. MR Zbl

Received March 1, 2019. Revised July 4, 2019.

MATT KERR
 DEPARTMENT OF MATHEMATICS
 WASHINGTON UNIVERSITY IN ST. LOUIS
 ST. LOUIS, MO
 UNITED STATES
 matkerr@math.wustl.edu

MUXI LI
 SCHOOL OF MATHEMATICAL SCIENCES
 UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA
 HEFEI
 CHINA
 limuxi@ustc.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 306 No. 2 June 2020

Cohomological kernels of purely inseparable field extensions	385
ROBERTO ARAVIRE, BILL JACOB and MANUEL O'RYAN	
Kuperberg and Turaev–Viro invariants in unimodular categories	421
FRANCESCO COSTANTINO, NATHAN GEER, BERTRAND PATUREAU-MIRAND and VLADIMIR TURAEV	
A new equivalence between super Harish-Chandra pairs and Lie supergroups	451
FABIO GAVARINI	
Generalized Mullineux involution and perverse equivalences	487
THOMAS GERBER, NICOLAS JACON and EMILY NORTON	
Isotypic multiharmonic polynomials and Gelbart–Helgason reciprocity	519
ANTHONY C. KABLE	
Two applications of the integral regulator	539
MATT KERR and MUXI LI	
Definability and approximations in triangulated categories	557
ROSANNA LAKING and JORGE VITÓRIA	
Remarks on the theta correspondence over finite fields	587
DONGWEN LIU and ZHICHENG WANG	
On the configurations of centers of planar Hamiltonian Kolmogorov cubic polynomial differential systems	611
JAUME LLIBRE and DONGMEI XIAO	
2-categories of symmetric bimodules and their 2-representations	645
VOLODYMYR MAZORCHUK, VANESSA MIEMIETZ and XIAOTING ZHANG	
The homotopy groups of the η -periodic motivic sphere spectrum	679
KYLE ORMSBY and OLIVER RÖNDIGS	
On the Noether Problem for torsion subgroups of tori	699
FEDERICO SCAVIA	
Explicit polynomial bounds on prime ideals in polynomial rings over fields	721
WILLIAM SIMMONS and HENRY TOWNSNER	
A new local gradient estimate for a nonlinear equation under integral curvature condition on manifolds	755
LIANG ZHAO and SHOUWEN FANG	



0030-8730(202006)306:2;1-J