

*Pacific  
Journal of  
Mathematics*

**REMARKS ON THE THETA CORRESPONDENCE  
OVER FINITE FIELDS**

DONGWEN LIU AND ZHICHENG WANG

Volume 306 No. 2

June 2020



## REMARKS ON THE THETA CORRESPONDENCE OVER FINITE FIELDS

DONGWEN LIU AND ZHICHENG WANG

**S.-Y. Pan decomposed the uniform projection of the Weil representation of a finite symplectic-odd orthogonal dual pair, in terms of Deligne–Lusztig virtual characters, assuming that the order of the finite field is large enough. We use Pan’s decomposition to study the theta correspondence for dual pairs of this kind, following the approach of Adams and Moy and Aubert, Michel and Rouquier. Our results give the theta correspondence between unipotent representations and certain quadratic unipotent representations.**

### 1. Introduction

The theory of Weil representations and theta correspondence over finite fields has been studied extensively (see, e.g., [Adams and Moy 1993; Aubert et al. 1996; 2016; Epequin Chavez 2019; Gérardin 1977; Gurevich and Howe 2017; Howe 1973; Pan 2016; 2019a; 2019b; Srinivasan 1968; 1979b]) and has many applications to the representation theory of  $p$ -adic groups, but there are still a few important open problems unsolved in the finite field case. In particular, there is a complete understanding for the theta correspondence of unipotent representations for finite reductive dual pairs, with only the exception of symplectic-odd orthogonal dual pairs. The aim of this paper is to fill this gap and give some answers for this missing case. In contrast to other dual pairs, it is well known that the theta lifting of unipotent representations may not be unipotent in general for symplectic-odd orthogonal dual pairs. It turns out that new phenomena occur and the correspondence involves certain quadratic unipotent representations, which are called  $\theta$ -representations in this paper.

To give some details, let us start from the Deligne–Lusztig characters [1976], which are a major tool for the study of representations of finite Lie groups. Let  $F$  be a finite field of  $q$  elements,  $G$  be a connected reductive group defined over  $F$ , and  $F$  be the corresponding Frobenius endomorphism. Let  $G = G^F$  be the group of rational points of  $G$ . For an  $F$ -stable maximal torus  $T$  of  $G$  and a character  $\theta$

---

*MSC2010:* primary 20C33; secondary 22E50.

*Keywords:* theta correspondence, finite classical groups, Deligne–Lusztig characters, unipotent cuspidal representations.

of  $T = T^F$ , let  $R_{T,\theta}^G$  denote the virtual character of  $G$  defined by P. Deligne and G. Lusztig in the seminal work [Deligne and Lusztig 1976].

Let  $T_0$  be a maximally split  $F$ -stable maximal torus of  $G$ , and  $W_G = N(T_0)/T_0$  be the Weyl group of  $G$ . For each  $w \in W_G$ , we can associate an  $F$ -stable maximal torus  $T_w$  of  $G$  and a specific character of  $T_w = T_w^F$  of order 2, denoted by  $\theta_w$  [Pan 2016]. If  $G = \mathbf{SO}_{2n+1}$  where  $n$  is a positive integer, then

$$\frac{1}{|W_{\mathbf{SO}_{2n+1}}|} \sum_{w \in W_{\mathbf{SO}_{2n+1}}} R_{T_w, \theta_w}^{\mathbf{SO}_{2n+1}}$$

is the unique linear character of  $\mathbf{SO}_{2n+1}(q)$  of order 2, which will be denoted by  $\chi_{\mathbf{SO}_{2n+1}}$ . For  $n = 0$ , by convention  $\chi_{\mathbf{SO}_{2n+1}}$  is the trivial character. An irreducible representation  $\pi$  of  $G$  is called a  $\theta$ -representation if it occurs in some  $R_{T_w, \theta_w}^G$ . The  $\theta$ -representations can be interpreted as certain quadratic unipotent representations of finite reductive Lie groups (see, e.g., [Mœglin 1995] for the notion of quadratic unipotent representations for  $p$ -adic groups). For  $\mathbf{SO}_{2n+1}$ ,  $\pi \rightarrow \chi_{\mathbf{SO}_{2n+1}} \pi$  gives a bijection between unipotent representations and  $\theta$ -representations.

Let  $\omega_{\mathbf{Sp}_{2n}}$  be the Weil representation of the finite symplectic group  $\mathbf{Sp}_{2n}(q)$  with  $q$  odd, which depends on a choice of a nontrivial additive character  $\psi$  of  $F$  (see [Gérardin 1977]). We will denote by  $\omega_{\mathbf{Sp}_{2n}}^\#$  the uniform projection of  $\omega_{\mathbf{Sp}_{2n}}$ , i.e., the projection onto the subspace of virtual characters spanned by all the Deligne–Lusztig characters. A pair  $(G, G')$  of mutually centralized reductive subgroups of  $\mathbf{Sp}_{2n}$  is called a reductive dual pair. Then the Weil representation  $\omega_{\mathbf{Sp}_{2n}}$  restricts to a representation  $\omega_{G, G'}$  of  $G \times G'$ , whose character will be denoted by the same notation.

J. Adams and A. Moy [1993] show that for unitary or symplectic-even orthogonal dual pairs, if  $\pi \otimes \pi'$  occurs in  $\omega_{G, G'}$ , then  $\pi$  is unipotent if and only if  $\pi'$  is unipotent, in which case they are in correspondence under the decomposition of the uniform projection  $\omega_{G, G'}^\#$ . They also study the first occurrence of cuspidal representations and give the explicit correspondence of cuspidal unipotent representations.

As we mentioned above, it is known that for symplectic-odd orthogonal dual pairs, theta correspondence no longer preserves unipotent representations in general. On the other hand, Pan [2016] gives the decomposition of  $\omega_{G, G'}^\#$  for such dual pairs. It turns out that based on Pan’s work we can obtain the following analog of the result in [Adams and Moy 1993], which fills the gap in the symplectic-odd orthogonal case. See Theorem 3.7 for details, and also Theorem 3.3 for a general statement.

**Theorem 1.1.** *Assume that  $q$  is odd and large enough such that the main result in [Srinivasan 1979b] holds. Let  $\lambda_k$  be the unique cuspidal unipotent representation of  $\mathbf{Sp}_{2n}$ ,  $n = k(k + 1)$ . Then its theta lifting to  $O_{2n+1}^\epsilon$ ,  $\epsilon = \pm$ , is the unique cuspidal  $\theta$ -representation of  $O_{2n+1}^\epsilon$  with trivial central character, and is the first occurrence in each Witt tower of odd orthogonal groups.*

The proof of this theorem invokes some ingredients, among which we would like to mention the so-called conservation relation (see, e.g., [Pan 2002; Sun and Zhu 2015]). Such a relation provides a link between the first occurrence of the theta lifting of a representation towards two different Witt towers. For finite dual pairs it is known only for cuspidal representations, which is sufficient for our purpose.

Applying Lusztig's bijection [1988] we are able to classify the cuspidal  $\theta$ -representations of symplectic groups (Theorem 3.11). Consequently we can switch the roles of symplectic and odd orthogonal groups in Theorem 1.1 and obtain an analogous result Theorem 3.12, the proof of which uses the conservation relations in an inductive manner.

The cuspidal representations are the building blocks of the so-called Harish-Chandra series. As a second step, A.-M. Aubert, J. Michel and R. Rouquier [Aubert et al. 1996, Théorème 3.7] study the theta correspondence of Harish-Chandra series of unipotent cuspidal representations for unitary and symplectic-even orthogonal dual pairs. According to Lusztig's bijection and [Pan 2002, Theorem 3.10], using their results we should be able to handle the theta correspondence of any irreducible representations. Based on Theorem 1.1, we are able to extend their results to the symplectic-odd orthogonal case using similar calculations to those in [Aubert et al. 1996].

Our second main result Theorem 4.2 establishes the theta correspondence between the Harish-Chandra series of cuspidal (quadratic) unipotent representations of symplectic-odd orthogonal dual pairs. We briefly describe its content as follows. Denote  $T_l$  the split torus over  $F$  of rank  $l$ , and let  $\theta_l$ ,  $\theta_{k,l}$  and  $\theta'_{k,l}$  be the quadratic characters of  $T_l$  defined in Section 4. Let  $(G_n, G'_{n'})$  be a dual pair of symplectic and odd orthogonal groups over  $F$ , and  $\lambda$  be an irreducible cuspidal representation of  $G_n$  such that its theta lifting  $\lambda'$  to  $G'_{n'}$  is irreducible cuspidal as well. Let  $\gamma$  be an irreducible constituent of the parabolic induction  $R_{G_n \times T_{m-n}}^{G_m}(\lambda \times \theta)$ , where  $\theta$  is some quadratic character specified below, and  $\Theta_{G'_{m'}}(\gamma)$  be the theta lifting of  $\gamma$  to  $G'_{m'}$ . Then Theorem 4.2 states that  $\Theta_{G'_{m'}}(\gamma) = 0$  if  $m' < n'$ , and is spanned by irreducible constituents of  $R_{G'_{n'} \times T_{m'-n'}}^{G'_{m'}}(\lambda' \otimes \theta')$  otherwise, where the pair  $(\theta, \theta')$  of quadratic characters is as follows:

- If  $G_n$  is symplectic and  $G'_{n'}$  is odd orthogonal, then  $(\theta, \theta')$  is equal to  $(\theta_{m-n}, \mathbf{1})$  or  $(\mathbf{1}, \theta'_{m-n, m'-n'})$ .
- If  $G_n$  is odd orthogonal and  $G'_{n'}$  is symplectic, then  $(\theta, \theta')$  is equal to  $(\mathbf{1}, \theta_{m'-n'})$  or  $(\theta_{m-n}, \theta_{m-n, m'-n'})$ .

We expect that our results may have some applications towards the theta lifting of quadratic unipotent representations of  $p$ -adic groups.

From Mackey theory one knows that constituents of such Harish-Chandra series can be parametrized by irreducible representations of certain Weyl groups. Thus the

theta correspondence reduces to a correspondence of Weyl group representations. In [Aubert et al. 1996], explicit formulas have been proved for the unitary case and conjectured for the symplectic-even orthogonal case, and the latter has been proved recently in [Pan 2019a]. We expect that similar conjectures should also hold for the symplectic-odd orthogonal case. We also remark that from the explicit correspondence we can observe that the conservation relations in general do not hold for noncuspidal representations of finite dual pairs. See [Pan 2019b, Section 9] for more results on the (non-)conservation relations.

The paper is organized as follows. In Section 2 we briefly recall the theory of Deligne–Lusztig characters as well as Pan’s decomposition formula [2016]. In Section 3 we prove the theta correspondence between cuspidal unipotent representations and cuspidal  $\theta$ -representations, following [Adams and Moy 1993] and using the conservation relations. Based on this result, in Section 4 we establish the theta correspondence of certain Harish-Chandra series, using some calculations of Jacquet and induction functors as in [Aubert et al. 1996] and [Mœglin et al. 1987]. We explain briefly the correspondence of Weyl group representations mentioned above at the end of Section 4.

**Notation.** In this paper the cardinality  $q$  of  $F$  is assumed to be odd and large enough (as in Theorem 3.1). The finite Lie group  $G = G^F$  is also written as  $G(q)$ . Representations are realized on vector spaces over  $\overline{\mathcal{Q}}_\ell$  where  $\ell \neq p$  is a prime. We do not distinguish a (virtual) representation and its character. Thus  $R(G)$  stands for the set of virtual representations of  $G$  as well as the set of integral combinations of irreducible characters. For  $\pi \in R(G)$ , let  $\text{Irr}(\pi)$  be the set of constituents of  $\pi$ . For  $\rho \in \text{Irr}(G)$ , we simply write  $\rho \in \pi$  if  $\rho \in \text{Irr}(\pi)$ . For  $\pi_1, \pi_2 \in R(G)$ , let

$$(\pi_1, \pi_2)_G = \frac{1}{|G|} \sum_{g \in G} \pi_1(g) \overline{\pi_2(g)},$$

where the conjugate is the restriction of the usual complex conjugate to  $\overline{\mathcal{Q}} \subset \overline{\mathcal{Q}}_\ell$ .

## 2. Deligne–Lusztig theory

Let  $F$  be a finite field of  $q$  elements,  $G$  be a connect reductive group defined over  $F$  and  $F$  be the corresponding Frobenius endomorphism. The group of rational points of  $G$  is denoted by  $G = G^F$ .

**2A. Conjugacy classes of maximal tori.** For an  $F$ -stable maximal torus  $T$  of  $G$ , let  $W_G(T) = N_G(T)/T$  be its Weyl group, where  $N_G(T)$  is the normalizer of  $T$  in  $G$ , and let  $T = T^F$  be the group of rational points of  $T$ .

Fix a maximally split  $F$ -stable maximal torus  $T_0$  of  $G$ , and denote its Weyl group by  $W_G$ . A conjugate  ${}^g T_0 = gT_0g^{-1}$  of  $T_0$  by an element  $g \in G$  is  $F$ -stable if and only if  $g^{-1}F(g) \in N_G(T_0)$ . By the Lang–Steinberg theorem, for each  $w \in W_G$  we may

choose an element  $g \in G$  such that  $g^{-1}F(g)$  is in  $N_G(T_0)$  whose image in  $W_G$  is equal to  $w$ . The  $F$ -stable maximal torus  ${}^sT_0$  will be denoted by  $T_w$ . Two elements  $w, w' \in W_G$  are called  $F$ -conjugate if there exists  $x \in W_G$  such that  $w' = x^{-1}wF(x)$ . The map  ${}^sT_0 \rightarrow g^{-1}F(g)$  gives a bijection between the  $G$ -conjugacy classes of  $F$ -stable maximal tori of  $G$  and the  $F$ -conjugacy classes of  $W_G$ .

**2B. Deligne–Lusztig characters.** In their celebrated paper, Deligne and Lusztig [1976] defined a virtual representation  $R_{T,\theta}^G$  of  $G$ , associated to a character  $\theta$  of  $T$ . The construction of Deligne–Lusztig characters makes use of the deep theory of  $\ell$ -adic cohomology. Here we only review some standard facts on these characters which will be used in this paper (see [Carter 1985, Chapter 7]).

If  $y = su$  is the Jordan decomposition of an element  $y \in G$ , then

$$(2-1) \quad R_{T,\theta}^G(y) = \frac{1}{|C^0(s)|} \sum_{g \in G, s^g \in T} \theta(s^g) Q_{sT}^{C^0(s)}(u)$$

where  $C^0(s) = C_G^0(s)$  is the connected component of the centralizer of  $s$  in  $G$ , and  $Q_{sT}^{C^0(s)} = R_{sT,1}^{C^0(s)}(u)$  is the Green function of  $C^0(s)$  associated to  ${}^sT$ . Note that  $s^g = g^{-1}sg \in T$  if and only if  ${}^sT = gTg^{-1} \subset C^0(s)$ .

It is known that

$$(2-2) \quad 1 = \frac{1}{|W_G|} \sum_{w \in W_G} R_{T_w,1}^G = \sum_{(T) \subset G} \frac{1}{|W_G(T)|} R_{T,1}^G$$

where  $(T) \subset G$  means that the summation is taken over the  $G$ -conjugacy classes of  $F$ -stable maximal tori in  $G$ .

Two virtual representations  $R_{T,\theta}^G$  and  $R_{T',\theta'}^G$  are disjoint if  $(T, \theta)$  and  $(T', \theta')$  are not geometrically conjugate. The elements in the subspace of class functions on  $G$  spanned by all the  $R_{T,\theta}^G$ 's are called uniform functions of  $G$ . For a class function  $f$  on  $G$ , denote by  $f^\#$  its orthogonal projection to the subspace of uniform functions.

**2C. Weyl group.** By [Srinivasan 1979b, Lemma 3.1], there is a natural bijection between conjugacy classes of maximal tori in  $\text{Sp}_{2n}(q)$  and the disjoint union of the  $O_{2n}^\epsilon(q)$ -conjugacy classes of maximal tori in  $\text{SO}_{2n}^\epsilon(q)$  for  $\epsilon = \pm$ . For  $w \in W_{\text{Sp}_{2n}}$ , we define  $\epsilon_w = \epsilon$  if the  $F$ -stable maximal torus  $T_w$  of  $\text{Sp}_{2n}(q)$  corresponds to an  $F$ -stable maximal torus in  $\text{SO}_{2n}^\epsilon(q)$ , and define

$$W_{\text{Sp}_{2n}}^\epsilon = \{w \in W_{\text{Sp}_{2n}} \mid \epsilon_w = \epsilon\}.$$

We shall identify the Weyl groups of  $\text{Sp}_{2n}$  and  $\text{SO}_{2n+1}$ , which will be denoted by  $W_n$ . Thus we also write  $W_n^\epsilon = W_{\text{Sp}_{2n}}^\epsilon$ . The Weyl group  $W_{\text{SO}_{2n}^+}$  of  $\text{SO}_{2n}^+$  can be regarded as a subgroup of  $W_n$  of index 2.

Clearly,  $\epsilon_w = +$  if  $T_w \simeq \text{GL}_1(q)$  in  $\text{Sp}_2(q)$ , and  $\epsilon_w = -$  if  $T_w \simeq U_1(q)$  in  $\text{Sp}_2(q)$ . Since  $\text{Sp}_{2n}(q')$  can be regarded as a subgroup of  $\text{Sp}_{2nt}(q)$  for  $t \in \mathbb{N}$ , an  $F$ -stable

maximal torus of  $\mathrm{Sp}_{2n}(q^t)$  is also an  $F$ -stable maximal torus of  $\mathrm{Sp}_{2nt}(q)$ . Hence  $W_{\mathrm{Sp}_{2n}(q^t)}$  can be embedded as a subset of  $W_{\mathrm{Sp}_{2nt}(q)}$ . For  $w \in W_{\mathrm{Sp}_{2n}(q^t)}$ , the value of  $\epsilon_w$  is the same no matter whether  $w$  is regarded as an element of  $W_{\mathrm{Sp}_{2n}(q^t)}$  or  $W_{\mathrm{Sp}_{2nt}(q)}$ , and therefore  $\epsilon_w = +$  (resp.  $\epsilon_w = -$ ) if  $T \cong \mathrm{GL}_1(q^t)$  (resp.  $T_w \cong U_1(q^t)$ ) in  $\mathrm{Sp}_{2t}(q)$ .

**2D. The character  $\theta_w$ .** If  $T \cong \mathrm{GL}_1(q^t)$  or  $U_1(q^t)$ , let  $\theta_T$  denote the unique character of  $T$  of order 2, i.e.,  $\theta_T(a) = \nu(a)^{(q^t-1)/2}$  (resp.  $\theta_T(a) = \nu(a)^{(q^t+1)/2}$ ) for  $a \in T$  if  $T \cong \mathrm{GL}_1(q^t)$  (resp.  $T \cong U_1(q^t)$ ), where  $\nu$  is an isomorphism of  $T$  onto a subgroup of  $\overline{\mathcal{Q}}_\ell^\times$ . Define  $\theta_T = \theta_{T_1} \otimes \cdots \otimes \theta_{T_r}$  if  $T \cong T_1 \times \cdots \times T_r$  is an  $F$ -stable maximal torus of a connected reductive group  $G$ , where  $T_i \cong \mathrm{GL}_1(q^{t_i})$  or  $U_1(q^{t_i})$  for some positive integer  $t_i$ ,  $i = 1, \dots, r$ . It is clear that  ${}^g\theta_T = \theta_{gT}$  for  $g \in G$ . If  $T = T_w$  for some  $w \in W_G$ , the character  $\theta_T$  will be denoted by  $\theta_w$ .

We can check that  $\theta_T = \chi_G|_T$  if  $G$  is equal to  $\mathrm{GL}_n$ ,  $U_n$  or  $\mathrm{SO}_n$  ( $n$  a positive integer), where  $\chi_G$  is the unique linear character of  $G$  of order 2. It is known as the spinor norm character in the case of orthogonal groups. For a representation or a character  $\pi$  of  $G$ , we abbreviate  $\chi_G \otimes \pi$  by  $\chi_G\pi$ .

The following identity can be found in [Pan 2016].

**Lemma 2.1.** *If  $G = \mathrm{GL}_n, U_n$  or  $\mathrm{SO}_n$ , then*

$$\chi_G = \frac{1}{|W_G|} \sum_{w \in W_G} R_{T_w, \theta_w}^G.$$

**Lemma 2.2.** *If  $u \in G$  is a unipotent element, then  $\chi_G(u) = 1$ .*

*Proof.* By (2-1), (2-2) and Lemma 2.1,

$$\begin{aligned} \chi_G(u) &= \frac{1}{|W_G|} \sum_{w \in W_G} R_{T_w, \theta_w}^G(u) \\ &= \frac{1}{|W_G|} \sum_{w \in W_G} \frac{1}{|G|} \sum_{g \in G} \theta_w(1) Q_{gT}^G(u) \\ &= \frac{1}{|W_G|} \sum_{w \in W_G} \frac{1}{|G|} \sum_{g \in G} Q_{gT}^G(u) = 1(u) = 1. \quad \square \end{aligned}$$

Recall that an irreducible representation  $\pi$  is called a unipotent representation if it occurs in some  $R_{T,1}^G$ . We have the following:

**Lemma 2.3.** *If  $G = \mathrm{GL}_n, U_n$  or  $\mathrm{SO}_n$ , then*

- (i)  $\pi$  is a unipotent representation if and only if  $\chi_G\pi$  occurs in some  $R_{T_w, \theta_w}^G$ ,
- (ii)  $R_{T_w, \theta_w}^G$  only consists of  $\chi_G\pi$  with  $\pi$  unipotent.

*Proof.* If  $y = su$  is the Jordan decomposition of an element  $y \in G$ , then

$$\begin{aligned} \chi_G R_{T_w,1}^G(y) &= \chi_G(y) \frac{1}{|C^0(s)|} \sum_{g \in G, s^G \in T_w} 1(s^g) \mathcal{Q}_{s^G T_w}^{C^0(s)}(u) \\ &= \frac{1}{|C^0(s)|} \sum_{g \in G, s^G \in T_w} \chi_G(s^g) \chi_G(u) \mathcal{Q}_{s^G T_w}^{C^0(s)}(u) \\ &= \frac{1}{|C^0(s)|} \sum_{g \in G, s^G \in T_w} \theta_w(s^g) \mathcal{Q}_{s^G T_w}^{C^0(s)}(u) = R_{T_w, \theta_w}^G(y). \quad \square \end{aligned}$$

Similar to the notion of unipotent representations, a representation  $\pi$  is called a  $\theta$ -representation if it occurs in some  $R_{T_w, \theta_w}^G$ . For  $G = \mathrm{Sp}_{2n}$ , we do not have nontrivial characters of  $G$ , but we can still talk about  $\theta$ -representations.

### 3. Weil representations and cuspidal unipotent representations

Adams and Moy [1993] (see also [Aubert et al. 1996; 2016; Epequin Chavez 2019]) consider the question of how the unipotent representations behave under theta correspondence. In particular they show that theta correspondence sends a cuspidal unipotent representation of  $G$  to a cuspidal unipotent representation of  $G'$  if  $(G, G') = (\mathrm{Sp}_{2n}, O_{2n}^\epsilon)$  or  $(U_n, U_{n'})$ . In the proof they use the decomposition of the Weil representation of  $(\mathrm{Sp}_{2n}, O_{2n}^\epsilon)$  and  $(U_n, U_{n'})$  given in [Srinivasan 1979b]. Pan [2016] gives a decomposition of the Weil representation of a finite symplectic-odd orthogonal dual pair, which we have recalled as Theorem 3.1. Thus we can give the correspondence of cuspidal unipotent representations of such dual pairs based on Pan's result, together with another ingredient called conservation relation.

**3A. Weil representation and theta lifting.** Let  $\omega_{\mathrm{Sp}_{2n}}$  be the character of the Weil representation (see [Gérardin 1977]) of the symplectic group  $\mathrm{Sp}_{2n}$  which depends on a nontrivial additive character  $\psi$  of  $F$ , and let  $\omega_{\mathrm{Sp}_{2n}}^\#$  denote the uniform projection of  $\omega_{\mathrm{Sp}_{2n}}$ , i.e., the projection onto the subspace of virtual characters spanned by all the Deligne–Lusztig characters. Let  $(G, G')$  be a reductive dual pair in  $\mathrm{Sp}_{2n}(q)$ . Write  $\omega_{G, G'}$  for the restriction of  $\omega_{\mathrm{Sp}_{2n}}$  to  $G \times G'$ . Then it decomposes into a direct sum

$$\omega_{G, G'} = \bigoplus_{\pi, \pi'} m_{\pi, \pi'} \pi \otimes \pi',$$

where  $\pi$  and  $\pi'$  run over irreducible representations of  $G$  and  $G'$ , respectively. We can rearrange this sum to get

$$\omega_{G, G'} = \bigoplus_{\pi} \pi \otimes \Theta_{G'}(\pi),$$

where  $\Theta_{G'}(\pi) = \bigoplus_{\pi'} m_{\pi, \pi'} \pi'$  is a (not necessarily irreducible) representation of  $G'$ , called the (big) theta lifting of  $\pi$  to  $G'$ . The theta lifting from  $G$  to  $G'$  will be written

as  $\pi \mapsto \Theta_{G'}(\pi)$ , and we have  $\pi' \in \Theta_{G'}(\pi)$  if and only if  $\pi \otimes \pi' \in \omega_{G,G'}$ . We should mention that it is an important open problem to find some “good” constituents from  $\Theta_{G'}(\pi)$  (see [Epequin Chavez 2019; Gurevich and Howe 2017] for different approaches and results in this direction). By abuse of notation we also write  $\omega_{G,G'}$  for its character, and write  $\omega_{G,G'}^\#$  for the uniform projection of  $\omega_{G,G'}$ .

It is often more convenient to work with the families of dual pairs associated to Witt towers instead of a single dual pair. Thus let us introduce systematically some notation for Witt towers. We will consider two Witt towers  $\mathcal{T}$  and  $\mathcal{T}'$  such that  $(G_n, G'_{n'})$  form a reductive dual pair of type I for any  $G_n \in \mathcal{T}$  and  $G'_{n'} \in \mathcal{T}'$ . There are the following types of Witt towers:

- For unitary groups there are two Witt towers  $U^+ = \{U_{2n}\}_{n \geq 0}$  and  $U^- = \{U_{2n+1}\}_{n \geq 0}$ .
- For symplectic groups there is only one Witt tower  $\text{Sp} = \{\text{Sp}_{2n}\}_{n \geq 0}$ .
- For even orthogonal groups there are two Witt towers  $O_{\text{even}}^+ = \{O_{2n}^+\}_{n \geq 0}$  and  $O_{\text{even}}^- = \{O_{2n}^-\}_{n \geq 1}$ .
- For odd orthogonal groups there are two Witt towers  $O_{\text{odd}}^\epsilon = \{O_{2n+1}^\epsilon\}_{n \geq 0}$ ,  $\epsilon = \pm$ .

For the dual pair  $(\text{Sp}_{2n}, O_{2n'}^\epsilon)$  where  $\epsilon = \pm$ , we write  $\omega_{n,n'}^\epsilon$  instead of  $\omega_{G,G'}$ . Since there are also two different nondegenerate symmetric bilinear forms on an odd-dimensional vector space over  $F$  (though their orthogonal groups are physically equal), we have two different types of symplectic-odd orthogonal dual pairs. Similarly we denote them by  $(\text{Sp}_{2n}, O_{2n'+1}^\epsilon)$  and the corresponding Weil representations by  $\omega_{n,n'}^\epsilon$ , where  $\epsilon = \pm$ . For unitary groups write  $G_n^\epsilon = U_{2n}$  for  $\epsilon = +$  or  $U_{2n+1}$  for  $\epsilon = -$ . Then we denote the Weil representation of  $(G_n^\epsilon, G_{n'}^{\epsilon'})$  by  $\omega_{n,n'}^{\epsilon,\epsilon'}$ . We will specify the dual pair we are working with when necessary, so it should cause no confusion when we use the notation  $\omega_{n,n'}^\epsilon$  or  $\omega_{n,n'}^{\epsilon,\epsilon'}$ , and occasionally we suppress the signs to ease notation.

Now we recall the main result of [Pan 2016], which will be a crucial tool used in this paper.

**Theorem 3.1.** *Suppose that  $q$  is sufficiently large such that the main theorem in [Srinivasan 1979b] holds. Then for a positive integer  $n$  and a nonnegative integer  $n'$ ,*

$$\omega_{\text{Sp}_{2n}, \text{SO}_{2n'+1}}^\# \cdot (1 \otimes \chi_{\text{SO}_{2n'+1}}) = \sum_{k=0}^{\min(n,n')} \frac{1}{|W_k|} \frac{1}{|W_{\text{Sp}_{2(n-k)}}|} \frac{1}{|W_{\text{SO}_{2(n'-k)+1}}|} \\ \sum_{v \in W_k} \sum_{\theta \in \text{Irr}(T_v)} \sum_{w \in W_{\text{Sp}_{2(n-k)}}} \sum_{w' \in W_{\text{SO}_{2(n'-k)+1}}} \epsilon_w R_{T_v \times T_w, \theta \otimes \theta_w}^{\text{Sp}_{2n}} \\ \otimes R_{T_v \times T_{w'}, \theta \otimes \theta_{w'}}^{\text{SO}_{2n'+1}}$$

where  $\text{Irr}(T_v)$  denotes the set of irreducible characters of the torus  $T_v$ .

We should point out that the above formula for the uniform projection is independent of the choice of  $\text{SO}_{2n'+1} = \text{SO}_{2n'+1}^\epsilon$ . In the rest of this paper the finite field  $F$  is assumed to be large enough as in Theorem 3.1.

**3B. Correspondence of unipotent and  $\theta$ -representations.** For finite unitary and symplectic-even orthogonal dual pairs, Adams and Moy showed that if  $\pi \otimes \pi'$  occurs in  $\omega_{n,n'}$ , then  $\pi$  is unipotent if and only if  $\pi'$  is unipotent. This does not hold for finite symplectic-odd orthogonal dual pairs, in which case one of the results we will prove says that  $\pi$  is unipotent if and only if  $\pi'$  is a  $\theta$ -representation when  $n = n'$ . Much as in [Adams and Moy 1993], our proof makes use of the decomposition of the Weil representation given by Theorem 3.1 in the symplectic-odd orthogonal case.

The following well-known result can be found in [Adams and Moy 1993; Mœglin et al. 1987], which says that the first occurrence of theta lifting of an irreducible cuspidal representation in a given Witt tower is also irreducible cuspidal, while the later occurrences are not cuspidal.

**Theorem 3.2.** *Let  $(G_n, G'_{n'})$  be a dual pair of finite classical groups. Assume that  $\pi$  is an irreducible cuspidal representation of  $G_n$ , and that*

- (i)  $\pi \mapsto \pi'$  in the theta correspondence for  $G_n \times G'_{n'}$ ,
- (ii)  $\pi$  does not occur in the theta correspondence for  $G_n \times G'_k$  for any  $k < n'$ .

*Then  $\pi'$  is irreducible cuspidal, and  $\Theta_{G'_k}(\pi)$  has no cuspidal constituents for  $k > n'$ .*

Recall that by Lemma 2.3, the map  $\pi \mapsto \chi_G \pi$  ( $G \neq \text{Sp}_{2n}$ ) gives a bijection between the unipotent representations and  $\theta$ -representations of  $G$ , and by Lemma 2.2 it preserves the set of cuspidal representations.

We have the following basic result.

**Theorem 3.3.** *Suppose that an irreducible representation  $\pi \otimes \pi'$  of  $(\text{Sp}_{2n}, \text{SO}_{2n'+1}^\epsilon)$  occurs in  $\omega_{n,n'}^\epsilon$ . Then the following hold.*

- (i) *If  $\pi$  is a unipotent representation of  $\text{Sp}_{2n}$ , then  $n' \geq n$  and  $\pi' \in R_{T_v \times T_{w'}, \theta_v \otimes 1}^{\text{SO}_{2n'+1}^\epsilon}$  for some  $v \in W_n$  and  $w' \in W_{n'-n}$ .*
- (ii) *If  $\pi'$  is a  $\theta$ -representation of  $\text{SO}_{2n'+1}^\epsilon$ , then  $n \geq n'$  and  $\pi \in R_{T_v \times T_w, 1 \otimes \theta_w}^{\text{Sp}_{2n}}$  for some  $v \in W_{n'}$  and  $w \in W_{n-n'}$ .*
- (iii)  *$\pi$  is a  $\theta$ -representation of  $\text{Sp}_{2n}$  if and only if  $\pi'$  is a unipotent representation of  $\text{SO}_{2n'+1}^\epsilon$ .*

*Proof.* The proof is similar in spirit to that of [Adams and Moy 1993, Theorem 3.5]. First of all, for  $G = \text{SO}_{2n'+1}$  from  $\theta_T = \chi_G|_T$  it follows that all the pairs  $(T_w, \theta_w)$ ,  $w \in W_n$ , are in the same geometric conjugacy class. More generally, two pairs of the form  $(T_{v_i} \times T_{w_i}, 1 \times \theta_{w_i})$ ,  $v_i \in W_{n_i}$ ,  $w_i \in W_{n'-n_i}$ ,  $i = 1, 2$ , are geometrically conjugate if and only if  $n_1 = n_2$ .

We need to recall some basic facts. Denote  $G \cdot G'$  by  $H$ , and the character  $R_T^G(\theta) \times R_{T'}^{G'}(\theta')$  by  $R_{T, T'}^H(\theta \cdot \theta')$ . These characters form an orthogonal basis for the space of uniform class functions of  $H$ . The set of irreducible characters  $\mathcal{E}(H)$  of  $H$  can be partitioned by geometric conjugacy classes (see, e.g., [Lusztig 1984])

$$\mathcal{E}(H) = \coprod_s \mathcal{E}(H, (s)),$$

where  $s = s_{G^*} \cdot s_{G'^*}$  runs over the semisimple conjugacy classes of the dual group  $H^* = G^* \cdot G'^*$ . The irreducible constituents of a single  $R_{T, T'}^H(\theta \times \theta')$  are all associated to one geometric conjugacy class  $(s)$ . Let  $\pi_H$  be a representation of  $H$ . Then we have

$$\pi_H = \bigoplus_s \pi_H(s),$$

where  $s$  runs over the semisimple conjugacy classes in  $H^*$  and  $\pi_H(s) \in \mathcal{E}(H, (s))$  is a subrepresentation of  $H$ . It is known that if  $\pi \in \mathcal{E}(H, (s))$  and  $\pi' \in \mathcal{E}(H, (s'))$  with  $(s) \neq (s')$ , then  $(\pi, \pi')_H = 0$ . We also have

$$\pi_H^\# = \bigoplus_s \pi_H(s)^\#.$$

If  $\pi_H(s) \neq 0$ , then it is a positive combination of representations in  $\mathcal{E}(H, (s))$ . It follows from the fact that the regular representation is uniform that  $\pi_H(s)^\#$  is nonzero.

Now let  $\pi_H = \pi \otimes \pi'$ , which occurs in  $\omega_{n, n'}^\epsilon$  by assumption. To prove (i), we note that if  $\pi$  is unipotent then  $\pi_H \in \mathcal{E}(H, (s))$  with  $s_{G^*} = 1$ . By the above argument,  $\pi_H^\# = \pi_H(s)^\# \neq 0$ ; hence it occurs in some

$$R_{T, 1}^{\text{Sp}_{2n}} \otimes R_{T', \theta}^{\text{SO}_{2n'+1}^\epsilon},$$

which has to appear in a summand of  $\omega_{n, n'}^\#$ . Then the conclusion follows from Theorem 3.1 and the first paragraph of the proof. The proof of (ii) and (iii) is similar and will be omitted. □

In particular, by Theorem 3.3 (i) and (ii), if  $n = n'$  then  $\pi$  is a unipotent representation of  $\text{Sp}_{2n}$  if and only if  $\pi'$  is a  $\theta$ -representation of  $\text{SO}_{2n+1}^\epsilon$ . Applying Theorems 3.2 and 3.3, we shall describe the theta correspondence between cuspidal unipotent representations of  $\text{Sp}_{2n}$  and cuspidal  $\theta$ -representations of  $\text{SO}_{2n'+1}$ . We remark that the latter notion is not vacuous thanks to the following simple lemma.

**Lemma 3.4.** *If  $\pi$  is a cuspidal unipotent representation of a finite classical group  $G$  which is not symplectic, then  $\chi_G \pi$  is a cuspidal  $\theta$ -representation.*

*Proof.* For any unipotent element  $u$ , by Lemma 2.2 we have

$$\chi_G \pi(u) = \chi_G(u) \pi(u) = \pi(u).$$

Hence  $\chi_G \pi$  is cuspidal. It is a  $\theta$ -representation by Lemma 2.3. □

**3C. First occurrence for symplectic-odd orthogonal pairs.** For an irreducible representation  $\pi$  of  $G_n$ , the smallest integer  $n'(\pi)$  such that  $\pi$  occurs in  $\omega_{n,n'(\pi)}$  is called the *first occurrence index* of  $\pi$  in the Witt tower  $\{G'_{n'}\}_{n' \geq 0}$ . By [Mœglin et al. 1987, Chapitre 3, Lemme IV.2], there exists  $n'$  such that  $\Theta_{G'_{n'}}(\pi) \neq 0$ , hence  $n'(\pi)$  is well-defined. In this section we compute the first occurrence of cuspidal unipotent and  $\theta$ -representations for symplectic-odd orthogonal dual pairs. We begin by reviewing Lusztig's results [1977] on the cuspidal unipotent representations of symplectic and orthogonal groups.

**Theorem 3.5.** *The groups*

- (i)  $U_n$ ,  $n = k(k+1)/2$ ,
- (ii)  $\mathrm{Sp}_{2n}$ ,  $n = k(k+1)$ ,
- (iii)  $\mathrm{SO}_{2n+1}$ ,  $n = k(k+1)$ ,
- (iv)  $\mathrm{SO}_{2n}^\epsilon$ ,  $n = k^2$ ,  $\epsilon = \mathrm{sgn}(-1)^k$

*are the only groups in their respective Lie families which possess a cuspidal unipotent representation. In each case, the specified group  $G$  has a unique cuspidal unipotent representation.*

By Lemma 3.4, we have the following immediate consequence for cuspidal  $\theta$ -representations.

**Corollary 3.6.** *The groups*

- (1)  $U_n$ ,  $n = k(k+1)/2$ ,
- (2)  $\mathrm{SO}_{2n+1}$ ,  $n = k(k+1)$ ,
- (3)  $\mathrm{SO}_{2n}^\epsilon$ ,  $n = k^2$ ,  $\epsilon = \mathrm{sgn}(-1)^k$

*are the only groups in their respective Lie families which possess a cuspidal  $\theta$ -representation. In each case, the specified group  $G$  has a unique cuspidal  $\theta$ -representation.*

We define a  $\theta$ -representation of  $O_{2n+1}$ , which is nonconnected, to be a constituent of  $\mathrm{Ind}_{\mathrm{SO}_{2n+1}}^{O_{2n+1}} \pi$  for a  $\theta$ -representation  $\pi$  of  $\mathrm{SO}_{2n+1}$ . Let  $\lambda'_k$  stand for the unique cuspidal unipotent representation of  $\mathrm{SO}_{2n+1}$ ,  $n = k(k+1)$ . Then

$$\lambda'_{k,\chi} := \chi_{\mathrm{SO}_{2n+1}} \lambda'_k$$

is the unique cuspidal  $\theta$ -representation of  $\mathrm{SO}_{2n+1}$ , and we have

$$\mathrm{Ind}_{\mathrm{SO}_{2n+1}}^{O_{2n+1}} \lambda'_{k,\chi} = \lambda'^+_{k,\chi} \oplus \lambda'^-_{k,\chi},$$

where  $\lambda'^+_{k,\chi} := \lambda'_{k,\chi} \otimes 1$  and  $\lambda'^-_{k,\chi} := \lambda'_{k,\chi} \otimes \mathrm{sgn}$  are cuspidal  $\theta$ -representations of  $O_{2n+1} \cong \mathrm{SO}_{2n+1} \times \{\pm I\}$ . Now we can give the first main result of this section, which is stated as Theorem 1.1 in the Introduction.

**Theorem 3.7.** *Let  $\lambda_k$  be the unique cuspidal unipotent representation of  $\mathrm{Sp}_{2n}$ ,  $n = k(k + 1)$ . Then the first occurrence index  $n^{\epsilon}(\lambda_k)$  in the Witt tower  $\{O_{2n'+1}^{\epsilon}\}_{n' \geq 0}$  is equal to  $n$ , and the theta lifting  $\Theta_{O_{2n+1}^{\epsilon}}(\lambda_k)$  equals  $\lambda_{k,\chi}^+$ .*

The proof of this theorem involves three ingredients. The first one is the following information which can be read from [Howe 1973].

**Proposition 3.8.** *Every irreducible representation of  $\mathrm{Sp}_{2n}$  is contained in  $\omega_{n,n}^+ \oplus \omega_{n,n}^-$ , where  $\omega_{n,n'}^{\epsilon} = \omega_{\mathrm{Sp}_{2n}, O_{2n'+1}^{\epsilon}}$ .*

The second ingredient is the following so-called conservation relation for cuspidal representations given in [Pan 2002].

**Proposition 3.9.** (i) *Let  $\pi$  be an irreducible cuspidal representation of  $\mathrm{Sp}_{2n}$  and  $n^{\epsilon}(\pi)$  be the first occurrence index of  $\pi$  in the Witt tower  $O_{\mathrm{odd}}^{\epsilon} = \{O_{2n'+1}^{\epsilon}\}_{n' \geq 0}$ . Then*

$$n^{'+}(\pi) + n^{-}(\pi) = 2n.$$

(ii) *Let  $\pi'$  be an irreducible cuspidal representation of  $O_{2n'+1}^{\epsilon}$  and  $n(\pi')$  be the first occurrence index of  $\pi'$  in the Witt tower  $\mathrm{Sp} = \{\mathrm{Sp}_{2n}\}_{n \geq 0}$ . Then*

$$n(\pi') + n(\pi' \otimes \mathrm{sgn}) = 2n' + 1.$$

A few remarks are in order. In the literature, the conservation relations are usually formulated in terms of the dimensions of vector spaces, but in this paper we will use the above form for convenience. Here we only recalled the statement for symplectic-odd orthogonal dual pairs and we refer the readers to [Pan 2002] for other cases. In the world of  $p$ -adic local fields, the conservation relations have been established in full generality in [Sun and Zhu 2015]. We should also mention that the standard formulation of conservation relations does not seem to hold in general for noncuspidal representations of finite dual pairs, and we will give some explanations at the end of the paper.

The last ingredient we need is that the action of  $-I$  under a unipotent representation is trivial. This fact might be known but we could not find a reference, so we include a detailed proof for completeness. It can be used to determine whether the theta lifting of a unipotent representation takes place in the even or odd part of the Weil representation, but we will not need this result.

**Proposition 3.10.** (i) *If  $\pi$  is a unipotent representation of  $\mathrm{GL}_n, U_n, \mathrm{Sp}_{2n}$  or  $\mathrm{SO}_{2n}^{\epsilon}$ , then  $\pi(-I)$  is trivial.*

(ii) *If  $\pi$  is a unipotent representation of  $\mathrm{Sp}_{2n}$  and  $\pi \otimes \pi'$  occurs in the Weil representation  $\omega_{n,n'}^{\epsilon}$  of  $(\mathrm{Sp}_{2n}, O_{2n'+1}^{\epsilon})$ , then  $\pi'(-I)$  is trivial.*

*Proof.* (ii) clearly follows from (i) because  $\pi(-I) = \pi'(-I)$  through the Weil representation  $\omega_{n,n'}^{\epsilon}$ . We shall prove (i) for all the classical groups (except odd

orthogonal groups) simultaneously by induction on  $n$ . We first observe that by (2-1),

$$(3-1) \quad R_{T,1}^G(-I) = R_{T,1}^G(I).$$

If  $n = 1$ , then  $G$  has only two unipotent representations  $1$  and  $\pi$ . Hence the conclusion follows from (3-1). Assume that  $n > 1$  and  $\pi \in R_{T,1}^G$  for some  $T$ . If  $T$  is not anisotropic, then it is contained in a Levi subgroup  $L$  of a proper parabolic subgroup  $P$  of  $G$ , i.e.,  $P = P^F$  for some  $F$ -stable proper parabolic subgroup  $P$  of  $G$ . By Frobenius reciprocity,

$$(\pi, R_{T,1}^G)_G = (\pi, \text{Ind}_P^G R_{T,1}^L)_G = (J_P^G(\pi), R_{T,1}^L)_L \neq 0,$$

where  $J_P^G$  is the Jacquet functor. We may write  $L = \text{GL}_{n_1} \times \text{GL}_{n_2} \times \cdots \times \text{GL}_{n_r} \times G_{n_0}$ , where  $n_i < n$ ,  $i = 0, \dots, r$ , and  $G_{n_0}$  is a classical group of the same type as  $G$ . There exists  $\pi_0 \in J_P^G(\pi)$  such that  $(\pi_0, R_{T,1}^L)_L \neq 0$ . Then by induction we have that  $\pi(-I) = \pi_0(-I)$  is trivial. By [Srinivasan 1979a, Theorem 6.25], if  $\pi \notin R_{T,1}^G$  for any  $T$  which is not anisotropic, then  $\pi$  is a cuspidal representation. If  $G = \text{GL}_n$  then we are done because it does not have cuspidal unipotent representations when  $n > 1$ . If  $G \neq \text{GL}_n$ , then it remains to consider the case that  $\pi$  is the unique cuspidal unipotent representation of  $G$  (if it exists). The uniqueness of  $\pi$  enables us to write

$$(\pi, R_{T,1}^G)\pi = R_{T,1}^G - \sum_{\pi_i \in R_{T,1}^G, \pi_i \neq \pi} (\pi_i, R_{T,1}^G)_G \pi_i,$$

where the  $\pi_i$ 's are all unipotent representations but not cuspidal. Then the conclusion follows from (3-1) and that  $\pi_i(-I)$  is trivial for each  $i$ . □

*Proof of Theorem 3.7.* By Proposition 3.8, the cuspidal unipotent representation  $\lambda_k$  of  $\text{Sp}_{2n}$  is contained in  $\omega_{n,n}^+ \oplus \omega_{n,n}^-$ . By Theorem 3.3,  $\lambda_k$  is not contained in any  $\omega_{n,n'}^\epsilon$  when  $n' < n$ , and for some  $\epsilon_0 \in \{+, -\}$  the theta lifting  $\Theta_{O_{2n+1}^{\epsilon_0}}(\lambda_k)$  to  $O_{2n+1}^{\epsilon_0}$  is the first occurrence in the Witt tower  $O_{\text{odd}}^{\epsilon_0}$ ; hence it is a cuspidal  $\theta$ -representation. Then it follows from Corollary 3.6 and Proposition 3.10 that  $\Theta_{O_{2n+1}^{\epsilon_0}}(\lambda_k) = \lambda_{k,\chi}^{\prime+}$ . The conservation relation Proposition 3.9(i) implies that  $n'^{-\epsilon_0}(\lambda_k) = 2n - n'^{\epsilon_0}(\lambda_k) = n$ , and one can similarly deduce that  $\Theta_{O_{2n+1}^{-\epsilon_0}}(\lambda_k) = \lambda_{k,\chi}^{\prime+}$  as well. □

We also study the first occurrence of the theta lifting of cuspidal  $\theta$ -representations of  $\text{Sp}_{2n}$  in  $O_{\text{odd}}^\epsilon$ , which by Theorem 3.3 will be cuspidal unipotent representations. By definition, a unipotent representation of  $O_{2n+1}$  is a constituent of  $\text{Ind}_{\text{SO}_{2n+1}}^{O_{2n+1}}(\pi)$  for a unipotent representation  $\pi$  of  $\text{SO}_{2n+1}$ . Let  $\lambda'_k$  be the unique cuspidal unipotent representation of  $\text{SO}_{2n+1}$ ,  $n = k(k+1)$ . Then we have

$$(3-2) \quad \text{Ind}_{\text{SO}_{2n+1}}^{O_{2n+1}}(\lambda'_k) = \lambda_k^{\prime+} \oplus \lambda_k^{\prime-},$$

where  $\lambda_k^{\prime+} := \lambda'_k \otimes 1$  and  $\lambda_k^{\prime-} := \lambda'_k \otimes \text{sgn}$  are cuspidal unipotent representations of  $O_{2n+1} \cong \text{SO}_{2n+1} \times \{\pm I\}$ .

Applying a result of Lusztig, we first obtain the following classification of cuspidal  $\theta$ -representations of finite symplectic groups.

**Theorem 3.11.** *The groups  $\mathrm{Sp}_{2n}$ ,  $n = k^2$ , are the only symplectic groups which possess cuspidal  $\theta$ -representations. Each  $\mathrm{Sp}_{2k^2}$ ,  $k \neq 0$ , has two cuspidal  $\theta$ -representations  $\lambda_{k,\alpha}$  and  $\lambda_{k,\beta}$ , which satisfy  $\lambda_{k,i}(-I) = (-1)^k$ ,  $i = \alpha, \beta$ .*

*Proof.* We first recall some general results. Assume that  $G$  is a simple simply connected linear algebraic group over  $F$ , and  $T$  is an  $F$ -stable maximal torus of  $G$ . Let  $G^*$  be the dual group of  $G$ , and  $T^*$  be the  $F$ -stable maximal torus of  $G^*$  which is the dual of  $T$ . Write  $G^* = G^{*F}$ ,  $T^* = T^{*F}$ . Then a character  $\theta$  of  $T$  corresponds to some element  $s \in T^*$ , which is well-defined up to conjugacy by  $W_{G^*}(T^*)^F$ . In this case we write  $R_{T^*,s}^G$  instead of  $R_{T,\theta}^G$ . Let  $\mathcal{E}(G, (s))$  be the set of irreducible representations which occur in  $R_{T^*,s}^G$  for some  $T^*$  such that  $s \in T^*$ . By [Lusztig 1988], there is a bijection

$$\mathcal{L} : \mathcal{E}(G, (s)) \rightarrow \mathcal{E}(C_{G^*}(s), (1)),$$

where  $\mathcal{E}(C_{G^*}(s), (1))$  is the set of irreducible representations of  $C_{G^*}(s)$  whose restrictions to  $C_{G^*}(s)^\circ$  are sums of unipotent representations of  $C_{G^*}(s)^\circ$ . Note that  $C_{G^*}(s)$  may not be connected in general. Moreover if the identity components of the centers of  $G$  and  $C_{G^*}(s)$  have the same  $F$ -rank, then  $\pi \in \mathcal{E}(G, (s))$  is cuspidal if and only if  $\mathcal{L}(\pi) \in \mathcal{E}(C_{G^*}(s), (1))$  is cuspidal (see, e.g., [Lusztig 1977, Chapter 9] and [Malle 2017, Lemma 2.7]).

For  $G = \mathrm{Sp}_{2n}$  we have  $G^* = \mathrm{SO}_{2n+1}$ . The pair  $(T_w, \theta_w)$  corresponds to  $s \in T_w$ , where  $s = (-I, 1)$  with  $I$  being the identity in  $\mathrm{SO}_{2n}^{\epsilon_w} \subset \mathrm{SO}_{2n+1}$ . Then

$$C_{G^*}(s) \cong O_{2n}^{\epsilon_w} \subset \mathrm{SO}_{2n+1};$$

hence by the above results we have a bijection between cuspidal  $\theta$ -representations of  $\mathrm{Sp}_{2n}$  and the union of cuspidal unipotent representations of  $O_{2n}^\epsilon$  over  $\epsilon = \pm$ . By Theorem 3.5, each  $O_{2k^2}^\epsilon$  with  $\epsilon := \mathrm{sgn}(-1)^k$  has two cuspidal unipotent representations, which are the two constituents induced from the unique cuspidal unipotent representation of  $\mathrm{SO}_{2k^2}^\epsilon$ , and these are the only even orthogonal groups which possess cuspidal unipotent representations. Hence, each  $\mathrm{Sp}_{2n}$ ,  $n = k^2$ , has two cuspidal  $\theta$ -representations  $\lambda_{k,1}$  and  $\lambda_{k,2}$ , and these are the only possibilities.

From [Howe 1973] we also know that every irreducible representation of  $\mathrm{Sp}_{2n}$  is contained in  $\omega_{\mathrm{Sp}_{2n}, O_{2n}^+} \oplus \omega_{\mathrm{Sp}_{2n}, O_{2n}^-}$ . Each  $\lambda_{k,i}$ ,  $i = \alpha, \beta$ , occurs in some  $R_{T_w, \theta_w}^G$  such that  $\epsilon_w = \epsilon = \mathrm{sgn}(-1)^k$ . Similar to the proof of Theorems 3.3 and 3.7, applying the main result of [Srinivasan 1979b] as well as conservation relations we can show that  $\lambda_{k,i}$  occurs in  $\omega_{\mathrm{Sp}_{2n}, O_{2n}^\epsilon}$  and  $\Theta_{O_{2n}^\epsilon}(\lambda_{k_i})$ ,  $i = \alpha, \beta$ , are the two cuspidal  $\theta$ -representations of  $O_{2n}^\epsilon$ . Since  $\chi_{\mathrm{SO}_{2n}^\pm}(-I) = \pm 1$ , we obtain that  $\lambda_{k,i}(-I) = \chi_{\mathrm{SO}^\epsilon}(-I) = (-1)^k$ . □

Finally, we can give the second main result of this section.

**Theorem 3.12.** *Let  $\lambda_{k,i}$ ,  $i = \alpha, \beta$ , be the cuspidal  $\theta$ -representations of  $\mathrm{Sp}_{2k^2}$ , and let  $n^{\epsilon}(\lambda_{k,i})$  be the first occurrence index of  $\lambda_{k,i}$  in the Witt tower  $\mathbf{O}_{\mathrm{odd}}^{\epsilon}$ . Then for some  $\epsilon_0 \in \{+, -\}$  one has  $n^{\epsilon_0}(\lambda_{k,\alpha}) = n'^{-\epsilon_0}(\lambda_{k,\beta}) = k(k-1)$  and  $n'^{-\epsilon_0}(\lambda_{k,\alpha}) = n^{\epsilon_0}(\lambda_{k,\beta}) = k(k+1)$ . Write  $\epsilon(k) = \mathrm{sgn}(-1)^k$ . Then the theta liftings are given by*

$$\begin{aligned} \Theta_{O_{2k(k-1)+1}^{\epsilon_0}}(\lambda_{k,\alpha}) &= \lambda'_{k-1}{}^{\epsilon(k)}, & \Theta_{O_{2k(k+1)+1}^{-\epsilon_0}}(\lambda_{k,\alpha}) &= \lambda'_k{}^{\epsilon(k)}, \\ \Theta_{O_{2k(k+1)+1}^{\epsilon_0}}(\lambda_{k,\beta}) &= \lambda'_k{}^{\epsilon(k)}, & \Theta_{O_{2k(k-1)+1}^{-\epsilon_0}}(\lambda_{k,\beta}) &= \lambda'_{k-1}{}^{\epsilon(k)}, \end{aligned}$$

where  $\lambda'_k{}^{\epsilon}$  is defined by (3-2).

We remark that the  $\epsilon_0$  above depends on the choice of the nontrivial character  $\psi$  of  $F$  which we used to define the Weil representations. If  $\psi$  is replaced by  $\psi_t(x) := \psi(tx)$  for some  $t \in F^{\times} \setminus F^{\times 2}$ , then  $\epsilon_0$  should be replaced by  $-\epsilon_0$ .

*Proof.* We use an inductive argument which is parallel to the proof in [Adams and Moy 1993]. By convention we let  $\lambda_0$  be the trivial representation of  $\mathrm{Sp}_0$ .

For  $\mathrm{Sp}_2$ ,  $\lambda_{1,\alpha}$  is contained in  $\omega_{1,1}^+ \oplus \omega_{1,1}^-$ . Since by Theorem 3.3(iii) the first occurrence of the theta lifting of  $\lambda_{1,\alpha}$  is unipotent cuspidal, we see that  $\lambda_{1,\alpha}$  occurs in  $\omega_{1,0}^{\epsilon_0}$  for some  $\epsilon_0$ . By conservation relation,  $\lambda_{1,\alpha}$  occurs in  $\omega_{1,2}^{-\epsilon_0}$  as the first occurrence in  $\mathbf{O}_{\mathrm{odd}}^{-\epsilon_0}$ . By Theorem 3.11, we have the theta liftings  $\Theta_{O_1^{\epsilon_0}}(\lambda_{1,\alpha}) = \lambda_0'^{-}$  and  $\Theta_{O_5^{-\epsilon_0}}(\lambda_{1,\alpha}) = \lambda_2'^{-}$ . Similarly  $\lambda_{1,\beta}$  is contained in  $\omega_{1,0}^+ \oplus \omega_{1,0}^-$ . It cannot happen that  $\Theta_{O_1^{\epsilon_0}}(\lambda_{1,\alpha}) = \Theta_{O_1^{\epsilon_0}}(\lambda_{1,\beta}) = \lambda_0'^{-}$ , hence  $\lambda_{1,\beta}$  must be contained in  $\omega_{1,0}^{-\epsilon_0}$ . By conservation again,  $\lambda_{1,\beta}$  occurs in  $\omega_{1,2}^{\epsilon_0}$  as the first occurrence in  $\mathbf{O}_{\mathrm{odd}}^{\epsilon_0}$ . The theta liftings of  $\lambda_{1,\beta}$  to  $O_1^{-\epsilon_0}$  and  $O_5^{\epsilon_0}$  are therefore  $\lambda_0'^{-}$  and  $\lambda_2'^{-}$ , respectively. We may exhaust the theta correspondence inductively through the following diagram:

$$\begin{array}{llll} \mathrm{Sp}_0 & \lambda_0 & \longrightarrow & \lambda_0^+ = 1 & \omega_{0,0}^{\epsilon_0} \\ \mathrm{Sp}_0 & \lambda_0 & \longrightarrow & \lambda_0^+ = 1 & \omega_{0,0}^{-\epsilon_0} \\ \mathrm{Sp}_2 & \lambda_{1,\alpha} & \longrightarrow & \lambda_0'^{-} = \mathrm{sgn} & \omega_{1,0}^{\epsilon_0} \\ \mathrm{Sp}_2 & \lambda_{1,\beta} & \longrightarrow & \lambda_0'^{-} = \mathrm{sgn} & \omega_{1,0}^{-\epsilon_0} \\ \mathrm{Sp}_2 & \lambda_{1,\alpha} & \longrightarrow & \lambda_1'^{-} & \omega_{1,2}^{-\epsilon_0} \\ \mathrm{Sp}_2 & \lambda_{1,\beta} & \longrightarrow & \lambda_1'^{-} & \omega_{1,2}^{\epsilon_0} \\ & & & \vdots & \\ \mathrm{Sp}_{2k^2} & \lambda_{k,\alpha} & \longrightarrow & \lambda_{k-1}'{}^{\epsilon(k)} & \omega_{k^2,k(k-1)}^{\epsilon_0} \\ \mathrm{Sp}_{2k^2} & \lambda_{k,\beta} & \longrightarrow & \lambda_{k-1}'{}^{\epsilon(k)} & \omega_{k^2,k(k-1)}^{-\epsilon_0} \\ \mathrm{Sp}_{2k^2} & \lambda_{k,\alpha} & \longrightarrow & \lambda_k'{}^{\epsilon(k)} & \omega_{k^2,k(k+1)}^{-\epsilon_0} \\ \mathrm{Sp}_{2k^2} & \lambda_{k,\beta} & \longrightarrow & \lambda_k'{}^{\epsilon(k)} & \omega_{k^2,k(k+1)}^{\epsilon_0} \end{array}$$

In general for  $\mathrm{Sp}_{2k^2}$ , by Proposition 3.8,  $\lambda_{k,i}$ ,  $i = \alpha, \beta$ , are contained in  $\omega_{k^2,k^2}^+ \oplus \omega_{k^2,k^2}^-$ . Since the first occurrences of their theta liftings are cuspidal unipotent, they are contained in  $\omega_{k^2,k(k-1)}^+ \oplus \omega_{k^2,k(k-1)}^-$ . By induction we have

$$\Theta_{O_{2k(k-1)+1}^{\epsilon_0}}(\lambda_{k-1,\beta}) = \lambda_{k-1}^{\epsilon(k-1)}.$$

Then by the conservation relation given by Proposition 3.9(ii) we may arrange the indices such that

$$\Theta_{O_{2k(k-1)+1}^{\epsilon_0}}(\lambda_{k,\alpha}) = \lambda_{k-1}^{\epsilon(k-1)} \otimes \mathrm{sgn} = \lambda_{k-1}^{\epsilon(k)}.$$

By conservation again,  $\lambda_{k,\alpha}$  occurs in  $\omega_{k^2,k(k+1)}^{-\epsilon_0}$ . In a similar way,  $\lambda_{k,\beta}$  occurs in  $\omega_{k^2,k(k-1)}^{-\epsilon_0}$  and  $\omega_{k^2,k(k+1)}^{\epsilon_0}$ . The remaining assertions about the theta liftings are clear. □

### 4. Theta correspondence and Harish-Chandra series

In this section we study the theta correspondence of representations in certain Harish-Chandra series in a way similar to [Aubert et al. 1996].

Let  $T_l$  be the split torus over  $F$  of rank  $l$ , and  $\theta_l := \theta_{T_l}$  the order 2 character of  $T_l$  defined in Section 2D. In particular  $(T_1, \theta_1) = (\mathrm{GL}_1, \chi_{\mathrm{GL}_1})$ . For later use, for  $k, l \geq 0$  we introduce a pair of quadratic characters  $\theta_{k,l}$  and  $\theta'_{k,l}$  of  $T_l$  by

$$(4-1) \quad \theta_{k,l} = \begin{cases} 1_k \otimes \theta_{l-k} & \text{if } k \leq l, \\ 1 & \text{otherwise,} \end{cases}$$

and

$$(4-2) \quad \theta'_{k,l} = \theta_{k,l} \theta_l = \begin{cases} \theta_k \otimes 1_{l-k} & \text{if } k \leq l, \\ \theta_l & \text{otherwise,} \end{cases}$$

where  $1_k$  and  $1_{l-k}$  are the trivial characters of  $T_k$  and  $T_{l-k}$ , respectively.

For two groups  $G_n \subset G_m$  in the same Witt tower, we can embed  $G_n$  into the Levi subgroup  $G_n \times T_{m-n}$  of a parabolic subgroup of  $G_m$ . Let  $\lambda$  be a cuspidal representation of  $G_n$ , and denote by  $R(G_m)_\lambda$  the subset of  $R(G_m)$  whose elements are spanned by

$$\mathrm{Irr}(G_m)_\lambda := \mathrm{Irr}(R_{G_n \times T_{m-n}}^{G_m}(\lambda \otimes 1)).$$

Similarly introduce the subset  $R(G_m)_{\lambda,\theta}$  of  $R(G_m)$  whose elements are spanned by

$$\mathrm{Irr}(G_m)_{\lambda,\theta} := \mathrm{Irr}(R_{G_n \times T_{m-n}}^{G_m}(\lambda \otimes \theta_{m-n})).$$

Note that in this context the functor  $R$  above stands for the usual parabolic induction, which coincides with the generalized Deligne–Lusztig induction.

The standard Levi subgroups  $L$  of  $G_m$  are of the form

$$L = \mathrm{GL}_{n_1}(q) \times \cdots \times \mathrm{GL}_{n_r}(q) \times G_l(q)$$

such that  $m = n_1 + \cdots + n_r + l$ . A representation  $\rho$  of  $L$  is a cuspidal, unipotent

or  $\theta$ -representation if it is of the form  $\rho = \rho_1 \otimes \cdots \otimes \rho_r \otimes \sigma$ , where  $\rho_i$  and  $\sigma$  are cuspidal, unipotent or  $\theta$ -representations of  $\text{GL}_{n_i}(q)$  and  $G_1(q)$ , respectively.

For a finite (reductive) group  $G$ , let  $(R^G, \overline{\mathcal{Q}}_\ell[G])$  be the regular representation of  $G \times G$  on the space of  $\overline{\mathcal{Q}}_\ell$ -valued functions  $f$  on  $G$  defined by

$$(R^G(g_1, g_2)f)(g) = f(g_1^{-1}gg_2), \quad \text{for all } g_1, g, g_2 \in G.$$

Then  $R^G$  has the well-known Peter–Weyl decomposition

$$R^G = \sum_{\pi \in \text{Irr}(G)} \pi \otimes \pi^\vee,$$

where  $\pi^\vee$  is the contragredient of  $\pi$ . For a quadratic character  $\chi$  of  $G$ , let  $\chi R^G$  be the representation of  $G \times G$  defined by

$$\chi R^G = \sum_{\pi \in \text{Irr}(G)} \chi \pi \otimes \pi^\vee = \sum_{\pi \in \text{Irr}(G)} \pi \otimes \chi \pi^\vee.$$

The following result given in [Aubert et al. 1996, Théorème 3.7] shows that the theta correspondence is compatible with the Harish-Chandra decomposition and that it preserves cuspidal unipotent representations for the dual pairs  $(U_n(q), U_{n'}(q))$  and  $(\text{Sp}_{2n}(q), O_{2n'}^\epsilon(q))$ .

**Theorem 4.1.** *Assume  $(\mathcal{T}, \mathcal{T}') = (U^\epsilon, U^{\epsilon'})$ ,  $(\text{Sp}, O_{\text{even}}^\epsilon)$  or  $(O_{\text{even}}^\epsilon, \text{Sp})$ ,  $\epsilon, \epsilon' = \pm$ . Let  $\lambda$  be an irreducible cuspidal representation of  $G_n \in \mathcal{T}$  and let  $n' = n'(\lambda)$  be its first occurrence index in  $\mathcal{T}'$ , so that  $\lambda' := \Theta_{G_{n'}}^\epsilon(\lambda)$  is an irreducible cuspidal representation of  $G_{n'}' \in \mathcal{T}'$ . Then for  $\gamma \in \text{Irr}(G_m)_\lambda$ ,  $\Theta_{G_{m'}}^{\epsilon'}(\lambda) = 0$  whenever  $m' < n'$  and  $\Theta_{G_{m'}}^{\epsilon'}(\gamma) \in R(G_{m'}')_{\lambda'}$  otherwise.*

We are going to extend this theorem to the Witt tower  $(\text{Sp}, O_{\text{odd}}^\epsilon)$ . Since  $O_{2n'+1}^+$  and  $O_{2n'+1}^-$  are isomorphic, we may simply write  $\text{Irr}(O_{2n'+1})$  instead of  $\text{Irr}(O_{2n'+1}^+)$  and  $\text{Irr}(O_{2n'+1}^-)$ . The main result of this section is the following theorem, which is compatible with Pan’s decomposition formula (Theorem 3.1).

**Theorem 4.2.** *Let  $\lambda$  be an irreducible cuspidal representation of  $\text{Sp}_{2n}$  and let  $n' = n'(\lambda)$  be its first occurrence index in  $O_{\text{odd}}^\epsilon$ , so that  $\lambda' := \Theta_{O_{2n'+1}}^\epsilon(\lambda)$  is an irreducible cuspidal representation of  $O_{2n'+1}^\epsilon$ . Then the following holds.*

- (i) *For  $\gamma \in \text{Irr}(\text{Sp}_{2m})_{\lambda, \theta}$ ,  $\Theta_{O_{2m'+1}}^\epsilon(\gamma) = 0$  if  $m' < n'$  and  $\Theta_{O_{2m'+1}}^\epsilon(\gamma) \in R(O_{2m'+1})_{\lambda'}$  otherwise.*
- (ii) *For  $\gamma' \in \text{Irr}(O_{2m'+1}^\epsilon)_{\lambda'}$ ,  $\Theta_{\text{Sp}_{2m}}(\gamma') = 0$  if  $m < n$  and  $\Theta_{\text{Sp}_{2m}}(\gamma') \in R(\text{Sp}_{2m})_{\lambda, \theta}$  otherwise.*
- (iii) *For  $\gamma \in \text{Irr}(\text{Sp}_{2m})_\lambda$ ,  $\Theta_{O_{2m'+1}}^\epsilon(\gamma) = 0$  if  $m' < n'$  and  $\Theta_{O_{2m'+1}}^\epsilon(\gamma) \in R(O_{2m'+1}^\epsilon)_{\lambda', \theta'}$  otherwise, where  $\theta' := \theta'_{m-n, m'-n'}$  is defined by (4-2).*

(iv) For  $\gamma' \in \text{Irr}(O_{2m'+1}^\epsilon)^{\lambda', \theta}$ ,  $\Theta_{\text{Sp}_{2m}}(\gamma') = 0$  if  $m < n$  and  $\Theta_{\text{Sp}_{2m}}(\gamma') \in R(\text{Sp}_{2m})_{\lambda, \theta'}$ , where  $\theta' := \theta_{m'-n', m-n}$  is defined by (4-1).

*Proof.* We will only prove (i) and (iii). The proofs of (ii) and (iv) are similar and will be left to the reader. The proofs are parallel to Theorem 4.1 and we shall follow the calculations in [Aubert et al. 1996] and [Mœglin et al. 1987] closely.

We first prove (i), by induction on  $m$ .

• Suppose that  $m = n$  (i.e.,  $\gamma = \lambda$  is cuspidal) and  $m' > n'$ . It is known (see [Mœglin et al. 1987, Chapitre 3]) that each constituent  $\gamma' \in \Theta_{O_{2m'+1}^\epsilon}(\gamma)$  is noncuspidal. Let  $j$  be the maximum integer such that

$$\gamma \in R_{O_{2(m'-j)+1} \times \text{GL}_j}^{O_{2m'+1}}(\lambda'_1 \otimes \rho')$$

with  $\lambda'_1 \in \text{Irr}(O_{2(m'-j)+1})$  cuspidal and  $\rho' \in \text{Irr}(\text{GL}_j)$ . Since  $\gamma' \in \Theta_{O_{2m'+1}^\epsilon}(\gamma)$ ,

$$\begin{aligned} 0 &< (\omega_{m,m'}^\epsilon, \gamma \otimes \gamma')_{\text{Sp}_{2m} \times O_{2m'+1}} \\ &\leq (\omega_{m,m'}^\epsilon, \gamma \otimes R_{O_{2(m'-j)+1} \times \text{GL}_j}^{O_{2m'+1}}(\lambda'_1 \otimes \rho'))_{\text{Sp}_{2m} \times O_{2m'+1}} \\ &= (*R_{O_{2(m'-j)+1} \times \text{GL}_j}^{O_{2m'+1}}(\omega_{m,m'}^\epsilon), \gamma \otimes \lambda'_1 \otimes \rho')_{\text{Sp}_{2m} \times O_{2(m'-j)+1} \times \text{GL}_j}. \end{aligned}$$

Here  $*R$  stands for the Jacquet functor, which is adjoint to the induction functor  $R$ . We have the following decomposition (see [Mœglin et al. 1987, Chapitre 3, IV Théorème 5]):

$$\begin{aligned} &*R_{O_{2(m'-j)+1} \times \text{GL}_j}^{O_{2m'+1}}(\omega_{m,m'}^\epsilon) \\ &= \bigoplus_{i=0}^{\min(m,j)} R_{\text{Sp}_{2(m-i)} \times \text{GL}_i \times (\text{GL}_{j-i} \times \text{GL}_i) \times O_{2(m'-j)+1}}^{\text{Sp}_{2m} \times \text{GL}_j \times O_{2(m'-j)+1}}(\omega_{m-i,m'-j}^\epsilon \otimes 1_{\text{GL}_{j-i}} \otimes \chi_{\text{GL}_i} R^{\text{GL}_i}). \end{aligned}$$

Hence  $(\omega_{m,m'}^\epsilon, \gamma \otimes \gamma')$  is bounded by

$$\sum_{i=0}^{\min(m,j)} \left( \omega_{m-i,m'-j}^\epsilon \otimes 1_{\text{GL}_{j-i}} \otimes \chi_{\text{GL}_i} R^{\text{GL}_i}, *R_{\text{Sp}_{2(m-i)} \times \text{GL}_i \times (\text{GL}_{j-i} \times \text{GL}_i) \times O_{2(m'-j)+1}}^{\text{Sp}_{2m} \times \text{GL}_j \times O_{2(m'-j)+1}}(\gamma \otimes \lambda'_1 \otimes \rho') \right),$$

where the scalar product in the  $i$ -th summand is taken over the group

$$\text{Sp}_{2(m-i)} \times O_{2(m'-j)+1} \times \text{GL}_{j-i} \times \text{GL}_i \times \text{GL}_i.$$

Since  $\gamma = \lambda$  is cuspidal, the only nonzero term corresponds to  $i = 0$ , which implies

$$(\omega_{m,m'-j}^\epsilon \otimes 1_{\text{GL}_j}, \gamma \otimes \lambda'_1 \otimes \rho') > 0.$$

It follows that  $\rho' = 1_{\text{GL}_j}$  and  $\lambda'_1 \in \Theta_{O_{2(m'-j)+1}^\epsilon}(\gamma)$ . Because  $\lambda'_1$  is cuspidal, we must have  $m' - j = n'$  and  $\lambda'_1 = \lambda'$ , i.e.,  $\gamma' \in \text{Irr}(O_{2m'+1})_{\lambda'}$ .

• Suppose that  $m > n$ . If  $\gamma \in \text{Irr}(\text{Sp}_{2m})_{\lambda, \theta}$ , then there exists  $\gamma_1 \in \text{Irr}(\text{Sp}_{2(m-1)})_{\lambda, \theta}$  such that  $\gamma \in R_{\text{Sp}_{2(m-1)} \times \text{GL}_1}^{\text{Sp}_{2m}}(\gamma_1 \otimes \chi_{\text{GL}_1})$ . For  $\gamma' \in \Theta_{O_{2m'+1}}^\epsilon(\gamma)$  we have

$$\begin{aligned} 0 &< (\omega_{m,m'}^\epsilon, \gamma \otimes \gamma')_{\text{Sp}_{2m} \times O_{2m'+1}} \\ &\leq (\omega_{m,m'}^\epsilon, R_{\text{Sp}_{2(m-1)} \times \text{GL}_1}^{\text{Sp}_{2m}}(\gamma_1 \otimes \chi_{\text{GL}_1}) \otimes \gamma')_{\text{Sp}_{2m} \times O_{2m'+1}} \\ &= (*R_{\text{Sp}_{2(m-1)} \times \text{GL}_1}^{\text{Sp}_{2m}}(\omega_{m,m'}^\epsilon), \gamma_1 \otimes \chi_{\text{GL}_1} \otimes \gamma')_{\text{Sp}_{2(m-1)} \times \text{GL}_1 \times O_{2m'+1}} \end{aligned}$$

We have the decomposition

$$\begin{aligned} &*R_{\text{Sp}_{2(m-1)} \times \text{GL}_1}^{\text{Sp}_{2m}}(\omega_{m,m'}^\epsilon) \\ &= \omega_{m-1,m'}^\epsilon \otimes \chi_{\text{GL}_1} + R_{\text{Sp}_{2(m-1)} \times O_{2m'+1} \times \text{GL}_1}^{\text{Sp}_{2(m-1)} \times O_{2m'-1} \times \text{GL}_1 \times \text{GL}_1}(\omega_{m-1,m'-1}^\epsilon \otimes \chi_{\text{GL}_1} R^{\text{GL}_1}). \end{aligned}$$

Hence  $(\omega_{m,m'}^\epsilon, \gamma \otimes \gamma')$  is bounded by

$$\begin{aligned} &(\omega_{m-1,m'}^\epsilon \otimes \chi_{\text{GL}_1}, \gamma_1 \otimes \gamma' \otimes \chi_{\text{GL}_1})_{\text{Sp}_{2(m-1)} \times O_{2m'+1} \times \text{GL}_1} \\ &+ (\omega_{m-1,m'-1}^\epsilon \otimes \chi_{\text{GL}_1} R^{\text{GL}_1}, \gamma_1 \otimes *R_{O_{2m'-1} \times \text{GL}_1}^{O_{2m'+1}}(\gamma') \otimes \chi_{\text{GL}_1})_{\text{Sp}_{2(m-1)} \times O_{2m'-1} \times \text{GL}_1 \times \text{GL}_1}. \end{aligned}$$

If the first term is nonzero, then  $\gamma' \in \Theta_{O_{2m'+1}}^\epsilon(\gamma_1)$ . By induction,  $\gamma' \in R(O_{2m'+1})_{\lambda'}$ . If the second term is nonzero, then there exist  $\gamma'_1 \in \text{Irr}(O_{2m'-1})$  and  $\rho' \in \text{Irr}(\text{GL}_1)$  such that  $\gamma'_1 \otimes \rho' \in *R_{O_{2m'-1} \times \text{GL}_1}^{O_{2m'+1}}(\gamma')$  and

$$(\omega_{m-1,m'-1}^\epsilon \otimes \chi_{\text{GL}_1} R^{\text{GL}_1}, \gamma_1 \otimes \gamma'_1 \otimes \rho' \otimes \chi_{\text{GL}_1})_{\text{Sp}_{2(m-1)} \times O_{2m'-1} \times \text{GL}_1 \times \text{GL}_1} \neq 0.$$

It follows that  $\rho' = 1$  and  $\gamma'_1 \in \Theta_{O_{2m'-1}}^\epsilon(\gamma_1)$ . By induction,  $\gamma'_1 \in \text{Irr}(O_{2m'-1})_{\lambda'}$  and therefore  $\gamma' \in \text{Irr}(O_{2m'+1})_{\lambda'}$  by Frobenius reciprocity. This proves (i).

Next we prove (iii), again by induction on  $m$ . Assume that  $\gamma \in \text{Irr}(\text{Sp}_{2m})_\lambda$  and  $\gamma' \in \Theta_{O_{2m'+1}}^\epsilon(\gamma)$ . The case  $m = n$  is covered by (i), so we may assume that  $m > n$ . Then there exists  $\gamma_1 \in \text{Irr}(\text{Sp}_{2(m-1)})_\lambda$  such that  $\gamma \in R_{\text{Sp}_{2(m-1)} \times \text{GL}_1}^{\text{Sp}_{2m}}(\gamma_1 \otimes 1)$ . Since  $\gamma' \in \Theta_{O_{2m'+1}}^\epsilon(\gamma)$ , we have

$$\begin{aligned} 0 &< (\omega_{m,m'}^\epsilon, \gamma \otimes \gamma')_{\text{Sp}_{2m} \times O_{2m'+1}} \leq (\omega_{m,m'}^\epsilon, R_{\text{Sp}_{2(m-1)} \times \text{GL}_1}^{\text{Sp}_{2m}}(\gamma_1 \otimes 1) \otimes \gamma')_{\text{Sp}_{2m} \times O_{2m'+1}} \\ &= (*R_{\text{Sp}_{2(m-1)} \times \text{GL}_1}^{\text{Sp}_{2m}}(\omega_{m,m'}^\epsilon), \gamma_1 \otimes 1 \otimes \gamma')_{\text{Sp}_{2(m-1)} \times \text{GL}_1 \times O_{2m'+1}}. \end{aligned}$$

Similar to the proof of (i),  $(\omega_{m,m'}^\epsilon, \gamma \otimes \gamma')$  is bounded by

$$\begin{aligned} &(\omega_{m-1,m'}^\epsilon \otimes \chi_{\text{GL}_1}, \gamma_1 \otimes \gamma' \otimes 1)_{\text{Sp}_{2(m-1)} \times O_{2m'+1} \times \text{GL}_1} \\ &+ (\omega_{m-1,m'-1}^\epsilon \otimes \chi_{\text{GL}_1} R^{\text{GL}_1}, \gamma_1 \otimes *R_{O_{2m'-1} \times \text{GL}_1}^{O_{2m'+1}}(\gamma') \otimes 1)_{\text{Sp}_{2(m-1)} \times O_{2m'-1} \times \text{GL}_1 \times \text{GL}_1}. \end{aligned}$$

The first term is zero. If the second term is nonzero, then there exists  $\gamma'_1 \in \text{Irr}(O_{2m'-1})$

and  $\rho' \in \text{Irr}(\text{GL}_1)$  such that  $\gamma'_1 \otimes \rho' \in {}^*R_{O_{2m'-1} \times \text{GL}_1}^{O_{2m'+1}}(\gamma')$  and

$$(\omega_{m-1, m'-1}^\epsilon \otimes \chi_{\text{GL}_1} R^{\text{GL}_1}, \gamma_1 \otimes \gamma'_1 \otimes \rho' \otimes 1)_{\text{Sp}_{2(m-1)} \times O_{2m'-1} \times \text{GL}_1 \times \text{GL}_1} \neq 0.$$

It follows that  $\rho' = \chi_{\text{GL}_1}$  and  $\gamma'_1 \in \Theta_{O_{2m'-1}}^\epsilon(\gamma_1)$ . By induction,

$$\gamma'_1 \in R_{O_{2n'+1} \times T_{m'-n'-1}}^{O_{2m'-1}}(\lambda' \otimes \theta'_{m-n-1, m'-n'-1}),$$

hence by Frobenius reciprocity

$$\gamma' \in R_{O_{2n'+1} \times T_{m'-n'}}^{O_{2m'+1}}(\lambda' \otimes \theta'_{m-n, m'-n'}).$$

This finishes the proof of (iii). □

In particular, if we consider the Harish-Chandra series of cuspidal unipotent representations, then combining Theorem 3.3 we obtain the following immediate consequence.

**Corollary 4.3.** *Let  $\lambda := \lambda_k$  be the unique cuspidal unipotent representation of  $\text{Sp}_{2n}$ ,  $n = k(k + 1)$ , and  $\lambda' := \lambda_{k, \chi}^{'+}$  be its theta lifting to  $O_{2n+1}^\epsilon$ . Then the following holds.*

(i) *For  $\gamma \in \text{Irr}(\text{Sp}_{2m})_\lambda$ ,  $\Theta_{O_{2m'+1}}^\epsilon(\gamma) = 0$  if  $m' < m$  and*

$$\text{Irr}(\Theta_{O_{2m'+1}}^\epsilon(\gamma)) \subset \text{Irr}(R_{O_{2n+1} \times T_{m-n} \times T_{m'-m}}^{O_{2m'+1}}(\lambda' \otimes \theta_{m-n} \otimes 1))$$

*otherwise.*

(ii) *For  $\gamma' \in \text{Irr}(O_{2m'+1}^\epsilon)_{\lambda', \theta}$ ,  $\Theta_{\text{Sp}_{2m}}^\epsilon(\gamma') = 0$  if  $m < m'$  and*

$$\text{Irr}(\Theta_{\text{Sp}_{2m}}^\epsilon(\gamma')) \subset \text{Irr}(R_{\text{Sp}_{2n} \times T_{m'-n} \times T_{m-m'}}^{\text{Sp}_{2m}}(\lambda \otimes 1 \otimes \theta_{m-m'}))$$

*otherwise.*

We end this paper by some remarks on the open problem of finding the explicit theta correspondence of Harish-Chandra series, which can be reformulated as a correspondence of Weyl group representations. Explicit descriptions have been proved for unitary dual pairs ([Aubert et al. 1996, Théorème 3.10]) and symplectic-even orthogonal dual pairs ([Aubert et al. 1996, Conjecture 3.11], proved recently in [Pan 2019a]). These results are also developed recently in [Epequin Chavez 2019] in order to pick up extremal components from the big theta lifting. We expect that a similar conjecture should hold for symplectic-odd orthogonal case as well according to the results in this section.

Let us give a few details. From Mackey theory we know that irreducible constituents in the Harish-Chandra series of a cuspidal unipotent representation are parametrized by irreducible representations of a Weyl group. This observation leads to a connection with the Springer correspondence [Aubert et al. 2016; Epequin Chavez 2019]. More precisely, let  $\lambda_k$  be the unique cuspidal unipotent

representation of a classical group  $G_{n_k}$ , where  $n_k$  is defined in Theorem 3.5. Namely, we have the table

$n_k$	type of $G$	sign of $G$
$\lfloor k(k+1)/4 \rfloor$	unitary	$\text{sgn}(-1)^{k(k+1)/2}$
$k(k+1)$	symplectic	NA
$k(k+1)$	odd orthogonal	$\pm$
$k^2$	even orthogonal	$\text{sgn}(-1)^k$

For  $G_{n_k} \subset G_m$  in the same Witt tower, the set  $\text{Irr}(G_m)_{\lambda_k}$  is parametrized by irreducible representations of the group

$$W_{G_m}(\lambda_k) = \{x \in N_{G_m}(L)/L \mid \lambda_k^x \cong \lambda_k\},$$

where  $L = G_{n_k} \times T_{m-n_k}$ . The uniqueness of  $\lambda_k$  implies that the defining condition of  $W_{G_m}(\lambda_k)$  is trivial, hence

$$W_{G_m}(\lambda_k) = N_{G_m}(L)/L \cong W_{m-n_k}$$

which is a Weyl group of type  $B_{m-n_k}$ .

By [Adams and Moy 1993], the theta correspondence takes  $\lambda_k$  to  $\lambda_{k'}$ , where  $k' = k$  for the symplectic-even orthogonal case and  $k' = k \pm 1$  for the unitary case. Hence in these cases the theta correspondence between Harish-Chandra series of cuspidal unipotent representations can be characterized as a correspondence between representations of a pair of Weyl groups  $(W_{m-n_k}, W_{m'-n_{k'}})$ . As we mentioned above, explicit correspondence of representations of such a pair  $(W_l, W_{l'})$  has been proved for unitary dual pairs and symplectic-even orthogonal dual pairs in [Aubert et al. 1996] and [Pan 2019a], respectively. The set  $\text{Irr}(W_l)$  has a well-known parametrization by bipartitions of  $l$  (see, e.g., [Geck and Pfeiffer 2000, Theorem 5.5.6]). J. Epequin Chavez [2019] further develops the above correspondence using bipartitions. Among other applications, an observation we make from the results in [Epequin Chavez 2019] is that in contrast to the  $p$ -adic case [Sun and Zhu 2015], the conservation relations do not hold in general for noncuspidal representations of finite dual pairs. It should be very interesting to understand the conservation type relations for finite dual pairs beyond the cuspidal case.

For the symplectic-odd orthogonal case, recall from Theorem 3.7 that  $\lambda_k$  is the unique cuspidal unipotent representation of  $\text{Sp}_{2n_k}$ , and  $\lambda_{k,\chi}^{'+}$  is the unique cuspidal  $\theta$ -representation of  $O_{2n_k+1}$  with trivial central character,  $n_k = k(k+1)$ . In a similar manner as in the unipotent case, the Harish-Chandra series  $\text{Irr}(\text{Sp}_{2m})_{\lambda_k,\theta}$  and  $\text{Irr}(O_{2m'+1})_{\lambda_{k,\chi}^{'+}}$  are in bijection with  $\text{Irr}(W_{m-n_k})$  and  $\text{Irr}(W_{m'-n_k})$ , respectively. By Theorem 4.2 (i) and (ii), we expect that a conjecture analogous to [Aubert et al. 1996, Conjecture 3.11] should hold for the theta correspondence between  $\text{Irr}(\text{Sp}_{2m})_{\lambda_k,\theta}$  and  $\text{Irr}(O_{2m'+1})_{\lambda_{k,\chi}^{'+}}$ .

## Acknowledgements

Liu would like to thank Jiajun Ma for many helpful and enlightening discussions on this topic. We thank the anonymous referee for many valuable comments.

## References

- [Adams and Moy 1993] J. Adams and A. Moy, “Unipotent representations and reductive dual pairs over finite fields”, *Trans. Amer. Math. Soc.* **340**:1 (1993), 309–321. MR Zbl
- [Aubert et al. 1996] A.-M. Aubert, J. Michel, and R. Rouquier, “Correspondance de Howe pour les groupes réductifs sur les corps finis”, *Duke Math. J.* **83**:2 (1996), 353–397. MR Zbl
- [Aubert et al. 2016] A.-M. Aubert, W. Kraškiewicz, and T. Przebinda, “Howe correspondence and Springer correspondence for dual pairs over a finite field”, pp. 17–44 in *Lie algebras, Lie superalgebras, vertex algebras and related topics*, edited by K. C. Misra et al., Proc. Sympos. Pure Math. **92**, Amer. Math. Soc., Providence, RI, 2016. MR Zbl
- [Carter 1985] R. W. Carter, *Finite groups of Lie type: conjugacy classes and complex characters*, Wiley, New York, 1985. MR Zbl
- [Deligne and Lusztig 1976] P. Deligne and G. Lusztig, “Representations of reductive groups over finite fields”, *Ann. of Math. (2)* **103**:1 (1976), 103–161. MR Zbl
- [Epequin Chavez 2019] J. Epequin Chavez, “Extremal unipotent representations for the finite Howe correspondence”, *J. Algebra* **535** (2019), 480–502. MR Zbl
- [Geck and Pfeiffer 2000] M. Geck and G. Pfeiffer, *Characters of finite Coxeter groups and Iwahori–Hecke algebras*, Lond. Math. Soc. Monogr. (N.S.) **21**, Oxford Univ. Press, 2000. MR Zbl
- [Gérardin 1977] P. Gérardin, “Weil representations associated to finite fields”, *J. Algebra* **46**:1 (1977), 54–101. MR Zbl
- [Gurevich and Howe 2017] S. Gurevich and R. Howe, “Small representations of finite classical groups”, pp. 209–234 in *Representation theory, number theory, and invariant theory* (New Haven, CT, 2015), edited by J. Cogdell et al., Progr. Math. **323**, Birkhäuser, Cham, 2017. MR Zbl
- [Howe 1973] R. Howe, “Invariant theory and duality for classical groups over finite fields with applications to their singular representation theory”, preprint, Yale Univ., 1973.
- [Lusztig 1977] G. Lusztig, “Irreducible representations of finite classical groups”, *Invent. Math.* **43**:2 (1977), 125–175. MR Zbl
- [Lusztig 1984] G. Lusztig, *Characters of reductive groups over a finite field*, Ann. of Math. Stud. **107**, Princeton Univ. Press, 1984. MR Zbl
- [Lusztig 1988] G. Lusztig, “On the representations of reductive groups with disconnected centre”, pp. 157–166 in *Orbites unipotentes et représentations, I: Groupes finis et algèbres de Hecke*, Astérisque **168**, Soc. Math. France, Paris, 1988. MR Zbl
- [Malle 2017] G. Malle, “Cuspidal characters and automorphisms”, *Adv. Math.* **320** (2017), 887–903. MR Zbl
- [Mœglin 1995] C. Mœglin, “Quadratic unipotent representations of  $p$ -adic groups”, pp. 245–262 in *Functional analysis on the eve of the 21st century, I* (New Brunswick, NJ, 1993), edited by S. Gindikin et al., Progr. Math. **131**, Birkhäuser, Boston, 1995. MR Zbl
- [Mœglin et al. 1987] C. Mœglin, M.-F. Vignéras, and J.-L. Waldspurger, *Correspondances de Howe sur un corps  $p$ -adique*, Lecture Notes in Math. **1291**, Springer, 1987. MR Zbl
- [Pan 2002] S.-Y. Pan, “Local theta correspondence of depth zero representations and theta dichotomy”, *J. Math. Soc. Japan* **54**:4 (2002), 793–845. MR Zbl

- [Pan 2016] S.-Y. Pan, “Weil representations of finite symplectic groups and finite odd-dimensional orthogonal groups”, *J. Algebra* **453** (2016), 291–324. MR Zbl
- [Pan 2019a] S.-Y. Pan, “Howe correspondence of unipotent characters for a finite symplectic/even-orthogonal dual pair”, preprint, 2019. arXiv
- [Pan 2019b] S.-Y. Pan, “Lusztig correspondence and Howe correspondence for finite reductive dual pairs”, preprint, 2019. arXiv
- [Srinivasan 1968] B. Srinivasan, “The characters of the finite symplectic group  $\mathrm{Sp}(4, q)$ ”, *Trans. Amer. Math. Soc.* **131** (1968), 488–525. MR Zbl
- [Srinivasan 1979a] B. Srinivasan, *Representations of finite Chevalley groups: a survey*, Lecture Notes in Math. **764**, Springer, 1979. MR Zbl
- [Srinivasan 1979b] B. Srinivasan, “Weil representations of finite classical groups”, *Invent. Math.* **51**:2 (1979), 143–153. MR Zbl
- [Sun and Zhu 2015] B. Sun and C.-B. Zhu, “Conservation relations for local theta correspondence”, *J. Amer. Math. Soc.* **28**:4 (2015), 939–983. MR Zbl

Received August 16, 2019. Revised December 13, 2019.

DONGWEN LIU  
SCHOOL OF MATHEMATICAL SCIENCES  
ZHEJIANG UNIVERSITY  
HANGZHOU  
CHINA  
malieu@zju.edu.cn

ZHICHENG WANG  
SCHOOL OF MATHEMATICAL SCIENCES  
ZHEJIANG UNIVERSITY  
HANGZHOU  
CHINA  
11735009@zju.edu.cn



# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Matthias Aschenbrenner  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[matthias@math.ucla.edu](mailto:matthias@math.ucla.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Wee Teck Gan  
Mathematics Department  
National University of Singapore  
Singapore 119076  
[matgwt@nus.edu.sg](mailto:matgwt@nus.edu.sg)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

---

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 306 No. 2 June 2020

---

Cohomological kernels of purely inseparable field extensions	385
ROBERTO ARAVIRE, BILL JACOB and MANUEL O'RYAN	
Kuperberg and Turaev–Viro invariants in unimodular categories	421
FRANCESCO COSTANTINO, NATHAN GEER, BERTRAND PATUREAU-MIRAND and VLADIMIR TURAEV	
A new equivalence between super Harish-Chandra pairs and Lie supergroups	451
FABIO GAVARINI	
Generalized Mullineux involution and perverse equivalences	487
THOMAS GERBER, NICOLAS JACON and EMILY NORTON	
Isotypic multiharmonic polynomials and Gelbart–Helgason reciprocity	519
ANTHONY C. KABLE	
Two applications of the integral regulator	539
MATT KERR and MUXI LI	
Definability and approximations in triangulated categories	557
ROSANNA LAKING and JORGE VITÓRIA	
Remarks on the theta correspondence over finite fields	587
DONGWEN LIU and ZHICHENG WANG	
On the configurations of centers of planar Hamiltonian Kolmogorov cubic polynomial differential systems	611
JAUME LLIBRE and DONGMEI XIAO	
2-categories of symmetric bimodules and their 2-representations	645
VOLODYMYR MAZORCHUK, VANESSA MIEMIETZ and XIAOTING ZHANG	
The homotopy groups of the $\eta$ -periodic motivic sphere spectrum	679
KYLE ORMSBY and OLIVER RÖNDIGS	
On the Noether Problem for torsion subgroups of tori	699
FEDERICO SCAVIA	
Explicit polynomial bounds on prime ideals in polynomial rings over fields	721
WILLIAM SIMMONS and HENRY TOWNSNER	
A new local gradient estimate for a nonlinear equation under integral curvature condition on manifolds	755
LIANG ZHAO and SHOUWEN FANG	



0030-8730(202006)306:2;1-J