

Pacific Journal of Mathematics

**EXISTENCE OF STEADY MULTIPLE VORTEX PATCHES TO
THE VORTEX-WAVE SYSTEM**

DAOMIN CAO AND GUODONG WANG

EXISTENCE OF STEADY MULTIPLE VORTEX PATCHES TO THE VORTEX-WAVE SYSTEM

DAOMIN CAO AND GUODONG WANG

We prove the existence of steady multiple vortex patch solutions to the vortex-wave system in a planar bounded domain. The construction is performed by solving a certain variational problem for the vorticity and studying the asymptotic behavior as the vorticity strength goes to infinity.

1. Introduction

The evolution of an incompressible inviscid fluid is described by the Euler equations

$$(1-1) \quad \begin{cases} \partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} = -\nabla P, \\ \nabla \cdot \mathbf{v} = 0, \end{cases}$$

where $\mathbf{v}$ is the velocity field and P is the pressure. In the planar case $\mathbf{v} = (v_1, v_2)$ and we introduce the scalar vorticity of the fluid as follows:

$$(1-2) \quad \omega := \partial_1 v_2 - \partial_2 v_1.$$

By taking the curl on both sides of the first equation of (1-1), we obtain the following vorticity form of the Euler equations:

$$(1-3) \quad \partial_t \omega + \mathbf{v} \cdot \nabla \omega = 0.$$

In the whole plane, the velocity can be recovered from the vorticity via the Biot–Savart law, that is,

$$(1-4) \quad \mathbf{v}(x, t) = K * \omega(x, t) := -\frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{J(x-y)}{|x-y|^2} \omega(y, t) dy,$$

where $J(a, b) := (b, -a)$ denotes clockwise rotation through $\pi/2$ of the planar vector $(a, b) \in \mathbb{R}^2$, and $K(x) = -(1/(2\pi))(Jx/|x|^2)$ is called the Biot–Savart kernel. Equations (1-3) and (1-4) mean that the vorticity ω is transported by a divergence-free velocity field induced by itself.

In some cases the vorticity is sharply concentrated in N small disjoint regions, where N is a positive integer, then its time evolution can be approximately described

MSC2010: 35J50, 35Q35.

Keywords: vortex-wave system, vortex patch, Euler equations, Kirchhoff–Routh function, variational problem.

by the point vortex model (see [Lin 1941] for example), an ODE system that can be written as

$$(1-5) \quad \frac{dx_i}{dt} = \sum_{j=1, j \neq i}^N \kappa_j K(x_i - x_j), \quad i = 1, \dots, N,$$

where x_i is the position of the i -th point vortex and κ_i is the corresponding vorticity strength. According to the point vortex model, for each point vortex located at x_i , the velocity it induces is $\kappa_i K(\cdot - x_i)$, and it moves in the velocity field induced by all the other $N - 1$ point vortices. The point vortex model and its relation with the Euler equations have been analyzed extensively; see [Cao et al. 2015; Marchioro and Papani 1986; Smets and Van Schaftingen 2010; Turkington 1987] for example.

Now it is natural to consider the mixed problem, that is, the vorticity consists of a continuously distributed part and a finite number of concentrated vortices. Marchioro and Pulvirenti [1991] first studied this problem and they called it the vortex-wave system. In the whole plane the vortex-wave system can be written as

$$(1-6) \quad \begin{cases} \partial_t \omega + \mathbf{v} \cdot \nabla \omega = 0, \\ \frac{dx_i}{dt} = K * \omega(x_i, t) + \sum_{j=1, j \neq i}^N \kappa_j K(x_i - x_j), \quad i = 1, \dots, N, \\ \mathbf{v} = K * \omega + \sum_{j=1}^k \kappa_j K(\cdot - x_j). \end{cases}$$

Throughout this paper we call ω the background vorticity. System (1-6) means that the background vorticity is transported by the velocity field induced by itself (the term $K * \omega$) and the N point vortices (the term $\sum_{j=1}^k \kappa_j K(\cdot - x_j)$), and each point vortex moves by the velocity induced by the background vorticity (the term $K * \omega(x_i, t)$) and all the other $N - 1$ point vortices (the term $\sum_{j \neq i, j=1}^N \kappa_j K(\cdot - x_j)$). By constructing Lagrangian paths Marchioro and Pulvirenti [1991] proved a theorem of existence for initial background vorticity belonging to $L^1(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$. More results on the existence and uniqueness can be found in [Bjorland 2011; Lacave and Miot 2009; Lopes Filho et al. 2011; Miot 2012].

In this paper, we will be focusing on the steady vortex-wave system in a bounded domain, the precise form of which will be given in the next section. In this case the Kirchhoff–Routh function (defined by (2-2) in the next section) plays an essential role. On the one hand, it is easy to see that any critical point of the Kirchhoff–Routh function is a stationary point of the vortex model, and it has been shown in [Cao et al. 2015] that if this critical point is nondegenerate, then there exists a family of steady vortex patch solutions (that is, the vorticity is a piecewise constant function) of the Euler equations shrinking to this critical point. Similar results can also be found in [Cao et al. 2014; Smets and Van Schaftingen 2010; Turkington 1983a; Wan 1988]. On the other hand, it is proved in [Cao et al. 2019a] that if there is a family of steady vortex patch solutions of the Euler equations shrinking to a point, then

this point must be in the interior of the domain and must be a critical point of the Kirchhoff–Routh function. In this way, the Kirchhoff–Routh function establishes connection between the Euler equations and the vortex model.

Our purpose in the present paper is to extend the result in [Cao et al. 2015] to the vortex-wave system. To be more precise, we prove that for any given strict local minimum point of the Kirchhoff–Routh function, there exists a family of steady vortex patch solutions to the vortex-wave system that shrinks to this point. Here by vortex patch solution of the vortex-wave system we mean that the background vorticity is a piecewise constant function.

The method we use in this paper to construct steady solutions is called the vorticity method, which was first established by Arnold [1978] and further developed by many authors [Burton 1989a; 1989b; Burton and McLeod 1991; Elcrat and Miller 1991; 1995; Turkington 1983a; 1983b]. Roughly speaking, the vorticity method is to maximize the kinetic energy of the fluid under some suitable constraints for the vorticity. For the Euler equations, the kinetic energy of the fluid with bounded vorticity is always finite, but for the vortex-wave system the kinetic energy is infinite due to the presence of point vortices. To overcome this difficulty, we drop the infinite self-energy term for each point vortex. We refer the interested reader to [Cao and Wang 2019] where the energy of the vortex-wave system with a single point vortex was calculated rigorously.

It is worth mentioning that the construction in [Cao et al. 2015] was based on a finite-dimensional reduction argument. The advantage of the method in [Cao et al. 2015] is that solutions concentrating at a given saddle point of the Kirchhoff–Routh function can be constructed. However, nondegeneracy of this saddle point is required in this situation. Using the vorticity method, we are able to construct solutions concentrating at a given strict local minimum point of the Kirchhoff–Routh function, even if the point is degenerate. Another advantage of the vorticity method is that we can analyze the energy of the solution, which is helpful to prove nonlinear stability; see [Burton 2005; Cao and Wang 2017] for example.

This paper is organized as follows. In Section 2, we give the formulation of the vortex-wave system in a bounded domain and state the main result. In Section 3, we solve a maximization problem for the vorticity and study the asymptotic behavior of the maximizers. In Section 4 we prove the main result.

2. Main result

Notation. Let $D \subset \mathbb{R}^2$ be a bounded and simply connected domain with smooth boundary. Green’s function for $-\Delta$ in D with zero Dirichlet data on ∂D can be written as

$$(2-1) \quad G(x, y) = \frac{1}{2\pi} \ln \frac{1}{|x - y|} - h(x, y), \quad x, y \in D,$$

where h is called the regular part of Green's function. Define

$$H(x) := \frac{1}{2}h(x, x),$$

which is usually called the Robin function. Let k be a positive integer, $\kappa_i \in \mathbb{R} \setminus \{0\}$, $i = 1, \dots, k$. Define the Kirchhoff–Routh function as

$$(2-2) \quad \mathcal{K}_k(x_1, \dots, x_k) = - \sum_{i \neq j, 1 \leq i, j \leq k} \kappa_i \kappa_j G(x_i, x_j) + \sum_{i=1}^k \kappa_i^2 h(x_i, x_i),$$

where $x_i \in D$ and $x_i \neq x_j$ if $i \neq j$. Note that if $k = 1$, then $\mathcal{K}_1 = 2\kappa_1^2 H$.

Throughout we will use the following notation. For any function g , $\text{supp}(g)$ denotes the support of g . For any real number a , $\text{sgn}(a)$ denotes the sign of a , that is,

$$(2-3) \quad \text{sgn}(a) := \begin{cases} 1 & \text{if } a > 0, \\ 0 & \text{if } a = 0, \\ -1 & \text{if } a < 0. \end{cases}$$

For any Lebesgue measurable set $A \subset \mathbb{R}^2$, $|A|$ denotes the two-dimensional Lebesgue measure of A ; I_A denotes the characteristic function of A , that is, $I_A(x) = 1$ if $x \in A$ and $I_A(x) = 0$ elsewhere; $\bar{A}$ denotes the closure of A in the Euclidean topology; and $\text{diam}(A)$ denotes the diameter of A , that is,

$$(2-4) \quad \text{diam}(A) = \sup_{x, y \in A} |x - y|.$$

Vortex-wave system in a bounded domain. We consider an incompressible steady flow confined in D with impermeability boundary condition. The evolution of the velocity field $\mathbf{v} = (v_1, v_2)$ and the pressure P is described by the Euler equations

$$(2-5) \quad \begin{cases} \partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} = -\nabla P & \text{in } D \times (0, +\infty), \\ \nabla \cdot \mathbf{v} = 0 & \text{in } D \times (0, +\infty), \\ \mathbf{v} \cdot \nu = 0 & \text{on } \partial D \times (0, +\infty), \end{cases}$$

where ν is the outward unit normal of ∂D .

Taking the curl on both sides of the first equation of (2-5), we get the vorticity equation

$$(2-6) \quad \partial_t \omega + \mathbf{v} \cdot \nabla \omega = 0.$$

Since $\mathbf{v}$ is divergence-free, there is a function ψ , called the stream function, such that

$$(2-7) \quad \mathbf{v} = J \nabla \psi = (\partial_2 \psi, -\partial_1 \psi).$$

By the definition of ω (recall (1-2)), it is easy to see that

$$(2-8) \quad -\Delta \psi = \omega.$$

The impermeability boundary condition given in the third equation of (2-5) implies that ψ is a constant on each connected component of ∂D . Since D is simply connected, after suitably adding a constant to ψ we can assume

$$(2-9) \quad \psi(x, t) = 0, \quad x \in \partial D.$$

By (2-8) and (2-9), we have

$$(2-10) \quad \psi(x, t) = (-\Delta)^{-1} \omega(x, t) := \int_D G(x, y) \omega(y, t) dy, \quad x \in D.$$

For brevity we introduce the notation

$$\partial(f, g) := \nabla f \cdot J \nabla g = \partial_1 f \partial_2 g - \partial_2 f \partial_1 g,$$

then the vorticity equation (2-6) can be written as

$$(2-11) \quad \begin{cases} \partial_t \omega + \partial(\omega, \psi) = 0, \\ \psi = (-\Delta)^{-1} \omega. \end{cases}$$

If the vorticity is a Dirac delta measure (also called a point vortex) located at $x \in D$, i.e., $\omega = \delta(x)$, then formally the velocity field it induces is

$$J \nabla (-\Delta)^{-1} \delta(x) = J \nabla G(x, \cdot) = \frac{1}{2\pi} \frac{J(x - \cdot)}{|x - \cdot|^2} - J \nabla h(x, \cdot).$$

Note that this velocity field is singular at x . Due to symmetry, we formally drop the term

$$\frac{1}{2\pi} \frac{J(x - \cdot)}{|x - \cdot|^2},$$

that is, we assume that the velocity at x is $-J \nabla h(x, \cdot)|_x = -J \nabla H(x)$, then the evolution of this point vortex is described by the following ODE:

$$(2-12) \quad \frac{dx}{dt} = -J \nabla H(x).$$

Similarly, the evolution of l point vortices can be described by the ODE system

$$(2-13) \quad \frac{dx_i}{dt} = -\kappa_i J \nabla H(x_i) + \sum_{j=1, j \neq i}^l \kappa_j J \nabla_{x_i} G(x_j, x_i), \quad i = 1, \dots, l,$$

where κ_i is the vorticity strength of the i -th point vortex. System (2-13) is also called the Kirchhoff–Routh equation. It is easy to see that the Kirchhoff–Routh function $\mathcal{K}_l$ is exactly the Hamiltonian of the system.

Now we consider the mixed problem. That is the vorticity consists of a continuously distributed part ω and l point vortices x_i with strength κ_i , $i = 1, \dots, l$. Then the evolution of ω and x_i will obey the equations

$$(2-14) \quad \begin{cases} \partial_t \omega + \partial(\omega, \psi + \sum_{j=1}^l \kappa_j G(x_j, \cdot)) = 0, \\ \frac{dx_i}{dt} = J \nabla(\psi + \sum_{j=1, j \neq i}^l \kappa_j G(x_j, \cdot) - \kappa_i H)(x_i), \quad i = 1, \dots, l, \\ \psi = (-\Delta)^{-1} \omega, \end{cases}$$

which together are called the vortex-wave system in D .

In this paper, we confine ourselves to the stationary case, that is, we consider the following system of equations:

$$(2-15) \quad \begin{cases} \partial(\omega, \psi + \sum_{j=1}^l \kappa_j G(x_j, \cdot)) = 0, \\ \nabla(\psi + \sum_{j=1, j \neq i}^l \kappa_j G(x_j, \cdot) - \kappa_i H)(x_i) = 0, \quad i = 1, \dots, l, \\ \psi = (-\Delta)^{-1} \omega. \end{cases}$$

Since we are going to deal with vortex patch solutions which are discontinuous, it is necessary to give the weak formulation of (2-15).

Definition 2.1. Let $\omega \in L^\infty(D)$, $x_i \in D$, $i = 1, \dots, l$, then $(\omega, x_1, \dots, x_l)$ is called a weak solution to (2-15) if it satisfies

$$(2-16) \quad \begin{cases} \int_D \omega(x) \partial(\psi(x) + \sum_{j=1}^l \kappa_j G(x_j, x), \phi(x)) dx = 0, \quad \forall \phi \in C_c^\infty(D), \\ \nabla(\psi(x) + \sum_{j=1, j \neq i}^l \kappa_j G(x_j, x) - \kappa_i H(x))|_{x=x_i} = 0, \quad i = 1, \dots, l, \end{cases}$$

where $\psi = (-\Delta)^{-1} \omega$.

Remark 2.2. Note that since $\omega \in L^\infty(D)$, by L^p estimate $\psi \in W^{2,p}(D)$ for any $1 < p < +\infty$, then by Sobolev embedding $\psi \in C^{1,\alpha}(\bar{D})$ for any $0 < \alpha < 1$.

Remark 2.3. Definition 2.1 can be derived formally from (2-15) by integration by parts; see [Cao and Wang 2019] for the detailed calculations.

Main result. Our main result in this paper is the following theorem:

Theorem 2.4. Let k, p, l be positive integers such that $p+l=k$, and κ_i , $i = 1, \dots, k$, be k real numbers such that $\kappa_i \neq 0$. Suppose that $(\bar{x}_1, \dots, \bar{x}_k)$ is a strict local minimum point of $\mathcal{K}_k$ defined by (2-2), where $\bar{x}_i \in D$ and $\bar{x}_i \neq \bar{x}_j$ for $i \neq j$. Then there exists $\lambda_0 > 0$ such that for $\lambda > \lambda_0$, (2-16) has a solution $(\omega^\lambda, x_{p+1}^\lambda, \dots, x_k^\lambda)$ satisfying

$$(2-17) \quad \omega^\lambda = \sum_{i=1}^p \omega_i^\lambda, \quad \int_D \omega_i^\lambda(x) dx = \kappa_i, \quad \omega_i^\lambda = \text{sgn}(\kappa_i) \lambda I_{A_i^\lambda}, \quad i = 1, \dots, p,$$

where A_i^λ has the form

$$(2-18) \quad A_i^\lambda = \left\{ x \in D \mid \operatorname{sgn}(\kappa_i) \left(\psi^\lambda(x) + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x) \right) > c_i^\lambda \right\} \cap B_\delta(\bar{x}_i)$$

for some $c_i^\lambda > 0$ and $\delta > 0$ (δ does not depend on λ). Moreover,

$$(2-19) \quad \operatorname{diam}(A_i^\lambda) \leq C\lambda^{-\frac{1}{2}}, \quad \lim_{\lambda \rightarrow +\infty} \left| \frac{1}{\kappa_i} \int_D x \omega_i^\lambda(x) dx - \bar{x}_i \right| = 0, \quad i = 1, \dots, p,$$

$$(2-20) \quad \lim_{\lambda \rightarrow +\infty} |x_j^\lambda - \bar{x}_j| = 0, \quad j = p+1, \dots, k,$$

where C is a positive number independent of λ .

Remark 2.5. By (2-19), we see that A_i^λ shrinks to $\bar{x}_i$ as $\lambda \rightarrow +\infty$, that is,

$$\lim_{\lambda \rightarrow +\infty} \sup_{x \in A_i^\lambda} |x - \bar{x}_i| = 0, \quad i = 1, \dots, p.$$

Consequently $\overline{A_i^\lambda} \subset B_\delta(\bar{x}_i)$ for sufficiently large λ .

3. Variational problem

Let $(\bar{x}_1, \dots, \bar{x}_k)$ be a strict local minimum point of $\mathcal{K}_k$, where $\bar{x}_i \in D$ and $\bar{x}_i \neq \bar{x}_j$ for $i \neq j$. Without loss of generality, we assume that $(\bar{x}_1, \dots, \bar{x}_k)$ is the unique minimum point of $\mathcal{K}_k$ on $\overline{B_{\delta_0}(\bar{x}_1)} \times \dots \times \overline{B_{\delta_0}(\bar{x}_k)}$, where $\delta_0 > 0$ is a small positive number such that $\overline{B_{\delta_0}(\bar{x}_i)} \subset D$ and $\overline{B_{\delta_0}(\bar{x}_i)} \cap \overline{B_{\delta_0}(\bar{x}_j)} = \emptyset$ for all $i, j = 1, \dots, k$ and $i \neq j$.

Remark 3.1. To our knowledge, there is no general result that guarantees the existence of a strict local minimum point of $\mathcal{K}_k$ for $k \geq 2$. Some special cases are as follows: if $k = 1$ and D is convex, by [Caffarelli and Friedman 1985] $\mathcal{K}_1$ is a strictly convex function in D , thus has a unique minimum point; if $k \geq 2$, $\kappa_i > 0$ for each i and D is convex, by [Grossi and Takahashi 2010] there does not exist any critical point of $\mathcal{K}_k$; if $k = 2$, some examples of strict local minimum points of $\mathcal{K}_2$ are given computationally in [Elcrat and Miller 1995]. More related results can also be found in [Bartsch and Pistoia 2015; Bartsch et al. 2010].

Let λ be a positive real number. Define

$$N_p^\lambda = \left\{ \omega \in L^\infty(D) \mid \omega = \sum_{i=1}^p \omega_i, \operatorname{supp}(\omega_i) \subset B_\delta(\bar{x}_i), \int_D \omega_i(x) dx = \kappa_i, 0 \leq \operatorname{sgn}(\kappa_i) \omega_i \leq \lambda \right\},$$

where $\delta < \frac{1}{2}\delta_0$ is a small positive number to be determined later. Hereafter we assume that λ is sufficiently large such that N_p^λ is not empty.

For $(\omega, x_{p+1}, \dots, x_k) \in N_p^\lambda \times \overline{B_\delta(\bar{x}_{p+1})} \times \dots \times \overline{B_\delta(\bar{x}_k)}$, define

$$\mathcal{E}(\omega, x_{p+1}, \dots, x_k)$$

$$:= E(\omega) + \sum_{j=p+1}^k \kappa_j \psi(x_j) + \frac{1}{2} \sum_{\substack{i \neq j \\ p+1 \leq i, j \leq k}} \kappa_i \kappa_j G(x_i, x_j) - \sum_{j=p+1}^k \kappa_j^2 H(x_j),$$

where

$$E(\omega) := \frac{1}{2} \int_D \int_D G(x, y) \omega(x) \omega(y) dx dy,$$

$$\psi(x) = (-\Delta)^{-1} \omega(x) := \int_D G(x, y) \omega(y) dy.$$

Let us explain the definition of $\mathcal{E}$ briefly. The first term $E(\omega)$ represents the self-interacting energy of the background vorticity ω , the second term represents the mutual interaction energy between the background vorticity and the l point vortices, the third term represents the total interaction energy between any two different point vortices, and the fourth term represents the interaction energy between the l point vortices and the boundary of D . As we have mentioned in [Section 1](#), we have dropped the infinite self-interacting energy for each point vortex.

Now we consider the maximization of $\mathcal{E}$ on $N_p^\lambda \times \overline{B_\delta(\bar{x}_{p+1})} \times \dots \times \overline{B_\delta(\bar{x}_k)}$.

Lemma 3.2. *For fixed λ , there exists*

$$(\omega^\lambda, x_{p+1}^\lambda, \dots, x_k^\lambda) \in N_p^\lambda \times \overline{B_\delta(\bar{x}_{p+1})} \times \dots \times \overline{B_\delta(\bar{x}_k)}$$

such that

$$(3-1) \quad \mathcal{E}(\omega^\lambda, x_{p+1}^\lambda, \dots, x_k^\lambda) = \sup_{(\omega, x_{p+1}, \dots, x_k) \in N_p^\lambda \times \overline{B_\delta(\bar{x}_{p+1})} \times \dots \times \overline{B_\delta(\bar{x}_k)}} \mathcal{E}(\omega, x_{p+1}, \dots, x_k).$$

Proof. First, for any $(\omega, x_{p+1}, \dots, x_k) \in N_p^\lambda \times \overline{B_\delta(\bar{x}_{p+1})} \times \dots \times \overline{B_\delta(\bar{x}_k)}$, since $G \in L^1(D \times D)$, $\text{dist}(B_{\delta_0}(\bar{x}_i), B_{\delta_0}(\bar{x}_j)) > 0$ for any $i \neq j$, $p+1 \leq i, j \leq k$, and $\text{dist}(B_{\delta_0}(\bar{x}_i), \partial D) > 0$, $p+1 \leq i \leq k$, we have

$$(3-2) \quad \mathcal{E}(\omega, x_{p+1}, \dots, x_k)$$

$$\begin{aligned} &= \frac{1}{2} \int_D \int_D G(x, y) \omega(x) \omega(y) dx dy + \sum_{j=p+1}^k \kappa_j \psi(x_j) \\ &\quad + \frac{1}{2} \sum_{\substack{i \neq j \\ p+1 \leq i, j \leq k}} \kappa_i \kappa_j G(x_i, x_j) - \sum_{j=p+1}^k \kappa_j^2 H(x_j) \\ &\leq \frac{\lambda^2}{2} \int_D \int_D |G(x, y)| dx dy + \sum_{j=p+1}^k |\kappa_j| \|\psi\|_{L^\infty(D)} + \sum_{j=p+1}^k \kappa_j^2 \|H\|_{L^\infty(B_\delta(\bar{x}_j))}. \end{aligned}$$

Thus

$$M := \sup_{(\omega, x_{p+1}, \dots, x_k) \in N_p^\lambda \times \overline{B_\delta(\bar{x}_{p+1})} \times \dots \times \overline{B_\delta(\bar{x}_k)}} \mathcal{E}(\omega, x_{p+1}, \dots, x_k) < +\infty.$$

Now we choose $(\omega^n, x_{p+1}^n, \dots, x_k^n) \in N_p^\lambda \times \overline{B_\delta(\bar{x}_{p+1})} \times \dots \times \overline{B_\delta(\bar{x}_k)}$ such that

$$\lim_{n \rightarrow +\infty} \mathcal{E}(\omega^n, x_{p+1}^n, \dots, x_k^n) = M.$$

Since N_p^λ is weakly star closed in $L^\infty(D)$ (for a detailed proof of this fact, see Theorem 2.1 in [Cao et al. 2019b]) and $\overline{B_\delta(\bar{x}_j)}$ is closed in the Euclidean topology for $j = p+1, \dots, k$, there exists $(\omega^\lambda, x_{p+1}^\lambda, \dots, x_k^\lambda) \in N_p^\lambda \times \overline{B_\delta(\bar{x}_{p+1})} \times \dots \times \overline{B_\delta(\bar{x}_k)}$ such that (up to a subsequence)

$$\begin{aligned} \omega^n &\rightarrow \omega^\lambda, & \text{weakly star in } L^\infty(D), \\ x_j^n &\rightarrow x_j^\lambda, & j = p+1, \dots, k. \end{aligned}$$

Then obviously

$$\mathcal{E}(\omega^\lambda, x_{p+1}^\lambda, \dots, x_k^\lambda) = \lim_{n \rightarrow +\infty} \mathcal{E}(\omega^n, x_{p+1}^n, \dots, x_k^n) = M,$$

which completes the proof. $\square$

Remark 3.3. It is easy to see that

$$(3-3) \quad E(\omega) + \sum_{j=p+1}^k \kappa_j \psi(x_j^\lambda) \leq E(\omega^\lambda) + \sum_{j=p+1}^k \kappa_j \psi^\lambda(x_j^\lambda)$$

for any $\omega \in N_p^\lambda$, and

$$\begin{aligned} (3-4) \quad & \sum_{j=p+1}^k \kappa_j \psi^\lambda(x_j) + \frac{1}{2} \sum_{\substack{i \neq j \\ p+1 \leq i, j \leq k}} \kappa_i \kappa_j G(x_i, x_j) - \sum_{j=p+1}^k \kappa_j^2 H(x_j) \\ & \leq \sum_{j=p+1}^k \kappa_j \psi^\lambda(x_j^\lambda) + \frac{1}{2} \sum_{\substack{i \neq j \\ p+1 \leq i, j \leq k}} \kappa_i \kappa_j G(x_i^\lambda, x_j^\lambda) - \sum_{j=p+1}^k \kappa_j^2 H(x_j^\lambda) \end{aligned}$$

for any $(x_{p+1}, \dots, x_k) \in \overline{B_\delta(\bar{x}_{p+1})} \times \dots \times \overline{B_\delta(\bar{x}_k)}$, where $\psi^\lambda = (-\Delta)^{-1} \omega^\lambda$ and $\psi = (-\Delta)^{-1} \omega$.

Since $\omega^\lambda \in N_p^\lambda$, we can write $\omega^\lambda = \sum_{i=1}^p \omega_i^\lambda$, where $\int_D \omega_i^\lambda(x) dx = \kappa_i$, $\text{supp}(\omega_i^\lambda) \subset B_\delta(\bar{x}_i)$ and $0 \leq \text{sgn}(\kappa_i) \omega_i^\lambda \leq \lambda$. The profile of each ω_i^λ is as follows.

Lemma 3.4. For $i = 1, \dots, p$, ω_i^λ has the form

$$\omega_i^\lambda = \operatorname{sgn}(\kappa_i) \lambda I_{A_i^\lambda},$$

where

$$A_i^\lambda := \left\{ x \in D \mid \operatorname{sgn}(\kappa_i) \left(\psi^\lambda(x) + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x) \right) > c_i^\lambda \right\} \cap B_\delta(\bar{x}_i)$$

for some $c_i^\lambda \in \mathbb{R}$ depending on λ and δ .

Proof. To make it clear, we divide the proof into two cases.

Case 1: $\kappa_i > 0$. For $s > 0$ we define a family of test functions $\omega_s^\lambda = \omega^\lambda + s(z_0 - z_1)$, where

$$(3-5) \quad \begin{cases} z_0, z_1 \in L^\infty(D), z_0, z_1 \geq 0, \int_D z_0(x) dx = \int_D z_1(x) dx, \\ \operatorname{supp}(z_0), \operatorname{supp}(z_1) \subset B_\delta(\bar{x}_i), \\ z_0 = 0 \text{ in } B_\delta(\bar{x}_i) \setminus \{\omega_i^\lambda \leq \lambda - \mu\}, \\ z_1 = 0 \text{ in } B_\delta(\bar{x}_i) \setminus \{\omega_i^\lambda \geq \mu\}, \end{cases}$$

where $\mu \in (0, \lambda)$. It is not hard to check that for fixed z_0, z_1 and μ , if s is sufficiently small (depending on z_0, z_1, μ), then $\omega_s^\lambda \in N_p^\lambda$. By (3-3) we have

$$\left. \frac{d}{ds} \left(E(\omega_s^\lambda) + \sum_{j=p+1}^k \kappa_j \psi_s^\lambda(x_j^\lambda) \right) \right|_{s=0^+} \leq 0,$$

where $\psi_s^\lambda = (-\Delta)^{-1} \omega_s^\lambda$. That is,

$$\int_D \left(\psi^\lambda(x) + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x) \right) z_0(x) dx \leq \int_D \left(\psi^\lambda(x) + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x) \right) z_1(x) dx.$$

Since z_0, z_1, μ are chosen arbitrarily as above we have

$$\sup_{\{\omega_i^\lambda < \lambda\} \cap B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right) \leq \inf_{\{\omega_i^\lambda > 0\} \cap B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right).$$

But $\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot)$ is a continuous function on $\overline{B_\delta(\bar{x}_i)}$ (notice that $x_j^\lambda \notin \overline{B_\delta(\bar{x}_i)}$ for $p+1 \leq j \leq k$), so we obtain

$$\sup_{\{\omega_i^\lambda < \lambda\} \cap B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right) = \inf_{\{\omega_i^\lambda > 0\} \cap B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right).$$

Define

$$(3-6) \quad \begin{aligned} c_i^\lambda &:= \sup_{\{\omega_i^\lambda < \lambda\} \cap B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right) \\ &= \inf_{\{\omega_i^\lambda > 0\} \cap B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right). \end{aligned}$$

It is easy to see that

$$(3-7) \quad \omega_i^\lambda \equiv \lambda \quad \text{a.e. on } \left\{ x \in D \mid \psi^\lambda(x) + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x) > c_i^\lambda \right\} \cap B_\delta(\bar{x}_i),$$

$$(3-8) \quad \omega_i^\lambda \equiv 0 \quad \text{a.e. on } \left\{ x \in D \mid \psi^\lambda(x) + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x) < c_i^\lambda \right\} \cap B_\delta(\bar{x}_i).$$

On the level set $\{x \in D \mid \psi^\lambda(x) + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x) = c_i^\lambda\} \cap B_\delta(\bar{x}_i)$, we have

$$(3-9) \quad \nabla(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot)) = 0, \quad \text{a.e.,}$$

which gives

$$(3-10) \quad \omega_i^\lambda = -\Delta(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot)) = 0, \quad \text{a.e..}$$

Equations (3-7), (3-8) and (3-10) together give

$$(3-11) \quad \omega_i^\lambda = \lambda I_{\{\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) > c_i^\lambda\} \cap B_\delta(\bar{x}_i)},$$

which completes the proof of Case 1.

Case 2: $\kappa_i < 0$. The argument is similar. For $s > 0$, define $\omega_s^\lambda = \omega^\lambda + s(z_1 - z_0)$, where

$$(3-12) \quad \begin{cases} z_0, z_1 \in L^\infty(D), z_0, z_1 \geq 0, \int_D z_0(x) dx = \int_D z_1(x) dx, \\ \text{supp}(z_0), \text{supp}(z_1) \subset B_\delta(\bar{x}_i), \\ z_0 = 0 \text{ in } B_\delta(\bar{x}_i) \setminus \{\omega_i^\lambda \geq \mu - \lambda\}, \\ z_1 = 0 \text{ in } B_\delta(\bar{x}_i) \setminus \{\omega_i^\lambda \leq -\mu\}, \end{cases}$$

where $\mu \in (0, \lambda)$. Then

$$\frac{d}{ds} \left(E(\omega_s^\lambda) + \sum_{j=p+1}^k \kappa_j \psi_s^\lambda(x_j^\lambda) \right) \Big|_{s=0^+} \leq 0.$$

Repeating the argument in Case 1, we obtain

$$\sup_{\{\omega_i^\lambda < 0\} \cap B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right) = \inf_{\{\omega_i^\lambda > -\lambda\} \cap B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right),$$

then we can define

$$\begin{aligned} (3-13) \quad c_i^\lambda &:= - \sup_{\{\omega_i^\lambda < 0\} \cap B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right) \\ &= - \inf_{\{\omega_i^\lambda > -\lambda\} \cap B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right). \end{aligned}$$

Similarly we have

$$\omega_i^\lambda = -\lambda I_{\{\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) < -c_i^\lambda\} \cap B_\delta(\bar{x}_i)}.$$

□

Lemma 3.5. *For the c_i^λ given in Lemma 3.4, we have the estimate*

$$c_i^\lambda > -\frac{|\kappa_i|}{2\pi} \ln \delta - C,$$

where $C > 0$ is independent of λ and δ .

Proof. We only prove the case $\kappa_i > 0$. The other case is similar. By the definition of c_i^λ (recall (3-6)), we have

$$\begin{aligned} (3-14) \quad c_i^\lambda &= \inf_{\{\omega_i^\lambda > 0\} \cap B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right) \\ &\geq \inf_{B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right) \quad (\text{by maximum principle}) \\ &\geq \inf_{\partial B_\delta(\bar{x}_i)} \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right) \\ &\geq \inf_{\partial B_\delta(\bar{x}_i)} \psi^\lambda - C \end{aligned}$$

for some $C > 0$ not depending on λ and δ . But for any $x \in \partial B_\delta(\bar{x}_i)$,

$$\begin{aligned} (3-15) \quad \psi^\lambda(x) &= -\frac{1}{2\pi} \int_D \ln |x - y| \omega_i^\lambda(y) dy - \int_D h(x, y) \omega_i^\lambda(y) dy \\ &\quad + \int_D G(x, y) \sum_{j=1, j \neq i}^p \omega_j^\lambda(y) dy \\ &\geq -\frac{\kappa_i}{2\pi} \ln |2\delta| - C \end{aligned}$$

for some $C > 0$ not depending on λ and δ . Here we use the fact that

$$\text{dist}(B_{\delta_0}(\bar{x}_i), B_{\delta_0}(\bar{x}_j)) > 0$$

for any $i \neq j$, $1 \leq i, j \leq p$, and $\text{dist}(B_{\delta_0}(\bar{x}_i), \partial D) > 0$, $1 \leq i \leq p$.

Combining (3-14) with (3-15) we get the desired result. $\square$

Lemma 3.6. *For δ sufficiently small, not depending on λ , the following assertion holds true:*

$$(3-16) \quad \text{sgn}(\kappa_i) \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right) - c_i^\lambda \leq 0 \quad \text{on } \partial B_{\delta_0}(\bar{x}_i)$$

for each $1 \leq i \leq p$.

Proof. Let $1 \leq i \leq p$ be fixed. By Lemma 3.5 it suffices to show that

$$\sup_{x \in \partial B_{\delta_0}(\bar{x}_i)} \left| \psi^\lambda(x) + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x) \right| \leq C$$

for some $C > 0$ not depending on λ and δ . In fact, since for any $j \neq i$, $j = p+1, \dots, k$, $\text{dist}(B_{\delta_0}(\bar{x}_i), B_{\delta_0}(\bar{x}_j)) > 0$, we have

$$\sup_{x \in \partial B_{\delta_0}(\bar{x}_i)} \left| \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x) \right| \leq C.$$

It remains to show that $\sup_{x \in \partial B_{\delta_0}(\bar{x}_i)} |\psi^\lambda(x)| \leq C$. For any $x \in \partial B_{\delta_0}(\bar{x}_i)$, we estimate $\psi^\lambda(x)$ as follows:

$$\begin{aligned} |\psi^\lambda(x)| &\leq \left| \int_D -\frac{1}{2\pi} \ln|x-y| \omega_i^\lambda(y) dy \right| + \left| \int_D h(x, y) \omega_i^\lambda(y) dy \right| \\ &\quad + \left| \int_D G(x, y) \sum_{j \neq i, j=1}^p \omega_j^\lambda(y) dy \right| \\ &\leq -\frac{|\kappa_i|}{2\pi} \ln(\delta_0 - \delta) + C \\ &\leq -\frac{|\kappa_i|}{2\pi} \ln\left(\frac{\delta_0}{2}\right) + C \quad \left(\text{recall that } \delta < \frac{\delta_0}{2} \right). \end{aligned}$$

Thus the proof is completed. $\square$

From now on we fix δ such that (3-16) holds true.

Now we analyze the asymptotic behavior of the maximizer $(\omega^\lambda, x_{p+1}^\lambda, \dots, x_k^\lambda)$ as $\lambda \rightarrow +\infty$. We will show that as $\lambda \rightarrow +\infty$, the support of ω_i^λ shrinks to $\bar{x}_i$ for

$i = 1, \dots, p$ and $x_j^\lambda \rightarrow \bar{x}_j$ for $j = p+1, \dots, k$. To achieve this goal, we estimate the energy of the background vorticity first.

For simplicity, hereafter we will use C to denote various positive numbers independent of λ , and $o(1)$ to denote various quantities that go to zero as $\lambda \rightarrow +\infty$.

Lemma 3.7. $E(\omega^\lambda) \geq -\frac{1}{4\pi} \sum_{i=1}^p \kappa_i^2 \ln \varepsilon_i - C$, where ε_i is the positive number such that $\lambda \pi \varepsilon_i^2 = |\kappa_i|$.

Proof. Choose the test function $\omega = \sum_{i=1}^p \omega_i$, where $\omega_i = \text{sgn}(\kappa_i) \lambda I_{B_{\varepsilon_i}(\bar{x}_i)}$. It is obvious that $\omega \in N_p^\lambda$, so by (3-3),

$$E(\omega) + \sum_{j=p+1}^k \kappa_j \psi(x_j^\lambda) \leq E(\omega^\lambda) + \sum_{j=p+1}^k \kappa_j \psi^\lambda(x_j^\lambda),$$

where $\psi = (-\Delta)^{-1} \omega$. It is easy to check that

$$\left| \sum_{j=p+1}^k \kappa_j \psi(x_j^\lambda) \right| \leq C, \quad \left| \sum_{j=p+1}^k \kappa_j \psi^\lambda(x_j^\lambda) \right| \leq C,$$

so

$$(3-17) \quad E(\omega^\lambda) \geq E(\omega) - C.$$

On the other hand,

$$\begin{aligned} (3-18) \quad E(\omega) &= \frac{1}{2} \int_D \int_D G(x, y) \omega(x) \omega(y) dx dy \\ &= \sum_{i=1}^p E(\omega_i) + \sum_{1 \leq i < j \leq p} \int_D \int_D G(x, y) \omega_i(x) \omega_j(y) dx dy \\ &\geq \sum_{i=1}^p E(\omega_i) - C. \end{aligned}$$

Each $E(\omega_i)$ can be calculated directly as follows

$$\begin{aligned} (3-19) \quad E(\omega_i) &= \frac{1}{2} \int_D \int_D G(x, y) \omega_i(x) \omega_i(y) dx dy \\ &= \frac{1}{2} \int_{B_{\varepsilon_i}(\bar{x}_i)} \int_{B_{\varepsilon_i}(\bar{x}_i)} -\frac{1}{2\pi} \ln |x - y| \omega_i(x) \omega_i(y) dx dy \\ &\quad - \frac{1}{2} \int_{B_{\varepsilon_i}(\bar{x}_i)} \int_{B_{\varepsilon_i}(\bar{x}_i)} h(x, y) \omega_i(x) \omega_i(y) dx dy \\ &\geq -\frac{\kappa_i^2}{4\pi} \ln(2\varepsilon_i) - \sum_{i=1}^p \kappa_i^2 H(\bar{x}_i) + o(1). \end{aligned}$$

Now (3-17), (3-18) and (3-19) together give the desired result. $\square$

Now define

$$(3-20) \quad T^\lambda := \sum_{i=1}^p \frac{\operatorname{sgn}(\kappa_i)}{2} \int_D \omega_i^\lambda(x) \left(\operatorname{sgn}(\kappa_i)(\psi^\lambda(x) + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x)) - c_i^\lambda \right) dx,$$

which represents the total kinetic energy of the fluid on the vorticity set $\bigcup_{i=1}^p A_i^\lambda$.

Lemma 3.8. $T^\lambda \leq C$.

Proof. For simplicity we denote

$$\zeta_i^\lambda := \operatorname{sgn}(\kappa_i) \left(\psi^\lambda + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, \cdot) \right) - c_i^\lambda.$$

It suffices to prove that for each $1 \leq i \leq p$,

$$\operatorname{sgn}(\kappa_i) \int_D \omega_i^\lambda \zeta_i^\lambda dx \leq C.$$

On the one hand, by Hölder's inequality and Sobolev embedding $W^{1,1}(B_\delta(\bar{x}_i)) \hookrightarrow L^2(B_\delta(\bar{x}_i))$, we have

$$\begin{aligned} (3-21) \quad & \operatorname{sgn}(\kappa_i) \int_D \omega_i^\lambda \zeta_i^\lambda dx = \lambda \int_{A_i^\lambda} \zeta_i^\lambda dx \\ & \leq \lambda |A_i^\lambda|^{\frac{1}{2}} \left(\int_{A_i^\lambda} |\zeta_i^\lambda|^2 dx \right)^{\frac{1}{2}} \quad (\zeta_i^\lambda \geq 0 \text{ on } A_i^\lambda) \\ & \leq \lambda |A_i^\lambda|^{\frac{1}{2}} \left(\int_{B_\delta(\bar{x}_i)} |(\zeta_i^\lambda)^+|^2 dx \right)^{\frac{1}{2}} \\ & \leq C \lambda |A_i^\lambda|^{\frac{1}{2}} \left(\int_{B_\delta(\bar{x}_i)} (\zeta_i^\lambda)^+ dx + \int_{B_\delta(\bar{x}_i)} |\nabla(\zeta_i^\lambda)^+| dx \right) \\ & \leq C \lambda |A_i^\lambda|^{\frac{1}{2}} \left(\int_{A_i^\lambda} \zeta_i^\lambda dx + \int_{A_i^\lambda} |\nabla \zeta_i^\lambda| dx \right) \\ & \leq C \lambda |A_i^\lambda|^{\frac{1}{2}} |A_i^\lambda|^{\frac{1}{2}} \left(\int_{A_i^\lambda} |\nabla \zeta_i^\lambda|^2 dx \right)^{\frac{1}{2}} + C |A_i^\lambda|^{\frac{1}{2}} \operatorname{sgn}(\kappa_i) \int_D \omega_i^\lambda \zeta_i^\lambda dx \\ & \leq C \left(\int_{A_i^\lambda} |\nabla \zeta_i^\lambda|^2 dx \right)^{\frac{1}{2}} + o(1) \operatorname{sgn}(\kappa_i) \int_D \omega_i^\lambda \zeta_i^\lambda dx, \end{aligned}$$

which implies

$$(3-22) \quad \operatorname{sgn}(\kappa_i) \int_D \omega_i^\lambda \zeta_i^\lambda dx \leq C \left(\int_{A_i^\lambda} |\nabla \zeta_i^\lambda|^2 dx \right)^{\frac{1}{2}}.$$

On the other hand, since $\zeta_i^\lambda \leq 0$ on $\partial B_{\delta_0}(\bar{x}_i)$ (recall (3-16)), integration by parts gives

$$(3-23) \quad \begin{aligned} \operatorname{sgn}(\kappa_i) \int_D \omega_i^\lambda \zeta_i^\lambda dx &= \operatorname{sgn}(\kappa_i) \int_{B_{\delta_0}(\bar{x}_i)} \omega_i^\lambda (\zeta_i^\lambda)^+ dx \\ &= \int_{B_{\delta_0}(\bar{x}_i)} |\nabla(\zeta_i^\lambda)^+|^2 dx \geq \int_{A_i^\lambda} |\nabla \zeta_i^\lambda|^2 dx. \end{aligned}$$

Combining (3-22) with (3-23) we complete the proof. $\square$

Lemma 3.9. $\sum_{i=1}^p c_i^\lambda |\kappa_i| \geq -\frac{1}{2\pi} \sum_{i=1}^p \kappa_i^2 \ln \varepsilon_i - C.$

Proof. By the definition of T^λ , the following identity holds true:

$$T^\lambda = E(\omega^\lambda) + \frac{1}{2} \sum_{i=1}^p \sum_{j=p+1}^k \int_D \omega_i^\lambda(x) \kappa_j G(x_j^\lambda, x) dx - \frac{1}{2} \sum_{i=1}^p c_i^\lambda |\kappa_i|.$$

Since $\operatorname{dist}(B_{\delta_0}(\bar{x}_i), B_{\delta_0}(\bar{x}_j)) > 0$ for any $i = 1, \dots, p$ and $j = p+1, \dots, k$, it is easy to see that

$$\left| \frac{1}{2} \sum_{i=1}^p \sum_{j=p+1}^k \int_D \omega_i^\lambda(x) \kappa_j G(x_j^\lambda, x) dx \right| \leq C,$$

then the desired result follows from Lemmas 3.7 and 3.8. $\square$

Now we are ready to estimate the size of $\operatorname{supp}(\omega_i^\lambda)$.

Lemma 3.10. *There exists $R_0 > 0$ independent of λ such that for $i = 1, \dots, p$,*

$$\operatorname{diam}(\operatorname{supp}(\omega_i^\lambda)) \leq R_0 \varepsilon_i.$$

Proof. For each $i = 1, \dots, p$, any $x_i \in \operatorname{supp}(\omega_i^\lambda)$, we have

$$\operatorname{sgn}(\kappa_i) \left(\psi^\lambda(x_i) + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x_i) \right) \geq c_i^\lambda.$$

It is easy to see that

$$\left| \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x_i) \right| \leq C,$$

so we have

$$\operatorname{sgn}(\kappa_i) \psi^\lambda(x_i) \geq c_i^\lambda - C,$$

which gives

$$(3-24) \quad \int_D -\frac{1}{2\pi} \ln |x_i - y| |\omega_i^\lambda(y)| dy \geq c_i^\lambda - C.$$

Combining (3-24) with Lemma 3.9 we obtain

$$\sum_{i=1}^p |\kappa_i| \int_D -\frac{1}{2\pi} \ln |x_i - y| |\omega_i^\lambda(y)| dy \geq -\frac{1}{2\pi} \sum_{i=1}^p \kappa_i^2 \ln \varepsilon_i - C,$$

or equivalently

$$\sum_{i=1}^p \frac{|\kappa_i|}{2\pi} \int_D \ln \frac{\varepsilon_i}{|x_i - y|} |\omega_i^\lambda(y)| dy \geq -C.$$

For any $R > 1$ to be determined, we have

$$(3-25) \quad \sum_{i=1}^p \frac{|\kappa_i|}{2\pi} \int_{D \setminus B_{R\varepsilon_i}(x_i)} \ln \frac{\varepsilon_i}{|x_i - y|} |\omega_i^\lambda(y)| dy \\ + \sum_{i=1}^p \frac{|\kappa_i|}{2\pi} \int_{B_{R\varepsilon_i}(x_i)} \ln \frac{\varepsilon_i}{|x_i - y|} |\omega_i^\lambda(y)| dy \geq -C.$$

The second integral in (3-25) is bounded (in fact, it can be calculated explicitly). That is,

$$\left| \sum_{i=1}^p \frac{|\kappa_i|}{2\pi} \int_{B_{R\varepsilon_i}(x_i)} \ln \frac{\varepsilon_i}{|x_i - y|} |\omega_i^\lambda(y)| dy \right| \leq C,$$

from which we deduce

$$(3-26) \quad \sum_{i=1}^p \frac{|\kappa_i|}{2\pi} \int_{D \setminus B_{R\varepsilon_i}(x_i)} \ln \frac{\varepsilon_i}{|x_i - y|} |\omega_i^\lambda(y)| dy \geq -C.$$

Therefore

$$(3-27) \quad \sum_{i=1}^p \frac{|\kappa_i|}{2\pi} \int_{D \setminus B_{R\varepsilon_i}(x_i)} |\omega_i^\lambda(y)| dy \leq \frac{C}{\ln R},$$

and consequently for each $1 \leq i \leq p$ there holds

$$(3-28) \quad \frac{|\kappa_i|}{2\pi} \int_{D \setminus B_{R\varepsilon_i}(x_i)} |\omega_i^\lambda(y)| dy \leq \frac{C}{\ln R}.$$

Since $\int_D |\omega_i^\lambda(y)| dy = |\kappa_i|$, we can choose R large enough such that

$$(3-29) \quad \int_{B_{R\varepsilon_i}(x_i)} |\omega_i^\lambda(y)| dy > \frac{|\kappa_i|}{2}, \quad i = 1, \dots, p.$$

Now we claim that

$$\text{diam}(\text{supp}(\omega_i^\lambda)) \leq 2R\varepsilon_i.$$

In fact, suppose that $\text{diam}(\text{supp}(\omega_i^\lambda)) > 2R\varepsilon_i$. Then we can choose $x_i, y_i \in \text{supp}(\omega_i^\lambda)$ such that $|x_i - y_i| > 2R\varepsilon_i$, then by (3-29) (recall that in (3-29) $x_i \in \text{supp}(\omega_i^\lambda)$ is arbitrary)

$$\int_D |\omega_i^\lambda(y)| dy \geq \int_{B_{R\varepsilon_i}(x_i)} |\omega_i^\lambda(y)| dy + \int_{B_{R\varepsilon_i}(y_i)} |\omega_i^\lambda(y)| dy > |\kappa_i|,$$

which is a contradiction.

Finally by choosing $R_0 = 2R$ we complete the proof. $\square$

By now we have constructed $\omega_i^\lambda \in N_p^\lambda, i = 1, \dots, p$, and $x_j^\lambda \in \overline{B_\delta(\bar{x}_j)}, j = p+1, \dots, k$. Moreover, we have shown that the diameter of $\text{supp}(\omega_i^\lambda)$ vanishes as $\lambda \rightarrow +\infty$. To analyze their limiting positions, we define the center of ω_i^λ as

$$z_i^\lambda := \frac{1}{\kappa_i} \int_D x \omega_i^\lambda dx, \quad i = 1, \dots, p.$$

It is easy to see that $z_i^\lambda \in \overline{B_\delta(\bar{x}_i)}$.

Lemma 3.11. $z_i^\lambda \rightarrow \bar{x}_i$ for $1 \leq i \leq p$ and $x_j^\lambda \rightarrow \bar{x}_j$ for $p+1 \leq j \leq k$ as $\lambda \rightarrow +\infty$.

Proof. Up to a subsequence we assume that $z_i^\lambda \rightarrow z_i \in \overline{B_\delta(\bar{x}_i)}, 1 \leq i \leq p$, and $x_j^\lambda \rightarrow z_j \in \overline{B_\delta(\bar{x}_j)}, p+1 \leq j \leq k$. It suffices to show that $(z_1, \dots, z_k) = (\bar{x}_1, \dots, \bar{x}_k)$.

Define $\omega = \sum_{i=1}^p \omega_i$, where $\omega_i = \text{sgn}(\kappa_i) \lambda I_{B_{\varepsilon_i}(\bar{x}_i)}$. It is easy to see that $\omega \in N_p^\lambda$, so we have

$$\mathcal{E}(\omega, \bar{x}_{p+1}, \dots, \bar{x}_k) \leq \mathcal{E}(\omega^\lambda, x_{p+1}^\lambda, \dots, x_k^\lambda),$$

that is,

$$\begin{aligned} (3-30) \quad E(\omega) + \sum_{j=p+1}^k \kappa_j \psi(\bar{x}_j) + \frac{1}{2} \sum_{\substack{i \neq j \\ p+1 \leq i, j \leq k}} \kappa_i \kappa_j G(\bar{x}_i, \bar{x}_j) - \sum_{j=p+1}^k \kappa_j^2 H(\bar{x}_j) \\ \leq E(\omega^\lambda) + \sum_{j=p+1}^k \kappa_j \psi^\lambda(x_j^\lambda) + \frac{1}{2} \sum_{\substack{i \neq j \\ p+1 \leq i, j \leq k}} \kappa_i \kappa_j G(x_i^\lambda, x_j^\lambda) - \sum_{j=p+1}^k \kappa_j^2 H(x_j^\lambda), \end{aligned}$$

where $\psi^\lambda = (-\Delta)^{-1} \omega^\lambda$ and $\psi = (-\Delta)^{-1} \omega$. It is easy to check that

$$\begin{aligned} (3-31) \quad E(\omega) = \frac{1}{2} \sum_{i=1}^p \int_D \int_D -\frac{1}{2\pi} \ln|x-y| \omega_i(x) \omega_i(y) dx dy - \sum_{i=1}^p \kappa_i^2 H(\bar{x}_i) \\ + \frac{1}{2} \sum_{\substack{i \neq j \\ 1 \leq i, j \leq p}} \kappa_i \kappa_j G(\bar{x}_i, \bar{x}_j) + o(1), \end{aligned}$$

and

$$(3-32) \quad \sum_{j=p+1}^k \kappa_j \psi(\bar{x}_j) = \sum_{i=1}^p \sum_{j=p+1}^k \kappa_i \kappa_j G(\bar{x}_i, \bar{x}_j) + o(1).$$

Similarly

$$(3-33) \quad E(\omega^\lambda) = \frac{1}{2} \sum_{i=1}^p \int_D \int_D -\frac{1}{2\pi} \ln |x-y| \omega_i^\lambda(x) \omega_i^\lambda(y) dx dy - \sum_{i=1}^p \kappa_i^2 H(z_i) \\ + \frac{1}{2} \sum_{\substack{i \neq j \\ 1 \leq i, j \leq p}} \kappa_i \kappa_j G(z_i, z_j) + o(1),$$

and

$$(3-34) \quad \sum_{j=p+1}^k \kappa_j \psi^\lambda(x_j^\lambda) = \sum_{i=1}^p \sum_{j=p+1}^k \kappa_i \kappa_j G(z_i, z_j) + o(1).$$

Hence from all the above we obtain

$$(3-35) \quad \frac{1}{2} \sum_{i=1}^p \int_D \int_D -\frac{1}{2\pi} \ln |x-y| \omega_i(x) \omega_i(y) dx dy - \sum_{i=1}^p \kappa_i^2 H(\bar{x}_i) \\ + \frac{1}{2} \sum_{\substack{i \neq j \\ 1 \leq i, j \leq p}} \kappa_i \kappa_j G(\bar{x}_i, \bar{x}_j) + \sum_{i=1}^p \sum_{j=p+1}^k \kappa_i \kappa_j G(\bar{x}_i, \bar{x}_j) \\ + \frac{1}{2} \sum_{\substack{i \neq j \\ p+1 \leq i, j \leq k}} \kappa_i \kappa_j G(\bar{x}_i, \bar{x}_j) - \sum_{j=p+1}^k \kappa_j^2 H(\bar{x}_j) \\ \leq \frac{1}{2} \sum_{i=1}^p \int_D \int_D -\frac{1}{2\pi} \ln |x-y| \omega_i^\lambda(x) \omega_i^\lambda(y) dx dy - \sum_{i=1}^p \kappa_i^2 H(z_i) \\ + \frac{1}{2} \sum_{\substack{i \neq j \\ 1 \leq i, j \leq p}} \kappa_i \kappa_j G(z_i, z_j) + \sum_{i=1}^p \sum_{j=p+1}^k \kappa_i \kappa_j G(z_i, z_j) \\ + \frac{1}{2} \sum_{\substack{i \neq j \\ p+1 \leq i, j \leq k}} \kappa_i \kappa_j G(z_i, z_j) - \sum_{j=p+1}^k \kappa_j^2 H(z_j) + o(1).$$

On the other hand, by Riesz's rearrangement inequality we have for each $1 \leq i \leq p$

$$(3-36) \quad \int_D \int_D -\frac{1}{2\pi} \ln |x-y| \omega_i^\lambda(x) \omega_i^\lambda(y) dx dy \\ \leq \int_D \int_D -\frac{1}{2\pi} \ln |x-y| \omega_i(x) \omega_i(y) dx dy.$$

Therefore by (3-35) and (3-36) we get

$$(3-37) \quad \frac{1}{2} \sum_{\substack{i \neq j \\ 1 \leq i, j \leq k}} \kappa_i \kappa_j G(\bar{x}_i, \bar{x}_j) - \sum_{j=1}^k \kappa_j^2 H(\bar{x}_j) \\ \leq \frac{1}{2} \sum_{\substack{i \neq j \\ 1 \leq i, j \leq k}} \kappa_i \kappa_j G(z_i, z_j) - \sum_{j=1}^k \kappa_j^2 H(z_j),$$

or equivalently

$$(3-38) \quad \mathcal{K}_k(z_1, \dots, z_k) \leq \mathcal{K}_k(\bar{x}_1, \dots, \bar{x}_k).$$

Since $(\bar{x}_1, \dots, \bar{x}_k)$ is the unique minimum point for $\mathcal{K}_k$ on $\overline{B_\delta(\bar{x}_1)} \times \dots \times \overline{B_\delta(\bar{x}_k)}$, we obtain

$$(z_1, \dots, z_k) = (\bar{x}_1, \dots, \bar{x}_k),$$

which completes the proof. $\square$

Remark 3.12. It is easy to see that $x_j^\lambda \in B_\delta(\bar{x}_j)$ for $j = p+1, \dots, k$, and $\text{dist}(\text{supp}(\omega_i^\lambda), \partial B_\delta(\bar{x}_i)) > 0$ for $i = 1, \dots, p$, provided that λ is sufficiently large.

4. Proof of Theorem 2.4

Having made all the preparations in the preceding sections, we are now ready to give the proof of Theorem 2.4.

Proof of Theorem 2.4. We only need to prove that $(\omega^\lambda, x_{p+1}^\lambda, \dots, x_k^\lambda)$ satisfies (2-16) if λ is sufficiently large, since the other assertions have been verified in Section 3.

First, by Lemma 3.11, $x_j^\lambda \in B_\delta(\bar{x}_j)$ for $j = p+1, \dots, k$, thus by (3-4) we have

$$\nabla \left(\psi^\lambda(x) + \sum_{j=p+1, j \neq i}^k \kappa_j G(x_j^\lambda, x) - \kappa_i H(x) \right) \Big|_{x=x_i^\lambda} = 0, \quad j = p+1, \dots, k.$$

Now for any given $\phi \in C_c^\infty(D)$, define a family of transformations $\Phi_t(x)$, $t \in \mathbb{R}$, from D to D by the following ordinary differential equation:

$$(4-1) \quad \begin{cases} \frac{d\Phi_t(x)}{dt} = J\nabla\phi(\Phi_t(x)), & t \in \mathbb{R}, \\ \Phi_0(x) = x. \end{cases}$$

Since $J\nabla\phi$ is a smooth vector field with compact support in D , (4-1) is solvable for all $t \in \mathbb{R}$. It is also easy to see that $J\nabla\phi$ is divergence-free, so by the Liouville theorem (see [Marchioro and Pulvirenti 1994, Appendix 1.1]) $\Phi_t(x)$ is area-preserving, or equivalently, for any measurable set $A \subset D$

$$(4-2) \quad |\Phi_t(A)| = |A| \quad \text{for all } t \in \mathbb{R}.$$

Now we define a family of test functions

$$(4-3) \quad \omega_t^\lambda(x) := \omega^\lambda(\Phi_{-t}(x)).$$

Since Φ_t is area-preserving and $\text{dist}(\text{supp}(\omega_i^\lambda), \partial B_\delta(\bar{x}_i)) > 0$ for each $1 \leq i \leq p$, we have $\omega_i^\lambda \in N_p^\lambda$ as long as $|t|$ is sufficiently small. Then by (3-3) we have

$$\left. \frac{d}{dt} \left(E(\omega_t^\lambda) + \sum_{j=p+1}^k \kappa_j \psi_t^\lambda(x_j^\lambda) \right) \right|_{t=0} = 0,$$

where $\psi_t^\lambda = (-\Delta)^{-1} \omega_t^\lambda$. It is easy to check that (see [Turkington 1983a] for example)

$$E(\omega_t^\lambda) = E(\omega^\lambda) + t \int_D \omega^\lambda(x) \partial(\psi^\lambda(x), \phi(x)) dx + o(t),$$

and

$$\psi_t^\lambda(x_j^\lambda) = \psi^\lambda(x_j^\lambda) + t \int_D \omega^\lambda(x) \partial(G(x_j^\lambda, x), \phi(x)) dx + o(t),$$

where $o(t)/t \rightarrow 0$ as $t \rightarrow 0$. Therefore we get

$$\int_D \omega^\lambda(x) \partial \left(\psi^\lambda(x) + \sum_{j=p+1}^k \kappa_j G(x_j^\lambda, x), \phi(x) \right) dx = 0,$$

which completes the proof of Theorem 2.4. □

Acknowledgements

Daomin Cao was supported by NNSF of China (No. 11831009) and Chinese Academy of Sciences (No. QYZDJ-SSW-SYS021). Guodong Wang was supported by China Postdoctoral Science Foundation (No. 2019M661261).

References

- [Arnold 1978] V. I. Arnold, *Mathematical methods of classical mechanics*, Springer, 1978. [MR](#) [Zbl](#)
- [Bartsch and Pistoia 2015] T. Bartsch and A. Pistoia, “Critical points of the N -vortex Hamiltonian in bounded planar domains and steady state solutions of the incompressible Euler equations”, *SIAM J. Appl. Math.* **75**:2 (2015), 726–744. [MR](#) [Zbl](#)
- [Bartsch et al. 2010] T. Bartsch, A. Pistoia, and T. Weth, “ N -vortex equilibria for ideal fluids in bounded planar domains and new nodal solutions of the sinh-Poisson and the Lane–Emden–Fowler equations”, *Comm. Math. Phys.* **297**:3 (2010), 653–686. [MR](#) [Zbl](#)
- [Bjorland 2011] C. Bjorland, “The vortex-wave equation with a single vortex as the limit of the Euler equation”, *Comm. Math. Phys.* **305**:1 (2011), 131–151. [MR](#) [Zbl](#)
- [Burton 1989a] G. R. Burton, “Rearrangements of functions, saddle points and uncountable families of steady configurations for a vortex”, *Acta Math.* **163**:3-4 (1989), 291–309. [MR](#) [Zbl](#)

- [Burton 1989b] G. R. Burton, “Variational problems on classes of rearrangements and multiple configurations for steady vortices”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **6**:4 (1989), 295–319. [MR](#) [Zbl](#)
- [Burton 2005] G. R. Burton, “Global nonlinear stability for steady ideal fluid flow in bounded planar domains”, *Arch. Ration. Mech. Anal.* **176**:2 (2005), 149–163. [MR](#) [Zbl](#)
- [Burton and McLeod 1991] G. R. Burton and J. B. McLeod, “Maximisation and minimisation on classes of rearrangements”, *Proc. Roy. Soc. Edinburgh Sect. A* **119**:3–4 (1991), 287–300. [MR](#) [Zbl](#)
- [Caffarelli and Friedman 1985] L. A. Caffarelli and A. Friedman, “Convexity of solutions of semilinear elliptic equations”, *Duke Math. J.* **52**:2 (1985), 431–456. [MR](#) [Zbl](#)
- [Cao and Wang 2017] D. Cao and G. Wang, “Nonlinear stability of planar vortex patches in an ideal fluid”, preprint, 2017. [arXiv](#)
- [Cao and Wang 2019] D. Cao and G. Wang, “Steady vortex patch solutions to the vortex-wave system”, *Nonlinearity* **32**:5 (2019), 1882–1904. [MR](#) [Zbl](#)
- [Cao et al. 2014] D. Cao, Z. Liu, and J. Wei, “Regularization of point vortices pairs for the Euler equation in dimension two”, *Arch. Ration. Mech. Anal.* **212**:1 (2014), 179–217. [MR](#) [Zbl](#)
- [Cao et al. 2015] D. Cao, S. Peng, and S. Yan, “Planar vortex patch problem in incompressible steady flow”, *Adv. Math.* **270** (2015), 263–301. [MR](#) [Zbl](#)
- [Cao et al. 2019a] D. Cao, Y. Guo, S. Peng, and S. Yan, “Local uniqueness for vortex patch problem in incompressible planar steady flow”, *J. Math. Pures Appl.* (9) **131** (2019), 251–289. [MR](#) [Zbl](#)
- [Cao et al. 2019b] D. Cao, J. Wan, and G. Wang, “Nonlinear orbital stability for planar vortex patches”, *Proc. Amer. Math. Soc.* **147**:2 (2019), 775–784. [MR](#) [Zbl](#)
- [Elcrat and Miller 1991] A. R. Elcrat and K. G. Miller, “Rearrangements in steady vortex flows with circulation”, *Proc. Amer. Math. Soc.* **111**:4 (1991), 1051–1055. [MR](#) [Zbl](#)
- [Elcrat and Miller 1995] A. R. Elcrat and K. G. Miller, “Rearrangements in steady multiple vortex flows”, *Comm. Partial Differential Equations* **20**:9–10 (1995), 1481–1490. [MR](#) [Zbl](#)
- [Grossi and Takahashi 2010] M. Grossi and F. Takahashi, “Nonexistence of multi-bubble solutions to some elliptic equations on convex domains”, *J. Funct. Anal.* **259**:4 (2010), 904–917. [MR](#) [Zbl](#)
- [Lacave and Miot 2009] C. Lacave and E. Miot, “Uniqueness for the vortex-wave system when the vorticity is constant near the point vortex”, *SIAM J. Math. Anal.* **41**:3 (2009), 1138–1163. [MR](#) [Zbl](#)
- [Lin 1941] C. C. Lin, “On the motion of vortices in two dimensions, I: Existence of the Kirchhoff–Routh function”, *Proc. Nat. Acad. Sci. U.S.A.* **27** (1941), 570–575. [MR](#) [Zbl](#)
- [Lopes Filho et al. 2011] M. C. Lopes Filho, E. Miot, and H. J. Nussenzweig Lopes, “Existence of a weak solution in L^p to the vortex-wave system”, *J. Nonlinear Sci.* **21**:5 (2011), 685–703. [MR](#) [Zbl](#)
- [Marchioro and Pagani 1986] C. Marchioro and E. Pagani, “Evolution of two concentrated vortices in a two-dimensional bounded domain”, *Math. Methods Appl. Sci.* **8**:3 (1986), 328–344. [MR](#) [Zbl](#)
- [Marchioro and Pulvirenti 1991] C. Marchioro and M. Pulvirenti, “On the vortex-wave system”, pp. 79–95 in *Mechanics, analysis and geometry: 200 years after Lagrange*, edited by M. Francaviglia, North-Holland, Amsterdam, 1991. [MR](#) [Zbl](#)
- [Marchioro and Pulvirenti 1994] C. Marchioro and M. Pulvirenti, *Mathematical theory of incompressible nonviscous fluids*, Applied Mathematical Sciences **96**, Springer, 1994. [MR](#) [Zbl](#)
- [Miot 2012] E. Miot, “Two existence results for the vortex-wave system”, *Riv. Math. Univ. Parma (N.S.)* **3**:1 (2012), 131–146. [MR](#) [Zbl](#)
- [Smets and Van Schaftingen 2010] D. Smets and J. Van Schaftingen, “Desingularization of vortices for the Euler equation”, *Arch. Ration. Mech. Anal.* **198**:3 (2010), 869–925. [MR](#) [Zbl](#)

- [Turkington 1983a] B. Turkington, “On steady vortex flow in two dimensions, I”, *Comm. Partial Differential Equations* **8**:9 (1983), 999–1030. [MR](#) [Zbl](#)
- [Turkington 1983b] B. Turkington, “On steady vortex flow in two dimensions, II”, *Comm. Partial Differential Equations* **8**:9 (1983), 1031–1071. [MR](#) [Zbl](#)
- [Turkington 1987] B. Turkington, “On the evolution of a concentrated vortex in an ideal fluid”, *Arch. Rational Mech. Anal.* **97**:1 (1987), 75–87. [MR](#) [Zbl](#)
- [Wan 1988] Y. H. Wan, “Desingularizations of systems of point vortices”, *Phys. D* **32**:2 (1988), 277–295. [MR](#) [Zbl](#)

Received June 11, 2018.

DAOMIN CAO
INSTITUTE OF APPLIED MATHEMATICS
CHINESE ACADEMY OF SCIENCES
BEIJING,
CHINA
dmcao@amt.ac.cn

GUODONG WANG
INSTITUTE FOR ADVANCED STUDY IN MATHEMATICS
HARBIN INSTITUTE OF TECHNOLOGY
HARBIN,
CHINA
wanguodong14@mails.ucas.ac.cn

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 308

No. 2

October 2020

Existence of steady multiple vortex patches to the vortex-wave system	257
DAOMIN CAO and GUODONG WANG	
Relations of rationality for special values of Rankin–Selberg L -functions of $GL_n \times GL_m$ over CM-fields	281
HARALD GROBNER and GUNJA SACHDEVA	
A bound for the conductor of an open subgroup of GL_2 associated to an elliptic curve	307
NATHAN JONES	
Topology of complexity one quotients	333
Yael KARSHON and SUSAN TOLMAN	
Flag Bott manifolds and the toric closure of a generic orbit associated to a generalized Bott manifold	347
SHINTARÔ KUROKI, EUNJEONG LEE, JONGBAEK SONG and DONG YOUP SUH	
Projective cases for the restriction of the oscillator representation to dual pairs of type I	393
SABINE J. LANG	
A remark on a trace Paley–Wiener theorem	407
GORAN MUIĆ	
Spectrum of the Laplacian and the Jacobi operator on rotational CMC hypersurfaces of spheres	419
OSCAR M. PERDOMO	
Mean curvature flow in a Riemannian manifold endowed with a Killing vector field	435
LIANGJUN WENG	
Green correspondence and relative projectivity for pairs of adjoint functors between triangulated categories	473
ALEXANDER ZIMMERMANN	