

*Pacific  
Journal of  
Mathematics*

**SPECTRUM OF THE LAPLACIAN  
AND THE JACOBI OPERATOR  
ON ROTATIONAL CMC HYPERSURFACES OF SPHERES**

OSCAR M. PERDOMO

# SPECTRUM OF THE LAPLACIAN AND THE JACOBI OPERATOR ON ROTATIONAL CMC HYPERSURFACES OF SPHERES

OSCAR M. PERDOMO

Let  $M \subset \mathbb{S}^{n+1} \subset \mathbb{R}^{n+2}$  be a compact CMC rotational hypersurface of the  $(n+1)$ -dimensional Euclidean unit sphere. Denote by  $|A|^2$  the square of the norm of the second fundamental form and  $J(f) = -\Delta f - nf - |A|^2 f$  the stability or Jacobi operator. In this paper we compute the spectra of their Laplace and Jacobi operators in terms of eigenvalues of second order Hill's equations. For the minimal rotational examples, we prove that the stability index — the numbers of negative eigenvalues of the Jacobi operator counted with multiplicity — is greater than  $3n+4$  and we also prove that there are at least 2 positive eigenvalues of the Laplacian of  $M$  smaller than  $n$ . When  $H$  is not zero, we have that every nonflat CMC rotational immersion is generated by rotating a planar profile curve along a geodesic called the axis of rotation. We assume that the coordinates of this plane has been set up so that the axis of rotation goes through the origin. The planar profile curve is made up of  $m$  copies, each one of them is a rigid motion of a single curve that we will call the fundamental piece. For this reason every nonflat rotational CMC hypersurface has  $Z_m$  in its group of isometries. If  $\theta$  denotes the change of the angle of the fundamental piece when written in polar coordinates, then  $l = \frac{m\theta}{2\pi}$  is a nonnegative integer. For unduloids (a subfamily of the rotational CMC hypersurfaces that include all the known embedded examples), we show that the number of negative eigenvalues of the operator  $J$  counted with multiplicity is at least  $(2l-1)n + (2m-1)$ .

## 1. Introduction

Rotational constant mean curvature hypersurfaces of spheres provide a variety of examples that can be used to understand the nature of CMC hypersurfaces in general. In this paper we derive a formula for the Laplacian on these hypersurfaces that allows us to compute the spectra of their Jacobi and Laplace operators to any desired degree of accuracy. Results on the stability index are relevant in both cases, the case  $H = 0$  and the case  $H \neq 0$ . Let us denote by  $\mathcal{M}^n$  any minimal, not necessarily

*MSC2010:* 53C42.

*Keywords:* stability operator, Jacobi operator, stability index, CMC, spheres, rotational hypersurfaces.

rotational, compact  $n$ -dimensional minimal hypersurface of the sphere. The spectra of the Laplacian and the stability operators on compact minimal hypersurfaces  $\mathcal{M}^n \subset \mathbb{S}^{n+1}$  have been one of the central topics in differential geometry. Let us denote by  $\lambda_1(\mathcal{M}^n)$ , the first nonzero eigenvalue of the Laplace operator of  $\mathcal{M}$ . For minimal hypersurfaces of spheres, the spectrum of the Laplacian is known only when  $\mathcal{M}^n$  is an Euclidean sphere,  $\mathcal{M}^n$  is the product of Euclidean spheres, or  $\mathcal{M}^n$  is a cubic isoparametric hypersurface, [Solomon 1990a, 1990b]. It is known that for any minimal hypersurface of the sphere,  $n$  is one of the eigenvalues of the Laplacian. Yau [1982] has conjectured that if  $\mathcal{M}^n$  is compact and embedded, then  $\lambda_1(\mathcal{M}^n) = n$ . A positive partial result of this conjecture was given by Choi and Wang [1983]. They showed that if  $\mathcal{M}^n$  is compact and embedded, then  $\lambda_1(\mathcal{M}^n) \geq \frac{n}{2}$ . In this paper we prove that if  $\mathcal{M}^n$  is rotational, then  $\lambda_1(\mathcal{M}^n) < n$ . Moreover we show that there are at least two positive eigenvalues of the Laplace operator smaller  $n$ .

Regarding the spectrum of the Jacobi operator, we have that the number of negative eigenvalues of the operator  $J$  counted with multiplicity is known as the stability index of  $\mathcal{M}^n$  and it is denoted as  $\text{ind}(\mathcal{M}^n)$ . It has been conjectured that if  $\mathcal{M}^n$  is not an Euclidean sphere, then  $\text{ind}(\mathcal{M}^n) = n + 3$  implies that  $\mathcal{M}^n$  is a product of Euclidean spheres. This conjecture was proven by Urbano [1990], when  $n = 2$ . For general  $n$  only partial results are known. In [Perdomo 2004a], the author shows that if  $\text{ind}(\mathcal{M}^n) = n + 3$ , then  $\int_{\mathcal{M}} |A|^2 \leq \int_{\mathcal{M}} (n - 1)$  with equality only if  $\mathcal{M}^n$  is a product of Euclidean spheres. On the other hand in [Perdomo 2004b], the author shows that if  $\mathcal{M}^n$  is rotational, then,  $\int_{\mathcal{M}} |A|^2 \leq \int_{\mathcal{M}} (n - 1)$ . Therefore, it may seem that the rotational minimal hypersurfaces are good candidates for a counterexample of the conjecture. This is not the case. Due to their symmetries, their stability index must be greater than  $n + 3$ . See [Perdomo 2001].

In this paper we show that there is a big jump in the stability index among the minimal rotational examples. We prove that if  $\mathcal{M}^n$  is rotational and minimal and it is not a product of spheres then  $\text{ind}(\mathcal{M}^n) \geq 3n + 5$ . Some other partial results on this conjecture are found in [Simons 1968; Perdomo 2002; Savo 2010]. One of the most important applications of this type of estimate on the stability index was given by Marques and Neves [2014], where, among other tools, they used Urbano's result to prove Willmore's conjecture.

To motivate the study of the stability index when  $H$  is a nonzero constant, we would like to point out that any time we have a one-parametric family of CMC hypersurfaces, a jump in the stability index as a function of the parameter indicates a potential bifurcation point, a hypersurface that is part of another one-parametric family of CMC hypersurfaces. As an example of this fact, the families of embedded rotational CMC hypersurfaces on the sphere explained by the author in [Perdomo 2010] can be viewed as families emerging from bifurcation points of the family  $S^1(r) \times S^{n-1}(\sqrt{1-r^2}) \subset \mathbb{S}^n$ . See [Alías and Piccione 2013].

To illustrate the way we can use our results to estimate the eigenvalues, we pick a 3-dimensional rotational minimal hypersurface in  $\mathbb{S}^4$  and prove that the first three eigenvalues of the Laplace operator are: 0, a number near 0.4404 with multiplicity 2, and 3 with multiplicity 5. We also show that the negative eigenvalues of the Jacobi operator are: a number near  $-8.6534$  with multiplicity 1, a number near  $-8.52078$  with multiplicity 2,  $-3$  with multiplicity 5, a number near  $-2.5596$  with multiplicity 6, and a number near  $-1.17496$  with multiplicity 1. The stability index of this hypersurface is thus 15.

When  $H$  is not zero, we prove (see [Theorem 3.7](#)) a lower bound for the number of negative eigenvalues of the Jacobi operator that generalizes to any dimension the result on the stability index of constant mean curvature tori of revolution in the 3-sphere proven by Rossman and Sultana [\[2007\]](#).

We will be using the *oscillation theorem* for the periodic problem on the Hill's equation (a proof can be found in [\[Magnus and Winkler 1966\]](#)).

**Theorem 1.1.** *Consider the differential equation*

$$(1-1) \quad z''(t) + (\lambda + Q(t))z(t) = 0$$

where  $Q$  is a smooth  $T$ -periodic function. For any  $\lambda$  let us define

$$\delta(\lambda) := z_1(T, \lambda) + z_2'(T, \lambda),$$

where  $z_1(t, \lambda)$  and  $z_2(t, \lambda)$  are solutions of (1-1) such that  $z_1(0, \lambda) = 1$ ,  $z_1'(0, \lambda) = 0$  and  $z_2(0, \lambda) = 0$ ,  $z_2'(0, \lambda) = 1$ . There exists an increasing infinite sequence of real numbers  $\lambda_1, \lambda_2, \dots$  such the differential equation (1-1) has a  $T$ -periodic solution if and only if  $\lambda = \lambda_j$ . Moreover the  $\lambda_j$  are the roots of the equation  $\delta(\lambda) = 2$ . The function  $\delta$  is called the discriminant function of the operator  $K[z] = z''(t) + Q(t)z(t)$ .

We will also be using the following theorem proven by Haupt [\[1914\]](#) from the Hill's equation theory. The next presentation of the Haupt theorem can be found in [\[Magnus and Winkler 1966\]](#).

**Theorem 1.2** [\[Haupt 1914\]](#). *Let us denote by  $\lambda_1 < \lambda_2 \leq \lambda_3 \leq \lambda_4 \dots$  the sequence of eigenvalues of the Hill's equation presented in [Theorem 1.1](#). If  $z(t)$  is a nonzero  $T$ -periodic solution of (1-1) with  $\lambda = \lambda_i$  then, the number of zeros of  $z(t)$  in the interval  $[0, T)$  is  $2 \lfloor \frac{i}{2} \rfloor$ .*

We would like to point out that Beekmann and Lökés [\[1990\]](#) have used the Hill equation to find bounds on the eigenvalues of the Laplacian on toroidal surfaces.

## 2. Describing rotational CMC hypersurfaces of spheres

Any compact CMC rotational hypersurface of  $\mathbb{S}^{n+1}$  is given by an immersion  $\phi : \mathbb{S}^{n-1} \times \mathbb{R} \rightarrow \mathbb{S}^{n+1}$  where,

$$(2-1) \quad \phi(y, t) = (r(t) y, \sqrt{1 - r(t)^2} \cos(\theta(t)), \sqrt{1 - r(t)^2} \sin(\theta(t)))$$

and  $r(t)$  is a positive  $T$ -periodic function that satisfies the conditions

$$(2-2) \quad (r')^2 + r^2(1 + \lambda^2) = 1,$$

with

$$(2-3) \quad \lambda = H + c^{-n/2} r^{-n}, \quad \theta(t) = \int_0^t \frac{r(\tau)\lambda(\tau)}{1 - r^2(\tau)} d\tau,$$

and  $c$  is a positive real number that satisfies

$$(2-4) \quad \theta(T) = 2\pi \frac{l}{m} \quad \text{where } l \text{ and } m \text{ are relative prime integers.}$$

The condition on  $c$  in (2-4) guarantees that the immersion  $\phi$  satisfies

$$\phi(y, t + mT) = \phi(y, t)$$

and makes  $M$  compact. Recall that the function  $r(t)$  depends on  $c$  since  $\lambda(t)$  depends on  $c$ . Also, since the function  $\theta(t)$  depends on  $r(t)$  and  $\lambda(t)$ , it follows that  $\theta(T)$  depends on  $c$  as well.

**Remark 2.1.** When  $M$  is minimal, Otsuki [1972; 1993], showed that the expression

$$\theta(T) = K(c) = \int_0^{T/2} \frac{c^{-1/n} r^{1-n}(\tau)}{1 - r^2(\tau)} d\tau$$

lies between  $\pi$  and  $\sqrt{2}\pi$ . As a consequence the equation  $\theta(T) = 2\pi \frac{l}{m}$  cannot be solved with  $l = 1$  and therefore no minimal rotational hypersurface is embedded.

The principal curvatures of  $M$  are  $\lambda$  with multiplicity  $(n - 1)$  and

$$\mu = H - (n - 1)c^{-n/2} r^{-n}$$

with multiplicity one. Differentiating (2-2) we obtain

$$(2-5) \quad r'' + r + r\lambda\mu = 0.$$

The next expression explicitly provides the Gauss map  $\nu$  of the immersion (2-1),

$$(2-6) \quad \nu(y, t) = \left( -r\lambda y, \frac{r^2\lambda \cos \theta - r' \sin \theta}{\sqrt{1 - r(t)^2}}, \frac{r^2\lambda \sin \theta + r' \cos \theta}{\sqrt{1 - r(t)^2}} \right)$$

All the details of the construction of these hypersurfaces can be found in [Perdomo 2010].

### 3. Main theorems

Before stating the following theorem, we recall that  $m$  is the integer given in (2-4),  $T$  is the period of the function  $r(t)$ , and  $mT$  is the period of the immersion  $\phi$ .

**Theorem 3.1.** *Let  $M$  be the rotational symmetric hypersurface defined in (2-1). For any function  $\bar{f} : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$  we define  $f : M \rightarrow \mathbb{R}$  as  $f(\phi(t, y)) = \bar{f}(y)$ . Likewise, for any  $mT$ -periodic function  $\bar{g} : \mathbb{R} \rightarrow \mathbb{R}$  we define  $g : M \rightarrow \mathbb{R}$  as  $g(\phi(t, y)) = \bar{g}(t)$ . We will denote by  $\bar{\Delta}$  the Laplacian operator on  $\mathbb{S}^{n-1}$ . With this notation we have*

$$(3-1) \quad \Delta(fg) = f\left(\bar{g}'' + (n-1)\frac{r'}{r}\bar{g}'\right) + \frac{\bar{\Delta}(\bar{f})}{r^2}g.$$

*Proof.* The proof is a direct computation using the fact that  $\nabla(fg) = \frac{g}{r}\bar{\nabla}\bar{f} + f\bar{g}'\frac{\partial}{\partial t}$  and that  $\operatorname{div}\left(\frac{\partial}{\partial t}\right) = (n-1)\frac{r'}{r}$ .  $\square$

**Remark 3.2.** We would like to point out that we are using the fact that the ambient space is  $\mathbb{R}^{n+2}$  in the argument above. In particular we are using the natural identification of all tangent spaces  $T_x\mathbb{R}^{n+2}$  with  $\mathbb{R}^{n+2}$ . If we decide to work intrinsically we will notice that the formula that compares the gradients will have an additional factor of  $r$ . From the point of view of intrinsic differential geometry, the formula in Theorem 3.1 can be generalized to warped products. See [Marrocos and Gomes 2019].

**Theorem 3.3.** *Let  $\alpha_1 = 0, \alpha_2 = (n-1), \dots, \alpha_k = (k-1)(n+k-3), \dots$  denote the spectrum of  $\mathbb{S}^{n-1}$ . The spectrum of the Laplace operator  $\Delta$  on  $M$  is given by  $\bigcup_{k=1}^{\infty} \Gamma_k$ , where,*

$$\Gamma_k = \{\lambda(k, 1), \lambda(k, 2), \dots\}$$

*is the ordered spectrum of the operator (defined on the set of  $mT$  periodic functions)*

$$K_{\Delta,k}[z] = z'' + (n-1)\frac{r'}{r}z' - \frac{\alpha_k}{r^2}z$$

*The spectrum of the Jacobi operator  $J$  on  $M$  is given by  $\bigcup_{k=1}^{\infty} \mathbb{F}_k$ , where*

$$\mathbb{F}_k = \{\tilde{\lambda}(k, 1), \tilde{\lambda}(k, 2), \dots\}$$

*is the ordered spectrum of the operator (defined on the set of  $mT$  periodic functions)*

$$K_{J,k}[z] = z'' + (n-1)\frac{r'}{r}z' + \left(n + nH^2 + n(n-1)c^{-n}r^{-2n} - \frac{\alpha_k}{r^2}\right)z$$

*Proof.* Let us prove the case of the Laplacian. By Theorem 3.1 we have that if  $\bar{f}_k$  is an eigenfunction of the Laplacian on  $\mathbb{S}^{n-1}$  with eigenvalue  $\alpha_k$  and  $\bar{g}_l(t)$  is an

eigenfunction of the operator  $K_{\Delta,k}$  with eigenvalue  $\lambda(k, l)$ , then  $h_{k,l} : M \rightarrow \mathbb{R}$  given by  $h_{kl} = f_k g_l$  is an eigenfunction of the Laplacian with eigenvalue  $\lambda(k, l)$ . Therefore  $\bigcup_{k=1}^{\infty} \Gamma_k$  is contained in the spectrum of  $\Delta$ . To prove the reverse inclusion, we only need to point out that every smooth function  $h : M \rightarrow \mathbb{R}$  can be written as a series of eigenfunctions of the form  $h_{kl}$ . There exists a basis

$$\bar{f}_{1,1}, \bar{f}_{2,1}, \bar{f}_{2,2}, \dots, \bar{f}_{2,n}, \bar{f}_{3,1}, \bar{f}_{3,2}, \dots, \bar{f}_{3,2n}, \bar{f}_{4,1}, \dots$$

for  $L^2(\mathbb{S}^{n-1})$  with  $\bar{\Delta} \bar{f}_{k,j} + \alpha_k \bar{f}_{k,j} = 0$ . So any  $h : M \rightarrow \mathbb{R}$  can be written as a sum of the form

$$a_{1,1}(t) \bar{f}_{1,1} + a_{2,1}(t) \bar{f}_{2,1} + \dots + a_{2,n}(t) \bar{f}_{2,n} + a_{3,1}(t) \bar{f}_{3,1} + \dots$$

We obtain the desired expression for the function  $h : M \rightarrow \mathbb{R}$ , by noticing that each  $a_{k,l}(t)$  can now be expanded in eigenfunctions of the operator  $K_{\Delta,k}$ . The proof for the Jacobi operator is similar and uses the following expression for  $|A|^2$ :

$$|A|^2 = n(H^2 + (n - 1)c^{-n}r^{-2n}). \quad \square$$

The following lemma allows us to use the oscillation theorem to compute the eigenvalues for the second order differential equations on [Theorem 3.3](#).

**Lemma 3.4.** *Let us denote by  $\lambda = H + c^{-n/2}r^{-n}$  and  $\mu = H - (n - 1)c^{-n/2}r^{-n}$ . The change of variables  $u = r^{(n-1)/2}z$  gives us,*

$$\begin{aligned} K_{\Delta,k} &= z'' + (n - 1)\frac{r'}{r}z' - \frac{\alpha_k}{r^2}z \\ &= \frac{1}{r^{(n-1)/2}} \left( u'' + \left( \frac{1}{4}\lambda^2((n - 1)(n - 3)) + \frac{1}{2}\lambda\mu(n - 1) - \frac{4\alpha_k + (n - 3)(n - 1)}{4r^2} \right. \right. \\ &\quad \left. \left. + \frac{1}{4}(n - 1)^2 \right) u \right) \end{aligned}$$

and

$$\begin{aligned} K_{J,k} &= z'' + (n - 1)\frac{r'}{r}z' + \left( n + nH^2 + n(n - 1)c^{-n}r^{-2n} - \frac{\alpha_k}{r^2} \right) z \\ &= \frac{1}{r^{(n-1)/2}} \left( u'' + \frac{1}{4} \left( \lambda^2(n^2 - 1) + 2\lambda\mu(n - 1) + 4\mu^2 + (n + 1)^2 \right. \right. \\ &\quad \left. \left. - \frac{4\alpha_k + (n - 3)(n - 1)}{r^2} \right) u \right) \end{aligned}$$

**Theorem 3.5.** *If  $\mathcal{M} \subset \mathbb{S}^{n+1}$  is a rotational minimal compact hypersurface, then  $\text{ind}(\mathcal{M}^n) \geq 3n + 5$  and there are at least two positive eigenvalues of the Laplace operator smaller than  $n$ . More precisely, if  $m$  and  $l$  are the relative prime integers in (2-4), then*

- $l \geq 2, m \geq 3,$

- $\text{ind}(\mathcal{M}^n) \geq (2l - 1)n + (2m - 1)$  and,
- there are exactly  $2l - 2$  positive eigenvalues of the Laplacian smaller than  $n$ .

*Proof.* As pointed out in Remark 2.1, the positive integer  $l$  that satisfies the equation  $\theta(T) = 2\pi \frac{l}{m}$  must be greater than 1. We also have that  $m \geq 3$ . Recall that Otsuki has shown that, for any  $c$ , the integral that defines  $\theta(T)$  is a number between  $\pi$  and  $\sqrt{2}\pi$ . In this proof we will be using the notation for  $\lambda(k, j)$  and  $\tilde{\lambda}(k, j)$  introduced in Theorem 3.3. More precisely, the eigenvalues of the operator  $K_{\Delta, k}$  are

$$\lambda(k, 1) < \lambda(k, 2) \leq \lambda(k, 3) \leq \dots$$

and the eigenvalues of the operator  $K_{J, k}$  are

$$\tilde{\lambda}(k, 1) < \tilde{\lambda}(k, 2) \leq \tilde{\lambda}(k, 3) \leq \dots$$

A direct verification shows that the functions  $f_1(t) = \sqrt{1 - r^2(t)} \cos(\theta)$  and  $f_2(t) = \sqrt{1 - r^2(t)} \sin(\theta)$  satisfy the equation  $K_{\Delta, 1}(f_i) + nf_i = 0$ , for  $i = 1, 2$ . Since the functions  $f_1$  and  $f_2$  have  $2l$  zeroes in the interval  $[0, mT)$  then, by Theorem 1.2 we conclude that  $n = \lambda(1, 2l) = \lambda(1, 2l + 1)$ . Therefore, the eigenvalues of  $K_{\Delta, 1}$  are

$$\lambda(1, 1) = 0 < \lambda(1, 2) \leq \lambda(1, 3) < \dots < \lambda(1, 2l) = \lambda(1, 2l + 1) = n < \dots$$

This shows the existence of  $2l - 2$  positive eigenvalues (counted with multiplicity) of the Laplace operator smaller than  $n$ . A direct verification shows that  $r(t)$  satisfies the equation  $K_{\Delta, 2}(r) + nr = 0$ . Since  $r(t)$  is positive, then  $\lambda(2, 1) = n$ . Therefore all the eigenvalues of the Laplace operator smaller than  $n$  comes from the operators  $K_{\Delta, 1}$ . We conclude that there are exactly  $2l - 2$  positive eigenvalues of the Laplacian smaller than  $n$ . Let us show the inequality for the stability index. A direct verification shows that the function  $r'$  satisfy the equation  $K_{J, 1}(r') = 0$ . The numbers of zeroes of  $r'$  in the interval  $[0, mT)$  is  $2m$ . By Theorem 1.2 we conclude that either  $0 = \tilde{\lambda}(1, 2m)$  or  $0 = \tilde{\lambda}(1, 2m + 1)$  Therefore the operator  $K_{J, 1}$  has at least  $2m - 1$  negative eigenvalues. Let us analyze the spectrum of the operator  $K_{J, 2}$ . A direct verification shows that the functions

$$f_3(t) = \frac{-c^{-n/2}r^{1-n} \cos(\theta) + rr' \sin(\theta)}{\sqrt{1 - r^2(t)}} \quad \text{and} \quad f_4(t) = \frac{c^{-n/2}r^{1-n} \sin(\theta) + rr' \cos(\theta)}{\sqrt{1 - r^2(t)}}$$

satisfy the equation  $K_{J, 2}(f_i) = 0$  for  $i = 3, 4$ . The numbers of zeroes  $f_3$  and  $f_4$  in the interval  $[0, mT)$  is at least  $2l$ . Let us check this property for the function  $f_3(t)$ . We have that  $\theta(t)$  changes from 0 to  $2\pi l$  when  $t$  moves from 0 to  $mT$ . Let  $t_1, t_2, \dots, t_{2l-1}$  be increasing values of  $t$  such that  $\theta(t_i) = i\pi$ . Recall that  $\theta(0) = 0$  and  $\theta(mT) = 2\pi l$ .

Since

$$f_3(0) = -\frac{c^{-n/2}r(0)^{1-n}}{\sqrt{1-r(0)^2}} < 0 \quad \text{and} \quad f_3(t_1) = \frac{c^{-n/2}r(0)^{1-n}}{\sqrt{1-r(0)^2}} > 0,$$

there is at least one zero of  $f_3$  between 0 and  $t_1$ . The same argument shows that there must be at least one zero of  $f_3(t)$  between  $t_1$  and  $t_2$ . By continuing with this argument we conclude that  $f_3$  must have at least  $2l$  zeroes. A similar argument shows that  $f_4$  has also at least  $2l$  zeroes. By [Theorem 1.2](#) we conclude that  $0 = \tilde{\lambda}(2, 2k) = \tilde{\lambda}(2, 2k + 1)$ , where  $2k$  is the number of zeroes of  $f_4$  and  $f_5$  on the interval  $[0, mT)$ . Recall that  $k \geq l$ . Therefore the operator  $K_{J,2}$  has at least  $2l - 1$  negative eigenvalues. Since the multiplicity of the eigenvalue  $\alpha_2 = (n - 1)$  of the Laplace operator on  $S^{n-1}$  is  $n$ , we conclude that  $\text{ind}(\mathcal{M}) \geq (2l - 1)n + (2m - 1)$ .  $\square$

**Remark 3.6.** There is a geometrical reason for the functions  $r'$  and  $f_i$ ,  $i = 1, 2, 3, 4$  to satisfy the respective Hill's equation mentioned in the proof of the previous theorem. If we write the immersion  $\phi$  in (2-1) as  $(\phi_1, \dots, \phi_{n+2})$  and the Gauss map  $\nu$  in (2-6) as  $(\nu_1, \dots, \nu_{n+2})$  then, the functions  $f_1$  and  $f_2$  are the last two coordinates of the immersion  $\phi$  and, in general, all the coordinates of an immersion of a minimal hypersurface of  $\mathbb{S}^{n+1}$  are eigenfunctions of the Laplacian associated with the eigenvalue  $n$ . The function  $r'$  is the simplification of the expression  $\phi_{n+2}\nu_{n+1} - \phi_{n+1}\nu_{n+2}$  and in general all the functions of the form  $\nu_i\phi_j - \nu_j\phi_i$  are either the zero function or they are eigenfunctions of the stability operator associated with the eigenvalue 0. Finally, we have that  $f_4$  and  $f_5$  are factors of the expression  $\phi_1\nu_{n+1} - \phi_{n+1}\nu_1$  and  $\phi_1\nu_{n+2} - \phi_{n+2}\nu_1$ .


**Theorem 3.7.** *Let  $M \subset \mathbb{S}^{n+1}$  be a rotational CMC compact hypersurface described in (2-1) with  $H \geq 0$ . If  $m$  and  $l$  are the relative prime integers in (2-4) then, the number of negative eigenvalues of the Jacobi operator counted with multiplicity is at least  $(2l - 1)n + (2m - 1)$*

*Proof.* The proof is similar to the one presented for the minimal case. A direct verification shows that  $J_{J,0}[r']$  vanishes and the numbers of zeroes of  $r'$  in the interval  $[0, mT)$  is  $2m$ . By [Theorem 1.2](#) we conclude that either  $0 = \tilde{\lambda}(1, 2m)$  or  $0 = \tilde{\lambda}(1, 2m + 1)$ . Therefore the operator  $K_{J,1}$  has at least  $2m - 1$  negative eigenvalues. Let us analyze the spectrum of the operator  $K_{J,2}$ . A direct verification shows that the functions

$$f_1(t) = \frac{c^{-n/2}r(t)(c^{n/2}(H \cos(\theta(t)) - r'(t) \sin(\theta(t))) + r(t)^{-n} \cos(\theta(t)))}{\sqrt{1-r(t)^2}}$$

and

$$f_2(t) = \frac{c^{-n/2}r(t)(c^{n/2}(r'(t) \cos(\theta(t)) + H \sin(\theta(t))) + r(t)^{-n} \sin(\theta(t)))}{\sqrt{1-r(t)^2}}$$



**Figure 1.** On the left we have the graph of the solution  $r(t)$  for the value  $ac = 2.8284247911397589$ . On the right we have the graph of the function  $\theta(t)$  defined in (2-3).

satisfy the equation  $K_{J,2}(f_i) = 0$  for  $i = 1, 2$ . The numbers of zeroes  $f_1$  and  $f_2$  in the interval  $[0, mT]$  is at least  $2l$ . By Theorem 1.2 we conclude that  $0 = \tilde{\lambda}(2, 2k) = \tilde{\lambda}(2, 2k + 1)$ , where  $2k$  is the number of zeroes of  $f_1$  and  $f_2$  on the interval  $[0, mT]$ . Recall that  $k \geq l$ . Therefore the operator  $K_{J,2}$  has at least  $2l - 1$  negative eigenvalues. Since the multiplicity of the eigenvalue  $\alpha_2 = (n - 1)$  of the Laplace operator on  $S^{n-1}$  is  $n$ , we conclude that the number of negative eigenvalue of the operator  $J$  is greater than  $(2l - 1)n + (2m - 2)$ .  $\square$

#### 4. An example that illustrates the method

In this section we pick an explicit rotational 3-dimensional minimal hypersurface  $M \subset S^4$  and we compute the first three eigenvalues of the Laplacian and its stability index.


**Construction of the particular example.** By the intermediate value theorem, it is easy to see that there is a value of  $c$  near  $ac = 2.82842479911$  such that the function  $r(t)$  has period  $T$  near  $aT = 2.6722005616$  and  $\theta(T) = \frac{4\pi}{3}$ . See Equation (2-4). In this case  $l = 2, m = 3$  and our manifold  $M$  is defined by this choice of  $c$ . Recall that the immersion  $\phi : S^2 \times \mathbb{R} \rightarrow S^4$  is given by

$$(4-1) \quad \phi(y, t) = (r(t)y, \sqrt{1 - r(t)^2} \cos(\theta(t)), \sqrt{1 - r(t)^2} \sin(\theta(t))).$$


Figure 1 shows the solution  $r(t)$  that produces the manifold  $M$ , and Figure 2 shows the profile curve of the rotational manifold  $M$ .

*Computing the first three eigenvalues of the Laplacian.* In order to use the oscillation theorem (Theorem 1.1) we notice that making  $u = rz$  we obtain (see also Lemma 3.4)

$$K_{\Delta,k}[z] = z'' + 2\frac{r'}{r}z' - \frac{\alpha_k}{r^2}z = \frac{1}{r} \left( u'' + \left( 1 - \frac{2}{c^3r^6} - \frac{\alpha_k}{r^2} \right) u \right).$$



**Figure 2.** Profile curve of the  $M$ . This curve is parametrized by  $(\sqrt{1 - r(t)^2} \cos(\theta(t)), \sqrt{1 - r(t)^2} \sin(\theta(t)))$ . The red piece represent the curve when  $t$  moves from 0 to  $T$ . Recall that the period of the immersion is  $3T$ .




**Figure 3.** Graph of the function  $\delta(\lambda)$ . The roots of the equation  $\delta(\lambda) = 2$  give us eigenvalues of the Laplacian of the form  $\lambda(1, j)$  defined on [Theorem 3.3](#).

Therefore  $\lambda(k, i)$  is an eigenvalue of the operator  $K_{\Delta,k}$  if and only if  $\lambda(k, i)$  is an eigenvalue of the operator

$$\bar{K}_{\Delta,k}[u] = u'' + \left(1 - \frac{2}{c^3 r^6} - \frac{\alpha_k}{r^2}\right)u.$$

For  $\alpha_1 = 0$ , [Figure 3](#) shows the discriminant function  $\delta(\lambda)$  for the operator  $\bar{K}_{\Delta,1}$ .



**Figure 4.** Graph of two solutions  $\xi_1$  and  $\xi_2$  of the equation  $K_{\Delta,1}[z] + 0.4404z = 0$ .

Figure 3 was made by taking 3400 values of  $\lambda$  between  $-0.1$  and  $3.3$ , one every  $0.001$ . For each value of  $\lambda$  we solve two differential equations to find the functions  $z_1(t, \lambda)$  and  $z_2(t, \lambda)$  defined in Theorem 1.1. Once we have  $z_1(t, \lambda)$  and  $z_2(t, \lambda)$  we computed  $\delta(\lambda)$ . The crossing of the graph of  $\delta(\lambda)$  with the horizontal line  $y = 2$  at  $\lambda(1, 1) = 0$  was expected since  $z(t) = 1$  is an eigenfunction and the crossing at  $\lambda = 3$  with multiplicity 2 was expected because the last two coordinates of the immersion, the functions  $\sqrt{1 - r^2} \cos(\theta)$  and  $\sqrt{1 - r^2} \sin(\theta)$ , are eigenfunctions of the Laplacian of  $M$ , see [Simons 1968]. Regarding the crossing near  $0.44$  we can check that  $|\delta(0.4404) - 2|$  is smaller than  $10^{-6}$ . Figure 4 shows two linearly independent solutions  $\xi_1$  and  $\xi_2$  of the equation  $K_{\Delta,1}[z] + 0.4404z = 0$ . All together we have 3 eigenvalues of  $K_{\Delta,1}$  smaller than 3, which agrees with Theorem 3.5.

We will move now to study the operator  $K_{\Delta,2}$ . Since the coordinates of the immersion  $\phi$  are eigenfunction of the Laplacian we have that the function  $r(t)$  satisfies the equation  $K_{\Delta,2}(r(t)) = -3r(t)$ . The previous equation also follows from (2-5). Since  $r(t)$  is positive,  $\lambda(2, 1) = 3$  is the first eigenvalue of  $K_{\Delta,2}$  and it has multiplicity 1.

**Remark 4.1.** Since the sequence  $\alpha_k$  is increasing, the sequence  $\lambda(k, 1)$  is also increasing.


From the previous remark we deduce that all other eigenvalues of the Laplacian of  $M$  are greater than 3.

*Computing the negative eigenvalues of the Jacobi operator.* Once again we use the oscillation theorem (Theorem 1.1). The change of variables  $u = rz$  gives us (see also Lemma 3.4)

$$K_{J,k}[z] = z'' + 2\frac{r'}{r}z' + \left(\frac{6}{c^3r^6} + 3 - \frac{\alpha_k}{r^2}\right)z = \frac{1}{r}\left(u'' + \left(4 + \frac{4}{c^3r^6} - \frac{\alpha_k}{r^2}\right)u\right).$$

Similar to the case of the Laplacian operator, we can compute the eigenvalues of the Jacobi operator by computing the eigenvalues of the operator

$$\bar{K}_{J,k}[u] = u'' + \left(4 + \frac{4}{c^3r^6} - \frac{\alpha_k}{r^2}\right)u.$$




**Figure 5.** Graph of the function  $\delta(\lambda)$  for the operator  $\bar{K}_{J,1}$ . The roots of the equation  $\delta(\lambda) = 2$  give us eigenvalues of the Jacobi operator of the form  $\lambda(1, j)$ . The graph on the center and on the right shows the function  $\delta$  on smaller intervals

Figure 5 shows the discriminant  $\delta$  for the operator  $\bar{K}_{J,1}$ . A closer look at the function tell us that the negative solutions of the equation  $\delta(\lambda) = 2$  are on the intervals  $[-8.7, -8.5]$  and  $[-3.1, 0]$ .


For the first eigenvalue of the Jacobi operator, it is easy to use the intermediate value theorem to show that  $\tilde{\lambda}(1, 1) = -8.6534$  within an error of  $10^{-4}$ . This eigenvalue has multiplicity one and Figure 6 shows a nonzero periodic eigenfunction of the operator  $K_{J,1}$ . For the next value we have that  $|\delta(-8.53078) - 2| < 10^{-5}$ . For this value of  $\lambda$  the two fundamental solutions of the equation  $K_{J,1}[z] + \lambda z = 0$  are shown in Figure 7. The next eigenvalue is  $-3$  with multiplicity 2, this eigenvalue was expected due to the fact that the coordinate functions of the Gauss map are eigenfunctions of the Jacobi operator. For the next eigenvalue we have that  $|\delta(-1.1749673) - 2| < 10^{-5}$ . The existence of an eigenvalue near  $-1.17496$  with multiplicity one is given by the Intermediate Value Theorem, see Figure 5. We know that this is the last negative eigenvalue because 0 is known to be an eigenvalue of  $K_{J,1}$ .

We now study the operator  $K_{J,2}$ . Figure 8 shows the discriminant  $\delta$  for the operator  $\bar{K}_{J,2}$ . We can directly check that  $K_{J,2}(r^{-2}) = 3r^{-2}$ . Since  $r(t)$  is positive, then we have that  $-3$  is the first eigenvalue of  $K_{J,2}$  with multiplicity one. Since we have that  $|\delta(-2.5596) - 2| < 10^{-5}$ , then there is an eigenvalue of  $K_{J,2}$  with multiplicity 2 near  $-2.5596$ . We can check that the first eigenvalue of the operator  $K_{J,3}$  is close to 4.3484453. Therefore we have gotten all negative eigenvalues of the Jacobi operator, in summary we have

**Remark 4.2.** Since the first two eigenvalues of  $\mathbb{S}^2$  are 0 with multiplicity 1 and 2 with multiplicity 3, then the stability index of  $M$  is 15. Counting with multiplicity we have that the first eigenvalue of the stability operator  $J$  on  $M$  is near  $-8.6534$  and has multiplicity one. We have two eigenvalues near  $-8.52078$ , it could be only one with multiplicity 2. We have  $-3$  with multiplicity 5. Even though this was known, it is interesting to point out that the multiplicity is 5 because  $-3$  is an eigenvalue with multiplicity 2 of  $K_{J,1}$  and  $-3$  is an eigenvalue of multiplicity 1 of



**Figure 6.** Graph of an eigenfunction associated with  $\tilde{\lambda}(1, 1) = -8.65\dots$ . This function also represents the first eigenfunction of the stability operator.



**Figure 7.** Two solutions of the equation  $K_{J,1}[z] - 8.53078z = 0$ .


$K_{J,2}$ , this multiplicity one needs to be multiply by 3 because the eigenvalue  $\alpha_2 = 2$  of the Laplace operator on  $\mathbb{S}^2$  has multiplicity 3. The next eigenvalues are six near  $-2.5596$ , they are either two with multiplicity 3 or one with multiplicity 6. The last negative eigenvalue is one near  $-1.17496$ .

### Acknowledgement

The author would like to thank Andrés Rivera, Bruce Solomon and Nelson Castañeda for their useful comments and suggestions.

### References

- [Alfías and Piccione 2013] L. J. Alfías and P. Piccione, “Bifurcation of constant mean curvature tori in Euclidean spheres”, *J. Geom. Anal.* **23**:2 (2013), 677–708. [MR](#) [Zbl](#)
- [Beekmann and Lökés 1990] B. Beekmann and H. Lökés, “Estimates for the eigenvalues of Hill’s equation and applications for the eigenvalues of the Laplacian on toroidal surfaces”, *Manuscripta Math.* **68**:3 (1990), 295–308. [MR](#) [Zbl](#)



**Figure 8.** Graph of the discriminant of the operator  $K_{J,2}$ .

- [Choi and Wang 1983] H. I. Choi and A. N. Wang, “A first eigenvalue estimate for minimal hypersurfaces”, *J. Differential Geom.* **18**:3 (1983), 559–562. [MR](#) [Zbl](#)
- [Haupt 1914] O. Haupt, “Über eine Methode zum Beweise von Oszillationstheoremen”, *Math. Ann.* **76**:1 (1914), 67–104. [MR](#) [Zbl](#)
- [Magnus and Winkler 1966] W. Magnus and S. Winkler, *Hill’s equation*, Interscience Tracts in Pure and Applied Mathematics **20**, Wiley, New York, 1966. [MR](#) [Zbl](#)
- [Marques and Neves 2014] F. C. Marques and A. Neves, “Min-max theory and the Willmore conjecture”, *Ann. of Math. (2)* **179**:2 (2014), 683–782. [MR](#) [Zbl](#)
- [Marrocos and Gomes 2019] M. A. M. Marrocos and J. N. V. Gomes, “Generic spectrum of warped products and  $G$ -manifolds”, *J. Geom. Anal.* **29**:4 (2019), 3124–3134. [MR](#) [Zbl](#)
- [Otsuki 1972] T. Otsuki, “On integral inequalities related with a certain nonlinear differential equation”, *Proc. Japan Acad.* **48** (1972), 9–12. [MR](#) [Zbl](#)
- [Otsuki 1993] T. Otsuki, “On a differential equation related with differential geometry”, *Mem. Fac. Sci. Kyushu Univ. Ser. A* **47**:2 (1993), 245–281. [MR](#) [Zbl](#)
- [Perdomo 2001] O. Perdomo, “Low index minimal hypersurfaces of spheres”, *Asian J. Math.* **5**:4 (2001), 741–749. [MR](#) [Zbl](#)
- [Perdomo 2002] O. Perdomo, “First stability eigenvalue characterization of Clifford hypersurfaces”, *Proc. Amer. Math. Soc.* **130**:11 (2002), 3379–3384. [MR](#) [Zbl](#)
- [Perdomo 2004a] O. Perdomo, “On the average of the scalar curvature of minimal hypersurfaces of spheres with low stability index”, *Illinois J. Math.* **48**:2 (2004), 559–565. [MR](#) [Zbl](#)
- [Perdomo 2004b] O. Perdomo, “Ridigity of minimal hypersurfaces of spheres with two principal curvatures”, *Arch. Math. (Basel)* **82**:2 (2004), 180–184. [MR](#) [Zbl](#)
- [Perdomo 2010] O. M. Perdomo, “Embedded constant mean curvature hypersurfaces on spheres”, *Asian J. Math.* **14**:1 (2010), 73–108. [MR](#) [Zbl](#)
- [Rossman and Sultana 2007] W. Rossman and N. Sultana, “Morse index of constant mean curvature tori of revolution in the 3-sphere”, *Illinois J. Math.* **51**:4 (2007), 1329–1340. [MR](#) [Zbl](#)
- [Savo 2010] A. Savo, “Index bounds for minimal hypersurfaces of the sphere”, *Indiana Univ. Math. J.* **59**:3 (2010), 823–837. [MR](#) [Zbl](#)
- [Simons 1968] J. Simons, “Minimal varieties in riemannian manifolds”, *Ann. of Math. (2)* **88** (1968), 62–105. [MR](#) [Zbl](#)

- [Solomon 1990a] B. Solomon, “The harmonic analysis of cubic isoparametric minimal hypersurfaces, I: Dimensions 3 and 6”, *Amer. J. Math.* **112**:2 (1990), 157–203. [MR](#) [Zbl](#)
- [Solomon 1990b] B. Solomon, “The harmonic analysis of cubic isoparametric minimal hypersurfaces, II: Dimensions 12 and 24”, *Amer. J. Math.* **112**:2 (1990), 205–241. [MR](#) [Zbl](#)
- [Urbano 1990] F. Urbano, “Minimal surfaces with low index in the three-dimensional sphere”, *Proc. Amer. Math. Soc.* **108**:4 (1990), 989–992. [MR](#) [Zbl](#)
- [Yau 1982] S. T. Yau, “Problem section”, pp. 669–706 in *Seminar on Differential Geometry*, edited by S. T. Yau, Ann. of Math. Stud. **102**, Princeton University Press, 1982. [MR](#) [Zbl](#)

Received October 5, 2019.

OSCAR M. PERDOMO  
DEPARTMENT OF MATHEMATICAL SCIENCES  
CENTRAL CONNECTICUT STATE UNIVERSITY  
NEW BRITAIN, CT  
UNITED STATES  
[perdomoosm@ccsu.edu](mailto:perdomoosm@ccsu.edu)

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Wee Teck Gan  
Mathematics Department  
National University of Singapore  
Singapore 119076  
[matgwt@nus.edu.sg](mailto:matgwt@nus.edu.sg)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Matthias Aschenbrenner  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[matthias@math.ucla.edu](mailto:matthias@math.ucla.edu)

Daryl Cooper  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[cooper@math.ucsb.edu](mailto:cooper@math.ucsb.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 308    No. 2    October 2020

---

Existence of steady multiple vortex patches to the vortex-wave system	257
DAOMIN CAO and GUODONG WANG	
Relations of rationality for special values of Rankin–Selberg $L$ -functions of $GL_n \times GL_m$ over CM-fields	281
HARALD GROBNER and GUNJA SACHDEVA	
A bound for the conductor of an open subgroup of $GL_2$ associated to an elliptic curve	307
NATHAN JONES	
Topology of complexity one quotients	333
Yael KARSHON and SUSAN TOLMAN	
Flag Bott manifolds and the toric closure of a generic orbit associated to a generalized Bott manifold	347
SHINTARÔ KUROKI, EUNJEONG LEE, JONGBAEK SONG and DONG YOUP SUH	
Projective cases for the restriction of the oscillator representation to dual pairs of type I	393
SABINE J. LANG	
A remark on a trace Paley–Wiener theorem	407
GORAN MUIĆ	
Spectrum of the Laplacian and the Jacobi operator on rotational CMC hypersurfaces of spheres	419
OSCAR M. PERDOMO	
Mean curvature flow in a Riemannian manifold endowed with a Killing vector field	435
LIANGJUN WENG	
Green correspondence and relative projectivity for pairs of adjoint functors between triangulated categories	473
ALEXANDER ZIMMERMANN	