

*Pacific
Journal of
Mathematics*

**THE FIRST NONZERO EIGENVALUE
OF THE p -LAPLACIAN ON DIFFERENTIAL FORMS**

SHOO SETO

Volume 309 No. 1

November 2020

THE FIRST NONZERO EIGENVALUE OF THE p -LAPLACIAN ON DIFFERENTIAL FORMS

SHOO SETO

We introduce a generalization of the p -Laplace operator to act on differential forms and generalize an estimate of Gallot and Meyer (1973) for the first nonzero eigenvalue on closed Riemannian manifolds.

1. Introduction

Let (M, g) be an n -dimensional closed Riemannian manifold. Motivated from the variational characterization of the Laplacian eigenvalue problem, we define the L^p -Dirichlet integral on k -forms (introduced in [Scott 1995]) by

$$(1) \quad \mathcal{F}[\alpha] := \int_M \|d\alpha\|^p + \|d^*\alpha\|^p, \quad \alpha \in \Omega^k(M),$$

where d^* is the L^2 -adjoint of the exterior derivative d . Note that $\mathcal{F}[\alpha] = 0$ if and only if $\alpha \in \mathcal{H}^k(M)$, that is, the minimum is zero and is attained for harmonic k -forms, i.e., $\alpha \in \ker(d) \cap \ker(d^*)$. For a nonzero infimum we consider the space

$$(2) \quad A_k := \left\{ \alpha \in \mathcal{W}^{1,p}(\Omega^k(M)) \mid \int_M \|\alpha\|^p = 1, \int_M \|\alpha\|^{p-2} \langle \alpha, \omega \rangle = 0, \omega \in \mathcal{H}^k(M) \right\},$$

where the space $\mathcal{W}^{1,p}(\Omega^k(M))$ is the $(1, p)$ -Sobolev space of differential k -forms defined in [Scott 1995]. See Section 3 for the precise definition. Computing the Euler–Lagrange equation leads us to the defining the following operator:

Definition 1.1 (p -Hodge Laplacian).

$$(3) \quad \Delta_p \alpha := d^*(\|d\alpha\|^{p-2} d\alpha) + d(\|d^*\alpha\|^{p-2} d^*\alpha), \quad \alpha \in \Omega^k(M).$$

When $p = 2$, this becomes the usual Hodge Laplacian. For $p \neq 2$ and $\alpha \in C^\infty(M)$ Δ_p becomes the usual p -Laplacian. The corresponding eigenvalue equation is given by

$$(4) \quad \Delta_p \alpha = \lambda \|\alpha\|^{p-2} \alpha, \quad \alpha \in \Omega^k(M)$$

MSC2010: 47J10, 53C65.

Keywords: differential forms, Hodge Laplacian, p -Laplacian, Weitzenböck curvature.

and the variational principle tells us that

$$\lambda_1 = \inf\{\mathcal{F}[\alpha] \mid \alpha \in A_k\}.$$

See Section 3 for details. When $p = 2$, there is much work on the spectrum of the Hodge Laplacian acting on differential forms. Among many others, we point out the work of Gallot and Meyer [1973; 1975] who show an estimate of the first eigenvalue using bounds from the Weitzenböck curvature on compact Riemannian manifolds. For manifolds with boundary, among many others, see the works [Kwong 2016; Savo 2009; Raulot and Savo 2011].

For $p \neq 2$, the p -Laplace eigenvalue problem on 0-forms (functions) has attracted much attention. See notes by Lindqvist [2006] for a general reference on the p -Laplace equation. For estimates on the first eigenvalue relating to the curvature, among many other works, see [Matei 2000; Naber and Valtorta 2014; Seto and Wei 2017] for eigenvalue estimates with $\text{Ric} \geq K$, $K \in \mathbb{R}$.

Remark 1.2. There is also a related notion of p -harmonic k -forms which looks at the minimizer in a cohomology class of k -forms with finite L^p -norm, i.e.,

$$\inf_{\alpha \in H_d^k(M)} \int_M \|\alpha\|^p.$$

The critical point of the variation leads to the following definition of p -harmonic, for closed k -forms α , if

$$d_p^* := d^*(\|\alpha\|^{p-2}\alpha) = 0$$

then α is p -harmonic. See [Dung 2017].

In this paper we prove the following lower bound estimate for the first eigenvalue:

Theorem 1.3. *Let M^n be a closed Riemannian manifold with the eigenvalues of the curvature operator bounded below by $H \in \mathbb{R}$ and $p \geq 2$. Then*

$$\lambda_1 \geq \left(\frac{k(n-k)}{2^{(2/p)-1}(C+(p-2)/2)} H \right)^{p/2},$$

where

$$C = \max\left\{ \frac{k}{k+1}, \frac{n-k}{n-k+1} \right\}.$$

Remark 1.4. When $p = 2$, the above recovers the estimate due to Gallot and Meyer [1973] (see also [Gallot and Meyer 1975]), for $1 \leq k \leq \frac{n}{2}$,

$$\lambda_1 \geq k(n-k+1)H.$$

The organization of this paper is as follows. In Section 2 we review some known estimates for differential k -forms. In Section 3 we show that the infimum can be characterized as an eigenvalue problem. In Section 4 we give the main estimate. In

Section 5 we give a brief discussion on boundary conditions for differential forms and possible future directions.

2. Some estimates on $\Omega^k(M)$

We first recall the Weitzenböck curvature

Definition 2.1. Let $p \in M$ and let $\{E_i\}_{i=1}^n$ be an orthonormal frame at p . Then for $\alpha \in \Omega^k(M)$, define the Weitzenböck curvature W_k by

$$W_k(\alpha)(X_1, \dots, X_k) := \sum (R(E_j, X_i)\alpha)(X_1, \dots, E_j, \dots, X_k).$$

Note that on 1-forms, this is simply the Ricci tensor.

If the eigenvalues of the curvature operator are bounded by $H \in \mathbb{R}$, we can show that

$$(5) \quad (W_k(\alpha), \alpha) \geq k(n-k)H\|\alpha\|^2.$$

The Weitzenböck curvature appears in the main tool we use in obtaining our estimate, which is the Bochner–Weitzenböck formula for k -forms:

$$(6) \quad \frac{1}{2}\Delta\|\alpha\|^2 = (\Delta\alpha, \alpha) - \|\nabla\alpha\|^2 - (W_k(\alpha), \alpha),$$

where $\Delta := \Delta_2 = dd^* + d^*d$. Note that for exact 1-form $\alpha = df$, since $\nabla df = \text{Hess } f$, the usual Cauchy–Schwarz inequality will give us an estimate on the middle term. For k -forms, we will need the following:

Lemma 2.2 [Gallot and Meyer 1973]. *Let $\alpha \in \Omega^k(M)$, $1 \leq k \leq n-1$. Then*

$$(7) \quad \|\nabla\alpha\|^2 \geq \frac{1}{k+1}\|d\alpha\|^2 + \frac{1}{n-k+1}\|d^*\alpha\|^2.$$

We give a proof for completeness. The proof we give is in the context of conformal Killing forms and can be found in various sources, for instance, [Moroianu and Semmelmann 2003].

Proof. Consider the two linear maps

$$\iota : TM \otimes \Omega^k(M) \rightarrow \Omega^{k-1}(M)$$

$$\iota(v, \alpha) = \iota_v\alpha$$

and

$$\wedge : \Omega^1(M) \otimes \Omega^k(M) \rightarrow \Omega^{k+1}(M)$$

$$\wedge(\beta, \alpha) = \beta \wedge \alpha.$$

Let ι^* and \wedge^* be their metric adjoint. Then

$$\wedge \circ \iota^*(\alpha) = 0 \quad \text{and} \quad \iota \circ \wedge^*(\alpha) = 0,$$

so that we get the decomposition

$$TM \otimes \Omega^k(M) \simeq \text{im}(\iota^*) \oplus \text{im}(\wedge^*) \oplus Y,$$

where Y is the orthogonal complement. By direct computation, for $\alpha \in \Omega^k(M)$, we have

$$\iota \circ \iota^*(\alpha) = (n - k + 1)\alpha \quad \text{and} \quad \wedge \circ \wedge^*(\alpha) = (k + 1)\alpha.$$

Viewing $\nabla\alpha \in \Gamma(TM \otimes \Omega^k(M))$, From the decomposition,

$$\nabla\alpha = \iota^*\beta + \wedge^*\gamma + \delta,$$

applying ι , we have

$$\iota\nabla\alpha = (n - k + 1)\beta.$$

So the projection operator onto $\text{im}(\iota^*)$ is given by

$$\pi_{\iota^*}\nabla\alpha = \frac{1}{n - k + 1}\iota^*\iota\nabla\alpha$$

and similarly

$$\pi_{\wedge^*}\nabla\alpha = \frac{1}{k + 1}\wedge^*\wedge\nabla\alpha.$$

Let $T\alpha := \pi_T(\nabla\alpha)$ the projection onto the orthogonal complement space. Since

$$d\alpha = \wedge(\nabla\alpha) \quad \text{and} \quad d^*\alpha = -\iota(\nabla\alpha),$$

we have the decomposition

$$T\alpha(X) = \nabla_X\alpha - \frac{1}{k + 1}\iota_X d\alpha + \frac{1}{n - k + 1}X^* \wedge d^*\alpha$$

and taking the norm gives us

$$\|\nabla\alpha\|^2 = \|T\alpha\|^2 + \frac{1}{k + 1}\|d\alpha\|^2 + \frac{1}{n - k + 1}\|d^*\alpha\|^2,$$

which implies (7). □

Remark 2.3. The projection operator T defined above is called the twistor operator and a form $\alpha \in \Omega^k(M)$ is called a conformal Killing form if $T\alpha = 0$.

The following lemma was pointed out by N.T. Dung and gives us a way to control the interior product by using an orthogonal decomposition of forms as the image under an interior product.

Lemma 2.4 [Dung and Sung 2019, Lemma 3.5]. *Let $V \in TM$, $\alpha \in \Omega^{k+1}$, $\beta \in \Omega^k$. Then*

$$|\langle \iota_V\alpha, \beta \rangle| \leq \|V\| \|\alpha\| \|\beta\|.$$

3. Variational characterization of the eigenvalue

In this section we will compute the Euler–Lagrange equation of (1) and show that the extremal problem can be reformulated as an eigenvalue problem. Analogous to the 0-form (function) case, we will look at weak solutions lying the $(1, p)$ -Sobolev space of differential k -forms first defined by Scott [1995] as

$$\mathcal{W}^{1,p}(\Omega^k(M)) := \{ \alpha \in W(\Omega^k(M)) \mid \alpha, d\alpha, d^*\alpha \in L^p(\Omega^*(M)) \},$$

where $W(\Omega^k(M))$ is the classical Sobolev space of k -forms, i.e., α is locally integrable and admits a generalized gradient.

Definition 3.1. We say that λ is an eigenvalue, if there exists a k -form $\alpha \in \mathcal{W}^{1,p}(\Omega^k(M))$ such that

$$\int_M \|d\alpha\|^{p-2} \langle d\alpha, d\beta \rangle + \int_M \|d^*\alpha\|^{p-2} \langle d^*\alpha, d^*\beta \rangle = \lambda \int_M \|\alpha\|^{p-2} \langle \alpha, \beta \rangle,$$

for any $\beta \in C^\infty(\Omega^k(M))$.

We will show the first nonzero eigenvalue λ_1 can be characterized as the infimum of the L^p -Dirichlet energy over the space A_k given in (2).

Proposition 3.2. For closed manifolds M and $p \geq 2$,

$$\lambda_1 = \inf \left\{ \int_M \|d\alpha\|^p + \|d^*\alpha\|^p \mid \alpha \in A_k \right\}.$$

Proof. Let ω be a fixed harmonic form and let $\beta(t) \in A$ for small $t > 0$ such that $\beta(0) = \alpha$. Computing the first variation of (1), we have

$$\begin{aligned} \left. \frac{d}{dt} \mathcal{F}[\beta(t)] \right|_{t=0} &= p \int_M \|d\alpha\|^{p-2} \langle d\alpha, d\beta'(0) \rangle + \|d^*\alpha\|^{p-2} \langle d^*\alpha, d^*\beta'(0) \rangle \\ &= p \int_M \langle \Delta_p \alpha, \beta'(0) \rangle. \end{aligned}$$

Next we compute the variation of the constraints so that

$$\left. \frac{d}{dt} \int_M \|\beta\|^p \right|_{t=0} = p \int_M \|\alpha\|^{p-2} \langle \alpha, \beta'(0) \rangle$$

and

$$\left. \frac{d}{dt} \int_M \|\beta\|^{p-2} \langle \beta, \omega \rangle \right|_{t=0} = (p-2) \int_M \|\alpha\|^{p-4} \langle \alpha, \beta'(0) \rangle \langle \alpha, \omega \rangle + \|\alpha\|^{p-2} \langle \beta'(0), \omega \rangle.$$

By the Lagrange multiplier method, there must be some λ and μ such that for $\beta \in \Omega^k(M)$,

$$\int_M \langle \Delta_p \alpha, \beta \rangle = \lambda \int_M \|\alpha\|^{p-2} \langle \alpha, \beta \rangle + \mu \int_M \|\alpha\|^{p-4} \langle \alpha, \beta \rangle \langle \alpha, \omega \rangle + \|\alpha\|^{p-2} \langle \beta, \omega \rangle.$$

Setting $\beta = \omega$, we have

$$0 = \mu \int_M \|\alpha\|^{p-4} \langle \alpha, \omega \rangle^2 + \|\alpha\|^{p-2} \|\omega\|^2$$

so that $\mu = 0$. Therefore,

$$\Delta_p \alpha = \lambda \|\alpha\|^{p-2} \alpha. \quad \square$$

4. Proof of Theorem 1.3

We will consider the following integral

$$\int_M \langle \Delta_p \alpha, \Delta \alpha \rangle = \int_M \langle \Delta_p \alpha, dd^* \alpha \rangle + \int_M \langle \Delta_p, d^* d \alpha \rangle.$$

Let $\alpha \in \Omega^k(M)$ be an eigenform satisfying (4). Then

$$\begin{aligned} (8) \quad \int_M \langle \Delta_p \alpha, d^* d \alpha \rangle &= \lambda \int_M \|\alpha\|^{p-2} \langle \alpha, d^* d \alpha \rangle \\ &= \lambda \int_M \langle d(\|\alpha\|^{p-2} \alpha), d \alpha \rangle \\ &= \lambda \int_M \langle d(\|\alpha\|^{p-2}) \wedge \alpha, d \alpha \rangle + \lambda \int_M \|\alpha\|^{p-2} \|d \alpha\|^2 \end{aligned}$$

and

$$\begin{aligned} (9) \quad \int_M \langle \Delta_p \alpha, dd^* \alpha \rangle &= \lambda \int_M \|\alpha\|^{p-2} \langle \alpha, dd^* \alpha \rangle \\ &= \lambda \int_M \langle d^*(\|\alpha\|^{p-2} \alpha), d^* \alpha \rangle \\ &= \lambda \int_M \|\alpha\|^{p-2} \|d^* \alpha\|^2 - \lambda \int_M \langle \iota_{\nabla \|\alpha\|^{p-2}} \alpha, d^* \alpha \rangle. \end{aligned}$$

On the other hand, by using the Bochner–Weitzenböck formula (6) we have

$$\begin{aligned} (10) \quad \int_M \langle \Delta_p \alpha, \Delta \alpha \rangle &= \lambda \int_M \|\alpha\|^{p-2} \langle \alpha, \Delta \alpha \rangle \\ &= \lambda \int_M ((p-2) \|\alpha\|^{p-2} |\nabla \|\alpha\|^2 + \|\alpha\|^{p-2} \|\nabla \alpha\|^2 + \|\alpha\|^{p-2} (W_k(\alpha), \alpha)). \end{aligned}$$

Combining (8), (9), and (10), we obtain

$$\begin{aligned} (11) \quad \int_M \langle d(\|\alpha\|^{p-2}) \wedge \alpha, d \alpha \rangle - \int_M \langle \iota_{\nabla \|\alpha\|^{p-2}} \alpha, d^* \alpha \rangle &+ \int_M \|\alpha\|^{p-2} \|d \alpha\|^2 \\ &+ \int_M \|\alpha\|^{p-2} \|d^* \alpha\|^2 \end{aligned}$$

$$(11 \text{ cont.}) = \int_M ((p-2)\|\alpha\|^{p-2}|\nabla\|\alpha\|^2 + \|\alpha\|^{p-2}\|\nabla\alpha\|^2 + \|\alpha\|^{p-2}(W_k(\alpha), \alpha)).$$

Using Lemma 2.4, the first term of (11) can be estimated as

$$\begin{aligned} \int_M \langle d(\|\alpha\|^{p-2}) \wedge \alpha, d\alpha \rangle &= \int_M \langle \alpha, \iota_{\nabla\|\alpha\|^{p-2}}(d\alpha) \rangle \\ &\leq \int_M \|\nabla\|\alpha\|^{p-2}\| \|d\alpha\| \|\alpha\| \\ &= (p-2) \int_M \|\alpha\|^{(p-2)/2} \|\nabla\|\alpha\| \| \|\alpha\|^{(p-2)/2} \|d\alpha\| \\ &\leq \frac{(p-2)}{2} \int_M \|\alpha\|^{p-2} \|\nabla\|\alpha\|^2 + \frac{(p-2)}{2} \int_M \|\alpha\|^{p-2} \|d\alpha\|^2 \end{aligned}$$

and similarly for the second term,

$$\begin{aligned} - \int_M \langle \iota_{\nabla\|\alpha\|^{p-2}}\alpha, d^*\alpha \rangle &\leq \int_M \|\nabla\|\alpha\|^{p-2}\|\alpha\| \|d^*\alpha\| \\ &= (p-2) \int_M \|\alpha\|^{(p-2)/2} \|\nabla\|\alpha\| \| \|\alpha\|^{(p-2)/2} \|d^*\alpha\| \\ &\leq \frac{(p-2)}{2} \int_M \|\alpha\|^{p-2} \|\nabla\|\alpha\|^2 + \frac{(p-2)}{2} \int_M \|\alpha\|^{p-2} \|d^*\alpha\|^2. \end{aligned}$$

Applying these estimates to (11), we get

$$\begin{aligned} &\frac{(p-2)+2}{2} \int_M \|\alpha\|^{p-2} \|d\alpha\|^2 + \frac{(p-2)+2}{2} \int_M \|\alpha\|^{p-2} \|d^*\alpha\|^2 \\ &\geq \int_M \|\alpha\|^{p-2} \|\nabla\alpha\|^2 + \int_M \|\alpha\|^{p-2} (W_k(\alpha), \alpha) \\ &\geq \frac{1}{k+1} \int_M \|\alpha\|^{p-2} \|d\alpha\|^2 + \frac{1}{n-k+1} \int_M \|\alpha\|^{p-2} \|d^*\alpha\|^2 + \int_M \|\alpha\|^{p-2} (W_k(\alpha), \alpha). \end{aligned}$$

Let

$$C := \max\left\{\frac{k}{k+1}, \frac{n-k}{n-k+1}\right\}.$$

Using

$$\int_M \|\alpha\|^{p-2} \|d\alpha\|^2 \leq \left(\int_M \|\alpha\|^p\right)^{1-2/p} \left(\int_M \|d\alpha\|^p\right)^{2/p}$$

and

$$\int_M \|\alpha\|^{p-2} \|d^*\alpha\|^2 \leq \left(\int_M \|\alpha\|^p\right)^{1-2/p} \left(\int_M \|d^*\alpha\|^p\right)^{2/p},$$

we have

$$\begin{aligned} \left(C + \frac{(p-2)}{2}\right) \left(\int_M \|\alpha\|^p\right)^{1-2/p} & \left[\left(\int_M \|d\alpha\|^p\right)^{2/p} + \left(\int_M \|d^*\alpha\|^p\right)^{2/p} \right] \\ & \geq \int_M \|\alpha\|^{p-2} (W_k(\alpha), \alpha). \end{aligned}$$

For $p \geq 2$, and using the lower bound of the Weitzenböck curvature (5), we have

$$\begin{aligned} 2^{(2/p)-1} \left(C + \frac{(p-2)}{2}\right) \left(\int_M \|\alpha\|^p\right)^{1-2/p} & \left(\int_M \|d\alpha\|^p + \|d^*\alpha\|^p\right)^{2/p} \\ & \geq k(n-k)H \int_M \|\alpha\|^p. \end{aligned}$$

Using the fact that $\int_M \|d\alpha\|^p + \|d^*\alpha\|^p = \lambda \int_M \|\alpha\|^p$ for eigenform α , we get

$$\lambda^{2/p} \geq \frac{k(n-k)}{2^{(2/p)-1}(C + (p-2)/2)}.$$

5. Boundary conditions

In this section we briefly discuss the situation of a compact manifold M with nonempty smooth boundary ∂M . Let n denote the unit outer normal vector and let $J : \partial M \rightarrow M$ be the inclusion. Then $J^*\alpha$ is the restriction of a form to the boundary. Then d and its adjoint d^* are related with an additional boundary term given by

$$\int_M \langle d\alpha, \beta \rangle = \int_M \langle \alpha, d^*\beta \rangle + \int_{\partial M} \langle J^*(\alpha), \iota_n \beta \rangle, \quad \alpha \in \Omega^k(M), \beta \in \Omega^{k+1}(M).$$

and the corresponding Green's formula for the p -Laplacian is

$$\begin{aligned} (\Delta_p \alpha, \beta) &= \int_M \|d\alpha\|^{p-2} \langle d\alpha, d\beta \rangle + \int_M \|d^*\alpha\|^{p-2} \langle d^*\alpha, d^*\beta \rangle \\ &\quad - \int_{\partial M} \langle \iota_n(\|d\alpha\|^{p-2} d\alpha), J^*(\beta) \rangle + \int_{\partial M} \langle \|d^*\alpha\|^{p-2} J^*(d^*\alpha), \iota_n \beta \rangle. \end{aligned}$$

The two most common boundary conditions for the classical Laplacian eigenvalue problem are the Dirichlet and Neumann boundary condition. For the Hodge Laplacian, the analogous boundary conditions are the absolute boundary condition

$$\begin{cases} \iota_n \alpha = 0, \\ \iota_n d\alpha = 0, \end{cases} \quad \text{on } \partial M$$

and the relative boundary condition

$$\begin{cases} J^*(\alpha) = 0, \\ J^*(d^*\alpha) = 0, \end{cases} \quad \text{on } \partial M.$$

The essential feature of the boundary condition is that if α satisfies either of the boundary conditions, then $\Delta_p \alpha = 0$ implies $d\alpha = 0$ and $d^* \alpha = 0$. The boundary terms that will be introduced to (11) are

$$\begin{aligned} & \int_M \langle d(\|\alpha\|^{p-2}) \wedge \alpha, d\alpha \rangle - \int_M \langle \iota_{\nabla \|\alpha\|^{p-2}} \alpha, d^* \alpha \rangle + \int_M \|\alpha\|^{p-2} \|d\alpha\|^2 \\ & + \int_M \|\alpha\|^{p-2} \|d^* \alpha\|^2 - \int_{\partial M} \|\alpha\|^{p-2} \langle J^*(\alpha), \iota_n(d\alpha) \rangle + \int_{\partial M} \|\alpha\|^{p-2} \langle J^*(d^* \alpha), \iota_n(\alpha) \rangle \\ & = \int_M ((p-2)\|\alpha\|^{p-2} |\nabla \|\alpha\|^2 + \|\alpha\|^{p-2} \|\nabla \alpha\|^2 + \|\alpha\|^{p-2} (W_k(\alpha), \alpha)). \end{aligned}$$

Since the boundary terms will vanish under either of the boundary conditions, we get the same estimate for the boundary value problem as well. It would be interesting to see what the Reilly formula, for instance a generalization of Theorem 3 in [Raulot and Savo 2011] would be in this context, however due to the asymmetry of the weight function in the p -Laplacian, it is not immediate what the appropriate Bochner–Weitzenböck type formula would be for Δ_p .

Acknowledgements

The author would like to thank Prof. Zhiqin Lu for very helpful discussions, suggestions and constant support, and Prof. Guofang Wei and Prof. Qi Zhang for discussions and encouragement on the problem. We also thank Prof. Nguyen Thac Dung for letting us know of an improvement on the constant, see Lemma 2.4.

References

- [Dung 2017] N. T. Dung, “ p -harmonic ℓ -forms on Riemannian manifolds with a weighted Poincaré inequality”, *Nonlinear Anal.* **150** (2017), 138–150. MR Zbl
- [Dung and Sung 2019] N. T. Dung and C.-J. A. Sung, “Analysis of weighted p -harmonic forms and applications”, *Int. J. Math.* **30**:11 (2019), art. id. 1950058. MR Zbl
- [Gallot and Meyer 1973] S. Gallot and D. Meyer, “Sur la première valeur propre du p -spectre pour les variétés à opérateur de courbure positif”, *C. R. Acad. Sci. Paris Sér. A-B* **276** (1973), A1619–A1621. MR Zbl
- [Gallot and Meyer 1975] S. Gallot and D. Meyer, “Opérateur de courbure et laplacien des formes différentielles d’une variété riemannienne”, *J. Math. Pures Appl.* (9) **54**:3 (1975), 259–284. MR Zbl
- [Kwong 2016] K.-K. Kwong, “Some sharp Hodge Laplacian and Steklov eigenvalue estimates for differential forms”, *Calc. Var. Partial Differential Equations* **55**:2 (2016), art. id. 38. MR Zbl
- [Lindqvist 2006] P. Lindqvist, *Notes on the p -Laplace equation*, Report. Univ. Jyväskylä Dept. Math. Stat. **102**, University of Jyväskylä, 2006. MR Zbl
- [Matei 2000] A.-M. Matei, “First eigenvalue for the p -Laplace operator”, *Nonlinear Anal.* **39**:8 (2000), 1051–1068. MR Zbl
- [Moroianu and Semmelmann 2003] A. Moroianu and U. Semmelmann, “Twistor forms on Kähler manifolds”, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) **2**:4 (2003), 823–845. MR Zbl

- [Naber and Valtorta 2014] A. Naber and D. Valtorta, “Sharp estimates on the first eigenvalue of the p -Laplacian with negative Ricci lower bound”, *Math. Z.* **277**:3-4 (2014), 867–891. MR Zbl
- [Raulot and Savo 2011] S. Raulot and A. Savo, “A Reilly formula and eigenvalue estimates for differential forms”, *J. Geom. Anal.* **21**:3 (2011), 620–640. MR Zbl
- [Savo 2009] A. Savo, “On the lowest eigenvalue of the Hodge Laplacian on compact, negatively curved domains”, *Ann. Global Anal. Geom.* **35**:1 (2009), 39–62. MR Zbl
- [Scott 1995] C. Scott, “ L^p theory of differential forms on manifolds”, *Trans. Amer. Math. Soc.* **347**:6 (1995), 2075–2096. MR Zbl
- [Seto and Wei 2017] S. Seto and G. Wei, “First eigenvalue of the p -Laplacian under integral curvature condition”, *Nonlinear Anal.* **163** (2017), 60–70. MR Zbl

Received April 2, 2019. Revised December 19, 2019.

SHOO SETO
DEPARTMENT OF MATHEMATICS
CALIFORNIA STATE UNIVERSITY, FULLERTON
FULLERTON, CA
UNITED STATES
shoseto@fullerton.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing


<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 309 No. 1 November 2020

- Lie 2-algebras of vector fields 1
DANIEL BERWICK-EVANS and EUGENE LERMAN
- Lower regularity solutions of the biharmonic Schrödinger equation in a quarter plane 35
ROBERTO DE A. CAPISTRANO-FILHO, MÁRCIO CAVALCANTE
and FERNANDO A. GALLEGO
- The arithmetic Hodge index theorem and rigidity of dynamical systems over function fields 71
ALEXANDER CARNEY
- On the vanishing of the theta invariant and a conjecture of Huneke and Wiegand 103
OLGUR CELIKBAS
- Algebraic and geometric properties of flag Bott–Samelson varieties and applications to representations 145
NAOKI FUJITA, EUNJEONG LEE and DONG YOUP SUH
- On a modular form of Zaremba’s conjecture 195
NIKOLAY G. MOSHCHEVITIN and ILYA D. SHKREDOV
- The first nonzero eigenvalue of the p -Laplacian on differential forms 213
SHOO SETO
- Global regularity of the Navier–Stokes equations on 3D periodic thin domain with large data 223
NA ZHAO



0030-8730(202011)309:1;1-J