

*Pacific
Journal of
Mathematics*

**THIN SUBGROUPS ISOMORPHIC TO
GROMOV-PIATETSKI-SHAPIRO LATTICES**

SAMUEL A. BALLAS

THIN SUBGROUPS ISOMORPHIC TO GROMOV–PIATETSKI-SHAPIRO LATTICES

SAMUEL A. BALLAS

In this paper we produce many examples of thin subgroups of special linear groups that are isomorphic to the fundamental group nonarithmetic hyperbolic manifolds. Specifically, we show that the nonarithmetic lattices in $\mathrm{SO}(n, 1, \mathbb{R})$ constructed by Gromov and Piatetski-Shapiro can be embedded into $\mathrm{SL}(n + 1, \mathbb{R})$ so that their images are thin subgroups.

Introduction

Let G be a semisimple Lie group and let Γ be a finitely generated subgroup. We say that Γ is a *thin subgroup* of G if there is a lattice $\Lambda \subset G$ containing Γ such that

- Γ has infinite index in Λ ,
- Γ is Zariski dense in G .

Intuitively, such groups are very sparse in the sense that they have infinite index in a lattice, but at the same time are dense in an algebraic sense. Note, that if one relaxes the first condition above, then Γ would be a lattice, so another way of thinking of thin groups is as infinite index analogues of lattices in semisimple Lie groups.

Over the last several years, thin groups have been the subject of much research, much of which has been motivated by the observation that many theorems and conjectures in number theory can be phrased in terms of counting primes in orbits of groups that are “abelian analogues of thin groups.” Here are two examples. First, let $G = \mathbb{R}$, $b, m \in \mathbb{N}$ such that $(b, m) = 1$, $\Delta = \mathbb{Z}$ and $\Gamma = m\mathbb{Z}$. The orbit $b + \Gamma$ is an arithmetic progression and Dirichlet’s theorem on primes in arithmetic progressions is equivalent to this orbit containing infinitely many primes. Next, let $G = \mathbb{R}^2$, $\Delta = \mathbb{Z}^2$, $\Gamma = \langle (1, 1) \rangle$ and $b = (1, 3) \in \mathbb{Z}^2$. The orbit $b + \Gamma = \{(m, m + 2) \mid m \in \mathbb{Z}\}$ and the twin prime conjecture is equivalent to the statement that this orbit contains infinitely many points whose components are both prime. Note that in the first case Γ is a lattice in G , but in the second case Γ has infinite index in Δ and is an analogue of a thin group (sans Zariski density) in G .

MSC2010: 22E40, 57M50.

Keywords: thin groups, nonarithmetic lattices.

This orbital perspective was used by Brun to attack the twin primes conjecture using “combinatorial sieving” techniques. Although the full conjecture remains unproven these techniques did yield some powerful results. For instance, using these methods, Chen [1978] was able to prove that there are infinitely many pairs n and $n + 2$ such that one is prime and the other is the product of at most 2 primes. More details of this perspective are explained in the excellent surveys of Bourgain [2014] and Lubotzky [2012].

Inspired by these results, Bourgain, Gamburd, and Sarnak [Bourgain et al. 2010] developed complementary “affine sieving” techniques to analyze thin group orbits. In this context, the thinness property of the group gives enough control of orbits to execute these counting arguments. Again, much of this is described in Lubotzky’s survey [2012].

Given these connections it is desirable to produce examples of thin groups and understand what types of groups are thin. Presently, there are many constructions of thin groups. For instance, in recent work of Fuchs and Rivin [2017] it is shown that if one “randomly” selects two matrices in $SL(n, \mathbb{Z})$ then with high probability, the group they generate is a thin subgroup of $SL(n, \mathbb{R})$. However, the groups constructed in this way are almost always free groups. There are also several constructions that allow one to produce thin subgroups isomorphic to fundamental groups of closed surfaces in a variety of algebraic groups (see [Cooper and Futer 2019; Kahn et al. 2018; Kahn and Markovic 2012; Kahn and Wright 2018], for instance). Given these examples one may ask which isomorphism classes of groups are thin? More precisely, if G is a semisimple Lie group and H is an abstract finitely generated group then we say that H can be *realized as a thin subgroup of G* if there is an embedding $\iota : H \rightarrow G$ whose image is a thin subgroup of G . With this definition in hand we can rephrase the previous question as: given a semisimple algebraic group G , what isomorphism types of groups can be realized as thin subgroups of G ? Recent work of the author and D. Long [Ballas and Long 2020] shows that there are many additional isomorphism types of groups that can arise as thin subgroups of special linear groups. More precisely, in [Ballas and Long 2020] it is shown that fundamental groups of arithmetic hyperbolic n -manifolds of “orthogonal type” can be realized as thin subgroups. In the present work, we extend the techniques of [Ballas and Long 2020] to produce infinitely many examples of nonarithmetic hyperbolic n -manifolds whose fundamental groups can be realized as thin subgroups of $SL_{n+1}(\mathbb{R})$. Our main result is:

Theorem 1. *For each $n \geq 3$, there is an infinite collection C_n of nonarithmetic hyperbolic n -manifolds with the property that if $M^n \in C_n$ then $\pi_1(M)$ can be realized as a thin subgroup of $SL_{n+1}(\mathbb{R})$. Furthermore, the collection C_n contains representatives from infinitely many commensurability classes of both compact and noncompact manifolds.*

It should be noted that the collection \mathcal{C}_n appearing in [Theorem 1](#) can be described fairly explicitly, and roughly speaking consists of the hyperbolic manifolds coming from the nonarithmetic lattices in $\mathrm{SO}(n, 1, \mathbb{R})$ constructed by Gromov–Piatetski-Shapiro in [[Gromov and Piatetski-Shapiro 1988](#)].

Outline of paper. In [Section 1](#) we recall the Gromov–Piatetski-Shapiro construction of nonarithmetic lattices in $\mathrm{SO}(n, 1, \mathbb{R})$ and define the collection \mathcal{C}_n appearing in [Theorem 1](#). In [Section 2](#) we show that the fundamental group of any element of \mathcal{C}_n can be embedded in several lattices in $\mathrm{SL}_{n+1}(\mathbb{R})$. Finally, in [Section 3](#) we prove [Theorem 1](#) by showing that the images of the previously mentioned embeddings are thin subgroups.

1. Gromov–Piatetski-Shapiro lattices

Gromov and Piatetski-Shapiro [[1988](#)], describe a method for constructing infinitely many nonarithmetic lattices in $\mathrm{SO}(n, 1, \mathbb{R})$. In this section we describe their construction and the construction of the lattices appearing in [Theorem 1](#).

Let K be a totally real number field of degree $d + 1$ with ring of integers \mathcal{O}_K . There are $d + 1$ embeddings $\{\sigma_0, \dots, \sigma_d\}$ of K into \mathbb{R} . Using the embedding σ_0 we will implicitly regard K as a subset of \mathbb{R} . In this way, it makes sense to say that elements of F are positive or negative. Let $s_K : K^\times \rightarrow \mathbb{Z}_{\geq 0}$, where $s_K(a) = |\{i \geq 1 \mid \sigma_i(a) > 0\}|$. In other words, $s_K(a)$ counts the nonidentity embeddings for which a has positive image.

Next, let $\alpha, \beta, a_2, \dots, a_{n+1} \in \mathcal{O}_K$ be positive elements such that

- β/α is not a square in K ,
- $s_K(\alpha) = s_K(\beta) = s_K(\alpha_i) = d$ for $1 \leq i \leq n$,
- $s_K(a_{n+1}) = 0$.

Next, define quadratic forms

$$(1-1) \quad J_1 = \alpha x_1^2 + \sum_{i=2}^n a_i x_i^2 - a_{n+1} x_{n+1}^2, \quad J_2 = \beta x_1^2 + \sum_{i=2}^n a_i x_i^2 - a_{n+1} x_{n+1}^2$$

If $A \subset \mathbb{R}$ is a subring containing 1 then we define

$$\mathrm{SO}(J_i, A) = \{B \in \mathrm{SL}_{n+1}(A) \mid J_i(Bv) = J_i(v) \forall v \in \mathbb{R}^{n+1}\}.$$

Using this notation, define $\Gamma_1 = \mathrm{SO}(J_1, \mathcal{O}_K)$ and $\Gamma_2 = h \mathrm{SO}(J_2, \mathcal{O}_K) h^{-1}$, where $h = \mathrm{Diag}(\sqrt{\beta/\alpha}, \dots, 1)$. Note that both Γ_1 and Γ_2 are lattices in $\mathrm{SO}(J_1, \mathbb{R})$, however, since β/α is not a square in K it follows from [[Gromov and Piatetski-Shapiro 1988](#), see [Corollary 2.7](#) and [§2.9](#)] that these lattices are not commensurable.

There is a model for hyperbolic n -space given by

$$\mathbb{H}^n = \{v \in \mathbb{R}^{n+1} \mid J_1(v) = -1, v_{n+1} > 0\}.$$

The identity component $\mathrm{SO}(J_1, \mathbb{R})^\circ$ of $\mathrm{SO}(J_1, \mathbb{R})$ consists of the orientation preserving isometries of \mathbb{H}^n (see [Ratcliffe 2006, §3.2] for details). By passing to finite index subgroups we can assume that $\Gamma_i \subset \mathrm{SO}(J_1, \mathbb{R})^\circ$, and so \mathbb{H}^n/Γ_i is a finite volume hyperbolic orbifold for $i = 1, 2$.

The lattice $\Gamma_2 \subset \mathrm{SO}(J_1, L)$, where $L = K(\sqrt{\beta/\alpha})$. Note that because α and β are positive and $s_K(\alpha) = s_K(\beta) = d$ it follows that L is also totally real. Furthermore, for every $\gamma \in \Gamma_2$, $\mathrm{tr}(\gamma) \in \mathcal{O}_K \subset \mathcal{O}_L$. The following lemma then shows that by passing to a subgroup of finite index we may assume that $\Gamma_2 \subset \mathrm{SO}(J_1, \mathcal{O}_L)$. This result seems well known to experts, but we include a proof for the sake of completeness.

Lemma 1.1. *Let $k \subset \mathbb{C}$ be a number field and let \mathcal{O}_k be the ring of integers of k . If $\Gamma \subset \mathrm{GL}_n(k)$ acts irreducibly on \mathbb{C}^n and has the property that $\mathrm{tr}(\gamma) \in \mathcal{O}_k$ for each $\gamma \in \Gamma$ then there is a finite index subgroup $\Gamma' \subset \Gamma$ such that $\Gamma' \subset \mathrm{GL}_n(\mathcal{O}_k)$.*

Proof. If $A \subset k$ is a subring then let $A\Gamma = \{\sum_i a_i \gamma_i \mid a_i \in A, \gamma_i \in \Gamma\}$. Note that in this definition all sums have finitely many terms. By [Bass 1980, Proposition 2.2], $\mathcal{O}_k\Gamma$ is an order in the central simple algebra $k\Gamma$. The order $\mathcal{O}_k\Gamma$ is contained in some maximal order \mathcal{D} in $M_n(k)$ ($n \times n$ matrices over k). Let $\mathcal{D}^1 \subset \mathrm{SL}_n(k)$ be the norm 1 elements of \mathcal{D} . Then $M_n(\mathcal{O}_k)$ is also an order in $M_n(k)$ whose group of norm 1 elements is $\mathrm{SL}_n(\mathcal{O}_k)$. It is a standard result using restriction of scalars that groups of norm 1 elements in maximal orders of $M_n(k)$ are commensurable. Roughly speaking this is a consequence of the fact that the intersection of two orders is again an order and the unit groups of these orders are irreducible lattices in $\mathrm{SL}_n(\mathbb{R}) \times \mathrm{SL}_n(\mathbb{R})$ (see [Morris 2015, §5.1 and Example 5.1 #7]). It follows that $\mathcal{D}^1 \cap \mathrm{SL}_n(\mathcal{O}_k)$ has finite index in \mathcal{D}^1 and so $\Gamma \cap \mathrm{SL}_n(\mathcal{O}_k)$ has finite index in Γ . \square

Note that since Γ_2 is a lattice in $\mathrm{SO}(J_1, \mathbb{R})$ it acts irreducibly on \mathbb{C}^{n+1} , and so by applying Lemma 1.1 we may assume that $\Gamma_2 \subset \mathrm{SO}(J_1, \mathcal{O}_L)$.

Denote by $\mathrm{SO}(n-1, 1, \mathbb{R})$ the subgroup of $\mathrm{SO}(J_1, \mathbb{R})$ that preserves both complementary components in \mathbb{R}^{n+1} of the hyperplane P given by the equation $x_1 = 0$. The intersection $P \cap \mathbb{H}^n$ is a model for hyperbolic $(n-1)$ -space, \mathbb{H}^{n-1} and the group $\mathrm{SO}(n-1, 1, \mathbb{R})$ can be identified with the subgroup of orientation preserving isometries of \mathbb{H}^{n-1} . Next, let $\hat{\Gamma} = \Gamma_1 \cap \Gamma_2 \cap \mathrm{SO}(n-1, 1, \mathbb{R})$. Since each $\Gamma_i \cap \mathrm{SO}(n-1, 1, \mathbb{R})$ is sublattice of the lattice $\mathrm{SO}(n-1, 1, \mathcal{O}_L)$ in $\mathrm{SO}(n-1, 1, \mathbb{R})$, it follows that $\hat{\Gamma}$ is also a lattice in $\mathrm{SO}(n-1, 1, \mathbb{R})$. It follows that $\mathbb{H}^{n-1}/\hat{\Gamma}$ is a hyperbolic $(n-1)$ -orbifold. By passing to finite index subgroups we may arrange the following properties:

- (1) Γ_i is torsion-free and contained in the identity component of $\mathrm{SO}(J_1, \mathbb{R})$. This component is isomorphic to $\mathrm{Isom}^+(\mathbb{H}^n)$, and so $M_i := \mathbb{H}^n/\Gamma_i$ is a finite volume

hyperbolic manifold (apply Selberg’s lemma and the fact that $\mathrm{SO}(J, \mathbb{R})^\circ$ has finite index in $\mathrm{SO}(J, \mathbb{R})$).

- (2) Since $\Sigma = \mathbb{H}^{n-1}/\hat{\Gamma}$ is a totally geodesic we may assume that Σ is a hyperbolic $(n-1)$ -manifold and this manifold is embedded in both M_1 and M_2 (see [Bergeron 2000, Theorem 1]).
- (3) If M_i is noncompact then all cusps of M_i are diffeomorphic to an $(n-1)$ -torus times an interval (apply [McReynolds et al. 2013, Theorem 3.1]).
- (4) The complement $\hat{M}_i = M_i \setminus \Sigma$ is connected for $i = 1, 2$ (see [Bergeron 2000, Theorem 2]).

The manifold \hat{M}_i is a convex submanifold of M_i and so $\hat{M}_i = V_i/\hat{\Gamma}_i$, where V_i is a component of the preimage of \hat{M}_i in \mathbb{H}^n under the universal covering projection $\mathbb{H}^n \rightarrow \mathbb{H}^n/\Gamma_i = M_i$, and $\hat{\Gamma}_i$ is a subgroup of Γ_i that stabilizes V_i . The manifold \hat{M}_i is a hyperbolic manifold with totally geodesic boundary equal to two isometric copies of Σ , and so it is possible to glue \hat{M}_1 and \hat{M}_2 along Σ to form the finite volume hyperbolic manifold N (see [Morris 2015, §6.5] for details). The manifold N can be realized as \mathbb{H}^n/Δ where, after appropriately conjugating $\hat{\Gamma}_i$ in Γ_i , we may assume that

$$(1-2) \quad \Delta = \langle \hat{\Gamma}_1, \hat{\Gamma}_2, s \rangle.$$

Here s comes from a “graph of spaces” description of N and can thus be written as a product $s = s_2s_1$, where s_i is the isometry corresponding to an appropriate lift to V_i a curve in M_i whose algebraic intersection with Σ is 1 (See Figure 1). In [Gromov and Piatetski-Shapiro 1988, §2.9] it is shown that Δ is a nonarithmetic lattice in $\mathrm{SO}(J_1, \mathbb{R})$. If $N = \mathbb{H}^n/\Delta$ then we call N an *interbreeding of M_1 and M_2* .

Since $\Gamma_1, \Gamma_2 \subset \mathrm{SO}(J_1, \mathcal{O}_L)$ it follows that $\Delta \subset \mathrm{SO}(J_1, \mathcal{O}_L)$. As a result, we call the field L the *field of definition of Δ* . Let \mathcal{C}_n be the collection of hyperbolic n -manifolds coming from the above interbreeding construction.

We close this section by proving the following result:

Proposition 1.2. *The collection \mathcal{C}_n contains representatives of infinitely many commensurability classes of both closed and noncompact hyperbolic n -manifolds satisfying the properties (1)–(4) from above.*

To prove this we will need the following invariant, originally due to Vinberg [1971]. Let Γ be a Zariski dense subgroup of a Lie group H with Lie algebra \mathfrak{h} . The adjoint action of Γ on \mathfrak{h} gives a representation $\mathrm{Ad} : \Gamma \rightarrow \mathrm{gl}(\mathfrak{h})$. In [Vinberg 1971] it is shown that the field $\mathbb{Q}(\{\mathrm{tr}(\mathrm{Ad}(\gamma)) \mid \gamma \in \Gamma\})$ is an invariant of the commensurability class of Γ in H . This field is called the *adjoint trace field of Γ* .

Next, let $N = \mathbb{H}^n/\Delta \in \mathcal{C}_n$, then Δ is a lattice in $\mathrm{SO}(J_1, \mathbb{R})$, which is Zariski dense by the Borel density theorem. The following lemma allows us to compute




Figure 1. An graph of spaces description of the manifold N .

the adjoint trace field of Δ . It is an immediate corollary of a theorem of Mila (see [Mila 2019, Theorem 4.7]) once it is observed that L is the smallest extension of K over which the forms J_1 and J_2 are isometric.

Lemma 1.3. *Let $N = \mathbb{H}^n / \Delta \in \mathcal{C}_n$ and let L be the field of definition of Δ . Then L is the adjoint trace field of Δ .*

Proof of Proposition 1.2. From [Gromov and Piatetski-Shapiro 1988], it follows that $N = \mathbb{H}^n / \Delta$ is compact if and only if the field K used to construct Δ is not equal to \mathbb{Q} . For each choice of a totally real field K and a pair $\alpha, \beta \in K$ so that α/β is not a square in K we can produce an element $N \in \mathcal{C}_n$ via the interbreeding construction. By varying the choices of α and β we can produce infinitely many distinct $L = K(\sqrt{\beta/\alpha})$ for each choice of K . It follows from Lemma 1.3 that the corresponding N are representatives of infinitely many commensurability classes of both compact and noncompact hyperbolic n -manifolds. \square

2. Lattices in $SL_{n+1}(\mathbb{R})$

In this section we describe the lattices $\Delta \subset SL_{n+1}(\mathbb{R})$ in which our thin groups will ultimately live. Let J_1 be one of the forms constructed in Section 1 and let L be the corresponding (totally real) field of definition. Let $M = L(\sqrt{r})$, where $r \in L$ is positive, square-free, and $s_L(r) = 0$. The number field M is a quadratic extension of L and we let $\tau : M \rightarrow M$ be the unique nontrivial Galois automorphism of M over L . In this context, we can extend the quadratic form J_1 on L^{n+1} to a ‘‘Hermitian’’ form on M^{n+1} . Let $N_{M/L} : M \rightarrow L$ given by $N_{M/L}(x) = x\tau(x)$ be the norm of the field extension M/L . Next let $x = (x_1, \dots, x_{n+1}) \in M^{n+1}$ and define $H_1 : M^{n+1} \rightarrow L$ as

$$H_1(x) = \alpha N_{M/L}(x_1) + \sum_{i=1}^n a_i N_{M/L}(x_i) - a_{n+1} N_{M/L}(x_{n+1}).$$

Note that this defines a Hermitian form in the sense that if $x \in M^{n+1}$ and $\lambda \in M$ then $H_1(\lambda x) = N_{M/L}(\lambda)H_1(x)$. Furthermore, since L is the fixed field of τ it follows that H_1 reduces to J_1 when restricted to L^{n+1} .

Next, we can define a unitary analogue of $\mathrm{SO}(J_1, \mathcal{O}_M)$ as

$$\mathrm{SU}(J_1, \tau, \mathcal{O}_M) = \{A \in \mathrm{SL}_{n+1}(\mathcal{O}_M) \mid H_1(Av) = H_1(v) \forall v \in M^{n+1}\}.$$

It is well known (see [Morris 2015, §6.8], for example) that $\mathrm{SU}(J_1, \tau, \mathcal{O}_M)$ is an arithmetic lattice in $\mathrm{SL}_{n+1}(\mathbb{R})$.

Let $N = \mathbb{H}^n/\Delta$ be one of the manifolds from \mathcal{C}_n . By construction, the manifold N contains the embedded totally geodesic hypersurface $\Sigma = \mathbb{H}^{n-1}/\hat{\Gamma}$, and so it is possible to deform Δ inside of $\mathrm{SL}_{n+1}(\mathbb{R})$ using the bending construction of Johnson and Millson [1987].

Specifically, let $c_t = \mathrm{Diag}(e^{-nt}, e^t, \dots, e^t) \in \mathrm{SL}_{n+1}(\mathbb{R})$. It is easy to check that c_t centralizes $\mathrm{SO}(n-1, 1, \mathbb{R})$. Since Σ is assumed to be nonseparating, we see that we can write Δ as an HNN extension $\Delta \cong \hat{\Delta} * s$, where $\hat{\Delta}$ is isomorphic to the fundamental group of $N \setminus \Sigma$ and s is a free letter. In this context, we may view $\hat{\Delta} \subset \mathrm{SO}(J_1, \mathcal{O}_L)$ and $s \in \mathrm{SO}(J_1, \mathcal{O}_L)$ and observe that as a subgroup of $\mathrm{SO}(J_1, \mathcal{O}_L)$ we can write $\Delta = \langle \hat{\Delta}, s \rangle$. We now define a new family of subgroups $\Delta_t = \langle \Delta, c_t s \rangle \subset \mathrm{SL}_{n+1}(\mathbb{R})$. Using basic theory of HNN extensions, it is easy to see that, since c_t centralizes the fundamental group of Σ , as an abstract group Δ_t is a quotient of Δ . However, by using the following result due to Benoist [2005] in the compact case and Marquis [2012] in the noncompact case, we can actually say much more.

Proposition 2.1. *For each t , the group Δ_t is isomorphic to Δ .*

Next, we show for certain values of t the group Δ_t is contained in one of the unitary lattices constructed above. Specifically, if $N = \mathbb{H}^n/\Delta$ is contained in \mathcal{C}_n , let J_1 and L be such that $\Delta \subset \mathrm{SO}(J_1, \mathcal{O}_L)$. Recall that the field L is totally real of degree $d+1$ over \mathbb{Q} and so there are $d+1$ embeddings $\{\sigma_0 = \mathrm{Id}, \dots, \sigma_d\}$ of L into \mathbb{R} . We can use Lemma 3.1 of [Ballas and Long 2020] to produce a unit $u \in \mathcal{O}_L^\times$ with the property that $|u| > 2$ and $0 < |\sigma_i(u)| < 1$ for $1 \leq i \leq d$. Let $p(x) = x^2 - ux + 1$ and let $M = L(v)$, where v is one of the roots of $p(x)$. It is easy to check that the discriminant of $p(x)$ is $u^2 - 4$ and so $M = L(\sqrt{u^2 - 4})$. By construction $s_L(u^2 - 4) = 0$, and so $\mathrm{SU}(J_1, \tau, \mathcal{O}_M)$ is an arithmetic lattice in $\mathrm{SL}_{n+1}(\mathbb{R})$, where $\tau : M \rightarrow M$ is the nontrivial Galois automorphism of M over L . The next lemma says that by carefully choosing t , we can arrange that $\Delta_t \subset \mathrm{SU}(J_1, \tau, \mathcal{O}_M)$.

Lemma 2.2. *Let u be as above. Then if $t = \log(u)$ then $\Delta_t \subset \mathrm{SU}(J_1, \tau, \mathcal{O}_M)$.*

This is basically Lemma 3.4 of [Ballas and Long 2020], but the proof is short so we include it here for the sake of completeness.

Proof. Recall from above that there is a subgroup $\hat{\Delta} \subset \text{SO}(J_1, \mathcal{O}_L)$ and $s \in \text{SO}(J_1, \mathcal{O}_L)$ so that $\Delta = \langle \hat{\Delta}, s \rangle$ and $\Delta_t = \langle \hat{\Delta}, c_t s \rangle$, where

$$c_t = \text{Diag}(e^{-nt}, e^t, \dots, e^t) \in \text{SL}_{n+1}(\mathbb{R}).$$

Since $\text{SO}(J_1, \mathcal{O}_L) \subset \text{SU}(J_1, \tau, \mathcal{O}_N)$ the proof will be complete if we can show that $c_t \in \text{SU}(J_1, \tau, \mathcal{O}_M)$.

If $t = \log(u)$ then $c_t = \text{Diag}(u^{-n}, u, \dots, u)$. Furthermore, since $\tau(u)$ is the other root of $p(x)$ it follows that $u\tau(u) = 1$, or in other words $\tau(u) = u^{-1}$. It follows that $c_t^* = \text{Diag}(u^n, u^{-1}, \dots, u^{-1})$. A simple computation then shows that for each $v \in M^{n+1}$, $H_1(c_t v) = H_1(v)$, and so $c_t \in \text{SU}(J_1, \tau, \mathcal{O}_M)$. \square

By combining [Lemma 2.2](#) and [Proposition 2.1](#) we get the following corollary:

Corollary 2.3. *For each $N = \mathbb{H}^n / \Delta \in \mathcal{C}_n$ there are infinitely many lattices $\Lambda \subset \text{SL}_{n+1}(\mathbb{R})$ that contain a subgroup Δ' isomorphic to Δ .*

3. Certifying thinness

The main goal of this section is to complete the proof of [Theorem 1](#). The proof consist of proving that the subgroups constructed in the previous section are thin.

Proof of Theorem 1. Recall, that if $N = \mathbb{H}^n / \Delta \in \mathcal{C}_n$ from [Corollary 2.3](#) it follows that we can find a lattice $\Lambda \subset \text{SL}_{n+1}(\mathbb{R})$ and a subgroup $\Delta' \subset \Lambda$ that is isomorphic to Δ .

Since Δ' was obtained from Δ via a bending construction it follows from [[Ballas and Long 2020](#), Proposition 4.1] that Δ' is Zariski dense in $\text{SL}_{n+1}(\mathbb{R})$. The proof will be complete if we can show that Δ' has infinite index in Λ . Suppose for contradiction that this index is finite. Since Λ is a lattice in $\text{SL}_{n+1}(\mathbb{R})$ this implies that Δ' is also a lattice in $\text{SL}_{n+1}(\mathbb{R})$. However, Δ' is isomorphic to Δ and Δ is a lattice in the Lie group $\text{SO}(n, 1)^\circ$. However, $\text{SO}(n, 1)^\circ$ and $\text{SL}_{n+1}(\mathbb{R})$ are not isomorphic and so this contradicts the Mostow rigidity theorem (see [[Morris 2015](#), Theorem 15.1.2]). \square

Acknowledgements

The author would like to thank Darren Long for several helpful conversations during the preparation of this work and Matt Stover for providing references that greatly simplified the proof of [Lemma 1.3](#). The author was partially supported by the NSF grant DMS-1709097.

References

[Ballas and Long 2020] S. A. Ballas and D. D. Long, “Constructing thin subgroups of $\text{SL}(n + 1, \mathbb{R})$ via bending”, *Algebr. Geom. Topol.* **20**:4 (2020), 2071–2093. [MR Zbl](#)

- [Bass 1980] H. Bass, “Groups of integral representation type”, *Pacific J. Math.* **86**:1 (1980), 15–51. [MR](#) [Zbl](#)
- [Benoist 2005] Y. Benoist, “Convexes divisibles, III”, *Ann. Sci. École Norm. Sup. (4)* **38**:5 (2005), 793–832. [MR](#) [Zbl](#)
- [Bergeron 2000] N. Bergeron, “Premier nombre de Betti et spectre du laplacien de certaines variétés hyperboliques”, *Enseign. Math. (2)* **46**:1-2 (2000), 109–137. [MR](#) [Zbl](#)
- [Bourgain 2014] J. Bourgain, “Some Diophantine applications of the theory of group expansion”, pp. 1–22 in *Thin groups and superstrong approximation*, edited by E. Breuillard and H. Oh, Math. Sci. Res. Inst. Publ. **61**, Cambridge Univ. Press, 2014. [MR](#) [Zbl](#)
- [Bourgain et al. 2010] J. Bourgain, A. Gamburd, and P. Sarnak, “Affine linear sieve, expanders, and sum-product”, *Invent. Math.* **179**:3 (2010), 559–644. [MR](#) [Zbl](#)
- [Chen 1978] J. R. Chen, “On the representation of a large even integer as the sum of a prime and the product of at most two primes, II”, *Sci. Sinica* **21**:4 (1978), 421–430. [MR](#)
- [Cooper and Futer 2019] D. Cooper and D. Futer, “Ubiquitous quasi-Fuchsian surfaces in cusped hyperbolic 3-manifolds”, *Geom. Topol.* **23**:1 (2019), 241–298. [MR](#) [Zbl](#)
- [Fuchs and Rivin 2017] E. Fuchs and I. Rivin, “Generic thinness in finitely generated subgroups of $SL_n(\mathbb{Z})$ ”, *Int. Math. Res. Not.* **2017**:17 (2017), 5385–5414. [MR](#) [Zbl](#)
- [Gromov and Piatetski-Shapiro 1988] M. Gromov and I. Piatetski-Shapiro, “Nonarithmetic groups in Lobachevsky spaces”, *Inst. Hautes Études Sci. Publ. Math.* **66** (1988), 93–103. [MR](#)
- [Johnson and Millson 1987] D. Johnson and J. J. Millson, “Deformation spaces associated to compact hyperbolic manifolds”, pp. 48–106 in *Discrete groups in geometry and analysis* (New Haven, CT, 1984), edited by R. Howe, Progr. Math. **67**, Birkhäuser, Boston, 1987. [MR](#) [Zbl](#)
- [Kahn and Markovic 2012] J. Kahn and V. Markovic, “Immersing almost geodesic surfaces in a closed hyperbolic three manifold”, *Ann. of Math. (2)* **175**:3 (2012), 1127–1190. [MR](#) [Zbl](#)
- [Kahn and Wright 2018] J. Kahn and A. Wright, “Nearly Fuchsian surface subgroups of finite covolume Kleinian groups”, preprint, 2018. [arXiv](#)
- [Kahn et al. 2018] J. Kahn, F. Labourie, and S. Mozes, “Surface subgroups in uniform lattices of some semi-simple Lie groups”, preprint, 2018, available at <http://flab.perso.math.cnrs.fr/preprints/SGL.pdf>.
- [Lubotzky 2012] A. Lubotzky, “Expander graphs in pure and applied mathematics”, *Bull. Amer. Math. Soc. (N.S.)* **49**:1 (2012), 113–162. [MR](#) [Zbl](#)
- [Marquis 2012] L. Marquis, “Exemples de variétés projectives strictement convexes de volume fini en dimension quelconque”, *Enseign. Math. (2)* **58**:1-2 (2012), 3–47. [MR](#) [Zbl](#)
- [McReynolds et al. 2013] D. B. McReynolds, A. W. Reid, and M. Stover, “Collisions at infinity in hyperbolic manifolds”, *Math. Proc. Cambridge Philos. Soc.* **155**:3 (2013), 459–463. [MR](#) [Zbl](#)
- [Mila 2019] O. Mila, *The trace field of hyperbolic gluings*, Ph.D. thesis, Universität Bern, 2019.
- [Morris 2015] D. W. Morris, *Introduction to arithmetic groups*, Deductive Press, 2015. [MR](#) [Zbl](#)
- [Ratcliffe 2006] J. G. Ratcliffe, *Foundations of hyperbolic manifolds*, 2nd ed., Graduate Texts in Mathematics **149**, Springer, 2006. [MR](#) [Zbl](#)
- [Vinberg 1971] E. B. Vinberg, “Rings of definition of dense subgroups of semisimple linear groups”, *Izv. Akad. Nauk SSSR Ser. Mat.* **35** (1971), 45–55. In Russian; translated in *Math. USSR-Izv.* **5**:1 (1971), 45–55. [MR](#) [Zbl](#)

SAMUEL A. BALLAS
DEPARTMENT OF MATHEMATICS
FLORIDA STATE UNIVERSITY
TALLAHASSEE, FL
UNITED STATES
ballas@math.fsu.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 309 No. 2 December 2020

Thin subgroups isomorphic to Gromov–Piatetski-Shapiro lattices	257
SAMUEL A. BALLAS	
Value distribution properties for the Gauss maps of the immersed harmonic surfaces	267
XINGDI CHEN, ZHIXUE LIU and MIN RU	
Scattered representations of $SL(n, \mathbb{C})$	289
CHAO-PING DONG and KAYUE DANIEL WONG	
Number of singular fibres of surface fibrations over \mathbb{P}^1	313
CHENG GONG and YI GU	
Asymptotic behavior of solutions for some elliptic equations in exterior domains	333
ZONGMING GUO and ZHONGYUAN LIU	
Ulrich elements in normal simplicial affine semigroups	353
JÜRGEN HERZOG, RAHELEH JAFARI and DUMITRU I. STAMATE	
Notes on equivariant homology with constant coefficients	381
SOPHIE KRIZ	
Local normal forms for multiplicity free $U(n)$ actions on coadjoint orbits	401
JEREMY LANE	
Functional determinant on pseudo-Einstein 3-manifolds	421
ALI MAALAOUI	
Distribution of distances in positive characteristic	437
THANG PHAM and LÊ ANH VINH	
Elliptic gradient estimates for a parabolic equation with V -Laplacian and applications	453
JIAN-HONG WANG and YU ZHENG	
Optimal L^2 extension of sections from subvarieties in weakly pseudoconvex manifolds	475
XIANGYU ZHOU and LANGFENG ZHU	