

*Pacific
Journal of
Mathematics*

NOTES ON EQUIVARIANT HOMOLOGY
WITH CONSTANT COEFFICIENTS

SOPHIE KRIZ

NOTES ON EQUIVARIANT HOMOLOGY WITH CONSTANT COEFFICIENTS

SOPHIE KRIZ

For a finite group, we discuss a method for calculating equivariant homology with constant coefficients. We apply this method to completely calculate the geometric fixed points of the equivariant spectrum representing equivariant (co)homology with constant coefficients. We also treat a more complicated example of inverting the standard representation in the equivariant homology of split extraspecial groups at the prime 2.

1. Introduction

Equivariant spectra are the foundation of equivariant generalized homology and cohomology theory of G -spaces (and ultimately, G -spectra) for a finite (or more generally compact Lie) group G , which has all the formal properties of generalized nonequivariant homology and cohomology theory, including duality, along with stability under suspensions by finite-dimensional real representations. They were introduced and developed in [Lewis et al. 1986] (see also [Adams 1984; Greenlees 1985]). Equivariant homology and cohomology $H\mathbb{A}_G$ with constant coefficients in an abelian group A , on the other hand, can be defined on the chain level, as a part of the theory of Bredon [1967]. Both contexts are reconciled in [Lewis et al. 1981], where, more generally, equivariant Eilenberg–MacLane spectra of Mackey functors are defined. In this paper, we discuss a spectral sequence (Propositions 4, 5) which can be used to compute generalized equivariant homology of a G -CW-complex from its subquotients of constant isotropy. This spectral sequence is especially efficient in the case of $H\mathbb{A}_G$. For example, we shall prove that for a (finite) p -group G , the spectral sequence computing $H\mathbb{Z}/p_*^G(X)$ always collapses to E^1 (see Theorem 7).

Note that this is false in cohomology. By [Bredon 1967], for a G -CW-complex X and an abelian group A ,

$$(1) \quad H_G^*(X; \underline{A}) = H^*(X/G; A).$$

Let $G = \mathbb{Z}/2$ and $X = S^\alpha$ where α denotes the sign representation of $\mathbb{Z}/2$. Then $X/(\mathbb{Z}/2) \simeq *$. Thus, by (1), $H_G^n(X; \underline{A})$ is only nontrivial for $n = 0$, while X has a 1-cell of isotropy $\{e\}$.

MSC2020: 55N91.

Keywords: equivariant homology, geometric fixed points, extraspecial groups.

One may think, therefore, that equivariant homology with constant coefficients carries less information than cohomology. It turns out, however, that equivariant E -homologies of certain spaces give important information about a spectrum E . For example, the *geometric fixed point spectra* [Greenlees 1985; Lewis et al. 1986, II §8, 9] $\Phi^H E$, where H runs through subgroups of G , completely characterize the spectrum E . The coefficients $\Phi^G E_*$, for a finite group G , are the reduced E -homology of the smash product $S^{\infty V}$ of infinitely many copies of the one-point compactification S^V of the reduced regular representation V . Geometric fixed point spectra proved very useful in applications, for example, in [tom Dieck 1970; Hesselholt and Madsen 1997; Hill et al. 2016].

We will see that our method allows a complete computation of the coefficients of the geometric fixed point spectrum of homology with constant coefficients $\Phi^G(H\underline{A})_*$, which we denote by $\Phi^G(\underline{A})_*$, by reducing it to the case where G is an elementary abelian group. We show (Proposition 11) that $\Phi^G H\underline{A}_G = 0$ if G is not a p -group, and that

$$\Phi^G H\underline{A}_G = \Phi^{G/G'_p} H\underline{A}_{G/G'_p},$$

where G is a p -group and G'_p is its Frattini subgroup (Proposition 9). In the elementary abelian case, the computation was carried out for $A = \mathbb{Z}/p$ in my previous paper [Kriz 2015] (see also [Holler and Kriz 2017; 2020]).

Let first $p = 2$. We have

$$H^*((\mathbb{Z}/2)^n; \mathbb{Z}/2) = \mathbb{Z}/2[x_1, \dots, x_n],$$

where x_i have cohomological dimension 1. Let, for $\alpha = (\alpha_1, \dots, \alpha_n) \in (\mathbb{Z}/2)^n \setminus \{0\}$,

$$x_\alpha = \alpha_1 x_1 + \dots + \alpha_n x_n.$$

For $p > 2$, following the notation of [Kriz 2015], we have

$$H^*((\mathbb{Z}/p)^n; \mathbb{Z}/p) = \mathbb{Z}/p[z_i] \otimes \Lambda_{\mathbb{Z}/p}[dz_i],$$

where z_i have cohomological dimension 2 and dz_i have cohomological dimension 1. Let, for $\alpha = (\alpha_1, \dots, \alpha_n) \in (\mathbb{Z}/p)^n \setminus \{0\}$,

$$z_\alpha = \alpha_1 z_1 + \dots + \alpha_n z_n,$$

$$dz_\alpha = \alpha_1 dz_1 + \dots + \alpha_n dz_n.$$

Theorem 1 [Holler and Kriz 2017; 2020; Kriz 2015]. (a) $\Phi^{(\mathbb{Z}/2)^n}(\underline{\mathbb{Z}/2})$ is the subring of

$$H^*((\mathbb{Z}/2)^n; \mathbb{Z}/2)[x_\alpha^{-1} \mid \alpha \neq 0]$$

generated by $y_\alpha = x_\alpha^{-1}$.

(b) For $p > 2$, $\Phi^{(\mathbb{Z}/p)^n}(\underline{\mathbb{Z}/p})$ is the subring of

$$H^*((\mathbb{Z}/p)^n; \mathbb{Z}/p)[z_\alpha^{-1} | \alpha \in (\mathbb{Z}/p)^n \setminus \{0\}]$$

generated by $t_\alpha = z_\alpha^{-1}$ and $u_\alpha = t_\alpha dz_\alpha$.

One can also explicitly describe these rings in generators and defining relations:

Theorem 2 [Holler and Kriz 2017; 2020; Kriz 2015]. (a) For $p = 2$, we have

$$\Phi_*^{(\mathbb{Z}/2)^n}(\underline{\mathbb{Z}/2}) \cong \mathbb{Z}/2[y_\alpha | \alpha \in (\mathbb{Z}/2)^n \setminus \{0\}] / \sim$$

where \sim denotes the relations

$$y_\alpha y_\beta + y_\alpha y_\gamma + y_\beta y_\gamma \sim 0$$

for $\alpha + \beta + \gamma = 0$. The elements y_α are in degree 1.

(b) For $p > 2$,

$$(2) \quad \Phi_*^{(\mathbb{Z}/p)^n}(\underline{\mathbb{Z}/p}) = \mathbb{Z}/p[t_\alpha] \otimes \Lambda_{\mathbb{Z}/p}[u_\alpha] / \sim,$$

where \sim denotes the relations

$$t_{i\alpha} \sim i^{-1}t_\alpha, \quad u_{i\alpha} \sim u_\alpha, \quad t_\beta t_{\alpha+\beta} + t_\alpha t_{\alpha+\beta} \sim t_\alpha t_\beta, \\ t_\beta u_{\alpha+\beta} - t_{\alpha+\beta} u_\beta + t_{\alpha+\beta} u_\alpha \sim u_\alpha t_\beta, \quad -u_\beta u_{\alpha+\beta} + u_\alpha u_{\alpha+\beta} \sim u_\alpha u_\beta,$$

where $i \in \mathbb{Z}/p \setminus \{0\}$, for $\alpha, \beta, \alpha + \beta \in (\mathbb{Z}/p)^n \setminus \{0\}$. The elements u_α are in degree 1 and the elements t_α are of degree 2.

This allows a complete answer for $\underline{\mathbb{Z}}$, the universal constant coefficients. Fix an element $\alpha_0 \in (\mathbb{Z}/p)^n \setminus \{0\}$ and put $\tilde{u}_\alpha = u_\alpha - u_{\alpha_0}$, ($\tilde{y}_\alpha = y_\alpha - y_{\alpha_0}$ for $p = 2$). For $p = 2$, we also set $t_{\alpha_0} = y_{\alpha_0}^2$.

Theorem 3. (a) For every prime p , we have that $\Phi_*^{(\mathbb{Z}/p)^n}(\underline{\mathbb{Z}})$ is the subring of the ring $\Phi_*^{(\mathbb{Z}/p)^n}(\underline{\mathbb{Z}/p})$ on which the Bockstein

$$\beta : \Phi_*^{(\mathbb{Z}/p)^n}(\underline{\mathbb{Z}/p}) \rightarrow \Phi_{*-1}^{(\mathbb{Z}/p)^n}(\underline{\mathbb{Z}/p})$$

vanishes.

(b) Explicitly, for $p = 2$, we have

$$\Phi_*^{(\mathbb{Z}/2)^n}(\underline{\mathbb{Z}}) \cong \mathbb{Z}/2[\tilde{y}_\alpha, t_{\alpha_0} | \alpha \in (\mathbb{Z}/2)^n \setminus \{0\}] / \sim,$$

where \sim denotes the relations

$$\tilde{y}_{\alpha_0} \sim 0, \quad \tilde{y}_\alpha \tilde{y}_\beta + \tilde{y}_\alpha \tilde{y}_\gamma + \tilde{y}_\beta \tilde{y}_\gamma + t_{\alpha_0} \sim 0,$$

where $\alpha + \beta + \gamma = 0$.

(c) For $p > 2$,

$$\Phi_*^{(\mathbb{Z}/p)^n}(\mathbb{Z}) \cong (\mathbb{Z}/p[t_\alpha | \alpha \in (\mathbb{Z}/p)^n \setminus \{0\}]) \otimes \Lambda_{\mathbb{Z}/p}[\tilde{u}_\alpha | \alpha \in (\mathbb{Z}/p)^n \setminus \{0\}] / \sim,$$

where \sim denotes the relations

$$(3) \quad \begin{aligned} t_{i\alpha} &\sim i^{-1}t_\alpha, & \tilde{u}_{i\alpha} &\sim \tilde{u}_\alpha, & t_\beta t_{\alpha+\beta} + t_\alpha t_{\alpha+\beta} &\sim t_\alpha t_\beta, & \tilde{u}_{\alpha_0} &\sim 0, \\ t_\beta \tilde{u}_{\alpha+\beta} - t_{\alpha+\beta} \tilde{u}_\beta + t_{\alpha+\beta} \tilde{u}_\alpha &\sim t_\beta \tilde{u}_\alpha, & -\tilde{u}_\beta \tilde{u}_{\alpha+\beta} + \tilde{u}_\alpha \tilde{u}_{\alpha+\beta} &\sim \tilde{u}_\alpha \tilde{u}_\beta, \end{aligned}$$

for $i \in \mathbb{Z}/p \setminus \{0\}$ and $\alpha, \beta, \alpha + \beta \in (\mathbb{Z}/p)^n \setminus \{0\}$.

The reductions contained in Propositions 9 and 11 are quite easy. However, if one considers the more general problem of calculating $\widetilde{H}A_*^G(S^{\infty\gamma})$ for a general finite-dimensional representation γ of G , one gets nontrivial examples. One such example is treated in Section 4, where G is a split extraspecial 2-group and γ is the irreducible representation nontrivial on the center.

The present paper is organized as follows: In Section 2, we discuss our spectral sequences. In Section 3, we discuss the application to geometric fixed points. Section 4 contains the extraspecial group example.

2. The spectral sequences

For a G -equivariant spectrum E , we will need to use the homotopy co-fixed point (Borel homology) spectrum

$$(4) \quad E_{hG} = (E \wedge EG_+)^G,$$

where EG is a nonequivariantly contractible free G -CW complex and for a space X , we write $X_+ = X \coprod \{*\}$. The formula (4) includes a key fact called the Adams isomorphism [Lewis et al. 1986, II §7]: G -equivariant cell spectra with G -free cells can be identified with naive (i.e., nonequivariant) cell spectra with a free cellular G -action. The Adams isomorphism says that for any cell G -spectrum E , E_{hG} is equivalent to $(E \wedge EG_+)/G$, where $E \wedge EG_+$ is considered as a naive G -spectrum.

Recall that a family \mathcal{F} is defined as a set of subgroups of G that is closed under subconjugacies. For a family \mathcal{F} , we have a G -CW complex $E_{\mathcal{F}}$ such that

$$\begin{aligned} E_{\mathcal{F}}^H &\simeq * & \text{for } H \in \mathcal{F}, \\ E_{\mathcal{F}}^G &\simeq \emptyset & \text{for } H \notin \mathcal{F}. \end{aligned}$$

If we denote by \widetilde{X} the unreduced suspension of a G -space X , we have

$$\begin{aligned} \widetilde{E}_{\mathcal{F}}^H &\simeq * & \text{for } H \in \mathcal{F}, \\ \widetilde{E}_{\mathcal{F}}^G &\simeq S^0 & \text{for } H \notin \mathcal{F}. \end{aligned}$$

If V is a real G -representation, denote by $S(V)$ the unit sphere of V and by S^V the union of the 1-point compactifications of S^W for finite-dimensional subrepresentations W of V . If we set $\infty V = \bigoplus_{\infty} V$, then $S(\infty V)$ is a model for $E_{\mathcal{F}_V}$, where

$\mathcal{F}_V = \{H \subseteq G \mid V^H \neq 0\}$. Thus $S^{\infty V}$ is a model for $\widetilde{E}\mathcal{F}_V$. Since $S^{\infty V} \wedge S^{\infty V} = S^{\infty V}$, for a commutative ring spectrum E ,

$$\widetilde{E}_* \widetilde{E}\mathcal{F}_V = \widetilde{E}_* S^{\infty V}$$

is a commutative ring. Here $\widetilde{E}_* X$, for a G -spectrum E and a based G -CW complex X (recall that the base point is required to be G -fixed), is the equivariant reduced homology of X , i.e., $\pi_*(E \wedge \Sigma^\infty X)$ (without adding a disjoint base point). Note that this is also the \mathbb{Z} -graded part of the $RO(G)$ -graded coefficient ring of $\alpha_V^{-1} E$, where $\alpha_V \in \pi_{-V} E$ is the class obtained from the inclusion $S^0 \rightarrow S^V$.

For a finite group G , a family \mathcal{F} , and an $H \in \mathcal{F}$, define the height of H inductively by

$$h_{\mathcal{F}}(H) = \max\{0, h_{\mathcal{F}}(K) \mid K \in \mathcal{F}, H \subsetneq K\} + 1.$$

Now let X be a G -CW-complex. Consider the family

$$\mathcal{F} = \mathcal{F}_X = \{H \subseteq G \mid X^H \neq \emptyset\}.$$

Let E be a G -spectrum. Then we have a spectral sequence converging to the E -homology of X , using Borel homology of parts of X of the same isotropy. There are two versions of the spectral sequence, one for the unreduced homology of X , the other for the reduced homology of its unreduced suspension. Keeping track of terms can be delicate, so we list both versions:

Proposition 4. *We have a spectral sequence*

$$E_{p,q}^1 \Rightarrow E_{p+q} X,$$

where

$$(5) \quad E_{p,q}^1 = \bigoplus_{(H), H \in \mathcal{F}_X, h_{\mathcal{F}_X}(H)=p} \left(\left(E^H \wedge \left(X^H / \bigcup_{H \subsetneq K} X^K \right) \right)_{hW(H)} \right)_{p+q}.$$

(Here $W(H) = N(H)/H$, where $N(H)$ is the normalizer of H in G , and (H) runs through the conjugacy classes of $H \in \mathcal{F}_X$.)

Proposition 5. *We have a spectral sequence*

$$E_{p,q}^1 \Rightarrow \widetilde{E}_{p+q} \widetilde{X},$$

where

$$(6) \quad E_{p,q}^1 = E_{0,q} = E_q(*),$$

$$(6) \quad E_{p,q}^1 = \bigoplus_{(H), H \in \mathcal{F}_X, h_{\mathcal{F}_X}(H)=p} \left(\left(E^H \wedge \left(X^H / \bigcup_{H \subsetneq K} X^K \right) \right)_{hW(H)} \right)_{p+q-1}.$$

The proofs of these statements are sufficiently similar to only give one of them. We prove [Proposition 5](#) which is more closely related to our applications.

Proof of Proposition 5. Define an increasing G -equivariant filtration of X by

$$F'_p X = \bigcup_{h_{\mathcal{F}}(H) \leq p} X^H = \bigcup_{h_{\mathcal{F}}(H)=p} X^H.$$

Then define an increasing G -equivariant filtration of \tilde{X} by

$$F_0 \tilde{X} = S^0, \\ F_p \tilde{X} = \bigcup_{h_{\mathcal{F}}(H) \leq p} (\tilde{X})^H = \bigcup_{h_{\mathcal{F}}(H)=p} (\tilde{X})^H.$$

We have a spectral sequence

$$E^1_{p,q} = \tilde{E}_{p+q}(F_p \tilde{X}/F_{p-1} \tilde{X}) \Rightarrow \tilde{E}_{p+q}(\tilde{X}) = (E \wedge \tilde{X})_{p+q}.$$

By definition, for $p \geq 1$,

$$F_p \tilde{X}/F_{p-1} \tilde{X} = \bigcup_{h_{\mathcal{F}}(H)=p} (\tilde{X})^H / \bigcup_{h_{\mathcal{F}}(H) \leq p-1} (\tilde{X})^H.$$

On the other hand,

$$F_p \tilde{X}/F_{p-1} \tilde{X} = (F_p \tilde{X}/F_0 \tilde{X}) / (F_{p-1} \tilde{X}/F_0 \tilde{X}) \\ = (\Sigma F'_p(X)_+) / (\Sigma F'_{p-1}(X)_+) \\ = \Sigma(F'_p(X)/F'_{p-1}(X)).$$

Thus,

$$\tilde{E}_{p+q}(F_p \tilde{X}/F_{p-1} \tilde{X}) = \tilde{E}_{p+q-1}(F'_p(X)/F'_{p-1}(X)).$$

Now, we have

$$F'_p(X)/F'_{p-1}(X) = \bigcup_{h_{\mathcal{F}}(H)=p} X^H / \bigcup_{h_{\mathcal{F}}(H) < p} X^H \\ = \bigvee_{h_{\mathcal{F}}(H)=p} \left(X^H / \bigcup_{H \subsetneq K \in \mathcal{F}} X^K \right).$$

Note that

$$X^H / \bigcup_{H \subsetneq K \in \mathcal{F}} X^K$$

is a free based $W(H)$ -CW complex.

On the other hand,

$$\begin{aligned}
 \tilde{E}_{p+q-1}^G \left(\bigvee_{h \in \mathcal{F}(H)=p} \left(X^H / \bigcup_{H \subsetneq K \in \mathcal{F}} X^K \right) \right) \\
 &= \bigoplus_{(H), h \in \mathcal{F}(H)=p} \tilde{E}_{p+q-1}^G \left(\bigvee_{H'=gHg^{-1}} \left(X^H / \bigcup_{H' \subsetneq K \in \mathcal{F}} X^K \right) \right) \\
 &= \bigoplus_{(H), h \in \mathcal{F}(H)=p} \tilde{E}_{p+q-1}^{N(H)} \left(X^H / \bigcup_{H \subsetneq K \in \mathcal{F}} X^K \right).
 \end{aligned}$$

To justify the third isomorphism above, note that G acts transitively on the conjugacy classes of H and thus

$$\bigvee_{H'=gHg^{-1}} \left(X^H / \bigcup_{H' \subsetneq K \in \mathcal{F}} X^K \right)$$

is the pushforward from $N(H)$ to G of

$$X^H / \bigcup_{H \subsetneq K \in \mathcal{F}} X^K.$$

Therefore,

$$\begin{aligned}
 E_{p,q}^1 &= \bigoplus_{(H), h \in \mathcal{F}(H)=p} \tilde{E}_{p+q-1}^{N(H)} \left(\left(X^H / \bigcup_{H \subsetneq K} X^K \right) \wedge EW(H)_+ \right) \\
 &= \bigoplus_{(H), h \in \mathcal{F}(H)=p} \left(\left(E^H \wedge \left(X^H / \bigcup_{H \subsetneq K} X^K \right) \right)_{hW(H)} \right)_{p+q-1},
 \end{aligned}$$

since $E^G = (E^K)^{G/K}$. □

We shall be especially interested in the case of classifying spaces of families. Consider, for a family \mathcal{F} and a group $H \in \mathcal{F}$, the poset

$$(7) \quad P_H^{\mathcal{F}} = \{K \in \mathcal{F} \mid K \supsetneq H\}$$

with respect to inclusion. Note that this poset has a $N(H)$ -action by conjugation.

For a poset P , denote by $|P|$ the nerve (also called the classifying space or bar construction) of P , which one defines as the geometric realization of the simplicial set whose n -simplices are chains

$$x_0 \leq \cdots \leq x_n$$

where faces are given by deletions and degeneracies by repetitions. This is a special case of the nerve of a category where n -simplices are composable n -tuples of morphisms.

Corollary 6. *We have a spectral sequence*

$$E^1_{p,q} \Rightarrow (E \wedge \widetilde{E\mathcal{F}})_{p+q} = \widetilde{E}_{p+q} \widetilde{E\mathcal{F}},$$

where

$$E^1_{0,q} = E_q(*)$$

and for $p > 0$,

$$(8) \quad E^1_{p,q} = \bigoplus_{(H), H \in \mathcal{F}, h_{\mathcal{F}}(H)=p} ((E^H \wedge |\widetilde{P_H^{\mathcal{F}}}|)_{hW(H)})_{p+q-1},$$

where (H) runs through the conjugacy classes of groups $H \in \mathcal{F}$.

Proof. Apply Proposition 5 to $X = E\mathcal{F}$. Note that $\mathcal{F}_{E\mathcal{F}} = \mathcal{F}$. Let $H \in \mathcal{F} = \mathcal{F}_X$. Then

$$E\mathcal{F}^H \simeq *.$$

We may realize the system of spaces $(E\mathcal{F}^K)_{H \subsetneq K \in \mathcal{F}}$ as a $N(H)$ -equivariant functor

$$F : P_H^{\mathcal{F}} \rightarrow \Delta^{Op}\text{-Set},$$

where the right-hand side denotes the category of simplicial sets, such that the canonical maps

$$\operatorname{colim}_{y < x} F(y) \rightarrow F(x)$$

are injective for all $x \in P_H^{\mathcal{F}}$. Now if we denote by $\pi : P_H^{\mathcal{F}} \rightarrow *$ the terminal map, $\bigcup_{H \subsetneq K \in \mathcal{F}} E\mathcal{F}^K$ is the left Kan extension $\pi_{\#}F$. However, by our injectivity assumption, the canonical $N(H)$ -equivariant map

$$(9) \quad L\pi_{\#}F \rightarrow \pi_{\#}F$$

(where L denotes the left derived functor) is a nonequivariant equivalence. Additionally, the left-hand side can be expressed as the 2-sided bar construction $B(*, P_H^{\mathcal{F}}, F)$. Since the values of F on objects are contractible, we have an $N(H)$ -equivariant map

$$(10) \quad B(*, P_H^{\mathcal{F}}, F) \rightarrow B(*, P_H^{\mathcal{F}}, *) = |P_H^{\mathcal{F}}|$$

which is a nonequivariant equivalence. The equivariant maps (9), (10) induce equivalences on homotopy fixed points, since they are nonequivariant equivalences. Thus, we have an equivalence

$$\left(E^H \wedge \left(E\mathcal{F}^H / \bigcup_{H \subsetneq K} E\mathcal{F}^K \right) \right)_{hW(H)} \sim (E^H \wedge |\widetilde{P_H^{\mathcal{F}}}|)_{hW(H)}$$

as required. □

Again, there is also an unbased version for $E\mathcal{F}$ instead of $\widetilde{E\mathcal{F}}$.

Theorem 7. *If G is a p -group and $E = H\mathbb{Z}/p$, then the spectral sequence (5) collapses to E^1 . Additionally, the spectral sequence (6) collapses to E^1 when $X^G = \emptyset$.*

Proof. Again, the proofs of the reduced and unreduced cases are similar. We treat the unreduced case this time.

Suppose G is a p -group and X is a G -CW-complex. Then $H\mathbb{Z}/p_*^G X$ can be calculated on the chain level. Let $C^G(X)$ be the cellular chain complex of X in the category of G -coefficient systems in the sense of Bredon [1967], i.e., functors $\mathcal{O}_G^{Op} \rightarrow Ab$ where \mathcal{O}_G is the orbit category. This is defined by

$$(C_n^G(X))(G/H) = C_n^{\text{cell}}(X^H).$$

Then we have

$$H\underline{A}_n(X) = H_n(C^G(X) \otimes_{\mathcal{O}_G} \underline{A}),$$

where \underline{A} is the constant co-coefficient system for an abelian group A , i.e., the functor $\mathcal{O}_G \rightarrow Ab$ where for $f : G/H \rightarrow G/K \in \text{Mor}(\mathcal{O}_G)$, f_* is multiplication by $|K|/|H|$.

We will show that

$$(11) \quad C^G(X) \otimes_{\mathcal{O}_G} \mathbb{Z}/p \cong \bigoplus_{(H)} \tilde{C}^{\text{cell}} \left(X^H / \bigcup_{H \subsetneq K} X^K \right) \otimes_{\mathbb{Z}[W(H)]} \mathbb{Z}/p.$$

In each degree separately, (11) holds as abelian groups, since all \mathcal{O}_G -identifications corresponding to nonisomorphisms are trivial. For any $f : H \subsetneq K$, consider the summand of the differential d^{tot} of $C^G(X) \otimes_{\mathcal{O}_G} \mathbb{Z}/p$

$$d_{H,K} : C_n(X^H) \otimes \mathbb{Z}/p \rightarrow C_{n-1}(X^K) \otimes \mathbb{Z}/p.$$

By conjugation, it suffices to show that these maps are 0.

The differential d^{tot} is given by

$$\bigoplus d / \sim : \bigoplus C_n(X^H) \otimes \mathbb{Z}/p / \sim \rightarrow \bigoplus C_{n-1}(X^H) \otimes \mathbb{Z}/p / \sim,$$

where \sim denotes the equivalence relation generated by

$$f^*a \otimes b \sim a \otimes f_*b.$$

In particular, for $q \in C_n(X^H)$, let $c \in C_{n-1}(X^H)$ be the sum of the terms of $d(q)$ on cells in X^K , where d is the differential of $C(X^H)$. Then we have

$$d_{H,K}(q) = f^*c \otimes 1 = c \otimes f_*1 = c \otimes \frac{|K|}{|H|} = 0.$$

Therefore, $d_{H,K} = 0$. Thus, we have proved (11), and hence the spectral sequence collapses to E^1 . □

3. Geometric fixed points

In this section, we shall apply the methods of the previous section to completely calculate the coefficients of the geometric fixed point spectrum

$$\Phi_*^G H\mathbb{A} = \widetilde{H}\mathbb{A}_* E\widetilde{\mathcal{F}}[G],$$

where $\mathcal{F}[G] = \{H \mid H \subsetneq G\}$ for any finite group G .

A basic fact about posets is useful for computing examples: For functors $F : \mathcal{C} \rightarrow \mathcal{D}$ between any categories, we get continuous maps

$$|F| : |\mathcal{C}| \rightarrow |\mathcal{D}|,$$

and for natural transformations $F \rightarrow G$, we get

$$|F| \simeq |G|.$$

In particular, if $f, g : P \rightarrow Q$ are morphisms of posets and $f \leq g$, then $|f| \simeq |g|$. Thus, in particular, if P has lowest or highest element x , $|\text{Id}| \simeq |\text{Const}_x|$. Therefore, then, $|P|$ is contractible.

Lemma 8. *For posets $Q \subseteq P$, if there exists a morphism $f : P \rightarrow Q$ satisfying both*

- (1) *for every $x \leq y \in P$, we have $f(x) \leq f(y)$,*
- (2) *for every $x \in P$, we have $x \leq f(x)$ (or alternatively $x \geq f(x)$),*

then the inclusion induces a homotopy equivalence $|Q| \simeq |P|$.

Proof. Suppose we have posets P, Q and a morphism $f : P \rightarrow Q$ satisfying the assumptions. We have an inclusion

$$\iota : Q \hookrightarrow P.$$

For an $x \in Q$, $\iota(x) = x$. So, for every $x \in Q$,

$$f \circ \iota(x) = f(x) \geq x = \text{Id}_Q(x).$$

Therefore $f \circ \iota \geq \text{Id}_Q$. So $|f||\iota| = |f \circ \iota| \simeq \text{Id}_{|Q|}$. On the other hand, for $x \in P$, $f(x) \in Q$, so

$$\iota \circ f(x) = f(x) \geq x = \text{Id}_P(x).$$

So, $f \circ \iota \geq \text{Id}_P$. So $|\iota||f| = |\iota \circ f| \simeq \text{Id}_{|P|}$. □

Denote by G'_p the Frattini subgroup of G , i.e., the subgroup generated by the commutator subgroup and p -th powers.

Proposition 9. *Suppose G is a p -group. Then for any G -spectrum E , we have*

$$\Phi^G(E) \simeq \Phi^{G^{ab}/p}(E^{G'_p}).$$

In particular, for a constant Mackey functor \underline{A} , we have

$$\Phi^G(\underline{A}) \simeq \Phi^{G^{ab}/p}(\underline{A}).$$

Comment. Note that one always has an equivalence

$$\Phi^G(E) \simeq \Phi^{G/H} \Phi^H E.$$

The special feature here is that $E^H \rightarrow \Phi^H E$ induces an equivalence on G/H -geometric fixed points if G is a p -group and H is the Frattini subgroup.

We shall first prove:

Lemma 10. *Let G be a p -group and let $H \subseteq G$ be a subgroup not containing G'_p . Then*

$$|P_H^{\mathcal{F}[G]}| \simeq *.$$

Proof. Denote by Q the poset of proper subgroups of G containing $G'_p \cdot H$. We have an inclusion $Q \subseteq P_H^{\mathcal{F}[G]}$. By the Burnside basis theorem, for any subgroup $K \subsetneq G$, we have $K \cdot G'_p \subsetneq G$. Thus, we have a map of posets $\varphi : P_H^{\mathcal{F}[G]} \rightarrow Q$ given by

$$K \mapsto K \cdot G'_p.$$

Also, for $K \in Q$, $\varphi(K) \supseteq K$. Thus, $|Q| \simeq |P_H^{\mathcal{F}[G]}|$ by [Lemma 8](#).

However, $|Q| \simeq *$ since Q has a minimal element. □

Note that this implies [Proposition 9](#), since the quotient map

$$(12) \quad \widetilde{E}^{\mathcal{F}}[G] \rightarrow \widetilde{E}^{\mathcal{F}}[G/G'_p]$$

induces an isomorphism on E^1 -terms of the spectral sequence [\(5\)](#). (The spectral sequence is not functorial in general with respect to change of groups. In the present case, however, we have a surjection of groups which preserves the height of the proper subgroup containing G'_p , while the remaining terms in the source are 0. Thus, a morphism of spectral sequences which is an isomorphism on E^1 arises.)

Also note that for the present purpose, the spectral sequence can be skipped entirely and one can simply argue that [\(12\)](#) is an equivalence by examining its H -fixed points for each H : The fixed point set of the left-hand side is contractible for $H \subsetneq G$ and equivalent to S^0 if $H = G$, while the right-hand side is contractible if $H \cdot G'_p \subsetneq G$ and equivalent to S^0 if $H \cdot G'_p = G$. By the Burnside basis theorem, both conditions are equivalent. This was pointed out to me during the process of revising this paper.

Proposition 11. *If G is not a p -group, then*

$$\Phi^G(\underline{A}) = 0.$$

Proof. First, suppose G is a finite group that is not a p -group.

The spectrum $\Phi^G(\mathbb{Z})$ is a commutative ring spectrum, since we have

$$S^{\infty V} \wedge S^{\infty V} \cong S^{\infty V}.$$

Choose a prime p . Then by the first Sylow theorem, there exists a p -Sylow subgroup P of G . Then there exists an H with $P \subseteq H \subsetneq G$ that is maximal (i.e., there does not exist K such that $P \subseteq H \subsetneq K \subsetneq G$). Therefore the contribution of H to the spectral sequence will include

$$(13) \quad \widetilde{H}_0^{W(H)}(\widetilde{\mathcal{O}}) \simeq H_0^{W(H)}(*) = \mathbb{Z}.$$

In (13), $1 \in \mathbb{Z}$ represents an element $\eta \in E_{1,0}^1$, where

$$d^1(\eta) = \pm \frac{|G|}{|H|} \in \mathbb{Z} = E_{0,0}^1.$$

Since we have $p \nmid |G|/|H|$, the gcd of all these numbers is 1, and thus, $1 \in \mathbb{Z} = E_{0,0}^1$ of the spectral sequence (8) is killed. Since $\Phi_*^G(\mathbb{Z})$ is a commutative ring, it must be 0. So

$$\Phi^G(\mathbb{Z}) = 0. \quad \square$$

Now, one can apply the results of my previous paper [Kriz 2015], as well as those of [Holler and Kriz 2017; 2020], to calculate $\Phi^{(\mathbb{Z}/p)^n}(\mathbb{Z})$.

Recall that for any space or spectrum X , we can obtain maps

$$(14) \quad H^n(X; \mathbb{Z}/p) \xrightarrow{\beta} H^{n+1}(X; \mathbb{Z}/p),$$

$$(15) \quad H^n(X; \mathbb{Z}/p) \xrightarrow{\beta} H^{n+1}(X; \mathbb{Z})$$

as the connecting maps of the long exact sequences from taking cohomology with coefficients in the following respective short exact sequences:

$$0 \rightarrow \mathbb{Z}/p \rightarrow \mathbb{Z}/(p^2) \rightarrow \mathbb{Z}/p \rightarrow 0,$$

$$0 \rightarrow \mathbb{Z} \xrightarrow{\cdot p} \mathbb{Z} \rightarrow \mathbb{Z}/p \rightarrow 0.$$

These are called the Bockstein maps, and the long exact sequence involving (15) forms an exact couple which gives rise to the Bockstein spectral sequence, in which (14) is d^1 .

Consider first the case of $p > 2$. Recalling the notation of Theorems 1 and 2, the Bockstein acts by

$$\beta(dz_i) = z_i, \quad \beta z_i = 0.$$

Also recall that we have

$$(16) \quad \beta(ab) = \beta a \cdot b + (-1)^{|a|} a \cdot \beta b.$$

Thus we get

$$\beta(t_\alpha) = 0, \quad \beta(u_\alpha) = 1$$

Note that β preserves the relations of (2). For example,

$$\begin{aligned} & \beta(-u_\beta u_{\alpha+\beta} + u_\alpha u_{\alpha+\beta} - u_\alpha u_\beta) \\ &= -\beta(u_\beta)u_{\alpha+\beta} + u_\beta \beta(u_{\alpha+\beta}) + \beta(u_\alpha)u_{\alpha+\beta} - u_\alpha \beta(u_{\alpha+\beta}) - \beta(u_\alpha)u_\beta + u_\alpha \beta(u_\beta) = 0. \end{aligned}$$

Now by computing directly on the chain level the equivariant $(\mathbb{Z}/p)^n$ -homology of $S^{\infty V}$, where V is the reduced regular representation, the standard $(\mathbb{Z}/p)^n$ -CW decomposition of $S^{\infty V}$ (thought of as a colimit of smash products of representation spheres of nontrivial 1-dimensional complex representations α) has only one 0-cell beside the base point, to which (using one α) p 1-cells are attached on which $(\mathbb{Z}/p)^n$ acts transitively. Thus, $\Phi_0^{(\mathbb{Z}/p)^n}(\mathbb{Z}) = \mathbb{Z}/p$. Since $\Phi_*^{(\mathbb{Z}/p)^n}(\mathbb{Z})$ is a ring, it has characteristic p (i.e., every element is annihilated by p). Thus the Bockstein spectral sequence collapses to E^2 , or in other words

$$(17) \quad H_*(\Phi_*^{(\mathbb{Z}/p)^n}(\mathbb{Z}/p), \beta) = 0.$$

Hence, we have an exact sequence

$$(18) \quad 0 \rightarrow \Phi_*^{(\mathbb{Z}/p)^n}(\mathbb{Z}) \rightarrow \Phi_*^{(\mathbb{Z}/p)^n}(\mathbb{Z}/p) \xrightarrow{\beta} \Phi_*^{(\mathbb{Z}/p)^n}(\mathbb{Z}/p).$$

Therefore, $\Phi_*^{(\mathbb{Z}/p)^n}(\mathbb{Z})$ contains the elements t_α and $\sum_i a_i u_{\alpha_i}$, where $\sum_i a_i = 0 \in \mathbb{Z}/p$. Choosing an $\alpha_0 \in (\mathbb{Z}/p)^n \setminus \{0\}$, since we are in characteristic p , the elements $\sum_i a_i u_{\alpha_i}$ where $\sum_i a_i = 0 \in \mathbb{Z}/p$ are linear combinations of $\tilde{u}_\alpha = u_\alpha - u_{\alpha_0}$. One easily verifies the relations (3). For example,

$$\begin{aligned} & -\tilde{u}_\beta \tilde{u}_{\alpha+\beta} + \tilde{u}_\alpha \tilde{u}_{\alpha+\beta} - \tilde{u}_\alpha \tilde{u}_\beta \\ &= -(u_\beta - u_{\alpha_0})(u_{\alpha+\beta} - u_{\alpha_0}) + (u_\alpha - u_{\alpha_0})(u_{\alpha+\beta} - u_{\alpha_0}) - (u_\alpha - u_{\alpha_0})(u_\beta - u_{\alpha_0}) \\ &= -u_\beta u_{\alpha+\beta} + u_{\alpha_0} u_{\alpha+\beta} + u_\beta u_{\alpha_0} - u_{\alpha_0}^2 + u_\alpha u_{\alpha+\beta} - u_{\alpha_0} u_{\alpha+\beta} - u_\alpha u_{\alpha_0} \\ & \quad + u_{\alpha_0}^2 - u_\alpha u_\beta + u_{\alpha_0} u_\beta + u_\alpha u_{\alpha_0} - u_{\alpha_0}^2 \\ &= 0. \end{aligned}$$

Let \tilde{R}_n denote the ring $\mathbb{Z}/p[t_\alpha, \tilde{u}_\alpha]$ modulo the relations (3). Write (18) as

$$0 \rightarrow R_{\mathbb{Z}} \rightarrow R_{\mathbb{Z}/p} \xrightarrow{\beta} R_{\mathbb{Z}/p}.$$

We therefore have a homomorphism of rings

$$\varphi : \tilde{R}_n \rightarrow R_{\mathbb{Z}}.$$

We want to prove that this is an isomorphism.

Now, let us consider $p = 2$. Choose again a representative $\alpha_0 \in (\mathbb{Z}/2)^n \setminus \{0\}$. Again, the $R_{\mathbb{Z}}$ contains elements of the form $\sum_i a_i y_i$ with $\sum_i a_i = 0$ which are

generated by $\tilde{y}_\alpha = y_\alpha - y_{\alpha_0}$. Also, the elements $t_\alpha = y_\alpha^2 \in R_{\mathbb{Z}}$ (by (16)), but note that $\tilde{y}_\alpha^2 = y_\alpha^2 + y_{\alpha_0}^2$ (since we are in characteristic 2), so we only need to include t_{α_0} in the generators. Now similarly as for $p > 2$, one proves the relations.

$$(19) \quad \tilde{y}_{\alpha_0} = 0, \quad \tilde{y}_\alpha \tilde{y}_\beta + \tilde{y}_\alpha \tilde{y}_{\alpha+\beta} + \tilde{y}_\beta \tilde{y}_{\alpha+\beta} + t_{\alpha_0} = 0.$$

Let \tilde{R}_n denote the quotient of the ring $\mathbb{Z}/2[\tilde{y}_\alpha, t_{\alpha_0}]$ by relations (19).

Again, we have a homomorphism of rings

$$\begin{aligned} \varphi : \tilde{R}_n &\rightarrow R_{\mathbb{Z}}, \\ \tilde{y}_\alpha &\mapsto y_\alpha - y_{\alpha_0}, \\ t_{\alpha_0} &\mapsto y_{\alpha_0}^2. \end{aligned}$$

We can prove that φ is an isomorphism by calculating the Poincaré series of \tilde{R}_n , checking that it is the same as the Poincaré series of $R_{\mathbb{Z}}$ and exhibiting an additive basis of \tilde{R}_n that is linearly independent in $R_{\mathbb{Z}/2}$.

Recall from [Holler and Kriz 2017; 2020] that the Poincaré series of $R_{\mathbb{Z}/p}$ is

$$P(R_{\mathbb{Z}/p}) = \frac{1}{(1-x)^n} \prod_{i=1}^n (1 + (p^{i-1} - 1)x).$$

Thus, by (17), we have an exact sequence of graded \mathbb{Z}/p -vector spaces

$$0 \rightarrow R_{\mathbb{Z}} \rightarrow R_{\mathbb{Z}/p} \xrightarrow{\beta} R_{\mathbb{Z}/p}[1] \xrightarrow{\beta} R_{\mathbb{Z}}[2] \rightarrow 0$$

and hence

$$(20) \quad P(R_{\mathbb{Z}}) = \frac{1}{1+x} P(R_{\mathbb{Z}/p}) = \frac{1}{(1-x^2)(1-x)^{n-1}} \prod_{i=1}^n (1 + (p^{i-1} - 1)x).$$

For $p = 2$, we know that the Poincaré series of \tilde{R}_1 is $1/(1-x^2)$. Now we can treat the α as elements of $(\mathbb{Z}/2)^n \setminus \{0\}$ and assume that $\alpha_0 = (1, 0, \dots, 0)$. For $n = 2$, we only have $\tilde{y}_{(0,1)}, \tilde{y}_{(1,1)}, \tilde{y}_{(1,0)} = 0$. By the relations, we have $\tilde{y}_{(0,1)}\tilde{y}_{(1,1)} = t_{\alpha_0}$. Therefore this ring has additive basis $\tilde{y}_{(0,1)}^{m \geq 1} t_{\alpha_0}^{m'}$, $\tilde{y}_{(1,1)}^{k \geq 1} t_{\alpha_0}^{k'}$, and $t_{\alpha_0}^\ell$ ($m', k', \ell \geq 0$), which give the terms $x/((1-x)(1-x^2))$ twice and $1/(1-x^2)$ in the Poincaré series. Therefore, the Poincaré series of the ring is

$$P(\tilde{R}_2) = \frac{1}{1-x^2} + 2 \cdot \frac{x}{(1-x)(1-x^2)} = \frac{1+x}{(1-x)(1-x^2)}.$$

After this, we can continue by induction since the relations imply

$$P(\tilde{R}_n) = P(\tilde{R}_{n-1}) \cdot (1 + (2^{n-1} - 1)x) \cdot \frac{1}{1-x}$$

similarly as in [Holler and Kriz 2020]: The additive basis is formed by the additive basis of \tilde{R}_{n-1} times $\tilde{y}_{(0, \dots, 0, 1)}^{\geq 0}$ or $\tilde{y}_{(\alpha', 1)}^{\geq 1}$, where $\alpha' \in (\mathbb{Z}/2)^{n-1} \setminus \{0\}$. These elements

are linearly independent in $R_{\mathbb{Z}/2}$ by performing a similar induction there (which was done in [Holler and Kriz 2020]).

The case of $p > 2$ is completely analogous. We define the homomorphism of rings

$$\begin{aligned} \varphi : \widetilde{R}_n &\rightarrow R_{\mathbb{Z}}, \\ \widetilde{u}_\alpha &\mapsto u_\alpha - u_{\alpha_0}, \\ t_\alpha &\mapsto t_\alpha, \end{aligned}$$

which again we can check is a ring homomorphism by computing the relations in the target. Again, we check φ is an isomorphism by checking that for every n , the Poincaré series of \widetilde{R}_n agrees with (20). We have \widetilde{R}_1 is generated by the basis of $(t_{(1)})^{m \geq 0}$. Thus, as before, $P(\widetilde{R}_1) = 1/(1-x^2)$. This time, for every $n \geq 2$, an additive basis of \widetilde{R}_n is given by the additive basis of \widetilde{R}_{n-1} times $t_{(0, \dots, 0, 1)}^{\geq 0} \cdot \widetilde{u}_{(0, \dots, 0, 1)}^\epsilon$, where $\epsilon \in \{0, 1\}$, or times $t_{(\alpha', 1)}^{\geq 1}$, or times $t_{(\alpha', 1)}^{\geq 0} \cdot \widetilde{u}_{(\alpha', 1)}$, where $\alpha' \in (\mathbb{Z}/p)^{n-1} \setminus \{0\}$. This gives

$$P(\widetilde{R}_n) = P(\widetilde{R}_{n-1}) \cdot (1 + (p^{n-1} - 1)x) \cdot \frac{1}{1-x},$$

and we can proceed by induction.

4. Another example

For the rest of the paper, we will consider equivariant homology with constant coefficients \mathbb{Z}/p for a prime p . If $G = (\mathbb{Z}/p)^n$ is an elementary abelian group, S is a set of 1-dimensional representations (real or complex depending on whether $p = 2$ or $p > 2$), and $\gamma = \bigoplus_{\alpha \in S} \alpha$, then we completely calculated in [Kriz 2015] the $(\mathbb{Z}$ -graded) coefficients of

$$(21) \quad \widetilde{H\mathbb{Z}/p}_*^G(S^{\infty\gamma}) = \widetilde{H\mathbb{Z}/p}_*^G \widetilde{E\mathcal{F}}_\gamma.$$

By the above method, for any p -group G and any set S of nontrivial irreducible 1-dimensional representations of G/G'_p , $\gamma = \bigoplus_{\alpha \in S} \alpha$, we have

$$\widetilde{H\mathbb{Z}/p}_*^G(S^{\infty\gamma}) = \widetilde{H\mathbb{Z}/p}_*^{G/G'_p}(S^{\infty\gamma}).$$

However, for a G -representation γ which does not factor through G/G'_p the calculation of (21) can be nontrivial. In this section, as an example, we consider the case where G is the split extraspecial group (as described below) at $p = 2$ and V is the irreducible real representation nontrivial on the center. This group is the central product of n copies of D_8 . We write $V_n = (\mathbb{Z}/2 \oplus \mathbb{Z}/2)^n$, where the generators of the i -th copy of $\mathbb{Z}/2 \oplus \mathbb{Z}/2$ are denoted by $v_{1,i}, v_{2,i}$. Define $q(v_{1,i}) = q(v_{2,i}) = 0$, $q(v_{1,i} + v_{2,i}) = 1$, and let q be additive between different i summands. This is a split quadratic form on the \mathbb{F}_2 -vector space V_n with associated symplectic form

$$b(x, y) = q(x + y) + q(x) + q(y).$$

A vector subspace $W \subseteq V$ is called *isotropic* when b is 0 on W and is called *q-isotropic* if q is 0 on W . The split extraspecial group \tilde{V}_n is an extension

$$1 \rightarrow \mathbb{Z}/2 \rightarrow \tilde{V}_n \rightarrow V_n \rightarrow 1,$$

where for $v \in V_n$, $2v \neq 0$ if and only if $q(v) \neq 0$ and for $v, w \in V_n$, $vwv^{-1}w^{-1} = b(v, w)$.

Clearly \tilde{V}_n is isomorphic to a central product of n copies of D_8 and the (real) irreducible representation γ nontrivial on the center is obtained as the tensor product of the dihedral representation of the n copies of D_8 . We shall apply [Proposition 4](#) to

$$\widetilde{H\mathbb{Z}/2}_*^{\tilde{V}_n}(S^{\infty\gamma}).$$

The family \mathcal{F}_γ consists of elementary abelian subgroups of \tilde{V}_n disjoint with the center which project to q -isotropic subspaces of V_n . Two subgroups are conjugate if and only if they project to the same q -isotropic subspaces U of V_n . We shall refer to these subgroups as *decorations* of U , and call them *decorated q-isotropic subspaces*.

We shall need to consider the following modular representations of the split extraspecial 2-group \tilde{V}_n : All these representations will factor through the Frattini quotient V_n . By the representation $\underline{2}_i$, we mean a tensor product of the regular representation of $\mathbb{Z}/2\{v_{1,i}\}$ with the trivial representation on $\mathbb{Z}/2\{v_{2,i}\}$, where $v_{1,j}, v_{2,j}$ act trivially for $j \neq i$. (For counting purposes, equivalently, 1 and 2 can be reversed). The representation $\underline{3}_i$ is the kernel of the augmentation from the regular representation on $\mathbb{Z}/2\{v_{1,i}, v_{2,i}\}$ to the trivial representation (with the other coordinates also acting trivially).

Let \underline{P}_n be the poset of elementary abelian subgroups of \tilde{V}_n which project to a nontrivial q -isotropic subspace of V_n . We will refer to these subgroups as *decorated q-isotropic subspaces* of V_n .

Theorem 12. For $n > 1$, $\tilde{H}_k(|\bar{P}_n|) = 0$ except for $k = n - 1$. As a \tilde{V}_n -representation, $\mathcal{H}_n := \tilde{H}_{n-1}(|\bar{P}_n|)$ is given recursively as follows:

$$\begin{aligned} \mathcal{H}_1 &= \underline{3}_1, \\ \mathcal{H}_{n+1} &= \underline{3}_{n+1} \otimes \mathcal{H}_n \\ (22) \quad &\oplus \underline{2}_{n+1} \otimes (2^{2n-1} - 2^{n-1})\mathcal{H}_n \\ &\quad \oplus \underline{2}_{n+1} \otimes (2^{2n-1} + 2^{n-1} - 1)(2\mathcal{H}_n - \underline{2}_n 2^{2n-2} \mathcal{H}_{n-1}). \end{aligned}$$

The subtraction in the last term is to be interpreted recursively as follows: We set $\mathcal{H}_0 = 1$. Then one copy of $\underline{2}_n \mathcal{H}_{n-1}$ is “subtracted” from the first summand of (22) with n replaced by $n - 1$, to leave a copy of \mathcal{H}_{n-1} . The remaining $2^{2n-2} - 1$ copies of $\underline{2}_n \mathcal{H}_{n-1}$ are subtracted from the second summand of the formula (22) with n replaced by $n - 1$ (thus, only one copy of the $2\mathcal{H}_n$ is involved in the subtraction).

Comment. The expression for \mathcal{H}_n given by the theorem is a direct sum of representations of the form

$$\bigotimes_{i \in S_1} \underline{2}_i \otimes \bigotimes_{j \in S_2} \underline{3}_j$$

for $S_1 \cap S_2 = \emptyset$. I do not know if homology groups of \tilde{V}_n with coefficients in these representations are all completely known. For $|S_2| \leq 2$, they can be deduced from the computation of Quillen [1971].

Proof. We proceed by induction on n . In the case of $n = 1$, the isotropic subspaces are $\langle v_1 \rangle$ and $\langle v_2 \rangle$, and there are two lifts to \tilde{V}_1 , and each pair of lifts is given by one $\mathbb{Z}/2$ -summand of $V_1 = \mathbb{Z}/2 \oplus \mathbb{Z}/2$. Note that we have $\underline{3}$, because we must consider reduced homology.

Now to pass from $|\bar{P}_n|$ to $|\bar{P}_{n+1}|$, consider first the poset \bar{Q}_n of decorated q -isotropic subspaces of V_{n+1} which intersect nontrivially with V_n . Then the inclusion $|\bar{P}_n| \subset |\bar{Q}_n|$ is an equivalence by Lemma 8, considering the map the other way given by

$$W \mapsto W \cap V_n.$$

Now a q -isotropic subspace W of V_{n+1} with $W \cap V_n = 0$ has $\dim(W) \leq 2$. Let \bar{R}_{n+1} denote the union of \bar{Q}_n and the set of decorated isotropic subspaces $W \subset V_{n+1}$ with $\dim(W) = 2$, $W \cap V_n = 0$. Then the poset $(\bar{R}_{n+1})_{\geq W} \cap \bar{Q}_n$ for any such space W consists of copies of the poset $\underline{4}\bar{P}_{n-1}$ (here we use $\underline{4}$ to denote 4 additional independent decorations). The factors (22) correspond to the choices of W , consisting of one nonzero q -isotropic vector $w \in V_n$, and one vector w' with $q(w') = 1$, $b(w, w') = 1$. There are 2^{2n-2} choices of w' for each w . Then $W = \langle v_1 + w, v_1 + v_2 + w' \rangle$. This leads to a based cofibration

$$(23) \quad |\bar{P}_n| \rightarrow |\bar{R}_{n+1}| \rightarrow \bigvee_{4(2^{2n-1}+2^{n-1}-1)2^{2n-2}} \Sigma|\bar{P}_{n-1}|.$$

Now \bar{P}_{n+1} is the union of \bar{R}_{n+1} with the set of all decorated q -isotropic subspaces $W \subset V_{n+1}$ with $W \cap V_n = 0$, $\dim(W) = 1$. For such a space W , $(\bar{P}_{n+1})_{\geq W} \cap \bar{R}_{n+1}$ consists of

$$(24) \quad \underline{2}(2(2^{2n-1} + 2^{n-1}) + (2^{2n-1} - 2^{n-1}))$$

copies of \bar{P}_n . The $\underline{2}$ corresponds to additional decorations. The $2(2^{2n-1} + 2^{n-1})$ summands correspond to q -isotropic vectors of the form $w + v_1$, $w + v_2$, for $w \in V_n$, the $2^{2n-1} - 2^{n-1}$ summand correspond to q -isotropic vectors of the form $w + v_1 + v_2$. Thus, we obtain a based cofibration sequence

$$(25) \quad |\bar{R}_{n+1}| \rightarrow |\bar{P}_{n+1}| \rightarrow \bigvee_{\underline{2}(2(2^{2n-1}+2^{n-1})+(2^{2n-1}-2^{n-1}))} \Sigma|\bar{P}_n|.$$

Now from (23) and (25), we can easily eliminate $|\bar{R}_{n+1}|$, as we see that the copies of $\Sigma|\bar{P}_n|$ in (25) corresponding to $w = 0$ project identically to $\Sigma|\bar{P}_n| \subset \Sigma|\bar{R}_{n+1}|$ under the connecting map. Thus, we obtain a cofibration sequence of the form

$$(26) \quad |\bar{P}_{n+1}| \rightarrow \bigvee_{\underline{2}(2(2^{2n-1}+2^{n-1}-1)+2^{2n-1}-2^{n-1})} \Sigma|\bar{P}_n| \rightarrow \bigvee_{\underline{4}(2^{2n-1}+2^{n-1}-1)2^{2n-2}} \Sigma^2|\bar{P}_{n-1}|.$$

The second map (26) is shown to be onto in reduced homology using the sums of terms indicated in the statement of the theorem. (In particular, we consider, for a q -isotropic vector $w + v$, with $0 \neq w \in V_{n+1}$, all choices of vectors w' such that $\langle w + v_1, w' + v_1 + v_2 \rangle$ is q -isotropic. Note that this includes but is not equal to, for $n > 1$, all $\langle w + v_1, u \rangle$ q -isotropic.)

The dichotomy between canceling the first or second summand in (22) comes from distinguishing whether the projection of w to V_{n-1} is 0 or not. □

Comment. The same method shows that the reduced homology of the poset of undecorated q -isotropic subspaces of V_n is concentrated in degree $n - 1$ and has rank $2^{n(n-1)}$. This poset (for $n \geq 2$) is the Tits building of $\Omega_{2n}^+(2)$ (the adjoint Chevally group of type D_n at the prime 2), and this fact therefore also follows from the Solomon–Tits theorem [Solomon 1969].

The number of q -isotropic subspaces U_k of dimension k of V_n is

$$v_{n,k} = 2^{k(k-1)/2} \cdot \frac{(2^{2n-1}+2^{n-1}-1)(2^{2n-3}+2^{n-2}-1) \dots (2^{2n-2k+1}+2^{n-k}-1)}{(2^k-1)(2^{k-1}-1) \dots (2-1)}$$

The Weyl group of U_k is \widetilde{V}_{n-k} . Thus, we have proved the following:

Theorem 13. *We have*

$$\widetilde{H\mathbb{Z}/2}_0^G(S^{\infty\gamma}) = \mathbb{Z}/2.$$

For $i > 0$,

$$\widetilde{H\mathbb{Z}/2}_i^G(S^{\infty\gamma}) = v_{n,k} \bigoplus_{k=0}^n H_{i-n+k-1}(\widetilde{V}_{n-k}, \mathcal{H}_{n-k}). \quad \square$$

References

[Adams 1984] J. F. Adams, “Prerequisites (on equivariant stable homotopy) for Carlsson’s lecture”, pp. 483–532 in *Algebraic topology* (Aarhus, 1982), edited by I. Madsen and B. Oliver, Lecture Notes in Math. **1051**, Springer, 1984. [MR](#) [Zbl](#)

[Bredon 1967] G. E. Bredon, *Equivariant cohomology theories*, Lecture Notes in Mathematics **34**, Springer, 1967. [MR](#) [Zbl](#)

[tom Dieck 1970] T. tom Dieck, “Bordism of G -manifolds and integrality theorems”, *Topology* **9** (1970), 345–358. [MR](#) [Zbl](#)

[Greenlees 1985] J. P. C. Greenlees, *Adams spectral sequences in equivariant topology*, Ph.D. thesis, University of Cambridge, 1985.

- [Hesselholt and Madsen 1997] L. Hesselholt and I. Madsen, “On the K -theory of finite algebras over Witt vectors of perfect fields”, *Topology* **36**:1 (1997), 29–101. [MR](#) [Zbl](#)
- [Hill et al. 2016] M. A. Hill, M. J. Hopkins, and D. C. Ravenel, “On the nonexistence of elements of Kervaire invariant one”, *Ann. of Math. (2)* **184**:1 (2016), 1–262. [MR](#) [Zbl](#)
- [Holler and Kriz 2017] J. Holler and I. Kriz, “On $RO(G)$ -graded equivariant “ordinary” cohomology where G is a power of $\mathbb{Z}/2$ ”, *Algebr. Geom. Topol.* **17**:2 (2017), 741–763. [MR](#)
- [Holler and Kriz 2020] J. Holler and I. Kriz, “On the coefficients of $(\mathbb{Z}/p)^n$ -equivariant ordinary cohomology with coefficients in \mathbb{Z}/p ”, 2020. [arXiv](#)
- [Kriz 2015] S. Kriz, “Equivariant cohomology and the super reciprocal plane of a hyperplane arrangement”, preprint, 2015, available at <http://www-personal.umich.edu/~skriz/hyperplanes20027.pdf>.
- [Lewis et al. 1981] G. Lewis, J. P. May, and J. McClure, “Ordinary $RO(G)$ -graded cohomology”, *Bull. Amer. Math. Soc. (N.S.)* **4**:2 (1981), 208–212. [MR](#) [Zbl](#)
- [Lewis et al. 1986] L. G. Lewis, Jr., J. P. May, M. Steinberger, and J. E. McClure, *Equivariant stable homotopy theory*, Lecture Notes in Mathematics **1213**, Springer, 1986. [MR](#)
- [Quillen 1971] D. Quillen, “The mod 2 cohomology rings of extra-special 2-groups and the spinor groups”, *Math. Ann.* **194** (1971), 197–212. [MR](#) [Zbl](#)
- [Solomon 1969] L. Solomon, “The Steinberg character of a finite group with BN -pair”, pp. 213–221 in *Theory of Finite Groups* (Cambridge, MA, 1968), Benjamin, New York, 1969. [MR](#) [Zbl](#)

Received July 5, 2020. Revised November 15, 2020.

SOPHIE KRIZ
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MICHIGAN
ANN ARBOR, MI
UNITED STATES
skriz@umich.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Matthias Aschenbrenner
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
matthias@math.ucla.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Wee Teck Gan
Mathematics Department
National University of Singapore
Singapore 119076
matgwt@nus.edu.sg

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2020 is US \$520/year for the electronic version, and \$705/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 309 No. 2 December 2020

Thin subgroups isomorphic to Gromov–Piatetski-Shapiro lattices	257
SAMUEL A. BALLAS	
Value distribution properties for the Gauss maps of the immersed harmonic surfaces	267
XINGDI CHEN, ZHIXUE LIU and MIN RU	
Scattered representations of $SL(n, \mathbb{C})$	289
CHAO-PING DONG and KAYUE DANIEL WONG	
Number of singular fibres of surface fibrations over \mathbb{P}^1	313
CHENG GONG and YI GU	
Asymptotic behavior of solutions for some elliptic equations in exterior domains	333
ZONGMING GUO and ZHONGYUAN LIU	
Ulrich elements in normal simplicial affine semigroups	353
JÜRGEN HERZOG, RAHELEH JAFARI and DUMITRU I. STAMATE	
Notes on equivariant homology with constant coefficients	381
SOPHIE KRIZ	
Local normal forms for multiplicity free $U(n)$ actions on coadjoint orbits	401
JEREMY LANE	
Functional determinant on pseudo-Einstein 3-manifolds	421
ALI MAALAOUI	
Distribution of distances in positive characteristic	437
THANG PHAM and LÊ ANH VINH	
Elliptic gradient estimates for a parabolic equation with V -Laplacian and applications	453
JIAN-HONG WANG and YU ZHENG	
Optimal L^2 extension of sections from subvarieties in weakly pseudoconvex manifolds	475
XIANGYU ZHOU and LANGFENG ZHU	