

*Pacific
Journal of
Mathematics*

**LOCAL MAASS FORMS AND EICHLER–SELBERG
RELATIONS FOR NEGATIVE-WEIGHT VECTOR-VALUED
MOCK MODULAR FORMS**

JOSHUA MALES AND ANDREAS MONO

Volume 322 No. 2

February 2023

LOCAL MAASS FORMS AND EICHLER–SELBERG RELATIONS FOR NEGATIVE-WEIGHT VECTOR-VALUED MOCK MODULAR FORMS

JOSHUA MALES AND ANDREAS MONO

By comparing two different evaluations of a modified (à la Borchers) higher Siegel theta lift on even lattices of signature (r, s) , we prove Eichler–Selberg relations for a wide class of negative-weight vector-valued mock modular forms. In doing so, we detail several properties of the lift, as well as showing that it produces an infinite family of local (and locally harmonic) Maaß forms on Grassmanians in certain signatures.

1. Introduction

Theta lifts have a storied history in the literature, receiving a vast amount of attention in the past few decades with applications throughout mathematics. We are concerned with generalizations of the Siegel theta lift originally studied by Borchers in the celebrated paper [2]. The classical Siegel lift maps half-integral weight modular forms to those of integral weight, and has seen a wide number of important applications. For example, in arithmetic geometry [14; 21], deep results in number theory [10], fundamental work of Bruinier and Funke [9], among many others.

More recently, Bruinier and Schwagenscheidt [12] investigated the Siegel theta lift on Lorentzian lattices (that is, even lattices of signature $(1, n)$), and in doing so provided a construction of recurrence relations for mock modular forms of weight $\frac{3}{2}$, as well as commenting as to how one could provide a similar structure for those of weight $\frac{1}{2}$, thereby including Ramanujan’s classical mock theta functions.

In the last few years, several authors have also considered so-called “higher” Siegel theta lifts of the shape $(k := \frac{1}{2}(1 - n), j \in \mathbb{N}_0)$

$$\int_{\mathcal{F}}^{\text{reg}} \langle R_{k-2j}^j f, \overline{\Theta_L(\tau, z)} \rangle v^k d\mu(\tau),$$

The research conducted by Males for this paper is supported by the Pacific Institute for the Mathematical Sciences (PIMS). The research and findings may not reflect those of the institute. Mono was supported by the CRC/TRR 191 “Symplectic Structures in Geometry, Algebra and Dynamics”, funded by the DFG (project number 281071066).

MSC2020: primary 11F27; secondary 11F37.

Keywords: higher Siegel theta lift, Eichler–Selberg relations, local Maass forms, vector-valued mock modular forms.

where $R_\kappa^n := R_{n-2} \circ R_{n-4} \circ \dots \circ R_\kappa$ is an iterated version of the Maaß raising operator $R_\kappa := 2i \frac{\partial}{\partial \tau} + \frac{\kappa}{v}$, f is weight $k - 2j$ harmonic Maaß form, and Θ_L is the standard Siegel theta function associated to an even lattice L of signature $(1, n)$. Here and throughout, $\tau = u + iv \in \mathbb{H}$ and $z \in \text{Gr}(L)$, the Grassmanian of L . Furthermore, $\langle \cdot, \cdot \rangle$ denotes the natural bilinear pairing. For example, they were considered by Bruinier and Ono (for $k = 0, j = 1$) in the influential work [11], by Bruinier, Ehlen and Yang in the breakthrough paper [8] in relation to the Gross–Zagier conjecture, and by Alfes-Neumann, Bringmann, Males and Schwagenscheidt in [1] for $n = 2$ and generic j .

In [32], Mertens investigated the classical Hurwitz class numbers, denoted by $H(n)$ for $n \in \mathbb{N}$. Using techniques in (scalar-valued) mock modular forms, he gave an infinite family of class number relations for odd n , two of which are

$$(1-1) \quad \sum_{s \in \mathbb{Z}} H(n - s^2) + \lambda_1(n) = \frac{1}{3} \sigma_1(n), \quad \sum_{s \in \mathbb{Z}} (4s^2 - n)H(n - s^2) + \lambda_3(n) = 0,$$

where $\lambda_k(n) = \frac{1}{2} \sum_{d|n} \min(d, \frac{n}{d})^k$ and σ_k is the usual k -th power divisor function. Because of their close similarity to the classical formula of Kronecker [28] and Hurwitz [24; 25]

$$\sum_{s \in \mathbb{Z}} H(n - s^2) - 2\lambda_1(n) = 2\sigma_1(n),$$

and those arising from the Eichler–Selberg trace formula, Mertens referred to the relationships (1-1) as *Eichler–Selberg relations*. More generally, let $[\cdot, \cdot]_\nu$ denote the ν -th Rankin–Cohen bracket (see Section 2). In general, the Rankin–Cohen bracket $[f, g]$ is a mixed mock modular form of degree ν . It is of inherent interest to determine its natural completion, say Λ , to a holomorphic modular form. Then following Mertens [33], we say that a (mock-) modular form f satisfies an Eichler–Selberg relation if there exists some holomorphic modular form g and some form Λ such that

$$[f, g]_\nu + \Lambda$$

is a holomorphic modular form. In the influential paper [33], Mertens showed the beautiful result that all mock-modular forms of weight $\frac{3}{2}$ with holomorphic shadow satisfy Eichler–Selberg relations, using the powerful theory of holomorphic projection and the Serre–Stark theorem stating that unary theta series form a basis for the spaces of holomorphic modular forms of the dual weight $\frac{1}{2}$.¹ In particular, Mertens explicitly describes the form Λ which completes the Rankin–Cohen brackets.

Following previous examples, to demonstrate the statement, let \mathcal{H} denote the generating function of Hurwitz class numbers, let $\vartheta = \sum_{n \in \mathbb{Z}} q^{n^2}$, where $\tau \in \mathbb{H}$, and

¹Mertens also provides results for mock theta functions in weight $\frac{1}{2}$, but since there is no analogue of Serre–Stark in the dual weight $\frac{3}{2}$ this is a real restriction.

$q^n = e^{2\pi i n \tau}$ throughout. Then Mertens’ results show that [33, p. 377]

$$[\mathcal{H}, \vartheta]_\nu + 2^{-2\nu-1} \binom{2\nu}{\nu} \left(\sum_{r \geq 1} 2 \sum_{\substack{m^2 - n^2 = r \\ m, n \geq 1}} (m - n)^{2\nu-1} q^r + \sum_{r \geq 1} r^{2\nu+1} q^r \right)$$

is a holomorphic modular form of weight $2\nu + 2$ for all $\nu \geq 1$, and a quasimodular form of weight 2 if $\nu = 0$.

In [31], Males combined techniques of [1; 12] during a further investigation of the higher Siegel lift on Lorentzian lattices. This lift was shown to be central in producing certain Eichler–Selberg relations in the vector-valued case, providing an analogue of the scalar-valued weight $\frac{3}{2}$ case of Mertens. We remark that the shape of the form Λ in the case of signature $(1, 1)$ is very close to that of Mertens (see [31, Theorem 1.1]), though we do not recall it here to save on complicated definitions in the introduction.

In the current paper, we develop the theory for even generic signature (r, s) lattices L and more general modified Siegel theta functions as in Borcherds [2], and consider the lift

$$\Psi_j^{\text{reg}}(f, z) := \int_{\mathcal{F}}^{\text{reg}} \langle R_{k-2j}^j(f)(\tau), \overline{\Theta_L(\tau, \psi, p)} \rangle v^k d\mu(\tau),$$

where Θ_L is a modified Siegel theta function as in Borcherds [2], essentially obtained by including a certain polynomial p in the summand of the usual vector-valued Siegel theta function. We require p to be homogenous and spherical of degree $d^+ \in \mathbb{N}_0$ in the first r variables, and $d^- \in \mathbb{N}_0$ in the last s variables (see (2-2) for precise definitions). Here, ψ is an isometry which in turn defines z ; see (2-3). Modifying the theta function in this way preserves modular properties of Θ_L , while allowing us to obtain different weights of output functions. Furthermore, since the case $j = 0$ is well-understood in the literature, we assume throughout that $j > 0$. We remark that the signature $(1, 2)$ with $j = 0$ case has also been studied in [16; 17].

In particular, we evaluate the higher lift in the now-standard ways of unfolding in Corollary 3.2, as well as recognizing it as a constant term in the Fourier expansion of the Rankin–Cohen bracket of a holomorphic modular form and a theta function (up to a boundary integral that vanishes for a certain class of input functions) in Theorem 3.3. For the second of these theorems, we use that at special points w , one may define positive- and negative-definite sublattices $P := L \cap w$ and $N := L \cap w^\perp$. In the simplest case, which we assume for the introduction, we have that $L = P \oplus N$. Then the theta series splits as $\Theta_L = \Theta_P \otimes \Theta_N$, where Θ_P is a positive-definite theta series and Θ_N a negative-definite one. Then we let \mathcal{G}_P^+ be the holomorphic part of a preimage of Θ_P under $\xi_\kappa := 2i v^\kappa \frac{\partial}{\partial \bar{\tau}}$. For the sake of simplicity, we assume that $\mathcal{G}_P^+ + g$ in the statement of Theorem 1.1 is bounded at $i\infty$ in the introduction; we overcome this assumption in Theorem 3.4 and offer a precise relation there.

Following the ideas of [31], by comparing these two evaluations of our lift and invoking Serre duality, we obtain the following theorem.

Theorem 1.1. *Let L be an even lattice of signature (r, s) , with associated Weil representation ρ_L . Let g be any holomorphic vector-valued modular form of weight $2 - (\frac{r}{2} + d^+)$ for ρ_L . Suppose that $\mathcal{G}_P^+ + g$ is bounded at $i\infty$. Then $\mathcal{G}_P^+ + g$ satisfies an explicit Eichler–Selberg relation. In particular, the form Λ is explicitly determined.*

The concept of so-called locally harmonic Maaß forms was introduced by Bringmann, Kane and Kohlen in [4]. These are functions that behave like classical harmonic Maaß forms, except for an exceptional set of density zero, where they have jump singularities. Since their inception, locally harmonic Maaß forms have seen applications throughout number theory, for example, in relation to central values of L -functions of elliptic curves [20], as well as traces of cycle integrals and periods of meromorphic modular forms [1; 30] among many others. Examples of such locally harmonic Maaß forms are usually achieved in the literature via similar theta lift machinery to that studied here. In addition to the direction of Theorem 1.1, we also discuss the action of the Laplace–Beltrami operator on the lift Ψ_j^{reg} in Theorem 4.2. In doing so, we prove the following theorem, thereby providing an infinite family of local Maaß forms (and locally harmonic Maaß forms) in signatures $(2, s)$. To state the result, we let $F_{m,k-2j,s}$ be a Maaß–Poincaré series as defined in (2-1).

Theorem 1.2. *Let L be an even isotropic lattice of signature $(2, s)$. Then the lift $\Psi_j^{\text{reg}}(F_{m,k-2j,s}, z)$ is a local Maaß form on $\text{Gr}(L)$ with eigenvalue $(\mathfrak{s} - \frac{k}{2})(1 - \mathfrak{s} - \frac{k}{2})$ under the Laplace–Beltrami operator.*

We provide an example of an input function to our lift. To this end, we specialize our setting to signature $(1, 2)$, in which case vector-valued modular forms can be identified with the usual scalar-valued framework on the complex upper half-plane, and in particular $\text{Gr}(L) \cong \mathbb{H}$. (We explain the required choices in Section 5.) In 1975, Cohen [15] defined the generalized class numbers

$$\begin{aligned}
 &H(\ell - 1, |D|) \\
 &:= \begin{cases} 0 & \text{if } D \not\equiv 0, 1 \pmod{4}, \\ \zeta(3 - 2\ell) & \text{if } D = 0, \\ L(2 - \ell, (\frac{D_0}{\cdot})) \sum_{d|j} \mu(d) (\frac{D_0}{d}) d^{\ell-2} \sigma_{2\ell-3}(\frac{j}{d}), & \text{else,} \end{cases}
 \end{aligned}$$

where $D = D_0 j^2$, as well as their generating functions

$$\mathcal{H}_\ell(\tau) := \sum_{n \geq 0} H(\ell, n) q^n, \quad \ell \in \mathbb{N} \setminus \{1\}.$$

Here, ζ refers to the Riemann zeta function, $L(s, \chi)$ to the Dirichlet L -function twisted by a Dirichlet character χ , and μ is the Möbius function. The functions \mathcal{H}_ℓ

are known as Cohen–Eisenstein series today, and can be viewed as half integral weight analogues of the classical integral weight Eisenstein series. Note that the numbers $H(2, n)$ are precisely the Hurwitz class numbers introduced above, and $\mathcal{H}_2 = \mathcal{H}$. Cohen proved that $\mathcal{H}_\ell \in M_{\ell-(1/2)}(\Gamma_0(4))$, the space of scalar-valued modular forms of weight $\frac{1}{2}$ on the usual congruence subgroup $\Gamma_0(4)$, and the coefficients satisfy Kohnen’s plus space condition by definition. (See [6, (2.13)–(2.15), Corollary 2.25] for more details on this.)

However, evaluating our lift requires negative weight and a nonconstant principal part of the input function. To overcome both obstructions, we let

$$f_{-2\ell, N}(\tau) = q^{-N} + \sum_{n>m} c_{-2\ell}(N, n) q^n, \quad N \geq -m,$$

$$m := \begin{cases} \lfloor \frac{-2\ell}{12} \rfloor - 1 & \text{if } -2\ell \equiv 2 \pmod{12}, \\ \lfloor \frac{-2\ell}{12} \rfloor, & \text{else} \end{cases}$$

be the unique weakly holomorphic modular form of weight -2ℓ for $SL_2(\mathbb{Z})$ with such a Fourier expansion, an explicit description of $f_{-2\ell, N}$ was given by Duke and Jenkins [18], and by Duke, Imamoğlu and Tóth [19, Theorem 1]. Our machinery now enables us to obtain Eichler–Selberg relations for the weakly holomorphic function $f_{-2\ell, N}(\tau) \mathcal{H}_\ell(\tau)$ along the lines of [15, Section 6], as well as the following variant of Theorem 1.2.

Theorem 1.3. *The lift $\Psi_j^{\text{reg}}(f_{-2\ell, N} \mathcal{H}_\ell, z)$ is a local Maaß form on \mathbb{H} for every $j \in \mathbb{N}$, $\ell \in \mathbb{N} \setminus \{1\}$, and $-m \leq N \in \mathbb{N}$ with exceptional set given by the net of Heegner geodesics*

$$\bigcup_{D=1}^N \{z = x + iy \in \mathbb{H} : \exists a, b, c \in \mathbb{Z}, b^2 - 4ac = D, a|z|^2 + bx + c = 0\}.$$

Remarks. (1) Theorem 1.3 generalizes immediately to any weakly holomorphic modular form g . The exceptional set is given by the union of geodesics of discriminant $D > 0$, for which the coefficient of g at q^{-D} is nonzero.

(2) Recently, Wagner [37] constructed a pullback of \mathcal{H}_ℓ under the ξ -operator, namely a harmonic Maaß form \mathcal{H}_ℓ of weight $-\ell + \frac{1}{2}$ on $\Gamma_0(4)$ that satisfies $\xi_{(1/2)-\ell} \mathcal{H}_\ell = \mathcal{H}_{\ell+2}$. An explicit definition of \mathcal{H}_ℓ can be found in [37, (1.5), (1.6)]. However, \mathcal{H}_ℓ is a harmonic Maaß form with noncuspidal image under ξ , and we restrict ourselves to a more restrictive growth condition in the discussion of Maaß forms (see Section 2) to ensure convergence of our lift. It would be interesting to investigate different regularizations of our lift, and in particular, lift the function \mathcal{H}_ℓ .

The paper is organized as follows. We establish the overall framework in Section 2. Section 3 is devoted to two evaluations of our theta lift and to the

proof of Theorem 1.1. In Section 4, we compute the action of the Laplace–Beltrami operator on our theta lift and prove Theorem 1.2. Lastly, Section 5 offers more details on the specialization to signature (1, 2), a proof of Theorem 1.3, and an indication on Eichler–Selberg relations for Cohen–Eisenstein series at the very end.

2. Preliminaries

We summarize some facts, which we require throughout.

The Weil representation. We recall the metaplectic double cover

$$\tilde{\Gamma} := \text{Mp}_2(\mathbb{Z}) := \left\{ (\gamma, \phi) : \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}), \phi : \mathbb{H} \rightarrow \mathbb{C} \text{ holomorphic, } \phi^2(\tau) = c\tau + d \right\}$$

of $\text{SL}_2(\mathbb{Z})$, which is generated by the pairs

$$\tilde{T} := \left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, 1 \right), \quad \tilde{S} := \left(\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \sqrt{\tau} \right),$$

where we fix a suitable branch of the complex square root throughout. Furthermore, we define $\tilde{\Gamma}_\infty$ as the subgroup generated by \tilde{T} .

We let L be an even lattice of signature (r, s) , and Q be a quadratic form on L with associated bilinear form $(\cdot, \cdot)_Q$. Moreover, we denote the dual lattice of L by L' , and the group ring of L'/L by $\mathbb{C}[L'/L]$. The group ring $\mathbb{C}[L'/L]$ has a standard basis, whose elements will be called ϵ_μ for $\mu \in L'/L$. We recall that there is a natural bilinear form $\langle \cdot, \cdot \rangle$ on $\mathbb{C}[L'/L]$ defined by $\langle \epsilon_\mu, \epsilon_\nu \rangle = \delta_{\mu, \nu}$.

Equipped with this structure, the Weil representation ρ_L of $\tilde{\Gamma}$ associated to L is defined on the generators by

$$\rho_L(\tilde{T})(\epsilon_\mu) := e(Q(\mu))\epsilon_\mu, \quad \rho_L(\tilde{S})(\epsilon_\mu) := \frac{e\left(\frac{1}{8}(s-r)\right)}{\sqrt{|L'/L|}} \sum_{\nu \in L'/L} e(-(\nu, \mu)_Q)\epsilon_\nu,$$

where we stipulate $e(x) := e^{2\pi i x}$ throughout. We let $L^- := (L, -Q)$ and call ρ_{L^-} the dual Weil representation of L .

The generalized upper half-plane and the invariant Laplacian. We follow the introduction in [7, Sections 3.2, 4.1], and let the signature of L be $(2, s)$ here. We assume that L is isotropic, i.e., it contains a nontrivial vector x of norm 0, and by rescaling we may assume that it is primitive, that is if $x = cy$ for some $y \in L$ and $c \in \mathbb{Z}$ then $c = \pm 1$. Note that for $s \geq 3$ all lattices contain such an isotropic vector (see [2, Section 8]).

Let $z \in L$ be a primitive norm 0 vector and $z' \in L'$ with $(z, z')_Q = 1$. Let $K := L \cap z^\perp \cap z'^\perp$. Let $d \in K$ be a primitive norm 0 vector, and $d' \in K'$ with

$(d, d')_Q = 1$. It follows that $D := K \cap d^\perp \cap d'^\perp$ is a negative-definite lattice, and we write

$$Z = (d' - Q(d')d)z_1 + z_2d + z_3d_3 + \cdots + z_\ell d_\ell =: (z_1, z_2, \dots, z_\ell) \in K \otimes \mathbb{C},$$

since $z_3d_3 + \cdots + z_\ell d_\ell \in D \otimes \mathbb{C}$. Each z_j has a real part x_j and an imaginary part y_j , and we note that

$$Q(Y) := Q(y_1, \dots, y_\ell) = y_1y_2 - y_3^2 - y_4^2 - \cdots - y_\ell^2.$$

This gives rise to the generalized upper half-plane

$$\mathbb{H}_\ell := \{Z \in K \otimes \mathbb{C} : y_1 > 0, Q(Y) > 0\} \cong \text{Gr}(L).$$

Letting

$$\partial_\mu := \frac{\partial}{\partial z_\mu} = \frac{1}{2} \left(\frac{\partial}{\partial x_\mu} - i \frac{\partial}{\partial y_\mu} \right), \quad \bar{\partial}_\mu := \frac{\partial}{\partial \bar{z}_\mu} = \frac{1}{2} \left(\frac{\partial}{\partial x_\mu} + i \frac{\partial}{\partial y_\mu} \right),$$

it can be shown that the invariant Laplacian on \mathbb{H}_ℓ has the coordinate representation [34]

$$\Omega := \sum_{\mu, \nu=1}^{\ell} y_\mu y_\nu \partial_\mu \bar{\partial}_\nu - Q(Y) \left(\partial_1 \bar{\partial}_2 + \bar{\partial}_1 \partial_2 - \frac{1}{2} \sum_{\mu=3}^{\ell} \partial_\mu \bar{\partial}_\mu \right).$$

Maaß forms. Let $\kappa \in \frac{1}{2}\mathbb{Z}$, $(\gamma, \phi) \in \tilde{\Gamma}$ and consider a function $f : \mathbb{H} \rightarrow \mathbb{C}[L'/L]$. The modular transformation in this setting is captured by the slash-operator

$$f|_{\kappa, \rho_L}(\gamma, \phi)(\tau) := \phi(\tau)^{-2\kappa} \rho_L^{-1}(\gamma, \phi) f(\gamma\tau),$$

which leads to vector-valued Maaß forms as follows [9].

Definition. Let $f : \mathbb{H} \rightarrow \mathbb{C}[L'/L]$ be smooth. Then f is a Maaß form of weight κ with respect to ρ_L if it satisfies the following three conditions.

- (1) We have $f|_{\kappa, \rho_L}(\gamma, \phi)(\tau) = f(\tau)$ for every $\tau \in \mathbb{H}$ and every $(\gamma, \phi) \in \tilde{\Gamma}$.
- (2) The function f is an eigenfunction of the weight κ hyperbolic Laplace operator, which is explicitly given by

$$\Delta_\kappa := -v^2 \left(\frac{\partial^2}{\partial u^2} + \frac{\partial^2}{\partial v^2} \right) + i\kappa v \left(\frac{\partial}{\partial u} + i \frac{\partial}{\partial v} \right).$$

- (3) There exists a polynomial² in q denoted by $P_f : \{0 < |w| < 1\} \rightarrow \mathbb{C}[L'/L]$ such that $f(\tau) - P_f(q) \in \mathcal{O}(e^{-\varepsilon v})$ as $v \rightarrow \infty$ for some $\varepsilon > 0$.

We call f a harmonic Maaß form if the eigenvalue equals 0.

²Such a polynomial is called the principal part of f .

We write $H_{\kappa,L}$ for the vector space of harmonic Maaß forms of weight κ with respect to ρ_L , and $M_{\kappa,L}^! \subseteq H_{\kappa,L}$ for the subspace of weakly holomorphic vector valued modular forms. The subspace $S_{\kappa,L}^! \subseteq M_{\kappa,L}^!$ collects all forms that vanish at all cusps, and such forms are referred to as weakly holomorphic cusp forms.

Bruinier and Funke [9] proved that a harmonic Maaß form f of weight $\kappa \neq 1$ decomposes as a sum $f = f^+ + f^-$ of a holomorphic and a nonholomorphic part, whose Fourier expansions are of the shape

$$f^+(\tau) = \sum_{\mu \in L'/L} \sum_{\substack{n \in \mathbb{Q} \\ n \gg -\infty}} c_f^+(\mu, n) q^n \mathbf{e}_\mu,$$

$$f^-(\tau) = \sum_{\mu \in L'/L} \sum_{\substack{n \in \mathbb{Q} \\ n < 0}} c_f^-(\mu, n) \Gamma(1 - \kappa, 4\pi |n|v) q^n \mathbf{e}_\mu,$$

where $\Gamma(t, x) := \int_x^\infty u^{t-1} e^{-u} du$ denotes the incomplete gamma function.

Harmonic Maaß forms can be inspected via the action of various differential operators. We require the antiholomorphic operator

$$\xi_\kappa := 2i v^\kappa \frac{\partial}{\partial \bar{\tau}},$$

as well as the Maaß raising and lowering operators

$$R_\kappa := 2i \frac{\partial}{\partial \tau} + \frac{\kappa}{v}, \quad L_\kappa := -2i v^2 \frac{\partial}{\partial \bar{\tau}}.$$

The operator ξ_κ defines a surjective map from $H_{\kappa,L}$ to $S_{2-\kappa,L}^!$ [9]. In particular, it intertwines with the slash operator introduced above, and the space $M_{\kappa,L}^!$ is precisely the kernel of ξ_κ when restricted to $H_{\kappa,L}$. Hence, every $f \in H_{\kappa,L}$ has a cuspidal shadow in our case.

The operators R_κ and L_κ increase and decrease the weight κ by 2 respectively, but do not preserve the eigenvalue under Δ_κ . For any $n \in \mathbb{N}_0$, we let

$$R_\kappa^0 := \text{id}, \quad R_\kappa^n := R_{\kappa+2n-2} \circ \cdots \circ R_{\kappa+2} \circ R_\kappa,$$

$$L_\kappa^0 := \text{id}, \quad L_\kappa^n := L_{\kappa-2n+2} \circ \cdots \circ L_{\kappa-2} \circ L_\kappa$$

be the iterated Maaß raising and lowering operators, which increase or decrease the weight κ by $2n$.

Remark. If one relaxes the growth condition (iii) to linear exponential growth, that is, $f(\tau) \in O(e^{\varepsilon v})$ as $v \rightarrow \infty$ for some $\varepsilon > 0$, then f^- is permitted to have an additional (constant) term of the form $c_f^-(\mu, 0) v^{1-\kappa} \mathbf{e}_\mu$. In this case, ξ_κ maps such a form to a weakly holomorphic modular form instead of a weakly holomorphic cusp form.

Local Maaß forms. Locally harmonic Maaß forms were introduced by Bringmann, Kane and Kohlen [4] for negative weights, and independently by Hövel [23] for weight 0. We generalize the exposition due to Bringmann, Kane and Kohlen here and provide a definition in our setting on Grassmannians and for arbitrary eigenvalues.

Definition. A local Maaß form of weight κ with closed exceptional set $X \subsetneq \mathbb{H}_\ell$ of measure zero is a function $f : \mathbb{H}_\ell \rightarrow \mathbb{C}[L'/L]$, which satisfies four properties:

- (1) For all $(\gamma, \phi) \in \tilde{\Gamma}$ and all $Z \in \mathbb{H}_\ell$ it holds that $f|_{\kappa, \rho_L}(\gamma, \phi)(Z) = f(Z)$.
- (2) For every $Z \in \mathbb{H}_\ell \setminus X$, there exists a neighborhood of Z , in which f is real analytic and an eigenfunction of Ω .
- (3) We have

$$f(Z) = \frac{1}{2} \lim_{\varepsilon \searrow 0} (f(Z + (i\varepsilon, 0, \dots, 0)^t) + f(Z - (i\varepsilon, 0, \dots, 0)^t))$$

for every $Z \in X$.

- (4) The function f is of at most polynomial growth towards all cusps.

Paralleling the definition of harmonic Maaß forms, we call a local Maaß form locally harmonic if the eigenvalue from the second condition is 0.

Poincaré series.

Weakly holomorphic Poincaré series. Following Knopp and Mason [27, Section 3], we let $m \in \mathbb{Z}$, $\kappa \in \frac{1}{2}\mathbb{N}$ satisfying $\kappa > 2$, $\mu \in L'/L$, and define

$$\mathbb{F}_{\mu, m, \kappa}(\tau) := \frac{1}{2} \sum_{(\gamma, \phi) \in \tilde{\Gamma}_\infty \setminus \tilde{\Gamma}} (e((m+1)\tau) \mathbf{e}_\mu)|_{\kappa, \rho_L}(\gamma, \phi).$$

Knopp and Mason [27] prove that $\mathbb{F}_{\mu, m, \kappa}$ converges absolutely, and that it defines a weakly holomorphic modular form of weight κ for ρ_L . In addition, they computed the Fourier expansion of $\mathbb{F}_{\mu, m, \kappa}$, which is of the shape

$$\mathbb{F}_{\mu, m, \kappa}(\tau) = \sum_{v \in L'/L} \left(\delta_{\mu, v} q^{m+1} + \sum_{n \geq 0} c(n) q^{n+1} \right) \mathbf{e}_v.$$

The Fourier coefficients $c(n)$ can be found in [27, Theorem 3.2] explicitly.

Maaß–Poincaré series. We recall an important example of harmonic Maaß forms. To this end, let $\kappa \in -\frac{1}{2}\mathbb{N}$, let $M_{\mu, v}$ be the usual M -Whittaker function (see [35, Section 13.14]), and define the auxiliary function

$$\mathcal{M}_{\kappa, s}(y) := |y|^{-\frac{\kappa}{2}} M_{\text{sgn}(y)\frac{\kappa}{2}, s-\frac{1}{2}}(|y|), \quad y \in \mathbb{R} \setminus \{0\}.$$

We average \mathcal{M}_κ over $\tilde{\Gamma}$ with respect to the parameters $\mu \in L'/L$, $m \in \mathbb{N} \setminus \{Q(\mu)\}$, and κ, \mathfrak{s} . This yields the vector-valued Maaß–Poincaré series [7]

$$(2-1) \quad F_{\mu,m,\kappa,\mathfrak{s}}(\tau) := \frac{1}{2\Gamma(2\mathfrak{s})} \sum_{(\gamma,\phi) \in \tilde{\Gamma}_\infty \setminus \tilde{\Gamma}} (\mathcal{M}_{\kappa,\mathfrak{s}}(4\pi m v) e(-mu) \mathfrak{e}_\mu)|_{\kappa,\rho_L}(\gamma, \phi).$$

By our choice of parameters and taking cosets, the series converges absolutely. The eigenvalue under Δ_κ is given by $(\mathfrak{s} - \frac{\kappa}{2})(1 - \mathfrak{s} - \frac{\kappa}{2})$. Hence if $\mathfrak{s} = \frac{\kappa}{2}$ or $\mathfrak{s} = 1 - \frac{\kappa}{2}$, then we have $F_{\mu,m,\kappa,\mathfrak{s}} \in H_{\kappa,L}$. The principal part of $F_{\mu,m,\kappa,\mathfrak{s}}$ is given by $e(-m\tau)(\mathfrak{e}_\mu + \mathfrak{e}_{-\mu})$ in this case, and $\xi_\kappa F_{\mu,-m,\kappa,\mathfrak{s}}$ is a weight $2 - \kappa$ cusp form.

Furthermore, the Maaß–Poincaré series have the following useful property thanks to their simple principal part.

Lemma 2.1. *Let $f \in H_{\kappa,L}$ with $\kappa \in -\frac{1}{2}\mathbb{N}$, and principal part*

$$P_f(\tau) = \sum_{\mu \in L'/L} \sum_{n < 0} c_f^+(\mu, n) e(n\tau) \mathfrak{e}_\mu \in \mathbb{C}[L'/L][e(-\tau)].$$

Then, we have

$$f(\tau) = \frac{1}{2} \sum_{\mu \in L'/L} \sum_{m > 0} c_f^+(\mu, -m) F_{\mu,m,\kappa,1-\frac{\kappa}{2}}(\tau).$$

Additionally, we require the following computational lemma, which is taken from [1, Lemma 2.1], and follows inductively from [8, Proposition 3.4].

Lemma 2.2. *For any $n \in \mathbb{N}_0$ it holds that*

$$R_\kappa^n(F_{\mu,m,\kappa,\mathfrak{s}})(\tau) = (4\pi m)^n \frac{\Gamma(\mathfrak{s} + n + \frac{\kappa}{2})}{\Gamma(\mathfrak{s} + \frac{\kappa}{2})} F_{\mu,m,\kappa+2n,\mathfrak{s}}(\tau).$$

Restriction, trace maps, and Rankin–Cohen brackets. As before, we fix an even lattice L . We let $A_{\kappa,L}$ be the space of smooth functions $f : \mathbb{H} \rightarrow \mathbb{C}[L'/L]$, which are invariant under the weight κ slash operator with respect to the representation ρ_L . Moreover, let $K \subseteq L$ be a finite index sublattice. Hence, we have $L' \subseteq K'$, and thus

$$L/K \subseteq L'/K \subseteq K'/K.$$

This induces a map

$$L'/K \rightarrow L'/L, \quad \mu \mapsto \bar{\mu}.$$

If $\mu \in K'/K$, $f \in A_{\kappa,L}$, $g \in A_{\kappa,K}$, and μ is a fixed preimage of $\bar{\mu}$ in L'/K , we define

$$(f_K)_\mu := \begin{cases} f_{\bar{\mu}} & \text{if } \mu \in L'/K, \\ 0 & \text{if } \mu \notin L'/K, \end{cases} \quad (g^L)_{\bar{\mu}} = \sum_{\alpha \in L/K} g_{\alpha+\mu}.$$

Lemma 2.3 [13, Section 3]. *In the notation above, there are two natural maps*

$$\begin{aligned} \text{res}_{L/K} : A_{\kappa,L} &\rightarrow A_{\kappa,K}, & \text{tr}_{L/K} : A_{\kappa,K} &\rightarrow A_{\kappa,L}, \\ f &\mapsto f_K, & g &\mapsto g^L, \end{aligned}$$

satisfying

$$\langle f, \bar{g}^L \rangle = \langle f_K, \bar{g} \rangle$$

for any $f \in A_{\kappa,L}$, $g \in A_{\kappa,K}$.

Let $\kappa, \ell \in \frac{1}{2}\mathbb{Z}$, $f \in A_{\kappa,K}$, $g \in A_{\ell,L}$. Writing

$$f = \sum_{\mu} f_{\mu} \mathbf{e}_{\mu}, \quad g = \sum_{\nu} g_{\nu} \mathbf{e}_{\nu}$$

and letting $n \in \mathbb{N}_0$, we define the tensor product of f and g as well as the n -th Rankin–Cohen bracket of f and g as

$$\begin{aligned} f \otimes g &:= \sum_{\mu, \nu} f_{\mu} g_{\nu} \mathbf{e}_{\mu+\nu} \in A_{\kappa+\ell, K \oplus L}, \\ [f, g]_n &:= \frac{1}{(2\pi i)^n} \sum_{\substack{r, s \geq 0 \\ r+s=n}} \frac{(-1)^r \Gamma(\kappa+n) \Gamma(\ell+n)}{\Gamma(s+1) \Gamma(\kappa+n-s) \Gamma(r+1) \Gamma(\ell+n-r)} f^{(r)} \otimes g^{(s)}, \end{aligned}$$

where $f^{(r)}$ and $g^{(s)}$ are usual higher derivatives of f and g . Then we have the following vector-valued analogue of [8, Proposition 3.6].

Lemma 2.4. *Let $f \in H_{\kappa, L_1}$ and $g \in H_{\ell, L_2}$. For $n \in \mathbb{N}_0$ it holds that*

$$\begin{aligned} (-4\pi)^n L_{\kappa+\ell+2n}([f, g]_n) &= \frac{\Gamma(\kappa+n)}{n! \Gamma(\kappa)} L_{\kappa}(f) \otimes R_{\ell}^n(g) + (-1)^n \frac{\Gamma(\ell+n)}{n! \Gamma(\ell)} R_{\kappa}^n(f) \otimes L_{\ell}(g). \end{aligned}$$

Finally, we have the following lemma, which can be verified straightforwardly (see [1, Proof of Theorem 4.1]).

Lemma 2.5. *Let h be a smooth function, g be holomorphic, and $\kappa, \ell \in \mathbb{R}$. Then it holds that*

$$R_{\ell-\kappa}(v^{\kappa} \bar{g} \otimes h) = v^{\kappa} \bar{g} \otimes R_{\ell} h.$$

Theta functions and special points. We fix an even lattice L of signature (r, s) and extend the quadratic form on L to $L \otimes \mathbb{R}$ in the natural way. We denote the orthogonal projection of $\lambda \in L + \mu$ onto the linear subspaces spanned by z and its orthogonal complement with respect to $(\cdot, \cdot)_Q$ by λ_z and $\lambda_{z^{\perp}}$ respectively. In other words, we have

$$L \otimes \mathbb{R} = z \oplus z^{\perp}, \quad \lambda = \lambda_z + \lambda_{z^{\perp}}.$$

Let $\text{Gr}(L)$ be the Grassmannian of r -dimensional subspaces of $L \otimes \mathbb{R}$. Let $Z \subseteq \text{Gr}(L)$ be the set of all such subspaces on which Q is positive definite. One can endow Z with the structure of a smooth manifold.

Let $p_r : \mathbb{R}^{r,0} \rightarrow \mathbb{C}$ and $p_s : \mathbb{R}^{0,s} \rightarrow \mathbb{C}$ be spherical polynomials, which are homogeneous of degree d^+ , $d^- \in \mathbb{N}_0$ respectively. Define

$$(2-2) \quad p := p_r \otimes p_s$$

and let $\psi : L \otimes \mathbb{R} \rightarrow \mathbb{R}^{r,s}$ be an isometry. We set

$$(2-3) \quad z := \psi^{-1}(\mathbb{R}^{r,0}) \in Z, \quad z^\perp = \psi^{-1}(\mathbb{R}^{0,s}).$$

For a positive-definite lattice (K, Q) of rank n and a homogeneous spherical polynomial p of degree d , we define the usual theta function

$$\Theta_K(\tau, \psi_K, p) := \sum_{\lambda \in K'} p(\psi_K(\lambda)) e(Q(\lambda)\tau),$$

where ψ_K is the isometry associated to K . It is a holomorphic modular form of weight $\frac{n}{2} + d$ for ρ_K . If the isometry is trivial, we write $\Theta_K(\tau, p)$.

Following Borchers [2] and Hövel [23], we define the general Siegel theta function as follows.³

Definition. Let $\tau \in \mathbb{H}$ and assume the notation above. Then we put

$$\Theta_L(\tau, \psi, p) := v^{\frac{s}{2}+d^-} \sum_{\mu \in L'/L} \sum_{\lambda \in L+\mu} p(\psi(\lambda)) e(Q(\lambda_z)\tau + Q(\lambda_{z^\perp})\bar{\tau}) \mathbf{e}_\mu.$$

One can check that the function Θ_L converges absolutely on $\mathbb{H} \times Z$. The following result is [23, Satz 1.55], which follows directly from [2, Theorem 4.1].

Lemma 2.6. *Let $(\gamma, \phi) \in \tilde{\Gamma}$. Then we have*

$$\Theta_L(\gamma\tau, \psi, p) = \phi(\tau)^{r+2d^+-(s+2d^-)} \rho_L(\gamma, \phi) \Theta_L(\tau, \psi, p).$$

Thus, we define

$$k := \frac{r-s}{2} + d^+ - d^-.$$

The following terminology is borrowed from [12].

Definition. An element $w \in \text{Gr}(L)$ is called a special point if it is defined over \mathbb{Q} , that is, $w \in L \otimes \mathbb{Q}$.

³In fact, Borchers considered a slightly more general theta function, where the polynomial p does not necessarily vanish under Δ_κ . For us however, this more general case would not yield spherical theta functions as we desire.

We observe that if w is a special point, then w^\perp is a special point as well. This yields the splitting

$$L \otimes \mathbb{Q} = w \oplus w^\perp,$$

which in turn yields the positive and negative-definite lattices

$$P := L \cap w, \quad N := L \cap w^\perp.$$

Clearly, $P \oplus N$ is a sublattice of L of finite index, and according to Lemma 2.3, the theta functions associated to both lattices are related by

$$\Theta_L = (\Theta_{P \oplus N})^L.$$

We identify $\mathbb{C}[(P \oplus N)' / (P \oplus N)]$ with $\mathbb{C}[P' / P] \otimes \mathbb{C}[N' / N]$, and let ψ_P, ψ_N be the restrictions of ψ onto P, N respectively. Consequently, we have the splitting

$$\Theta_{P \oplus N}(\tau, \psi, p) = \Theta_P(\tau, \psi_P, p_r) \otimes v^{\frac{s}{2} + d^-} \overline{\Theta_{N^-}(\tau, \psi_N, p_s)}$$

at a special point w , which can be verified straightforwardly. Furthermore, we observe that $\Theta_P(\tau, \psi_P, p_r)$ is holomorphic and of weight $\frac{r}{2} + d^+$ as a function of τ , while $v^{\frac{s}{2} + d^-} \overline{\Theta_{N^-}(\tau, \psi_N, p_s)}$ is of weight $-\frac{s}{2} - d^-$ with respect to τ .

Serre duality.

Proposition 2.7 [29, Proposition 2.5, Serre duality]. *Let L be an even lattice and $\kappa \in \frac{1}{2}\mathbb{Z}$. Assume that*

$$g(\tau) = \sum_{h \in L'/L} \sum_{n \geq 0} c_g(h, n) e(n\tau) \mathfrak{e}_h$$

is bounded at the cusp $i\infty$. Then g is a holomorphic modular form of weight κ for the Weil representation ρ_L if and only if we have

$$\sum_{h \in L'/L} \sum_{n \geq 0} c_g(h, n) c_f(h, -n) = 0$$

for every weakly holomorphic modular form f of weight $2 - \kappa$ for $\bar{\rho}_L$.

3. The theta lift

We consider the theta lift $\Psi_j^{\text{reg}}(f, z)$ and evaluate it in two different ways. Using Serre duality goes back to Borcherds [3].

Evaluation in terms of ${}_2F_1$. We begin by evaluating the higher modified lift as a series involving Gauß hypergeometric functions as follows.

Evaluating the theta lift of Maaß–Poincaré series for general spectral parameters.

Let $\mathfrak{s} \in \mathbb{C}$ be such that

$$F_{m,\kappa,\mathfrak{s}}(\tau) := \sum_{\mu \in L'/L} F_{\mu,m,\kappa,\mathfrak{s}}(\tau)$$

converges absolutely, that is, $\operatorname{Re}(\mathfrak{s}) > 1 - \frac{\kappa}{2}$.

Theorem 3.1. *We have*

$$\begin{aligned} \Psi_j^{\operatorname{reg}}(F_{m,k-2j,\mathfrak{s}}, z) &= (4\pi m)^{j+1-k-\frac{\mathfrak{s}}{2}-d^-} \frac{\Gamma(\mathfrak{s} + \frac{k}{2}) \Gamma(\frac{k+\mathfrak{s}}{2} + d^- - 1 + \mathfrak{s})}{2\Gamma(2-k+2j)\Gamma(\mathfrak{s} + \frac{k}{2} - j)} \\ &\quad \times \sum_{\mu \in L'/L} \sum_{\substack{\lambda \in L+\mu \\ Q(\lambda)=-m}} \overline{p(\psi(\lambda))} \left(\frac{Q(\lambda)}{Q(\lambda_{z^\perp})} \right)^{\frac{k+\mathfrak{s}}{2} + d^- - 1 + \mathfrak{s}} \\ &\quad \times {}_2F_1\left(k + \mathfrak{s}, \frac{k + \mathfrak{s}}{2} + d^- - 1 + \mathfrak{s}; 2\mathfrak{s}; \frac{Q(\lambda)}{Q(\lambda_{z^\perp})}\right). \end{aligned}$$

Remark. Choosing the homogeneous polynomial in the theta kernel function to be the constant function 1 and computing the action of R_{k-2j}^j on $F_{m,k-2j,\mathfrak{s}}$ by Lemma 2.2, this result becomes [7, Theorem 2.14].

Proof. We summarize the argument from [7, Theorem 2.14] for convenience of the reader. We need to evaluate

$$\Psi_j^{\operatorname{reg}}(F_{m,k-2j,\mathfrak{s}}, z) = \int_{\mathcal{F}} \langle R_{k-2j}^j(F_{m,k-2j,\mathfrak{s}})(\tau), \overline{\Theta_L(\tau, \psi, p)} \rangle v^k d\mu(\tau).$$

Consequently, we compute the action of the raising operator first, and have

$$\Psi_j^{\operatorname{reg}}(F_{m,k-2j,\mathfrak{s}}, z) = (4\pi m)^j \frac{\Gamma(\mathfrak{s} + \frac{k}{2})}{\Gamma(\mathfrak{s} + \frac{k}{2} - j)} \int_{\mathcal{F}} \langle (F_{m,k,\mathfrak{s}})(\tau), \overline{\Theta_L(\tau, \psi, p)} \rangle v^k d\mu(\tau)$$

by Lemma 2.2. Secondly, we insert the definitions of both functions and unfold the integral, obtaining

$$\begin{aligned} \Psi_j^{\operatorname{reg}}(F_{m,k-2j,\mathfrak{s}}, z) &= \frac{(4\pi m)^j \Gamma(\mathfrak{s} + \frac{k}{2})}{2\Gamma(2-k+2j)\Gamma(\mathfrak{s} + \frac{k}{2} - j)} \sum_{\mu \in L'/L} \sum_{\lambda \in L+\mu} \overline{p(\psi(\lambda))} \\ &\quad \times \int_0^1 \int_0^\infty (4\pi m v)^{-\frac{k}{2}} M_{-\frac{k}{2}, \mathfrak{s}-\frac{1}{2}}(4\pi m v) e(-mu) \\ &\quad \times \overline{e(Q(\lambda_z)\tau + Q(\lambda_{z^\perp})\bar{\tau})} v^{\frac{\mathfrak{s}}{2} + d^- + k - 2} dv du. \end{aligned}$$

Third, we compute the integral over u using that $\overline{e(w)} = e(-\bar{w})$ and that

$$\int_0^1 e(-mu) e(-Q(\lambda_z)u - Q(\lambda_{z^\perp})u) du = \begin{cases} 1 & \text{if } Q(\lambda_z) + Q(\lambda_{z^\perp}) = -m, \\ 0, & \text{else.} \end{cases}$$

Hence, we obtain

$$\begin{aligned} \Psi_j^{\text{reg}}(F_{m,k-2j,s}, z) &= \frac{(4\pi m)^{j-\frac{k}{2}} \Gamma(\mathfrak{s} + \frac{k}{2})}{2\Gamma(2-k+2j)\Gamma(\mathfrak{s} + \frac{k}{2} - j)} \sum_{\mu \in L'/L} \sum_{\substack{\lambda \in L + \mu \\ Q(\lambda) = -m}} \overline{p(\psi(\lambda))} \\ &\quad \times \int_0^\infty M_{-\frac{k}{2}, \mathfrak{s} - \frac{1}{2}}(4\pi m v) e^{-2\pi v(Q(\lambda_z) - Q(\lambda_{z^\perp}))} v^{\frac{s+k}{2} + d^- - 2} dv. \end{aligned}$$

The integral is a Laplace transform. Using that

$$\frac{m}{2m} + \frac{Q(\lambda_z) - Q(\lambda_{z^\perp})}{2m} = \frac{Q(\lambda_{z^\perp})}{Q(\lambda)}$$

along with [35, (13.23.1)], it evaluates

$$\begin{aligned} &\int_0^\infty M_{-\frac{k}{2}, \mathfrak{s} - \frac{1}{2}}(4\pi m v) e^{-2\pi v(Q(\lambda_z) - Q(\lambda_{z^\perp}))} v^{\frac{k+s}{2} + d^- - 2} dv \\ &= \frac{(4\pi m)^{1-\frac{k+s}{2}-d^-} \Gamma(\frac{k+s}{2} + d^- - 1 + \mathfrak{s})}{\left(\frac{Q(\lambda_z) - Q(\lambda_{z^\perp})}{2m} + \frac{1}{2}\right)^{\frac{k+s}{2} + d^- - 1 + \mathfrak{s}}} \\ &\quad \times {}_2F_1\left(k + \mathfrak{s}, \frac{k+s}{2} + d^- - 1 + \mathfrak{s}; 2\mathfrak{s}; \frac{1}{\frac{1}{2} + \frac{Q(\lambda_z) - Q(\lambda_{z^\perp})}{2m}}\right). \end{aligned}$$

We recall $Q(\lambda) = Q(\lambda_z) + Q(\lambda_{z^\perp}) = -m$ and rewrite the argument of the hypergeometric function to

$$\frac{m}{2m} + \frac{Q(\lambda_z) - Q(\lambda_{z^\perp})}{2m} = \frac{Q(\lambda_{z^\perp})}{Q(\lambda)}.$$

Thus, we arrive at

$$\begin{aligned} \Psi_j^{\text{reg}}(F_{m,k-2j,s}, z) &= (4\pi m)^{j+1-k-\frac{s}{2}-d^-} \frac{\Gamma(\mathfrak{s} + \frac{k}{2}) \Gamma(\frac{k+s}{2} + d^- - 1 + \mathfrak{s})}{2\Gamma(2-k+2j)\Gamma(\mathfrak{s} + \frac{k}{2} - j)} \\ &\quad \times \sum_{\mu \in L'/L} \sum_{\substack{\lambda \in L + \mu \\ Q(\lambda) = -m}} \overline{p(\psi(\lambda))} \left(\frac{Q(\lambda)}{Q(\lambda_{z^\perp})}\right)^{\frac{k+s}{2} + d^- - 1 + \mathfrak{s}} \\ &\quad \times {}_2F_1\left(k + \mathfrak{s}, \frac{k+s}{2} + d^- - 1 + \mathfrak{s}; 2\mathfrak{s}; \frac{Q(\lambda)}{Q(\lambda_{z^\perp})}\right), \end{aligned}$$

as claimed. □

Combining the previous result with Lemma 2.1 yields the following consequence.

Corollary 3.2. *Let $j \in \mathbb{N}_0$ and $f \in H_{k-2j,L}$. Assume that $k - 2j < 0$. Then we have*

$$\Psi_j^{\text{reg}}(f, z) = \frac{(4\pi)^{j+1-k-\frac{s}{2}-d^-} j! \Gamma\left(\frac{s}{2} + d^- + j\right)}{4\Gamma(2 - k + 2j)} \sum_{\substack{\lambda \in L' \\ Q(\lambda) < 0}} c_f^+(\lambda, Q(\lambda)) \overline{p(\psi(\lambda))} \\ \times \frac{|Q(\lambda)|^{2j+1-k}}{|Q(\lambda_{z^\perp})|^{\frac{s}{2}+j+d^-}} {}_2F_1\left(1 + j, \frac{s}{2} + d^- + j; 2 - k + 2j; \frac{Q(\lambda)}{Q(\lambda_{z^\perp})}\right).$$

Proof. Since the weight of f is negative, we have

$$f(\tau) = \frac{1}{2} \sum_{h \in L'/L} \sum_{m \geq 0} c_f^+(h, -m) F_{h,m,k-2j,1-\frac{k}{2}+j}(\tau)$$

according to Lemma 2.1. We observe that the term corresponding to $m = 0$ will vanish due to $c_f^+(h, 0) = 0$ by our more restrictive growth condition on Maaß forms. Consequently, we have

$$\Psi_j^{\text{reg}}(f, z) = \frac{1}{2} \sum_{\mu \in L'/L} \sum_{m > 0} c_f^+(\mu, -m) \Psi_j^{\text{reg}}(F_{\mu,m,k-2j,1-\frac{k}{2}+j}, z).$$

We insert the spectral parameter $\mathfrak{s} = 1 - \frac{k-2j}{2}$ into Theorem 3.1, which yields the claim. □

Evaluation in terms of the constant term in a Fourier expansion. Next we determine the lift as a constant term in a Fourier expansion plus a certain boundary integral that vanishes for a certain class of input function.

Theorem 3.3. *Let $f \in H_{k-2j,L}$ and w be a special point, and \mathcal{G}_P^+ be the holomorphic part of a preimage of Θ_P under $\xi_{2-(\frac{r}{2}+d^+)}$. Then we have*

$$\Psi_j^{\text{reg}}(f, w) = \frac{j! (4\pi)^j \Gamma\left(2 - \frac{r}{2} - d^+\right)}{\Gamma\left(2 - \frac{r}{2} - d^+ + j\right)} \\ \times \left(\text{CT}(\langle f_{P \oplus N}(\tau), [\mathcal{G}_P^+(\tau), \Theta_{N^-}(\tau)]_j \rangle) \right. \\ \left. - \int_{\mathcal{F}}^{\text{reg}} \langle L_{k-2j}(f_{P \oplus N})(\tau), [\mathcal{G}_P^+(\tau), \Theta_{N^-}(\tau)]_j \rangle v^{-2} d\tau \right).$$

Remark. In general, the coefficients of \mathcal{G}_P^+ are expected to be transcendental. However, in weights $\frac{1}{2}$ and $\frac{3}{2}$ the function \mathcal{G}_P^+ may be chosen to have rational coefficients — a situation which is expected to also hold for ξ -preimages of CM modular forms. It is therefore expected that one obtains rationality (up to powers of π) of the modified higher lift only in these cases, and stipulating that f is weakly holomorphic meaning that the final integral vanishes.

By a slight abuse of notation, we write $\Theta_L(\tau, w, p)$ for the theta function evaluated at an isometry ψ that produces a special point w .

Proof of Theorem 3.3. We restrict to special points $w \in \text{Gr}(L)$. So we can write

$$\langle R_{k-2j}^j(f)(\tau), \overline{\Theta_L(\tau, w, p)} \rangle = \langle R_{k-2j}^j(f_{P \oplus N})(\tau), \overline{\Theta_{P \oplus N}(\tau, w, p)} \rangle.$$

Next, we use that the raising and lowering operator are adjoint to each other (see [7, Lemma 4.2]), which gives

$$\Psi_j^{\text{reg}}(f, w) = \int_{\mathcal{F}}^{\text{reg}} \langle f_{P \oplus N}(\tau), L_k^{j-1}(\overline{\Theta_{P \oplus N}(\tau, w, p)}) \rangle v^{k-2} d\tau.$$

We observe that the boundary terms disappear in the same fashion as during the proof of [7, Lemma 4.4]. Next, we rewrite

$$\Psi_j^{\text{reg}}(f, w) = (-1)^j \int_{\mathcal{F}}^{\text{reg}} \langle f_{P \oplus N}(\tau), R_{-k}^j(\overline{\Theta_{P \oplus N}(\tau, w, p)} v^k) \rangle v^{-2} d\tau$$

and recall that

$$\Theta_{P \oplus N}(\tau, w, p) = \Theta_P(\tau, p_r) \otimes v^{\frac{s}{2}+d^-} \overline{\Theta_{N^-}(\tau, p_s)} = v^{\frac{s}{2}+d^-} \Theta_P(\tau, p_r) \otimes \overline{\Theta_{N^-}(\tau, p_s)}.$$

Consequently, we obtain

$$\begin{aligned} R_{-k}^j(\overline{\Theta_{P \oplus N}(\tau, w, p)} v^k) &= R_{-k}^j(v^{k+\frac{s}{2}+d^-} \overline{\Theta_P(\tau, p_r)} \otimes \Theta_{N^-}(\tau, p_s)) \\ &= v^{k+\frac{s}{2}+d^-} \overline{\Theta_P(\tau, p_r)} \otimes (R_{\frac{s}{2}+d^-}^j \Theta_{N^-}(\tau, p_s)) \end{aligned}$$

by Lemma 2.5. In particular, we note that $v^{k+\frac{s}{2}+d^-} \overline{\Theta_P(\tau, p_r)}$ has weight

$$-k - \frac{s}{2} - d^- = -\frac{r}{2} - d^+.$$

We choose a preimage \mathcal{G}_P of $\Theta_P(\tau, p_r)$ under $\xi_{2-(\frac{r}{2}+d^+)}$, namely

$$\Theta_P(\tau, p_r) = \xi_{2-\frac{r}{2}-d^+} \mathcal{G}_P(\tau) = v^{-\frac{r}{2}-d^+} \overline{L_{2-\frac{r}{2}-d^+} \mathcal{G}_P},$$

which yields

$$R_{-k}^j(\overline{\Theta_{P \oplus N}(\tau, w, p)} v^k) = L_{2-\frac{r}{2}-d^+} \mathcal{G}_P(\tau) \otimes (R_{\frac{s}{2}+d^-}^j \Theta_{N^-}(\tau, p_s)).$$

We apply the computation of the Rankin–Cohen brackets given in Lemma 2.4 noting that $L_\ell \Theta_{N^-} = 0$, and that it suffices to deal with the holomorphic part \mathcal{G}_P^+ of \mathcal{G}_P (both by virtue of holomorphicity in computing the Rankin–Cohen bracket). Thus,

$$\begin{aligned} &R_{-k}^j(\overline{\Theta_{P \oplus N}(\tau, w, p)} v^k) \\ &= \frac{j!(-4\pi)^j \Gamma(2-k)}{\Gamma(2-k+j)} v^{-\frac{s}{2}-d^-} L_{2-k+\frac{s}{2}+d^-+2j} [\mathcal{G}_P^+(\tau), \Theta_{N^-}(\tau, p_s)]_j. \end{aligned}$$

Hence, the theta lift becomes

$$\begin{aligned} \Psi_j^{\text{reg}}(f, w) &= \frac{j!(4\pi)^j \Gamma(2 - \frac{r}{2} - d^+)}{\Gamma(2 - \frac{r}{2} - d^+ + j)} \int_{\mathcal{F}}^{\text{reg}} \langle f_{P \oplus N}(\tau), L_{2-k+2j}[\mathcal{G}_P^+(\tau), \Theta_{N^-}(\tau, p_s)]_j \rangle v^{-2} d\tau. \end{aligned}$$

The last step is to apply Stokes’ theorem, compare the proof of [7, Lemma 4.2] for example, which yields

$$\begin{aligned} \Psi_j^{\text{reg}}(f, w) &= \frac{j!(4\pi)^j \Gamma(2 - \frac{r}{2} - d^+)}{\Gamma(2 - \frac{r}{2} - d^+ + j)} \\ &\quad \times \left(\lim_{T \rightarrow \infty} \int_{iT}^{1+iT} \langle f_{P \oplus N}(\tau), [\mathcal{G}_P^+(\tau), \Theta_{N^-}(\tau, p_s)]_j \rangle v^{-2} d\tau \right. \\ &\quad \left. - \int_{\mathcal{F}}^{\text{reg}} \langle L_{k-2j}(f_{P \oplus N})(\tau), [\mathcal{G}_P^+(\tau), \Theta_{N^-}(\tau, p_s)]_j \rangle v^{-2} d\tau \right), \end{aligned}$$

utilizing again that boundary terms vanish. We observe that the left integral can be regarded as the Fourier coefficient of index 0 in the Fourier expansion of the integrand, see the bottom of page 14 in [12]. This proves the claim. \square

We end this section by noting that to obtain recurrence relations, as in [12], one would need to compute the Fourier expansion of the lift. In general, this is a lengthy but straightforward process, and since we do not require it in this paper we omit the details. In essence, one follows the calculations of Borcherds [2] by using Lemma 2.2. A resulting technicality is to then take care of the different spectral parameter. One may overcome this by relating the coefficients of Maaß–Poincaré series to those with other spectral parameters, again using the action of the iterated Maaß raising operator as in Lemma 2.2.

Eichler–Selberg relations. We now prove a refined version of Theorem 1.1. To this end, we define the function

$$\begin{aligned} (3-1) \quad \Lambda_L(\psi, p, j) &:= \frac{(4\pi)^{1-\frac{r}{2}-d^+} \Gamma(\frac{s}{2} + j + d^-) \Gamma(2 - \frac{r}{2} - d^+ + j)}{4\Gamma(2 - k + 2j) \Gamma(2 - \frac{r}{2} - d^+)} \\ &\quad \times \sum_{\substack{m \geq 1, \lambda \in L' \\ Q(\lambda) = -m}} \overline{p(\psi(\lambda))} \frac{|Q(\lambda)|^{2j+1-k}}{|Q(\lambda_{z^\perp})|^{\frac{s}{2}+j+d^-}} \\ &\quad \times {}_2F_1\left(1 + j, \frac{s}{2} + j + d^-; 2 - k + 2j; \frac{Q(\lambda)}{Q(\lambda_{z^\perp})}\right) q^m \end{aligned}$$

for $j > 0$. We write

$$\mathcal{G}_P^+(\tau) = \sum_{\mu \in L'/L} \sum_{n \gg -\infty} a(n) q^n \mathbf{e}_\mu$$

and furthermore define

$$\mathcal{G}_P^+(\tau) := \mathcal{G}_P^+(\tau) - \sum_{\mu \in L'/L} \sum_{n < 0} a(n) \mathbb{F}_{\mu, n-1, 2j+2-k}(\tau).$$

Since one may add any weakly holomorphic modular form of appropriate weight for ρ_L to \mathcal{G}_P^+ , Theorem 1.1 follows directly from the following result (noting that the linear combination of Maaß–Poincaré series may change).

Theorem 3.4. *Let L be an even lattice of signature (r, s) , let p be as before, and w be a special point defined by the isometry ψ . Let $j > 0$ and k be such that $2j + 2 - k > 2$. Then the function*

$$[\mathcal{G}_P^+(\tau), \Theta_{N^-}(\tau, p_s)]_j^L - \Lambda_L(\psi, p, j)$$

is a holomorphic vector-valued modular form of weight $2j + 2 - k$ for ρ_L .

Remarks. (1) This provides the general vector-valued analogue, assuming that the lattice is chosen such that $2j + 2 - k > 2$, of Mertens’ scalar-valued results in weights $\frac{1}{2}$ and $\frac{3}{2}$ [33].

(2) Note that the slight correction of \mathcal{G}_P^+ by Poincaré series was missing in [31].

(3) In certain cases the hypergeometric function may be simplified (for example, the $n = 1$ case as in [12; 31], which yields a form very similar to Mertens’ scalar-valued result). It appears to be possible that one should be able to prove the same results via holomorphic projection acting on vector-valued modular forms (see [26]) in much the same way as Mertens’ original scalar valued proofs in [33].

Proof of Theorem 3.4. Let f be a weakly holomorphic form of weight $k - 2j$ with Fourier coefficients c_f^+ . By construction, the form \mathcal{G}_P^+ is holomorphic at $i\infty$, and hence

$$\text{CT}(\langle f_{P \oplus N}(\tau), [\mathcal{G}_P^+(\tau), \Theta_{N^-}(\tau, p_s)]_j^L \rangle)$$

contains only the Fourier coefficients of nonpositive index of f . We note that $L_{k-2j}f = 0$, and subtract the resulting expressions of the lift from Corollary 3.2 and Theorem 3.3. We obtain

$$\begin{aligned} 0 = & \text{CT}(\langle f_{P \oplus N}(\tau), [\mathcal{G}_P^+(\tau), \Theta_{N^-}(\tau, p_s)]_j^L \rangle) \\ & - \frac{(4\pi)^{1-\frac{r}{2}-d^+} \Gamma(\frac{s}{2} + j + d^-) \Gamma(2 - \frac{r}{2} - d^+ + j)}{4\Gamma(2 - k + 2j) \Gamma(2 - \frac{r}{2} - d^+)} \\ & \times \sum_{\substack{m \geq 1, \lambda \in L' \\ Q(\lambda) = -m}} c_f^+(\lambda, -m) \overline{p(\psi(\lambda))} \frac{|Q(\lambda)|^{2j+1-k}}{|Q(\lambda_{z^\perp})|^{\frac{s}{2}+j+d^-}} \\ & \times {}_2F_1\left(1 + j, \frac{s}{2} + j + d^-; 2 - k + 2j; \frac{Q(\lambda)}{Q(\lambda_{z^\perp})}\right). \end{aligned}$$

The Rankin–Cohen bracket is bilinear and a linear combination of vector-valued Poincaré series is modular itself. We apply Proposition 2.7 and the claim follows. \square

In a similar way to [33, Corollary 5.4], we obtain the following structural corollary by rewriting Theorem 3.4, keeping the same notation as throughout this paper.

Corollary 3.5. *Let θ denote the space generated by all Θ_{N^-} functions of weight $\frac{s}{2} + d^-$ for ρ_{N^-} . Then the equivalence classes $\Lambda_L(\psi, p, j) + M_{2j+2-k,L}^!$ generate the \mathbb{C} -vector space*

$$[\mathcal{M}_{2j+2-k,P}^{mock}, \theta]_j^L / M_{2j+2-k,L}^!$$

4. The action of the Laplace–Beltrami operator

In this section, we prove Theorem 1.3. To this end, we compute the action of the Laplace–Beltrami operator on the lift, and show that for certain spectral parameters, we obtain a local Maaß form. We recall that the signature of L is assumed here to be $(2, s)$. Moreover, we observe that our Siegel theta function Θ_L and the Siegel theta function inspected by Bruinier depend in the same way on Z , and thus the following result applies.

Proposition 4.1 [7, Proposition 4.5]. *The Siegel theta function $\Theta_L(\tau, Z, p)$ considered as a function on $\mathbb{H} \times \mathbb{H}_\ell$ satisfies the differential equation*

$$\Omega \Theta_L(\tau, Z, p) v^{\frac{\ell}{2}} = -\frac{1}{2} \Delta_k \Theta_L(\tau, Z, p) v^{\frac{\ell}{2}}.$$

Our next step is to inspect the action of Ω on our theta lift. By Lemma 2.1 it suffices to investigate

$$\Psi_j^{\text{reg}}(F_{m,k-2j,s}, Z) = \int_{\mathcal{F}}^{\text{reg}} \langle R_{k-2j}^j(F_{m,k-2j,s})(\tau), \overline{\Theta_L(\tau, Z, p)} \rangle v^k d\mu(\tau).$$

Let

$$H(m) := \bigcup_{\mu \in L'/L} \bigcup_{\substack{\lambda \in \mu + L \\ Q(\lambda) = -m}} \lambda^\perp \subseteq \text{Gr}(L),$$

which collects the singularities of $\Psi_j^{\text{reg}}(F_{m,k-2j,s}, Z)$ as a function of Z . We apply the previous proposition to our theta lift, which yields a variant of [7, Theorem 4.6].

Theorem 4.2. *Let $Z \in \mathbb{H}_\ell \setminus H(m)$ and $\text{Re}(s) > 1 - \frac{k}{2}$. Then it holds that*

$$\Omega \Psi_j^{\text{reg}}(F_{m,k-2j,s}, Z) = (s - \frac{k}{2})(1 - s - \frac{k}{2}) \Psi_j^{\text{reg}}(F_{m,k-2j,s}, Z).$$

Proof. First, we note that

$$\Omega \Psi_j^{\text{reg}}(F_{m,k-2j,s}, Z) = \int_{\mathcal{F}}^{\text{reg}} \langle R_{k-2j}^j(F_{m,k-2j,s})(\tau), \Omega \overline{\Theta_L(\tau, Z, p)} v^{\frac{\ell}{2}} \rangle v^{k-\frac{\ell}{2}} d\mu(\tau),$$

because all partial derivatives with respect to Z converge locally uniformly in Z as $T \rightarrow \infty$ (see [7, p. 99]). By the previous proposition, we infer that

$$\begin{aligned} \Omega\Psi_j^{\text{reg}}(F_{m,k-2j,s}, Z) &= -\frac{1}{2} \int_{\mathcal{F}}^{\text{reg}} \langle R_{k-2j}^j(F_{m,k-2j,s})(\tau), \Delta_k \overline{\Theta_L(\tau, Z, p)} v^{\frac{\ell}{2}} \rangle v^{k-\frac{\ell}{2}} d\mu(\tau). \end{aligned}$$

By the splitting $\Delta_k = R_{k-2} L_k$ and the adjointness of both operators (see [7, Lemmas 4.2–4.4]), we obtain

$$\begin{aligned} \Omega\Psi_j^{\text{reg}}(F_{m,k-2j,s}, Z) &= -\frac{1}{2} \int_{\mathcal{F}}^{\text{reg}} \langle \Delta_k R_{k-2j}^j(F_{m,k-2j,s})(\tau), \overline{\Theta_L(\tau, Z, p)} v^{\frac{\ell}{2}} \rangle v^{k-\frac{\ell}{2}} d\mu(\tau). \end{aligned}$$

Lastly, we observe that Δ_k and R_{k-2j}^j commute by virtue of Lemma 2.2, namely

$$\Delta_k R_{k-2j}^j(F_{m,k-2j,s})(\tau) = \left(s - \frac{k}{2}\right) \left(1 - s - \frac{k}{2}\right) R_{k-2j}^j(F_{m,k-2j,s})(\tau),$$

and this establishes the claim by rewriting

$$\begin{aligned} \langle R_{k-2j}^j(F_{m,k-2j,s})(\tau), \overline{\Theta_L(\tau, Z, p)} v^{\frac{\ell}{2}} \rangle v^{k-\frac{\ell}{2}} &= \langle R_{k-2j}^j(F_{m,k-2j,s})(\tau), \overline{\Theta_L(\tau, Z, p)} \rangle v^k \end{aligned}$$

again. □

Proof of Theorem 1.2. By Theorem 4.2, the lift is an eigenfunction of the Laplace–Beltrami operator with the quoted eigenvalue. Since $\Psi_j^{\text{reg}}(F_{m,k-2j,s}, Z)$ is an eigenfunction of an elliptic differential operator, it is real-analytic in $\text{Gr}(L)$ outside of $H(m)$. The other conditions for the lift to be a vector-valued local Maaß form can be easily seen by applying the proof of [5, Theorem 1.1] *mutatis mutandis*. When $s = \frac{k}{2}$ or $s = \frac{k}{2} - 1$, we obtain locally harmonic Maaß forms. □

5. Cohen–Eisenstein series

We specialize the framework from Section 2 following [12, Section 4.4] (or [36, Section 2.2]). We fix the signature $(1, 2)$ as mentioned in the introduction, and the rational quadratic space

$$V := \left\{ X = \begin{pmatrix} x_2 & x_1 \\ x_3 & -x_2 \end{pmatrix} \in \mathbb{Q}^{2 \times 2} \right\},$$

with quadratic form $Q(X) = \det(X)$. The Grassmannian of positive lines in $V \otimes \mathbb{R}$ can be identified with \mathbb{H} via

$$\lambda(x+iy) = \frac{1}{\sqrt{2}y} \begin{pmatrix} -x & x^2+y^2 \\ -1 & x \end{pmatrix}.$$

We choose the lattice

$$L := \left\{ \begin{pmatrix} b & c \\ -a & -b \end{pmatrix} : a, b, c \in \mathbb{Z} \right\},$$

with dual lattice

$$L' = \left\{ \begin{pmatrix} \frac{b}{2} & c \\ -a & -\frac{b}{2} \end{pmatrix} : a, b, c \in \mathbb{Z} \right\}.$$

We observe that L' can be identified with the set of integral binary quadratic forms of discriminant

$$\det \begin{pmatrix} \frac{b}{2} & c \\ -a & -\frac{b}{2} \end{pmatrix} = -\frac{1}{4}(b^2 - 4ac).$$

Furthermore, $L'/L \cong \mathbb{Z}/2\mathbb{Z}$ with quadratic form $x \mapsto -\frac{1}{4}x^2$. According to [12, p. 22], it holds that

$$\begin{aligned} \mathcal{Q} \left(\begin{pmatrix} \frac{b}{2} & c \\ -a & -\frac{b}{2} \end{pmatrix}_{x+iy} \right) &= \frac{1}{4y^2} (a(x^2 + y^2) + bx + c)^2, \\ \mathcal{Q} \left(\begin{pmatrix} \frac{b}{2} & c \\ -a & -\frac{b}{2} \end{pmatrix}_{(x+iy)^\perp} \right) &= -\frac{1}{4y^2} |[a, b, c](x + iy, 1)|^2. \end{aligned}$$

We remark that both are invariant under modular substitutions. By a result from Eichler and Zagier [22, Theorem 5.4], the space of vector-valued modular forms of weight k for ρ_L is isomorphic to the space $M_k^+(\Gamma_0(4))$ of scalar-valued modular forms satisfying the Kohnen plus space condition via the map

$$f_0(\tau) \mathbf{e}_0 + f_1(\tau) \mathbf{e}_1 \mapsto f_0(4\tau) + f_1(4\tau).$$

This enables us to use scalar-valued forms as inputs for our theta lift.

Proof of Theorem 1.3. As outlined in the introduction, the function $f(\tau) := f_{-2\ell, N}(\tau) \mathcal{H}_\ell(\tau)$ is of weight $-\ell - \frac{1}{2} < 0$ for $\Gamma_0(4)$, has nonconstant principal part at the cusp $i\infty$, and its image under ξ is trivial, and hence in particular cuspidal. This enables us to apply Corollary 3.2 to f . To this end, we have the parameters

$$k = -\frac{1}{2} + d^+ + d^-, \quad k - 2j = -\ell - \frac{1}{2}.$$

Rewriting those yields

$$j = \frac{\ell + d^+ + d^-}{2},$$

and the hypergeometric function from Theorem 3.1 becomes

$${}_2F_1 \left(\frac{\ell + 2 + d^+ + d^-}{2}, \frac{\ell + 2 + d^+ + 3d^-}{2}, \frac{5}{2} + \ell, \frac{4my^2}{|[a, b, c](z, 1)|^2} \right).$$

Inspecting the parameters, we have the condition $\ell + d^+ + d^- \in 2\mathbb{N}$ by $j \in \mathbb{N}$, and combining with $d^+, d^- \in \mathbb{N}_0$, $\ell \in \mathbb{N} \setminus \{1\}$, the smallest possible values are

$(\ell, d^+, d^-) = (2, 0, 0), (2, 2, 0), (2, 1, 1), (2, 0, 2)$. For example, the corresponding hypergeometric functions for the cases $(\ell, d^+, d^-) = (2, 0, 0), (2, 1, 1)$ are

$$\begin{aligned}
 {}_2F_1\left(2, 2, \frac{9}{2}, \tilde{z}\right) &= -\frac{35(11\tilde{z} - 15)}{12\tilde{z}^3} - \frac{35(2\tilde{z}^2 - 7\tilde{z} + 5) \arcsin(\sqrt{\tilde{z}})}{4\tilde{z}^{\frac{7}{2}}\sqrt{1-\tilde{z}}}, \\
 {}_2F_1\left(3, 4, \frac{9}{2}, \tilde{z}\right) &= -\frac{35(8\tilde{z}^2 - 26\tilde{z} + 15)}{128\tilde{z}^3(\tilde{z} - 1)^2} + \frac{105(8\tilde{z}^2 - 12\tilde{z} + 5) \arcsin(\sqrt{\tilde{z}})}{128\tilde{z}^{\frac{7}{2}}\sqrt{1-\tilde{z}}(\tilde{z} - 1)^2},
 \end{aligned}$$

and the other cases are of similar shape. Analogous expressions can be obtained for higher integer parameters via Gauß’ contiguous relations for the hypergeometric function, which can be found in [35, Section 15.5(ii)] for instance.

We infer a local behavior as sketched in the introduction by virtue of $(4m = D = b^2 - 4ac)$

$$\arcsin(\sqrt{\tilde{z}}) = \arcsin\left(\frac{\sqrt{D}y}{|az^2 + bz + c|}\right) = \arctan\left|\frac{\sqrt{D}y}{a|z|^2 + bx + c}\right|,$$

which in turn follows by

$$(b^2 - 4ac)y^2 + (a|z|^2 + bx + c)^2 = |az^2 + bz + c|^2,$$

compare [4, Section 3]. The denominator $a|z|^2 + bx + c$ vanishes if and only if z is located on the Heegner geodesic associated to $Q = [a, b, c]$. Since the principal part of f is given by

$$\sum_{n=0}^N H(\ell, n) q^{n-N} + O(q^{m+1}), \quad m = \begin{cases} \lfloor \frac{-2\ell}{12} \rfloor - 1 & \text{if } -2\ell \equiv 2 \pmod{12}, \\ \lfloor \frac{-2\ell}{12} \rfloor, & \text{else,} \end{cases}$$

we conclude that f has the exceptional set

$$\bigcup_{D=1}^N \{z = x + iy \in \mathbb{H} : \exists a, b, c \in \mathbb{Z}, b^2 - 4ac = D, a|z|^2 + bx + c = 0\}.$$

In other words, the exceptional set of f is a finite union of nets of Heegner geodesics. Furthermore, we recall that the spectral parameter in Corollary 3.2 is $\mathfrak{s} = 1 - \frac{k-2j}{2}$, and hence the eigenvalue under $\Delta_{-\ell-(1/2)}$ is

$$\left(\mathfrak{s} - \frac{k}{2}\right)\left(1 - \mathfrak{s} - \frac{k}{2}\right) = (1 - k + j)(-j) = j(j - \ell - \frac{3}{2}). \quad \square$$

Eichler–Selberg relations for Cohen–Eisenstein series. Eichler–Selberg relations for Cohen–Eisenstein series could be obtained as follows. On one hand, the input function $f(\tau) = f_{-2\ell, N}(\tau) \mathcal{H}_\ell(\tau)$ is weakly holomorphic, thus we do not need to deal with the additional term

$$\int_{\mathcal{F}}^{\text{reg}} \langle L_{k-2j}(f_{P \oplus N})(\tau), [\mathcal{G}_P^+(\tau), \Theta_{N^-}(\tau)]_j \rangle v^{-2} d\tau$$

arising from Theorem 3.3. Further, the function Λ_L from (3-1) simplifies to

$$\begin{aligned} \Lambda_L(\psi, p, j) &= \frac{4^{3d^-} \pi^{\frac{1}{2}-d^+} \Gamma(j+1+d^-) \Gamma(\frac{3}{2}-d^++j)}{\Gamma(\ell+\frac{1}{2}) \Gamma(\frac{3}{2}-d^+)} \\ &\quad \times \sum_{D \geq 1} \sum_{Q \in \mathcal{Q}_D} \frac{p(\psi(Q)) D^{\ell+\frac{3}{2}} y^{2+2j+2d^-}}{|Q(z, 1)|^{2+2j+2d^-}} \\ &\quad \times {}_2F_1\left(\frac{\ell+2+d^++d^-}{2}, \frac{\ell+2+d^++3d^-}{2}, \frac{5}{2}+\ell, \frac{Dy^2}{|Q(z, 1)|^2}\right) q^D, \end{aligned}$$

where \mathcal{Q}_D denotes the set of integral binary quadratic forms of discriminant D . After evaluating the hypergeometric function as in the previous proof, one may follow our proof of Theorem 3.4, namely subtract the two evaluations of $\Psi_j^{\text{reg}}(f, z)$ from each other and apply Serre duality to the resulting expression. Computing the principal part of \mathcal{G}_p^+ in addition, this yields the desired result. However, we do not pursue this here explicitly as the resulting expression is rather lengthy.

Acknowledgments

The authors thank Markus Schwagenscheidt for fruitful discussions on the topics of the paper, and for many valuable comments on a previous version. The authors would also like to thank Ken Ono for suggesting the example of Cohen–Eisenstein series, and Kathrin Bringmann as well as the referee for helpful comments on a previous version of the paper.

References

- [1] C. Alfes-Neumann, K. Bringmann, J. Males, and M. Schwagenscheidt, “Cycle integrals of meromorphic modular forms and coefficients of harmonic Maass forms”, *J. Math. Anal. Appl.* **497**:2 (2021), art. id. 124898. MR Zbl
- [2] R. E. Borcherds, “Automorphic forms with singularities on Grassmannians”, *Invent. Math.* **132**:3 (1998), 491–562. MR Zbl
- [3] R. E. Borcherds, “The Gross–Kohnen–Zagier theorem in higher dimensions”, *Duke Math. J.* **97**:2 (1999), 219–233. MR Zbl
- [4] K. Bringmann, B. Kane, and W. Kohnen, “Locally harmonic Maass forms and the kernel of the Shintani lift”, *Int. Math. Res. Not.* **2015**:11 (2015), 3185–3224. MR Zbl
- [5] K. Bringmann, B. Kane, and M. Viazovska, “Theta lifts and local Maass forms”, *Math. Res. Lett.* **20**:2 (2013), 213–234. MR Zbl
- [6] K. Bringmann, A. Folsom, K. Ono, and L. Rolin, *Harmonic Maass forms and mock modular forms: theory and applications*, Amer. Math. Soc. Colloq. Publ. **64**, Amer. Math. Soc., Providence, RI, 2017. MR Zbl
- [7] J. H. Bruinier, *Borcherds products on $O(2, l)$ and Chern classes of Heegner divisors*, Lecture Notes in Math. **1780**, Springer, 2002. MR Zbl

- [8] J. H. Bruinier, S. Ehlen, and T. Yang, “CM values of higher automorphic Green functions for orthogonal groups”, *Invent. Math.* **225**:3 (2021), 693–785. MR Zbl
- [9] J. H. Bruinier and J. Funke, “On two geometric theta lifts”, *Duke Math. J.* **125**:1 (2004), 45–90. MR Zbl
- [10] J. H. Bruinier and K. Ono, “Heegner divisors, L -functions and harmonic weak Maass forms”, *Ann. of Math. (2)* **172**:3 (2010), 2135–2181. MR Zbl
- [11] J. H. Bruinier and K. Ono, “Algebraic formulas for the coefficients of half-integral weight harmonic weak Maass forms”, *Adv. Math.* **246** (2013), 198–219. MR Zbl
- [12] J. H. Bruinier and M. Schwagenscheidt, “Theta lifts for Lorentzian lattices and coefficients of mock theta functions”, *Math. Z.* **297**:3-4 (2021), 1633–1657. MR Zbl
- [13] J. H. Bruinier and T. Yang, “Faltings heights of CM cycles and derivatives of L -functions”, *Invent. Math.* **177**:3 (2009), 631–681. MR Zbl
- [14] J. H. Bruinier and S. Zemel, “Special cycles on toroidal compactifications of orthogonal Shimura varieties”, *Math. Ann.* **384**:1-2 (2022), 309–371. MR Zbl
- [15] H. Cohen, “Sums involving the values at negative integers of L -functions of quadratic characters”, *Math. Ann.* **217**:3 (1975), 271–285. MR Zbl
- [16] J. K. Crawford, *A singular theta lift and the Shimura correspondence*, Ph.D. thesis, Durham University, 2015, available at <http://etheses.dur.ac.uk/11269/>.
- [17] J. Crawford and J. Funke, “The Shimura–Shintani correspondence via singular theta lifts and currents”, preprint, 2021. arXiv 2112.11379
- [18] W. Duke and P. Jenkins, “On the zeros and coefficients of certain weakly holomorphic modular forms”, *Pure Appl. Math. Q.* **4**:4 (2008), 1327–1340. MR Zbl
- [19] W. Duke, Ö. Imamoğlu, and Á. Tóth, “Rational period functions and cycle integrals”, *Abh. Math. Semin. Univ. Hambg.* **80**:2 (2010), 255–264. MR Zbl
- [20] S. Ehlen, P. Guerzhoy, B. Kane, and L. Rolén, “Central L -values of elliptic curves and local polynomials”, *Proc. Lond. Math. Soc. (3)* **120**:5 (2020), 742–769. MR Zbl
- [21] S. Ehlen and S. Sankaran, “On two arithmetic theta lifts”, *Compos. Math.* **154**:10 (2018), 2090–2149. MR Zbl
- [22] M. Eichler and D. Zagier, *The theory of Jacobi forms*, Progr. Math. **55**, Birkhäuser, Boston, 1985. MR Zbl
- [23] M. Hövel, *Automorphe Formen mit Singularitäten auf dem hyperbolischen Raum*, Ph.D. thesis, Technische Universität Darmstadt, 2012. Zbl
- [24] A. Hurwitz, “Ueber Relationen zwischen Klassenzahlen binärer quadratischer Formen von negativer Determinante”, *Leipzig Ber.* **36** (1884), 193–197. Zbl
- [25] A. Hurwitz, “Ueber Relationen zwischen Klassenzahlen binärer quadratischer Formen von negativer Determinante”, *Math. Ann.* **25**:2 (1885), 157–196. MR
- [26] Ö. Imamoğlu, M. Raum, and O. K. Richter, “Holomorphic projections and Ramanujan’s mock theta functions”, *Proc. Natl. Acad. Sci. USA* **111**:11 (2014), 3961–3967. MR Zbl
- [27] M. Knopp and G. Mason, “Vector-valued modular forms and Poincaré series”, *Illinois J. Math.* **48**:4 (2004), 1345–1366. MR Zbl
- [28] L. Kronecker, “Ueber die Anzahl der verschiedenen Classen quadratischer Formen von negativer Determinante”, *J. Reine Angew. Math.* **57** (1860), 248–255. MR Zbl
- [29] Y. Li and M. Schwagenscheidt, “Mock modular forms with integral Fourier coefficients”, *Adv. Math.* **399** (2022), art. id. 108264. MR Zbl

- [30] S. Löbrich and M. Schwagenscheidt, “Locally harmonic Maass forms and periods of meromorphic modular forms”, *Trans. Amer. Math. Soc.* **375**:1 (2022), 501–524. MR Zbl
- [31] J. Males, “Higher Siegel theta lifts on Lorentzian lattices, harmonic Maass forms, and Eichler–Selberg type relations”, *Math. Z.* **301**:4 (2022), 3555–3569. MR Zbl
- [32] M. H. Mertens, “Mock modular forms and class number relations”, *Res. Math. Sci.* **1** (2014), art. id. 6. MR Zbl
- [33] M. H. Mertens, “Eichler–Selberg type identities for mixed mock modular forms”, *Adv. Math.* **301** (2016), 359–382. MR Zbl
- [34] S. Nakajima, “On invariant differential operators on bounded symmetric domains of type IV”, *Proc. Japan Acad. Ser. A Math. Sci.* **58**:6 (1982), 235–238. MR Zbl
- [35] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark (editors), *NIST handbook of mathematical functions*, Cambridge Univ. Press, 2010. MR Zbl
- [36] M. Schwagenscheidt, *Regularized theta lifts of harmonic Maass forms*, Ph.D. thesis, Technische Universität Darmstadt, 2018, available at <https://tuprints.ulb.tu-darmstadt.de/7192/>. Zbl
- [37] I. Wagner, “Harmonic Maass form eigencurves”, *Res. Math. Sci.* **5**:2 (2018), art. id. 24. MR Zbl

Received October 19, 2021. Revised January 9, 2023.

JOSHUA MALES
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MANITOBA
WINNIPEG
CANADA
joshua.males@umanitoba.ca

ANDREAS MONO
DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE
DIVISION OF MATHEMATICS
UNIVERSITY OF COLOGNE
COLOGNE
GERMANY
amono@math.uni-koeln.de

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2023 is US \$605/year for the electronic version, and \$820/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 322 No. 2 February 2023

Elements of higher homotopy groups undetectable by polyhedral approximation	221
JOHN K. ACETI and JEREMY BRAZAS	
Regularity for free multiplicative convolution on the unit circle	243
SERBAN T. BELINSCHI, HARI BERCOVICI and CHING-WEI HO	
Invariant theory for the free left-regular band and a q -analogue	251
SARAH BRAUNER, PATRICIA COMMINS and VICTOR REINER	
Irredundant bases for finite groups of Lie type	281
NICK GILL and MARTIN W. LIEBECK	
Local exterior square and Asai L -functions for $GL(n)$ in odd characteristic	301
YEONGSEONG JO	
On weak convergence of quasi-infinitely divisible laws	341
ALEXEY KHARTOV	
C^* -irreducibility of commensurated subgroups	369
KANG LI and EDUARDO SCARPARO	
Local Maass forms and Eichler–Selberg relations for negative-weight vector-valued mock modular forms	381
JOSHUA MALES and ANDREAS MONO	
Representations of orientifold Khovanov–Lauda–Rouquier algebras and the Enomoto–Kashiwara algebra	407
TOMASZ PRZEŹDZIECKI	