

*Pacific  
Journal of  
Mathematics*

PEGO THEOREM ON COMPACT GROUPS

MANOJ KUMAR

Volume 328 No. 1

January 2024



## PEGO THEOREM ON COMPACT GROUPS

MANOJ KUMAR

**The Pego theorem characterizes the precompact subsets of the square-integrable functions on  $\mathbb{R}^n$  via the Fourier transform. We prove the analogue of the Pego theorem on (not necessarily abelian) compact groups.**

### 1. Introduction

Characterizing precompact subsets is one of the classical topics in function space theory. It is well known that the Arzelà–Ascoli theorem characterizes a precompact subset of the space of continuous functions over a compact Hausdorff space. The celebrated Riesz–Kolmogorov theorem provides a characterization of precompact subsets of  $L^p(\mathbb{R}^n)$ . We refer to [8] for a historical account of it. Weil [14, page 52] extended it to the Lebesgue spaces over locally compact groups. See [7] for its extension to the Banach function spaces over locally compact groups.

In 1985, Pego [13] used the Riesz–Kolmogorov theorem to find a characterization of precompact subsets of  $L^2(\mathbb{R}^n)$  via certain decay of the Fourier transform.

**Theorem 1.1.** [13, Theorems 2 and 3] *Let  $K$  be a bounded subset of  $L^2(\mathbb{R}^n)$ . Then, the following are equivalent:*

- (i)  $K$  is precompact.
- (ii)  $\int_{|x|>r} |f(x)|^2 dx \rightarrow 0$  and  $\int_{|\xi|>r} |\hat{f}(\xi)|^2 d\xi \rightarrow 0$  as  $r \rightarrow \infty$ , both uniformly for  $f$  in  $K$ .
- (iii)  $\int_{\mathbb{R}^n} |f(x+y) - f(x)|^2 dx \rightarrow 0$  as  $y \rightarrow 0$ , and  $\int_{\mathbb{R}^n} |\hat{f}(\xi + \omega) - \hat{f}(\xi)|^2 d\xi \rightarrow 0$  as  $\omega \rightarrow 0$ , both uniformly for  $f$  in  $K$ .

An application of this theorem to information theory has been provided in [13].

Pego-type theorems have also been studied via the short-time Fourier and wavelet transforms [2], the Laplace transform [11] and the Laguerre and Hankel transforms [10]. The Pego theorem has been extended to the locally compact abelian groups with some technical assumptions [5]. Using the Pontryagin duality and the Arzelà–Ascoli theorem, the authors in [6] showed that the technical assumptions

*MSC2020:* primary 43A30, 43A77; secondary 22C05.

*Keywords:* compact group, Fourier transform, compactness.

are redundant. For the  $L^1$ -space analogue of the Pego theorem over locally compact abelian groups, see [12].

In Section 2, we present preliminaries on compact groups. In Section 3, using Weil’s compactness theorem, we extend Theorem 1.1 to (not necessarily abelian) compact groups; see Theorem 3.4.

### 2. Fourier analysis on compact groups

Let  $G$  be a compact Hausdorff group. Let  $m_G$  denote the normalized positive Haar measure on  $G$ . Let  $L^p(G)$  denote the  $p$ -th Lebesgue space w.r.t. the measure  $m_G$ . The norm on the space  $L^p(G)$  is denoted by  $\|\cdot\|_p$ .

We denote by  $\widehat{G}$  the space consisting of all irreducible unitary representations of  $G$  up to the unitary equivalence. The set  $\widehat{G}$  is known as the unitary dual of  $G$  and is equipped with the discrete topology. Note that the representation space  $\mathcal{H}_\pi$  of  $\pi \in \widehat{G}$  is a complex Hilbert space and finite-dimensional. Denote by  $d_\pi$  the dimension of  $\mathcal{H}_\pi$ .

Let  $\Lambda \subset \widehat{G}$ . Assume that  $\{(X_\pi, \|\cdot\|_\pi) : \pi \in \Lambda\}$  is a family of Banach spaces. For  $1 \leq p < \infty$ , we denote by  $\ell^p\text{-}\bigoplus_{\pi \in \Lambda} X_\pi$  the Banach space

$$\left\{ (x_\pi) \in \prod_{\pi \in \Lambda} X_\pi : \sum_{\pi \in \Lambda} d_\pi \|x_\pi\|_\pi^p < \infty \right\}$$

endowed with the norm  $\|(x_\pi)\|_{\ell^p\text{-}\bigoplus_{\pi \in \Lambda} X_\pi} := \left(\sum_{\pi \in \Lambda} d_\pi \|x_\pi\|_\pi^p\right)^{1/p}$ . Denote by  $\ell^\infty\text{-}\bigoplus_{\pi \in \Lambda} X_\pi$  the Banach space

$$\left\{ (x_\pi) \in \prod_{\pi \in \Lambda} X_\pi : \sup_{\pi \in \Lambda} \|x_\pi\|_\pi < \infty \right\}$$

endowed with the norm  $\|(x_\pi)\|_{\ell^\infty\text{-}\bigoplus_{\pi \in \Lambda} X_\pi} := \sup_{\pi \in \Lambda} \|x_\pi\|_\pi$ . Similarly, denote by  $c_0\text{-}\bigoplus_{\pi \in \Lambda} X_\pi$  the space consisting of  $(x_\pi)$  from  $\ell^\infty\text{-}\bigoplus_{\pi \in \Lambda} X_\pi$  such that  $x_\pi \rightarrow 0$  as  $\pi \rightarrow \infty$ , i.e., for any given  $\epsilon > 0$  there exists a finite set  $\Lambda_\epsilon \subset \Lambda$  such that  $\|x_\pi\|_\pi < \epsilon$  for all  $\pi \in \Lambda \setminus \Lambda_\epsilon$ . Note that  $c_0\text{-}\bigoplus_{\pi \in \Lambda} X_\pi$  is a closed subspace of  $\ell^\infty\text{-}\bigoplus_{\pi \in \Lambda} X_\pi$ .

For  $1 \leq p < \infty$ , let  $\mathcal{B}_p(\mathcal{H}_\pi)$  denote the space of all bounded linear operators  $T$  on  $\mathcal{H}_\pi$  such that  $\|T\|_{\mathcal{B}_p(\mathcal{H}_\pi)} := (\text{tr}(|T|^p))^{1/p} < \infty$ . The space  $\mathcal{B}_2(\mathcal{H}_\pi)$  is called the space of the Hilbert–Schmidt operators on the Hilbert space  $\mathcal{H}_\pi$ . The space  $\mathcal{B}_2(\mathcal{H}_\pi)$  is a Hilbert space endowed with the inner product

$$\langle T, S \rangle_{\mathcal{B}_2(\mathcal{H}_\pi)} := \text{tr}(TS^*).$$

Let  $\mathcal{B}(\mathcal{H}_\pi)$  denote the space consisting of all bounded linear operators on  $\mathcal{H}_\pi$  endowed with the operator norm.

Let  $f \in L^1(G)$ . The Fourier transform of  $f$  is defined by

$$\hat{f}(\pi) = \int_G f(t)\pi(t)^* dm_G(t), \quad \pi \in \widehat{G}.$$

The Fourier transform operator  $f \mapsto \hat{f}$  from  $L^1(G)$  into  $\ell^\infty\text{-}\bigoplus_{\pi \in \widehat{G}} \mathcal{B}(\mathcal{H}_\pi)$  is injective and bounded. By the Riemann–Lebesgue lemma, we know that  $\hat{f} \in c_0\text{-}\bigoplus_{\pi \in \widehat{G}} \mathcal{B}(\mathcal{H}_\pi)$ . The convolution of  $f, g \in L^1(G)$  is given by

$$f * g(x) = \int_G f(xy^{-1})g(y) dm_G(y).$$

Then,  $\widehat{f * g}(\pi) = \hat{g}(\pi)\hat{f}(\pi)$ ,  $\pi \in \widehat{G}$ . For  $y \in G$ , the right translation  $R_y$  of  $f \in L^p(G)$  is given by  $R_y(f)(x) = f(xy)$ ,  $x \in G$ . Then,  $\widehat{R_y f}(\pi) = \pi(y)\hat{f}(\pi)$ ,  $\pi \in \widehat{G}$ .

For more information on compact groups, we refer to [4; 9].

Throughout the paper,  $G$  will denote a (not necessarily abelian) compact Hausdorff group. The identity of  $G$  is denoted by  $e$ . We will denote by  $I_{d_\pi}$  the  $d_\pi \times d_\pi$  identity matrix.

### 3. Pego theorem on compact groups

We discuss the characterization of precompact subsets of square-integrable functions on  $G$  in terms of the Fourier transform. We need the following definitions.

Let  $K \subset L^p(G)$ . Define  $\widehat{K} := \{\hat{f} : f \in L^p(G)\}$ .  $K$  is said to be *uniformly  $L^p(G)$ -equicontinuous* if for any given  $\epsilon > 0$  there exists an open neighborhood  $O$  of  $e$  such that

$$\|R_y f - f\|_p < \epsilon, \quad f \in K \text{ and } y \in O.$$

Let  $F \subset \ell^p\text{-}\bigoplus_{\pi \in \widehat{G}} \mathcal{B}_p(\mathcal{H}_\pi)$ .  $F$  is said to have *uniform  $\ell^p\text{-}\bigoplus_{\pi \in \widehat{G}} \mathcal{B}_p(\mathcal{H}_\pi)$ -decay* if for any given  $\epsilon > 0$  there exists a finite set  $A \subset \widehat{G}$  such that

$$\|\phi\|_{\ell^p\text{-}\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_p(\mathcal{H}_\pi)} < \epsilon, \quad \phi \in F.$$

Let us begin with some important lemmas.

**Lemma 3.1.** *Let  $K \subset L^p(G)$ , where  $p \in [1, 2]$ . If  $K$  is uniformly  $L^p(G)$ -equicontinuous then  $\widehat{K}$  has uniform  $\ell^{p'}\text{-}\bigoplus_{\pi \in \widehat{G}} \mathcal{B}_{p'}(\mathcal{H}_\pi)$ -decay.*

*Proof.* Let  $(e_U)_{U \in \Lambda}$  be a Dirac net on  $G$ ; see [1, page 24]. By the Riemann–Lebesgue lemma [9, Theorem 28.40],  $\widehat{e_U} \in c_0\text{-}\bigoplus_{\pi \in \widehat{G}} \mathcal{B}(\mathcal{H}_\pi)$ . Then, there exists a finite set  $A \subset \widehat{G}$  such that

$$\|\widehat{e_U}(\pi)\|_{\mathcal{B}(\mathcal{H}_\pi)} \leq \frac{1}{2}, \quad \pi \in \widehat{G} \setminus A.$$

Let  $f \in K$ . We denote by  $\widehat{e_U} \hat{f}$  the pointwise product of  $\widehat{e_U}$  and  $\hat{f}$ . Now,

$$\begin{aligned} & \|\hat{f}\|_{\ell^{p'}-\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_{p'}(\mathcal{H}_\pi)} \\ & \leq \|\hat{f} - \widehat{e_U} \hat{f}\|_{\ell^{p'}-\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_{p'}(\mathcal{H}_\pi)} + \|\widehat{e_U} \hat{f}\|_{\ell^{p'}-\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_{p'}(\mathcal{H}_\pi)} \\ & \leq \|\widehat{f - f * e_U}\|_{\ell^{p'}-\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_{p'}(\mathcal{H}_\pi)} + \|\hat{f}\|_{\ell^{p'}-\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_{p'}(\mathcal{H}_\pi)} \sup_{\pi \in \widehat{G} \setminus A} \|\widehat{e_U}(\pi)\|_{\mathcal{B}(\mathcal{H}_\pi)} \\ & \leq \|\widehat{f - f * e_U}\|_{\ell^{p'}-\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_{p'}(\mathcal{H}_\pi)} + \frac{1}{2} \|\hat{f}\|_{\ell^{p'}-\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_{p'}(\mathcal{H}_\pi)}. \end{aligned}$$

Then, applying the Hausdorff–Young inequality [9, Theorem 31.22], we get

$$\begin{aligned} \|\hat{f}\|_{\ell^{p'}-\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_{p'}(\mathcal{H}_\pi)} & \leq 2 \|\widehat{f - f * e_U}\|_{\ell^{p'}-\bigoplus_{\pi \in \widehat{G}} \mathcal{B}_{p'}(\mathcal{H}_\pi)} \\ & \leq 2 \|f - f * e_U\|_p \\ & = 2 \left( \int_G |f(x) - f * e_U(x)|^p dm_G(x) \right)^{1/p} \\ & = 2 \left( \int_G \left| \int_G (f(x) - f(xy^{-1})) e_U(y) dm_G(y) \right|^p dm_G(x) \right)^{1/p}. \end{aligned}$$

Therefore, using the Minkowski integral inequality, we obtain

$$\begin{aligned} \|\hat{f}\|_{\ell^{p'}-\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_{p'}(\mathcal{H}_\pi)} & \leq 2 \int_G \left( \int_G |f(x) - f(xy^{-1})|^p dm_G(x) \right)^{1/p} e_U(y) dm_G(y) \\ & \leq 2 \sup_{y \in U} \left( \int_G |f(x) - f(xy^{-1})|^p dm_G(x) \right)^{1/p}. \end{aligned}$$

Let  $\epsilon > 0$ . Since  $K$  is uniformly  $L^p(G)$ -equicontinuous, there exists an open neighborhood  $O$  of  $e$  such that

$$\|R_y f - f\|_p < \frac{\epsilon}{2}, \quad f \in K \text{ and } y \in O.$$

By [1, Lemma 1.6.5, page 24], we get that there exists  $U \in \Lambda$  such that

$$\|R_y f - f\|_p < \frac{\epsilon}{2}, \quad f \in K \text{ and } y \in U.$$

Hence,

$$\|\hat{f}\|_{\ell^{p'}-\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_{p'}(\mathcal{H}_\pi)} < \epsilon, \quad f \in K. \quad \square$$

**Lemma 3.2.** *Let  $K$  be a subset of  $L^{p'}(G)$ , where  $p \in [1, 2]$ . If  $\widehat{K}$  has uniform  $\ell^p-\bigoplus_{\pi \in \widehat{G}} \mathcal{B}_p(\mathcal{H}_\pi)$ -decay then  $K$  is uniformly  $L^{p'}(G)$ -equicontinuous.*

*Proof.* Let  $\epsilon > 0$ . Since  $\widehat{K}$  has uniform  $\ell^p-\bigoplus_{\pi \in \widehat{G}} \mathcal{B}_p(\mathcal{H}_\pi)$ -decay, there exists a finite set  $A \subset \widehat{G}$  such that

$$\|\hat{f}\|_{\ell^p-\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_p(\mathcal{H}_\pi)} < \frac{\epsilon}{4}, \quad f \in K.$$

Let  $f \in K$  and  $y \in G$ . Then, applying [9, Corollary 31.25], we obtain

$$\begin{aligned}
 \|R_y f - f\|_{p'} &\leq \|\widehat{R_y f - f}\|_{\ell^p \cdot \bigoplus_{\pi \in \widehat{G}} \mathcal{B}_p(\mathcal{H}_\pi)} \\
 &= \left( \sum_{\pi \in \widehat{G}} d_\pi \|\widehat{R_y f}(\pi) - \widehat{f}(\pi)\|_{\mathcal{B}_p(\mathcal{H}_\pi)}^p \right)^{1/p} \\
 &\leq \left( \sum_{\pi \in A} d_\pi \|\pi(y) \widehat{f}(\pi) - \widehat{f}(\pi)\|_{\mathcal{B}_p(\mathcal{H}_\pi)}^p \right)^{1/p} \\
 &\quad + \left( \sum_{\pi \in \widehat{G} \setminus A} d_\pi \|\pi(y) \widehat{f}(\pi) - \widehat{f}(\pi)\|_{\mathcal{B}_p(\mathcal{H}_\pi)}^p \right)^{1/p} \\
 &\leq \sup_{\pi \in A} \|\pi(y) - I_{d_\pi}\|_{\mathcal{B}(\mathcal{H}_\pi)} \left( \sum_{\pi \in A} d_\pi \|\widehat{f}(\pi)\|_{\mathcal{B}_p(\mathcal{H}_\pi)}^p \right)^{1/p} \\
 &\quad + \sup_{\pi \in \widehat{G} \setminus A} \|\pi(y) - I_{d_\pi}\|_{\mathcal{B}(\mathcal{H}_\pi)} \left( \sum_{\pi \in \widehat{G} \setminus A} d_\pi \|\widehat{f}(\pi)\|_{\mathcal{B}_p(\mathcal{H}_\pi)}^p \right)^{1/p} \\
 &\leq M \sup_{\pi \in A} \|\pi(y) - I_{d_\pi}\|_{\mathcal{B}(\mathcal{H}_\pi)} + \frac{\epsilon}{2},
 \end{aligned}$$

where  $M$  is a positive number such that  $(\sum_{\pi \in A} d_\pi \|\widehat{f}(\pi)\|_{\mathcal{B}_p(\mathcal{H}_\pi)}^p)^{1/p} \leq M$ .

Let  $\pi \in A$ . Using continuity of  $\pi$ , we obtain that there exists a neighborhood  $O_\pi$  of  $e$  such that

$$\|\pi(y) - I_{d_\pi}\|_{\mathcal{B}(\mathcal{H}_\pi)} < \frac{\epsilon}{2M}, \quad y \in O_\pi.$$

Assume that  $O = \bigcap_{\pi \in A} O_\pi$ . Then,

$$\|\pi(y) - I_{d_\pi}\|_{\mathcal{B}(\mathcal{H}_\pi)} < \frac{\epsilon}{2M}, \quad \pi \in A \text{ and } y \in O.$$

Hence,

$$\|R_y f - f\|_{p'} < \epsilon, \quad f \in K \text{ and } y \in O. \quad \square$$

The following corollary is a generalization of [13, Theorem 1] studied on  $\mathbb{R}^n$ , and [5, Theorem 1] and [3, Lemma 2.5] studied on locally compact abelian groups. This is also an improvement of the corresponding result on compact abelian groups in the sense that we do not assume boundedness of the subset of  $L^2(G)$ .

**Corollary 3.3.** *Let  $K \subset L^2(G)$ . Then,  $K$  is uniformly  $L^2(G)$ -equicontinuous if and only if  $\widehat{K}$  has uniform  $\ell^2 \cdot \bigoplus_{\pi \in \widehat{G}} \mathcal{B}_2(\mathcal{H}_\pi)$ -decay.*

*Proof.* This is a direct consequence of Lemmas 3.1 and 3.2.  $\square$

Now, we present our main result, that is, the Pego theorem over compact groups. It is a consequence of the Weil theorem and above corollary.

**Theorem 3.4.** *Let  $K$  be a bounded subset of  $L^2(G)$ . Then, the following are equivalent:*

- (i)  $K$  is precompact.

- (ii)  $K$  is uniformly  $L^2(G)$ -equicontinuous.  
 (iii)  $\widehat{K}$  has uniform  $\ell^2\text{-}\bigoplus_{\pi \in \widehat{G}} \mathcal{B}_2(\mathcal{H}_\pi)$ -decay.

*Proof.* For any given  $\epsilon > 0$  we have that

$$\sup_{f \in K} \|f \chi_{G \setminus \epsilon}\|_2 = 0 < \epsilon.$$

Therefore, (i) and (ii) are equivalent by the Weil theorem [14, page 52] (or see [7, Theorems 3.1 and 3.3]). Further, (ii) and (iii) are equivalent by Corollary 3.3.  $\square$

The following gives an example of a set  $K \subset L^2(G)$  which is not precompact but  $K$  is uniformly  $L^2(G)$ -equicontinuous and  $\widehat{K}$  has uniform  $\ell^2\text{-}\bigoplus_{\pi \in \widehat{G}} \mathcal{B}_2(\mathcal{H}_\pi)$ -decay.

**Example 3.5.** Consider the set  $K = \{n \chi_G : n \in \mathbb{N}\} \subset L^2(G)$  as given in [7, Example 4.2]. Since  $K$  consists of only constant functions, it is clear that  $K$  is uniformly  $L^2(G)$ -equicontinuous. By Corollary 3.3,  $\widehat{K}$  has uniform  $\ell^2\text{-}\bigoplus_{\pi \in \widehat{G}} \mathcal{B}_2(\mathcal{H}_\pi)$ -decay. Since  $K$  is not bounded,  $K$  is not precompact.

Now, with the help of our main result Theorem 3.4, we show that certain subsets of  $L^2(G)$  are precompact.

**Example 3.6.** (i) Let  $r \in \mathbb{R}$ . Consider the set  $K = \{\frac{r}{n} \chi_G : n \in \mathbb{N}\} \subset L^2(G)$ . Since  $\{\frac{r}{n} : n \in \mathbb{N}\}$  is bounded and  $K$  consists of only constant functions, it follows that  $K$  is bounded and uniformly  $L^2(G)$ -equicontinuous. Therefore, by Theorem 3.4,  $K$  is precompact.

(ii) Let  $A$  be a finite subset of  $\widehat{G}$ . Assume that  $K$  is a bounded subset of the linear span of the set consisting of matrix entries [4, page 139] of elements in  $A$ . Note that the matrix entries are bounded functions. For  $f \in K$ , using the Schur orthogonality relations [4, Theorem 5.8] we obtain that

$$\|\widehat{f}\|_{\ell^2\text{-}\bigoplus_{\pi \in \widehat{G} \setminus A} \mathcal{B}_2(\mathcal{H}_\pi)} = 0.$$

Thus,  $\widehat{K}$  has uniform  $\ell^2\text{-}\bigoplus_{\pi \in \widehat{G}} \mathcal{B}_2(\mathcal{H}_\pi)$ -decay. Hence, by Theorem 3.4,  $K$  is precompact. In particular, the convex hull of the set consisting of matrix entries of elements in  $A$  is precompact.

### Acknowledgements

The author is supported by the NBHM postdoctoral fellowship with reference number 0204/3/2021/R&D-II/7356 from the Department of Atomic Energy (DAE), Government of India. The author is grateful to Sundaram Thangavelu for his useful comments.



## References

- [1] A. Deitmar and S. Echterhoff, *Principles of harmonic analysis*, 2nd ed., Springer, 2014. MR Zbl
- [2] M. Dörfler, H. G. Feichtinger, and K. Gröchenig, “Compactness criteria in function spaces”, *Colloq. Math.* **94**:1 (2002), 37–50. MR Zbl
- [3] H. G. Feichtinger, “Compactness in translation invariant Banach spaces of distributions and compact multipliers”, *J. Math. Anal. Appl.* **102**:2 (1984), 289–327. MR Zbl
- [4] G. B. Folland, *A course in abstract harmonic analysis*, 2nd ed., CRC Press, Boca Raton, FL, 2016. MR
- [5] P. Górká, “Pego theorem on locally compact abelian groups”, *J. Algebra Appl.* **13**:4 (2014), art. id. 1350143. MR Zbl
- [6] P. Górká and T. Kostrzewa, “Pego everywhere”, *J. Algebra Appl.* **15**:4 (2016), art. id. 1650074. MR Zbl
- [7] P. Górká and P. Pośpiech, “Banach function spaces on locally compact groups”, *Ann. Funct. Anal.* **10**:4 (2019), 460–471. MR Zbl
- [8] H. Hanche-Olsen and H. Holden, “The Kolmogorov–Riesz compactness theorem”, *Expo. Math.* **28**:4 (2010), 385–394. MR Zbl
- [9] E. Hewitt and K. A. Ross, *Abstract harmonic analysis, II: Structure and analysis for compact groups: analysis on locally compact Abelian groups*, Grundlehren der Math. Wissenschaften **152**, Springer, 1970. MR Zbl
- [10] A. P. Horváth, “Compactness criteria via Laguerre and Hankel transformations”, *J. Math. Anal. Appl.* **507**:2 (2022), art. id. 125852. MR Zbl
- [11] M. Krukowski, “Characterizing compact families via the Laplace transform”, *Ann. Acad. Sci. Fenn. Math.* **45**:2 (2020), 991–1002. MR Zbl
- [12] M. Krukowski, “How Arzelà and Ascoli would have proved Pego theorem for  $L^1(G)$  (if they lived in the 21<sup>st</sup> century)?”, preprint, 2020. arXiv 2006.12130
- [13] R. L. Pego, “Compactness in  $L^2$  and the Fourier transform”, *Proc. Amer. Math. Soc.* **95**:2 (1985), 252–254. MR Zbl
- [14] A. Weil, *L’intégration dans les groupes topologiques et ses applications*, Actualités Scientifiques et Industrielles **869**, Hermann, Paris, 1940. MR Zbl

Received February 21, 2023. Revised December 14, 2023.

MANOJ KUMAR  
DEPARTMENT OF MATHEMATICS  
INDIAN INSTITUTE OF SCIENCE BANGALORE  
BANGALORE  
INDIA  
manojk9t3@gmail.com



# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Paul Balmer  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[balmer@math.ucla.edu](mailto:balmer@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Matthias Aschenbrenner  
Fakultät für Mathematik  
Universität Wien  
Vienna, Austria  
[matthias.aschenbrenner@univie.ac.at](mailto:matthias.aschenbrenner@univie.ac.at)

Atsushi Ichino  
Department of Mathematics  
Kyoto University  
Kyoto 606-8502, Japan  
[atsushi.ichino@gmail.com](mailto:atsushi.ichino@gmail.com)

Dimitri Shlyakhtenko  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[shlyakht@ipam.ucla.edu](mailto:shlyakht@ipam.ucla.edu)

Robert Lipshitz  
Department of Mathematics  
University of Oregon  
Eugene, OR 97403  
[lipshitz@uoregon.edu](mailto:lipshitz@uoregon.edu)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Ruixiang Zhang  
Department of Mathematics  
University of California  
Berkeley, CA 94720-3840  
[ruixiang@berkeley.edu](mailto:ruixiang@berkeley.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2024 is US \$645/year for the electronic version, and \$875/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 328    No. 1    January 2024

---

|   |     |
|---|-----|
| Categorification of the internal braid group action for quantum groups,<br>I: 2-functoriality | 1   |
| MICHAEL T. ABRAM, LAFFITE LAMBERTO-EGAN, AARON D.<br>LAUDA and DAVID E. V. ROSE               |     |
| Fused Hecke algebra and one-boundary algebras   | 77  |
| LOÏC POULAIN D'ANDECY and MERI ZAIMI  |     |
| On the generic behavior of the spectral norm  | 119 |
| ERMAN ÇINELI, VIKTOR L. GINZBURG and BAŞAK Z. GÜREL   |     |
| Pego theorem on compact groups  | 137 |
| MANOJ KUMAR   |     |
| Maximal degree of a map of surfaces   | 145 |
| ANDREY RYABICHEV  |     |
| A Viro–Zvonilov-type inequality for Q-flexible curves of odd degree                           | 157 |
| ANTHONY SAINT-CRIQ  |     |