

*Pacific
Journal of
Mathematics*

**ON THE COEFFICIENT INEQUALITIES
FOR SOME CLASSES OF HOLOMORPHIC MAPPINGS
IN COMPLEX BANACH SPACES**

QINGHUA XU, XIAOHUA YANG AND TAISHUN LIU

ON THE COEFFICIENT INEQUALITIES FOR SOME CLASSES OF HOLOMORPHIC MAPPINGS IN COMPLEX BANACH SPACES

QINGHUA XU, XIAOHUA YANG AND TAISHUN LIU

Let \mathcal{C} be the familiar class of normalized close-to-convex functions in the unit disk. Koepf (1987) proved that for a function $f(z) = z + \sum_{k=2}^{\infty} a_k z^k$ in the class \mathcal{C} ,

$$|a_3 - \lambda a_2^2| \leq \begin{cases} 3 - 4\lambda, & \lambda \in [0, \frac{1}{3}], \\ \frac{1}{3} + \frac{4}{9\lambda}, & \lambda \in [\frac{1}{3}, \frac{2}{3}], \\ 1, & \lambda \in [\frac{2}{3}, 1] \end{cases} \quad \text{and} \quad ||a_3| - |a_2|| \leq 1.$$

Recently, Xu et al. (2023) generalized the above results to a subclass of close-to-quasiconvex mappings of type B defined on the open unit polydisc in \mathbb{C}^n , and to a subclass of close-to-starlike mappings defined on the open unit ball of a complex Banach space, respectively. In the first part of this paper, by using different methods, we obtain the corresponding results of norm type and functional type on the open unit ball in a complex Banach space. We next give the coefficient inequalities for a subclass of g -starlike mappings of complex order λ on the open unit ball of a complex Banach space, which generalize many known results. Moreover, the proofs presented here are simpler than those given in the related papers.

1. Introduction

Let \mathcal{S} be the class of functions of the form

$$(1-1) \quad f(\xi) = \xi + \sum_{m=2}^{\infty} a_m \xi^m,$$

which are univalent in the open unit disk

$$\mathbb{U} = \{\xi \in \mathbb{C} : |\xi| < 1\}.$$

This work was supported by National Natural Science Foundation of China (grant 11971165) and Natural Science Foundation of Zhejiang Province (grant LY21A010003).

MSC2020: primary 32H02; secondary 30C55.

Keywords: Fekete and Szegő problem, close-to-quasiconvex mapping of type B , close-to-starlike mapping, g -starlike mapping of complex order λ .

Let X be a complex Banach space with norm $\|\cdot\|$, \mathbb{B} be the open unit ball of X . Let $L(X, Y)$ denote the set of continuous linear operators from X into a complex Banach space Y . Let I be the identity in $L(X, X)$. For each $x \in X \setminus \{0\}$, we define

$$T(x) = \{T_x \in L(X, \mathbb{C}) : \|T_x\| = 1, T_x(x) = \|x\|\}.$$

According to the Hahn–Banach theorem, $T(x)$ is nonempty.

Let $H(\mathbb{B})$ denote the set of all holomorphic mappings from \mathbb{B} into X . It is well known that if $f \in H(\mathbb{B})$, then

$$f(y) = \sum_{m=0}^{\infty} \frac{1}{m!} D^m f(x)((y-x)^m)$$

for all y in some neighborhood of $x \in \mathbb{B}$, where $D^m f(x)$ is the m -th Fréchet derivative of f at x , and for $m \geq 1$,

$$D^m f(x)((y-x)^m) = D^m f(x) \underbrace{(y-x, \dots, y-x)}_m.$$

Furthermore, $D^m f(x)$ is a bounded symmetric m -linear mapping from

$$X^m = \underbrace{X \times \dots \times X}_m \text{ into } X.$$

A holomorphic mapping $f : \mathbb{B} \rightarrow X$ is said to be biholomorphic if the inverse f^{-1} exists and is holomorphic on the open set $f(\mathbb{B})$. A mapping $f \in H(\mathbb{B})$ is called locally biholomorphic if the Fréchet derivative $Df(x)$ has a bounded inverse for each $x \in \mathbb{B}$. If $f : \mathbb{B} \rightarrow X$ is a holomorphic mapping, then f is called normalized if $f(0) = 0$ and $Df(0) = I$, where I represents the identity operator from X into X . A mapping $f \in H(\mathbb{B})$ is called starlike if f is biholomorphic on \mathbb{B} and $f(\mathbb{B})$ is a starlike domain. Let $\mathcal{S}^*(\mathbb{B})$ denote the class of normalized starlike mappings on \mathbb{B} , when $X = \mathbb{C}$, $\mathbb{B} = \mathbb{U}$, the class $\mathcal{S}^*(\mathbb{U})$ is denoted by \mathcal{S}^* . Suppose $f, g \in H(\mathbb{U})$. If there exists a Schwarz function φ (i.e., $\varphi \in H(\mathbb{U})$, $\varphi(0) = 0$, $\varphi(\mathbb{U}) \subseteq \mathbb{U}$) such that $f = g \circ \varphi$, then we say that f is subordinate to g (written $f < g$).

Now, we introduce the class of quasiconvex mappings of type B on \mathbb{B} in X , which has been introduced by Roper and Suffridge [31] on the unit ball $\mathbb{B} \subset \mathbb{C}^n$.

Definition 1.1. Let $h : \mathbb{B} \rightarrow X$ be a normalized locally biholomorphic mapping. If

$$(1-2) \Re\{T_x[(Dh(x))^{-1}(D^2h(x)(x^2)+Dh(x)x)]\} > 0, \quad x \in \mathbb{B} \setminus \{0\}, T_x \in T(x),$$

then h is called a quasiconvex mapping of type B on \mathbb{B} .

Let $\mathcal{Q}_B(\mathbb{B})$ denote the class of quasiconvex mappings of type B on \mathbb{B} . When $X = \mathbb{C}$, $\mathbb{B} = \mathbb{U}$, we deduce easily that relation (1-2) is equivalent to

$$\Re\left(1 + \frac{\xi h''(\xi)}{h'(\xi)}\right) > 0, \quad \xi \in \mathbb{U},$$

which is the well-known criterion of convex functions on \mathbb{U} . Let \mathcal{K} denote the class of normalized convex functions on \mathbb{U} .

Xu et al. [35] introduced the following class of mappings on the open unit ball of a complex Banach space.

Definition 1.2 [35]. Suppose that $f : \mathbb{B} \rightarrow X$ is a normalized holomorphic mapping. If there exists a mapping $h \in \mathcal{Q}_B(\mathbb{B})$ such that

$$(1-3) \quad \Re\{T_x[(Dh(x))^{-1}Df(x)x]\} > 0, \quad x \in \mathbb{B} \setminus \{0\}, T_x \in T(x),$$

then f is called a close-to-quasiconvex mapping of type B on \mathbb{B} .

If $X = \mathbb{C}^n$, $\mathbb{B} = \mathbb{U}^n$, then it is obvious that the relation (1-3) is equivalent to

$$\Re \frac{p_j(z)}{z_j} > 0, \quad z \in \mathbb{U}^n \setminus \{0\},$$

where $p(z) = (p_1(z), \dots, p_n(z))' = (Dh(z))^{-1}Df(z)z$ is a column vector in \mathbb{C}^n , and j satisfies $|z_j| = \|z\| = \max_{1 \leq k \leq n} \{|z_k|\}$.

The following definition has been introduced by Pfaltzgraff and Suffridge [29] on the unit ball with respect to an arbitrary norm in \mathbb{C}^n .

Definition 1.3. Suppose that $f : \mathbb{B} \rightarrow X$ is a normalized locally biholomorphic mapping. If there exists a mapping $h \in \mathcal{S}^*(\mathbb{B})$ such that

$$(1-4) \quad \Re\{T_x[(Df(x))^{-1}h(x)]\} > 0, \quad x \in \mathbb{B} \setminus \{0\}, T_x \in T(x),$$

then f is called a close-to-starlike mapping on \mathbb{B} .

Remark 1.4. Clearly, if $X = \mathbb{C}$, $\mathbb{B} = \mathbb{U}$, then the relation (1-3) (respectively, the relation (1-4)) is equivalent to $\Re \frac{f'(\xi)}{h'(\xi)} > 0$, $\xi \in \mathbb{U}$, here $h \in \mathcal{K}$ (respectively, $\Re \frac{\xi f'(\xi)}{h(\xi)} > 0$, $\xi \in \mathbb{U}$, here $h \in \mathcal{S}^*$), which is the usual definition of close-to-convex functions on \mathbb{U} .

Koepf [23] obtained the following Fekete and Szegő inequality for the class \mathcal{C} .

Theorem 1.5 [23]. Let the function $f(\xi)$ be defined by (1-1). If $f \in \mathcal{C}$, then

$$|a_3 - \lambda a_2^2| \leq \begin{cases} 3 - 4\lambda, & \lambda \in [0, \frac{1}{3}], \\ \frac{1}{3} + \frac{4}{9\lambda}, & \lambda \in [\frac{1}{3}, \frac{2}{3}], \\ 1, & \lambda \in [\frac{2}{3}, 1]. \end{cases}$$

As an interesting application of Theorem 1.5, it was proved that $||a_3| - |a_2|| \leq 1$ for the class \mathcal{C} .

In recent years, the Fekete and Szegő inequality for subclass of biholomorphic mappings in several complex variables has been studied by some authors (see [3; 4; 5; 6; 7; 15; 20; 21; 32; 36; 38]).

Xu et al. [39] obtained the following Fekete and Szegő inequality for the subclass of close-to-convex mappings of type B on the open unit polydisk \mathbb{U}^n in \mathbb{C}^n with respect to $H \in \mathcal{Q}_B(\mathbb{U}^n)$, which could be regarded as a generalization of Theorem 1.5 to several complex variables.

Theorem 1.6 [39]. *Let $f : \mathbb{U}^n \rightarrow \mathbb{C}$, $h : \mathbb{U}^n \rightarrow \mathbb{C}$ be holomorphic functions, and $H(z) = zh(z) \in \mathcal{Q}_B(\mathbb{U}^n)$. Suppose that $F(z) = zf(z)$ is a close-to-convex mapping of type B with respect to $H(z)$. Then, for $\lambda \in [0, 1]$, $z \in \mathbb{U}^n$, we have*

$$\left\| \frac{1}{3!} D^3 F(0)(z^3) - \lambda \frac{1}{2!} D^2 F(0) \left(z, \frac{D^2 F(0)(z^2)}{2!} \right) \right\| \leq \begin{cases} (3 - 4\lambda) \|z\|^3, & \lambda \in [0, \frac{1}{3}], \\ (\frac{1}{3} + \frac{4}{9\lambda}) \|z\|^3, & \lambda \in [\frac{1}{3}, \frac{2}{3}], \\ \|z\|^3, & \lambda \in [\frac{2}{3}, 1] \end{cases}$$

and

$$\left\| \frac{D^3 F(0)(z^3)}{3!} \right\| - \left\| \frac{D^2 F(0)(z^2) \|z\|}{2!} \right\| \leq \|z\|^3.$$

The above estimates are sharp.

Hamada [15] generalized Theorem 1.6 to the open unit ball of a complex Banach space under weaker assumptions than in Theorem 1.6. Moreover, in the same paper, Hamada also obtained the Fekete and Szegő inequality of functional type for the subclasses of close-to-convex mappings of type B on the open unit ball \mathbb{B} in a complex Banach space.

Theorem 1.7 [15]. *Let G be a convex mapping of type B on \mathbb{B} such that*

$$\frac{1}{2!} D^2 G(0)(x^2) = L_G(x) x, \quad x \in X,$$

where $L_G(\cdot) \in L(X, \mathbb{C})$. Let F be a close-to-convex mapping of type B on \mathbb{B} with respect to G such that

$$\frac{1}{2!} D^2 F(0)(x^2) = L_F(x) x, \quad x \in X,$$

where $L_F(\cdot) \in L(X, \mathbb{C})$ and

$$\frac{1}{3!} D^3 F(0)(x^3) = Q_F(x) x, \quad x \in X,$$

where $Q_F(x)$ is a homogeneous polynomial of degree 2 with values in \mathbb{C} . Let $x_0 \in X$ with $\|x_0\| = 1$. Then, for $\lambda \in [0, 1]$, it holds that

$$\left\| \frac{1}{3!} D^3 F(0)(x_0^3) - \lambda \frac{1}{2!} D^2 F(0) \left(x_0, \frac{D^2 F(0)(x_0^2)}{2!} \right) \right\| \leq \begin{cases} 3 - 4\lambda, & \lambda \in [0, \frac{1}{3}], \\ \frac{1}{3} + \frac{4}{9\lambda}, & \lambda \in [\frac{1}{3}, \frac{2}{3}], \\ 1, & \lambda \in [\frac{2}{3}, 1]. \end{cases}$$

The above estimates are sharp.

More recently, Xu et al. [37] gave another extension of Theorem 1.5 to higher dimensions, and established the following Fekete and Szegő inequality for the subclass of close-to-starlike mappings on the open unit ball \mathbb{B} in a complex Banach space with respect to $H \in \mathcal{S}^*(\mathbb{B})$.

Theorem 1.8 [37]. Let $f : \mathbb{B} \rightarrow \mathbb{C}$, $h : \mathbb{B} \rightarrow \mathbb{C}$ be holomorphic functions, and $H(x) = xh(x) \in \mathcal{S}^*(\mathbb{B})$. Suppose that $F(x) = xf(x)$ is a close-to-starlike mapping with respect to $H(x)$. Then, for $x \in \mathbb{B} \setminus \{0\}$, $T_x \in T(x)$, $\lambda \in [0, 1]$, we have

$$\left| \frac{T_x(D^3 F(0)(x^3))}{3! \|x\|^3} - \lambda \left(\frac{T_x(D^2 F(0)(x^2))}{2! \|x\|^2} \right)^2 \right| \leq \begin{cases} 3 - 4\lambda, & \lambda \in [0, \frac{1}{3}], \\ \frac{1}{3} + \frac{4}{9\lambda}, & \lambda \in [\frac{1}{3}, \frac{2}{3}], \\ 1, & \lambda \in [\frac{2}{3}, 1] \end{cases}$$

and

$$\left| \left| \frac{T_x(D^3 F(0)(x^3))}{3! \|x\|^3} \right| - \left| \frac{T_x(D^2 F(0)(x^2))}{2! \|x\|^2} \right| \right| \leq 1.$$

The above estimates are sharp.

Zhang et al. [40] introduced the following class of g -starlike mapping of complex order λ on \mathbb{B} in X , which has been introduced by Hu et al. [22] on \mathbb{B}^n .

Definition 1.9 [40]. Let $g : \mathbb{U} \rightarrow \mathbb{C}$ be a biholomorphic function such that $g(0) = 1$, $\Re g(\xi) > 0$ on \mathbb{U} . Let $\lambda \in \mathbb{C}$ with $\Re \lambda \leq 0$ and let $f : \mathbb{B} \rightarrow X$ be a normalized locally biholomorphic mapping. If

$$(1 - \lambda) \frac{\|x\|}{T_x(Df(x))^{-1} f(x)} + \lambda \in g(\mathbb{U}), \quad x \in \mathbb{B} \setminus \{0\}, T_x \in T(x),$$

then f is called a g -starlike mapping of complex order λ .

Let $\mathcal{S}_{g,\lambda}^*(\mathbb{B})$ denote the class of g -starlike mapping of complex order λ on \mathbb{B} . In particular, when $X = \mathbb{C}$, $\mathbb{B} = \mathbb{U}$, the above relation implies that

$$f \in \mathcal{S}_{g,\lambda}^*(\mathbb{U}) \quad \text{if and only if} \quad (1 - \lambda) \frac{\xi f'(\xi)}{f(\xi)} + \lambda \prec g, \quad \xi \in \mathbb{U}.$$

In view of Remark 2.4 of [40], we know that some important subclasses of $\mathcal{S}(\mathbb{B})$ coincide with the classes of $\mathcal{S}_{g,\lambda}^*(\mathbb{B})$ for certain choices of g and λ .

Since the authors [37; 39] used Definitions 1.2 and 1.3 directly, the proofs of Theorems 1.6 and 1.8 are long and rather complicated. In Section 3, under the same

conditions as in Theorems 1.6 and 1.8, we will establish the corresponding inequalities of norm type and functional type for the subclasses of close-to-convex mappings of type B and close-to-starlike mappings on the open unit ball of a Banach space, respectively. Next, In Section 4, we obtain the coefficient inequalities for a subclass of g -starlike mappings of complex order λ on the open unit ball of a complex Banach space. The various results of this paper would generalize many known results. Moreover, the proof methods presented here simplify those appeared in some earlier papers [25; 26; 28; 33; 36; 37; 39].

Some investigations concerning the coefficient estimates for subclasses of holomorphic mappings in several variables have been obtained by Bracci et al. [1; 2], Graham et al. [8; 9; 10; 11; 12; 13; 14], Hamada et al. [16; 18; 19], Kohr [24], Liu and Wu [27], Liu et al. [28], and Xu et al. [34; 35].

2. Some lemmas

In order to prove the desired results, we need to provide the following lemmas.

Lemma 2.1 [17]. *Let $h : \mathbb{B} \rightarrow X$ be a normalized locally biholomorphic mapping. Then h is a starlike mapping on \mathbb{B} if and only if*

$$\Re(T_x(Dh(x)^{-1}h(x))) > 0, \quad x \in \mathbb{B} \setminus \{0\}, T_x \in T(x).$$

Comparing Lemma 2.1 with Definition 1.3, we remark that any normalized starlike mapping on \mathbb{B} is close-to-starlike (with respect to itself).

Lemma 2.2. *Let $g : \mathbb{U} \rightarrow \mathbb{C}$ satisfy the conditions of Definition 1.9, $f \in H(\mathbb{B}, \mathbb{C})$, $f(0) = 1$, $F(x) = xf(x)$. Fix $x \in \mathbb{B} \setminus \{0\}$ and denote $x_0 = \frac{x}{\|x\|}$. Let $l(\xi) = T_x(F(\xi x_0))$, $\xi \in \mathbb{U}$. Then*

$$l \in \mathcal{S}_{g,\lambda}^*(\mathbb{U}) \Leftrightarrow F \in \mathcal{S}_{g,\lambda}^*(\mathbb{B}).$$

Proof. Since $F \in \mathcal{S}_{g,\lambda}^*(\mathbb{B})$, we deduce from Definition 1.9 that

$$(1 - \lambda) \frac{\|x\|}{T_x((DF(x))^{-1}F(x))} + \lambda \in g(\mathbb{U}), \quad x \in \mathbb{B} \setminus \{0\}, T_x \in T(x).$$

It follows that F is locally biholomorphic on \mathbb{B} , and thus $f(x) \neq 0$, $x \in \mathbb{B}$. Using a similar method to that in [8] (also see [9, Theorem 7.1.14]), we have

$$(2-1) \quad [DF(x)]^{-1} = \frac{1}{f(x)} \left(I - \frac{\frac{x Df(x)}{f(x)}}{1 + \frac{Df(x)x}{f(x)}} \right).$$

Hence,

$$(DF(x))^{-1}F(x) = x \left(\frac{1}{1 + \frac{Df(x)x}{f(x)}} \right) = \frac{xf(x)}{f(x) + Df(x)x}, \quad x \in \mathbb{B}.$$

We find from the above equality that

$$(2-2) \quad (1 - \lambda) \frac{\|x\|}{T_x((DF(x))^{-1}F(x))} + \lambda = (1 - \lambda) \frac{f(x) + Df(x)x}{f(x)} + \lambda \in g(\mathbb{U}).$$

Since $l(\xi) = T_x(F(\xi x_0)) = T_x(\xi x_0 f(\xi x_0)) = \xi f(\xi x_0)$, we have

$$l'(\xi) = f(\xi x_0) + Df(\xi x_0) \xi x_0$$

and

$$(2-3) \quad (1 - \lambda) \frac{\xi l'(\xi)}{l(\xi)} + \lambda = (1 - \lambda) \frac{f(\xi x_0) + Df(\xi x_0) \xi x_0}{f(\xi x_0)} + \lambda \in g(\mathbb{U}),$$

which implies that $l \in \mathcal{S}_{g,\lambda}^*(\mathbb{U})$.

Conversely, we assume $l \in \mathcal{S}_{g,\lambda}^*(\mathbb{U})$. Then it is clear that $\frac{\xi l'(\xi)}{l(\xi)} \neq 0$, $\xi \in \mathbb{U}$. Hence we have

$$1 + \frac{Df(x)x}{f(x)} \neq 0, \quad x \in \mathbb{B}.$$

It is not hard to deduce from this and (2-1) that F is locally biholomorphic on \mathbb{B} .

On the other hand, in view of (2-2) and (2-3), we can conclude that

$$(1 - \lambda) \frac{\xi}{T_{\xi x_0}((DF(\xi x_0))^{-1}F(\xi x_0))} + \lambda = (1 - \lambda) \frac{\xi l'(\xi)}{l(\xi)} + \lambda \in g(\mathbb{U}).$$

Taking $\xi = \|x\|$ in the above relation, we obtain

$$(1 - \lambda) \frac{\|x\|}{T_x((DF(x))^{-1}F(x))} + \lambda \in g(\mathbb{U}),$$

as desired. □

Lemma 2.3 [20]. *Let $g(\xi) = 1 + g'(0)\xi + \frac{g''(0)}{2}\xi^2 + \dots$ be a holomorphic function on \mathbb{U} such that $g'(0) \neq 0$. Let $s(\xi) = 1 + s'(0)\xi + \frac{s''(0)}{2}\xi^2 + \dots$ be a holomorphic function on \mathbb{U} such that $s \prec g$. Then for every $\mu \in \mathbb{C}$, it holds that*

$$\left| \frac{s''(0)}{2} - \mu(s'(0))^2 \right| \leq \max \left\{ |g'(0)|, \left| \frac{g''(0)}{2} - \mu(g'(0))^2 \right| \right\}.$$

This estimate is sharp.

Lemma 2.4. *Let $g : \mathbb{U} \rightarrow \mathbb{C}$ satisfy the conditions of Definition 1.9 and*

$$l(\xi) = \xi + \sum_{m=2}^{\infty} l_m \xi^m \in \mathcal{S}_{g,\lambda}^*(\mathbb{U}).$$

Then

$$|l_3 - \nu l_2^2| \leq \frac{|g'(0)|}{2|1-\lambda|} \max \left\{ 1, \left| \frac{1}{2} \frac{g''(0)}{g'(0)} + \frac{1-2\nu}{1-\lambda} g'(0) \right| \right\}, \quad \nu \in \mathbb{C}.$$

The above estimate is sharp for the function

$$l(\xi) = \xi \exp \frac{1}{1-\lambda} \int_0^\xi (g(t) - 1) \frac{1}{t} dt$$

if $\left| \frac{1}{2} \frac{g''(0)}{g'(0)} + \frac{1-2\nu}{1-\lambda} g'(0) \right| \geq 1$, and for

$$l(\xi) = \xi \exp \frac{1}{1-\lambda} \int_0^\xi (g(t^2) - 1) \frac{1}{t} dt$$

if $\left| \frac{1}{2} \frac{g''(0)}{g'(0)} + \frac{1-2\nu}{1-\lambda} g'(0) \right| \leq 1$.

Proof. Since $l \in \mathcal{S}_{g,\lambda}^*(\mathbb{U})$, we have

$$b(\xi) = (1-\lambda) \frac{\xi l'(\xi)}{l(\xi)} + \lambda \in g(\mathbb{U}), \quad \xi \in \mathbb{U}, \quad b \prec g.$$

A computation shows that

$$b'(0) = (1-\lambda) l_2, \quad \frac{b''(0)}{2} = 2(1-\lambda) l_3 - (1-\lambda) l_2^2.$$

By using Lemma 2.3, we have

$$\left| \frac{b''(0)}{2} - \mu(b'(0))^2 \right| \leq \max \left\{ |g'(0)|, \left| \frac{g''(0)}{2} - \mu(g'(0))^2 \right| \right\}, \quad \mu \in \mathbb{C}.$$

From the above relations, we obtain that

$$|l_3 - \nu l_2^2| \leq \frac{|g'(0)|}{2|1-\lambda|} \max \left\{ 1, \left| \frac{1}{2} \frac{g''(0)}{g'(0)} + \frac{1-2\nu}{1-\lambda} g'(0) \right| \right\}, \quad \nu \in \mathbb{C}. \quad \square$$

Remark 2.5. Lemma 2.4 generalizes Theorem 3.1 of [36], when $\lambda = 0$, Lemma 2.4 obtained by Xu et al. [36]. Moreover, the proof presented here is simpler than that in [36, Theorem 3.1].

Lemma 2.6. Let $g : \mathbb{U} \rightarrow \mathbb{C}$ be a convex function which satisfies the conditions of Definition 1.9 and

$$l(\xi) = \xi + \sum_{m=2}^{\infty} l_m \xi^m \in \mathcal{S}_{g,\lambda}^*(\mathbb{U}).$$

Then

$$|l_m| \leq \frac{1}{(m-1)!} \prod_{k=2}^m \left((k-2) + \frac{|g'(0)|}{|1-\lambda|} \right), \quad m = 2, 3, 4, \dots$$

Proof. Since $l \in \mathcal{S}_{g,\lambda}^*(\mathbb{U})$, the function p is defined by

$$p(\xi) = (1-\lambda) \frac{\xi l'(\xi)}{l(\xi)} + \lambda, \quad \xi \in \mathbb{U},$$

in view of Definition 1.9, we have $p \prec g$. If

$$p(\xi) = 1 + p_1 \xi + p_2 \xi^2 + \dots + p_m \xi^m + \dots, \quad \xi \in \mathbb{U},$$

then from Rogosinski's theorem [30], we obtain that $|p_m| \leq |g'(0)|$, $m = 1, 2, 3, \dots$. Comparing the coefficients in the power series of $l(\xi)p(\xi) = (1 - \lambda)\xi l'(\xi) + \lambda l(\xi)$, we deduce that $(1 - \lambda)l_2 = p_1$ and

$$(1 - \lambda)(m - 1)l_m = p_{m-1} + l_2 p_{m-2} + l_3 p_{m-3} + \dots + l_{m-1} p_1, \quad m = 2, 3, 4, \dots$$

Thus by the mathematical induction, we obtain

$$|l_m| \leq \frac{1}{(m - 1)!} \prod_{k=2}^m \left((k - 2) + \frac{|g'(0)|}{|1 - \lambda|} \right), \quad m = 2, 3, 4, \dots,$$

as desired. □

3. Simplified proofs of Fekete–Szegő inequalities for close-to-quasiconvex mappings of type B and close-to-starlike mappings

In this section, by using the proof methods different from those appeared in [39] and [37], we obtain the corresponding results of norm type and functional type for subclasses of close-to-quasiconvex mappings of type B and close-to-starlike mappings defined on the open unit ball in a complex Banach space (see Theorem 1.7).

Theorem 3.1. *Let $f : \mathbb{B} \rightarrow \mathbb{C}$, $h : \mathbb{B} \rightarrow \mathbb{C}$ be holomorphic functions, and let $H(x) = xh(x) \in \mathcal{Q}_B(\mathbb{B})$. Suppose that $F(x) = xf(x)$ is a close-to-quasiconvex mapping of type B with respect to $H(x)$. Then, for $x \in X \setminus \{0\}$, $T_x \in T(x)$ and $\lambda \in [0, 1]$, we have*

$$\left\| \frac{D^3 F(0)(x^3)}{3! \|x\|^3} - \lambda \frac{1}{2 \|x\|^3} D^2 F(0) \left(x, \frac{D^2 F(0)(x^2)}{2!} \right) \right\| \leq \begin{cases} 3 - 4\lambda, & \lambda \in [0, \frac{1}{3}], \\ \frac{1}{3} + \frac{4}{9\lambda}, & \lambda \in [\frac{1}{3}, \frac{2}{3}], \\ 1, & \lambda \in [\frac{2}{3}, 1], \end{cases}$$

$$\left| \frac{T_x(D^3 F(0)(x^3))}{3! \|x\|^3} - \lambda \left(\frac{T_x(D^2 F(0)(x^2))}{2! \|x\|^2} \right)^2 \right| \leq \begin{cases} 3 - 4\lambda, & \lambda \in [0, \frac{1}{3}], \\ \frac{1}{3} + \frac{4}{9\lambda}, & \lambda \in [\frac{1}{3}, \frac{2}{3}], \\ 1, & \lambda \in [\frac{2}{3}, 1], \end{cases}$$

$$\left| \frac{T_x(D^3 F(0)(x^3))}{3! \|x\|^3} \right| - \left| \frac{T_x(D^2 F(0)(x^2))}{2! \|x\|^2} \right| \leq 1$$

and

$$\left\| \frac{D^3 F(0)(x^3)}{3! \|x\|^3} \right\| - \left\| \frac{D^2 F(0)(x^2)}{2! \|x\|^2} \right\| \leq 1.$$

The above estimates are sharp.

Proof. Fix $x \in X \setminus \{0\}$ and denote $x_0 = \frac{x}{\|x\|}$. Let $p : \mathbb{U} \rightarrow \mathbb{C}$ be given by

$$p(\xi) = \begin{cases} \frac{T_x((DH(\xi x_0))^{-1}(D^2 H(\xi x_0)(\xi x_0)^2 + DH(\xi x_0)(\xi x_0)))}{\xi}, & \xi \neq 0, \\ 1, & \xi = 0. \end{cases}$$

Then $p \in H(\mathbb{U})$, $p(0) = 1$, and

$$\begin{aligned} p(\xi) &= \frac{T_x((DH(\xi x_0))^{-1}(D^2H(\xi x_0)(\xi x_0)^2 + DH(\xi x_0)(\xi x_0))}{\xi} \\ &= \frac{T_{\xi x_0}((DH(\xi x_0))^{-1}(D^2H(\xi x_0)(\xi x_0)^2 + DH(\xi x_0)(\xi x_0))}{\|\xi x_0\|}, \quad \xi \in \mathbb{U} \setminus \{0\}. \end{aligned}$$

Since $H \in \mathcal{Q}_B(\mathbb{B})$, using Definition 1.1, we obtain

$$(3-1) \quad \Re e(p(\xi)) > 0, \quad \xi \in \mathbb{U}.$$

On the other hand, by using an elementary computation, we have

$$(DH(x))^{-1} = \frac{1}{h(x)} \left(I - \frac{\frac{x Dh(x)}{h(x)}}{1 + \frac{Dh(x)x}{h(x)}} \right)$$

and

$$(DH(x))^{-1}(D^2H(x)(x^2) + DH(x)x) = \frac{D^2h(x)(x^2) + 3Dh(x)x + h(x)}{h(x) + Dh(x)x}x.$$

This allows us to rewrite p in the form

$$(3-2) \quad p(\xi) = \frac{D^2h(\xi x_0)((\xi x_0)^2) + 3Dh(\xi x_0)(\xi x_0) + h(\xi x_0)}{h(\xi x_0) + Dh(\xi x_0)(\xi x_0)}, \quad \xi \in \mathbb{U}.$$

Let $k(\xi) = T_x(H(\xi x_0)) = \xi h(\xi x_0)$ for $\xi \in \mathbb{U}$. Elementary computations using this inequality yield that

$$(3-3) \quad k'(\xi) = h(\xi x_0) + Dh(\xi x_0)(\xi x_0)$$

and

$$(3-4) \quad 1 + \frac{\xi k''(\xi)}{k'(\xi)} = \frac{D^2h(\xi x_0)((\xi x_0)^2) + 3Dh(\xi x_0)(\xi x_0) + h(\xi x_0)}{h(\xi x_0) + Dh(\xi x_0)(\xi x_0)}.$$

Using (3-1), (3-2) and (3-4), we obtain $k \in \mathcal{K}$.

Let $s(x) = (DH(x))^{-1}DF(x)x$, and let

$$r(\xi) = \begin{cases} \frac{T_x(s(\xi x_0))}{\xi}, & \xi \in \mathbb{U} \setminus \{0\}, \\ 1, & \xi = 0. \end{cases}$$

Then r is holomorphic on \mathbb{U} , $r(0) = 1$ and

$$r(\xi) = \frac{T_x(s(\xi x_0))}{\xi} = \frac{T_{x_0}(s(\xi x_0))}{\xi} = \frac{T_{\xi x_0}(s(\xi x_0))}{\|\xi x_0\|}, \quad \xi \in \mathbb{U} \setminus \{0\}.$$

Since $F(x)$ is a close-to-convex mapping of type B with respect to $H(x)$, from Definition 1.2, we obtain

$$(3-5) \quad \Re e(r(\xi)) > 0, \quad \xi \in \mathbb{U}.$$

A simple computation shows that

$$(3-6) \quad r(\xi) = \frac{T_x(s(\xi x_0))}{\xi} = \frac{T_x((DH(\xi x_0))^{-1}DF(\xi x_0)\xi x_0)}{\xi} \\ = \frac{f(\xi x_0) + Df(\xi x_0)\xi x_0}{h(\xi x_0) + Dh(\xi x_0)\xi x_0}, \quad \xi \in \mathbb{U} \setminus \{0\}.$$

Letting $l(\xi) = T_x(F(\xi x_0))$, $\xi \in \mathbb{U}$, we have

$$(3-7) \quad l'(\xi) = T_x(DF(\xi x_0)x_0) = f(\xi x_0) + Df(\xi x_0)\xi x_0.$$

Hence from (3-3), (3-5), (3-6) and (3-7), we obtain

$$\Re e\left(\frac{l'(\xi)}{k'(\xi)}\right) = \Re e\left(\frac{f(\xi x_0) + Df(\xi x_0)\xi x_0}{h(\xi x_0) + Dh(\xi x_0)\xi x_0}\right) > 0,$$

which means that $l \in \mathcal{C}$. Thus from Theorem 1.5, we have

$$(3-8) \quad \left| \frac{l'''(0)}{3!} - \lambda \left(\frac{l''(0)}{2!} \right)^2 \right| \leq \begin{cases} 3 - 4\lambda, & \lambda \in [0, \frac{1}{3}], \\ \frac{1}{3} + \frac{4}{9\lambda}, & \lambda \in [\frac{1}{3}, \frac{2}{3}], \\ 1, & \lambda \in [\frac{2}{3}, 1] \end{cases}$$

and

$$(3-9) \quad \left| \left| \frac{l'''(0)}{3!} \right| - \left| \frac{l''(0)}{2!} \right| \right| \leq 1.$$

Furthermore, since $l(\xi) = T_x(F(\xi x_0)) = T_x(\xi x_0 f(\xi x_0)) = \xi f(\xi x_0)$ for $\xi \in \mathbb{U}$, a simple computation yields that

$$\frac{l'''(0)}{3!} = \frac{T_x(D^3F(0)(x_0^3))}{3!} = \frac{D^2f(0)(x_0^2)}{2}, \\ \left(\frac{l''(0)}{2!} \right)^2 = \left(\frac{T_x(D^2F(0)(x_0^2))}{2!} \right)^2 = (Df(0)(x_0))^2,$$

and

$$\frac{D^3F(0)(x_0^3)}{3!} = \frac{D^2f(0)(x_0^2)}{2!}x_0, \quad \frac{1}{2}D^2F(0)\left(x_0, \frac{D^2F(0)(x_0^2)}{2!}\right) = (Df(0)(x_0))^2x_0.$$

Using the above equalities, (3-8) and (3-9), we obtain all of the desired conclusions about Theorems 3.1. The example which shows the sharpness of Theorem 3.1 is the same as the mapping defined in [37]. This completes the proof of Theorem 3.1. \square

Theorem 3.2. *Let $f : \mathbb{B} \rightarrow \mathbb{C}$, $h : \mathbb{B} \rightarrow \mathbb{C}$ be holomorphic functions, and let $H(x) = xh(x) \in \mathcal{S}^*(\mathbb{B})$. Suppose that $F(x) = xf(x)$ is a close-to-starlike mapping with respect to $H(x)$. Then, for $x \in X \setminus \{0\}$, $T_x \in T(x)$ and $\lambda \in [0, 1]$, we have the same conclusions as in Theorem 3.1.*

Proof. Fix $x \in X \setminus \{0\}$ and denote $x_0 = \frac{x}{\|x\|}$. Let $p : \mathbb{U} \rightarrow \mathbb{C}$ be given by

$$p(\xi) = \begin{cases} \frac{T_x((DH(\xi x_0))^{-1}H(\xi x_0))}{\xi}, & \xi \neq 0, \\ 1, & \xi = 0. \end{cases}$$

Then $p \in H(\mathbb{U})$, $p(0) = 1$, and

$$p(\xi) = \frac{T_x((DH(\xi x_0))^{-1}H(\xi x_0))}{\xi} = \frac{T_{\xi x_0}((DH(\xi x_0))^{-1}H(\xi x_0))}{\|\xi x_0\|}, \quad \xi \in \mathbb{U} \setminus \{0\}.$$

Since $H \in \mathcal{S}^*(\mathbb{B})$, by Lemma 2.1, we have

$$(3-10) \quad \Re e(p(\xi)) > 0, \quad \xi \in \mathbb{U}.$$

At the same time, a short computation yields the relation

$$(DH(x))^{-1}H(x) = \frac{h(x)}{h(x) + Dh(x)x}x.$$

Hence the above relations imply that

$$(3-11) \quad p(\xi) = \frac{h(\xi x_0)}{h(\xi x_0) + Dh(\xi x_0)(\xi x_0)}, \quad \xi \in \mathbb{U}.$$

Let

$$k(\xi) = T_x(H(\xi x_0)) = \xi h(\xi x_0), \quad \xi \in \mathbb{U}.$$

Then, we have

$$k'(\xi) = h(\xi x_0) + Dh(\xi x_0)(\xi x_0)$$

and

$$(3-12) \quad \frac{\xi k'(\xi)}{k(\xi)} = \frac{h(\xi x_0) + Dh(\xi x_0)(\xi x_0)}{h(\xi x_0)}.$$

By using (3-10), (3-11) and (3-12), we obtain $k \in \mathcal{S}^*$.

Let $s(x) = (DF(x))^{-1}H(x)$, and let

$$r(\xi) = \begin{cases} \frac{T_x(s(\xi x_0))}{\xi}, & \xi \in \mathbb{U} \setminus \{0\}, \\ 1, & \xi = 0. \end{cases}$$

Then r is holomorphic on \mathbb{U} , $r(0) = 1$ and

$$r(\xi) = \frac{T_x(s(\xi x_0))}{\xi} = \frac{T_{x_0}(s(\xi x_0))}{\xi} = \frac{T_{\xi x_0}(s(\xi x_0))}{\|\xi x_0\|}, \quad \xi \in \mathbb{U} \setminus \{0\}.$$

Since $F(x)$ is a close-to-starlike mapping with respect to $H(x)$, from Definition 1.3, we obtain

$$\Re e(r(\xi)) > 0, \quad \xi \in \mathbb{U}.$$

A simple computation shows that

$$\begin{aligned} r(\xi) &= \frac{T_x(s(\xi x_0))}{\xi} = \frac{T_x((DF(\xi x_0))^{-1}H(\xi x_0))}{\xi} \\ &= \frac{h(\xi x_0)}{f(\xi x_0) + Df(\xi x_0)\xi x_0}, \quad \xi \in \mathbb{U} \setminus \{0\}. \end{aligned}$$

Letting $l(\xi) = T_x(F(\xi x_0))$, $\xi \in \mathbb{U}$ again, we obtain

$$l'(\xi) = T_x(DF(\xi x_0)x_0) = f(\xi x_0) + Df(\xi x_0)\xi x_0.$$

Consequently, combining this observation with the preceding relations, we have

$$\Re e\left(\frac{\xi l'(\xi)}{k(\xi)}\right) = \Re e\left(\frac{f(\xi x_0) + Df(\xi x_0)\xi x_0}{h(\xi x_0)}\right) > 0,$$

which implies that $l \in \mathcal{C}$. The remaining part of the proof of Theorem 3.2 is similar to that in the proof of Theorem 3.1, so we omit the details. □

4. Simplified proofs of coefficient inequalities for a subclass of g -starlike mappings of complex order λ

By using Lemmas 2.2, 2.4 and 2.6, we establish bounds of all terms of homogeneous expansions and the Fekete–Szegő inequality for a subclass of g -starlike mappings of complex order λ on the open unit ball of a complex Banach space, which generalize the corresponding results appeared in [25; 26; 28; 33; 36].

Theorem 4.1. *Let $g : \mathbb{U} \rightarrow \mathbb{C}$ be a convex function which satisfies the conditions of Definition 1.9, and $f \in H(\mathbb{B}, \mathbb{C})$, $f(0) = 1$. Suppose that $F(x) = xf(x) \in \mathcal{S}_{g,\lambda}^*(\mathbb{B})$. Then for $x \in \mathbb{B}$, we have*

$$\frac{\|D^m F(0)(x^m)\|}{m!} \leq \frac{\prod_{r=2}^m \left[r - 2 + \frac{1}{1-\lambda} |g'(0)|\right]}{(m-1)!} \|x\|^m, \quad m = 2, 3, 4, \dots$$

Proof. Fix $x \in \mathbb{B} \setminus \{0\}$ and denote $x_0 = \frac{x}{\|x\|}$. Let $l(\xi) = T_x(F(\xi x_0))$, $\xi \in \mathbb{U}$. In view of Lemma 2.2, we have $l \in \mathcal{S}_{g,\lambda}^*(\mathbb{U})$. Since $l(\xi) = T_x(F(\xi x_0)) = \xi f(\xi x_0)$, we obtain

$$\frac{D^m F(0)(x_0^m)}{m!} = x_0 \frac{D^{m-1} f(0)(x_0^{m-1})}{(m-1)!} = \frac{l^{(m)}(0)}{m!}.$$

By using Lemma 2.6, we have

$$\frac{\|D^m F(0)(x^m)\|}{m!} \leq \frac{\prod_{r=2}^m \left[r - 2 + \frac{1}{1-\lambda} |g'(0)|\right]}{(m-1)!} \|x\|^m, \quad m = 2, 3, 4, \dots,$$

as desired. □

Remark 4.2. Theorem 4.1 generalizes many known results. In Theorem 4.3 if we set $\lambda = 0$ and $g(\xi) = \frac{1+\xi}{1-\xi}$, $\xi \in \mathbb{U}$, $\lambda = 0$ and $\lambda = -\frac{\alpha}{1-\alpha}$, $0 \leq \alpha < 1$, we can readily deduce the corresponding results of [28], [26] and [33], respectively. Moreover, the proofs presented here are simpler than those given in [28], [26] and [33].

Theorem 4.3. *Let $g : \mathbb{U} \rightarrow \mathbb{C}$ satisfy the conditions of Definition 1.9, and let $f \in H(\mathbb{B}, \mathbb{C})$, $f(0) = 1$. Suppose that $F(x) = xf(x) \in \mathcal{S}_{g,\lambda}^*(\mathbb{B})$. Then for $v \in \mathbb{C}$, $x \in \mathbb{B} \setminus \{0\}$, we have*

$$\begin{aligned} \left| \frac{D^3 F(0)(x^3)}{3! \|x\|^3} - v \frac{1}{2 \|x\|^3} D^2 F(0) \left(x, \frac{D^2 F(0)(x^2)}{2!} \right) \right| \\ \leq \frac{|g'(0)|}{2|1-\lambda|} \max \left\{ 1, \left| \frac{1}{2} \frac{g''(0)}{g'(0)} + \frac{1-2v}{1-\lambda} g'(0) \right| \right\} \end{aligned}$$

and

$$\begin{aligned} \left| \frac{T_x(D^3 F(0)(x^3))}{3! \|x\|^3} - v \left(\frac{T_x(D^2 F(0)(x^2))}{2! \|x\|^2} \right)^2 \right| \\ \leq \frac{|g'(0)|}{2|1-\lambda|} \max \left\{ 1, \left| \frac{1}{2} \frac{g''(0)}{g'(0)} + \frac{1-2v}{1-\lambda} g'(0) \right| \right\}. \end{aligned}$$

The above estimates are sharp.

Proof. Fix $x \in \mathbb{B} \setminus \{0\}$ and denote $x_0 = \frac{x}{\|x\|}$. Let $l(\xi) = T_x(F(\xi x_0))$, $\xi \in \mathbb{U}$. From Lemma 2.2, we have $l \in \mathcal{S}_{g,\lambda}^*(\mathbb{U})$. Since $l(\xi) = T_x(F(\xi x_0)) = \xi f(\xi x_0)$, we have

$$\begin{aligned} \frac{l'''(0)}{3!} &= \frac{T_x(D^3 F(0)(x_0^3))}{3!} = \frac{D^2 f(0)(x_0^2)}{2}, \\ \left(\frac{l''(0)}{2!} \right)^2 &= \left(\frac{T_x(D^2 F(0)(x_0^2))}{2!} \right)^2 = (Df(0)(x_0))^2, \end{aligned}$$

and

$$\frac{D^3 F(0)(x_0^3)}{3!} = \frac{D^2 f(0)(x_0^2)}{2!} x_0, \quad \frac{1}{2} D^2 F(0) \left(x_0, \frac{D^2 F(0)(x_0^2)}{2!} \right) = (Df(0)(x_0))^2 x_0.$$

Using the above equalities and Lemma 2.4, we obtain the desired conclusion.

In order to prove that the estimates of Theorem 4.3 are sharp, it suffices to consider the following examples.

If $\left| \frac{1}{2} \frac{g''(0)}{g'(0)} + \frac{1-2v}{1-\lambda} g'(0) \right| \geq 1$, we consider the example

$$F(x) = x \exp \frac{1}{1-\lambda} \int_0^{T_u(x)} (g(t) - 1) \frac{dt}{t}, \quad x \in \mathbb{B}, \|u\| = 1.$$

If $\left| \frac{1}{2} \frac{g''(0)}{g'(0)} + \frac{1-2v}{1-\lambda} g'(0) \right| \leq 1$, we consider the example

$$F(x) = x \exp \frac{1}{1-\lambda} \int_0^{T_u(x)} (g(t^2) - 1) \frac{dt}{t}, \quad x \in \mathbb{B}, \|u\| = 1. \quad \square$$

Remark 4.4. If we set $\lambda = -i \tan \beta$, $-\frac{\pi}{2} < \beta < \frac{\pi}{2}$ and $\lambda = 0$ in Theorem 4.3, we obtain the corresponding results of [25] and [36], respectively. Moreover, the proofs presented here are simpler than those given in [25] and [36].

References

- [1] F. Bracci, “Shearing process and an example of a bounded support function in $S^0(\mathbb{B}^2)$ ”, *Comput. Methods Funct. Theory* **15**:1 (2015), 151–157. MR Zbl
- [2] F. Bracci, I. Graham, H. Hamada, and G. Kohr, “Variation of Loewner chains, extreme and support points in the class S^0 in higher dimensions”, *Constr. Approx.* **43**:2 (2016), 231–251. MR Zbl
- [3] R. Długosz and P. Liczberski, “Some results of Fekete–Szegő type for Bavrin’s families of holomorphic functions in \mathbb{C}^n ”, *Ann. Mat. Pura Appl.* (4) **200**:4 (2021), 1841–1857. MR Zbl
- [4] R. Długosz and P. Liczberski, “Fekete–Szegő problem for Bavrin’s functions and close-to-starlike mappings in \mathbb{C}^n ”, *Anal. Math. Phys.* **12**:4 (2022), art. id. 103. MR Zbl
- [5] M. Elin and F. Jacobzon, “The Fekete–Szegő problem for spirallike mappings and non-linear resolvents in Banach spaces”, *Stud. Univ. Babeş-Bolyai Math.* **67**:2 (2022), 329–344. MR Zbl
- [6] M. Elin and F. Jacobzon, “Note on the Fekete–Szegő problem for spirallike mappings in Banach spaces”, *Results Math.* **77**:3 (2022), art. id. 137. MR Zbl
- [7] M. Elin, S. Reich, and D. Shoikhet, *Numerical range of holomorphic mappings and applications*, Birkhäuser, Cham, 2019. MR Zbl
- [8] I. Graham and G. Kohr, *Geometric function theory in one and higher dimensions*, Monographs and Textbooks in Pure and Applied Mathematics **255**, Marcel Dekker, New York, 2003. MR Zbl
- [9] I. Graham, H. Hamada, and G. Kohr, “Parametric representation of univalent mappings in several complex variables”, *Canad. J. Math.* **54**:2 (2002), 324–351. MR Zbl
- [10] I. Graham, G. Kohr, and M. Kohr, “Loewner chains and parametric representation in several complex variables”, *J. Math. Anal. Appl.* **281**:2 (2003), 425–438. MR Zbl
- [11] I. Graham, H. Hamada, T. Honda, G. Kohr, and K. H. Shon, “Growth, distortion and coefficient bounds for Carathéodory families in \mathbb{C}^n and complex Banach spaces”, *J. Math. Anal. Appl.* **416**:1 (2014), 449–469. MR Zbl
- [12] I. Graham, H. Hamada, G. Kohr, and M. Kohr, “Support points and extreme points for mappings with A -parametric representation in \mathbb{C}^n ”, *J. Geom. Anal.* **26**:2 (2016), 1560–1595. MR Zbl
- [13] I. Graham, H. Hamada, G. Kohr, and M. Kohr, “Bounded support points for mappings with g -parametric representation in \mathbb{C}^2 ”, *J. Math. Anal. Appl.* **454**:2 (2017), 1085–1105. MR Zbl
- [14] I. Graham, H. Hamada, and G. Kohr, “Extremal problems for mappings with g -parametric representation on the unit polydisc in \mathbb{C}^n ”, pp. 141–167 in *Complex analysis and dynamical systems*, Birkhäuser, Cham, 2018. MR Zbl
- [15] H. Hamada, “Fekete–Szegő problems for spirallike mappings and close-to-quasi-convex mappings on the unit ball of a complex Banach space”, *Results Math.* **78**:3 (2023), art. id. 109. MR Zbl
- [16] H. Hamada and T. Honda, “Sharp growth theorems and coefficient bounds for starlike mappings in several complex variables”, *Chinese Ann. Math. Ser. B* **29**:4 (2008), 353–368. MR Zbl
- [17] H. Hamada and G. Kohr, “ Φ -like and convex mappings in infinite dimensional spaces”, *Rev. Roumaine Math. Pures Appl.* **47**:3 (2002), 315–328. MR Zbl

- [18] H. Hamada and G. Kohr, “Support points for families of univalent mappings on bounded symmetric domains”, *Sci. China Math.* **63**:12 (2020), 2379–2398. MR Zbl
- [19] H. Hamada, T. Honda, and G. Kohr, “Growth theorems and coefficients bounds for univalent holomorphic mappings which have parametric representation”, *J. Math. Anal. Appl.* **317**:1 (2006), 302–319. MR Zbl
- [20] H. Hamada, G. Kohr, and M. Kohr, “The Fekete–Szegő problem for starlike mappings and nonlinear resolvents of the Carathéodory family on the unit balls of complex Banach spaces”, *Anal. Math. Phys.* **11**:3 (2021), art. id. 115. MR Zbl
- [21] H. Hamada, G. Kohr, and M. Kohr, “Fekete–Szegő problem for univalent mappings in one and higher dimensions”, *J. Math. Anal. Appl.* **516**:2 (2022), art. id. 126526. MR Zbl
- [22] C. Y. Hu, T. S. Liu, and J. F. Wang, “ g -starlike mappings of complex order λ on the unit ball \mathcal{B}^n ”, *Acta Math. Sinica (Chinese Ser.)* **66**:1 (2023), 149–160. MR
- [23] W. Koepf, “On the Fekete–Szegő problem for close-to-convex functions”, *Proc. Amer. Math. Soc.* **101**:1 (1987), 89–95. MR Zbl
- [24] G. Kohr, “On some best bounds for coefficients of several subclasses of biholomorphic mappings in \mathbb{C}^n ”, *Complex Variables Theory Appl.* **36**:3 (1998), 261–284. MR Zbl
- [25] Y. Lai and Q. Xu, “On the coefficient inequalities for a class of holomorphic mappings associated with spirallike mappings in several complex variables”, *Results Math.* **76**:4 (2021), art. id. 191. MR Zbl
- [26] X. S. Liu and T. S. Liu, “The estimates of all homogeneous expansions for a subclass of biholomorphic mappings which have parametric representation in several complex variables”, *Acta Math. Sin. (Engl. Ser.)* **33**:2 (2017), 287–300. MR Zbl
- [27] M.-S. Liu and F. Wu, “Sharp inequalities of homogeneous expansions of almost starlike mappings of order α ”, *Bull. Malays. Math. Sci. Soc.* **42**:1 (2019), 133–151. MR Zbl
- [28] X. Liu, T. Liu, and Q. Xu, “A proof of a weak version of the Bieberbach conjecture in several complex variables”, *Sci. China Math.* **58**:12 (2015), 2531–2540. MR Zbl
- [29] J. A. Pfaltzgraff and T. J. Suffridge, “Close-to-starlike holomorphic functions of several variables”, *Pacific J. Math.* **57**:1 (1975), 271–279. MR Zbl
- [30] W. Rogosinski, “On the coefficients of subordinate functions”, *Proc. London Math. Soc. (2)* **48** (1943), 48–82. MR Zbl
- [31] K. A. Roper and T. J. Suffridge, “Convexity properties of holomorphic mappings in \mathbb{C}^n ”, *Trans. Amer. Math. Soc.* **351**:5 (1999), 1803–1833. MR Zbl
- [32] Z. Tu and L. Xiong, “Unified solution of Fekete–Szegő problem for subclasses of starlike mappings in several complex variables”, *Math. Slovaca* **69**:4 (2019), 843–856. MR Zbl
- [33] L. Xiong, X. Sima, and D. Ouyang, “Bounds of all terms of homogeneous expansions for a subclass of g -parametric biholomorphic mappings in \mathbb{C}^n ”, *Anal. Math. Phys.* **13**:2 (2023), art. id. 34. MR Zbl
- [34] Q. Xu and T. Liu, “On coefficient estimates for a class of holomorphic mappings”, *Sci. China Ser. A* **52**:4 (2009), 677–686. MR Zbl
- [35] Q. Xu, T. Liu, and X. Liu, “The sharp estimates of homogeneous expansions for the generalized class of close-to-quasi-convex mappings”, *J. Math. Anal. Appl.* **389**:2 (2012), 781–791. MR Zbl
- [36] Q. Xu, T. Liu, and X. Liu, “Fekete and Szegő problem in one and higher dimensions”, *Sci. China Math.* **61**:10 (2018), 1775–1788. MR Zbl
- [37] Q. Xu, W. Fang, W. Feng, and T. Liu, “The Fekete–Szegő inequality and successive coefficients difference for a subclass of close-to-starlike mappings in complex Banach spaces”, *Acta Math. Sci. Ser. B (Engl. Ed.)* **43**:5 (2023), 2075–2088. MR Zbl

- [38] Q. Xu, T. Jiang, and T. Liu, “The refinement of Fekete and Szegő problems for close-to-convex functions and close-to-quasi-convex mappings”, *J. Math. Anal. Appl.* **527**:1 (2023), art. id. 127428. MR Zbl
- [39] Q. Xu, T. Liu, and J. Lu, “The Fekete and Szegő inequality for a class of holomorphic mappings on the unit polydisk in \mathbb{C}^n and its application”, *Complex Var. Elliptic Equ.* **68**:1 (2023), 67–80. MR Zbl
- [40] X. Zhang, S. Feng, T. Liu, and J. Wang, “Loewner chains applied to g -starlike mappings of complex order of complex Banach spaces”, *Pacific J. Math.* **323**:2 (2023), 401–431. MR Zbl

Received August 2, 2023. Revised March 12, 2024.

QINGHUA XU
SCHOOL OF SCIENCE
ZHEJIANG UNIVERSITY OF SCIENCE AND TECHNOLOGY
HANGZHOU
CHINA
xuqh@mail.ustc.edu.cn

XIAOHUA YANG
SCHOOL OF SCIENCE
ZHEJIANG UNIVERSITY OF SCIENCE AND TECHNOLOGY
HANGZHOU
CHINA
yangxhh2024@163.com

TAISHUN LIU
DEPARTMENT OF MATHEMATICS
HUZHOU UNIVERSITY
HUZHOU
CHINA
lts@ustc.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2024 is US \$645/year for the electronic version, and \$875/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 329 No. 1 March 2024

Sp(1)-symmetric hyperkähler quantisation	1
JØRGEN ELLEGAARD ANDERSEN, ALESSANDRO MALUSÀ and GABRIELE REMBADO	
Combinatorics of the tautological lamination	39
DANNY CALEGARI	
Limit theorems and wrapping transforms in bi-free probability theory	63
TAKAHIRO HASEBE and HAO-WEI HUANG	
Tame quasiconformal motions and monodromy	105
YUNPING JIANG, SUDEB MITRA and ZHE WANG	
A characterization and solvability of quasihomogeneous singularities	121
GUORUI MA, STEPHEN S.-T. YAU, QIWEI ZHU and HUIQING ZUO	
Stable value of depth of symbolic powers of edge ideals of graphs	147
NGUYEN CONG MINH, TRAN NAM TRUNG and THANH VU	
Collapsed limits of compact Heisenberg manifolds with sub-Riemannian metrics	165
KENSHIRO TASHIRO	
On the coefficient inequalities for some classes of holomorphic mappings in complex Banach spaces	183
QINGHUA XU, XIAOHUA YANG and TAISHUN LIU	